1 **Title: Approximate Bayesian computation untangles signatures of**
2 **contemporary and historical hybridization between two endangered species**
3

4 Dittberner Hannes[1], Tellier Aurelien[2] and de Meaux Juliette[1*]

5 [1] Institute of Plant Sciences, Zülpicher str. 47b, University of Cologne, Germany.

6 [2] Department of Life Science Systems, Technical University of Munich, Freising,
7 Germany

8 [*] Correspondance to jdemeaux@uni-koeln.de

10 ABSTRACT

Contemporary gene flow, when resumed after a period of isolation, can have
crucial consequences for endangered species, as it can both increase the supply of
adaptive alleles and erode local adaptation. Determining the history of gene flow and
thus the importance of contemporary hybridization, however, is notoriously difficult.
Here, we focus on two endangered plant species, *Arabis nemorensis* and *A. sagittata*,
which hybridize naturally in a sympatric population located on the banks of the Rhine.
Using reduced genome sequencing, we determined the phylogeography of the two taxa
but report only a unique sympatric population. Molecular variation in chloroplast DNA
indicated that *A. sagittata* is the principal receiver of gene flow. Applying classical D-
statistics and its derivatives to whole-genome data of 35 accessions, we detect gene
flow not only in the sympatric population but also among allopatric populations. Using
an Approximate Bayesian computation approach, we identify the model that best
describes the history of gene flow between these taxa. This model shows that low levels
of gene flow have persisted long after speciation. Around 10 000 years ago, gene flow
stopped and a period of complete isolation began. Eventually, a hotspot of
contemporary hybridization was formed in the unique sympatric population. Occasional
sympatry may have helped protect these lineages from extinction in spite of their
extremely low diversity.

29
30

31 INTRODUCTION

Individual taxa do not always evolve in isolation. Interspecific hybridization, when it leads to fertile offspring, can result in the transfer of alleles across species barriers and even allow new species to emerge (Seehausen 2004; Abbott 2017). The footprints of gene flow are detectable in many genera and demes (Seehausen 2004; Marcet-Houben and Gabaldón 2015; Ackermann et al. 2019; Taylor and Larson 2019). Genomic analyses have revealed that species barriers are established progressively, depending on the size and degree of reproductive isolation of the species, with genetic variation being shared over periods of time that are longer than previously thought (Brandvain et al. 2014; Novikova et al. 2016; Edelman et al. 2019; Small et al. 2020). The climatic oscillations of quaternary glaciation cycles have likely contributed to multiple opportunities for hybridization in many taxa (Hewitt 2000).

The evolutionary consequences of hybridization can be manifold, offering a powerful channel for some taxa to capture and quickly fix alleles that have been subjected to selection in other species (Hedrick 2013; Goulet et al. 2017; Suarez-Gonzalez et al. 2018a; Vallejo-Marín and Hiscock 2018). The potent adaptive potential unleashed by hybridization have been confirmed in a number of species (Rieseberg et al. 2003; Baduel et al. 2018; Ma et al. 2019; Marburger et al. 2019). For example, introgression appeared to accelerate local adaptation via the transfer of alleles contributing to the success of a relative of the sunflower (Todesco et al. 2020). In *Heliconius* butterflies, an inversion associating genes that control color patterns has been repeatedly exchanged between species (Edelman et al. 2019). In addition, transgressive phenotypic variation can emerge from the recombination of resident and incoming alleles throughout the genome (Rieseberg et al. 1999; Seehausen 2004). Positive selection is therefore expected to increase introgression rates specifically in and around adaptive loci. Heterogeneous introgression rates, however, can also arise along the genome in the absence of adaptive gene flow (Schumer, Rosenthal, et al. 2018). Covariation between the rate of introgression, recombination and gene density, for example, indicates that selections acts to limit introgression and points to the existence of barriers to gene flow (Brandvain et al. 2014; Schumer, Xu, et al. 2018). In humans, the impact of deleterious alleles was shown to correlate negatively with the frequency of alleles introgressed from the related species *Homo neanderthalis,* whose

population size was low and eventually collapsed (Juric et al. 2016; Steinrücken et al. 2018). Ultimately, high rates of gene flow can introduce maladapted alleles and ultimately impede adaptation (Lenormand 2002; Yeaman 2015; Tigano and Friesen 2016). These alleles, in turn, can select for alleles that will reinforce species barriers to prevent resources from being wasted by producing poorly performing hybrids (Hopkins and Rausher 2012). Gene flow, regardless of whether it promotes or erodes adaptation, is expected to create a heterogeneous pattern of introgression throughout the genome (Martin and Jiggins 2017; Schumer, Rosenthal, et al. 2018).

If gene flow can be either a blessing or a curse, it is particularly crucial to understand its importance for species that have to be protected from extinction. In a context of global climate change, the number of threatened species is expected to increase (Díaz et al. 2019; Eichenberg et al. 2020). At the same time, many species barriers that have so far been maintained by habitat, phenological or behavioral separation are likely to disintegrate, generating unprecedented opportunities for novel episodes of hybridization (Anderson et al. 2012; Chunco 2014). Contemporary hybridization will most affect endangered species if the taxa involved have been separated long enough to acquire distinct ecological and population genomics characteristics. For example, the hybridization of long separated continental sub-species of salmon has been associated with the manifestation of Dobzhansky-Muller incompatibilities (Rougemont and Bernatchez 2018). In *Mimulus*, barriers to gene flow are strong today despite an ancient history of hybridization, but contemporary hybridization has been reported at specific locations (Brandvain et al. 2014; Kenney and Sweigart 2016). The potential of hybridization to create a novel genetic make-up today that may impact future evolution and support the evolutionary rescue of biodiversity depends, therefore, on how much time has elapsed during the period of isolation.

Determining the history of gene flow between taxa is a complex task. Dating gene flow cannot be achieved with a single summary statistic of genomic variation, because effective population sizes, migration rates and/or the time since species formation jointly influence patterns of standing variation, divergence and expected allele sharing. The widely used D-statistics tests for gene flow, while accounting for

incomplete lineage sorting at the time of speciation (Durand et al. 2011). This and
related statistics have, for example, revealed the extent of gene flow between species
thought to be separated by differences in ploidy (Arnold et al. 2016; Paape et al. 2018;
Kryvokhyzha et al. 2019). Yet such statistics do not allow speciation or gene flow to be
dated, nor do they evaluate the duration of isolation periods (Schumer, Rosenthal, et al.
2018; Hibbins and Hahn 2019). Failing to understand the correct history of speciation
and gene flow can lead to confounding neutral patterns of gene flow with convergent
evolution, sympatric speciation or even adaptive divergence (Bierne et al. 2013; Ravinet
et al. 2017).

The evolutionary and demographic histories of hybridizing species are often too
complex to be determined analytically. Intensive simulation methods, such as
Approximate Bayesian Computation (ABC), offer a powerful alternative (Csilléry et al.
2010). It not only makes it possible to choose among competing models for the one best
able to explain the data, but also enables population parameter (population sizes,
migration rates) and their fluctuation over time to be estimated (Rougemont and
Bernatchez 2018). However, ABC approaches have rarely been applied to all genome
data and when, then only to infer relatively simple demographic events such as
bottlenecks or expansions (Jay et al. 2019). The current limitation stems from the
inherent statistical complexity of the approach (high dimensionality of whole-genome
data), the necessity for an educated guess regarding the choice of summary statistics to
use, and the lack of user-friendly ready-to-use software capable of handling all genome
data. The potential of ABC approaches to determine the timing, intensity and duration of
hybridization episodes based on all genome data thus remains to be fully leveraged.

Here, we focused on a documented case of contemporary hybridization
between two endangered plant species and reconstructed the history of gene flow.
The species we examined, *Arabis nemorensis* and *A. sagittata,* are selfing biennial
forbs from the Brassicaceae family. *A. nemorensis*, which is strictly confined to
floodplain environments in Central and Northern Europe, harbors extremely low levels
of nucleotide diversity (Dittberner et al. 2019). Since the 1950s, conversion to arable
land and management intensification have reduced the area covered by species–rich
floodplain meadows, the ecosystem in which *A. nemorensis* thrives, by more than 80

%. Only small remnant patches have persisted within protected areas (Hölzel 2005).
The unique ecology of *A. nemorensis*, its shrinking habitat and its selfing mating
system all contribute to the acute danger of extinction this species finds itself in (Hölzel
2005; Burmeier et al. 2010; Mathar et al. 2015). *A. nemorensis*, however, was found to
occur in sympatry with its relative, *A. sagittata*, in a small set of pristine habitat patches
located on the banks of the Rhine near Mainz, Germany (Dittberner et al. 2019). *A.*
sagittata, known to thrive in relatively dry environments, is also endangered, but its
presence in floodplain meadows is novel (Hand and Gregor 2006; Dittberner et al.
2019). Interestingly, approximately 10% of the individuals of the sympatric population
appeared to be hybrids between these two species (Dittberner et al. 2019).

Using a combination of reduced sequencing and whole-genome sequencing,
we asked the following questions: Is hybridization local or is there a large contact
zone? Is gene flow symmetrical? Do rates of introgression vary along the genome? Is
hybridization contemporary or was gene flow continuous throughout the history of the
species? Our study confirms the existence of a contemporary hotspot of hybridization
where asymmetric gene flow has resumed after approximately 10 000 years of
complete isolation between taxa.

RESULTS

Hybrids detected only in a single sympatric population

We combined previously published and newly generated RAD-sequencing data for a total of 231 accessions to describe the phylogeographic distribution of *A. nemorensis* and assess the distribution of *A. sagittata* and interspecific hybrids in the area of distribution of *A. nemorensis* (Table S1). This collection of genotypes covered all sites where the presence of *A. nemorensis* had been documented (see methods). We performed an admixture analysis (Alexander and Lange 2011a) and identified two genetic clusters that were previously assigned to *A. nemorensis* and *A. sagittata* (Figure S1-5, Dittberner et al. 2019). We found individuals of the two species growing in sympatry in only one of the eleven sites (the Rhine population), indicating that opportunities for natural hybridization are restricted (Figure 1A, Dittberner et al. 2019). Of the 140 individuals sampled in the sympatric population, 75 were *A. sagittata*, 42 were *A. nemorensis* and 23 were individuals we identified as having a hybrid origin because they showed less than 95% purity in the output of the ADMIXTURE analysis. Phenotypic observations, such as low seed number per silique and elongated stems, had suggested that hybridization may occur in some rare instances (Novotná and Czapik 1974; Titz 1979). In contrast, genetic analyses indicate that the two species frequently cross-hybridize within the sympatric population (Dittberner et al. 2019). Interestingly, *A. sagittata* ancestry predominates in most hybrid individuals (Figure 1B), indicating the frequent backcross of hybrids with *A. sagittata* individuals, which are more frequent than *A.nemorensis* individuals in the sympatric population.

Chloroplast DNA indicates that *A. nemorensis* is the maternal parent of hybrids

Chloroplast DNA is maternally inherited and its sequence variation can provide information about the maternal genotype of hybrids (McCauley 1995). To determine the maternal genotypes of hybrid individuals, we focused on RADseq stacks that mapped to

a total of 10 kb (5%) of the chloroplast sequence of the close relative *Arabis hirsuta*

(Kawabe et al. 2018). Haplotype network analysis showed that the chloroplast

sequences of *A. sagittata* formed four closely related haplotype groups and two for *A.*

nemorensis (Figure 3C). As all but one interspecific hybrid had an *A. nemorensis*

haplotype, we conclude that *A. nemorensis* is the maternal genotype of most hybrid

individuals. This observation confirms that gene flow between the two clusters is

asymmetrical. Surprisingly, 24 *A. sagittata* individuals were found to carry the

A. nemorensis chloroplast haplotype although the admixture analysis found no trace of

A. nemorensis in their nuclear genome. This suggests that some individuals may have a

hybrid ancestry but their *A. nemorensis* ancestry is undetectable in the admixture

analysis of RAD-seq data, showing in turn that the examination of whole-genome

sequences was necessary.

Figure 1: Hybridization between A. nemorensis *and* A. sagittata

A) Map of sampled populations. Each population is represented by a pie chart showing the average

ancestry proportions of A. nemorensis *and* A. sagittata *in the given population, based on RAD-*

sequencing data. B) Representation of individual ancestry components in the sympatric ("Rhine")

population, based on RAD-seq data. Each bar represents one individual and is colored according to its

genomic ancestry. Bars are ordered by decreasing A. sagittata *ancestry. Hybrid admixed individuals are*

Gene flow between *A. nemorensis* and *A. sagittata* is not restricted to the sympatric population

To quantify interspecific gene flow between the two genetic clusters, we
resequenced 35 whole genomes of accessions from both allopatric and sympatric
populations for the two species, as well as one individual of the closest diploid relative,
A. androsacea (Table S2). In order to understand the history of gene flow, we
specifically excluded individuals that were predicted to be admixed hybrids (Fig. 1B),
which were obviously formed a handful of generations ago. Analyses presented
hereafter were all conducted on data generated by whole-genome sequencing in this
set of 35 accessions. We first computed Patterson's D (ABBA-BABA) statistic (Durand
et al. 2011; Green et al. 2010), a statistic that quantifies gene flow after accounting for
incomplete allele sorting. This statistic is computed over the whole genome for
phylogenies of the form (((P1,P2),P3),O), where O is the outgroup *A. androsacea* and
P1 and P2 two populations of the same species. D was highest for the phylogeny with
P1 = *A. sagittata* allopatric, P2 = *A. sagittata* sympatric and P3 = all *A. nemorensis*
individuals with 0.2 (p<<0.001;block jackknife test). This significant and positive D value
shows that gene flow is higher between P3 *A. nemorensis* and P2, the sympatric
A. sagittata population, than between *A. nemorensis* and P1, the allopatric *A. sagittata*
individuals (Figure 2A). Although this test showed elevated rates of gene flow in the
sympatric population, it did not rule out introgressions in the allopatric populations.
Thus, we also compared the two allopatric *A. sagittata* populations Lob and Adl-1
(Figure 1A) with P1 = *A. sagittata* Lob, P2 = *A. sagittata* Adl-1 and P3 = all
A. nemorensis, and found *D* was 0.09 (p<<0.001; block jackknife test) indicating
significant gene flow outside of the sympatric population. Finally, we tested the
phylogeny with P1 = *A. nemorensis* sympatric, P2 = *A. nemorensis* allopatric and P3 =
A. sagittata Lob (allopatric), and found that *D* was 0.17 (p<<0.001, block jackknife test).

This positive value indicated that gene flow from *A. sagittata* into *A. nemorensis* was
stronger in allopatry than in sympatry, demonstrating that the history of hybridization in
this system predates the formation of the sympatric population.

The frequency of introgression is heterogeneous along the genome

The *D*-statistic provides information about genome-wide rates of gene flow but
not about its distribution across the genome. Thus, we calculated the f_D-statistic (Martin
et al. 2015) across the genome for five phylogenies covering all possible scenarios of
interspecific gene flow (Figure 2B). Distributions of f_D were highly zero-inflated for all
populations, indicating that introgressions were generally rare across the genome. Yet,
in all populations, we found genomic regions with elevated f_D values. We did not find
strong differences in the distribution of these regions among chromosomes. The D and
f_D statistics rely on the assumption that there was no gene flow between P3 and P1
(Green et al. 2010; Durand et al. 2011; Martin et al. 2015). As we found introgressions
in all populations, this assumption was violated, meaning any introgression shared by
P1 and P2 would remain undetected. Our estimates of gene flow are therefore likely to
be conservative.

Introgressed fragments are on average largest in the sympatric population

To locate introgressed fragments and their boundaries in the genome, we
calculated the ratio of average intra- and interspecific genetic distance for each
individual in 10 kb genomic windows (see Methods). This ratio is expected to be smaller
than the ratio throughout the genome, except in introgressed regions. We used a value
equal or greater than 2 as a conservative threshold for the ratio value calling
introgressed fragments. The average number of introgressions per individual was
highest in the sympatric *A. sagittata* population, with 24.7 introgressed fragments per
genome (standard deviation (sd) = 3.6), followed by the allopatric *A. sagittata*
populations, with an overall mean of 13.7 fragments per genome (sd = 3.5, Table S3). In
contrast, in *A. nemorensis,* we observed an average of 9.6 introgressed fragments per
genome (sd = 1.47) in the sympatric population and 2.6 (sd = 0.55) in the allopatric.
These results confirmed that interspecific gene flow from *A. nemorensis* to *A. sagittata*

was stronger than vice versa. Furthermore, the presence of introgressions in allopatric
populations suggests that gene flow is at least partly historical.

The high occurrence of introgression fragments in the sympatric *A. sagittata*
population indicates that some of these fragments could have been introduced more
recently by contemporary gene flow. We thus assessed whether the introgression was
likely to have occurred in the sympatric population (recent gene flow) or whether it
predated the separation of local populations   (historical gene flow). For this, we
determined the individual in the donor species, whose orthologous region was most
closely related to the introgressed fragment identified in the receiver species. We then
observed that the proportion of introgressed fragments that were most similar to alleles
of the sympatric *A. nemorensis* population was almost twice as high in the sympatric *A.*
sagittata population (38.5%) as in the allopatric *A. sagittata* population (20%), a
difference that was marginally significant (Chi square= 3.76, p=0.05, Figure 2C). This
trend was reversed in *A. nemorensis,* with 25% of introgressions originating from a
sympatric *A. sagittata* lineage in the sympatric *A. nemorensis* population as opposed to
43% in the allopatric *A. sagittata* population. However, as there were only four
introgressions in the sympatric population, this reversal is likely due to chance.

Figure 2: Patterns of introgression in sympatric and allopatric populations of parental

species.

A) D-statistic results calculated over the whole genome for different phylogenies. Bar color represents the
species of the P3 / donor population. Asterisks represent the result of a Jackknife test: *** p<0.001, ** p <
0.01. B) Genome-wide distribution of fD calculated in 50-kbp windows for different phylogenies, as shown
on the right. Plots are ordered by decreasing overall genome-wide fD. C) Fractions of introgression
origins for each population, inferred based on minimum genetic distances (see Methods). Numbers on top
show the total number of introgressions in each population. This plot does not take into account the
frequency of the identified introgression tracts. D) Distribution of introgression size in each population,
represented by dots and boxplots. Dots are colored according to the introgression origin. This plot does
not take into account the frequency of the identified introgression tracts. The compact letter display
indicates significant differences among groups. Abbreviations in the legends: nemo. - A. nemorensis, sag.
- A. sagittata, allo. - allopatric, symp. - sympatric

We further reasoned that recent introgressions originating from sympatric donor
lineages should be on average larger than introgressions originating from allopatric
donor lineages because the latter being more ancient, they should have had more time
to be broken down by recombination. Thus, we compared the distributions of

introgression size among populations. In *A. sagittata*, introgressions in the sympatric

population were significantly larger than in the allopatric population (Z=-4.47, p<0.001),

with median values of 37,900 bp and 10,000 bp, respectively (Figure 2D). We did not

find this difference in the *A. sagittata* allopatric populations (Kruskal-Wallis chi-squared

= 0.5174, df = 1, p = 0.472) or in the two *A. nemorensis* populations (symp.: Kruskal-

Wallis chi-squared = 0.6, df = 1, p = 0.4386; allop.: Kruskal-Wallis chi-squared = 1.772,

df = 1, p = 0.1831).

304   The best fit model excludes constant gene flow between the two taxa

Based on the above, we observed i) asymmetrical gene flow between species, ii)

evidence for gene flow both within and outside of the sympatric population. Our data

thus indicate that gene flow may have been continuous throughout the history of the two

species. However, we also observed that gene flow from *A. nemorensis* into *A. sagittata*

might have intensified in the sympatric population. Gene flow could thus have occurred

in the past and stopped during a phase of complete isolation, only to resume recently in

the sympatric population. We took a modelling ABC approach to date gene flow

between *A. nemorensis* and *A. sagittata* and to estimate its strength and past

fluctuation.

We modelled the history of interspecific gene flow using a random-forest-based

approximate Bayesian computation (ABC) approach (Raynal et al. 2019). We started

with five demographic models, which explored different modes of intra- and interspecific

gene flow (Figure 3A), and generated 50,000 coalescent simulations under each one

(see Methods). To choose the model that best fitted the data, we first trained a random-

forest classifier on the data all five models use. The accuracy of the trained classifier

was evaluated with a confusion matrix that described how many simulations were

assigned to the model under which they were generated (Figure 3B). Based on this

matrix, we identified two subsets of models that summary statistics could accurately

distinguish (max. error 5.2 %). The first subset comprised the two weak-migration

models: the model without any migration and the model with early ancestral interspecific

migration. The second subset contained the three models with strong interspecific

migration: ancestral migration, ancestral and recent sympatric migration, and

continuous migration. Within this last subset, the first model was classified with a high
accuracy of 84 %, but the second and third models were only classified with accuracies
of 53 % and 61 %, respectively. These results show that i) models with both high and
low migration rates could be distinguished easily by the classifier; and ii) determining
the exact model within each of the two groups was less straightforward. This result was
not surprising because when recent migration is low, the ancestral and recent migration
model gives results similar to those generated under the ancestral migration model,
which does not allow migration in the sympatric population.

Next, we used the trained random-forest classifier to determine the model that
best explained the observed data. To this end, we randomly selected 100 datasets,
each consisting of 500 loci of individual length 75 kbp sampled from the observed
genotype (full genome) data and summarized with 200 summary statistics (Table S4).
Models with non-constant gene flow always received the majority of the votes. The
model with ancestral & recent sympatric migration was chosen as the best model for 78
% of the observed genomic subsamples (Figure 3C). This was followed by the model
assuming ancestral migration only for 22 % of observed genomic subsamples. The
other three models were not selected as the most likely model.

Ancestral and recent sympatric migration separated by a period of isolation most
aptly characterize the history of this plant system (Figure 3A). A history of low but
uninterrupted gene flow (Model 5) can be clearly ruled out. We further quantified the
goodness-of-fit of the model we identified as the best. The distance between observed
and simulated datasets was not significantly different, confirming that the observed data
lay well within expectations of the model (Figure 3D). Next, we used this model to
estimate the timing, strength and direction of gene flow between species and
populations as well as its fluctuation across genome sub-samples.

Figure 3: Choosing the best demographic model.

A) Schematic representation of the tested demographic models. Populations are colored according to the bottom-right code. Arrows represent migration rates. Interspecific migration rates can be

 asymmetric (even with double-sided arrows) and intraspecific migration is symmetric. Dashed lines are
timepoints at which populations split and/or migration rates change. B) Confusion matrix of the model-
choice random forest model. For each model, simulated (out-of-bag) samples were classified by the
trained random forest. Correct classifications are on the diagonal. Results are represented as
percentages. C) Mean model classification results, i.e. random forest votes, for 100 observed datasets,
with 500 randomly selected genomic windows each. Error bars represent the standard deviation. Bars are
colored according to the percentage of observed datasets for which the respective model was selected as
the best fit. D) Evaluation of fit for the best model. For each of the 100 observed datasets, the median
normalized summary statistic distance between observed data and the closest 1% of simulated datasets
was calculated (orange). As a null distribution, the same calculation was done with 1000 simulated
datasets of pseudo-observed data. For better clarity, the x-axis was trimmed at 300, but no observed
distance was larger than that.

Estimation of demographic parameters indicates very low Ne in both taxa

To estimate demographic parameters of the best model, we generated a total of
360,000 coalescent simulations under this demographic model. We used two methods
for parameter estimation -- random-forest-based ABC (Marin et al. 2019), which also
generated confidence intervals, and extreme gradient boosting (XGBoost; Chen et al.
2020), which provided the most accurate point estimates -- and estimated each
parameter independently (see Methods for details). We achieved highest accuracy for
the estimation of present population sizes and lowest accuracy for the time and
population size at speciation (Figure S6-7). Of the two methods, XGBoost tended to
have the lowest RMSE and the highest R2, indicating that it could best reconstruct the
parameters under which simulated data was generated (Figure S8).

We then estimated the demographic parameters of the two species based on
observed data. We present in detail the estimates obtained for one of the observed
datasets we randomly picked among those assigned to the selected model. The
estimated contemporary population sizes for *A. nemorensis* were 6,775 individuals (sd =
1,988) for the allopatric and 2,964 (sd = 615) for the sympatric population (Figure 4,
Table S5). For *A. sagittata*, 14,895 individuals (sd = 2,407) made up the effective
allopatric population and 6,425 (sd = 1,649) individuals, the effective sympatric
population. These results agree with those of the previous report, namely, that in the

sympatric population, *A. sagittata* harbored levels of genetic diversity higher than those
of *A. nemorensis* (Dittberner et al. 2019). That the estimated sizes of the allopatric
populations was larger in both species could be because the allopatric samples
included individuals collected in several sites. We further estimated that the ancestral
population in *A. sagittata* was more than twice as large as that in *A. nemorensis*, with
59,031 (sd = 12,529) and 25,206 (sd = 9,777), respectively. The magnitude of the
population size declined after the split in both species, with a mean factor of 6.11 for
A. nemorensis, which was similar to the factor of 6.57 estimated for *A. sagittata*. A
decline of effective population sizes in the recent past has also been observed in the
closely related species *Arabidopsis thaliana,* which is also selfing (Durvasula et al.
2017). However, contemporary population sizes in this species are still larger by at least
an order of magnitude. This difference indicates that *A. nemorensis* and *A. sagittata*
have not been particularly successful in their natural landscape.
401
The period of isolation dates back to the last glaciation
The split between the species occurred approximately 894,801 (sd = 119,548)
generations ago, when the effective size of the ancestral population was approximately
150,281 individuals (sd = 39,960). Standard deviations were high, presumably because
our model also allowed for migration between populations (Figure S7). Within species,
the populations split almost simultaneously: 10,441 (sd = 3,910) generations ago in
A. nemorensis and 10,858 (sd = 3,417) generations ago in *A. sagittata*, and their split
time coincides with the last glacial maximum. Our analysis thus suggests that the
genetic landscape of the two taxa was established after the last glaciation, a period
during which they became completely isolated.
412
Estimates of migration rate confirmed gene flow resumed in sympatry
The low but significant estimates of gene flow showed that the model with
ancestral and recent migration explained the data better than a model assuming only
ancestral migration (Figure 3). We report the migration rates as the log10-transformed
fraction of a population migrating per generation. Migration from *A. nemorensis* to
A. sagittata in the ancestral population was approximately five times higher than vice

versa: -6.34 (sd = 0.36) and -7.00 (sd = 0.85), respectively. This amounts to about one
migrant every 33 generations. Point estimates indicated that, in the sympatric
population, migration rates were lower than in the ancestral population. Yet, the rate
was still approximately ten times higher from *A. nemorensis* to *A. sagittata* than vice
versa, with a rate of -7.72 (sd = 1.39) and -8.74 (sd = 1.33). The mode of the posterior
distribution for these two parameters differed even more strongly (-6.36 for migration
from *A. nemorensis* to *A. sagittata* and -9.80 for the opposite direction). This result
aligns well with the asymmetry of gene flow revealed by the analysis of chloroplast
genome variation. As expected estimates of intraspecific migration rate were higher
than interspecific migration rates. We noted that the estimate was two orders of
magnitude higher in *A. nemorensis* compared to *A. sagittata* (-5.61, sd = 1.61 and -7.09,
sd = 1.60 in *A. nemorensis* and *A. sagittata*, respectively).

449   *Figure 4: Model parameter estimation*

Variation in the observed genomic sample affects ancestral population size and sympatric migration rates

Since we observed that the introgression rate varies across the genome, we
examined whether estimates of gene flow varied depending on the genomic regions
included in the observed sample. To test this, we compared the distributions of random-
forest-based and xgboost-based point estimates of all demographic parameters for the
100 observed datasets (Table S6). Estimates were fairly robust to genome subsamples
because they were always in the same order of magnitude (Figure 5).

Yet, we made two notable observations. First, the abcrf estimate of sympatric
migration from *A. nemorensis* to *A. sagittata* was 4.6 times more variable than the
estimate of migration in the opposite direction (Figure 5). The distribution of point
estimates for recent sympatric migration from *A. nemorensis* to *A. sagittata* ranged from
-8.3 to -6.5, with a mean of -7.5 (Figure S9). Since the estimate was positively
correlated with the proportion of introgression regions included in each genomic sub-
sample, it partly reflects the heterogeneity of introgression rates along the genome
(Spearman correlation coefficient Rho=0.2, p= 0.03). However, migration rates
estimated by gradient boosting, a machine learning method that does not take the
variance of the coalescent process into account, appeared more robust to genomic
heterogeneity in introgression rate. Second, while both methods indicated the same
asymmetry in gene flow between *A. nemorensis* and *A. sagittata* in the sympatric
population, they yielded opposite conclusions for ancestral gene flow. Drawing a firm
conclusion on the direction of ancestral gene flow is therefore not possible.

Figure 5: Distribution of parameter estimates for the 100 observed datasets formed by sub-sampling 500 75kb-long regions in the genome. M_N and M_S: intraspecific migration rate among A. nemorensis *and* A. sagittata *populations, respectively. M_{NS} and M_{SN}: interspecific migration rate from* A. nemorensis *into* A. sagittata*, respectively. Sym: sympatric population, Anc. : ancient gene flow. N_{N1} : Effective population size of allopatric* A. nemorensis *populations, N_{N2} : Effective population of sympatric* A. nemorensis *population, N_{S1}: Effective population of sympatric* A. sagittata *population, N_{S2}– Effective population size of allopatric* A. sagittata *population, N_{Nanc}: Ancestral population size of* A. nemorensis N_{Nanc}, N_{Sanc} *: Ancestral population size of* A. sagittata*, T_N : time elapsed since split of* A. nemorensis *population, T_S : Time elapsed since split of* A. sagittata *populations, T_{Anc}: time of speciation, N_{Anc} : ancestral population size prior to speciation.*

DISCUSSION

ABC modelling untangles contemporary and ancestral gene flow between two taxa

Explorations of both genome-wide and local estimates of interspecific introgressions, as well as allelic distributions within and between species at introgressed loci, indicated that these two Arabis species share a history of gene flow that predates the formation of the sympatric population. The potential of ABC approaches had been revealed previously by the study of population isolation in *Salmo salar* (Rougemont and Bernatchez 2018). Patterns of gene flow in *S. salar* populations, however, were

resolved by genotyping more than 2,000 individuals at some 5,000 SNPs (Rougemont
and Bernatchez 2018). This study now shows that ABC approaches have the power to
resolve complex demographic histories even with a relatively small sample of
genotypes. Indeed, we provide compelling evidence that, although low levels of gene
flow were maintained long after speciation, interspecific gene flow was not continuous in
the history of these two plant species. Models postulating a phase of complete isolation
initiated approximately 10,000 generations ago, were markedly better at explaining the
data than those assuming constant gene flow.

We can further confirm that the Rhine population forms a contemporary hotspot
of hybridization. The migration rates estimated for past gene flow were low in absolute
number but they clearly supported a model in which gene flow has resumed specifically
in the sympatric population. This inference aligns well with the observation that, in the
sympatric population of *A. sagittata*, introgressed fragments are longer than in the
allopatric population. We note that the estimate of gene flow in the sympatric population
should be considered as the lower bound since the sample used for ABC demographic
inference excluded individuals with obvious signatures of admixture. The low but
significant rate of gene flow estimated between the two species in the sympatric
populations suggests that gene flow might have happened in the past as it happens
today: through the occasional formation of sympatric demes leading to localized bursts
of interspecific gene flow.

Interspecific gene flow as a chance for evolutionary rescue?

A. nemorensis and *A. sagittata* are both considered endangered in Germany and
in many parts of Europe (Dittberner et al. 2019). The demographic model confirms that
effective population sizes are particularly small in these two low-diversity taxa (Figure
S2-S5). Genetic diversity in both species was approximately an order of magnitude
lower than in more common selfing relatives such as *Arabidopsis thaliana* (Alonso-
Blanco et al. 2016) or *Capsella rubella* (Onge et al. 2011). These low levels of diversity,
however, do not necessarily point to a low adaptive potential that would increase their
risk of extinction. Indeed, the hotspot of contemporary gene flow we uncovered has
formed after a period of isolation that might have been sufficiently long to allow

functional divergence. Variance in abcrf estimates associated with the sub-sampling of genomic variation indicates that the signature of contemporary gene flow is not homogeneous throughout the genome, in contrast to that of ancient gene flow. Such a heterogeneity suggests that the contemporary contact is not without fitness consequences, and, indeed, we found several high-frequency introgressions that could be a signature of adaptive gene flow (Racimo et al. 2017). Studies in organisms such as yeast or sunflower have shown that hybrid populations can adapt efficiently to changing environmental conditions (Todesco et al. 2020; Mitchell et al. 2019; Stelkens et al. 2014). Moreover, gene flow can also resolve fitness tradeoffs limiting adaptation in the parental species (Walter et al. 2020).

At this stage, however, it is premature to conclude that introgressions will positively affect the receiving taxon (Suarez-Gonzalez et al. 2018; Shumer et al. 2018). First, we do not know whether the two species possess alleles of potential fitness relevance for each other. Second, F1 hybrids display low fertility, indicating that gene flow causes fertility reduction, at least initially (HD pers. com.; Titz 1979). The simple removal of incompatibility alleles could be sufficient to explain the pattern of heterogeneous introgression documented here (Schumer et al. 2014). Introgressions may also carry deleterious alleles, either due to fitness trade-offs between the parental species (Arnegard et al. 2014) or due to slightly deleterious variants in the parental genomes that have not been purged by selection, i.e., genetic load. This phenomenon is known to have shaped the landscape of introgression from Neanderthals to humans (Harris and Nielsen 2016; Juric et al. 2016), resulting in more introgressions outside of either functionally important regions or of regions with low recombination rates (Schumer, Xu, et al. 2018). Novel methods for the detection of adaptive gene flow have been proposed, but this task remains arduous in selfing species, where recombination events are rare (Setter et al. 2020). Further work is required to identify and disentangle the positive and negative fitness effects of gene flow between *A. sagittata* and *A. nemorensis,* Although Having identified a local hotspot of contemporary hybridization, it is now possible to monitor these consequences in situ.

MATERIAL AND METHODS

Plant material and DNA extraction

In 2016 and 2017, we identified 30 sites reported to host *A. nemorensis*
populations on the Deutschland Flora Database. Of these, 24 could be visited, and *A.*
nemorensis/A. sagittata populations were observed in 11 of them. We sampled seeds
from at least 10 *A. nemorensis* and/or *A. sagittata* plants per site for a total of 231
accessions (Table S1). Populations were located in Southern Germany and Austria
(Figure 1A). One of these sites, referred to as "Rhine," was previously described in
(Dittberner et al. 2019) and consists of multiple proximate pristine habitat patches at one
site. Individuals from these patches were assembled in one sympatric population. We
extracted DNA for genotyping as previously described (Dittberner et al. 2019).

RAD-seq genotyping for phylogeography

In addition to the 140 accessions originating from the Rhine populations that
were previously genotyped in Dittberner et al. (2019), we genotyped 91 accessions
using the original RAD-seq protocol (Etter et al. 2011) with the modification described in
(Dittberner et al. 2019). Libraries were sequenced at the Cologne Center for Genomics
on three Illumina HiSeq 4000 lanes with 2 × 150bp. We used *FastQC* (Andrews 2010)
to check the raw reads. We trimmed adapters and removed reads shorter than 100bp
using *Cutadapt* (Martin 2011). We removed PCR duplicates based on a 5 bp stretch of
random nucleotides at the end of the adapter, using the *clone_filter* module of *Stacks*
version 1.37 (Catchen et al. 2013). We de-multiplexed samples using the
process_radtags module from *Stacks*. We filtered reads with ambiguous barcodes
(allowed distance 2) and cut-sites, reads with uncalled bases and low-quality reads
(default threshold).

We used our previously described pipeline to call nuclear genotypes in all
samples (Dittberner et al. 2019). Briefly, we used BWA (Li and Durbin 2009) to map
reads against the high-quality reference genome we had sequenced and assembled for
A. nemorensis (Dittberner et al. 2019). We filtered mapped reads using *SAMtools* (Li et
al. 2009) to remove regions with excessively low (30%) or high (2-fold) coverage,

compared to the mean coverage. We called genotypes using samtools mpileup and
VarScan2 (Koboldt et al. 2012). We filtered genotyped loci using VCFtools (Danecek et
al. 2011). Knowing that both species are predominantly selfing, we filtered out SNPs
with more than 20% overall heterozygosity and SNPs with more than 75%
heterozygosity within a population, because heterozygous SNPs tended to cluster on a
few single RAD-seq fragments, indicating inaccuracies in mapping that were not picked
up by filtering on coverage. In total, we genotyped 2.6 million sites (~ 1% of the
genome) (excluding sites with more than 5% missing data in our sample) and identified
25,634 single nucleotide polymorphisms (SNPs).

To obtain SNPs in the chloroplast sequence, we mapped the RAD-seq reads
against a chloroplast reference genome of *Arabis hirsuta* (Kawabe et al. 2018), using
the *mem* algorithm of BWA (Li and Durbin 2009) with default settings. We called
genotypes and SNPs as described above. We set heterozygous genotype calls to
missing data, as these calls most likely resulted from mapping errors (the plastome is
effectively haploid). Furthermore, we used VCFtools to remove SNPs with either more
than 20% missing data or more than two alleles, which resulted in 23 chloroplast SNPs.

Determining hybrids with RAD-seq and chloroplast data

To determine the genomic make-up of each population and identify the presence
of hybrids, we first analyzed SNP data using ADMIXTURE (Durand et al. 2011), varying
K from 2 to 10. First we converted VCF files to bed format using PLINK (Purcell et al.
2007; Purcell 2009). We ran ADMIXTURE analysis (Alexander and Lange 2011a) for
K=1 to K=6, with 10 iterations of cross-validation each. We normalized clusters across
runs using CLUMPAK (Kopelman et al. 2015). We used K=2 for further analysis, as we
were analyzing two species and the value was well supported by the cross-validation
error (Figure S1). Individuals were defined as hybrids if they had less than 95%
ancestry from either species. We created plots using the libraries *ggplot2* (Wickham
2009), *ggmap* (Kahle and Wickham 2013), *scatterpie* (Yu 2018) and *ggsn* (Baquero
2017). Finally, we used the library *pegas* (Paradis et al. 2016b), to determine
chloroplast haplotypes and build a haplotype network.

Analysis of genetic diversity

We used the RAD-seq data to determine the level of genetic diversity within and between species. Using the vcfR package (Knaus and Grünwald 2017), the genotype data were loaded into R and converted to DNAbin format. We used the *pegas* package (Paradis et al. 2016b) to calculate pairwise genetic distances among all individuals. Based on the resulting distance matrix, we calculated average genetic distances within and between both populations and species. We calculated the diversity in the hybrid complex as the average genetic distance within the whole Rhine population.

Whole genome resequencing for gene flow quantification and ABC

To achieve greater resolution for introgression detection, we randomly selected five *A. nemorensis* and 14 *A. sagittata* accessions from the sympatric population, 4 and 6 accessions from two allopatric *A. sagittata* populations and one accession of each of the six allopatric *A. nemorensis* populations. In total, the whole genome of 35 accessions was sequenced (Table S2). To provide an outgroup species for estimating gene flow, we also sequenced one accession of *Arabis androsacea*, provided by Jean-Gabriel Valay (Jardin Alpin du Lautaret, France). DNA for these accessions was extracted as described above. Libraries were prepared using Illumina TruSeq DNA PCR-free kits at the Cologne Center for Genomics. Six samples were sequenced on a HiSeq 4000 with 50 million 2x75 bp reads per sample. The remaining samples were sequenced on a NovaSeq6000 with 25 million 2x150 bp reads per sample, resulting in a depth of ~ 25x.

We used *FastQC* (Andrews 2010) to quality-check the resulting reads. We filtered the reads using the *process_shortreads* module from *Stacks* v2.2 (Catchen et al. 2013), removing reads shorter than 70 bp or 100 bp (threshold was adapted to the sequencing device) and reads with uncalled bases or low-quality scores (default threshold). At this point, we included reads of the *A. nemorensis* reference genome accession from the sympatric population (Dittberner et al. 2019). We mapped the reads against the *A. nemorensis* reference genome (Dittberner et al. 2019) using the *mem* algorithm of BWA (Li and Durbin 2009) with default settings. We filtered out poorly mapped reads using the following criteria: mapping quality < 30, read not mapped in

proper pairs, > 50% of the read is soft-clipped. We called genotypes as described above
but with the default strand-filter of *VarScan2* (Koboldt et al. 2012). Genotypes were
filtered using *VCFtools* (Danecek et al. 2011). As in the above, filters were set to allow
maximum 20% missing data, maximum two alleles and not more than 20%
heterozygosity per site (see above) and to remove indels and low quality genotype calls
marked "filtered" by *VarScan2*. Additionally, we removed sites with a mean depth
greater than 45 (twice the mean depth of all sites) and smaller than 7.5 (mean depth of
all sites divided by 3).   From the resulting genotype dataset, which comprised
78.976.767 nucleotide positions, we extracted a total of 2.954.526 SNPs using
VCFtools. To verify sample identity, we performed a PCA using the *adegenet* package
(Jombart et al. 2016). Based on this PCA and the RAD-seq data above, we re-assigned
one *A. sagittata* mis-labeled sample (174) from one allopatric population to the other.

Estimation of the population recombination rate

In order to take within-locus recombination into account in our ABC estimations
of the demographic history, we estimated the population recombination rate R; we used
the sympatric *A. sagittata* population because we had sequenced the most individuals in
this population, which should maximize the accuracy of the estimate. To this end, we
first phased the reads using the read-aware phasing algorithm implemented in
SHAPEIT (Delaneau et al. 2013). We then used FastEPRR (Gao et al. 2016) to
estimate the local recombination rate in non-overlapping 75 kbp windows along the
genome, with otherwise default settings. We used the median of the distribution
(R=11.5) as the per locus population recombination rate (i.e. R=0.15 per kb).

Detection of interspecific gene flow

To investigate signatures of gene flow among non-hybrid individuals, we
analyzed the whole-genome resequencing (WGS) data using two four-taxon statistics:
we calculated D (Durand et al. 2011) over the whole genome to detect interspecific
gene flow, and we calculated f_D (Martin et al. 2015) in 50kbp sliding windows to analyze
the heterogeneity of gene flow along the chromosomes. These tests assumed the
general phylogeny: (P1,P2)(P3) outgroup; where P1 was the control population (no

gene flow), P2 was the gene-flow target population and P3 was the donor population of
gene flow. We used *A. androsacea* as the outgroup to determine the ancestral allelic
state. We calculated f_D in 50 kbp sliding windows with 25 kbp overlap, following Martin
et al. (2015). We skipped sites with derived allele frequency of 0 in P3, as these could
bias values of f_D. We calculated these statistics for the phylogenies shown in Figure 2.

To complement the four-taxon analysis, we applied a method based on analyses
of genetic distance to locate and assign boundaries to putative introgressed fragments
in each accession. Our reasoning was as follows: if there is no introgression, the mean
genetic distance of a given accession to members of the same species should be
smaller than the distance to members of other species; for introgression, the opposite is
true. Thus, in 10 kbp windows along the genome, we calculated the mean intra- and
interspecific genetic distance for each accession, excluding individuals from the same
population for the intraspecific case, as they are likely to carry the same introgression.
We added a constant of 0.0001 to all interspecific distances to avoid zero-division and
removed windows with zero interspecific distance and non-extreme intraspecific
distance (< 90^{th} percentile), as these could lead to spurious signals of introgression. For
each window, we calculated the ratio of intraspecific and interspecific genetic distance
(hereafter, distance ratio), which we used to identify introgressions.

For each accession, we identified introgressions by first removing all windows
with a distance ratio smaller than or equal to 1. We then classified windows as
introgressions if their distance ratio was > 2 or if both of their neighboring windows
fulfilled this condition. Adjacent introgression windows were connected to a single
introgression. We applied several filters to the resulting set of candidate introgressions:
We excluded candidate introgressions from further analysis if they contained too many
repetitive elements (i.e those with mean number of genotyped positions less than the
25% quantile of the number of genotyped position in all introgressions). Furthermore,
we overlapped candidate introgressions with D and f_D values (in 50 kbp windows) for
the respective target population; for introgressions larger than 50 kbp, we calculated the
mean of all windows within the introgression. We kept introgressions with values of D >
0.1 and f_D > 0.01. Subsequently, we removed introgressions with > 30% mean
heterozygosity, as these often gave inaccurate signals: the expectation for this distance

ratio is not the same as for the homozygous case (detecting heterozygous
introgressions with this method would require haplotype data). To define the boundaries
of each introgression accurately within each of the remaining candidate introgressions,
we recalculated the genetic distances in sliding 2 kbp windows in 200 bp steps. We
applied the same thresholds as described above to define our final set of introgression
regions in all these windows.

Finally, we aimed to determine the likely origin of each introgression by
calculating genetic distances of the introgression region between all accessions of the
donor species and the target accession. We then determined the accession(s) with the
minimum genetic distance to a given introgression. We standardized genetic distances
by dividing them with the approximate number of genotyped sites for each pair of
accessions (we used an approximation to reduce computational requirements). The
approximate number was estimated by taking the mean of all 2 kbp windows (200 bp
steps) within an introgression and scaling this value by the length of the introgression. If
all accessions originated from the same population (i.e. sympatric or allopatric for the
donor species), we inferred that the introgression likely originated from this population
(or a closely related one). If donor accessions originated from multiple populations, the
inferred origin was ambiguous.

Statistical analysis was performed in R (R Development Core Team 2008).
Visualization was done using the ggplot2 library (Wickham 2009). To account for the
introgression frequency in statistical analyses of origin and introgression size, we
counted each unique introgression (identical start- and end-points) only once,
regardless of its frequency. We tested whether introgression size differed among
populations using a Kruskal-Wallis test followed by a pairwise Dunn-Test, implemented
in the *FSA* package (Ogle et al. 2020).

Modelling the history of interspecific gene flow

We used coalescent simulations and an ABC algorithm with random forest-
based Bayesian parameter inference to determine the most likely demographic history
of the two species, focusing especially on the history of interspecific gene flow (Raynal
et al. 2019). Modelling was performed in two steps: i) we compared data obtained from

each postulated demographic model to our observed genetic data to identify the best-
fitting one and ii) we estimated the posterior distribution for each demographic
parameter of the best model.

The summary statistics for the observed data were computed from the 35 fully
sequenced genomes. We randomly selected 500 genomic windows of 75 kbp each, all
of which were at least 150 kbp apart from one another and contained at least 20,000
genotyped sites, which allowed excluding regions that were excessively repetitive. We
computed the following summary statistics using a custom python script: overall θ_W, θ_W
and θ_π within each population, Tajima's D for each population, d_{XY} and F_{ST} between all
pairs of populations, the fraction of shared, private, invariant and fixed SNPs for all pairs
of populations, Patterson's D and f_D (ABBA-BABA) for 8 possible phylogenies. All
summary statistics, except Patterson's D, were calculated per locus and the resulting
distributions were further summarized as the mean and variance as well as the 5% and
95% quantile, resulting in a total of 236 summary statistics. To estimate the variance in
model selection and parameter inference due to window choice, we assembled 100
random sets of these 500 genomic windows and considered each as one independent
observation.

Simulated data for each of the five model described in Figure 3A were obtained
with the software *escrm* (Sellinger et al. 2020), which is a modification of the widely
used coalescent simulation software scrm (Staab et al. 2015) that accommodates
selfing (we set a 90% selfing rate for all populations). Fixed parameter values and prior
distributions for all demographic parameters are found in Table 1. For each model, we
simulated 50,000 datasets, each consisting of five hundred 75 kbp-long loci. Summary
statistics for the observed data are computed using the same custom python script.

Model choice procedure

In the first step, we specified five different demographic models, all of which
differed in their mode of intra- and interspecific gene flow (Figure 3A). In the first model,
we assumed no migration at all, either intra- or interspecific. In the second model,
interspecific migration stopped 100,000 generations after the species split. In the third

model, ancestral interspecific migration continued until the first intraspecific population-

split. The fourth model was an extension of the third model that additionally allowed

migration between the sympatric populations, after populations in both species had split.

In the fifth model, interspecific migration continued throughout the history of the species,

allowing a change in intensity after the intraspecific population had split in both species.

All interspecific migration rates were allowed to be asymmetrical, while symmetric

intraspecific migration was assumed in all models (except model # 1, which assumed no

migration at all). Population sizes were constant for each population but were allowed to

change at all population split points (Table 1).

Table 1: Overview of demographic parameters (see Figure 3A) and their prior

distributions or fixed values. M_{Sym} and M_{Anc} had the same prior distributions for both

directions of migration. 500 loci of length 75bp were simulated assuming a per site

mutation rate of 7.10^{-9} (Ossowski et al. 2010) and a population recombination rate of

0.15 per kb.

Parameter	Minimum	Maximum	Distribution
N_{N1}, N_{N2}, N_{S1}, N_{S4}	500	100,000	log-uniform
N_{Nanc}, N_{Sanc}	500	100,000	Uniform
N_{Anc}	50,000	200,000	Uniform
T_N, T_S	0	20,000	Uniform
T_{Anc}	500,000	1,500,000	Uniform
M_{Sym}, M_{Anc}	1e-11	1e-3	log-uniform
M_N, M_S	1e-11	1e-1	log-uniform

We used *abcrf,* an R package designed for ABC-based demographic modelling

based on random forests, to choose the model that best fit our data and to infer their

parameters (Marin et al. 2019; Raynal et al. 2019). We trained a random forest classifier

to distinguish between the five models based on a set of summary statistics, i.e., the
model was the dependent variable and the 236 summary statistics were independent
variables. We trained the random forest algorithm on 50,000 simulated datasets for
each of the five models drawing from the priors in Table 1 and using *abcrf* default
parameters, except that parameter "lda" was set to false, disabling linear discriminant
analysis. To estimate the classification error for model choice, we used the so-called
out-of-bag method implemented in *abcrf*. Due to bootstrap aggregating, each sample
was used only to build a subset of trees in the random forest, and the remaining trees
could be used to predict the best model for this sample. The results were represented
as a confusion matrix showing how frequently the predicted model matches (or does not
match) the true model (Figure 3 B). All samples were expected to fall on the diagonal of
such a matrix. We then used the trained classifier to assign the most likely demographic
model to each of the 100 observed samples (see above). The model that was most
frequently selected as the best model for the observed samples was selected as the
model fitting our data best.

We further quantified the goodness-of-fit for the best model using a method
implemented in the R package *abc* (Katalin et al. 2015), which is based on a rejection
algorithm. Briefly, the normalized distance of summary statistics between the observed
and all 50,000 simulated datasets was calculated; then the median of the lowest 1% of
distances was extracted (the rejection step). The same procedure was performed for
1,000 simulated datasets under this model, drawing parameter values from the prior
distributions of distances (so-called pseudo-observed distances). If the model fit is
good, the observed distance (1% best-fitting simulations to the observed data) should
lie well within the distribution of pseudo-observed distances (1% best simulations fit
each of the pseudo-observed dataset). We conducted this analysis for all 100 observed
datasets. For each dataset, a p-value was calculated as the fraction of pseudo-
observed distances, which are higher than the observed distance.

Estimation of demographic parameters

After identifying the best-fitting demographic model, we estimated the posterior
distributions of the different parameters. Demographic parameters with log-uniform

prior distributions were log-transformed prior to training to make them uniform. To
reduce computational load, we first selected a subset of the most informative summary
statistics for each parameter; we then trained a random forest for each demographic
parameter based on 115,000 simulated datasets and extracted the top ten most
important summary statistics. This approach reduced the 236 summary statistics to a
set of 111 unique summary statistics for training the full models.

The accuracy of random-forest predictions can be greatly increased by tuning
training parameters. We tuned the following training parameters by training random
forests using all possible combinations: the number of variables to possibly split in each
node (mtry; search space: 11, 22, 44, 66, 88); and the minimal node size (search space:
5, 10, 30, 50). This training was conducted for a subset of representative demographic
parameters (N_{N1}, N_{Nanc}, T_N, $M_{SN,anc}$, M_S, $M_{SN,sym}$). We plotted the out-of-bag prediction
error for each combination of training and demographic parameters (Figure S10). Based
on these results, we took mtry=44 as the optimal number of randomly sampled
variables, as higher values strongly increased training time while increasing accuracy
only minimally. The influence of minimal node size was generally small, so we kept it at
the default value of 5.

Finally, we trained a random forest for each parameter on 359,000 simulated
datasets, keeping 1,000 pseudo-observed datasets out of the training dataset to use as
a test. We quantified the estimation error for each model parameter by computing the
parameter estimates for each pseudo-observed dataset and calculating R² between true
and estimated values. Additionally, we computed the root-mean-square error (RMSE)

per model parameter as: $\sqrt{\frac{\sum_{t=1}^{T}(\hat{y}_t - y_t)^2}{T}}$, where T is the number of pseudo-observed
datasets, y_t is the true value and $\hat{y}_t$ is the estimated value. To make RMSE more
comparable among parameters, we further converted RMSE to percentages (%RMSE)
by scaling it to the prior range of each parameter (Figure S8).

To calculate parameter estimates for the observed data, we first randomly
selected one of the 100 genomic sub-samples (observed samples) that was previously
classified as fitting the best model For this dataset, point estimate, standard deviation,
posterior mode and the 5% and 95% quantiles of the posterior distributions were

determined for each demographic parameter. Additionally, we determined point
estimates for the remaining 99 observed datasets to assess the variation of
demographic parameters depending on the chosen genomic sub-sample.

To complement *abcrf,* which estimates a posterior distribution, we also used
XGBoost (Chen et al. 2020), a widely used tree-based machine-learning method that
gives only a point estimate but often achieves greater accuracy than random forests.
We included all summary statistics in XGBoost models. We split the 50 000 simulated
datasets into a test dataset of 20,000 samples and a training dataset with the remaining
30,000 samples. Tree depth (dp; values: 3, 6, 9) and minimum child weight (values: 5,
10), are the two training parameters that control the complexity of the trees. We tuned
these parameters for the same demographic parameters as described above,
chooseing those that minimized RMSE (Figure S11). The algorithm was stopped if the
test statistic did not improve for 100 rounds. For tree depth, we chose six as the optimal
value for training models for T_S, T_N, N_{Nanc} and N_{Sanc}, and nine for all other model
parameters. We chose 10 as the optimal value for minimum child weight. Using these
settings, we trained an XGBoost model for each demographic parameter, again
stopping after 100 rounds if there was no improvement. We assessed accuracy using
the test dataset with 20,000 simulated observations, as described for *abcrf*. We then
performed the estimation of the demographic parameters for the 100 genomic sub-
samples forming the observed datasets.

888
889

ACKNOWLEDGMENTs

This research was funded by the European Research Council (ERC) through the
"AdaptoSCOPE" grant 648617, and by the Deutsche Forschungsgemeinschaft through
DFG Grant ME2742/13-1. HD acknowledges funding by the International Max Planck
Research School. We thank Dr. Markus Stetter and Dr. Gregor Schmitz and Prof. N.
Hölzel for helpful discussions. Sequence data are available at ENA under ERS5040446-
ERS5040483. Markdown scripts are available as supplementary information.

897
898

899 **SUPPLEMENTARY MATERIAL**

900
Suppl_Fig.pdf: Supplementary figures S1-S11 with legends.
RADseq.html: R Markdown of RAD-seq genotyping script and analyses.

Introg_Det_final.html: R Markdown of Introgression detection in whole genome data
Demography_estimation_final2.html: R markdown for ABC modeling
Suppl_Tables: Table S1 Description of genotypes included in the RAD-seq analysis,
Table S2: Description of genotypes chosen for whole-genome sequencing –Table S3:
list of introgressed fragments identified in the genomes of individuals described in Table
S2. Table S4: Summary statistics describing the 100 observed datasets. Table S5:
ABCRF parameter distribution and XGBoost point estimate for the representative
observed dataset shown in Fig. 4. Table S6: Variation in parameter point estimates for
the 100 observed datasets.
912

REFERENCES

Abbott RJ. 2017. Plant speciation across environmental gradients and the occurrence

and nature of hybrid zones. *J. Syst. Evol.* 55:238–258.

Ackermann RR, Arnold ML, Baiz MD, Cahill JA, Cortés-Ortiz L, Evans BJ, Grant BR, Grant

PR, Hallgrimsson B, Humphreys RA, et al. 2019. Hybridization in human evolution: Insights from

other organisms. *Evol. Anthropol. Issues News Rev.* 28:189–209.

Alexander DH, Lange K. 2011a. Enhancements to the ADMIXTURE algorithm for

individual ancestry estimation. *BMC Bioinformatics* 12:246.

Alexander DH, Lange K. 2011b. Enhancements to the ADMIXTURE algorithm for

individual ancestry estimation. *BMC Bioinformatics* 12:246.

Alonso-Blanco C, Andrade J, Becker C, Bemm F, Bergelson J, Borgwardt KM, Cao J, Chae

E, Dezwaan TM, Ding W, et al. 2016. 1,135 Genomes Reveal the Global Pattern of

Polymorphism in Arabidopsis thaliana. *Cell* 166:481–491.

Anderson JT, Panetta AM, Mitchell-Olds T. 2012. Evolutionary and Ecological Responses

to Anthropogenic Climate Change: Update on Anthropogenic Climate Change. *Plant Physiol.*

160:1728–1740.

Andrews S. 2010. FastQC: a quality control tool for high throughput sequence data.

Available from: http://www.bioinformatics.babraham.ac.uk/projects/fastqc

Arnegard ME, McGee MD, Matthews B, Marchinko KB, Conte GL, Kabir S, Bedford N,

Bergek S, Chan YF, Jones FC, et al. 2014. Genetics of ecological divergence during speciation.

Nature 511:307–311.

Arnold BJ, Lahner B, DaCosta JM, Weisman CM, Hollister JD, Salt DE, Bomblies K, Yant L.

2016. Borrowed alleles and convergence in serpentine adaptation. *Proc. Natl. Acad. Sci. U. S. A.*

113:8320–8325.

Baduel P, Hunter B, Yeola S, Bomblies K. 2018. Genetic basis and evolution of rapid

cycling in railway populations of tetraploid Arabidopsis arenosa. *PLOS Genet.* 14:e1007510.

Baquero OS. 2017. ggsn: North Symbols and Scale Bars for Maps Created with "ggplot2"

or "ggmap." Available from: https://CRAN.R-project.org/package=ggsn

Bierne N, Gagnaire P-A, David P. 2013. The geography of introgression in a patchy

environment and the thorn in the side of ecological speciation. *Curr. Zool.* 59:72–86.

Brandvain Y, Kenney AM, Flagel L, Coop G, Sweigart AL. 2014. Speciation and

Introgression between Mimulus nasutus and Mimulus guttatus. *PLOS Genet.* 10:e1004410.

Burmeier S, Eckstein RL, Donath TW, Otte A. 2010. Plant Pattern Development during

Early Post-Restoration Succession in Grasslands-A Case Study of Arabis nemorensis. *Restor.*

Ecol. 19:648–659.

Catchen J, Hohenlohe PA, Bassham S, Amores A, Cresko WA. 2013. Stacks: an analysis

tool set for population genomics. *Mol. Ecol.* 22:3124–3140.

Chen T, He T, Benesty M, Khotilovich V, Tang Y, Cho H, Chen K, Mitchell R, Cano I, Zhou

T, et al. 2020. xgboost: Extreme Gradient Boosting. Available from: https://CRAN.R-

project.org/package=xgboost

Chunco AJ. 2014. Hybridization in a warmer world. *Ecol. Evol.* 4:2019–2031.

Csilléry K, Blum MGB, Gaggiotti OE, François O. 2010. Approximate Bayesian

Computation (ABC) in practice. *Trends Ecol. Evol.* 25:410–418.

Danecek P, Auton A, Abecasis G, Albers CA, Banks E, DePristo MA, Handsaker RE, Lunter

G, Marth GT, Sherry ST, et al. 2011. The variant call format and VCFtools. *Bioinformatics*

27:2156–2158.

Delaneau O, Howie B, Cox AJ, Zagury J-F, Marchini J. 2013. Haplotype Estimation Using

Sequencing Reads. *Am. J. Hum. Genet.* 93:687–696.

Díaz S, Settele J, Brondízio ES, Ngo HT, Agard J, Arneth A, Balvanera P, Brauman KA,

Butchart SHM, Chan KMA, et al. 2019. Pervasive human-driven decline of life on Earth points to

the need for transformative change. *Science* [Internet] 366. Available from:

https://science.sciencemag.org/content/366/6471/eaax3100

Dittberner H, Becker C, Jiao W-B, Schneeberger K, Hölzel N, Tellier A, Meaux J de. 2019.
Strengths and potential pitfalls of hay transfer for ecological restoration revealed by RAD-seq
analysis in floodplain Arabis species. *Mol. Ecol.* 28:3887–3901.

Durand EY, Patterson N, Reich D, Slatkin M. 2011. Testing for Ancient Admixture
between Closely Related Populations. *Mol. Biol. Evol.* 28:2239–2252.

Durvasula A, Fulgione A, Gutaker RM, Alacakaptan SI, Flood PJ, Neto C, Tsuchimatsu T,
Burbano HA, Picó FX, Alonso-Blanco C, et al. 2017. African genomes illuminate the early history
and transition to selfing in *Arabidopsis thaliana*. *Proc. Natl. Acad. Sci.* 114:5213–5218.

Edelman NB, Frandsen PB, Miyagi M, Clavijo B, Davey J, Dikow RB, García-Accinelli G,
Belleghem SMV, Patterson N, Neafsey DE, et al. 2019. Genomic architecture and introgression
shape a butterfly radiation. *Science* 366:594–599.

Eichenberg D, Bowler DE, Bonn A, Bruelheide H, Grescho V, Harter D, Jandt U, May R,
Winter M, Jansen F. 2020. Widespread decline in Central European plant diversity across six
decades. *Glob. Change Biol.* [Internet] n/a. Available from:
https://onlinelibrary.wiley.com/doi/abs/10.1111/gcb.15447

Etter PD, Bassham S, Hohenlohe PA, Johnson EA, Cresko WA. 2011. SNP discovery and
genotyping for evolutionary genetics using RAD sequencing. *Methods Mol. Biol. Clifton NJ*
772:157–178.

Gao F, Ming C, Hu W, Li H. 2016. New Software for the Fast Estimation of Population
Recombination Rates (FastEPRR) in the Genomic Era. *G3 GenesGenomesGenetics* 6:1563–1571.

Goudet J, Jombart T. 2015. hierfstat: Estimation and Tests of Hierarchical F-Statistics.
Available from: https://cran.r-project.org/web/packages/hierfstat/index.html

Goulet BE, Roda F, Hopkins R. 2017. Hybridization in Plants: Old Ideas, New Techniques.
Plant Physiol. 173:65–78.

Green RE, Krause J, Briggs AW, Maricic T, Stenzel U, Kircher M, Patterson N, Li H, Zhai W,
Fritz MH-Y, et al. 2010. A Draft Sequence of the Neandertal Genome. *Science* 328:710–722.

Hand R, Gregor T. 2006. Die Verbreitung von Arabis sagittata In Deutschland. Ergebnisse
einer Herbarstudie. *Kochia* 1:21–31.

Harris K, Nielsen R. 2016. The Genetic Cost of Neanderthal Introgression. *Genetics*

203:881–891.

Hedrick PW. 2013. Adaptive introgression in animals: examples and comparison to new

mutation and standing variation as sources of adaptive variation. *Mol. Ecol.* 22:4606–4618.

Hewitt G. 2000. The genetic legacy of the Quaternary ice ages. *Nature* 405:907–913.

Hibbins MS, Hahn MW. 2019. The Timing and Direction of Introgression Under the

Multispecies Network Coalescent. *Genetics* [Internet]. Available from:

https://www.genetics.org/content/early/2019/01/21/genetics.118.301831

Hölzel N. 2005. Seedling recruitment in flood-meadow species: The effects of gaps, litter

and vegetation matrix. *Appl. Veg. Sci.* 8:115–124.

Hopkins R, Rausher MD. 2012. Pollinator-Mediated Selection on Flower Color Allele

Drives Reinforcement. *Science* 335:1090–1092.

Jay F, Boitard S, Austerlitz F. 2019. An ABC Method for Whole-Genome Sequence Data:

Inferring Paleolithic and Neolithic Human Expansions. *Mol. Biol. Evol.* 36:1565–1579.

Jombart T, Kamvar ZN, Lustrik R, Collins C, Beugin M-P, Knaus B, Solymos P, Schliep K,

Ahmed I, Cori A, et al. 2016. adegenet: Exploratory Analysis of Genetic and Genomic Data.

Available from: https://cran.r-project.org/web/packages/adegenet/index.html

Juric I, Aeschbacher S, Coop G. 2016. The Strength of Selection against Neanderthal

Introgression. *PLOS Genet.* 12:e1006340.

Kahle D, Wickham H. 2013. ggmap: Spatial Visualization with ggplot2. *R J.* 5:144–161.

Katalin C, Louisiane L, Olivier F, Michael B. 2015. abc: Tools for Approximate Bayesian

Computation (ABC). Available from: https://CRAN.R-project.org/package=abc

Kawabe A, Nukii H, Furihata HY. 2018. Exploring the History of Chloroplast Capture in

Arabis Using Whole Chloroplast Genome Sequencing. *Int. J. Mol. Sci.* [Internet] 19. Available

from: https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5855824/

Kenney AM, Sweigart AL. 2016. Reproductive isolation and introgression between

sympatric *Mimulus* species. *Mol. Ecol.* 25:2499–2517.

Knaus BJ, Grünwald NJ. 2017. VCFR: a package to manipulate and visualize variant call

format data in R. *Mol. Ecol. Resour.* 17:44–53.

Koboldt DC, Zhang Q, Larson DE, Shen D, McLellan MD, Lin L, Miller CA, Mardis ER, Ding

L, Wilson RK. 2012. VarScan 2: Somatic mutation and copy number alteration discovery in

cancer by exome sequencing. *Genome Res.* 22:568–576.

Kopelman NM, Mayzel J, Jakobsson M, Rosenberg NA, Mayrose I. 2015. Clumpak: a

program for identifying clustering modes and packaging population structure inferences across

K. *Mol. Ecol. Resour.* 15:1179–1191.

Kryvokhyzha D, Salcedo A, Eriksson MC, Duan T, Tawari N, Chen J, Guerrina M, Kreiner

JM, Kent TV, Lagercrantz U, et al. 2019. Parental legacy, demography, and admixture influenced

the evolution of the two subgenomes of the tetraploid Capsella bursa-pastoris (Brassicaceae).

PLoS Genet. 15:e1007949.

Lenormand T. 2002. Gene flow and the limits to natural selection. *Trends Ecol. Evol.*

17:183–189.

Li H, Durbin R. 2009. Fast and accurate short read alignment with Burrows–Wheeler

transform. *Bioinformatics* 25:1754–1760.

Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, Marth G, Abecasis G, Durbin

R, 1000 Genome Project Data Processing Subgroup. 2009. The Sequence Alignment/Map

format and SAMtools. *Bioinforma. Oxf. Engl.* 25:2078–2079.

Ma Y, Wang J, Hu Q, Li J, Sun Y, Zhang L, Abbott RJ, Liu J, Mao K. 2019. Ancient

introgression drives adaptation to cooler and drier mountain habitats in a cypress species

complex. *Commun. Biol.* 2:1–12.

Marburger S, Monnahan P, Seear PJ, Martin SH, Koch J, Paajanen P, Bohutínská M,

Higgins JD, Schmickl R, Yant L. 2019. Interspecific introgression mediates adaptation to whole

genome duplication. *Nat. Commun.* 10:5218.

Marcet-Houben M, Gabaldón T. 2015. Beyond the Whole-Genome Duplication:

Phylogenetic Evidence for an Ancient Interspecies Hybridization in the Baker's Yeast Lineage.

PLOS Biol. 13:e1002220.

Marin J-M, Raynal L, Pudlo P, Robert CP, Estoup A. 2019. abcrf: Approximate Bayesian

Computation via Random Forests. Available from: https://CRAN.R-project.org/package=abcrf

Martin M. 2011. Cutadapt removes adapter sequences from high-throughput

sequencing reads. *EMBnet.journal* 17:10–12.

Martin SH, Davey JW, Jiggins CD. 2015. Evaluating the use of ABBA-BABA statistics to
locate introgressed loci. *Mol. Biol. Evol.* 32:244–257.

Martin SH, Jiggins CD. 2017. Interpreting the genomic landscape of introgression. *Curr.*
Opin. Genet. Dev. 47:69–74.

Mathar W, Kleinebecker T, Hölzel N. 2015. Environmental variation as a key process of
co-existence in flood-meadows.Collins B, editor. *J. Veg. Sci.* 26:480–491.

Mattila TM, Tyrmi J, Pyhäjärvi T, Savolainen O. 2017. Genome-Wide Analysis of
Colonization History and Concomitant Selection in Arabidopsis lyrata. *Mol. Biol. Evol.* 34:2665–
2677.

McCauley DE. 1995. The use of chloroplast DNA polymorphism in studies of gene flow in
plants. *Trends Ecol. Evol.* 10:198–202.

Mitchell N, Owens GL, Hovick SM, Rieseberg LH, Whitney KD. 2019. Hybridization speeds
adaptive evolution in an eight-year field experiment. *Sci. Rep.* 9:6746.

Novikova PY, Hohmann N, Nizhynska V, Tsuchimatsu T, Ali J, Muir G, Guggisberg A,
Paape T, Schmid K, Fedorenko OM, et al. 2016. Sequencing of the genus Arabidopsis identifies a
complex history of nonbifurcating speciation and abundant trans-specific polymorphism. *Nat.*
Genet. 48:1077–1082.

Novotná I, Czapik R. 1974. Studies on the Progenies of Hybrids from the Arabis hirsuta
Complex. *Folia Geobot. Phytotaxon.* 9:341–357.

Ogle D, Wheeler P, Dinno A. 2020. FSA: Simple Fisheries Stock Assessment Methods.
Available from: https://CRAN.R-project.org/package=FSA

Onge KRS, Källman T, Slotte T, Lascoux M, Palmé AE. 2011. Contrasting demographic
history and population structure in Capsella rubella and Capsella grandiflora, two closely
related species with different mating systems. *Mol. Ecol.* 20:3306–3320.

Ossowski S, Schneeberger K, Lucas-Lledó JI, et al. The rate and molecular spectrum of
spontaneous mutations in Arabidopsis thaliana. *Science*. 2010;327(5961):92-94.

Paape T, Briskine RV, Halstead-Nussloch G, Lischer HEL, Shimizu-Inatsugi R, Hatakeyama
M, Tanaka K, Nishiyama T, Sabirov R, Sese J, et al. 2018. Patterns of polymorphism and selection

in the subgenomes of the allopolyploid Arabidopsis kamchatica. *Nat. Commun.* 9:3909.

Paradis E, Jombart T, Schliep K, Potts A, Winter D. 2016a. pegas: Population and

Evolutionary Genetics Analysis System. Available from: https://cran.r-

project.org/web/packages/pegas/index.html

Paradis E, Jombart T, Schliep K, Potts A, Winter D. 2016b. pegas: Population and

Evolutionary Genetics Analysis System. Available from: https://cran.r-

project.org/web/packages/pegas/index.html

Purcell S. 2009. PLINK.

Purcell S, Neale B, Todd-Brown K, Thomas L, Ferreira MAR, Bender D, Maller J, Sklar P,

de Bakker PIW, Daly MJ, et al. 2007. PLINK: a tool set for whole-genome association and

population-based linkage analyses. *Am. J. Hum. Genet.* 81:559–575.

R Development Core Team. 2008. R: A Language and Environment for Statistical

Computing. Vienna, Austria: R Foundation for Statistical Computing Available from:

http://www.R-project.org

Racimo F, Marnetto D, Huerta-Sánchez E. 2017. Signatures of Archaic Adaptive

Introgression in Present-Day Human Populations. *Mol. Biol. Evol.* 34:296–317.

Ravinet M, Faria R, Butlin RK, Galindo J, Bierne N, Rafajlović M, Noor M a. F, Mehlig B,

Westram AM. 2017. Interpreting the genomic landscape of speciation: a road map for finding

barriers to gene flow. *J. Evol. Biol.* 30:1450–1477.

Raynal L, Marin J-M, Pudlo P, Ribatet M, Robert CP, Estoup A. 2019. ABC random forests

for Bayesian parameter inference. *Bioinformatics* 35:1720–1728.

Rieseberg LH, Archer MA, Wayne RK. 1999. Transgressive segregation, adaptation and

speciation. *Heredity* 83 (Pt 4):363–372.

Rieseberg LH, Raymond O, Rosenthal DM, Lai Z, Livingstone K, Nakazato T, Durphy JL,

Schwarzbach AE, Donovan LA, Lexer C. 2003. Major Ecological Transitions in Wild Sunflowers

Facilitated by Hybridization. *Science* 301:1211–1216.

Rougemont Q, Bernatchez L. 2018. The demographic history of Atlantic salmon (*Salmo*

salar) across its distribution range reconstructed from approximate Bayesian computations*:

ATLANTIC SALMON HISTORY AND LINKED SELECTION. *Evolution* 72:1261–1277.

Schumer M, Cui R, Powell DL, Dresner R, Rosenthal GG, Andolfatto P. 2014. High-

resolution mapping reveals hundreds of genetic incompatibilities in hybridizing fish

species.McVean G, editor. *eLife* 3:e02535.

Schumer M, Rosenthal GG, Andolfatto P. 2018. What do we mean when we talk about

hybrid speciation? *Heredity* 120:379–382.

Schumer M, Xu C, Powell DL, Durvasula A, Skov L, Holland C, Blazier JC, Sankararaman S,

Andolfatto P, Rosenthal GG, et al. 2018. Natural selection interacts with recombination to

shape the evolution of hybrid genomes. *Science* 360:656–660.

Seehausen O. 2004. Hybridization and adaptive radiation. *Trends Ecol. Evol.* 19:198–207.

Sellinger TPP, Awad DA, Moest M, Tellier A. 2020. Inference of past demography,

dormancy and self-fertilization rates from whole genome sequence data. *PLOS Genet.*

16:e1008698.

Setter D, Mousset S, Cheng X, Nielsen R, DeGiorgio M, Hermisson J. 2020.

VolcanoFinder: Genomic scans for adaptive introgression. PLOS Genetics 16(6): e1008867

Small ST, Labbé F, Lobo NF, Koekemoer LL, Sikaala CH, Neafsey DE, Hahn MW, Fontaine

MC, Besansky NJ. 2020. Radiation with reticulation marks the origin of a major malaria vector.

Proc. Natl. Acad. Sci. 117:31583–31590.

Staab PR, Zhu S, Metzler D, Lunter G. 2015. scrm: efficiently simulating long sequences

using the approximated coalescent with recombination. *Bioinformatics* 31:1680–1682.

Steinrücken M, Spence JP, Kamm JA, Wieczorek E, Song YS. 2018. Model-based

detection and analysis of introgressed Neanderthal ancestry in modern humans. *Mol. Ecol.*

27:3873–3888.

Stelkens RB, Brockhurst MA, Hurst GDD, Greig D. 2014. Hybridization facilitates

evolutionary rescue. *Evol. Appl.* 7:1209–1217.

Suarez-Gonzalez A, Lexer C, Cronk QCB. 2018a. Adaptive introgression: a plant

perspective. *Biol. Lett.* 14:20170688.

Suarez-Gonzalez A, Lexer C, Cronk QCB. 2018b. Adaptive introgression: a plant

perspective. *Biol. Lett.* 14:20170688.

Takou M, Hämälä T, Steige KA, Koch E, Dittberner H, Yant L, Genete M, Sunyaev S,

Castric V, Vekemans X, et al. 2019. Maintenance of adaptive dynamics in a bottlenecked range-

edge population that retained outcrossing. *bioRxiv*:709873.

Taylor SA, Larson EL. 2019. Insights from genomes into the evolutionary importance and

prevalence of hybridization in nature. *Nat. Ecol. Evol.* 3:170–177.

Tigano A, Friesen VL. 2016. Genomics of local adaptation with gene flow. *Mol. Ecol.*

25:2144–2164.

Titz W. 1979. Die Interfertilitätsbeziehungen europäischer Sippen derArabis hirsuta-

Gruppe (Brassicaceae). *Plant Syst. Evol.* 131:291–310.

Todesco M, Owens GL, Bercovich N, Légaré J-S, Soudi S, Burge DO, Huang K, Ostevik KL,

Drummond EBM, Imerovski I, et al. 2020. Massive haplotypes underlie ecotypic differentiation

in sunflowers. *Nature* 584:602–607.

Vallejo-Marín M, Hiscock SJ. 2018. Hybridization and hybrid speciation under global

change. *New Phytol.* 211:1170–1187.

Walter GM, Richards TJ, Wilkinson MJ, Blows MW, Aguirre JD, Ortiz-Barrientos D. 2020.

Loss of ecologically important genetic variation in late generation hybrids reveals links between

adaptation and speciation. *Evol. Lett.* 4:302–316.

Wickham H. 2009. Ggplot2: elegant graphics for data analysis. New York: Springer

Yeaman S. 2015. Local Adaptation by Alleles of Small Effect*. *Am. Nat.* 186:S74–S89.

Yu G. 2018. scatterpie: Scatter Pie Plot. Available from: https://CRAN.R-

project.org/package=scatterpie

1159
1160
1161