Differences in gait stability and acceleration characteristics between healthy young and older females

Yuge Zhang[1], Xinglong Zhou[2], Mirjam Pijnappels[1], Sjoerd M. Bruijn [1,2,4*]

[1]Department of Human Movement Sciences, Faculty of Behavioural and Movement Sciences, Vrije

Universiteit Amsterdam, Amsterdam, The Netherlands

[2]Department of Sports Human Science, Beijing Sport University, Beijing, PR China

[3]Institute of Brain and Behavior Amsterdam, Amsterdam, The Netherlands

[4]Biomechanics Laboratory, Fujian Medical University, Quanzhou, Fujian, PR China

[*] **Corresponding:**
Sjoerd M. Bruijn
s.m.bruijn@gmail.com

Abstract

Our aim was to evaluate differences in gait acceleration intensity, variability and stability of feet
and trunk between older females and young females using inertial sensors. Twenty older females
(OF; mean age 68.4, SD 4.1 years) and eighteen young females (YF; mean age 22.3, SD 1.7 years)
were asked to walk straight for 100 meters at their preferred speed, while wearing inertial sensors
on heels and lower back. We calculated spatiotemporal measures, foot and trunk acceleration
characteristics and their variability, as well as trunk stability using the local divergence exponent
(LDE). Two-way analysis of variance, Student's t-test, and Mann–Whitney test were used to
compare statistical difference of measures between groups. Cohen's d effects were calculated for
each variable. Foot maximum vertical acceleration and amplitude, trunk-foot vertical acceleration
attenuation, as well as their variability were significantly smaller in OF than in YF. In contrast,
trunk mediolateral acceleration amplitude, maximum vertical acceleration, and amplitude, as well
as their variability were significantly larger in OF than in YF. Moreover, OF showed lower stability
(i.e., higher LDE values) in ML acceleration, ML and VT angular velocity of the trunk. Even though
we measured healthy older females, these participants showed lower vertical foot accelerations with
higher vertical trunk acceleration, lower trunk-foot vertical acceleration attenuation, less gait
stability, and more variability of the trunk, and hence, were more likely to fall. Moreover, the
acceleration of trunk was sensitive to age effects, both in variability and stability.

Keywords: walking, aging, wearable system, motor control, balance

1 Introduction

Falls among older adults are the leading indirect cause of disability and death (Tang et al., 2017). Epidemiological studies have shown that the 30% of people aged 65 years and older fall, with an increase in incidence to 40% in people over 80 years (Weber et al., 2018). This is due to poorer physiological function and control of stability with ageing (Winter, 1995). In China, 53% of falls occur while walking (Xia et al., 2010), and hence, it is particularly important to pay attention to gait performance of older adults for early identification of stability problems to prevent falls. Moreover, many studies have shown that among people over 60 years, females were more likely to fall (Tinetti et al., 1988;Daley and Spinks, 2000;De Rekeneire et al., 2003), as about 65% of women and 44% of men fell in their usual place of residence (Masud and Morris, 2001). Therefore, we focused on gait stability of females in our study.

There are several ways to evaluate gait, such as clinical function tests, questionnaires and measurements in a biomechanics laboratory (Hamacher et al., 2011). Questionnaires and clinical tests cannot reflect gait performance outside the laboratory, and sometimes have poor objectivity (van Schooten et al., 2015a). Gait assessment in a biomechanical laboratory is accurate but costly, time-consuming and limited to space and time (Terrier and Deriaz, 2011). Nowadays, portable ambulatory measurement systems such as inertial sensors can be used to collect gait data in people's own environment, to identify gait stability problems (Tao et al., 2012;Zhu et al., 2012;Weiss et al., 2013).

Gait stability reflects the ability to keep walking in the face of perturbations (Pai and Bhatt, 2007;Bruijn et al., 2013). Dynamical systems and non-linear time series analysis can been used to evaluate gait stability by quantifying the complex and chaotic characteristics of the human body (Bressel, 2004). One of these measures, the local divergence exponent (LDE) has been shown to have good reliability and validity (England and Granata, 2007;Son et al., 2009;Hu et al., 2012). The LDE quantifies the average exponential rate of divergence of neighboring trajectories in state space, and provides a direct measure of the sensitivity of a system to small perturbations (Dingwell and Marin, 2006).

Internal perturbations of the human body cause variability and randomness in gait (Zhang et al., 2011). If gait is within a stable range, people would not need to correct this variability. Increased variability likely reflects a less automatic gait pattern, instability and increased susceptibility to falls (Weiss et al., 2013). Studies also confirmed that variability in some gait characteristics (such as stride length, stride width, stride time) is highly related to the risk of falling (O'Loughlin et al., 1994;Chau et al., 2005). However, some studies suggested that variability is not equal to stability, as the level of variability was not necessarily negatively related to the level of stability (Li et al., 2005;van Emmerik et al., 2016).

As the control of stability in gait declines with ageing, we aimed to use inertial sensors to assess differences in gait stability and variability between healthy young and older females. In doing so, we focused on data obtained from trunk as well as foot sensors and calculated acceleration intensity, stability, and variability measures. We hypothesized that older females have a lower gait stability and increase variability on trunk accelerations compared with younger females.

2 Materials and Method

2.1 Participants

A total of 20 healthy older females (OF) and 18 younger females (YF) were recruited from the campus of Beijing Sport university, China (Table 1). None of our participants had any orthopedic or neurological disorders, acute pain or other complaints that might have affected gait and they were all able to walk at least 20 m. All participants were informed about the research procedures and the protocol was approved by the Ethics Committee of Sports Science Experiment of Beijing Sport University (approval number: 2021010H).

2.2 Data Acquisition

Participants wore three inertial sensors (Xsens MTw Awinda, the Netherland) on heels and on the lumbar region of the trunk, using the supplied elastic belt. These sensors had a sample rate of 100 samples/s and a range of -160 m/s^2 and +160 m/s^2. Data collection was synchronized between sensors. All participants wore the same model of shoes. They were asked to walk 100 meters on a straight running track at a self-selected speed, since gait variability is expected minimal at this speed for healthy people (Hausdorff et al., 1997).

2.3 Gait measures

MATLAB (R2019b, MathWorks Inc, Natick, MA, USA) was used to analyze data without the first and last steps. Each gait cycle was identified from the sagittal plane angular velocity of foot sensors with three gait events: heel-strike (T_{heel_strike}), toe-off (T_{toe_off}) and foot-flat (T_{foot_flat}) (Mariani et al., 2010). Stride time was defined as the duration between two consecutive T_{heel_strike}. Combined with the gait events of both feet, we got the initial double support period (IDS) and the terminal double support period (TDS).

For the trunk sensor, sensor data were realigned to a coordinate system based on the accelerometer's orientation with respect to gravity (vertical axis) and optimization of left-right symmetry (mediolateral axis) (Rispens et al., 2015; Van Schooten et al., 2015b).

For the foot sensors, initial displacements were calculated by integrating linear accelerations twicer for each gait cycle (in the global coordinate system), using zero-velocity-update method to eliminate drift, assuming linearity of the drift (Skog et al., 2010). The hence obtained direction of displacement was not necessary along the x or y axis of the global coordinate system. To obtain meaningful stride lengths, we thus rotated the obtained positions, the acceleration and angular velocity of the feet to a coordinate system that was aligned with the direction of walking (i.e., end position minus starting position), with the vertical axis being vertical. Then, walking speed was obtained by dividing the distance of the walking direction by the time.

For acceleration measures, maximum vertical acceleration of feet and trunk were calculated to reflect the intensity of ground contact (Gill and O'Connor, 2003). Another study also indicated that by using real-time tibial acceleration data as visual feedback, runners reduced ground reaction force loading to lower the risk of stress fractures (Crowell et al., 2010). Therefore, we defined trunk-foot vertical acceleration attenuation as the difference in maximum vertical acceleration between trunk and foot, which represents the impact absorption of the lower limbs. Acceleration amplitude (in the coordinate system prescribed by the walking direction, see above) for each direction (AP, ML and VT) was calculated as the range of acceleration in a gait cycle.

For above measures of each person, after getting the mean and standard deviation (SD) over
all cycles (see Table 2, Table3), we obtained coefficient of variation (CV) by dividing the standard
deviation by the mean (Kavanagh and Menz, 2008) (see Supplementary Table 1).
We calculated the LDE of acceleration and angular velocity of each dimension separately (in
the coordinate system prescribed by the walking direction, see above). The time series of 50 gait
cycles was normalized into 5000 samples, with the average of 100 samples per cycle. From these
data, state spaces were reconstructed using the method of correlation integral (C-C method), which
not only can determine both embedding dimension and delay time, but also has a good robustness
to the noise in small amount of data (Kim et al., 1999) (see Supplementary Tables 2 and 3 for
dimension and delay values). LDE was expressed as the mean logarithmic rate of divergence per
stride using Rosenstein's method (Rosenstein et al., 1993). Higher values of the LDE indicate a
lower local stability.

2.4 Statistical Analysis

Normality was assessed using the Kolmogorov–Smirnov test. For measures of the left and right
feet, differences were tested using two-way ANOVAs, with within-subject factor Foot (left and
right) and between-subject factor Group (YF and OF). For other measures, we used Student's t-
tests to compare between age groups. For LDE, which appeared not distributed normally, we
compared between groups using the Mann–Whitney test. For all measures, $p<0.05$ was considered
as a significant effect. Cohen's d effects were calculated for each variable as the difference between
group means divided by the group pooled standard deviation. Magnitudes of $d = 0.01, 0.20, 0,50,$
0.80, 1.20 and 2.0 were considered very small, small, medium, large, very large and huge,
separately (Sawilowsky, 2009;Charach et al., 2011;Cohen, 2013).

3   Results

Descriptive characteristics of the participants are summarized in Table 1. OF were significantly
older, shorter and had a higher weight and higher BMI than YF. The mean age of OF and YF was
68.4 and 22.3, respectively.
Table 2 shows the mean values for all measures. We found no interaction between Foot and
Group for any of the outcome measures, and no significant effect of Foot. Hence, all variables that
were calculated for both feet are displayed as averages over both feet. OF had higher maximum
vertical acceleration of the trunk than YF, with medium effect size (0.75), but smaller maximum
vertical acceleration of the feet than YF, with very large effect size (1.34). As a result, OF had
significantly smaller trunk-foot vertical acceleration attenuation, with a very large effect size of 1.8.
In addition, OF's vertical accelerations amplitude of the feet were significantly smaller than YF,
with medium effect size (-0.59). For the trunk, OF's ML and VT acceleration amplitude were
significantly larger than YF, and the effect size of the latter was largest (0.76). The LDE of trunk
from ML acceleration, and from ML and VT angular velocity were significantly larger (less stable)
for OF than for YF, with large (1.01), very large (1.48) and medium effect size (0.66), respectively.
Table 3 shows the variability of all measures. No significant difference in variability of spatial-
temporal gait measures were found between groups. The variability of maximum vertical
acceleration of the feet was significantly smaller for OF than YF, and its effect size was 1.70. While
for trunk, the variability of the maximum vertical acceleration was significantly larger for the OF

(medium effect size 0.72). The variability of trunk-foot vertical acceleration attenuation was smaller
in OF than in YF, (effect size very large, 1.87). OF had significantly smaller variability of
acceleration amplitude of the feet in three directions than YF, with huge effect size in ML direction
(2.53) and a large effect size in the VT direction (1.01). For the trunk, OF's variability of
acceleration amplitude was significantly larger than YF in ML and VT direction, with effect sizes
of 0.41 and 0.86, respectively. The CV of gait measures showed largely the same pattern as the SD
(see Supplementary Table 1).

4 Discussion

4.1 Mean gait measures

In this study, we used inertial sensors to evaluate differences in acceleration intensity, variability
and stability of feet and trunk during gait between healthy young and older females. Although older
adults generally were suggested to walk slower due to physical limitations like muscle weakness or
loss of flexibility (Hamacher et al., 2014), the OF in our study walked at similar preferred speed
and stride length as the YF.
We found a reduction in foot vertical maximum acceleration in OF, which probably reflected
a reduction of peak ground reaction forces. Such a reduction of ground reaction forces could result
from a crouch-like gait, which has been shown in young adults to lead to a reduction of the peak
ground reaction force (Li et al., 1996;Grasso et al., 2000). Such a crouch like gait may increase the
metabolic cost of locomotion in the elderly (Carey and Crompton, 2005). Although the trunk
segment plays a key role in damping gait-related oscillations (Kavanagh et al., 2006), the damping
of oscillations by the trunk in the vertical direction has been suggested to be minor (Prince et al.,
1994;Kavanagh et al., 2004). In our study, we found a lower trunk-foot vertical acceleration
attenuation and a higher trunk acceleration amplitude in OF, which implies a decreased cushioning
(impact absorption) and hence less preservation of the head's stability (Menz et al., 2003). Even
though foot (vertical) accelerations were lower in OF, suggesting less impact, the OF were not able
to attenuate the higher accelerations in the trunk. Considering that two-thirds of the weight of the
human body is in the upper body, such higher trunk accelerations may be destabilizing, which may
cause falls (Woollacott and Tang, 1997).
For stability, LDE calculated from trunk time-series data have been shown to better reflect
differences in gait stability due to age than LDE calculated from data of other segments (Punt et al.,
2015). In our study, OF showed significantly lower local dynamic stability (higher LDE) in ML
acceleration, ML and VT angular velocity. Among these, the LDE calculated from trunk ML
angular velocity had the largest effect size. As stability in the ML direction needs more control than
stability in the AP direction during gait (Bruijn et al., 2009;O'Connor and Kuo, 2009), decreased
LDE of trunk angular velocity in ML direction could be an early indicator of gait stability problems.

4.2 Variability measures

All participants in this study walked under the same environmental conditions. Thus, any between-
subject differences in variability arose from differences in (internal) neuromotor noise and not
(external) environmental noise. No differences were found in the variability of spatiotemporal
measures, which was consistent with a previous study showing that temporal gait variability of

older non-fallers was not significantly different from young adults in terms of standard deviation
and coefficient of variation (Hausdorff et al., 1997).
Our OF walked with similar variability of maximum vertical acceleration of feet variability
compared to YF (Table 3 and Supplementary Table 4). However, the variability of ML and VT
acceleration amplitude of the trunk was larger for the OF, which could suggest OF are at a higher
risk of balance loss and falling (Kavanagh and Menz, 2008).
All in all, our findings suggests that stability of the trunk might be a more sensitive indicator
of locomotor impairment and potential future risk of falls than changes in variability of the trunk,
as the LDE had higher effect sizes (Kang and Dingwell, 2008). Measures of variability of
acceleration of the feet showed even higher effect sizes and might thus be even more useful.
However, here, it should be noted that these effects were opposite from theoretically expected, with
the OF having lower (means and variability) acceleration of the foot.

4.3 Limitations

All tests in our study were aimed at testing the same hypothesis, that is, OF are less stable and more
variable than YF, hence, we did not use a correction for multiple testing. Nonetheless, not correcting
may lead to Type I errors, and thus, some caution is warranted. Furthermore, the older participants
in our study were quite fit and additional studies are needed to further investigate the applicability
of acceleration attenuation when studying older adults. Future research can expand the sample size
and conduct a multi-center study to obtain more representative results.

5   Conclusions

Although healthy older females had similar walking speed and spatiotemporal parameters as young
females during steady state walking, they showed lower vertical foot accelerations and higher
vertical trunk accelerations, suggesting less impact, and less absorption of the impact. In addition,
lower gait stability and higher variability of trunk movements for older females also indicated they
were more likely to fall. The measures derived from the accelerations of the trunk were sensitive to
reflect the gait instability as expected, especially trunk-foot vertical acceleration attenuation and its
variability. While the variability of foot acceleration amplitudes was also sensitive to age, these
differences were opposite from expected, making it harder to draw any conclusion as to their
usefulness for fall prediction.

6   Conflicts of Interest

221 The authors declare no conflict of interest.

7   Author Contributions

Yuge Zhang: Methodology, Software, Formal analysis, Investigation, Data curation, Writing-
Original Draft, Writing-Review and Editing, Visualization. **Xinglong Zhou:** Conceptualization,
Resources. **Mirjam Pijnappels:** Formal analysis, Writing-review and Editing, Supervision,
Funding acquisition. **Sjoerd M. Bruijn:** Methodology, Software, Formal analysis, Writing-
Review and Editing, Supervision, Funding acquisition. All authors have read and agreed to the
published version of the manuscript.

8   Acknowledgments

YZ was funded by a CSC Scholarship Council (CSC) fellowship (202009110145). MP funded by
a VIDI Grant (no. 91714344) from the Dutch Organization for Scientific Research (NWO). SMB
was funded by a VIDI grant (016.Vidi.178.014) from the Dutch Organization for Scientific
Research (NWO).

9   References

Bressel, E. (2004). Innovative Analyses of Human Movement: Analytical Tools for Human
Movement Research. *Medicine & Science in Sports & Exercise* 36**,** 1834.
Bruijn, S.M., Meijer, O., Beek, P., and Van Dieen, J.H. (2013). Assessing the stability of human
locomotion: a review of current measures. *Journal of the Royal Society Interface* 10**,**
20120999.
Bruijn, S.M., Van Dieën, J.H., Meijer, O.G., and Beek, P.J. (2009). Is slow walking more stable?
Journal of biomechanics 42**,** 1506-1512.
Carey, T.S., and Crompton, R.H. (2005). The metabolic costs of 'bent-hip, bent-knee' walking in
humans. *Journal of Human Evolution* 48**,** 25-44.
Charach, A., Dashti, B., Carson, P., Booker, L., Lim, C.G., Lillie, E., Yeung, E., Ma, J., Raina, P.,
and Schachar, R. (2011). Attention deficit hyperactivity disorder: effectiveness of
treatment in at-risk preschoolers; long-term effectiveness in all ages; and variability in
prevalence, diagnosis, and treatment.
Chau, T., Young, S., and Redekop, S. (2005). Managing variability in the summary and
comparison of gait data. *Journal of neuroengineering and rehabilitation* 2**,** 22.
Cohen, J. (2013). *Statistical power analysis for the behavioral sciences.* Academic press.
Crowell, H.P., Milner, C.E., Hamill, J., and Davis, I.S. (2010). Reducing impact loading during
running with the use of real-time visual feedback. *journal of orthopaedic & sports*
physical therapy 40**,** 206-213.
Daley, M.J., and Spinks, W.L. (2000). Exercise, mobility and aging. *Sports medicine* 29**,** 1-12.
De Rekeneire, N., Visser, M., Peila, R., Nevitt, M.C., Cauley, J.A., Tylavsky, F.A., Simonsick,
E.M., and Harris, T.B. (2003). Is a fall just a fall: correlates of falling in healthy older
persons. The Health, Aging and Body Composition Study. *Journal of the American*
Geriatrics Society 51**,** 841-846.
Dingwell, J.B., and Marin, L.C. (2006). Kinematic variability and local dynamic stability of upper
body motions when walking at different speeds. *Journal of biomechanics* 39**,** 444-452.
England, S.A., and Granata, K.P. (2007). The influence of gait speed on local dynamic stability of
walking. *Gait & posture* 25**,** 172-178.
Gill, H., and O'connor, J. (2003). Heelstrike and the pathomechanics of osteoarthrosis: a pilot gait
study. *Journal of biomechanics* 36**,** 1625-1631.
Grasso, R., Zago, M., and Lacquaniti, F. (2000). Interactions between posture and locomotion:
motor patterns in humans walking with bent posture versus erect posture. *Journal of*
Neurophysiology 83**,** 288-300.

Hamacher, D., Hamacher, D., and Schega, L. (2014). Towards the importance of minimum toe
clearance in level ground walking in a healthy elderly population. *Gait & posture* 40**,** 727-
729.
Hamacher, D., Singh, N., Van Dieën, J.H., Heller, M., and Taylor, W.R. (2011). Kinematic
measures for assessing gait stability in elderly individuals: a systematic review. *Journal of*
The Royal Society Interface 8**,** 1682-1698.
Hausdorff, J.M., Edelberg, H.K., Mitchell, S.L., Goldberger, A.L., and Wei, J.Y. (1997).
Increased gait unsteadiness in community-dwelling elderly fallers. *Archives of physical*
medicine and rehabilitation 78**,** 278-283.
Hu, F., Gu, D., Dai, K., An, B., Chen, J., and Wu, Y. (2012). Nonlinear time series analysis of
gait stability during walking. *Journal of Medical Biomechanics* 27**,** 51-57.
Kang, H.G., and Dingwell, J.B. (2008). Effects of walking speed, strength and range of motion on
gait stability in healthy older adults. *Journal of biomechanics* 41**,** 2899-2905.
Kavanagh, J., Barrett, R., and Morrison, S. (2006). The role of the neck and trunk in facilitating
head stability during walking. *Experimental brain research* 172**,** 454.
Kavanagh, J.J., Barrett, R.S., and Morrison, S. (2004). Upper body accelerations during walking
in healthy young and elderly men. *Gait & Posture* 20**,** 291-298.
Kavanagh, J.J., and Menz, H.B. (2008). Accelerometry: a technique for quantifying movement
patterns during walking. *Gait & posture* 28**,** 1-15.
Kim, H.S., Eykholt, R., and Salas, J. (1999). Nonlinear dynamics, delay times, and embedding
windows. *Physica D: Nonlinear Phenomena* 127**,** 48-60.
Li, L., Haddad, J.M., and Hamill, J. (2005). Stability and variability may respond differently to
changes in walking speed. *Human movement science* 24**,** 257-267.
Li, Y., Crompton, R., Alexander, R.M., Günther, M., and Wang, W. (1996). Characteristics of
ground reaction forces in normal and chimpanzee-like bipedal walking by humans. *Folia*
Primatologica 66**,** 137-159.
Mariani, B., Hoskovec, C., Rochat, S., Bula, C., Penders, J., and Aminian, K. (2010). 3D gait
assessment in young and elderly subjects using foot-worn inertial sensors. *J Biomech* 43**,**
2999-3006.
Masud, T., and Morris, R.O. (2001). Epidemiology of falls. *Age and ageing* 30**,** 3-7.
Menz, H.B., Lord, S.R., and Fitzpatrick, R.C. (2003). Acceleration patterns of the head and pelvis
when walking on level and irregular surfaces. *Gait & posture* 18**,** 35-46.
O'connor, S.M., and Kuo, A.D. (2009). Direction-dependent control of balance during walking
and standing. *Journal of neurophysiology* 102**,** 1411-1419.
O'loughlin, J.L., Boivin, J.-F., Robitaille, Y., and Suissa, S. (1994). Falls among the elderly:
distinguishing indoor and outdoor risk factors in Canada. *Journal of epidemiology and*
community health 48**,** 488.
Pai, Y.-C., and Bhatt, T.S. (2007). Repeated-slip training: an emerging paradigm for prevention of
slip-related falls among older adults. *Physical therapy* 87**,** 1478-1491.
Prince, F., Winter, D.A., Stergiou, P., and Walt, S.E. (1994). Anticipatory control of upper body
balance during human locomotion. *Gait & Posture* 2**,** 19-25.
Punt, M., Bruijn, S.M., Wittink, H., and Van Dieën, J.H. (2015). Effect of arm swing strategy on
local dynamic stability of human gait. *Gait & posture* 41**,** 504-509.

Rispens, S.M., Van Schooten, K.S., Pijnappels, M., Daffertshofer, A., Beek, P.J., and Van Dieen,
J.H. (2015). Identification of fall risk predictors in daily life measurements: gait
characteristics' reliability and association with self-reported fall history.
Neurorehabilitation and Neural Repair 29**,** 54-61.

Rosenstein, M.T., Collins, J.J., and De Luca, C.J. (1993). A practical method for calculating
largest Lyapunov exponents from small data sets. *Physica D: Nonlinear Phenomena* 65**,**
117-134.

Sawilowsky, S.S. (2009). New effect size rules of thumb. *Journal of modern applied statistical*
methods 8**,** 26.

Skog, I., Handel, P., Nilsson, J.-O., and Rantakokko, J. (2010). Zero-velocity detection—An
algorithm evaluation. *IEEE transactions on biomedical engineering* 57**,** 2657-2666.

Son, K., Park, J., and Park, S. (2009). Variability analysis of lower extremity joint kinematics
during walking in healthy young adults. *Medical engineering & physics* 31**,** 784-792.

Tang, Y., Guo, X., Qiao, Z., and Qiu, P. (2017). Analysis on prevalence and risk factors for falls
among the elderly in communities of Beijing and Shanghai. *Chinese Journal of disease*
control & prevention 21**,** 72-76.

Tao, W., Liu, T., Zheng, R., and Feng, H. (2012). Gait analysis using wearable sensors. *Sensors*
12**,** 2255-2283.

Terrier, P., and Deriaz, O. (2011). Kinematic variability, fractal dynamics and local dynamic
stability of treadmill walking. *Journal of Neuroengineering and Rehabilitation* 8**,** 12-12.

Tinetti, M.E., Speechley, M., and Ginter, S.F. (1988). Risk factors for falls among elderly persons
living in the community. *New England journal of medicine* 319**,** 1701-1707.

Van Emmerik, R.E., Ducharme, S.W., Amado, A.C., and Hamill, J. (2016). Comparing dynamical
systems concepts and techniques for biomechanical analysis. *Journal of Sport and Health*
Science 5**,** 3-13.

Van Schooten, K.S., Pijnappels, M., Rispens, S.M., Elders, P.J., Lips, P., and Van Dieen, J.H.
(2015a). Ambulatory fall-risk assessment: amount and quality of daily-life gait predict
falls in older adults. *Journals of Gerontology Series A: Biomedical Sciences and Medical*
Sciences 70**,** 608-615.

Van Schooten, K.S., Pijnappels, M., Rispens, S.M., Elders, P.J.M., Lips, P., and Van Dieen, J.H.
(2015b). Ambulatory Fall-Risk Assessment: Amount and Quality of Daily-Life Gait
Predict Falls in Older Adults. *Journals of Gerontology Series A-biological Sciences and*
Medical Sciences 70**,** 608-615.

Weber, M., Van Ancum, J.M., Bergquist, R., Taraldsen, K., Gordt, K., Mikolaizak, A.S., Nerz,
C., Pijnappels, M., Jonkman, N.H., and Maier, A.B. (2018). Concurrent validity and
reliability of the Community Balance and Mobility scale in young-older adults. *BMC*
Geriatrics 18**,** 1-10.

Weiss, A., Brozgol, M., Dorfman, M., Herman, T., Shema, S., Giladi, N., and Hausdorff, J.M.
(2013). Does the evaluation of gait quality during daily life provide insight into fall risk?
A novel approach using 3-day accelerometer recordings. *Neurorehabilitation and neural*
repair 27**,** 742-752.

Winter, D.A. (1995). Human balance and posture control during standing and walking. *Gait &*
posture 3**,** 193-214.

Woollacott, M.H., and Tang, P.-F. (1997). Balance Control During Walking in the Older Adult:
Research and Its Implications. *Physical Therapy* 77**,** 646-660.
Xia, Q., Jiang, Y., Tang, C., and Niu, C. (2010). Study on the epidemiologic characteristics and
medical burden of falls among adults in community. *Chinese Journal of disease control &*
prevention 14**,** 647-649.
Zhang, B., Jiang, S., Yan, K., Wei, D., and Smigorski, K. (2011). Human walking analysis,
evaluation and classification based on motion capture system. *Heal. Manag.–Differ.*
*Approaches Solut***,** 361-398.
Zhu, S., Anderson, H., and Wang, Y. (Year). "A real-time on-chip algorithm for IMU-Based gait
measurement", in: *Pacific-Rim Conference on Multimedia*: Springer), 93-104.

10   Tables

365   **Table 1.** Participant characteristics

Groups	YF	OF	p-Value (T-test)
Age (years)	22.3 (1.7)	68.4 (4.1)	<0.001***
Height (m)	1.65 (0.04)	1.59 (0.05)	<0.001***
Body mass (kg)	54.66 (3.93)	63.2 (7.95)	<0.001***
BMI (kg/m^2)	20.19 (1.53)	24.96 (2.60)	<0.001***
Leg length (cm)	88.21 (3.53)	87.28 (3.19)	0.29

374   **Table 2.** Mean (and SD) of all gait measures

	OF	YF	p-value	Effect size
Spatial-temporal measures				
Stride time (s)	1.04 (0.07)	1.07(0.05)	0.06	-0.69
Initial double support period, IDS (%)	14.30(1.70)	14.00(1.20)	0.50	0.00
Terminal double support period, TDS (%)	14.10(1.60)	14.30(1.40)	0.76	0.02
Swing (%)	35.80(1.70)	35.90(1.20)	0.78	-0.01
Velocity of feet (m/s)	1.35(0.17)	1.37(0.14)	0.88	-0.06
Stride length of feet (m)	1.28(0.11)	1.35(0.12)	0.06	-0.61
VT Acceleration Maximum (m/s^2)				
VT maximum acceleration of feet	24.81(4.32)	33.52(8.78)	<0.001***	-1.34
VT maximum acceleration of trunk	18.04(2.75)	16.43(1.09)	0.011*	0.75
Trunk-foot vertical acceleration attenuation (m/s^2)	17.00(3.73)	28.62(8.61)	0.023*	-1.80
Acceleration Amplitude (m/s^2)				
AP acceleration amplitude of feet	86.53(14.35)	94.11(13.45)	0.13	0.53
ML acceleration amplitude of feet	31.25(7.01)	28.37(4.53)	0.29	-0.86
VT acceleration amplitude of feet	60.11(10.34)	68.30(9.74)	0.007**	-0.59
AP acceleration amplitude of trunk	8.97(2.69)	7.35(1.69)	0.07	0.71
ML acceleration amplitude of trunk	9.11(2.89)	8.52(1.28)	0.01*	0.26
VT acceleration amplitude of trunk	11.94(3.88)	9.65(1.39)	0.01*	0.76
Local Divergence Exponent (LDE)				
AP acceleration of trunk	1.15(0.46)	1.00(0.37)	0.28	0.35

ML acceleration of trunk	0.84(0.18)	0.68(0.12)	0.005**	1.01
VT acceleration of trunk	0.92(0.26)	0.87(0.21)	0.89	0.19
AP angular velocity of trunk	0.80(0.31)	0.69(0.30)	0.06	0.38
ML angular velocity of trunk	0.97(0.23)	0.69(0.11)	<0.001***	1.48
VT angular velocity of trunk	0.73(0.32)	0.56(0.12)	0.048*	0.66

375   Notes: AP = anteroposterior direction; ML = mediolateral direction; VT = vertical direction.
376

377 **Table 3.** Variability (and SD) of all gait measures

	OF	YF	p-value	Effect Size
Spatial-temporal Gait Measures				
Stride time (s)	0.02(0.01)	0.02(0.01)	0.64	-0.23
Initial double support period, IDS (%)	0.74(0.11)	0.71(0.11)	0.48	0.35
Terminal double support period, TDS (%)	0.80(0.20)	0.79(0.15)	0.93	0.17
Swing (%)	0.75(0.16)	0.75(0.19)	0.97	0.10
Velocity of feet (m/s)	0.06(0.03)	0.08(0.04)	0.05	-0.72
Stride length of feet (m)	0.02(0.01)	0.05(0.05)	0.13	-0.38
VT Acceleration Maximum (m/s^2)				
VT maximum acceleration of feet	4.05(1.39)	7.90(3.26)	<0.001**	-1.70
VT maximum acceleration of trunk	1.02(0.44)	0.78(0.13)	0.007**	0.72
Trunk-foot vertical acceleration attenuation (m/s^2)	4.02(1.09)	7.99(2.91)	0.008**	-1.87
Acceleration Amplitude (m/s^2)				
AP acceleration amplitude of feet	5.97(1.77)	7.52(1.91)	0.026*	-0.85
ML acceleration amplitude of feet	4.64(1.20)	5.84(1.88)	0.045*	-2.53
VT acceleration amplitude of feet	4.92(1.38)	9.69(2.72)	<0.001**	-1.01
AP acceleration amplitude of trunk	1.26(0.45)	1.05(0.31)	0.350	0.53
ML acceleration amplitude of trunk	0.98(0.48)	0.82(0.21)	0.017*	0.41
VT acceleration amplitude of trunk	1.31(0.67)	0.08(0.15)	<0.001**	0.86

378   Notes: AP = anteroposterior direction; ML = mediolateral direction; VT = vertical direction.