Working title:

Late-life fitness gains explain the absence of a selection shadow in ants

Authors: Jaimes Nino LM, Heinze, J., Oettler, J

Affiliation: Zoologie/Evolutionsbiologie, Universität Regensburg

Abstract

A key hypothesis for the occurrence of senescence is a decrease in the selection strength because of low late-life fitness – the so-called selection shadow. However, in social insects, aging is considered a plastic trait and senescence seems to be absent. By life-long tracking of 102 ant colonies, we find that queens increase sexual productivity in late life regardless of their absolute lifespan or worker investment. This indicates a genetically accommodated adaptive shift towards increasingly queen-biased caste ratios over the course of a queens' life. Furthermore, mortality decreased with age, supporting the hypothesis that aging is adaptive. We argue that selection for late life reproduction diminishes the selection shadow of old age and leads to the apparent absence of senescence in ants, in contrast to most iteroparous species.

Introduction

The near-ubiquitous occurrence of senescence, a trait without any apparent fitness benefit, has been explained by two classic prevailing theories, mutation accumulation (MA) and antagonistic pleiotropy (AP) (Haldane 1941; Kirkwood 1977; Hughes and Reynolds 2005; Maklakov et al. 2015; Flatt and Partridge 2018). These theories rely on the basic assumption of the existence of a "selection shadow" (Williams 1957). Random extrinsic mortality decreases selection pressure after organisms have reached reproductive maturity, which in turn can lead to loss-of-function and senescence. Therefore, mutations with an effect late in life could accumulate (MA) and/or genes with beneficial effect early in life can be selected for even if they are detrimental to the organism later in life (AP) (Maklakov and Chapman 2019). Such antagonistic pleiotropy is visible in the conflict over resource allocation to either reproduction or somatic maintenance (i.e. disposable soma theory, or energy trade-offs) (Kirkwood 1977; Kirkwood and Austad 2000) and/or in suboptimal gene regulation after maturation (functional trade-offs) (López-Otín et al. 2013; Maklakov and Chapman 2019). However, the huge diversity of aging patterns across metazoan species (Jones et al. 2014; Cohen 2018) calls for a critical evaluation of the general validity of these theories.

When considering social insect reproductives (e.g. ant queens), the basic assumptions underlying MA and AP may not be met (Monroy Kuhn and Korb 2016; Toth et al. 2016; Lucas and Keller 2017). Extrinsic mortality is minimized as queens live in a sheltered fortress. Keller and Genoud (Keller and Genoud 1997) argue that low extrinsic mortality of queens maintains a high selection pressure with age, while the costs of reproduction (energy intake, brood care) are outsourced to the workers. In addition, it is posited that selection at high ages is reinforced if there is no marked decrease in fecundity after maturity (Keller and Genoud 1997), predicting that classic energy trade-offs may be shaped differently or may not even exist. This has found support in studies of the model ant *Cardiocondyla obscurior* (Schrempf et al. 2005; Schrempf et al. 2011; Oettler and Schrempf 2016) where queens exhibit a positive correlation between lifespan and reproduction (Kramer et al. 2015). Furthermore, experimental manipulation of the fecundity of *C. obscurior* queens does not affect lifespan (Schrempf et al. 2017) and queens appear to have only a very brief period of senescence (Heinze and Schrempf 2012). Additional evidence for the lack of senescence comes from a gene expression study, which showed a lack of transcriptional ageing in old *C. obscurior* queens, possibly facilitated by strong selection at old age and by well-regulated anti-ageing mechanisms (Harrison et al. 2021). So how did this exceptional pattern of positive correlation between fecundity and lifespan, and the absence of senescence, evolve, and are there other trade-offs that shape queen lifespan instead?

In order to disentangle aging and the investment into somatic maintenance and sexual reproduction of ant queens it is necessary to consider the colony as a superorganism (Boomsma and Gawne 2018), comprising a soma- (i.e. workers) and a germline (i.e. queens and males), where the longevity and mortality of the individual affects the fitness of the queen (Bourke 2007; Kramer and Schaible 2013). Therefore, by manipulating the investment into caste ratio by lifelong standardization of worker number, we expected to find trade-offs at the level of the germline (i.e. lifespan and productivity of queens) and/or at the level of the colony (investment in next-generation soma- or germline), if they exist. To this end, we monitored the lifetime production of 102 individual queens in colonies, which were standardized weekly to 10, 20 or 30 workers (Fig. S1A and Fig. S1B), corresponding to natural colony size variation (Schrader et al. 2014, Fig. S2). In order to compare queen mortality with worker aging patterns, we tracked the survival of 40 workers kept in colonies with 10 or 20 marked nestmate workers.

Results and Discussion

We hypothesized that queens which experienced a worker shortage would compensate by investing less into queen production. This assumes queen control over caste fate in *C. obscurior*, in contrast to many species where caste is environmentally or genetically determined (Corona et al. 2016). Indeed, queens with 10 workers (n=31) produced significantly fewer queen pupae than queens with 20 (n=34)

(glmmTMB z-value = 2.81, IRR = 1.97, p = 0.005) and 30 workers (n=34) (z-value = 2.58, IRR = 1.78, p = 0.009, Fig. 1A) with no significant differences between 20 and 30 workers (z-value = -0.49, p = 0.877). We used a correction factor (c) based on dry weight of individuals to estimate the energy allocated to castes, considering the higher costs of queen compared to worker production, following the rationale for estimating investment sex ratios (Boomsma et al. 1995). Queens' total investment was higher in colonies with 20 and 30 workers (defined as queen*c/(queen*c + worker pupae), 10 vs. 20 workers: z-value = 4.42, IRR= 0.095, p <0.001 and 10 vs. 30: z-value = 4.08, IRR=0.086, p < 0.001, Fig. 1B), with no significant differences between queens with 20 and 30 workers (z-value = -0.45, p = 0.90). The factor c takes differences in investment into the two phenotypes into account and is sensitive to a power-conversion factor for differences in respiratory rates (see Methods). The treatment did not affect total egg production (Fig. 1C, 10 vs. 20 workers: z-value = -0.38, p = 0.70 and 10 vs. 30: z-value = -0.96, p = 0.34) or worker pupae production (Fig 1D, 10 vs. 20 workers: z-value = 0.09, p = 0.93 and 10 vs. 30: z-value = -0.39, p = 0.70). We explored whether the quality of workers was affected by measuring the head width of workers produced over months 3 to 6 of the queen's lifetime (~5 workers per month). Head width was 7.5 μm and 11.2 μm significantly smaller in workers produced in small colonies than in colonies with 20 (lme $F_{6,76}$ = 2.36, p = 0.026) and 30 workers respectively (lme $F_{6,76}$ = 3.52, p < 0.002, Fig. 2).

We could not detect a "terminal investment" strategy where queens increased their productivity until death as described for *C. obscurior* queens from a Brazilian population (Kramer et al. 2015), in queens from a Japanese population used for this experiment (Fig. 3A). Worker production tightly matched the curve of egg production (Fig 4A). Importantly, queen-biased caste allocation increased late in life irrespective of treatment (Fig 3B), in both short-lived and long-lived queens (queens with a lifespan below and above the mean lifespan of 25 weeks, Fig. 3C and D respectively). Variation in caste investment is not correlated with queen lifespan, similarly to fecundity (Schrempf et al. 2017). In support of this, the treatment did not affect lifespan (Fig. S3A, Cox proportional hazard regression model, Likelihood ratio test, X^2=1.57, p = 0.46), which was highly variable across treatments (Variation coefficient: 32.2%, Fig. S3B).

To compare mortality and fecundity patterns we mean-standardized queen age-specific mortality and fecundity (Jones et al. 2014). After an earlier increase in relative mortality, *C. obscurior* ant queens exhibit a decrease below the average level of adult mortality late in life (Fig. 4A), indicating selection against senescence. Interestingly, the mean-standardized mortality of workers is very similar to the queen's aging pattern, where relative mortality also first increased then decreased (Fig 4B). Furthermore, in contrast to the prediction that there is no marked decrease in queen fecundity after maturity (Keller and Genoud 1997), relative fecundity reaches a maximum (~16 weeks) before the median lifespan (~26 weeks). However, the relative investment in queen pupae reaches a maximum

late in life (~28 weeks). This pattern is not due to the delay in development from egg to pupa, because queen development only lasts ~5 weeks (Schrempf and Heinze 2006). A first peak in queen-biased investment occurs at an earlier age, followed by an increasing queen bias with increasing age (Fig 3A).

This caste ratio shift is independent of queen lifespan (Fig 3B, C), implying that queens are selected for late life productivity. Founder queens invest first in growing numbers of workers (ergonomic phase) and subsequently in the production of new sexuals, when the colony has reached the threshold required to enter the reproductive phase (Macevicz and Oster 1976; Oster and Wilson 1978; Beekman et al. 1998). Our data suggest that this switch in caste allocation is a fixed trait, independent of colony size and absolute lifespan of queens.

Senescence is defined as an increase in relative mortality with age due to a decrease in the strength of selection, and a decrease in relative fecundity (Hughes and Reynolds 2005). Strikingly, *C. obscurior* queens exhibit a decrease in relative mortality with age. Further, while fecundity decreases, investment into queen pupae reaches a maximum late in life. In addition, because *C. obscurior* is a polygynous species, effective removal of senescent individuals with diminishing fecundity from the population, and replacement with young queens, might lead to additional selection for a short senescent phase. Thus, *C. obscurior* queens continue to experience strong selection at high ages. In turn, the short selection shadow might have led to the absence of a distinguishable senescence phase. Whether this holds true for social insects in general, and whether this underlies the evolution of the extraordinary lifespans found in social insects, remains to be studied.

Acknowledgements
This study was funded by the Deutsche Forschungsgemeinschaft (OE549/2-2). We thank Vera Ermer, Benjamin Dofka, Julia Haschlar, Judith Weber and Lena-Marie Süß for help with the experiment, and Eva Schultner and Tomer Czaczkes for comments. **Funding:** This study was funded by the Deutsche Forschungsgemeinschaft (OE549/2-2). **Author contributions:** JH and JO designed the study, LJ and JO produced the data, LJ analyzed the data, LJ and JO wrote the original draft, all authors reviewed and edited the final version of the draft. **Competing interests:** Authors declare no competing interests. **Data and materials availability:** All data is available in the main text or the supplementary materials.

Figures

Fig. 1. Productivity of *C. obscurior* colonies across treatments, A) Total number of queen pupae, B) Investment in queen over the total caste investment (as queen and worker pupae), C) Total number of eggs, and D) of worker pupae. Significant differences are given with ** for p<0.01 and *** for p<0.001. Significant pattern of investment in graph B) is robust to different power conversion values (Table S1).

Fig. 2. Worker quality per treatment. Head width measurements of workers produced in queens of colonies with 10, 20 and 30 workers. Significant differences are given with * for p < 0.05 and ** for p < 0.01.

Fig. 3. Life-time caste allocation A) Number of worker and queen pupae produced over time, B) Queen/(Queen + Worker) Pupae ratio for all ant queens, C) Queens with lifespan below (N= 44), and D) Above the mean of 25 weeks (N = 55). After queen death, eggs and larvae were allowed to develop into pupae for a final count. Therefore, smooth splines extend ca. 4 weeks after queen death.

Fig. 4. Relative mortality and fecundity as a function of age. Mean-standardization of age by dividing age-specific mortality and fecundity by their mean after maturation (Jones et al. 2014) in A) the queen, and B) worker. The graph uses a Loess smoothing method (span = 0.75) and a confidence interval of 95%. The dashed line at y = 1, indicates when relative mortality and fertility are equivalent to mean mortality and fertility.

Materials and Methods

We set up 102 freshly eclosed queens from stock colonies of a Japanese population (OypB, from the Oonoyama Park in Naha, Okinawa) established in the laboratory since 2011. The experiment took place between January 2019 and January 2020. Queens were allowed to mate with a single wingless male and were placed in nest boxes with either 10, 20 or 30 workers from the maternal colony (N = 34). This number of workers represents the naturally occurring number in the field and corresponds closely to the first, median and third quantile of number of workers of this population (N = 62, median = 28.5, Fig. S2). The colony was set up with half of the workers selected from inside of the nest near the brood (younger nurses) of the stock colonies, and the other half from outside the nest (older foragers) in order to minimize a putative effect of worker age on the queen (Giehr et al. 2017). Colonies were kept under a 12 h of dark 22°C/12 h light 26°C cycle and fed *ad libitum* three times per week with diluted honey (0.6:1 honey: distilled water), cockroaches and flies. Once per week workers, eggs, and all pupae (worker, queen, winged and wingless male) were counted. Additionally, the number of workers was standardized to the assigned treatment, and newly produced reproductive pupae produced were removed. *C. obscurior* workers are sterile, and all

produced offspring originated from the focal queen. Queen control over caste fate was assumed, as caste fate can be determined as early as the last embryonic stage. The number of counted eggs correlates with the production of workers, queens, and the workers and queens together (Fig. S4.A-C, Kendall's Tau correlation test, p < 0.001: eggs-worker pupae= 0.59, eggs-queen pupae= 0.70, eggs-worker and queen pupae=0.73). Pupae might have been counted more precisely than eggs, especially when larger number of eggs were produced. Pupae are hardly missed, compared to eggs which tend to cluster together. Eggs and worker pupae might have been counted more than once, as development lasts a median of 8 and 18 days for eggs and worker pupae, respectively. Finally, three colonies (10 worker treatment) were not considered in the analysis as they were accidentally killed.

To examine the worker aging patterns, 40 focal worker pupae were set up in individual colonies with 10 or 20 marked workers. These two treatments were selected, as no significant differences were observed between the 20 workers and 30 workers colonies in terms of queen productivity. Marking was done by clipping the tarsae of the middle right leg. The colony was set up with brood (5 larvae in the 10 workers colonies, and 10 in the 20 workers colonies), and two wingless queens to avoid a queenless period. The number of marked workers, queens and larvae was standardized weekly to the assigned treatment, and newly produced pupae produced were removed. Dead marked workers were replaced with fresh worker pupae and clipped one or two days after eclosion to avoid confusion with the focal worker.

Offspring investment

Freshly eclosed adult workers were sampled monthly for head width measurements (from the 3rd to 6th month of the queen's life, and up to five workers depending on availability). Workers were dried, pinned, and blindly measured using a Keyence Microscope 200X. A single worker was chosen randomly and measured 10 times to obtain a proxy for measurement error (Mean = 383.61 μm, standard deviation=5.05 μm). Additionally, newly eclosed worker and queen adults from the stock population were dried at 60°C for 72h and five individuals were pooled to weigh to the nearest 0.1 μg with a fine scale (Sartorius Micro SC2) for biomass investment calculations, until 50 individuals from each were measured. Caste allocation was measured including biomass weight adapting the formula for sexual investment as $c = (\frac{q}{w})^k$, where q and w are the queen and worker dry weight and the power

conversion power is k=0.7 (Boomsma et al. 1995), as commonly used for differences in investment among queens and males. We calculated caste investment (I) as $I_i = c * q_i + w_i$ (Boomsma 1989) and explored differences in caste allocation for a range of k values between 0.6-1. This power of conversion might be species specific, and the exact value for *C. obscurior* is unknown but is likely to be within this range.

Statistical tests

To test for significant differences between treatments, we used generalized linear mixed effects models within the R package glmmTMB (R version 3.5.2, (Pinheiro et al. 2011)) and a negative binomial distribution for count data. If the count data and caste investment ratios were log transformed, a Gaussian family distribution was used. The dependent variable was analyzed as a function of the fixed effects: treatment (Number of workers as a factor), and random effects: stock nest and box of origin, box of set up, set up date. All models were also graphically checked for consistency and model diagnostics were performed using the DHARMA package (R version 0.3.3.0, (Hartig 2020)). To test for differences in head width size, we used the average of the head width measurements of the workers per time point (each month). Data is available as external database S1, and the R-script is filed under database S2.

References

Beekman M, Lingeman R, Kleijne FM, Sabelis MW. 1998. Optimal timing of the production of sexuals in bumblebee colonies. Entomol. Exp. Appl. 88:147–154.

Boomsma JJ. 1989. Sex-investment ratios in ants: has female bias been systematically overestimated? Am. Nat. 133:517–532.

Boomsma JJ, Gawne R. 2018. Superorganismality and caste differentiation as points of no return: how the major evolutionary transitions were lost in translation. Biol. Rev. 93:28–54.

Boomsma JJ, Keller L, Nielsen MG. 1995. A comparative analysis of sex ratio investment parameters in ants. Funct. Ecol. 9:743–753.

Bourke AFG. 2007. Kin selection and the evolutionary theory of aging. Annu. Rev. Ecol. Evol. Syst. 38:103–128.

Cohen AA. 2018. Aging across the tree of life: The importance of a comparative perspective for the use of animal models in aging. Biochim. Biophys. Acta - Mol. Basis Dis. 1864:2680–2689.

Corona M, Libbrecht R, Wheeler DE. 2016. Molecular mechanisms of phenotypic plasticity in social insects. Curr. Opin. Insect Sci. [Internet] 13:55–60. Available from: http://dx.doi.org/10.1016/j.cois.2015.12.003

Flatt T, Partridge L. 2018. Horizons in the evolution of aging. BMC Biol. 16:1–13.

Giehr J, Heinze J, Schrempf A. 2017. Group demography affects ant colony performance and individual speed of queen and worker aging. BMC Evol. Biol. 17:1–9.

Haldane J. 1941. New Paths in Genetics. London : George Alien & Unwin, Ltd.

Harrison MC, Jaimes Nino LM, Rodrigues MA, Ryll J, Flatt T, Oettler J, Bornberg-bauer E. 2021. Gene co-expression network reveals highly conserved, well-regulated anti-ageing mechanisms in old ant queens.

Hartig F. 2020. DHARMa : Residual Diagnostics for Hierarchical (Multi-Level / Mixed) Regression Models. Available from: https://cran.r-project.org/package=DHARMa

Heinze J, Schrempf A. 2012. Terminal investment: Individual reproduction of ant queens increases with age. PLoS One 7:1–4.

Hughes KA, Reynolds RM. 2005. Evolutionary and mechanistic theories of aging. Annu. Rev. Entomol. 50:421–445.

Jones OR, Scheuerlein A, Salguero-Gómez R, Camarda CG, Schaible R, Casper BB, Dahlgren JP, Ehrlén J, García MB, Menges ES, et al. 2014. Diversity of ageing across the tree of life. Nature 505:169–173.

Keller L, Genoud M. 1997. Extraordinary lifespans in ants: a test of evolutionary theories of ageing. Nature 389:3–5.

Kirkwood TBL. 1977. Evolution of ageing. Nature 270:301–303.

Kirkwood TBL, Austad SN. 2000. Why do we age? Nature 408:233–238.

Kramer BH, Schaible R. 2013. Colony size explains the lifespan differences between queens and

workers in eusocial Hymenoptera. Biol. J. Linn. Soc. 109:710–724.

Kramer BH, Schrempf A, Scheuerlein A, Heinze J. 2015. Ant colonies do not trade-off reproduction against maintenance. PLoS One 10:1–13.

López-Otín C, Blasco MA, Partridge L, Serrano M, Kroemer G. 2013. The hallmarks of aging. Cell 153:1194–1217.

Lucas E, Keller L. 2017. Explaining extraordinary lifespans : the proximate and ultimate causes of differential lifespan in social insects. In: Jones OR, Salguero-Gómez R, editors. The Evolution of Senescence in the tree of life. Shefferson. Cambridge: Cambridge University Press. p. 198–219.

Macevicz S, Oster G. 1976. Modeling social insect populations II: Optimal reproductive strategies in annual eusocial insect colonies. Behav. Ecol. Sociobiol. 1:265–282.

Maklakov AA, Chapman T. 2019. Evolution of ageing as a tangle of trade-offs: energy versus function. Proc. R. Soc. B Biol. Sci.:1–8.

Maklakov AA, Rowe L, Friberg U. 2015. Why organisms age: Evolution of senescence under positive pleiotropy? BioEssays 37:802–807.

Monroy Kuhn JM, Korb J. 2016. Editorial overview: Social insects: aging and the re-shaping of the fecundity/longevity trade-off with sociality. Curr. Opin. Insect Sci. 16:vii–x.

Oettler J, Schrempf A. 2016. Fitness and aging in *Cardiocondyla obscurior* ant queens. Curr. Opin. Insect Sci. 16:58–63.

Oster GF, Wilson EO. 1978. Caste and ecology in the social insects. Princeton, New Jersey, USA: Princeton University Press.

Pinheiro J, Bates D, Debroy S, Sarkar D. 2011. Linear and Nonlinear Mixed Effects Models.

Schrader L, Kim JW, Ence D, Zimin A, Klein A, Wyschetzki K, Weichselgartner T, Kemena C, Stökl J, Schultner E, et al. 2014. Transposable element islands facilitate adaptation to novel environments in an invasive species. Nat. Commun. 5:1–10.

Schrempf A, Cremer S, Heinze J. 2011. Social influence on age and reproduction: Reduced lifespan and fecundity in multi-queen ant colonies. J. Evol. Biol. 24:1455–1461.

Schrempf A, Giehr J, Röhrl R, Steigleder S, Heinze J. 2017. Royal Darwinian Demons: enforced changes in reproductive efforts do not affect the life expectancy of ant queens. Am. Nat. 189:436–442.

Schrempf A, Heinze J. 2006. Proximate mechanisms of male morph determination in the ant Cardiocondyla obscurior. Evol. Dev. 8:266–272.

Schrempf A, Heinze J, Cremer S. 2005. Sexual cooperation: mating increases longevity in ant queens. Curr. Biol. 15:267–270.

Toth AL, Sumner S, Jeanne RL. 2016. Patterns of longevity across a sociality gradient in vespid wasps. Curr. Opin. Insect Sci. [Internet] 16:28–35. Available from: http://dx.doi.org/10.1016/j.cois.2016.05.006

Williams GC. 1957. Pleiotropy, Natural Selection, and the Evolution of Senescence. Evolution (N. Y).:398–411.