1   Aug 31st, 2021

Intra-helical salt bridge contribution to membrane protein insertion

Gerard Duart [1#], John Lamb [2#], Juan Ortiz-Mateu [1], Arne Elofsson [2*] and Ismael

Mingarro [1*]

[1] Departament de Bioquímica i Biologia Molecular, Institut Universitari de

Biotecnologia i Biomedicina (BioTecMed), Universitat de València. E-46100

Burjassot, Spain.

[2] Science for Life Laboratory and Department of Biochemistry and biophysics,

Stockholm University, 171 21 Solna, Sweden

[#] Equal contribution

[*] Corresponding authors

ABSTRACT

Salt bridges between negatively (D, E) and positively charged (K, R, H) amino acids play an important role in protein stabilization. This has a more prevalent effect in membrane proteins where polar amino acids are exposed to a very hydrophobic environment. In transmembrane (TM) helices the presence of charged residues can hinder the insertion of the helices into the membrane. This can sometimes be avoided by TM region rearrangements after insertion, but it is also possible that the formation of salt bridges could decrease the cost of membrane integration. However, the presence of intra-helical salt bridges in TM domains and their effect on insertion has not been properly studied yet. In this work, we use an analytical pipeline to study the prevalence of charged pairs of amino acid residues in TM α-helices, which shows that potentially salt-bridge forming pairs are statistically over-represented. We then selected some candidates to experimentally determine the contribution of these electrostatic interactions to the translocon-assisted membrane insertion process. Using both *in vitro* and *in vivo* systems, we confirm the presence of intra-helical salt bridges in TM segments during biogenesis and determined that they contribute between 0.5-0.7 kcal/mol to the apparent free energy of membrane insertion (ΔG_{app}). Our observations suggest that salt bridge interactions can be stabilized during translocon-mediated insertion and thus could be relevant to consider for the future development of membrane protein prediction software.

KEYWORDS

electrostatic interactions; membrane insertion; salt bridge; translocon; transmembrane helix;

INTRODUCTION

Most integral membrane proteins have to insert their transmembrane (TM) segments into the lipid bilayer in a helical conformation and then acquire a defined three-dimensional structure by packaging their helices (Martínez-Gil et al., 2011). α-Helical TM segments are largely composed of apolar residues because of the hydrophobic nature of the membrane environment. Nevertheless, in some cases, it is necessary for the protein activity to include polar amino acids in a TM region in order to develop a functional or structural role (Baeza-Delgado et al., 2012). This fact is sometimes not contemplated in modern membrane topology prediction tools (Tsirigos et al., 2018; 2015), in which the presence of charged amino acids in a sequence automatically suppose a penalty increase in the predicted free energy (ΔG_{pred}) of insertion. The presence of polar amino acids in TM regions is more frequent than what would be expected (Bañó-Polo et al., 2012), especially when these are in pairs on the same face of an α-helix.

Salt bridges are electrostatic interactions between negatively (D, E) and positively charged (K, R, H) amino acids that play an important role in protein stabilization (Marqusee and Baldwin, 1987). Many studies have shown that pairs of charged residues that form potential salt-bridges stabilize soluble α-helices (Donald et al., 2011). Salt bridges play an important role in the folding of globular proteins and, despite their low occurrence in TM domains, it seems that the contribution in membrane protein stability could be even more determinant. This contribution is especially important in membrane protein biogenesis, as salt bridges help to bury the polarity of charged residues in a hydrophobic environment (Mbaye et al., 2019). Apart from that, it has been suggested that potential salt bridges could help in the

insertion of TM α-helices (Baeza-Delgado et al., 2016; Bañó-Polo et al., 2012), even

though their predicted ΔG penalty is well above what is usually seen for TM

segments.

To investigate the potential formation of intra-helical salt bridges in TM α-

helices, we analyzed the composition of the TM domains from membrane proteins

of known structures looking for preferences in the pairing of charged amino acids.

This analysis showed that charged residue pairing is more prevalent than expected

for pairs located on the same face of α-helices within membranes. Likely, salt bridge

formation on the same face of α-helices reduces the unfavorable energetics of

inserting ionizable residues into the hydrophobic membrane core (Chin and Heijne,

2000). We use this knowledge together with the known membrane protein

structures to generate a list of potential candidates for further *in vitro* and *in vivo*

experiments.

In this work, we have studied the presence of intra-helical salt bridges in TM

domains using *in silico*, *in vitro* and *in vivo* systems, showing that despite this being

an unexpected phenomenon in nature, salt bridges can be crucial for membrane

insertion. Also, we determined in a quantitative manner that the apparent free

energy (ΔG_{app}) of membrane insertion through the translocon machinery can be

decreased between 0.5-0.7 kcal/mol by position-specific charge pair interaction,

which is not contemplated in the commonly used ΔG predictors. These findings will

lead to a better understanding of the insertion mechanism of TM helices and to

improve prediction tools that would more accurately be able to model the presence

of charged residues in these helices.

RESULTS

Charge pair interactions in model transmembrane helices.

To test the contribution of potential salt bridges to the translocon-mediated membrane insertion, we used the vehicle protein leader peptidase (Lep) from *Escherichia coli* (Fig.1a). The Lep protein consists of two TM segments (H1 and H2) connected by a cytoplasmic loop (P1) and a large C-terminal (P2) domain, which inserts into endoplasmic reticulum (ER)-derived rough microsomes with both termini located in the microsomes lumen. The designed TM segments were inserted into the luminal P2 domain and flanked by two acceptor sites (G1 and G2) for N-linked glycosylation. Glycosylation occurs exclusively in the lumen of the ER (or microsomes) because of the location of the oligosaccharyltransferase (OST) active site (a translocon-associated enzyme responsible for the oligosaccharide transfer) (Braunger et al., 2018). In this case, the engineered glycosylation sites can be used as membrane insertion reporters because G1 will always be glycosylated due to its native luminal localization, but G2 will be glycosylated only upon translocation of the analyzed sequence across the microsomal membrane. A singly glycosylated construct in which a tested sequence is inserted into the membrane has a molecular mass ~2.5 kDa higher than the molecular mass of Lep molecule expressed in the absence of microsomes; the molecular mass shifts by ~5 kDa upon double glycosylation, which facilitates its identification by gel electrophoresis when expressed in the presence of [^{35}S-labeled] amino acids. Then, *in vitro* transcription/translation of these chimeric proteins in the presence of rough microsomal membranes (RMs) allows for accurate and quantitative description of membrane insertion of designed sequences (Bañó-Polo et al., 2019; Hessa et al.,

2005; 2007; Tamborero et al., 2011). The degree of membrane insertion is quantified

by analyzing the fractions of singly glycosylated (i.e., membrane inserted) and

doubly glycosylated (i.e., non-inserted) molecules, which can be expressed as an

experimental apparent free energy of membrane insertion, ΔG_{exp} (see Materials and

Methods) (Hessa et al., 2005).

We first compared the effects of oppositely charged Lys and Asp residues

on the insertion of a 19-residue-long hydrophobic stretch (L4/A15 scaffold, 4

leucines and 15 alanines), which was designed to insert stably into the RM

membranes (Hessa et al., 2005), including charges centered in the TM segment at

different positions (Fig. 1b) and "insulated" from the surrounding sequence by N-

and C-terminal GGPG- and -GPGG tetrapeptides. Single Lys and Asp residues

were placed in positions 8 and 12 respectively, and pairs of Lys-Asp residues were

designed to cover positions 7-12 (that is, more than one helical turn). When pairs of

charged residues are present, our results showed a tendency to insert more

efficiently when pair charges were placed in positions (*i*, *i*+1; *i*, *i*+3; *i*, *i*+4) that are

permissive with salt bridge formation (Fig. 1c), actually an effect not observed in the

predictions (Fig. 1b). Similar results were obtained on a different Leu/Ala

background with a slightly higher insertion efficiency (L5/A14, 5 leucines and 14

alanines), those mutants harbouring charged pairs compatible with salt bridge

formation (i.e. *i*, *i*+3; *i*, *i*+4) insert more efficiently than the non-compatible one *i*, *i*+5

(Figure S1).   Being the insertion of charged residues a thermodynamically

inconvenient phenomenon within the membrane environment, it is expected that

the sequence context and the amino acid composition of the TM helix would be

determinant for salt bridge formation. Accordingly, we scrutinized a large dataset of

membrane proteins of known three-dimensional structures in order to focus on

natural salt bridges present within TM segments.

Charged pairs in transmembrane helices.

Alpha helices are a common secondary structure in both globular and membrane

proteins. We created two main datasets, TM dataset with helical membrane

proteins of known structure from the PDBTM-dataset (Kozma et al., 2013), and

GLOB dataset with globular alpha helical proteins selected from the SCOP-

database (Andreeva et al., 2013; 2019), see methods for the full creation steps.

Table 1 shows a breakdown of alpha-helices, charged residues and salt bridges in

the two datasets. What is clear is that long alpha helices (≥ 17 residues) form a

larger proportion in the TM dataset than in the globular helices (GLOB) dataset,

which is logical as most TM helices need to span through the hydrophobic core of

the lipid bilayer of thickness ~30 Å.

Our TM dataset showed the same distribution of polar charges as previous

studies (Illergård et al., 2011), with about 10 % of polar residues in the core

membrane regions (see Table 1), and about one-third of these are charged. Over

half of the charged residues could form pairs with other charged residues at

intervals of i, $i+1$; i, $i+3$ and i, $i+4$. In contrast, the GLOB dataset had a much higher

proportion of charged polar residues. The GLOB dataset also contained about 15

times as many charged pairs relative to its size compared to the TM dataset, again

indicating that charged residues are more common in globular than in TM helices.

The two datasets, TM and GLOB, were extended with homologous

sequences identified by searching with jackhammer against UniProt. All sequences

with an E-value lower than 10^{-3} were included. These datasets are named TM-MSA

and GLOB-MSA. Using data in the TM-MSA dataset to produce the log odds ratios,

we identified periodicity patterns of charged residue pairs that are more common

than what would have been expected from the underlying amino acid composition

(see Figure 2). We found that polar residues at pairs i, $i+3$; i, $i+4$ and i, $i+7$ are

significantly enhanced (Fig. 2). This feature is strengthened when we examine the

same plot for the GLOB-MSA dataset, where these patterns were not observed

(Figure S2). This was also clear when statistical significance was taken into account,

see Figures S3 and S4.

Table 2 lists the log odds ratios together with errors and familywise error-

corrected p-values for all pairs at separation up to seven residues. It is clear that

pair residues placed at i, $i+1$; i, $i+3$ and i, $i+4$ positions are most significant,

especially in the case of oppositely charged pairs.

Structural analysis of charged residues in transmembrane helices.

The abundance of salt bridges also shows a remarkable difference between TM and

globular proteins. In the GLOB dataset over 13% of the proteins contains at least

one local salt bridge in an alpha helix whereas just over 5% of the membrane

proteins contains a local salt bridge, see Table 1. This does conform to our current

understanding of soluble versus TM alpha-helices and their different environments,

and suggests that salt bridges in TM regions perform a vital functional and/or

structural role.

As seen in Figure 2, charged pairs of amino acids are especially prevalent at

positions i, $i+1$; i, $i+3$ and i, $i+4$. Oppositely charged residues stand out, especially

Glu-Arg at *i*, *i*+1, Glu-Lys at *i*, *i*+3 and Asp-Lys at *i*, *i*+4. Also, several same charged

pairings at *i*, *i*+3 and *i*, *i*+6 are more frequent than expected. Other known structural

features can also be hinted at, including aromatic ring stacking by His-Trp pair

(Samanta et al., 1999) at *i*, *i*+6, and contacting with prosthetic groups by His-His

pair at *i*, *i*+7 (Illergård et al., 2011).

Charged pairs placed at *i*, *i*+1; *i*, *i*+3 and *i*, *i*+4 could potentially form salt

bridges as they are all on the same relative face of the alpha helix and are close

enough in vertical separation on the helix (see Figure 3, top). Although oppositely

charged pairs at *i*, *i*+7 are also on the same face of the alpha helix, unless the alpha

helix has a bend, both residues are too far separated to form a salt bridge. This was

clearly seen in Figure 3 where the TM dataset was used. In both absolute count and

log odds ratios (Fig. 3a) it is clear that residues at *i*, *i*+1; *i*, *i*+3 and *i*, *i*+4 are by far

the most common and overrepresented pairings. Figure 3a also shows that

oppositely and same charged pairs have about the same overrepresentation at *i*,

i+3, whereas oppositely charged pairs are stronger than same charged pairs at *i*,

i+1and *i*, *i*+4, both within salt bridge range, and same charged pairs are stronger at

positions *i*, *i*+7 and *i*, *i*+8, too far to form salt bridges.

When structure-observed salt bridges in the different positions were

compared to the oppositely charged pairs a clear image aroused, see Figure 3b.

Even though there are more oppositely charged pairs at position *i*, *i*+1 than in

positions *i*, *i*+3 ~~and *i*, *i*+4~~ with both log odds ratios over 1.0 (Fig. 3a), only about 15%

of the oppositely charged pairs at *i*, *i*+1 form salt bridge (Fig. 3b). This is in contrast

to *i*, *i*+3 where almost 40 % of the pairs form salt bridges, and just under 25% at *i*,

i+4 (Fig. 3b).

Selection of natural salt bridges from membrane protein structures.

To further look for candidate proteins containing membrane-spanning helices with salt bridges we started with the redundant (TM-Red) dataset of known structures (see Materials and Methods). Each of the 8,687 proteins in our dataset were scanned for oppositely charged pairs in positions *i, i*+1*; i, i*+3 and *i, i*+4 within any TM segments core region. For each of these, we only kept proteins with at least one TM segment that contains such a pair and where this pair was within salt bridging distance.

To select for potential candidates, we also look at the estimated ΔG_{pred} values and choose helices with a positive value above one, as per our hypothesis, a salt bridge helps TM insertion of sequences for which the hydrophobic force would not be enough to insert. This stricter definition results in a set of 426 candidates of a total of 431 salt bridges with a wide range of estimated ΔG_{pred} penalty values (Figure S5). As shown in Figure S5, most TM segments with a salt bridge exhibit a surprisingly high ΔG_{pred} value above 0 that in normal circumstances are not expected to insert into a membrane. Then, we selected TM7 (helix G) from halorhodopsin protein (PDB ID: 3QBG) with an estimated ΔG_{pred} value above +1.7 kcal/mol, and helix A from calcium ATPase (PDB ID: 1SU4) with a higher estimated ΔG_{pred} value (above +4.1 kcal/mol), as candidates for systematic studies to cover a wide range of insertion penalties. See the github repository (https://github.com/ElofssonLab/salt_bridges) for the full lists.

Intra-helical salt bridge stabilizes the insertion of helix G from halorhodopsin.

Halorhodopsin (hR) from *Natronomonas pharaonis* (3QBG) is a protein made up of

seven TM helices (helix A through G) and a retinal chromophore that is bound via a

protonated Schiff base to the ε-amino group of a lysine (K258) residue located

roughly in the middle of helix G (Kanada et al., 2011). *In silico* analysis of 3QBG

structure, the anion-free form of the protein, showed a charged pair of amino acids

(Asp-Lys), involving the functional K258 and the D254 in the center of helix G (Fig.

4). The position (*i*, *i*+4) and the distance between the anionic carboxylate ($RCOO^-$)

from the Asp residue and the cationic ammonium (RNH_3^+) from the lysine residue,

in the crystal structure, was about 3.5 Å, a permissive distance for a salt bridge

formation (Fig. 4g), which has been established as being lower than 4 Å (Kumar and

Nussinov, 2002). To get insights into this interaction, we designed three mutants

that were supposed to perturb the salt bridge interaction by different ways: K258D

mutant by placing two charged residues with the same polarity at positions *i*, *i*+4;

K258A mutant by replacing one of the charged residues by a non-polar amino acid,

and K258Y/Y259K double mutant by placing the charged pair at a non-permissive

salt bridge position (*i*, *i*+5; Fig. 4g), while keeping the same amino acid composition

(Fig. 4a).

Halorhodopsin is a trimeric protein in which the helix G is neither exposed at

the monomer-monomer interface nor oriented to the inner part of the trimeric

structure (Fig. 4, panels b-d). In fact, the charged pair found in helix G is oriented

toward the core of the 'globular' structure in each monomer. Then, the insertion of

helix G was studied using the Lep-based assay. When constructs harboring helix G

wild type sequence were translated *in vitro* in the presence of RMs singly-

glycosylated (reporting insertion) forms were found (Fig. 4e, lane 2), despite its

positive (suggesting non-insertion) ΔG_{pred} value (Fig. 4a). The nature of the higher

molecular weight polypeptide species was analysed by endoglycosidase H (EndoH)

treatment, a highly specific enzyme that cleaves N-linked oligosaccharides.

Treatment with EndoH of the samples eliminated higher molecular mass bands (Fig.

4E, lane 1), confirming the sugar source of the retarded electrophoretic mobility

bands and suggesting helix G insertion into the microsomal membrane. However,

locating the Asp-Lys pair at *i, i*+5, which is non-compatible with salt bridge

interaction (Fig. 4g), strongly reduced the experimental insertion efficiency (Fig. 4e,

lane 5). Interestingly, replacing the positively charged lysine residue by a negatively

charged aspartic acid residue (K258D) rendered similarly low levels of insertion

efficiency (Fig. 4e, lane 4). As expected, replacement of the ionizable lysine residue

by the aliphatic alanine increased the insertion efficiency (Fig. 4e, lane 3). In this

later mutant, the most likely cause for the increased insertion is the absence of the

positively charged lysine amino side chain and by the presence of the methyl side

chain group of the mutant alanine residue. The results of the Lep-based

glycosylation assay indicated that wild type (wt) and DA sequences (two stabilized

charges or only one charge, respectively) are inserted properly into the microsomal

membrane (ΔG_{exp} values -0.24 and -0.88 kcal/mol, respectively), but when the salt

bridge is disrupted, either by having two charged amino acids with the same polarity

(DD) or by placing oppositely charged residues at a non-permissive distance (*i, i*+5)

in the center of the helix, the translocation of the segment increases substantially.

It should be mentioned that the K258Y/Y259K double mutant has the same amino

acid composition than the original helix G, but insertion efficiency is remarkably

decreased (ΔG_{exp} = +0.31 kcal/mol). Together these results show that the interaction

(salt bridge) between Asp and Lys residues in the center of the helix G from 3QBG

is essential for its proper insertion into the microsomal membrane. The salt bridge

contributes approximately ~0,5 kcal/mol to the apparent experimental free energy

of microsomal membrane insertion, as this is the difference found between the

ΔG_{exp} values for the wt and *i, i+5* mutant.

Next, to ensure that the *in vitro* results are relevant to the *in vivo* situation, wt

and *i, i+5* constructs were also expressed *in vivo* in HEK-293T cells. To this end, a

c-myc tag was engineered at the C-terminus of the Lep chimera to allow immune-

detection of our constructs in cell extracts. As shown in Fig. 4f, transfected cells

with the chimera containing helix G wt sequence rendered singly glycosylated

molecules, indicating *in vivo* membrane insertion. In contrast, cells transfected with

the construct harboring *i, i+5* sequence rendered almost exclusively doubly

glycosylated forms, as proved by EndoH treatment (Fig. 4f, lane 2), suggesting

membrane translocation. These results emphasized the relevance of salt bridge

interactions in translocon-mediated TM insertion, especially in the *in vivo* (cellular)

environment.

Salt bridge contribution to the insertion of a heavily charged helix.

In order to challenge salt bridge interactions in a more hydrophilic TM helix (Figure

S5), we focus on helix A from the sarcoplasmic/ER calcium ATPase 1 (Toyoshima

et al., 2000). Calcium ATPase (PDB ID: 1SU4, *Oryctolagus cuniculus*) is a member

of the P-type ATPases that transport ions across the membrane against a

concentration gradient involving 10 α-helices (helix A through J) in the membrane-

embedded region (Toyoshima et al., 2000). *In silico* analysis of 1SU4 structure, the

crystal structure of the calcium ATPase with two bound calcium ions, showed Asp-

Arg (DR) pair (D59 and R63) in the center of the helix A. This helix extends beyond

the membrane (Fig. 5a, from Leu49 to Phe78) and shows some particularities. On

the one hand, in the structure the membrane-embedded stretch (the N-terminal

region of helix A) encompasses from Leu49 to Ala69 residues, and includes several

charged amino acids, probably involved in the Ca^{2+} transport across the membrane

(Glu51, Glu55, Glu58, Asp59 and Arg63). Therefore, the ΔG_{pred} value for this

segment (L49-A69) is remarkably higher (and positive, +4.12 kcal/mol) than

expected for a TM helix (Fig. 5b). On the other hand, the C-terminal region of this

helix contains a high prevalence of non-polar amino acids that is more compatible

with the hydrophobicity of the membrane core. Nevertheless, the presence of the

functional amino acids (e.g. Glu58) in the membrane-embedded region reinforce

the idea that the more hydrophilic N-terminal region must ultimately be embedded

within the membrane (Toyoshima et al., 2000). It has been previously shown that

the position in the membrane of TM helices in protein folded structures does not

always correspond to the thermodynamically favored positions in the membrane of

the isolated helices (Kauko et al., 2010). Instead, after translocon-mediated insertion

of the more hydrophobic region, repositioning of TM helices relative to the lipid

bilayer provides a convenient way for non-hydrophobic polypeptide segments to

become buried within the membrane. Then, the nature of helix A suggested the

possibility that initial insertion of the hydrophobic region can be followed by

subsequent repositioning of the charged region into the membrane hydrophobic

core, which will be the final segment embedded in the lipid bilayer (Fig. 5a).

Interestingly, the ΔG Predictor server (https://dgpred.cbr.su.se/index.php?p=home)

selected the adjacent (C-terminus, L60-F78) hydrophobic region as TM (Fig. 5b),

instead of the charged region found at the high-resolution structure (Uniprot code:

P04191).

Focusing on the potential salt bridge residues (D59 and R63 pair), the

distance between the anionic carboxylate ($RCOO^-$) from the D59 and the cationic

guanidinium ($RC(NH_2)_2^+$) from the R63 was about 3.0 Å in the crystal structure

(Figure 5a), clearly within the permissive range for salt bridge formation. To

investigate the contribution of this potential salt bridge interaction in the translocon-

mediated membrane insertion of this region, we worked with two different scaffold

sequences: the full helix A involving the residues 49-78 (Long, L); and a shorter

membrane-embedded version including residues 49-69 (Short, S) as found in the

solved structure. We also challenged the D59-R63 charge pair interaction in both

sequences by increasing the separation between the ionizable residues from the

native *i*, *i*+4 to non-permissive *i*, *i*+5, while maintaining amino acid composition

(R63I/I64R double mutant). *In vitro* transcription/translation of these sequences in

the presence of microsomes rendered singly glycosylated molecules for the

construct containing full-length helix A (Fig. 5c, lane 2). In contrast, when only the

membrane-embedded sequence was included, the Lep chimera was mainly doubly

glycosylated (Fig. 5c, lane 3), suggesting that in the full-length protein helix A inserts

initially through the more hydrophobic L60-F78 region and then, after protein

rearrangements, repositions the more hydrophilic L49-A69 region at the membrane

core, as found in the solved structure. Accordingly, the translocon inefficiently

inserted the isolated membrane-embedded (L49-A69) region (Fig. 5c, lane 3),

properly inserting helix A only when the full helical sequence is present (Fig. 5c, lane

2). When the charge paired residues were placed at a non-permissive distance in

terms of salt bridge interaction (i, $i+5$; Fig. 5e) the insertion efficiency was reduced

(Fig. 5c, lane 4), with a ΔG_{app} decrease (absolute values) of ~0,7 kcal/mol relative to

the wild type sequence (Fig. 5b). As expected, this effect was not observed when

the same mutations were grafted on the membrane-embedded (S) sequence (Fig.

5c, lane 5).

Next, we analysed the salt bridge interaction in HEK-293T cells to study the

translocon performance *in vivo*. When cell cultures were transfected with a Lep-

derived chimera containing the helix A wild type sequence only singly glycosylated

molecules were observed (Fig. 5d, lanes 1 and 2). However, a construct harboring

double mutant (R63I/I64R; i, $i+5$) sequence showed doubly glycosylated molecules

to a measurable extent (Fig. 5d, lanes 3 and 4), indicating a lower insertion efficiency

that can be attributed to the altered salt bridge interaction.

Effect of salt bridge formation in the absence of previous TM regions.

To investigate the effect of salt bridge formation in translocon-mediated membrane

insertion in the absence of precedent TM regions we used a different glycosylation-

based reporter system in which the TM sequences of interest (bR helix G and

ATPase helix A) were connected to the well-folded constant domain of the antibody

λ light chain, C_L (Feige and Hendershot, 2013). The C-terminal of the TM sequence

carries an Asn-Val-Thr glycosylation site (G2), and we engineered an extra

glycosylation site (G1) within the C_L sequence (Figure 6a, top). As in the case of the

Lep system, G1 will always be glycosylated due to the native translocation of the λ

light chain, but G2 will be glycosylated only upon translocation of the analyzed

(tested) sequence across the ER membrane (Figure 6a, bottom). When constructs

harboring either hR helix G or ATPase helix A wild type sequences were transfected

into Hek293T cells, singly glycosylated (reporting insertion) forms were

predominantly found (Fig. 6b, lane 1 and Fig. 6c, lane 2, respectively), discarding

any potential contribution to membrane insertion of precedent TM segments. On

the contrary, sequences containing non-permissive (i, i+5) salt-bridge forming pairs

were more efficiently doubly glycosylated (Figures 6b, lane 3 and 6c, lane 4,

respectively). Thus, similar to what was observed using the Lep system (Figures 4f

and 5d), the presence of chair pairs at permissive salt bridge formation distances

strongly promotes helix integration into the ER membrane.

DISCUSSION

Charged residues found in alpha-helices can both give a stabilizing effect

(Armstrong and Baldwin, 1993) as well as being important for interaction and

function such as in zinc finger motifs (Lin and Lin, 2018). Charged residues have

also been found to be more common in globular (soluble) helices compared to TM

helices (Kauko et al., 2008). This can be explained by the fact that globular helices

reside in a more hydrophilic environment compared to the hydrophobic

environment of the lipid bilayer. An intermediate between these is the existence of

amphipathic alpha-helix in contact with the surface of a bilayer, where one face

contains mainly polar residues facing an aqueous environment and the opposite

face with mostly nonpolar residues facing a hydrophobic environment (Giménez-

Andrés et al., 2018).

Whereas pairs of charged residues of the same charge facing the inside

pores in TM regions fill an essential functional role, pairs of oppositely charged

residues can form salt bridges that can stabilize the helix and play an important role

in function such as for transportation (Walther and Ulrich, 2014). Salt bridges might

also be important for hiding the charges during translocon-mediated TM helix

insertion, as the charged residues are hidden from the hydrophobic environment.

Previous work has shown that charged and polar residues are conserved within TM

segments (Illergård et al., 2011), indicating they are crucial for function and/or

stability.

Asp-Lys pairs at position i, $i+4$ and Glu-Lys pairs at position i, $i+3$ are the

most prevalent as seen previously in Figure 2. They are both among the most

prevalent oppositely charged pairs and the charged pairs that form the highest

number of salt bridges in membrane protein structures. This is in stark contrast to

Glu-Arg pair at position i, $i+1$ that although as frequent in pairs as Asp-Lys and Glu-

Lys at positions i, $i+4$ and i, $i+3$ respectively, only form salt bridges in one-fourth of

the cases as found in Fig. 3b.

An interesting observation is that positively charged Arg-Arg pairs at position

i, $i+3$ is numerically the most common pair at this position. Positively charged pairs

at i, $i+6$ are also more prevalent than expected and although Arg-Arg pairs

numerically only show up half as often at i, $i+6$ as in i, $i+3$, 80% of the Arg-Arg pairs

at position i, $i+6$ also contain an arginine at $i+3$, as found in helices from voltage-

gated ion channels that contain three or more periodically aligned Arg residues with

two intervening hydrophobic residues (Okamura et al., 2015).

The fraction of salt bridges at positions i, $i+5$ (Fig. 3b) is an anomaly due to

bend alpha-helices, as found in the helix E from bacterial translocon (PDB ID: 5MG3)

and in the helix A from a lipid flippase (PDB ID: 6CC4) due in both cases to the

presence of a glycine residue (Figure S6). Without the observed helix bending, these

salt bridges would not be formed.

By analyzing two native helices containing intra-helical salt bridges we now

find that the free energy of insertion (ΔG_{app}) is significantly reduced if both oppositely

charged residues are spaced at a permissive distance. These results indicate that

intra-helix salt bridge can form during translocon-assisted insertion or even earlier,

since in contrast to globular (soluble) helices, TM helices can be compacted inside

the ribosome exit tunnel (Bañó-Polo et al., 2018). The maximal reduction in ΔG_{app}

seen with Asp-Lys and Asp-Arg pairs in both hR and Ca^{2+} ATPase helices is 0.5-0.7

kcal/mol, which is in good agreement with the 0-1 kcal/mol estimated for these two

pairs from thermodynamic peptide partition into octanol experiments (Jayasinghe

et al., 2001). As found in the case of hR helix G (Figures 4f and 6b), this reduction

might be even higher in the cell context, since some auxiliary components of the

membrane insertion machinery (Chitwood and Hegde, 2020; Shurtleff et al., 2018;

Tamborero et al., 2011) can be in suboptimal conditions in the microsomal vesicles.

As mentioned above, in the case of hR helix G the lysine residue involved in

the salt bridge (K258) is bound to the retinal chromophore via a protonated Schiff

base as found in the crystal structure of a close homologue (Kolbe et al., 2000).

Then, the lysine residue plays a fundamental role for protein function but at the

same time introduces a penalty for membrane insertion. Interestingly, our data

suggest that helix G translocon-mediated insertion efficiency could be increased by

salt bridge formation between K258 and D254, and once in the membrane, retinol

binding to the apoprotein could occur by covalently binding the retinal as a

protonated Schiff base to K258 and perturbing D254 salt bridge interaction.

Beyond the conceptual issues involving the membrane insertion process, we

note that the availability of quantitative experimental data on the contribution of salt

bridge interactions to the free energy of insertion (ΔG_{app}) will make it possible to fine-

tune current membrane protein topology-prediction methods based on free energy

calculations. Although today's state of the art topology prediction tools uses

amphiphatic biologically derived scales, they do not take these types of salt bridge

interaction into account. Current algorithms tend to overestimate the free energy

of insertion due to the penalty acquainted by charged residues in the TM region.

However, distinguishing between charged residues of the same or opposite

polarity, i.e., incorporating the effect of potential salt bridges in the reduction of

ΔG_{app} during membrane integration should help to make prediction tools even more

accurate.

MATERIALS AND METHODS

Enzymes and chemicals. TNT T7 Quick for PCR DNA was from Promega

(Madison, WI, USA). Dog pancreas ER column washed rough microsomes were

from tRNA Probes (College Station, TX, USA). EasyTag™ EXPRESS^{35}S Protein

Labeling Mix, [^{35}S]-L-methionine and [^{35}S]-L-cysteine, for *in vitro* labeling was

purchased from Perkin Elmer (Waltham, MA, USA). Restriction enzymes were from

New England Biolabs (Massachusetts, USA) and endoglycosidase H was from

Roche Molecular Biochemicals (Basel, Switzerland). PCR and plasmid purification

kits were from Thermo Fisher Scientific (Ulm, Germany). All oligonucleotides were

purchased from Macrogen (Seoul, South Korea).

DNA Manipulation. The sequences of interest were introduced into the modified

Lep sequence from the pGEM1 plasmid (Hessa et al., 2005) between the *Spe*I and

*Kpn*I sites using two double-stranded oligonucleotides with overlapping overhangs

at the ends. The complementary oligonucleotides pairs were first annealed at 85 °C

for 10 min followed by gradual cooling to 30 °C and ligated into the vector (a kind

gift from G. von Heijne's lab). Mutations were obtained by site-directed mutagenesis

using the QuikChange kit (Stratagene, La Jolla, California). Lep system including the

sequences of interest in the P2 region were subcloned into *Kpn*I linearized pCAGGS

in-house version using In-Fusion HD cloning Kit (Takara) according to the

manufacturer's instructions. An engineered glycosylation site (Q36N) was added to

the C_L-TM plasmid (a kind gift from L. Hendershot's lab), in which the sequences

from hR helix G and ATPase helix A were introduced flanked by 'insulating' Gly-Pro

tetrapeptides. A *c*-myc tag (EQKLISEEDL) at the C-terminus of the Lep- and C_L-

derived sequences was added by PCR before cloning. For *in vitro* assays, DNA was

amplified by PCR adding the T7 promoter during the process. All sequences were

confirmed by sequencing the plasmid DNA at Macrogen Company (Seoul, South

Korea).

Translocon-mediated insertion into microsomal membranes. Lep constructs in

pGEM with L4/A15, L5/A14, 3QBG and 1SU4 segments and its variations were

transcribed and translated using the TNT T7 Quick Coupled System (#L1170,

Promega). Each reaction containing 1 µL of PCR product, 0.5 of EasyTag^{TM}

EXPRESS 35S Protein Labeling Mix (Perkin Elmer) (5.5 µCi) and 0.3 µL of

microsomes (tRNA Probes) was incubated at 30ºC for 90 min. Endo H treatment

was done following the manufacturer's instructions. Samples were analysed by

SDS-PAGE (12-14% polyacrylamide). The bands were quantified using a Fuji FLA-

3000 phosphoimager and the Image Reader 8.1j software. Free energy was

calculated using: ΔG_{app}=-RT lnK_{app}, where K_{app}=f_{2g}/f_{1g} being f_{1g} and f_{2g} the fraction of

singly glycosylated and double glycosylated protein, respectively.

Free apparent insertion energy, ΔG. The free insertion energy of a TM region, ΔG,

is calculated as per the experimentally defined Biological hydrophobicity scale

(Hessa et al., 2005). This scoring is amphiphilic with hydrophobic residues

contributing a lower (negative) ΔG while hydrophilic contributes a higher (positive)

ΔG. The total ΔG of a region is the sum of individual position specific scores. This

scoring can give an indication of how favorable the amino acid composition of a TM

region is to be inserted in a lipid bilayer membrane. To note is that hydrophobicity

alone is not the only driving force and that the positive inside rule (Heijne, 1989;

Lerch-Bader et al., 2008) and help from proceeding TM regions (Bañó-Polo et al.,

2013; Hedin et al., 2010) can assist insertion in polytopic TM proteins especially.

Core segment definition. We define core segments as a TM region minus the first and last 5 residues. This is to ignore the interface regions which are known to contain polar residues.

Salt bridge definition. Salt bridges are defined as per (Kumar et al., 2000), where a salt bridge is defined if a side chain carbonyl oxygen atom in Asp-Glu is within 4.0 Å from the nitrogen atom in Arg-Lys. This conforms to other works (Bosshard et al., 2004; Donald et al., 2011) with the definition that the atoms are within hydrogen bond distance. We also define local salt bridges as being bridges that are separated by at most 7 residues in the sequence. This is to separate long salt bridge interactions, which can occur between spatially close residues that are separated in sequence, such as coiled coils where salt bridges can be between separate alpha-helices.

Transmembrane helices dataset (TM dataset). The full pipeline is available as a Makefile together with supporting scripts in the github repository. The full PDBTM database (Kozma et al., 2013) was downloaded together with their list of non-redundant protein pdb ids. This list is used to generate both sequence and topology of the proteins by extracting both from the PDBTM-xml. For each protein in the non-redundant list, the membrane regions are extracted as per the PDBTM database annotation. All non-membrane regions are annotated 'i' for convenience. To support future analysis, membrane regions longer than 10 were run through DSSP and annotated with 'M' if all residues in the core segment were defined as alpha-helix ('H'), otherwise, the full membrane region is annotated 'm'. Observe that this creates fasta-like 3line files, that only contain topological annotation with ambiguous TM regions annotated as 'm' instead of the normal 'M'. These proteins are then cluster

using cd-hit (Fu et al., 2012; Li and Godzik, 2006) at 40% identity using the

parameter -c 0.4 -n 2 -T 0 -M 0 -d 0.

During the extraction of TM regions, the corresponding structure file from

RCSB was used to calculate all salt bridges within the current protein and any salt

bridge that has at least one residue within any TM region was saved. Additionally,

all salt bridges whose both residues were within the same segment and within 7

residues of each other were annotated as local as per the salt bridge definition

above.

Extraction of charged residues. From all annotated TM regions of length 17 or

longer (Baeza-Delgado et al., 2012), the core segment was extracted. All these core

regions were then scanned and when a charged residue was encountered, we

recorded any other charged residue from 7 residues before the current one to 7

residues after. This results in charged residues that can contain a charged pairing

partner outside of the TM segment and will therefore differ slightly from charged

pairs which are defined next.

Extraction of charged pairs. From all annotated TM regions of length 17 or longer

(Baeza-Delgado et al., 2012), the core segment was extracted. All these core

regions were then scanned and when a charged residue was encountered, we look

at up to 7 residues in front of it or to the end of that core region, whichever came

first. All occurrences of charged pairs were recorded, resulting in charged pairs

where both residues were fully within the core segment of a TM helix.

MSA-dataset extension (TM-MSA). Using the TM dataset, we extended it by

creating an MSA alignment of each protein using jackhmmer (Eddy, 2011) against

Uniref90 with one iteration and an E-value cut off of 10^{-3} with the following

parameters:

-N 1 -E 1e-3 --incE 1e-3 --cpu 14

From each alignment, we then sampled up to 200 hits, including the initial

seed sequence. If there were fewer than 200 hits, we used them all. We then used

the original topology for each alignment to extract all TM regions, only to include

parts where the sequence covers the full TM region and where the sequence did

not contain any insertions or deletions.

Dataset of helices from globular proteins (GLOB and GLOB-MSA). To create a

reference dataset of globular α-helical proteins we extracted all globular all-alpha

protein domains from SCOP. As SCOP classifies domains of proteins resulting in

that one domain of a protein can be annotated as globular whereas another domain

as TM (see 1PPJ chain D as an example) we reduced the SCOP list against the

redundant list of all PDBTM (Kozma et al., 2013) chains to clear any overlap. This

results in 4,500 proteins in total.

Topological files with sequence and membrane topology are created with the

help of the RCSB secondary structure file and only membrane segments whose

core region (*i.e.*, central 15 residues) is annotated as pure (canonical) α-helices were

retained, *i.e.*, those TM segments containing any residues within the core annotated

as loops or other types of secondary structures were removed. This file was further

homology reduced and alignments prepared in the same manner and using the

same parameters as the TM dataset described above.

The GLOB-MSA dataset was created in the same way as the TM-MSA

dataset using jackhmmer to extend the sequences to alignments and then to extract

helix sequences.

Redundant (TM-Red) dataset. The full PDBTM database was used as in the

preparation of the TM dataset. We skip the clustering step and instead use all

redundant proteins to generate their respective topology files. We added in the

constraint that each selected TM region must fully contain at least one potential salt

bridge. This means a local salt bridge where both residues are within the core

segment. This dataset was only used to find potential candidates for further *in vitro*

and *in vivo* experiments. See section of natural salt bridges above.

Calculation of log odds ratio. The log odds ratios for each amino acid pair for

steps 1 through 10 are calculated as follows:

logOddsRatio = log((A/B)/(C/D))

Where for two amino acids p_1 and p_2:

A = number of pairs of p_1 to p_2

B = number of total pairs

C = number of p_1 times number of p_2

D = number of total pairs squared

The standard error, SE, and z-value is calculated as follows allowing for a two-sided

test:

$$SE = \sqrt{(1/A + 1/B + 1/C + 1/D)}$$

z = abs(logOddsValue/SE)

The survival function, sf, from the scipy python packages is used to calculate the

p-values. To correct for multiple hypothesis, the Bonferroni Correction is used

based on the number of hypothesis, 20 * 20 * 10, number of amino acids square

times the number of steps.

Expression in mammalian cells. Lep or C$_L$-derived constructs containing 3QBG

or 1SU4 segments and its variations were tagged with *c*-myc epitope at their Ct

(EQKLISEEDL) and inserted in the appropriate plasmids. Once the sequence was

verified, plasmids were transfected into HEK293-T cells using Lipofectamine 2000

(Life Technologies) according to the manufacturer's protocol. Approximately 24 h

post-transfection cells were harvested and washed with PBS buffer. After a short

centrifugation (1000 rpm for 5 min on a table-top centrifuge) cells were lysed by

adding 100 μL of lysis buffer (30 mM Tris-HCl, 150 mM NaCl, 0.5% Nonidet P-40)

were sonicated in an ice bath in a bioruptor (Diagenode) during 10min and

centrifuged. After protein quantification, equal amounts of protein were submitted

to Endo H treatment or mock-treated followed by SDS-PAGE analysis and

transferred into a PVDF transfer membrane (ThermoFisher Scientific) as previously

described (Duart et al., 2020). Protein glycosylation status was analysed by Western

Blot using an anti-*c*-myc antibody (Sigma), anti-rabbit IgG-peroxidase conjugated

(Sigma), and with ECL developing reagent (GE Healthcare). Chemiluminescence

was visualized using an ImageQuantTM LAS 4000mini Biomolecular Imager (GE

Healthcare).

Acknowledgments

We thank Pilar Selvi and Beatriz Iborra for excellent technical and administrative

assistance, respectively. The C$_L$-TM plasmid was a kind gift from Prof. Linda M.

Hendershot (St. Jude Children's Research Hospital). This work was supported by

grants PID2020-119111GB-100 from the Spanish Ministry of Science and

Innovation and PROMETEU/2019/065 from Generalitat Valenciana (to I.M.), and by

grant from the Swedish Research Council (VR-NT 2016-03798 to A.E.). G.D. was

recipient of a predoctoral contract (FPU18/05771) from the Spanish Ministry of

Education.

Conflict of Interest Statement

None declared.

Table 1. About 10 % of the core residues in the TM dataset are polar and just under 4% are charged residues. This is significantly less than 37.5% and 25.5% respectively in the globular dataset. In the transmembrane dataset, just over half of the charged core residues form a charged pair at distance 1, 3 or 4. *In the case of the globular set, there are more potential charged pairs than charged core residues as multiple residues are counted more than once as they form more than one potential pairing with separation of 1, 3 or 4 steps. It is clearly seen that although the TM dataset contains more helices per protein and has a higher proportion of long (≥ 17 residues) helices it contains significantly fewer charged pairs.

Dataset	TM	TM-MSA	GLOB	GLOB-MSA
Charge statistics				
Core residues	40116	3990464	74299	7388277
Polar core residues	4080 (10.2%)	490861 (12.3%)	27851 (37.5%)	2696564 (36.5%)
Charged core residues	1467 (3.7%)	204544 (5.1%)	18959 (25.5%)	1819047 (24.6%)
Pairs of charged core residues (+1, +3 or +4)	825	88380	23755*	2271405*
# of same charged pairs	373	45220	10835	1117270
# of oppositely charged pairs	452	43160	12920	1154135
Charge Pairs statistics				
Proteins	925	130934	2475	363610
Total alpha helices	5435	481729	23991	2118158
Alpha helices ≥17	3755	306632	5895	485797
Proteins with charged pairs	204	25187	1780	193534
Helices with charged pairs	265	34029	4251	344142

Proteins with any salt bridge	241	-	*751*	-
Proteins with local salt bridge in helix	56	-	*335*	-
Total number of local salt bridges in helices	59	-	*577*	-

Table 2. Log odds ratios for all charged pairs, same charged pairs and oppositely
charged pairs with calculated errors and multiple hypotheses corrected p-values. It
is clear that charged pairs occur more often than predicted, evidenced by the
positive log odds ratios in all cases. It is clear that positions +1, +3 and +4 are the
most prevalent pairings, with the bolded values highlighting log odds ratio above
0.8. It is also clear that oppositely charged residues which have the potential to form
salt bridges are prevalent in all three positions whereas the same charge is mainly
prevalent in position 3. Most likely these same charges facing the same face of the
helix are involved in functions such as ion transport.

Spacing	All Log odds			Same charged Log odds			Oppositely charged Log odds		
	ratio	error	p-value	ratio	error	p-value	ratio	error	p-value
+1	**0.879**	0.025	$1.31e^{-257}$	0.548	0.042	$4.03e^{-41}$	**1.107**	0.032	$1.65e^{-256}$
+2	0.252	0.037	$2.02e^{-08}$	0.185	0.054	$1.88e^{-00}$	0.316	0.050	$1.09e^{-06}$
+3	**1.073**	0.026	$<0.00e^{-300}$	**1.118**	0.035	$1.15e^{-214}$	**1.026**	0.037	$5.78e^{-165}$
+4	**0.965**	0.028	$1.14e^{-253}$	0.696	0.046	$2.88e^{-49}$	**1.177**	0.036	$2.74e^{-233}$
+5	0.517	0.037	$4.88e^{-42}$	0.390	0.055	$4.15e^{-09}$	0.630	0.049	$1.21e^{-34}$
+6	0.540	0.038	$1.66e^{-43}$	0.545	0.053	$2.33e^{-21}$	0.536	0.053	$2.10e^{-20}$
+7	0.615	0.038	$1.21e^{-55}$	0.764	0.050	$7.78e^{-50}$	0.439	0.058	$1.97e^{-10}$

 FIGURE LEGENDS

Figure 1. Effects on membrane insertion of single or pairs of Asp and Lys

residues in a model TM segment. (a) Schematic representation of the leader

peptidase (Lep) model protein. G1 and G2 denote artificial glycosylation acceptor

sites. The sequence under investigation was introduced in the P2 region after H2.

Recognition of the tested sequence as a TM by the translocon machinery

(highlighted in green) results in the modification of the G1 site but not G2. The Lep

chimera will be double glycosylated if the sequence being tested is not recognized

as a TMD and thus translocated into the microsomes lumen (shown in red). **(b)** The

tested sequences from L4/A15 model TM (including the charged residues, bold),

the gap distance, and the predicted ΔG (ΔG_{pred}) values in kcal/mol are shown. Amino

acids with positive and negative charge are highlighted in blue (K) and red (D)

respectively. **(c)** Experimental ΔG (ΔG_{exp}) in kcal/mol of each tested sequence in the

Lep-based microsomal assay. The mean and standard deviation of at least 3

independent experiments is represented (*n* values: 4 [from 1 to 7] and 3 [8 and 9]).

The individual value of each experiment is represented by a solid dot, *p*-values

(ordinary one-way ANOVA test with Dunnett correction) are indicated above the

corresponding bars with values <0.005 highlighted in green. In addition, a green

square represents the experimental ΔG value for the L4/A15 sequence from an

earlier study [12]. The wt and single mutants are shown in white bars. Charges at

compatible distances with salt bridge formation (*i, i*+1; *i, i*+3; and *i, i*+4) are shown

in brown, orange and yellow, respectively. Not compatible distances with salt bridge

formation (*i, i*+2; and *i, i*+5) are shown in dark gray. The inset shows a representative

SDS-PAGE gel for L4/A15 construct. The construct was expressed in rabbit

reticulocyte lysed in the presence (+RM) or absence (–RM) of rough microsomes.

Bands of non-glycosylated proteins are indicated by a white dot; mono and double

glycosylated proteins are indicated by one and two black dots, respectively.

Figure 2. Log odds ratios of each pair of amino acids for '*i, i*+1' through '*i, i*+8'

for the TM-MSA dataset. The rows on the y-axis indicate the first amino acid in

the pair and the columns on the x-axis the second. The residues are ordered by

hydrophobicity according to the Engelman order (Engelman et al., 1986). See figure

S1 for the equivalent of the globular dataset. S2 and S3 show the same plots

masked for statistical significance.

Figure 3. Charge pairs in TM helical sequences and structures. Helical wheel

projection and lateral views of an α-helix are shown on top. The initial position *i* and

the following 8 residues are numbered. Residues in positions *i*+3 (orange), *i*+4

(yellow), and *i*+7 (light brown) are mainly on the same face of the helix, but *i*+7 is

placed too far for a salt bridge interaction. **a)** The log odds ratios of charged pairs

for '*i, i*+1' through '*i, i*+8' in the TM dataset. Plain filled bars refer to oppositely

charged pairs and the forward slash are the and same charged pairs, all with error

bars. **b)** Fraction of oppositely charged pairs that form local salt bridges. The small

bump at *i* +5 are the two proteins 6CC4 (helix A) and 5MG3 (helix E), which both

exhibit a bend in the alpha helix due to the presence of glycine residues (see Fig.

S6).

Figure 4. Insertion of halorhodopsin helix G from *Natronomonas pharaonis*

(3QBG) into microsomal and cellular membranes. (a) Tested sequences from

3QBG including the gap distance, and the predicted (ΔG_{pred}) and experimental (*in*

vitro $\Delta G_{exp}^{in\ vitro}$ and in vivo $\Delta G_{exp}^{in\ vivo}$, respectively) ΔG values in kcal/mol are shown.

Amino acids with a positive or negative charge are highlighted in blue (K) and red

(D), respectively. Green numbers indicate negative ΔG (insertion) values, while red

numbers denote ΔG values above 0 (translocation). **(b)** Frontal view of 3QBG

monomer structure. The helix G is highlighted in orange with the D254 and K258

shown in sticks colored red and blue respectively. The membrane position is

indicated by a red (outer) and blue (inner) discontinuous line, according to OPM

dataset [22]. Lateral **(c)** and upper **(d)** views of the 3QBG trimeric structure. The

helix G is highlighted in orange with the D and K shown in sticks colored red and

blue, respectively. The different monomers are shown in transparent blue, pink and

green. Representative examples (*n*=3) of *in vitro* protein translations in the presence

of ER-derived microsomes **(e)** and Western blots (*n*=3) of *in vivo* protein translations

in HEK-293T cells **(f)** in the presence (+) or absence (-) of Endoglycosidase H

(EndoH), a glycan-removing enzyme. The absence of glycosylation of G1 and G2

acceptor sites is indicated by two white dots, single glycosylation by one white and

one black dot, and double glycosylation by two black dots. **(g)** Zoom view centered

on the salt bridge between D254 and K257 at *i*, *i*+4 (left) and *i*, *i*+5 (right) gaps. D

and K residues are shown in sticks colored red and blue, respectively, while the

dashed line indicates the $RCOO^-$ to RNH_3^+ distance.

Figure 5. Insertion of Calcium ATPase (1SU4) helix A into microsomal and

cellular membranes. (a) Lateral view of 1SU4 structure. Zoom view of the A helix

(right panel). The membrane-embedded region of helix A is highlighted in yellow.

Charged amino acids are shown as sticks in blue (R), red (D) and pink (E),

respectively. L49, A69 and F78 are also shown as sticks to define helix's

subdomains. The membrane location is indicated by red (outer) and blue (inner)

discontinuous lines according to OPM dataset [22] and the distance between the R

and D charges is indicated in Å. **(b)** Helix A-derived sequences from 1SU4 including

the gap between charged residues, and the predicted (ΔG_{pred}) and experimental (*in*

vitro $\Delta G_{exp}^{in\ vitro}$ and *in vivo* $\Delta G_{exp}^{in\ vivo}$, respectively) ΔG values in kcal/mol are shown.

Amino acids with a positive charge are highlighted in blue (K) while negatively

charged are marked in red (D) and pink (E). The residues predicted as TM by the ΔG

Prediction server are underlined. Green numbers indicate negative ΔG (insertion)

values while red numbers denote ΔG values above 0 (translocation). Representative

examples (n=3) of *in vitro* protein translations in the presence of ER-derived

microsomes **(c)** and Western blots (n=3) of *in vivo* protein translation in HEK-293T

cells **(d)** in the presence (+) or absence (-) of Endoglycosidase H (EndoH), a glycan-

removing enzyme. The absence of glycosylation of G1 and G2 acceptor sites is

indicated by two white dots, single glycosylation by one white and one black dot,

and double glycosylation by two black dots. **(e)** 1SU4 helix A *i*, *i*+5 mutant. The

membrane-embedded region of helix A is highlighted in yellow. Charged residues

are shown as sticks in blue (R), red (D) and pink (E). L49, A69 and F78 are also

shown as sticks to define helix's subdomains. The membrane is indicated by red

(outer) and blue (inner) discontinuous lines as in (a), and the dashed line indicates

the $RCOO^-$ and $RC(NH_2)_2^+$ distance.

Figure 6. Salt bridge effect on the insertion in the absence of preceding TM segments. (a) (Top) Schematic of the C_LTM construct used. It is composed of the domain of the antibody λ light chain (C_L) containing a glycosylation site (G1) connected by flexible linkers to the sequence of analysis followed by a C-terminal glycosylation site (G2) and a *c*-myc tag. (Bottom) Scheme depicting the main features of the C_LTM insertion assay. Black dots represent glycosylated sites while white dots represent non-glycosylated sites. **(b)** Representative halorhodopsin (3QBG) helix G western blot (*n*=3) of *in vivo* protein translations in HEK-293T cells in the presence (+) or absence (-) of Endoglycosidase H (EndoH), a glycan-removing enzyme. **(c)** Representative Ca^{2+} ATPase (1SU4) helix A western blot (*n*=3) of *in vivo* protein translations in HEK-293T cells in the presence (+) or absence (-) of EndoH. Non-glycosylated proteins are indicated by two white dots, singly-glycosylated proteins are indicated by one white and one black dot, and doubly-glycosylated proteins are indicated by two black dots. Experimental ΔG values (kcal/mol) are shown above each sample (*n*=3).

Figure S1. Effects on membrane insertion of single or pairs of Asp and Lys residues in L5/A14. a) The tested sequences from L5/A14 model TM (including the charged residues, bold), the gap distance, and the predicted ΔG ($ΔG_{pred}$) and experimental ($ΔG_{exp}$) values in kcal/mol are shown. Amino acids with positive and negative charge are highlighted in blue (K) and red (D) respectively. **b)** Experimental ΔG ($ΔG_{exp}$) in kcal/mol of each tested sequence in the Lep-based microsomal assay. The mean and standard deviation of 3 independent experiments are represented.

The individual value of each experiment is represented by a solid dot, *p*-values are

indicated above. In addition, a green dot represents the ΔG_{pred} value for the L5/A14

sequence. The wt and single mutants are shown in white bars. Charges at

compatible distances with salt bridge formation (*i*, *i*+3; and *i*, *i*+4) are shown in

orange and yellow, respectively. Not compatible distances with salt bridge

formation (*i*, *i*+5) is shown in gray. The inset shows a representative SDS-PAGE gel

for L4/A15 and L5/A14 constructs. The construct was expressed in rabbit

reticulocyte lysed in the presence (+RM) or absence (–RM) of column washed rough

microsomes. Bands of non-glycosylated proteins are indicated by a white dot;

mono and double glycosylated proteins are indicated by one and two black dots,

respectively.

Figure S2. Log odds ratios of each pair of amino acids for '*i*, *i*+1' through '*i*, *i*+8'

for the GLOB-MSA dataset. Log odds ratios for the middle core residues of α-

helices of at least 17 residues in length in the GLOB-MSA dataset. The rows on the

y-axis indicate the first amino acid in the pair and the columns on the x-axis the

second. The residues are ordered as in Fig. 2.

Figure S3. Log odds ratios of each pair of amino acids for '*i*, *i*+1' through '*i*, *i*+8'

for the GLOB-MSA dataset. Log odds ratios as in S2 but all pairs with P-

value >0.05 have been masked.

Figure S4. Masked log odds ratios of each pair of amino acids for '*i*, *i*+1'

through '*i*, *i*+8' for the TM-MSA dataset. Log odds ratios as in Fig. 2 but all pairs

with P-value >0.05 have been masked.

Figure S5. Histogram of ΔG_{pred} values for the salt bridge containing TM segments.

ΔG values represented with a bin size of 0.8 kcal/mol, with negative (indicative of

insertion) and positive (indicative of non-inserted) values highlighted with a green

and red background, respectively. Halorhodopsin 3QBG helix G (ΔG_{pred} of 1.73

kcal/mol) and Ca2+ ATPase helix A (ΔG_{pred} of 4.15 kcal/mol) are part of the left and

right blue highlighted bars, respectively.

Figure S6. Salt bridges at pair *i*, *i*+5. Left: helix E from bacterial translocon (PDB

ID: 5MG3). Glycine (orange) residue causes a kink facilitating salt bridge interaction

between Asp50 (red) and Arg55 (blue). Right: helix A from a lipid flippase (PDB ID:

6CC4). Glycine (orange) residue causes a kink facilitating salt bridge interaction

between Arg153 (blue) and Glu158 (red). Distances are shown in Ångström.

REFERENCES

Andreeva, A., Howorth, D., Chothia, C., Kulesha, E., and Murzin, A.G. (2013).
SCOP2 prototype: a new approach to protein structure mining. Nucleic Acids Res.
42, D310–D314.

Andreeva, A., Kulesha, E., Gough, J., and Murzin, A.G. (2019). The SCOP
database in 2020: expanded classification of representative family and
superfamily domains of known protein structures. Nucleic Acids Res. *48*, D376–
D382.

Armstrong, K.M., and Baldwin, R.L. (1993). Charged histidine affects alpha-helix
stability at all positions in the helix by interacting with the backbone charges.
Proc. Natl. Acad. Sci. U.S.a. *90*, 11337–11340.

Baeza-Delgado, C., Heijne, von, G., Marti-Renom, M.A., and Mingarro, I. (2016).
Biological insertion ofcomputationally designed shorttransmembrane segments.
Sci. Rep. 1–9.

Baeza-Delgado, C., Marti-Renom, M.A., and Mingarro, I. (2012). Structure-based
statistical analysis of transmembrane helices. Eur Biophys J *42*, 199–207.

Bañó-Polo, M., Baeza-Delgado, C., Orzáez, M., Marti-Renom, M.A., Abad, C., and
Mingarro, I. (2012). Polar/Ionizable Residues in Transmembrane Segments: Effects
on Helix-Helix Packing. PLoS ONE *7*, e44263.

Bañó-Polo, M., Baeza-Delgado, C., Tamborero, S., Hazel, A., Grau, B., Nilsson, I.,
Whitley, P., Gumbart, J.C., Heijne, von, G., and Mingarro, I. (2018).
Transmembrane but not soluble helices fold inside the ribosome tunnel. Nat
Commun *9*, 5246.

Bañó-Polo, M., Martínez-Gil, L., Barrera, F.N., and Mingarro, I. (2019). Insertion of
Bacteriorhodopsin Helix C Variants into Biological Membranes. ACS Omega *5*,
556–560.

Bañó-Polo, M., Martínez-Gil, L., Wallner, B., Nieva, J.L., Elofsson, A., and
Mingarro, I. (2013). Charge Pair Interactions in Transmembrane Helices and Turn
Propensity of the Connecting Sequence Promote Helical Hairpin Insertion. J Mol
Biol *425*, 830–840.

Bosshard, H.R., Marti, D.N., and Jelesarov, I. (2004). Protein stabilization by salt
bridges: concepts, experimental approaches and clarification of some
misunderstandings. J. Mol. Recognit. *17*, 1–16.

Braunger, K., Pfeffer, S., Shrimal, S., Gilmore, R., Berninghausen, O., Mandon,
E.C., Becker, T., Förster, F., and Beckmann, R. (2018). Structural basis for
coupling protein transport and N-glycosylation at the mammalian endoplasmic
reticulum. Science *360*, 215–219.

Chin, C.N., and Heijne, von, G. (2000). Charge pair interactions in a model
transmembrane helix in the ER membrane. J Mol Biol *303*, 1–5.

Chitwood, P.J., and Hegde, R.S. (2020). An intramembrane chaperone complex
facilitates membrane protein biogenesis. Nature *584*, 1–22.

Donald, J.E., Kulp, D.W., and DeGrado, W.F. (2011). Salt bridges: Geometrically
specific, designable interactions. Proteins *79*, 898–915.

Duart, G., García-Murria, M.J., Grau, B., Acosta-Cáceres, J.M., Martínez-Gil, L.,
and Mingarro, I. (2020). SARS-CoV-2 envelope protein topology in eukaryotic
membranes. Open Biol. *10*, 200209–6.

Eddy, S.R. (2011). Accelerated Profile HMM Searches. PLoS Comput Biol *7*,
e1002195.

Engelman, D.M., Steitz, T.A., and Goldman, A. (1986). Identifying nonpolar
transbilayer helices in amino acid sequences of membrane proteins. Annu Rev
Biophys Biophys Chem *15*, 321–353.

Feige, M.J., and Hendershot, L.M. (2013). Quality Control of Integral Membrane
Proteins by Assembly-Dependent Membrane Integration. Mol Cell *51*, 297–309.

Fu, L., Niu, B., Zhu, Z., Wu, S., and Li, W. (2012). CD-HIT: accelerated for
clustering the next-generation sequencing data. Bioinformatics *28*, 3150–3152.

Giménez-Andrés, M., Čopič, A., and Antonny, B. (2018). The Many Faces of
Amphipathic Helices. Biomolecules *8*, 45–14.

Hedin, L.E., Ojemalm, K., Bernsel, A., Hennerdal, A., Illergård, K., Enquist, K.,
Kauko, A., Cristobal, S., Heijne, von, G., Lerch-Bader, M., et al. (2010). Membrane
insertion of marginally hydrophobic transmembrane helices depends on sequence
context. J Mol Biol *396*, 221–229.

Heijne, von, G. (1989). Control of topology and mode of assembly of a polytopic
membrane protein by positively charged residues. Nature *341*, 456–458.

Hessa, T., Kim, H., Bihlmaier, K., Lundin, C., Boekel, J., Andersson, H., Nilsson, I.,
White, S.H., and Heijne, von, G. (2005). Recognition of transmembrane helices by
the endoplasmic reticulum translocon. Nature *433*, 377–381.

Hessa, T., Meindl-Beinker, N.M., Bernsel, A., Kim, H., Sato, Y., Lerch-Bader, M.,
Nilsson, I., White, S.H., and Heijne, von, G. (2007). Molecular code for
transmembrane-helix recognition by the Sec61 translocon. Nature *450*, 1026–
1030.

Illergård, K., Kauko, A., and Elofsson, A. (2011). Why are polar residues within the
membrane core evolutionary conserved? Proteins *79*, 79–91.

Jayasinghe, S., Hristova, K., and White, S.H. (2001). Energetics, stability, and
prediction of transmembrane helices. J Mol Biol *312*, 927–934.

Kanada, S., Takeguchi, Y., Murakami, M., Ihara, K., and Kouyama, T. (2011).
Crystal Structures of an O-Like Blue Form and an Anion-Free Yellow Form of
pharaonis Halorhodopsin. J Mol Biol *413*, 162–176.

Kauko, A., Hedin, L.E., Thebaud, E., Cristobal, S., Elofsson, A., and Heijne, von, G.
(2010). Repositioning of transmembrane alpha-helices during membrane protein
folding. J Mol Biol *397*, 190–201.

Kauko, A., Illergård, K., and Elofsson, A. (2008). Coils in the Membrane Core Are
Conserved and Functionally Important. J Mol Biol *380*, 170–180.

Kolbe, M., Besir, H., Essen, L.O., and Oesterhelt, D. (2000). Structure of the light-
driven chloride pump halorhodopsin at 1.8 A resolution. Science *288*, 1390–1396.

Kozma, D., Simon, I., and Tusnády, G.E. (2013). PDBTM: Protein Data Bank of
transmembrane proteins after 8 years. Nucleic Acids Res. *41*, D524–D529.

Kumar, S., Ma, B., Tsai, C.J., and Nussinov, R. (2000). Electrostatic strengths of
salt bridges in thermophilic and mesophilic glutamate dehydrogenase monomers.
Proteins *38*, 368–383.

Kumar, S., and Nussinov, R. (2002). Relationship between Ion Pair Geometries
and Electrostatic Strengths in Proteins. Biophys J *83*, 1595–1612.

Lerch-Bader, M., Lundin, C., Kim, H., Nilsson, I., and Heijne, von, G. (2008).
Contribution of positively charged flanking residues to the insertion of
transmembrane helices into the endoplasmic reticulum. *105*, 4127–4132.

Li, W., and Godzik, A. (2006). Cd-hit: a fast program for clustering and comparing
large sets of protein or nucleotide sequences.

Lin, C.-Y., and Lin, L.-Y. (2018). The conserved basic residues and the charged
amino acid residues at the α-helix of the zinc finger motif regulate the nuclear
transport activity of triple C2H2 zinc finger proteins. PLoS ONE *13*, e0191971–20.

Marqusee, S., and Baldwin, R.L. (1987). Helix stabilization by Glu-...Lys+ salt
bridges in short peptides of de novo design. Proc. Natl. Acad. Sci. U.S.a. *84*,
8898–8902.

Martínez-Gil, L., Saurí, A., Marti-Renom, M.A., and Mingarro, I. (2011). Membrane
protein integration into the endoplasmic reticulum. Febs J *278*, 3846–3858.

Mbaye, M.N., Hou, Q., Basu, S., Teheux, F., Pucci, F., and Rooman, M. (2019). A
comprehensive computational study of amino acid interactions in membrane
proteins. Sci. Rep. *9*, 12043–14.

Okamura, Y., Fujiwara, Y., and Sakata, S. (2015). Gating Mechanisms of Voltage-
Gated Proton Channels. Annu Rev Biochem *84*, 685–709.

Samanta, U., Pal, D., and Chakrabarti, P. (1999). Packing of aromatic rings against
tryptophan residues in proteins. Acta Cryst (1999). D55, 1421-1427
[Doi:10.1107/S090744499900726X] 1–7.

Shurtleff, M.J., Itzhak, D.N., Hussmann, J.A., Schirle Oakdale, N.T., Costa, E.A.,
Jonikas, M., Weibezahn, J., Popova, K.D., Jan, C.H., Sinitcyn, P., et al. (2018). The
ER membrane protein complex interacts cotranslationally to enable biogenesis of
multipass membrane proteins. eLife *7*, 382.

Tamborero, S., Vilar, M., Martínez-Gil, L., Johnson, A.E., and Mingarro, I. (2011).
Membrane insertion and topology of the translocating chain-associating
membrane protein (TRAM). J Mol Biol *406*, 571–582.

Toyoshima, C., Nakasako, M., Nomura, H., and Ogawa, H. (2000). Crystal
structure of the calcium pump of sarcoplasmic reticulum at 2.6 Å resolution.
Nature *405*, 647–655.

Tsirigos, K.D., Govindarajan, S., Bassot, C., Västermark, Å., Lamb, J., Shu, N., and
Elofsson, A. (2018). ScienceDirect Topology of membrane proteins — predictions,
limitations and variations. Curr Opin Struct Biol *50*, 9–17.

Tsirigos, K.D., Peters, C., Shu, N., Käll, L., and Elofsson, A. (2015). The TOPCONS
web server for consensus prediction of membrane protein topology and signal
peptides. Nucleic Acids Res. *43*, W401–W407.

Walther, T.H., and Ulrich, A.S. (2014). Transmembrane helix assembly and the role
of salt bridges. Curr Opin Struct Biol *27*, 63–68.

a)

b)

AA	Gap	Sequence	ΔG$_{pred.}$	ΔG$_{exp.}$
L4/A15	-	AAAAA**L**A**L**AAAAA**L**A**L**AAAA	-0.49	-0.40 ± 0.11
K	-	AAAAA**L**A**L**AAAAA**L**A**L**AAAA	+1.41	+0.15 ± 0.07
D	-	AAAAA**L**A**L**AAAA**D**A**L**AAAA	+1.39	+0.14 ± 0.07
KD	1	AAAAA**L**A**L**AAA**KD**A**L**AAAA	+3.22	+0.18 ± 0.04
KD	2	AAAAA**L**A**L**AA**K**A**D**A**L**AAAA	+2.93	+0.27 ± 0.05
KD	3	AAAAA**L**A**L**A**K**AA**D**A**L**AAAA	+3.35	+0.22 ± 0.03
KD	4	AAAAA**L**A**L****K**AAA**D**A**L**AAAA	+3.34	+0.13 ± 0.05
KD	5	AAAAA**L**A**K**LAAA**D**A**L**AAAA	+2.98	+0.31 ± 0.02
DD	4	AAAAA**L**A**L**D**AAAA**D**A**L**AAAA	+3.55	+0.23 ± 0.01

c)

Figure 2

Figure 3

Figure 4

a)

Name	Sequence				Gap	$\Delta G_{pred.}$	$\Delta G_{exp.}^{in\ vitro}$	$\Delta G_{exp.}^{in\ vivo}$
wt	²⁴³VGVTSWGYSFL**D**IVA**K**YIFAFLLLNYLTS²⁷¹				4	+1.73	-0.24	-1.54
DD	VGVTSWGYSFL**D**IVA**D**YIFAFLLLNYLTS				4	+1.66	+0.11	-
DA	VGVTSWGYSFL**D**IVA**A**YIFAFLLLNYLTS				4	-0.38	-0.88	-
DK i, i+5	VGVTSWGYSFL**D**IVAY**K**IFAFLLLNYLTS				5	+1.25	+0.31	+1.25

In the wt sequence, position 254 is marked, and 258 is marked.

e)

EndoH	+	-	-	-	-
	wt		DA	DD	i, i+5
	1	2	3	4	5

●●
●○
○○

f)

EndoH

kDa

	wt	i, i+5	wt	i, i+5
EndoH	+	+	-	-
	1	2	3	4

●●
●○
○○

g)

i, i+4   i, i+5

D254
3.5 Å
K258

Figure 5

S, Short (membrane embedded)

L, Long (full helix A)

F78

A69

3.0 Å

L49

b)

Name	Sequence	Gap	ΔG_pred.	ΔG$_{exp.}^{in\ vitro}$	ΔG$_{exp.}^{in\ vivo}$
L (L49-F78)	49 59 63 78 LWELVIEQFEDLLVRILLLAACISFVLAWF	4	+1.06	-1.05	-2.06
L i, i+5 (L49-F78)	LWELVIEQFEDLLVIRLLLAACISFVLAWF	5	+0.44	-0.34	-1.05
S (L49-A69)	LWELVIEQFEDLLVRILLLAA	4	+4.12	+0.22	-
S i, i+5 (L49-A69)	LWELVIEQFEDLLVIRLLLAA	5	+3.46	+0.06	-

c)

| S wt EndoH | L wt | S wt | L i, i+5 | S i, i+5 |

●●
●○
○○

1   **2**   **3**   **4**   **5**

d)

kDa   EndoH   L wt   L i, i+5

+   -   +   -

●●
●○
○○

1   **2**   **3**   **4**

e)   *i, i+5*

F78

A69

10 Å

L49

Figure 6

a)

b)

	3QBG wt		3QBG *i, i+5*
EndoH	−	+	−
ΔG$_{exp.}$	-0.81	−	+0.60

1	2	3

●●
●○
○○

c)

	1SU4 L wt		1SU4 L *i, i+5*	
EndoH	+	−	+	−
ΔG$_{exp.}$	−	-1.24	−	-0.50

1	2	3	4

●●
●○
○○

Figure S1

a)

AA	Gap	Sequence	ΔG_{pred.}	ΔG_{exp.}
L5/A14	-	AAAALALAALAALALAAAA	-1.00	-1.17 ± 0.12
K	-	AAAALALKALAALALAAAA	1.00	0.31 ± 0.08
D	-	AAAALALAALADLALAAAA	0.98	0.57 ± 0.08
KD	3	AAAALALAKLADLALAAAA	2.97	0.69 ± 0.03
KD	4	AAAALALKAAADLALAAAA	2.93	0.58 ± 0.05
KD	5	AAAALAKLAAADLALAAAA	2.14	0.88 ± 0.04

b)

Figure S2

Figure S3

Figure S4

Figure S5

ΔG_{pred} (Kcal/mol)

Figure S6

5MG3
helix E

3.1

2.6

6CC4
helix A

3.2

3.6