Intra-helical salt bridge contribution to membrane protein insertion

Gerard Duart [1] [#], John Lamb [2] [#], Arne Elofsson [2] [*] and Ismael Mingarro [1] [*]

[1] Departament de Bioquímica i Biologia Molecular, Insitut Universitari de Biotecnologia i Biomedicina (BioTecMed), Universitat de València. E-46100 Burjassot, Spain.

[2] Science for Life Laboratory and Department of Biochemistry and biophysics, Stockholm University, 171 21 Solna, Sweden

[#] Equal contribution (alphabetical order)

[*] Corresponding authors

ABSTRACT

Salt bridges between negatively (D, E) and positively charged (K, R, H) amino acids play an important role in protein stabilization. This has a more prevalent effect in membrane proteins where polar amino acids are exposed to a very hydrophobic environment. In transmembrane (TM) helices the presence of charged residues can hinder the insertion of the helices into the membrane. This can sometimes be avoided by TM region rearrangements after insertion, but it is also possible that the formation of salt bridges could decrease the cost of membrane integration. However, the presence of intra-helical salt bridges in TM domains and their effect on insertion has not been properly studied yet. In this work, we use an analytical pipeline to study the prevalence of charged pairs of amino acid residues in TM α-helices, which shows that potentially salt-bridge forming pairs are statistically over-represented. We then selected some candidates to experimentally determine the contribution of these electrostatic interactions to the translocon-assisted membrane insertion process. Using both *in vitro* and *in vivo* systems, we confirm the presence of intra-helical salt bridges in TM segments during biogenesis and determined that they contribute between 0.5-0.7 kcal/mol to the apparent free energy of membrane insertion (ΔG_{app}). Our observations suggest that salt bridge interactions can be stabilized during translocon-mediated insertion and thus could be relevant to consider for the future development of membrane protein prediction software.

KEYWORDS

39   electrostatic interactions; membrane insertion; salt bridge; translocon; transmembrane

40   helix;

INTRODUCTION

Most integral membrane proteins have to insert their transmembrane (TM) segments into the lipid bilayer in a helical conformation and then acquire a defined three-dimensional structure by packaging their helices [1]. α-Helical TM segments are largely composed of apolar residues because of the hydrophobic nature of the membrane environment. Nevertheless, in some cases, it is necessary for the protein activity to include polar amino acids in a TM region in order to develop a functional or structural role [2]. This fact is sometimes not contemplated in modern membrane topology prediction tools [3,4], in which the presence of charged amino acids in a sequence automatically suppose a penalty increase in the predicted free energy (ΔG_{pred}) of insertion. The presence of polar amino acids in TM regions is more frequent than what would be expected [5], especially when these are in pairs on the same face of an α-helix.

Salt bridges are electrostatic interactions between negatively (D, E) and positively charged (K, R, H) amino acids that play an important role in protein stabilization [6]. Many studies have shown that pairs of charged residues that form potential salt-bridges stabilize soluble α-helices [7]. Salt bridges play an important role in the folding of globular proteins and, despite their low occurrence in TM domains, it seems that the contribution in membrane protein stability could be even more determinant. This contribution is especially important in membrane protein biogenesis, as salt bridges help to bury the polarity of charged residues in a hydrophobic environment [8]. Apart from that, it has been suggested that potential salt bridges could help in the insertion of TM α-

helices [5,9], even though their predicted ΔG penalty is well above what is usually seen

for TM segments.

To investigate the potential formation of intra-helical salt bridges in TM α-helices,

we analyzed the composition of the TM domains from membrane proteins of known

structures looking for preferences in the pairing of charged amino acids. This analysis

showed that charged residue pairing is more prevalent than expected for pairs located on

the same face of α-helices within membranes. Likely, salt bridge formation on the same

face of α-helices reduces the unfavorable energetics of inserting ionizable residues into

the hydrophobic membrane core [10]. We use this knowledge together with the known

membrane protein structures to generate a list of potential candidates for further *in vitro*

and *in vivo* experiments.

In this work, we have studied the presence of intra-helical salt bridges in TM

domains using *in silico*, *in vitro* and *in vivo* systems, showing that despite this being an

unexpected phenomenon in nature, salt bridges can be crucial for membrane insertion.

Also, we determined in a quantitative manner that the apparent free energy (ΔG_{app}) of

membrane insertion through the translocon machinery can be decreased between 0.5-

0.7 kcal/mol by position-specific charge pair interaction, which is not contemplated in

the commonly used ΔG predictors. These findings will lead to a better understanding of

the insertion mechanism of TM helices and to improve prediction tools that would more

accurately be able to model the presence of charged residues in these helices.

RESULTS

Charge pair interactions in a model transmembrane helix.

To test the contribution of potential salt bridges to the translocon-mediated membrane insertion, we used the vehicle protein leader peptidase (Lep) from *Escherichia coli* (Fig.1a). The Lep protein consists of two TM segments (H1 and H2) connected by a cytoplasmic loop (P1) and a large C-terminal (P2) domain, which inserts into endoplasmic reticulum (ER)-derived rough microsomes with both termini located in the microsomes lumen. The designed TM segments were inserted into the luminal P2 domain and flanked by two acceptor sites (G1 and G2) for N-linked glycosylation. Glycosylation occurs exclusively in the lumen of the ER (or microsomes) because of the location of the oligosaccharyltransferase (OST) active site (a translocon-associated enzyme responsible for the oligosaccharide transfer) [11]. In this case, the engineered glycosylation sites can be used as membrane insertion reporters because G1 will always be glycosylated due to its native luminal localization, but G2 will be glycosylated only upon translocation of the analyzed sequence across the microsomal membrane. A singly glycosylated construct in which a tested sequence is inserted into the membrane has a molecular mass ~2.5 kDa higher than the molecular mass of Lep molecule expressed in the absence of microsomes; the molecular mass shifts by ~5 kDa upon double glycosylation which facilitates its identification by gel electrophoresis when expressed in the presence of [^{35}S-labeled] amino acids. Then, *in vitro* transcription/translation of these chimeric proteins in the presence of rough microsomal membranes (RMs) allows for accurate and quantitative description of membrane insertion of designed sequences [12-15]. The degree of

membrane insertion is quantified by analyzing the fractions of singly glycosylated (i.e.,

membrane inserted) and doubly glycosylated (i.e., non-inserted) molecules, which can

be expressed as an experimental apparent free energy of membrane insertion, ΔG_{exp} (see

Materials and Methods) [12].

We first compared the effects of oppositely charged Lys and Asp residues on the

insertion of a 19-residue-long hydrophobic stretch (L4/A15 scaffold, 4 leucines and 15

alanines), which was designed to insert stably into the RM membranes [12], including

charges centered in the TM segment at different positions (Fig. 1b) and "insulated" from

the surrounding sequence by N- and C-terminal GGPG- and -GPGG tetrapeptides. Single

Lys and Asp residues were placed in positions 8 and 12 respectively, and pairs of Lys-Asp

residues were designed to cover positions 7-12 (that is, more than one helical turn). Our

results showed a tendency to better insertion when pair charges were placed in positions

(i, i+1; i, i+3; i, i+4) that are permissive with salt bridge formation (Fig. 1c), actually an

effect not observed in the predictions (Fig. 1b). Being a thermodynamically inconvenient

phenomenon within the membrane environment, it is expected that the sequence

context and the amino acid composition of the TM helix would be determinant for salt

bridge formation. Accordingly, we scrutinized a large dataset of membrane proteins of

known three-dimensional structures in order to focus on natural salt bridges present

within TM segments.

Charged pairs in transmembrane helices

Alpha helices are a common secondary structure in both globular and membrane

proteins. We created two main datasets, TM dataset with helical membrane proteins of

known structure from the PDBTM-dataset [16], and GLOB dataset with globular alpha

helical proteins selected from the SCOP-database [17,18], see methods for the full

creation steps. Table 1 shows a breakdown of alpha-helices, charges and salt bridges in

the two datasets. What is clear is that long alpha helices (≥ 17 residues) form a larger

proportion in the TM dataset than in the globular helices (GLOB) dataset, which is logical

as most TM helices need to span through the hydrophobic core of the lipid bilayer of

thickness ~30 Å.

Our TM dataset showed the same distribution of polar charges as previous studies

[19], with about 10 % of polar residues in the core membrane regions (see Table 1), and

about one-third of these are charged. Over half of the charged residues could form pairs

with other charged residues at intervals of i, $i+1$; i, $i+3$ and i, $i+4$. In contrast, the GLOB

dataset had a much higher proportion of charged polar residues. The GLOB dataset also

contained about 15 times as many charged pairs relative to its size compared to the TM

dataset, again indicating that charged residues are more common in globular than in TM

helices.

The two datasets, TM and GLOB, were extended with homologous sequences

identified by searching with jackhammer against UniProt. All sequences with an E-value

lower than 10^{-3} were included. These datasets are named TM-MSA and GLOB-MSA.

Using data in the TM-MSA dataset to produce the log odds ratios, we identified

periodicity patterns of charged residue pairs that are more common than what would

have been expected from the underlying amino acid composition, see Figure 2). We found

that polar residues at pairs i, i+3; i, i+4 and i, i+7 are significantly enhanced (Fig. 2). This

feature is strengthened when we examine the same plot for the GLOB-MSA dataset,

where these patterns were not observed (Figure S1). This was also clear when statistical

significance was taken into account, see Figures S2 and S3.

Table 2 lists the log odds ratios together with errors and familywise error-

corrected p-values for all pairs at separation up to seven residues. It is clear that pair

residues placed at i, i+1; i, i+3 and i, i+4 positions are most significant, especially in the

case of oppositely charged pairs.

Structural analysis of charged residues in transmembrane helices

The abundance of salt bridges also shows a remarkable difference between

transmembrane and globular proteins. In the GLOB dataset over 13% of the proteins

contains at least one local salt bridge in an alpha helix whereas just over 5% of the

membrane proteins contains a local salt bridge, see Table 1. This does conform to our

current understanding of soluble versus TM alpha-helices and their different

environments, and suggests that salt bridges in TM regions perform a vital functional

and/or structural role.

As seen in Figure 2, charged pairs of amino acids are especially prevalent at

positions i, i+1; i, i+3 and i, i+4. Oppositely charged residues stand out, especially Glu-Arg

at i, i+1, Glu-Lys at i, i+3 and Asp-Lys at i, i+4. Also, several same charged pairings at i,

i+3 and i, i+6 are more frequent than expected. Other known structural features can also

be hinted at, including aromatic ring stacking by His-Trp pair [20] at i, i+6.

Charged pairs placed at i, i+1; i, i+3 and i, i+4 could potentially form salt bridges

as they are all on the same relative face of the alpha helix and are close enough in vertical

separation on the helix (see Figure 3, top). Although oppositely charged pairs at i, i+7 are

also on the same face of the alpha helix, unless the alpha helix has a bend, both residues

are too far separated to form a salt bridge. This was clearly seen in Figure 3 where the TM

dataset was used. In both absolute count (Fig. 3 panel a) and log odds ratios (Fig. 3 panel

b) it is clear that residues at i, i+1; i, i+3 and i, i+4 are by far the most common and

overrepresented pairings. Figure 3b also shows that oppositely and same charged pairs

have about the same overrepresentation at i, i+3, whereas oppositely charged pairs are

stronger than same charged pairs at i, i+1and i, i+4, both within salt bridge range, and

same charged pairs are stronger at positions i, i+7 and i, i+8, too far to form salt bridges.

When structure-observed salt bridges in the different positions were compared to

the oppositely charged pairs a clear image aroused, see Figure 3c. Even though there are

more oppositely charged pairs at position i, i+1 than in positions i, i+3 and i, i+4 with all

log odds ratios over 1.0 (Fig. 3b), only about 15% of the oppositely charged pairs at i, i+1

form salt bridge (Fig. 3c). This is in contrast to i, i+3 where almost 40 % of the pairs form

salt bridges, and just under 25% at i, i+4 (Fig. 3c).

Selection of natural salt bridges from membrane protein structures

To further look for candidate proteins containing membrane-spanning helices with salt

bridges we started with the redundant (TM-Red) dataset of known structures. Each of

the 8,687 proteins in our dataset were scanned for oppositely charged pairs in positions i,

$i+1$; i, $i+3$ and i, $i+4$ within any TM segments core region. For each of these, we only kept

proteins with at least one TM segment that contains such a pair and where this pair was

within salt bridging distance (183 protein chains with a total of 188 salt bridges).

When we now make a selection for potential candidates, we also look at the

estimated ΔG_{pred} values and choose helices with a positive value above one, as per our

hypothesis, a salt bridge helps TM insertion of sequences for which the hydrophobic force

would not be enough to insert. This stricter definition results in a set of 136 candidates

of a total of 138 salt bridges and we selected TM7 (helix G) from Halorhodopsin protein

(PDB ID: 3QBG) with an estimated ΔG_{pred} value above +1.7 kcal/mol, and helix A from

calcium ATPase (PDB ID: 1SU4) with a higher estimated ΔG_{pred} value (above 4.1

kcal/mol), as candidates for systematic studies. See the github repository

(https://github.com/ElofssonLab/salt_bridges) for the full lists.

209 Intra-helical salt bridge stabilizes the insertion of helix G from halorhodopsin.

Halorhodopsin (hR) from *Natronomonas pharaonis* (3QBG) is a protein made up of

seven TM helices (helix A through G) and a retinal chromophore that is bound via a

protonated Schiff base to the ε-amino group of a lysine (K258) residue located

roughly in the middle of helix G [21]. *In silico* analysis of 3QBG structure, the anion-

free form of the protein, showed a charged pair of amino acids (Asp-Lys), involving the

functional K258 and the D254 in the center of helix G (Fig. 4). The position (i, i+4) and

the distance between the anionic carboxylate (RCOO⁻) from the Asp residue and the

cationic ammonium (RNH₃⁺) from the lysine residue, in the crystal structure, was about

3.5 Å, a permissive distance for a salt bridge formation (Fig. 4g), which has been

established as being lower than 4 Å [22]. To get insights into this interaction, we designed

three mutants that were supposed to perturb the salt bridge interaction by different ways:

K258D mutant by placing two charged residues with the same polarity at positions i, i+4;

K258A mutant by replacing one of the charged residues by a non-polar amino acid, and

K258Y/Y259K double mutant by placing the charged pair at a non-permissive salt bridge

position (i, i+5; Fig. 4g), while keeping the same amino acid composition (Fig. 4a).

hR is a trimeric protein in which the helix G is neither exposed at the monomer-

monomer interface nor oriented to the inner part of the trimeric structure (Fig. 4, panels

b-d). In fact, the charged pair found in helix G is oriented toward the core of the 'globular'

structure in each monomer. Then, the insertion of helix G was studied using the Lep-

based assay. When constructs harboring helix G wild type sequence were translated *in*

vitro in the presence of RMs singly-glycosylated (reporting insertion) forms were found

(Fig. 4e, lane 2), despite its positive (suggesting non-insertion) ΔG_{pred} value (Fig. 4a). The

nature of the higher molecular weight polypeptide species was analysed by

endoglycosidase H (EndoH) treatment, a highly specific enzyme that cleaves N-linked

oligosaccharides. Treatment with EndoH of the samples eliminated higher molecular

mass bands (Fig. 4a, lane 1), confirming the sugar source of the retarded electrophoretic

mobility bands and suggesting helix G insertion into the microsomal membrane.

However, locating the Asp-Lys pair at *i*, *i*+5, which is non-compatible with salt bridge

interaction (Fig. 4g), strongly reduced the experimental insertion efficiency (Fig. 4e, lane

5). Interestingly, replacing the positively charged lysine residue by a negatively charged

aspartic acid residue (K258D) rendered similarly low levels of insertion efficiency (Fig.

4e, lane 4). As expected, replacement of the ionizable lysine residue by the aliphatic

alanine increased the insertion efficiency (Fig. 4e, lane 3). In this later mutant, the most

likely cause is the absence of the positively charged lysine amino side chain and by the

presence of the methyl side chain group of the mutant alanine residue. The results of the

Lep-based glycosylation assay indicated that wild type (wt) and DA sequences (two

stabilized charges or only one charge, respectively) are inserted properly into the

microsomal membrane (ΔG_{exp} values -0.24 and -0.88 kcal/mol, respectively), but when

the salt bridge is disrupted, either by having two charged amino acids with the same

polarity (DD) or by placing oppositely charged residues at a non-permissive distance (*i*,

i+5) in the center of the helix, the translocation of the segment increases substantially. It

should be mentioned that the K258Y/Y259K double mutant has the same amino acid

composition than the original helix G, but insertion efficiency is remarkably decreased

(ΔG_{exp} = +0.31 kcal/mol). Together these results show that the interaction (salt bridge)

between Asp and Lys residues in the center of the helix G from 3QBG is essential for its

proper insertion into the microsomal membrane. The salt bridge contributes

approximately ~0,5 kcal/mol to the apparent experimental free energy of membrane

insertion.

Next, to ensure that the *in vitro* results are relevant to the *in vivo* situation, wt

and *i*+5 constructs were also expressed *in vivo* in HEK-293T cells. To this end, a *c*-myc

tag was engineered at the C-terminus of the Lep chimera to allow immune-detection of

our constructs in cell extracts. As shown in Fig. 4f, transfected cells with the chimera

containing helix G wt sequence rendered singly-glycosylated molecules, indicating *in*

vivo membrane insertion. In contrast, cells transfected with the construct harboring *i*,

i+5 sequence rendered almost exclusively doubly-glycosylated forms, as proved by

EndoH treatment (Fig. 4f, lane 2), suggesting membrane translocation. These results

emphasized the relevance of salt bridge interactions in translocon-mediated TM

insertion, especially in the *in vivo* (cellular) environment.

Salt bridge contribution to the insertion of a heavily charged helix.

In order to challenge salt bridge interactions in a more hydrophilic TM helix we focus on

helix A from the sarcoplasmic/ER calcium ATPase 1 [23]. Calcium ATPase (PDB ID:

1SU4, *Oryctolagus cuniculus*) is a member of the P-type ATPases that transport ions

across the membrane against a concentration gradient involving 10 α-helices (helix A

through J) in the membrane-embedded region [23]. *In silico* analysis of 1SU4 structure,

the crystal structure of the calcium ATPase with two bound calcium ions, showed Asp-

Arg (DR) pair (D59 and R63) in the center of the helix A. This helix extends beyond the

membrane (Fig. 5a, from Leu49 to Phe78) and shows some particularities. On the one

hand, in the structure the membrane-embedded stretch (the N-terminal region of helix

A) encompasses from Leu49 to Ala69 residues, and includes several charged amino acids,

probably involved in the Ca^{2+} transport across the membrane (Glu51, Glu55, Glu58,

Asp59 and Arg63). Therefore, the ΔG_{pred} value for this segment (L49-A69) is remarkably

higher (and positive) than expected for a TM helix (Fig. 5b). On the other hand, the C-

terminal region of this helix contains a high prevalence of non-polar amino acids that is

more compatible with the hydrophobicity of the membrane core (Fig. 5b). Nevertheless,

the presence of the functional amino acids (e.g. Glu58) in the membrane-embedded

region reinforce the idea that the more hydrophilic N-terminal region must ultimately

be embedded within the membrane [23]. It has been previously shown that the position

in the membrane of TM helices in protein folded structures does not always correspond

to the thermodynamically favored positions in the membrane of the isolated helices [24].

Instead, after translocon-mediated insertion of the more hydrophobic region,

repositioning of TM helices relative to the lipid bilayer provides a convenient way for

non-hydrophobic polypeptide segments to become buried within the membrane. Then,

the nature of helix A suggested the possibility that initial insertion of the hydrophobic

region can be followed by subsequent repositioning of the charged region into the

membrane hydrophobic core, which will be the final segment embedded in the lipid

bilayer   (Fig.   5a).   Interestingly,   the ΔG Predictor   server

(https://dgpred.cbr.su.se/index.php?p=home) selected the adjacent (C-terminus, L60-

F78) hydrophobic region as TM (Fig. 5b), instead of the charged region found at the high-

resolution structure (Uniprot code: P04191).

Focusing on the potential salt bridge residues (D59 and R63 pair), the distance

between the anionic carboxylate (RCOO$^-$) from the D59 and the cationic guanidinium

$(RC(NH_2)_2^+)$ from the R63 was about 3.0 Å in the crystal structure (Figure 5a), clearly

within the permissive range for salt bridge interaction. To investigate the contribution of

this potential salt bridge interaction in the translocon-mediated membrane insertion of

this region, we worked with two different scaffold sequences: the full helix A involving

the residues 49-78 (Long, L); and a shorter membrane-embedded version including

residues 49-69 (Short, S) as found in the solved structure. We also challenged the D59-

R63 charge pair interaction in both sequences by increasing the separation between the

ionizable residues from the native i, $i+4$ to non-permissive i, $i+5$, while maintaining amino

acid composition (R63I/I64R double mutant). *In vitro* transcription/translation of these

sequences in the presence of microsomes rendered singly-glycosylated molecules for the

construct containing full-length helix A (Fig. 5c, lane 2). In contrast, when only the

membrane-embedded sequence was included, the Lep chimera was mainly doubly-

glycosylated (Fig. 5c, lane 3), suggesting that helix A is properly inserted when the full

helical sequence is present. Still, the translocon inefficiently inserted the isolated

membrane-embedded region. When the charge paired residues were placed at a non-

permissive distance in terms of salt bridge interaction (i, $i+5$; Fig. 5e) the insertion

efficiency was reduced (Fig. 5c, lane 4), with a ΔG_{app} decrease (absolute values) of ~0,7

kcal/mol relative to the wild type sequence (Fig. 5b). As expected, this effect was not

observed when the same mutations were grafted on the membrane-embedded (S)

sequence (Fig. 5c, lane 5).

Finally, we analysed the salt bridge interaction in HEK-293T cells to study the

translocon performance *in vivo*. When cell cultures were transfected with a Lep-derived

chimera containing the helix A wild type sequence only singly-glycosylated molecules

were observed (Fig. 5d, lanes 1 and 2). However, a construct harboring double mutant

(R63I/I64R; i, i+5) sequence showed doubly-glycosylated molecules to a measurable

extent (Fig. 5d, lanes 3 and 4), indicating a lower insertion efficiency that can be

attributed to the altered salt bridge interaction.

DISCUSSION

Charged residues found in alpha-helices can both give a stabilizing effect [25] as well as

being important for interaction and function such as in zinc finger motifs [26]. Charged

residues have also been found to be more common in globular (soluble) helices compared

to TM helices [27]. This can be explained by the fact that globular helices reside in a more

hydrophilic environment compared to the hydrophobic environment of the lipid bilayer.

An intermediate between these is the existence of amphipathic alpha-helix in contact

with the surface of a bilayer, where one face contains mainly polar residues facing an

aqueous environment and the opposite face with mostly nonpolar residues facing a

hydrophobic environment [28].

340   Whereas pairs of charged residues of the same charge facing the inside pores in

TM regions fill an essential functional role, pairs of oppositely charged residues can form

salt bridges that can stabilize the helix and play an important role in function such as for

transportation [29]. Salt bridges might also be important for hiding the charges during

translocon-mediated TM helix insertion, as the charged residues are hidden from the

hydrophobic environment. Previous work has shown that charged and polar residues are

conserved within TM segments [19], indicating they are crucial for function and/or

stability.

Asp-Lys pairs at position i, $i+4$ and Glu-Lys pairs at position i, $i+3$ are the most

prevalent as seen previously in Figure 2. They are both among the most prevalent

oppositely charged pairs and the charged pairs that form the highest number of salt

bridges in membrane protein structures. This is in stark contrast to Glu-Arg pair at

position i, $i+1$ that although as frequent in pairs as Asp-Lys and Glu-Lys at positions i, $i+4$

and i, $i+3$ respectively, only form salt bridges in one-fourth of the cases.

An interesting observation is that positively charged Arg-Arg pairs at position i,

$i+3$ is numerically the most common pair at this position. Positively charged pairs at i, $i+6$

are also more prevalent than expected and although Arg-Arg pairs numerically only show

up half as often at i, $i+6$ as in i, $i+3$, 80% of the Arg-Arg pairs at position i, $i+6$ also contain

an arginine at $i+3$, as found in helices from voltage-gated ion channels that contain three

or more periodically aligned Arg residues with two intervening hydrophobic residues

[30].

The fraction of salt bridges at positions i, $i+5$ (Fig. 3c) is an anomaly due to bend

alpha-helices, as found in the helix E from bacterial translocon (PDB ID: 5MG3) and in

the helix A from a lipid flippase (PDB ID: 6CC4) due to the presence of glycine residues

(Supplementary Fig. S4). Without the observed helix bending, these salt bridges would

not be formed.

By analyzing two native helices containing intra-helical salt bridges we now find

that the free energy of insertion (ΔG_{app}) is significantly reduced if both oppositely charged

residues are spaced at a permissive distance. These results indicate that intra-helix salt

bridge can form during translocon-assisted insertion or even earlier, since in contrast to

globular (soluble) helices, TM helices can be compacted inside the ribosome exit tunnel

[31]. The maximal reduction in ΔG_{app} seen with Asp-Lys and Asp-Arg pairs in both hR

and Ca ATPase helices is 0.5-0.7 kcal/mol, which is in good agreement with the 0-1

kcal/mol estimated for these two pairs from thermodynamic peptide partition into

octanol experiments [32]. As found in the case of hR helix G (Fig. 4f), this reduction might

be even higher in the cell context, since some auxiliary components of the membrane

insertion machinery [15,33,34] can be in suboptimal conditions in the microsomal

vesicles.

Beyond the conceptual issues involving the membrane insertion process, we note

that the availability of quantitative experimental data on the contribution of salt bridge

interactions to the free energy of insertion (ΔG_{app}) will make it possible to fine-tune

current membrane protein topology-prediction methods based on free energy

calculations.

MATERIALS AND METHODS

Enzymes and chemicals. TNT T7 Quick for PCR DNA was from Promega (Madison, WI, USA). Dog pancreas ER rough microsomes were from tRNA Probes (College Station, TX, USA). EasyTag™ EXPRESS^{35}S Protein Labeling Mix, [^{35}S]-L-methionine and [^{35}S]-L-cysteine, for *in vitro* labeling was purchased from Perkin Elmer (Waltham, MA, USA). Restriction enzymes were from New England Biolabs (Massachusetts, USA) and endoglycosidase H was from Roche Molecular Biochemicals (Basel, Switzerland). PCR and plasmid purification kits were from Thermo Fisher Scientific (Ulm, Germany). All oligonucleotides were purchased from Macrogen (Seoul, South Korea).

DNA Manipulation. The sequences of interest were introduced into the modified Lep sequence from the pGEM1 plasmid [12] between the *Spe*I and *Kpn*I sites using two double-stranded oligonucleotides with overlapping overhangs at the ends. The complementary oligonucleotides pairs were first annealed at 85 °C for 10 min followed by gradual cooling to 30 °C and ligated into the vector (a kind gift from G. von Heijne's lab). Mutations were obtained by site-directed mutagenesis using the QuikChange kit (Stratagene, La Jolla, California). Lep system including the sequences of interest in the P2 region were subcloned into *Kpn*I linearized pCAGGS in-house version using In-Fusion HD cloning Kit (Takara) according to the manufacturer's instructions. *c*-myc tag (Glu-Gln-Lys-Leu-Ile-Ser-Glu-Glu-Asp-Leu, EQKLISEEDL) was added in Ct in hanging with de PCR primer before cloning. For *in vitro* assays, DNA was amplified by PCR adding the T7 promoter during the process. All sequences were confirmed by sequencing the plasmid DNA at Macrogen Company (Seoul, South Korea).

Translocon-mediated insertion into microsomal membranes. Lep constructs in pGEM

with L4/A15, 3QBG and 1SU4 segments and its variations were transcribed and translated

using the TNT T7 Quick Coupled System (#L1170, Promega). Each reaction containing 1

μL of PCR product, 0.5 of EasyTag™ EXPRESS 35S Protein Labeling Mix (Perkin Elmer)

(5.5 μCi) and 0.3 μL of microsomes (tRNA Probes) was incubated at 30ºC for 90 min. Endo

H treatment was done following the manufacturer's instructions. Samples were analysed

by SDS-PAGE. The bands were quantified using a Fuji FLA-3000 phosphoimager and the

Image Reader 8.1j software. Free energy was calculated using: $\Delta G_{app} = -RT \ln K_{app}$, where

$K_{app} = f_{2g}/f_{1g}$ being f_{1g} and f_{2g} the fraction of singly glycosylated and double glycosylated

protein, respectively.

Free apparent insertion energy, ΔG. The free insertion energy of a TM region, ΔG, is

calculated as per the experimentally defined Biological hydrophobicity scale [12]. This

scoring is amphiphilic with hydrophobic residues contributing a lower (negative) ΔG

while hydrophilic contributes a higher (positive) ΔG. A region's total ΔG is the sum of

these individual scores weighted on where in the region the residue, a residue in the

middle of the helix has a higher weight than residues at the ends. This scoring can give

an indication of how favourable the amino acid composition of a TM region is to be

inserted in a lipid bilayer membrane. To note is that hydrophobicity alone is not the only

driving force and that the positive inside rule [35,36] and help from proceeding TM

regions [37,38] can help insertion in polytopic TM proteins especially.

Expression in mammalian cells. Lep constructs containing 3QBG or 1SU4 segments and

its variations were tagged with c-myc epitope at their Ct (Glu-Gln-Lys-Leu-Ile-Ser-Glu-

Glu-Asp-Leu, EQKLISEEDL) and inserted in a pCAGGS-ampicillin plasmid. Once the

sequence was verified, plasmids were transfected into HEK293-T cells using

Lipofectamine 2000 (Life Technologies) according to the manufacturer's protocol.

Approximately 24 h post-transfection cells were harvested and washed with PBS buffer.

After a short centrifugation (1000 rpm for 5 min on a table-top centrifuge) cells were

lysed by adding 100 μL of lysis buffer (30 mM Tris-HCl, 150 mM NaCl, 0.5% Nonidet P-

40) were sonicated in an ice bath in a bioruptor (Diagenode) during 10min and

centrifugated. Total protein was quantified and equal amounts of protein submitted to

Endo H treatment or mock-treated, followed by SDS-PAGE analysis and transferred into

a PVDF transfer membrane (ThermoFisher Scientific). Protein glycosylation status was

analysed by Western Blot using an anti-c-myc antibody (Sigma), anti-rabbit IgG-

peroxidase conjugated (Sigma), and with ECL developing reagent (GE Healthcare).

Chemiluminescence was visualized using an ImageQuantTM LAS 4000mini

Biomolecular Imager (GE Healthcare).

Core segment definition. We define core segments as a transmembrane region minus

the first and last 5 residues. This is to ignore the interface regions which are known to

contain polar residues.

Salt bridge definition. Salt bridges are defined as per [39], where a salt bridge is defined

if a side chain carbonyl oxygen atom in Asp-Glu is within 4.0 Å from the nitrogen atom

in Arg-Lys. This conforms to other works [7,40] with the definition that the atoms are

within hydrogen bond distance. We also define local salt bridges as being bridges that

are separated by at most 7 residues in the sequence. This is to separate long salt bridge

interactions, which can occur between spatially close residues that are separated in

sequence, such as coiled coils where salt bridges can be between separate alpha-helices.

Transmembrane helices dataset (TM dataset). The full pipeline is available as a Makefile

together with supporting scripts in the github repository. The full PDBTM database [16]

was downloaded together with their list of non-redundant protein pdb ids. This list is

used to generate both sequence and topology of the proteins by extracting both from the

PDBTM-xml. For each protein in the non-redundant list, the membrane regions are

extracted as per the PDBTM database annotation. All non-membrane regions are

annotated 'i' for convenience. To support future analysis, membrane regions longer than

10 were run through DSSP and annotated with 'M' if all residues in the core segment

were defined as alpha-helix ('H'), otherwise, the full membrane region is annotated 'm'.

Observe that this creates fasta-like 3line files, that only contain topological annotation

with ambiguous TM regions annotated as 'm' instead of the normal 'M'. These proteins

are then clustered using cd-hit [41,42] at 40% identity using the parameter -c 0.4 -n 2 -T

0 -M 0 -d 0.

During the extraction of TM regions, the corresponding structure file from RCSB

was used to calculate all salt bridges within the current protein and any salt bridge that

has at least one residue within any TM region was saved. Additionally, all salt bridges

whose both residues were within the same segment and within 7 residues of each other

were annotated as local as per the salt bridge definition above.

Extraction of charged residues. From all annotated TM regions of length 17 or longer [2],

the core segment was extracted. All these core regions were then scanned and when a

charged residue was encountered, we recorded any other charged residue from 7 residues

before the current one to 7 residues after. This results in charged residues that can contain

a charged pairing partner outside of the TM segment and will therefore differ slightly

from charged pairs which are defined next.

Extraction of charged pairs. From all annotated TM regions of length 17 or longer [2], the

core segment was extracted. All these core regions were then scanned and when a

charged residue was encountered, we look at up to 7 residues in front of it or to the end

of that core region, whichever came first. All occurrences of charged pairs were recorded,

resulting in charged pairs where both residues were fully within the core segment of a

TM helix.

MSA-dataset extension (TM-MSA). Using the TM dataset, we extended it by creating an

MSA alignment of each protein using jackhmmer [43] against Uniref90 with one iteration

and an E-value cut off of 10^{-3} with the following parameters:

-N 1 -E 1e-3 --incE 1e-3 --cpu 14

From each alignment, we then sampled up to 200 hits, including the initial seed

sequence. If there were fewer than 200 hits, we used them all. We then used the

original topology for each alignment to extract all TM regions, only to include parts

where the sequence covers the full TM region and where the sequence did not contain

any insertions or deletions.

Dataset of helices from globular proteins (GLOB and GLOB-MSA). To create a reference

dataset of globular α-helical proteins we extracted all globular all-alpha protein domains

from SCOP. As SCOP classifies domains of proteins resulting in that one domain of a

protein can be annotated as globular whereas another domain as TM (see 1PPJ chain D

as an example) we reduced the SCOP list against the redundant list of all PDBTM [16]

chains to clear any overlap. This results in 4,500 proteins in total.

Topological files with sequence and membrane topology are created with the help

of the RCSB secondary structure file and only membranes annotated as pure α-helices

were retained. This file was further homology reduced and alignments prepared in the

same manner and using the same parameters as the TM dataset described above.

The GLOB-MSA dataset was created in the same way as the TM-MSA dataset

using jackhmmer to extend the sequences to alignments and then to extract helix

sequences.

Redundant (TM-Red) dataset. The full PDBTM database was used as in the preparation

of the TM dataset. In this case we used all redundant sequences and generate topology

files, but we did not clustered the proteins. We added in the constraint that each selected

TM region must fully contain at least one potential salt bridge. This means a local salt

bridge where both residues are within the core segment. This dataset was only used to

find potential candidates for further *in vitro* and *in vivo* experiments.

Acknowledgments

We thank Pilar Selvi for excellent technical assistance. This work was supported by grants

BFU2016-79487-P form the Spanish Ministry of Economy and Competitiveness and

PROMETEU/2019/065 from Generalitat Valenciana (to I.M.), and by grant from the

Conflict of Interest Statement

None declared.

Table 1. About 10 % of the core residues in the TM dataset are polar and just under 4% are charged residues. This is significantly less than 37.5% and 25.5% respectively in the globular dataset. In the transmembrane dataset, just over half of the charged core residues form a charged pair at distance 1, 3 or 4. *In the case of the globular set, there are more potential charged pairs than charged core residues as multiple residues are counted more than once as they form more than one potential pairing with separation of 1, 3 or 4 steps. It is clearly seen that although the TM dataset contains more helices per protein and has a higher proportion of long ($\geq$ 17 residues) helices it contains significantly fewer charged pairs.

Dataset	TM	TM-MSA	GLOB	GLOB-MSA
Charge statistics				
Core residues	40116	3990464	74299	7388277
Polar core residues	4080 (10.2%)	490861 (12.3%)	27851 (37.5%)	2696564 (36.5%)
Charged core residues	1467 (3.7%)	204544 (5.1%)	18959 (25.5%)	1819047 (24.6%)
Pairs of charged core residues (+1, +3 or +4)	825	88380	23755*	2271405*
# of same charged pairs	373	45220	10835	1117270
# of oppositely charged pairs	452	43160	12920	1154135
Charge Pairs statistics				
Proteins	925	130934	2475	363610
Total alpha helices	5435	481729	23991	2118158
Alpha helices $\geq$17	3755	306632	5895	485797
Proteins with charged pairs	204	25187	1780	193534
Helices with charged	265	34029	4251	344142

pairs				
Proteins with any salt bridge	241	-	*751*	-
Proteins with local salt bridge in helix	56	-	*335*	-
Total number of local salt bridges in helices	59	-	*577*	-

Table 2. Log odds ratios for all charged pairs, same charged pairs and oppositely charged

pairs with calculated errors and multiple hypotheses corrected p-values. It is clear that

charged pairs occur more often than predicted, evidenced by the positive log odds ratios

in all cases. It is clear that positions +1, +3 and +4 are the most prevalent pairings, with

the bolded values highlighting log odds ratio above 0.8. It is also clear that oppositely

charged residues which have the potential to form salt bridges are prevalent in all three

positions whereas the same charge is mainly prevalent in position 3. Most likely these

same charges facing the same face of the helix are involved in functions such as ion

transport.

Spacing	All Log odds			Same charged Log odds			Oppositely charged Log odds		
	ratio	error	p-value	ratio	error	p-value	ratio	error	p-value
+1	**0.879**	0.025	$1.31e^{-257}$	0.548	0.042	$4.03e^{-41}$	**1.107**	0.032	$1.65e^{-256}$
+2	0.252	0.037	$2.02e^{-08}$	0.185	0.054	$1.88e^{-00}$	0.316	0.050	$1.09e^{-06}$
+3	**1.073**	0.026	$0.00e^{+00}$	**1.118**	0.035	$1.15e^{-214}$	**1.026**	0.037	$5.78e^{-165}$
+4	**0.965**	0.028	$1.14e^{-253}$	0.696	0.046	$2.88e^{-49}$	**1.177**	0.036	$2.74e^{-233}$
+5	0.517	0.037	$4.88e^{-42}$	0.390	0.055	$4.15e^{-09}$	0.630	0.049	$1.21e^{-34}$
+6	0.540	0.038	$1.66e^{-43}$	0.545	0.053	$2.33e^{-21}$	0.536	0.053	$2.10e^{-20}$
+7	0.615	0.038	$1.21e^{-55}$	0.764	0.050	$7.78e^{-50}$	0.439	0.058	$1.97e^{-10}$

FIGURE LEGENDS

Figure 1. Effects on membrane insertion of single or pairs of Asp and Lys residues in a

model TM segment. (a) Schematic representation of the leader peptidase (Lep) model

protein. G1 and G2 denote artificial glycosylation acceptor sites. The sequence under

investigation was introduced in the P2 region after H2. Recognition of the tested

sequence as a TM by the translocon machinery (highlighted in green) results in the

modification of the G1 site but not G2. The Lep chimera will be double glycosylated if

the sequence being tested is not recognized as a TMD and thus translocated into the

microsomes lumen (shown in red). **(b)** The tested sequences from L4/A15 model TM

(including the charged residues, bold), the gap distance, and the predicted ΔG (ΔG_{pred})

values in kcal/mol are shown. Amino acids with positive and negative charge are

highlighted in blue (K) and red (D) respectively. **(c)** Experimental ΔG (ΔG_{exp}) in kcal/mol

of each tested sequence in the Lep assay. The mean and standard deviation of at least 3

independent experiments are represented (*n* values: 4 [from 1 to 7] and 3 [8 and 9]). The

individual value of each experiment in represented by a solid dot being represented as a

green square the experimental ΔG value for the L4/A15 sequence from [2]. The wt and

simple mutants are shown in white bars. Charges at compatible distances with salt bridge

formation (*i*, *i*+1; *i*, *i*+3; and *i*, *i*+4) are shown in yellow. Not compatible distances with

salt bridge formation (*i*, *i*+2; and *i*, *i*+5) are shown in dark gray. Compatible distances but

not compatible amino acid pair (*i*, *i*+4 DD pair) is shown in clear gray.

Figure 2. Log odds ratios of each pair of amino acids for '*i*, *i*+1' through '*i*, *i*+8' for the TM-

MSA dataset. The rows on the y-axis indicate the first amino acid in the pair and the

columns on the x-axis the second. The residues are ordered by hydrophobicity according

to the Engelman order [44]. See figure S1 for the equivalent of the globular dataset. S2

and S3 show the same plots masked for statistical significance.

Figure 3. Charge pairs in TM helical sequences and structures. Helical wheel projection

and lateral views of an α-helix are shown on top. The initial position *i* and the following

8 residues are numbered. Residues in positions *i*+3 (orange), *i*+4 (yellow), and *i*+7 (light

brown) are mainly on the same face of the helix, but *i*+7 is placed too far for a salt bridge

interaction. **a)** The absolute count of charged pairs for '*i*, *i*+1' through '*i*, *i*+8' in the TM

dataset. The forward slash are the oppositely charged pairs and backward slash are the

same charged pairs. **b)** The log odds ratios of the same data for all charges, oppositely

charged and same charged pairs, all with error bars. **c)** Fraction of oppositely charged

pairs that form local salt bridges. The small bump at *i*+5 are the two proteins 6CC4 (helix

A) and 5MG3 (helix E), which both exhibit a bend in the alpha helix due to the presence

of glycine residues (see Fig. S4).

Figure 4. Insertion of halorhodopsin helix G from *Natronomonas pharaonis* (3QBG) into

microsomal and cellular membranes. (a) Tested sequences from 3QBG including the gap

distance, the distance between charges in Å, and the predicted and experimental ΔG

values (ΔG$_{pred}$ and Δg$_{exp}$ respectively) in kcal/mol are shown. Amino acids with a positive

or negative charge are highlighted in blue (K) and red (D), respectively. Green numbers

indicate negative ΔG (insertion) values, while red numbers denote ΔG values above 0

(translocation). **(b)** Frontal view of 3QBG monomer structure. The helix G is highlighted

in orange with the D254 and K258 shown in sticks colored red and blue respectively. The

membrane position is indicated by a red (outer) and blue (inner) discontinuous line,

according to OPM dataset [22]. Lateral **(c)** and upper **(d)** views of the 3QBG trimeric

structure. The helix G is highlighted in orange with the D and K shown in sticks colored

red and blue, respectively. The different monomers are shown in transparent blue, purple

and indigo. Representative examples (*n*=3) of *in vitro* protein translations in the presence

of ER-derived microsomes **(e)** and Western blots (*n*=3) of *in vivo* protein translations in

HEK-293T cells **(f)** in the presence (+) or absence (-) of Endoglycosidase H (EndoH), a

glycan-removing enzyme. The absence of glycosylation of G1 and G2 acceptor sites is

indicated by two white dots, single glycosylation by one white and one black dot, and

double glycosylation by two black dots. **(g)** Zoom view centered on the salt bridge

between D254 and K257 at *i*, *i*+4 (left) and *i*, *i*+5 (right) gaps. D and K residues are shown

in sticks colored red and blue, respectively, while the dashed line indicates the $RCOO^-$ to

RNH_3^+ distance.

Figure 5. Insertion of Calcium ATPase (1SU4) helix A into microsomal and cellular

membranes. (a) Lateral view of 1SU4 structure. Zoom view of the A helix (right panel).

The membrane-embedded region of helix A is highlighted in yellow. Charged amino

acids are shown as sticks in blue (R), red (D) and pink (E), respectively. L49, A69 and F78

are also shown as sticks to define helix's subdomains. The membrane location is indicated

by red (outer) and blue (inner) discontinuous lines according to OPM dataset [22] and

the distance between the R and D charges is indicated in Å. **(b)** Helix A-derived sequences

from 1SU4 including the gap between charged residues, the distance between charges in

Å, and the predicted and experimental ΔG values (ΔG_{pred} and Δg_{exp} respectively) in

kcal/mol are shown. Amino acids with a positive charge are highlighted in blue (K) while

negatively charged are marked in red (D) and pink (E). Green numbers indicate negative

ΔG (insertion) values while red numbers denote ΔG values above 0 (translocation).

Representative examples (n=3) of *in vitro* protein translations in the presence of ER-

derived microsomes **(c)** and Western blots (n=3) of *in vivo* protein translation in HEK-

293T cells **(d)** in the presence (+) or absence (-) of Endoglycosidase H (EndoH), a glycan-

removing enzyme. The absence of glycosylation of G1 and G2 acceptor sites is indicated

by two white dots, single glycosylation by one white and one black dot, and double

glycosylation by two black dots. **(e)** 1SU4 helix A *i*, *i*+5 mutant. The membrane-embedded

region of helix A is highlighted in yellow. Charged residues are shown as sticks in blue

(R), red (D) and pink (E). L49, A69 and F78 are also shown as sticks to define helix's

subdomains. The membrane is indicated by red (outer) and blue (inner) discontinuous

lines as in (a), and the dashed line indicates the $RCOO^-$ and $RC(NH_2)_2^+$ distance.

Figure S1. Log odds ratios of each pair of amino acids for '*i*, *i*+1' through '*i*, *i*+8' for the

GLOB-MSA dataset. Log odds ratios for the middle core residues of α-helices of at least

17 residues in length in the GLOB-MSA dataset. The rows on the y-axis indicate the first

amino acid in the pair and the columns on the x-axis the second. The residues are ordered

as in Fig. 2.

Figure S2. Log odds ratios of each pair of amino acids for '*i*, *i*+1' through '*i*, *i*+8' for the

GLOB-MSA dataset. Log odds ratios as in S1 but all pairs with P-value >0.05 have been

masked.

Figure S3. Masked log odds ratios of each pair of amino acids for '*i*, *i*+1' through '*i*, *i*+8'

for the TM-MSA dataset. Log odds ratios as in Fig. 1 but all pairs with P-value >0.05 have

been masked.

Figure S4. Salt bridges at pair *i*, *i*+5. Left: helix E from bacterial translocon (PDB ID:

5MG3). Glycine (orange) residue causes a kink facilitating salt bridge interaction between

Asp50 (red) and Arg55 (blue). Right: helix A from a lipid flippase (PDB ID: 6CC4).

Glycine (orange) residue causes a kink facilitating salt bridge interaction between Arg153

(blue) and Glu158 (red). Distances are shown in Ångström.

REFERENCES

652
1. Martínez-Gil L, Saurí A, Marti-Renom MA, Mingarro I (2011) Membrane protein
integration into the endoplasmic reticulum. *FEBS J* **278**: 3846–3858.
2. Baeza-Delgado C, Marti-Renom MA, Mingarro I (2012) Structure-based statistical
analysis of transmembrane helices. *Eur Biophys J* **42**: 199–207.

3. Tsirigos KD, Peters C, Shu N, Käll L, Elofsson A (2015) The TOPCONS web server
for consensus prediction of membrane protein topology and signal peptides. *Nucleic*
Acids Res **43**: W401–W407.
4. Tsirigos KD, Govindarajan S, Bassot C, Västermark Å, Lamb J, Shu N, Elofsson A
(2018) ScienceDirect Topology of membrane proteins — predictions, limitations
and variations. *Curr Opin Struct Biol* **50**: 9–17.
5. Bañó-Polo M, Baeza-Delgado C, Orzáez M, Marti-Renom MA, Abad C, Mingarro I
(2012) Polar/Ionizable Residues in Transmembrane Segments: Effects on Helix-
Helix Packing. *PLoS ONE* **7**: e44263.
6. Marqusee S, Baldwin RL (1987) Helix stabilization by Glu-...Lys+ salt bridges in
short peptides of de novo design. *Proc Natl Acad Sci USA* **84**: 8898–8902.
7. Donald JE, Kulp DW, DeGrado WF (2011) Salt bridges: Geometrically specific,
designable interactions. *Proteins* **79**: 898–915.
8. Mbaye MN, Hou Q, Basu S, Teheux F, Pucci F, Rooman M (2019) A comprehensive
computational study of amino acid interactions in membrane proteins. *Sci Rep* **9**:
12043–14.
9. Baeza-Delgado C, Heijne von G, Marti-Renom MA, Mingarro I (2016) Biological
insertion ofcomputationally designed shorttransmembrane segments. *Sci Rep* 1–9.
10. Chin CN, Heijne von G (2000) Charge pair interactions in a model transmembrane
helix in the ER membrane. *J Mol Biol* **303**: 1–5.
11. Braunger K, Pfeffer S, Shrimal S, Gilmore R, Berninghausen O, Mandon EC, Becker
T, Förster F, Beckmann R (2018) Structural basis for coupling protein transport and
N-glycosylation at the mammalian endoplasmic reticulum. *Science* **360**: 215–219.
12. Hessa T, Kim H, Bihlmaier K, Lundin C, Boekel J, Andersson H, Nilsson I, White
SH, Heijne von G (2005) Recognition of transmembrane helices by the endoplasmic
reticulum translocon. *Nature* **433**: 377–381.
13. Hessa T, Meindl-Beinker NM, Bernsel A, Kim H, Sato Y, Lerch-Bader M, Nilsson I,
White SH, Heijne von G (2007) Molecular code for transmembrane-helix
recognition by the Sec61 translocon. *Nature* **450**: 1026–1030.
14. Bañó-Polo M, Martínez-Gil L, Barrera FN, Mingarro I (2019) Insertion of
Bacteriorhodopsin Helix C Variants into Biological Membranes. *ACS Omega* **5**:
556–560.
15. Tamborero S, Vilar M, Martínez-Gil L, Johnson AE, Mingarro I (2011) Membrane
insertion and topology of the translocating chain-associating membrane protein
(TRAM). *J Mol Biol* **406**: 571–582.
16. Kozma D, Simon I, Tusnády GE (2013) PDBTM: Protein Data Bank of
transmembrane proteins after 8 years. *Nucleic Acids Res* **41**: D524–D529.
17. Andreeva A, Howorth D, Chothia C, Kulesha E, Murzin AG (2013) SCOP2
prototype: a new approach to protein structure mining. *Nucleic Acids Res* **42**:
D310–D314.
18. Andreeva A, Kulesha E, Gough J, Murzin AG (2019) The SCOP database in 2020:
expanded classification of representative family and superfamily domains of known
protein structures. *Nucleic Acids Res* **48**: D376–D382.

19. Illergård K, Kauko A, Elofsson A (2011) Why are polar residues within the membrane core evolutionary conserved? *Proteins* **79**: 79–91.

20. Samanta U, Pal D, Chakrabarti P (1999) Packing of aromatic rings against tryptophan residues in proteins. *Acta Cryst (1999) D55, 1421-1427 [doi:101107/S090744499900726X]* 1–7.

21. Kanada S, Takeguchi Y, Murakami M, Ihara K, Kouyama T (2011) Crystal Structures of an O-Like Blue Form and an Anion-Free Yellow Form of pharaonis Halorhodopsin. *J Mol Biol* **413**: 162–176.

22. Kumar S, Nussinov R (2002) Relationship between Ion Pair Geometries and Electrostatic Strengths in Proteins. *Biophys J* **83**: 1595–1612.

23. Toyoshima C, Nakasako M, Nomura H, Ogawa H (2000) Crystal structure of the calcium pump of sarcoplasmic reticulum at 2.6 Å resolution. *Nature* **405**: 647–655.

24. Kauko A, Hedin LE, Thebaud E, Cristobal S, Elofsson A, Heijne von G (2010) Repositioning of transmembrane alpha-helices during membrane protein folding. *J Mol Biol* **397**: 190–201.

25. Armstrong KM, Baldwin RL (1993) Charged histidine affects alpha-helix stability at all positions in the helix by interacting with the backbone charges. *Proc Natl Acad Sci USA* **90**: 11337–11340.

26. Lin C-Y, Lin L-Y (2018) The conserved basic residues and the charged amino acid residues at the α-helix of the zinc finger motif regulate the nuclear transport activity of triple C2H2 zinc finger proteins. *PLoS ONE* **13**: e0191971–20.

27. Kauko A, Illergård K, Elofsson A (2008) Coils in the Membrane Core Are Conserved and Functionally Important. *J Mol Biol* **380**: 170–180.

28. Giménez-Andrés M, Čopič A, Antonny B (2018) The Many Faces of Amphipathic Helices. *Biomolecules* **8**: 45–14.

29. Walther TH, Ulrich AS (2014) ScienceDirectTransmembrane helix assembly and the role of salt bridges. *Curr Opin Struct Biol* **27**: 63–68.

30. Okamura Y, Fujiwara Y, Sakata S (2015) Gating Mechanisms of Voltage-Gated Proton Channels. *Annu Rev Biochem* **84**: 685–709.

31. Bañó-Polo M, Baeza-Delgado C, Tamborero S, Hazel A, Grau B, Nilsson I, Whitley P, Gumbart JC, Heijne von G, Mingarro I (2018) Transmembrane but not soluble helices fold inside the ribosome tunnel. *Nat Commun* **9**: 5246.

32. Jayasinghe S, Hristova K, White SH (2001) Energetics, stability, and prediction of transmembrane helices. *J Mol Biol* **312**: 927–934.

33. Shurtleff MJ, Itzhak DN, Hussmann JA, Schirle Oakdale NT, Costa EA, Jonikas M, Weibezahn J, Popova KD, Jan CH, Sinitcyn P, et al. (2018) The ER membrane protein complex interacts cotranslationally to enable biogenesis of multipass membrane proteins. *eLife* **7**: 382.

34. Chitwood PJ, Hegde RS (2020) An intramembrane chaperone complex facilitates membrane protein biogenesis. *Nature* **584**: 1–22.

35. Heijne von G (1989) Control of topology and mode of assembly of a polytopic membrane protein by positively charged residues. *Nature* **341**: 456–458.

36. Lerch-Bader M, Lundin C, Kim H, Nilsson I, Heijne von G (2008) Contribution of
positively charged flanking residues to the insertion of transmembrane helices into
the endoplasmic reticulum. **105**: 4127–4132.
37. Hedin LE, Ojemalm K, Bernsel A, Hennerdal A, Illergård K, Enquist K, Kauko A,
Cristobal S, Heijne von G, Lerch-Bader M, et al. (2010) Membrane insertion of
marginally hydrophobic transmembrane helices depends on sequence context. *J
Mol Biol* **396**: 221–229.
38. Bañó-Polo M, Martínez-Gil L, Wallner B, Nieva JL, Elofsson A, Mingarro I (2013)
Charge Pair Interactions in Transmembrane Helices and Turn Propensity of the
Connecting Sequence Promote Helical Hairpin Insertion. *J Mol Biol* **425**: 830–840.
39. Kumar S, Ma B, Tsai CJ, Nussinov R (2000) Electrostatic strengths of salt bridges in
thermophilic and mesophilic glutamate dehydrogenase monomers. *Proteins* **38**:
368–383.
40. Bosshard HR, Marti DN, Jelesarov I (2004) Protein stabilization by salt bridges:
concepts, experimental approaches and clarification of some misunderstandings. *J
Mol Recognit* **17**: 1–16.
41. Fu L, Niu B, Zhu Z, Wu S, Li W (2012) CD-HIT: accelerated for clustering the next-
generation sequencing data. *Bioinformatics* **28**: 3150–3152.
42. Li W, Godzik A (2006) *Cd-hit: a fast program for clustering and comparing large
sets of protein or nucleotide sequences*.
43. Eddy SR (2011) Accelerated Profile HMM Searches. *PLoS Comput Biol* **7**: e1002195.
44. Engelman DM, Steitz TA, Goldman A (1986) Identifying nonpolar transbilayer
helices in amino acid sequences of membrane proteins. *Annu Rev Biophys Biophys
Chem* **15**: 321–353.
766

Figure 1

b)

AA	Gap	Sequence	$\Delta G_{pred.}$
L4/A15	-	AAAAA**L**A**L**AAAAA**L**A**L**AAAA	-0.490
K	-	AAAAA**L**A**L**K**AAAAA**L**A**L**AAAA	1.414
D	-	AAAAA**L**A**L**AAAA**D**LA**L**AAAA	1.389
KD	1	AAAAA**L**A**L**AAA**KD**LA**L**AAAA	3.222
KD	2	AAAAA**L**A**L**AA**K**A**D**LA**L**AAAA	2.927
KD	3	AAAAA**L**A**L**A**K**AA**D**LA**L**AAAA	3.349
KD	4	AAAAA**L**A**L**K**AAA**D**LA**L**AAAA	3.339
KD	5	AAAAA**L**A**K**LAAA**D**LA**L**AAAA	2.980
DD	4	AAAAA**L**A**L**D**AAA**D**LA**L**AAAA	3.548

The positions 8 and 12 are marked above the L4/A15 sequence.

c)

Figure 2

Figura 3

Figure 4

a)

Name	Sequence		Charges	Gap	Å	ΔG$_{pred.}$	ΔG$_{exp.}$
wt	243 254 258 271 VGVTSWGYSFL**D**IVA**K**YIFAFLLLNYLTS		DK	4	3.5	+1.728	−0.243
DD	VGVTSWGYSFL**D**IVA**D**YIFAFLLLNYLTS		DD	4	-	+1.664	+0.112
DA	VGVTSWGYSFL**D**IVA**A**YIFAFLLLNYLTS		DA	4	-	−0.376	−0.880
DK *i, i*+5	VGVTSWGYSFL**D**IVAY**K**IFAFLLLNYLTS		DK	5	-	+1.252	+0.307

Figura 5

a)

b)

Name	Sequence	Gap	(Å)	$\Delta G_{pred.}$	$\Delta G_{exp.}$
L (L49-F78)	49 59 63 78 LWELVIEQFEDLLVRILLLAACISFVLAWF	4	3.0	+1.063	-1.050
S (L49-A69)	LWELVIEQFEDLLVRILLLAA	4	3.0	+4.122	+0.216
ΔG_{pred} TM (L60-F78)	LLVRILLLAACISFVLAWF	-	-	-2.478	-
S *i, i+5* (L49-A69)	LWELVIEQFEDLLVIRLLLAA	5	-	+3.459	+0.060
L *i, i+5* (L49-F78)	LWELVIEQFEDLLVIRLLLAACISFVLAWF	5	-	+0.442	-0.336

c)

d)

e) *i, i+5*

Figure S1

Figure S2

Figure S3

Figure S4

5MG3
helix E

3.1

2.6

6CC4
helix A

3.2

3.6