Rats spontaneously perceive global motion direction of drifting plaids

Giulio Matteucci[1,2,*], Benedetta Zattera[1,3,*], Rosilari Bellacosa Marotti[1], and Davide Zoccolan[1,#]

1. Visual Neuroscience Lab, International School for Advanced Studies (SISSA), Trieste, 34136, Italy

2. Current address: Department of Basic Neurosciences, University of Geneva, Geneva, 1206, Switzerland

3. Current address: Friedrich Miescher Institute for Biomedical Research (FMI), Basel, 4058, Switzerland

*These authors contributed equally to this work

Corresponding author

Davide Zoccolan
International School for Advanced Studies (SISSA)
Via Bonomea, 265   34136 Trieste (TS) ITALY
Office: +39 040 3787 701   Fax: +39 040 3787 702
E-mail: zoccolan@sissa.it

Summary

Computing global motion direction of extended visual objects is a hallmark of primate high-level vision. Although neurons selective for global motion have also been found in mouse visual cortex, we still do not know whether rodents can combine multiple motion signals into global, integrated percepts. Here, we trained two groups of rats to discriminate motion direction of either gratings or plaids (i.e., superpositions of two gratings) and we tested whether these visual patterns were able to prime rat perception consistently with an integrated representation of plaid direction. We found that, depending on the identity of the training stimuli, rats displayed either a shared representation of gratings and plaids based on global motion, or a non-shared one. As shown by computational modeling, these findings are consistent with the existence of a population of pattern-like cells, playing a functional role similar to that demonstrated in primates.

Introduction

Extracting global motion direction of visual objects is crucial to guide behavior in many ethological contexts (Lisberger, 2015; Nishida et al., 2018). Such computation has been widely studied in the dorsal stream of primates, and, in particular, in the middle temporal area (MT) of the macaque (Born and Bradley, 2005). This area is known to receive direct input from neurons in primary visual cortex (V1), which signal the "local" direction of moving elements (e.g., edges) in their small, localized, receptive fields (RF). MT units appear to combine the afferent V1 inputs in such a way to acquire selectivity for the "global" motion direction of visual objects (or patterns) made of multiple local edges (Orban, 2008). The leading hypothesis emerging from the primate literature is that this is achieved by integrating the local motion signals carried by V1 afferents over different spatial positions, directions and frequencies (Adelson and Movshon, 1982; Born and Bradley, 2005; Orban, 2008). Such integration is necessary, because the local output of any V1-like edge detector is intrinsically ambiguous insofar global motion is concerned. Any local output can be generated by infinite combinations of global object directions and speeds. This ambiguity is at the core of what is known, in the psychophysics literature, as the "aperture problem" (Fennema and Thompson, 1979; Wuerger et al., 1996; Marr, 2010): only by combining information from multiple, local, moving-edge detectors, it is possible to infer global motion direction. Achieving a circuit level understanding of such computation remains a key question in visual neuroscience.

Over the past 10 years, the development of a wide array of tools for the dissection of neural circuits (Luo et al., 2008, 2018; Fenno et al., 2011; Chen et al., 2013; Madisen et al., 2015), combined with the demonstration of advanced visual behaviors in mice and rats (Carandini and Churchland, 2013; Zoccolan, 2015), has fostered the use of rodents as model systems to study visual cortical processing. This led some investigators to look for the signature of motion integration in the rodent brain (Juavinett and Callaway, 2015; Muir et al., 2015; Palagina et al., 2017). Inspired by monkey studies (Simoncelli and Heeger, 1998; Smith et al., 2005; Rust et al., 2006; Solomon et al., 2011), they recorded the responses of mouse visual cortical neurons to drifting gratings and coherent plaids − i.e., complex visual patterns made of two overlapping gratings with the same contrast and speed, but moving along independent directions. These stimuli have been widely used to investigate motion integration in human and non-human primates since they enable to clearly distinguish local- from global-based motion responses. A plaid, in fact, has two well-defined local motion components and a global direction that is different from the local ones, while, for a grating,

the local and global directions coincide (Adelson and Movshon, 1982). By employing these

stimuli, mouse visual neurons were classified as "pattern" (i.e., responsive to global motion)

or "component" (i.e., responsive to local motion). The majority of neurons amenable to such

classification fell into the component category, but a small fraction of pattern units was

reported in V1 (Muir et al., 2015; Palagina et al., 2017) and in two extrastriate areas, the

lateromedial (LM) and rostrolateral (RL) areas (Juavinett and Callaway, 2015). However, the

relevance of such tiny population of pattern cells in determining mouse motion perception

was left untested [but see further discussion of (Palagina et al., 2017) below].

Even if a few studies tested the ability of rats and mice to discriminate the dominant

motion direction of random dot kinematograms (Douglas et al., 2006; Stirman et al., 2016;

Marques et al., 2018; Samonds et al., 2018) (RDKs: a task linked to motion integration, since

it requires spatial pooling of local motion cues), no study ever probed the ability of rodents to

integrate into a global motion percept multiple oriented elements moving along different

directions. In addition, all the studies employing RDKs explicitly trained the animals in the

direction discrimination task. Critically, this prevented assessing the spontaneous, purely

perceptual motion integration abilities of naïve rodents. In fact, if feedback (e.g., reward) is

provided to an animal about the outcome of his perceptual choices, it may learn to associate a

given stimulus to a given response category without necessarily perceiving the sensory

quality that the investigator intends to associate to that category (Tafazoli et al., 2012;

Zoccolan, 2015; Zoccolan and Di Filippo, 2018). It would thus be impossible to infer from

such kind of experiments whether the choices of the animal are driven by a general-purpose

representation of global motion direction at the level of visual cortex (as the one found in

monkey MT), or by the ability to learn abstract representations of arbitrary stimulus

categories in higher-order association areas involved in perceptual decision making, such as

posterior parietal cortex (PPC) (Raposo et al., 2014; Licata et al., 2017; Nikbakht et al., 2018;

Odoemene et al., 2018).

Only in (Palagina et al., 2017) this confound was avoided, by monitoring the direction

of the optokinetic nystagmus (OKN) while presenting plaids to naïve mice. The resulting

bimodal distribution of OKN movements tracking both the local-components and the overall

global directions of the stimulus is suggestive of mouse ability to spontaneously perceive

global motion. However, this conclusion is affected by the intrinsic limitation of OKN-based

experiments to probe cortical processing. In fact, OKN is a reflexive phenomenon that is

known to be largely controlled by subcortical structures (Smith and Harris, 1991; Harvey et

al., 1997; Stahl, 2004), while motion perception in direction discrimination tasks has been
shown to rely heavily on visual cortex (Newsome and Pare, 1988; Salzman et al., 1990;
Marques et al., 2018). As a result, it remains unclear whether rodents spontaneously combine
independent motion signals (as those found in plaids) into integrated percepts of global
motion direction.

Our study was designed to address this question and provide a thorough
psychophysical assessment of rat ability to spontaneously perceive the global direction of
drifting plaids. To this aim, we relied on a visual priming paradigm (Tafazoli et al., 2012;
Zoccolan and Di Filippo, 2018) that allowed measuring the perceptual similarity between
gratings and plaids with the same global direction, without explicitly training the animals to
associate these stimuli to the same response category.

Results

A visual priming paradigm to probe spontaneous perception of global motion direction

In the first phase of the study (the *training phase*), a group of 11 male Long-Evans rats
(referred to as the G group in what follows) was trained to discriminate leftward- from
rightward-drifting gratings (Figure 1A, left), using the high-throughput behavioral rig
described in (Zoccolan, 2015) and previously employed in several investigations of rat object
recognition (Zoccolan et al., 2009; Tafazoli et al., 2012; Alemi-Neissi et al., 2013; Rosselli et
al., 2015; Djurdjevic et al., 2018). Another group of 10 rats (referred to as the P group in
what follows) was trained to discriminate leftward- from rightward-drifting plaids with a
cross-angle of 120º – i.e. composite patterns made of the superposition of two gratings
drifting along directions that were 120º apart (Figure 1B, left). Rats that maintained at least a
criterion performance of 75% correct choices over a four consecutive day period were moved
to the second phase of the study to be tested with the priming paradigm (all rats starting the
training phase managed to reach such criterion).

Figure 1. Visual stimuli and discrimination tasks. (**A**) Discriminations tasks administered to the rats of the *grating* (G) *group*. During both the training (left) and priming (right) phases, a trial started when an animal triggered the central port of a 3-way licking sensor (see Materials and methods). During the training phase, a target grating (drifting either leftward or rightward) was presented immediately at the onset of each trial. To receive the reward, a rat had to lick the lateral port corresponding to the direction of motion of the presented target. During the priming phase, the structure of the task was similar, but at the onset of each trial, a drifting prime stimulus was presented for 75 ms, followed by a 75 ms blank screen and finally by the presentation of the target grating (lasting 750 ms). The prime stimulus could move in 19 possible directions (from 0° = right to 180° = left) and could be either a grating or a plaid. (**B**) Discriminations tasks administered to the rats of the *plaid* (P) *group*. The structure of the task was the same as in A, with the key difference that the target stimulus was a plaid instead of a grating (again drifting either leftward or rightward). In this case, the reward was delivered when a rat licked the lateral port corresponding to the global motion direction of the plaid.

In the priming paradigm, the "target" stimulus remained the same as in the training phase (i.e., G and P rats still had to report the direction of drifting gratings and plaids, respectively), but now the presentation of the target was preceded by the brief presentation (75 ms) of either a grating or a plaid drifting along one of 19 possible directions (from 0° = rightward to 180° = leftward, in steps of 10°; Figure 1 A-B, right). The identity (i.e., either grating or plaid) and the motion direction of such "prime" stimulus was randomly selected in each trial. In what follows, we refer to trials in which the identities of prime and target coincide as *identity-priming* condition (e.g., a grating prime stimulus followed by a grating target stimulus), whereas we refer to the opposite case as *cross-priming* condition (e.g., a plaid prime stimulus followed by a grating target stimulus). Critically, the rats kept receiving feedback (i.e., either reward or a time-out period in case of correct and incorrect choices,

respectively) only about the correctness of their responses to the target stimuli. The identity and direction of the prime stimuli were never paired to either the leftward or rightward response categories. As such, the extent to which the prime stimuli were able to affect the choices of the rats was purely due to the spontaneous, perceived similarity between primes and targets. Assessing such similarity was the ultimate goal of our experiments, since it allowed understanding whether, in the cross-priming condition, the perceived motion direction of the plaids was the global one. Based on previous motion adaptation studies in humans (Kanai and Verstraten, 2005) and on former priming/masking experiments carried out in rats (Tafazoli et al., 2012; Alwis et al., 2016; Dell et al., 2018, 2019), we choose the duration of the prime and of the inter-stimulus interval (ISI) between prime and target (75 ms each) in the attempt of inducing a strong priming effect – i.e., in the attempt of biasing rats' responses to the target towards the motion direction of the prime.

In what follows, we will report the accuracy of rats at discriminating the motion direction of a target stimulus as a function of its angular distance from the direction of the prime stimulus that preceded it. As such, the resulting *priming curve* will range between two extremes: 1) the case in which the direction of the prime and target stimuli was the same (e.g., both drifting rightward), referred to as *coherent prime* condition; and 2) the case in which the prime and target stimuli had opposite directions (e.g., the prime drifting rightward and the target drifting leftward), referred to as *incoherent prime* condition. The case in which the prime stimulus drifted vertically (midpoint of the priming curve) will be referred to as *neutral prime* condition. This terminology will be used for both the identity- and cross-priming conditions and for both the P and G groups.

The first step in our analysis was to verify whether the prime stimuli were indeed capable of biasing rat choices depending on the level of coherence between their direction and that of the target stimuli, in case of the identity-priming condition. This was the case for 10 out of 11 rats of the G group and 9 out of 10 rats of the P group, which were therefore included in the subsequent analysis (see Materials and Methods). For the G group, rat group average accuracy at discriminating the motion direction of the target gratings was strongly modulated, as function of the direction of the prime grating stimulus (Figure 2A, black curve). In the coherent prime condition, rat classification of target direction was facilitated, as compared to the reference neutral prime condition (dashed line). Conversely, in the incoherent prime condition, rat classification accuracy was substantially lowered. Overall, the identity-priming curve was approximately sigmoidal. Even for the P group, in the identity-

priming condition, a strong priming was observed (Figure 2B, black curve). Also in this case,
the magnitude and sign of the priming effect depended on the similarity between the
directions of the prime and target plaid stimuli, resulting in an approximately monotonic drop
of rat classification accuracy from the coherent priming condition towards the incoherent one.

The strong modulation of rat classification accuracy in the identity-priming condition
has two important implications. First, it demonstrates the effectiveness of the priming
paradigm in capturing the perceived similarity between the learned (horizontal) motion
directions of the target stimuli and the untrained (previously unseen) directions of the prime
stimuli, in the case in which targets and primes are either both gratings or both plaids (i.e.
identity-priming condition). Second, and more importantly, it serves as a reference against
which to compare the modulation of the accuracy observed in the cross-priming condition. In
fact, the degree of similarity between the priming curves measured in the identity- and cross-
priming conditions (i.e., between the black and gray curves in Figure 2) allows answering
two key questions about the neuronal representations of the grating and plaid stimuli.

Figure 2. Identity- and cross-priming curves. (**A**) The identity-priming curve (black) reports the mean
accuracy of the rats of the G group at discriminating the leftward from the rightward drifting gratings,
as a function of the difference between the direction of these target stimuli and the gratings that were
used as prime stimuli. The cross-priming curve (gray) reports rat mean accuracy when the plaids were
used instead as prime stimuli. The shaded areas are 95% confidence intervals obtained by bootstrap
(see Materials and methods). The dashed line shows the accuracy achieved with the neutral identity-
prime stimulus (i.e., the grating drifting upward). The inset quantifies the absolute magnitude of
priming obtained in the two cases, as the mean absolute difference between the four leftmost (and
rightmost) points in a curve and the point corresponding to the neutral condition. The error bars show

95% confidence intervals obtained by bootstrap (see Materials and methods). (**B**) Same as in A, but
for the rats of the P group. That is, in this case, the curves report rat mean accuracy at discriminating
the leftward from the rightward drifting plaids. The identity-priming curve (black) refers to trials where
the plaids were used as prime stimuli, while the cross-priming curve (gray) refers to trials where the
gratings were used as primes.
215

Evidence of a shared representation of gratings' and plaids' motion direction

The first question is whether these representations are "shared" (i.e., the same neural

population encodes both gratings and plaids) or "non-shared" (i.e., independent, non-

overlapping populations represent the two types of stimuli). In the former case, we would

expect a substantial modulation of rat classification accuracy also in the cross-priming

condition, possibly as large as in the identity-priming condition. In the latter case, no cross-

priming effect would be observable, given the independence of the two populations

underlying the perception of gratings and plaids. In case of shared representations, then a

second crucial question could be answered by comparing the shapes of the priming curves

obtained in the identity- and cross-priming conditions. If rats spontaneously represented the

stimuli according to their global direction, then the priming curves produced by gratings and

plaids would be very similar, given that, by construction, the global direction of the prime

stimuli was the same for matching grating and plaid primes. If the opposite were true, then

the priming curves produced by gratings and plaids would be different.

In the G group (Figure 2A), the cross-priming curve (gray) was almost flat and

significantly different from the identity-priming curve (black), as shown by the non-

overlapping 95% confidence intervals (shaded areas). The priming magnitude produced by

the plaid stimuli was found to be significantly smaller than that produced by the gratings

(Figure 2A, inset: gray vs. black bar; non-overlapping 95% confidence intervals). This

inability of the plaids (when used as prime stimuli) to affect the discrimination (hence the

perception) of the target gratings strongly suggests that, for the rats of the G group, the

neuronal population representing the direction of the gratings was largely inactive during the

presentation of the plaids.

By contrast, in the P group, the cross-priming and identity-priming curves largely

overlapped (Figure 2B: gray vs. black curve). As a result, the priming magnitude observed in

the two conditions was equally large (Figure 2B, inset: gray vs. black bar; overlapping 95%

confidence intervals). In other words, the gratings and plaids (when used as prime stimuli)

were equally effective at biasing rat discrimination of the target plaids. This indicates that the

same neuronal population was active during the presentation of both kinds of stimuli, as expected in case of a shared representation of gratings and plaids. More importantly, the priming produced by gratings and plaids was not only equivalent in terms of overall magnitude, but it was also very similar for matching (global) directions of the two stimuli (i.e., overlapping curves in Figure 2B). This strongly suggests that the representation of plaids in rats trained to discriminate such stimuli (P group) is not only shared with the representation of gratings, but is structured in such a way to encode plaids' global motion direction.

Rat perception of drifting plaids is more consistent with a pattern- than a component-based representation of motion signals

To test more formally the extent to which the priming curves observed for the P group are consistent with a representation of global motion direction of the plaid stimuli, we performed a decoding analysis using linear classifiers. We simulated two alternative scenarios: a representation made of pattern cells (i.e., neurons tuned to the global direction of the plaids) and a representation made of component cells (i.e., neurons tuned to the local direction of the constituent gratings of the plaids).

Figure 3. Simulating the discrimination accuracy afforded by component- and a pattern-based representations. (**A**) Tuning curves (rows) for a simulated population of 24 components cells, whose preferred directions homogeneously spanned the [0º 360º] range, starting from 0º (first row), up to 345º (last row), in steps of 15º. The responses of the units are shown as a function of 24 possible directions (columns) of the grating (top) and plaid (bottom) stimuli. The gray scale bar indicates the intensity of the response. (**B**) Same as in A but for a population of 24 simulated pattern cells. (**C**) Weights assigned to each of the 24 component cells in A (the x axis reports the preferred direction of each unit) by a logistic classifier that was trained to discriminate a 0º- (rightward) from a 180º-drifting (leftward) plaid. (**D**) Fraction of times that the logistic classifier fed with the component-based representation (whose weights are shown in C) classified the test stimuli as drifting rightward (i.e., at 0º). The test stimuli were either plaids (solid black curve) or gratings (dashed gray curve) spanning the whole [0º 360º] range of directions. (**E**) Weights assigned to each of the 24 pattern cells in B by a logistic classifier that was trained to discriminate a 0º- (rightward) from a 180º-drifting (leftward) plaid. (**F**) Same as in D, but for the logistic classifier fed with the pattern-based representation (whose weights are shown in E). Figure 3–figure supplement 1 reports an extended version of these simulations, where a range of widths of the tuning curves (from very broad to very narrow) was tested.

In both cases, we simulated a population of 24 units, whose preferred directions homogeneously spanned the [0º 360º] range (i.e., one unit tuned to every direction in steps of 15º; Figure 3A-B). The tuning curves of the simulated units were defined by Von Mises functions (Kohn and Movshon, 2004; Matteucci et al., 2019), which are the circular analog of Gaussian functions and whose width is controlled by the parameter k, which is equivalent to the inverse of the variance of a Gaussian distribution (Berens, 2009) (see Methods for details). In our simulations, k was set to 7, which roughly corresponds to a peak of the tuning curve with a full width at half maximum (FWHM) of about 50º (see grayscale matrix in Figure 3A-B). By construction, for a component cell, the response to the drifting plaids as a function of their direction featured two peaks, separated by 120º (Figure 3A, bottom). These peaks correspond to the global directions of the plaid that bring one of the constituent gratings to be aligned to the preferred (local) direction of the unit (i.e., the one shown in Figure 3A, top). Conversely, the tuning curves of the simulated pattern cells featured a single peak, corresponding the global direction of the stimulus (no matter whether grating or plaid) they were tuned to (Figure 3B) (see Methods for further details).

We used these populations to simulate how a rat trained to discriminate leftward from rightward drifting plaids (i.e., same task of the animals in the P group; see Figure 1B) would spontaneously categorize the novel, previously unseen gratings and plaids used as prime

stimuli, under the assumption of either a purely pattern-based or a purely component-based
representation. Rat perceptual decisions were simulated by training a logistic classifier to
discriminate the leftward from the rightward drifting plaids, based on their representation in
the simulated neural population space. As a result of the training, each classifier learned to
rely on a specific subset of the units of the representation, as shown by the weights'
distribution of Figure 3C and E. We then tested how each classifier categorized the sets of
drifting plaids and gratings used in the identity- and cross-priming experiments. This yielded
generalization curves (Figure 3D and F) that, in the [0º 180º] range, are equivalent to the
identity- and cross-priming curves of Figure 2B.

The shape of the generalization curves can be understood by considering the
corresponding weights' distribution resulting from training with the 0º- and 180º-drifting
plaids. In the case of the component representation, these training stimuli activated mostly the
units tuned for directions at ±60º with respect to the direction of the stimuli themselves. As a
result, the classifier learned to assign large, positive weights to the units tuned at ~60º and
~300º (i.e., those providing the strongest evidence of a rightward, 0º-drifting plaid), and
large, negative weights to the units tuned at ~120º and ~240º (i.e., those providing the
strongest evidence of a leftward, 180º-drifting plaid) – hence, the twin positive and negative
peaks in the weights' distribution (Figure 3C).

When the classifier was presented with test plaids spanning the whole [0º 360º] range
of directions, the proportion of rightward choices (solid curve in Figure 3D) was obviously
maximal (i.e., 100%) at 0º (since this was the rightward training direction), but then dropped
quickly, becoming minimal (i.e., 0%) for a plaid drifting at 60º. In fact, for this stimulus, one
of the constituent gratings activated the units tuned at ~0º, where the weights were close to
zero (see Figure 3C), while the other grating activated the units tuned at ~120º, where the
weights were large but negative, since those units signaled the presence of the leftward, 180º-
drifting plaid. When the direction of the test plaid further increased, the proportion of
rightward choices grew again, reaching a new positive peak at 120º. This was because one of
the constituent gratings of the plaid activated the units tuned at 60º (with large, positive
weights), while the other grating activated those tuned at 180º (where the weights were close
to zero). The generalization curve reached again another minimum when the test plaid drifted
at 180º (the leftward training direction), with the constituent gratings that activated the units
tuned at 120º and 240º, with large, negative weights. As shown in Figure 3D, a mirror trend
was produced for plaids drifting along directions larger than 180º.

As for the generalization curve obtained when feeding to the classifier isolated
drifting gratings (dashed, gray curve in Figure 3D, which is equivalent to the cross-priming
curve of Figure 2B), it largely followed the shape of the weights' distribution. This can be
easily understood, given that a grating provided as much evidence of rightward (leftward)
motion as close its direction was to the positive (negative) peaks of the weights' distribution.
As a result, the generalization curves produced by plaids and gratings (solid vs. dashed curve
in Figure 3D) followed completely different trends, with no overlap. In addition, none of
them matched the monotonic decrease observed for the corresponding identity- and cross-
priming curves of Figure 2B (solid vs. dashed curve). Together, this strongly suggests that a
representation made of component cells is incompatible with the priming curves measured for
the rats of the P group.

A very different result was found for the simulated representation of pattern cells. In
this case, the weights' distribution had a large, positive peak at 0º, corresponding to the units
coding for the rightward motion, as well as a large, negative peak at 180º, corresponding to
the units coding for the leftward motion (Figure 3E). Given that, by construction, the pattern
cells responded in the same way to isolated gratings and plaids having the same (global)
direction, the generalization curves corresponding to the identity- and cross-priming
conditions fully overlapped (Figure 3F, black, solid line vs. gray, dashed line). More
importantly, they both followed a monotonic, decreasing trend in the [0º 180º] range, which
largely trailed the shape of the weights' distribution. Crucially, these trends were highly
consistent with those observed for the priming curves of Figure 2B. This shows that,
differently from the case of the component representation, a representation made of pattern
cells would support the identity- and cross-priming curves obtained for the rats of the P
group.

To verify the extent to which this conclusion depended on the width of the tuning
curves of the simulated units, we allowed the parameter k (that defines the width of the Von
Mises functions) to range from 0.5 to 11. Examples of the resulting tuning curves obtained
for the simulated component cells are shown in Figure 3–figure supplement 1A. At very low
k, as a result of the increased broadness of the tuning, the two peaks in the response to the
plaid stimuli (solid curves) tended to merge. This translated in a broadening of the weights'
distribution of the logistic classifier (Figure 3–figure supplement 1B) and in a concomitant
increase of the similarity between the generalization curves obtained for gratings and plaids
(Figure 3–figure supplement 1C; dashed vs. solid curves). To quantify this similarity, we

computed the absolute differences between the proportions of rightward choices in the two generalization curves across all tested directions and we averaged them. The resulting average difference was large (as already shown in Figure 3D) for $k > 5$ but became negligible for $k < 3$ (Figure 3–figure supplement 1D). This corresponds to peaks in the tuning curves with FWHM ~ 80° or wider (see Figure 3–figure supplement 1A). Therefore, a population of component cells with such a broad tuning could in principle account for the overlapping identity- and cross-priming curves obtained for the P group in our experiments (Figure 2B).

To understand whether this is a plausible scenario, one would need to measure the width of tuning of component cells in rodent visual cortex. Unfortunately, the few studies investigating component and pattern cells in mouse visual cortex did not perform a statistical characterization of the sharpness of their orientation tuning. This prevented our simulated units from quantitatively matching a physiological ground-truth in terms of orientation tuning. Nevertheless, the tuning curves of the example component cells reported in rodent studies investigating this topic feature such sharp and narrow peaks over the orientation axis to decisively fall in the regime corresponding to $k \gg 5$ in our simulations (i.e., with FWHM $\ll$ 60°) (Juavinett and Callaway, 2015; Palagina et al., 2017). A similar conclusion is supported by the general tendency of orientation tuning to be quite sharp in mouse and rat visual cortex (Girman et al., 1999; Niell and Stryker, 2008; Gao et al., 2010; Marshel et al., 2011; Vermaercke et al., 2014; Ghodrati et al., 2016; Matteucci et al., 2019; Matteucci and Zoccolan, 2020). As such, the most likely hypothesis is that the stimulus representation underlying the priming curves of the P group (Figure 2B) consisted of pattern, rather than component cells. Further neurophysiological studies will be necessary to verify this hypothesis – e.g., by directly testing, as done here in our simulations, the extent to which populations of recorded component and pattern cells yield direction classification curves that are consistent with the priming curves measured in our experiments.

Discussion

The results presented in Figure 2B show that rats are capable of spontaneously combining local motion cues into integrated percepts of global motion direction of a complex visual pattern. Our simulations show that this ability is consistent with a representation of global motion signals, as the one provided by pattern cells (Figure 3 and Figure 3–figure supplement 1). As such, our findings strongly suggest that rats are capable of processing motion stimuli

by relying on pattern-like representations. This, in turn, establishes a candidate perceptual

correlate of the neuronal selectivity for global motion previously reported across rodent

visual cortical areas (Juavinett and Callaway, 2015; Muir et al., 2015; Palagina et al., 2017),

suggesting that, despite being a tiny fraction of the overall visual cortical population, rodent

pattern cells do play an important (possibly preferential) role in the processing of visual

motion. Critically, such motion integration ability only emerged in rats trained to discriminate

the plaid stimuli (P group). The animals trained in the grating discrimination task (G group)

were instead virtually insensitive to the plaids (Figure 2A), indicating that the neuronal

population they relied upon to encode the gratings failed to effectively represent the plaids. In

the following sections we provide two tentative explanations of such observations based on

different sets of assumptions. Simulations of component- and pattern-based representations

and linear read-out mechanisms (similar to those shown in Figure 3) allowed testing the

ability of such assumptions to possibly explain our findings.

Simulating a scenario where training with either gratings or plaids leads to a decoding

pool enriched of either component or pattern cells.

The first explanation rests on the hypothesis that the specific discrimination being

reinforced during the training biased the recruitment of the visual cortical pool that was read

out by downstream decision neurons towards either a more component-enriched or a more

pattern-enriched cell population (referred to as the "decoding pool" in what follows). Such a

task-dependent selection of the decoding pool may stem from differences between

component and pattern cells in terms of two key properties: i) their relative proportion within

visual cortex; and ii) the intensity of their responses to gratings and plaids.

With regard to the first property, although the number of studies investigating motion

integration in rodent visual cortical areas is still limited (Juavinett and Callaway, 2015; Muir

et al., 2015; Palagina et al., 2017), a finding that is consistent among them is that component

cells by far outnumber pattern cells, with the former being ~25% of direction-tuned units and

the latter being about ~5% (with the rest being unclassified). With respect to the second

property, a well-known phenomenon likely affecting the strength of grating and plaid

responses is cross-orientation suppression (DeAngelis et al., 1992; Priebe and Ferster, 2006;

Li et al., 2006; Busse et al., 2009), which has been recently documented also in rodents (Self

et al., 2014; Alwis et al., 2016; Samonds et al., 2017). It consists in a reduction of the

response to the preferred oriented stimulus when this is presented together with other stimuli

with different orientations. This means that cross-orientation suppression could, in principle, lead to diminished responses to plaids as compared to gratings. However, most studies documenting this phenomenon mainly targeted V1, without distinguishing between pattern and component cells. In addition, when focusing on studies of motion integration in rodents (Juavinett and Callaway, 2015; Muir et al., 2015; Palagina et al., 2017), it was difficult to draw any insight about the incidence of cross-orientation suppression in pattern and component cells, because of methodological limitations (e.g., tuning curves normalization and calcium imaging lacking spike-count-level information). This is why we turned to the primate literature, finding hints of cross-orientation suppression being stronger in component cells than in pattern cells (Smith et al., 2005; Rust et al., 2006).

Our hypothesis is that the combined effect of component cells being more numerous than pattern cells, but also more sensitive to cross-orientation suppression, led the decoding pool to become dominated by component cells in the case of training with gratings (G group) and by pattern cells in the case of training with plaids (P group). In turn, this would qualitatively explain the difference in terms of cross-priming found for the rats of the G and P groups (Figure 2), as graphically illustrated in Figure 4A, B. Briefly, in the case of training with gratings (Figure 4A, top), these stimuli would strongly activate both pattern (orange) and component (green) cells (filled and unfilled neurons indicate, respectively, responsive and unresponsive units). As such, the selection of the neuronal decoding pool by a downstream decision neuron would not be biased towards either class of neurons (because of no differences in their responsiveness). However, since the component cells substantially outnumber the pattern cells, the decoding pool would contain more of the former (black-boundary neurons) than of the latter (red-boundary neurons). This follows from the assumption that the selection of the decoding pool is the result of a random sampling from the subpopulation of all responsive units. This, in turn, is consistent with the idea that such recruiting process is mediated by some form of plasticity similar to the reward-gated Hebbian learning reviewed in (Roelfsema and Holtmaat, 2018). In the priming experiment, all the neurons in the decoding pool would be active during presentation of the grating primes (Figure 4A, middle), leading to the strong identity priming observed in Figure 2A (solid curve). By contrast, most component cells in the pool would be strongly depressed by cross-orientation suppression during presentation of the plaid primes (Figure 4A, bottom; empty black-boundary neurons), and the few pattern cells in the pool (orange-filled, red-boundary neurons), although active, would not be enough to affect the perception of the target stimulus.

461   Hence, the lack of cross-priming in the group of rats trained with gratings (Figure 2A, gray

462   curve).

Figure 4. Simulating a scenario where training with either gratings or plaids leads to a decoding pool enriched of either component or pattern cells. (**A**) Cartoon illustrating how training with gratings would lead to the selection of a decoding pool enriched of component cells, under the assumption that: 1) component cells are more numerous than pattern cells; and 2) cross-orientation suppression is stronger for component than pattern cells. The sketched pyramidal neurons represent either component (green) or pattern (orange) cells. Filled and unfilled neurons indicate, respectively, responsive and unresponsive units. Black- and red-boundary neurons indicate the cells that the decision neuron learned to rely upon to discriminate the rightward- from the leftward-drifting gratings (i.e., the cells in the decoding pool). The grating stimuli activate equally strongly the component and pattern cells, but, being the former more numerous, the decoding pool is mainly composed of component cells (top). When tested with gratings, the cells in the decoding pool are strongly activated (middle), but, when they are tested with plaids, they are only weakly activated (bottom), because of cross-orientation suppression. (**B**) Cartoon illustrating how training with plaids would lead to the selection of a decoding pool enriched of pattern cells. Because of cross-orientation suppression, component cells are weakly activated by plaids, thus the decoding pool is enriched of pattern cells (top). As a result, the decoding pool is strongly activated when presented with both plaids (middle) or gratings (bottom). (**C**) Top: weights assigned to a mixed population of component (green) and pattern (orange) cells, as a function of their preferred direction, by a logistic classifier that was trained to discriminate a 0°- (rightward) from a 180°-drifting (leftward) grating. Bottom: The proportion of component (green) and pattern cells (orange), initially set to, respectively, 80% and 20% of the total, remained unchanged when weights with increasingly larger magnitude were progressively set to zero (i.e., pruned from the decoding pool). (**D**) Top: same as in C (top) but for a logistic classifier trained to discriminate a 0°- (rightward) from a 180°-drifting (leftward) plaid. Because of cross-orientation suppression, the classifier learned to assign weights with lower magnitude to the component cells. Bottom: as a result, when the input connections to the classifier were made increasingly sparser (by pruning weights with progressively larger magnitude), the proportion of pattern cells in the decoding pool (orange) reached and overtook the proportion of component cells (green).

In the case of training with plaids (Figure 4B, top), these stimuli would strongly activate the population of pattern cells (orange-filled neurons) but only weakly excite the component cells (green-filled neurons), because of cross-orientation suppression. This would lead to a strong bias towards the recruitment of pattern cells in the decoding pool (red-boundary neurons) by a downstream decision neuron. When tested with prime stimuli, such pattern-enriched pool would respond with a similar intensity to both kinds of primes (grating and plaids; Figure 4B, middle and bottom panels respectively), leading to similar priming curves in both identity- and cross-priming conditions, as shown in Figure 2B (solid vs. gray curve). This would account for the shared representation of global motion direction observed for rats trained with plaids.

To quantitatively test whether the mechanistic account hypothesized in the previous paragraphs could explain our findings, we built a computational model where a decision neuron had to select his afferent units from a mixed population of component and pattern cells, in order to discriminate the direction of either drifting gratings or plaids. The relative proportion of simulated component (80%) and pattern (20%) cells roughly matched that

found, on average, across rodent visual areas V1, LM and RL (Juavinett and Callaway, 2015; Muir et al., 2015; Palagina et al., 2017). Each unit was simulated as a Poisson spiking neuron, whose average firing rate as a function of direction was defined by a von Mises function (Kohn and Movshon, 2004; Matteucci et al., 2019) having a peak that was randomly sampled across the [0º 350º] direction axis (see Materials and methods for details).

In addition, to simulate cross-orientation suppression in component cells, their peak responses to plaids were set to half their peak responses to gratings. On the other hand, pattern cells were assumed to be unaffected by cross-orientation suppression. As for the simulation of Figure 3, the decision neuron consisted in a logistic classifier that was trained to discriminate between 0º- and 180º-drifting stimuli (i.e., leftward vs. rightward motion). This classifier simply performed a weighted sum over the inputs provided by all the simulated component and pattern cells, followed by the application of a sigmoidal nonlinearity. As a result of training, the weights became adjusted in such a way to maximize the classification accuracy (eventually attaining 100% in such a simple motion discrimination task).

When trained in the grating discrimination task (to simulate the training regime of the G group), the classifier learned to rely more heavily on those motion detectors (no matter whether component or pattern cells) whose preferred direction was close to that of the training stimuli. This is shown in Figure 4C (top), where the weights of the simulated units as a function of their preferred direction are reported. For both the component (green dots) and pattern (orange dots) cell populations, the most informative neurons were those with preferred direction close to 0º (rightward-drifting gratings) and 180º (leftward-drifting gratings), while the units tuned around the vertical drift direction (i.e., 90º and 270º) were assigned close-to-zero weights. Given these weight distributions, we simulated different degrees of sparsity constraints on the connectivity of the decision neuron. This was done by pruning input connections with progressively larger weight magnitude, thus gradually reducing the size of the decoding pool. However, given the closely matching weight magnitudes of the component and pattern cells, the level of pruning did not alter their relative proportion in the decoding pool – this proportion remained the same as in the original population (i.e., 80% vs. 20%; Figure 4C, bottom). This confirmed the intuition (see Figure 4A) that training with gratings, even in presence of asymmetric cross-orientation suppression (i.e., affecting component but not pattern cells), leads to a decoding pool that simply reflects the proportion of component and pattern cells of the overall cortical population and, as such, is component-enriched.

A very different scenario emerged when we simulated the training of the P group in

the plaid discrimination task (see Figure 4D). The weight distribution learned by the classifier

for the pattern cell population (top, orange dots) was the same as in the grating discrimination

task. This is because our simulated pattern cells were by definition immune to cross-

orientation suppression. On the other hand, the weight distribution for the component cell

population (green dots) was very different from that obtained in the grating discrimination

task. First, since these cells responded to the components of the plaids (i.e., their constituent

gratings), the classifier learned to assign larger weights to those units with preferred direction

at ±60º (i.e., half plaid cross-angle) with respect to the global directions of the leftward- and

rightward-drifting plaids (same as for the simulations shown in Figure 3C). In addition,

because the simulated components cells were affected by cross-orientation suppression, the

classifier learned to rely less on component than on pattern cells, thus assigning lower

weights to the former than to the latter (compare the absolute height of the peaks in the

distributions of green and orange dots in Figure 4D, top). As a result, when the sparsity of the

connections to the decision neuron was increased, by leaving in the decoding pool only inputs

with progressively larger weight magnitude, the proportions of component and pattern cells

in the pool followed opposite, mirror trends (Figure 4D, bottom). While the former decreased

(green curve), the latter increased (orange curve), eventually leading to a reversal of the

fraction of component and pattern cells in the decoding pool when the decision neuron was

pruned of 90% of its potential inputs. This confirmed the intuition (see Figure 4B) that cross-

orientation suppression can, at least in principle, strongly bias the recruitment of pattern cells

in the decoding pool, under the constraint of sparse connectivity to the decision neurons, thus

leading to a pattern-enriched representation despite the prevalence of component cells in the

overall cortical population. As a control experiment, we repeated the whole simulation

without any cross-orientation suppression. As expected, this yielded to a component-enriched

representation for both the simulated G and P groups (data not shown).

Simulating a scenario where a decision neuron reading out a pattern representation

fails to be activated by plaids, if originally trained to discriminate drifting gratings.

Despite the plausibility of the mechanism illustrated in the previous section, the key

assumption of having weaker cross-orientation suppression on pattern cells than on

component cells rests on hints from the primate literature only and has never been directly

demonstrated in rodents. In addition, in our simulations, the decision neurons drew their

inputs from a mixed population of pattern and component cells. That is, we assumed the
decision neuron to pool from different stages of a putative motion processing hierarchy.
Considering primates, this assumption is clearly at odd with the higher functional and
anatomical rank of MT, with respect to V1, along the dorsal stream (Born and Bradley, 2005;
Orban, 2008). In the case of rodents, it is still unclear whether a similar hierarchy of dorsal
areas exists, but at least one study has reported extra-striate area RL to be richer of pattern
cells than V1 (Juavinett and Callaway, 2015). In addition, although an anatomical hierarchy
of motion processing stages as structured as the one found in primates may not exist in
rodents, pattern cells distributed over striate and extrastriate cortex could still sit above
component cells along the functional motion processing chain. In fact, pattern cells, given
their ability to encode motion direction in a shape-independent way, would behave as better
motion detectors than component cells in virtually any visually guided behavior. As such,
motion information could still be preferentially routed from visual cortical areas (e.g., V1,
RL, LM) to higher-order decision centers (i.e., frontal and parietal cortices) mainly via
pattern cells. This possibility is consistent with the well-established ability of distinct
subpopulations, within mouse V1, to selectively make (receive) target-specific projections
with (from) downstream (upstream) areas (Gao et al., 2010; Matsui and Ohki, 2013;
Glickfeld et al., 2013; Ji et al., 2015; D'Souza et al., 2019). In addition, it is consistent with a
very recent study showing that multiple visual areas (V1, LM and AL) are causally involved
in coding stimulus orientation and contrast (Jin and Glickfeld, 2020) – a demonstration that
the representation of key visual features is distributed among neuronal populations within
distinct visual areas.

Figure 5. Simulating a scenario where a decision neuron reading out a population of pattern cells fails to be strongly activated by drifting plaids, when originally trained to discriminate drifting gratings. (**A**) L2 norms of the population vectors fed as inputs to a decision neuron (simulated as a logistic classifier with L2 regularization) that was trained to discriminate either a 0°- from 180°-drifting grating (black) or a 0°- from 180°-drifting plaid (gray). The classifier received inputs from a population of pattern cells. Each bar reports the mean of the norms of the input populations vectors produced by the 0°- and 180°-drifting stimuli. (**B**) Weights assigned by the logistic classifier to the population of pattern cells as a function of their preferred direction, in case of training with gratings (top) and plaids (bottom). (**C**) L2 norms of the weights shown in B. (**D**) Weighted inputs to the logistic classifier, originally trained to discriminate either gratings (black curves) or plaids (gray curves), when the population of simulated pattern cells was presented with drifting stimuli, homogeneously spanning the [0° 360°] range of directions. Such stimuli were either of the same kind (solid curves) or of a different kind (dashed curves) of those used during training. For instance, a classifier originally trained with plaids was tested with either plaids (gray, solid curve) or gratings (gray, dashed curve).

In light of these considerations, an alternative scenario to interpret our findings is one where: i) decision neurons receive inputs mainly from pattern cells, regardless of the motion

discrimination task to be performed (i.e., the decoding pool is, by "construction", pattern

enriched); and ii) pattern cells are as sensitive to cross-orientation suppression as component

cells are. The question is whether these assumptions are consistent with the difference in the

magnitude of cross-priming produced by plaids on the rats of the G group (Figure 2A) and by

gratings on the rats of the P group (Figure 2B). To account for this difference, we reasoned

that pattern cells connected to a given decision neuron could fire less in response to plaids

(because of cross-orientation suppression) than in response to gratings. This could lead the

decision neuron to form stronger synaptic connections with its afferent pattern cells in the

case of training with plaids than with gratings, under the assumption that some compensatory

mechanisms actively keep the weighted input to the decision neuron around a target setpoint.

Such compensatory process would be fully consistent with the well-known role played by

homeostatic plasticity in cortex, where the overall synaptic strength scales as a function of the

magnitude of the input, so as to maintain the average firing rate of cortical neurons within a

given range (Turrigiano and Nelson, 2000; Turrigiano, 2008; Hengen et al., 2013).

To verify the plausibility of these intuitions, we simulated a downstream decision

neuron reading out the representation conveyed by a population of pattern cells with strong

cross-orientation suppression. As done in the previous section, we implemented the decision

neuron as a logistic classifier with a regularization term on the L2 norm of its "synaptic"

weights, so as to simulate the effect of homeostatic plasticity. The classifier was trained to

discriminate either gratings or plaids drifting in opposite directions (i.e., $0º$ vs. $180º$ motion

direction). As a result of the cross-orientation suppression, the L2 norm of the population

vectors fed as inputs to the classifier was larger for the gratings than for the plaids (Figure

5A). This led the magnitude of the weights to become larger in the case of training with the

plaids than with the gratings (Figure 5B), with the L2 norm of the weights' vector being more

than twice as large in the former case (Figure 5C).

This had the "homeostatic" effect of nearly equalizing the magnitude of the weighted

inputs to the classifier, when this was fed with the same kind of stimuli that were used for

training. As shown in Figure 5D, the weighted inputs resulting from presenting drifting

gratings to the classifier that was trained to discriminate grating motion direction (solid black

curve) were very close to the weighted inputs resulting from presenting drifting plaids to the

classifier that was trained to discriminate plaid motion direction (solid gray curve). This

would explain the strong identity priming observed for the rats of both the G and P groups in

our experiment (Figure 2, black curves). At the same time, the large weights' magnitude

resulting from the training with the plaids also guaranteed a very strong weighted input in the

case of presentation of the grating stimuli (dashed gray curve in Figure 5D). This would

explain the strong cross-priming observed for the rats of the P group (Figure 2B, gray curve).

By contrast, the much smaller weights resulting from training with the gratings yielded a

near-zero weighted input in the case of presentation of the plaid stimuli (dashed black curve

in Figure 5D). This would explain the very small cross-priming observed for the rats of the G

group (Figure 2A, gray curve). A plaid prime stimulus would simply be too weak (because of

cross-orientation suppression) to activate the decision neuron through the weak, grating-

adapted synaptic weights.

Conclusions and implications

Although speculative in nature, the two explanations detailed in the previous paragraphs and

backed up by our simulations propose an account for the behavioral data reported in our

study. Importantly, they both rest on well-defined physiological assumptions. As such, they

give rise to alternative hypotheses about the neural mechanisms underlying the processing of

motion information that could be tested experimentally. For instance, the first explanation

(Figure 4) would be supported by observing cross-orientation suppression acting way more

strongly on component than on pattern cells in rodent visual cortex. On the other hand, the

second explanation (Figure 5) would be corroborated by observing that cross-orientation

suppression affects equally strongly pattern and component cells, and by finding a projection-

specific enrichment in the proportion of pattern cells relaying visual information to decision

making centers, such as posterior parietal cortex (PPC) (Raposo et al., 2014; Licata et al.,

2017; Nikbakht et al., 2018; Odoemene et al., 2018).

In conclusion, our study not only yields behavioral evidence of motion integration in

rodents, but also provides solid grounding for future investigations aimed at dissecting the

neuronal circuits underlying integration of local motion cues into global motion percepts. In

fact, as highlighted in a recent perspective (Rasmussen and Yonehara, 2017), a behavioral

task involving direction discrimination of gratings and plaids is a necessary ingredient of

future experiments aimed at establishing a correlational and/or causal link between pattern-

like responses observed in rodent visual cortex and integrated motion perception.

Materials and methods

Animal procedures

We trained 21 male Long Evans rats (Charles River Laboratory) in a motion direction discrimination task. Upon arrival in the lab, rats weighed ~250 g, and they grew up to ~500 g. Animals started training during ~ 7th postnatal week. During the experimental period, they had free access to food, while their access to water was restricted. Their daily liquid intake included 5 – 15 ml of a 1:4 juice-water solution, plus an ad libitum water access for one hour after training. Training sessions lasted 50 – 70 minutes and took place 5 days per week. All animal procedures were conducted in accordance with the international and institutional standards for the care and use of animals in research and were approved by the Italian Ministry of Health and after consulting with a veterinarian (Project DGSAF 25271, submitted on December 1, 2014 and approved on September 4, 2015, approval 940/2015-PR).

Behavioral apparatus and visual stimuli

The behavioral rig consisted of two racks, each equipped with three operant boxes to allow training a batch of six rats simultaneously (Zoccolan, 2015; Zoccolan and Di Filippo, 2018). In each box, one of the walls bore a 3 cm-diameter hole, so that a rat could extend its head outside the box and face the stimulus display (a LCD monitor ASUS® Ve228 located at 30 cm from the viewing hole). The box was equipped with three stainless-steel feeding needles (Cadence Science®), located right outside the viewing hole, and serving as response ports and for reward delivery. To this end, each needle was connected to a led-photodiode pair to detect when the nose of the animal approached and touched it. The two ports positioned to the right and to the left of the hole were connected to two computer-controlled syringe pumps (New Era Pump Systems NE-500®) for delivery of liquid reward, while the central port served exclusively to trigger the onset of each behavioral trial. Each rat learnt to extend its head through the viewing hole, lick the central needle to trigger stimulus presentation and then lick either the left or right port to report the identity of the visual stimulus presented in the current trial. In case of correct response, the reward was delivered through the feeding needle. In case of an incorrect choice, a 1-3 s timeout period started, with the stimulus display flickering from black to middle gray at a rate of 15 Hz (during this period, a "failure" sound was also played). Stimulus presentation, response collection and reward delivery were controlled via workstations running open source MWorks software (https://mworks.github.io/).

As explained in the Results, the rats were divided in two groups. The animals in the G

group were trained to discriminate gratings drifting horizontally in opposite directions, i.e.,

180° (leftward) vs. 0° (rightward). The rats of the P group were trained to discriminate two

drifting plaids, whose global motion direction was, again, either 180° or 0º. The grating

stimuli consisted in full-field, full-contrast, sine wave drifting gratings with a temporal

frequency of 2 Hz and a spatial frequency of 0.04 cycles/°. The plaid stimuli were

constructed by superimposition of two half-contrast gratings (again 0.04 cycles/° and 2 Hz)

with a motion direction difference of 120° (i.e., 120° plaid cross-angle). In the experiments

with the priming paradigm (see below), the drift direction of the prime stimuli (either gratings

or plaids) was randomly sampled in each trial from 19 possible directions (i.e., from 0° to

180° in steps of 10; see Figure 1).

Experimental design - Training phase

As soon as a rat licked the central response port, the target stimulus appeared (either a grating

or a plaid depending on the animal's experimental group) and remained on screen for 2 s. To

prevent rats from making very quick, impulsive responses, a trial was aborted if the animal's

reaction time was lower than 300 ms. In such a case, the animal's response was not evaluated

(neither reward or time-out was administered), the stimulus was immediately turned off, and

a brief tone was played. Such "too fast" trials were also excluded from the analysis.

Similarly, trials in which the rat responded more than 2 s after the offset of the stimulus were

considered "ignored" and excluded from the analysis. No reward was delivered in these

invalid trials. In order to accomplish the task and receive the reward, the rat had to reach and

lick the response port matching the global motion direction of the target stimulus. Correct

execution of such task was rewarded equally for both directions during each experimental

session. In correct trials, the target stimulus remained on the screen for the whole reward

delivery period, so as to strengthen the operant association between stimulus response and

reward. On the other hand, when the animal provided an incorrect response, reward was not

delivered, the stimulus was immediately turned off and the timeout period started (see

previous section).

Experimental design - Priming paradigm

In the priming paradigm, a prime stimulus was shown to the animal before the target

stimulus. The prime duration was 75 ms, the inter-stimulus interval (ISI, i.e., the interval

separating the offset of the prime from the onset of the target) was also 75 ms and the target

duration was 750 ms. As in the training phase, "too fast" or "ignored" trials were excluded
from the analysis. Only, a trial was defined as "ignored" if the rat responded more than 1 s
after the onset of the target stimulus. As a result, the allowed response window extended from
300 to 1000 ms from target onset. As in the training phase, an animal, in order to obtain the
reward, had to reach and lick the response port (right or left) corresponding to the global
motion direction of the target. Accordingly, no feedback was ever provided to the rat
regarding the motion direction of prime stimuli. In this paradigm, the target was not kept on
screen during the reward, because there was no longer any need to favor the association
between response, stimulus, and reward. Only animals maintaining a performance in the
training task greater that 75% for 4 consecutive days were tested with this priming paradigm.

The rationale of this design rests on well-established findings in the human
psychophysics literature (Bar and Biederman, 1998; Leopold et al., 2001; Kanai and
Verstraten, 2005; Kravitz et al., 2010), where presenting an "adapter" before a "test" stimulus
has been shown to bias the perception of the test, depending on two key factors: the duration
of the adapter and the inter stimulus interval (ISI) between adapter and target presentation.
When the bias induced by the adapter attracts the perceptual choices towards the identity of
the adapter itself, this effect is called "priming". Vice versa, when it repels the perceptual
choices away from the identity of the adapter, it is named "adaptation after-effect". Our
group previously shown that brief presentation of a static shape (~50 ms), followed by a short
ISI (66 ms), is able to induce a strong and robust priming effect on rat response to a target
shape (Tafazoli et al., 2012). Here, we adapted this paradigm to moving stimuli, and we
relied on previous motion priming studies in humans (Kanai and Verstraten, 2005) to select
the timing parameters of the task, in the attempt of inducing a strong priming effect.

Analysis of behavioral data

Only rats displaying a strong identity-priming effect where included in the analysis described
in the main text. Quantitatively, this corresponded to setting a threshold on the absolute
priming magnitude in the identity priming condition amounting to 5%. Priming magnitude
was computed averaging together the mean values of the absolute difference between the left
and right extremes (4 points per side) of each priming curve with respect to the neutral prime
condition (insets in Figure 2). The enforcement of such criterion led to the rejection of one rat
per group, bringing the number of rats included in the analysis to 10 animals for the G group

and to 9 animals for the P group out of the original 11 animals trained in grating direction

discrimination and 10 trained in plaid direction discrimination.

To statistically compare the identity- and cross-priming curves obtained for each

group of rats (solid and dashed curves in Figure 2) we performed a bootstrap analysis, in

which the recorded behavioral sessions were resampled 50 times with replacement for each

animal independently. This gave us the possibility to estimate a bootstrap standard deviation

for both the average identity- and cross-priming curves of Figure 2, as well as for the average

priming magnitudes computed as described above (insets in Figure 2). Bootstrap standard

deviations were then used to compute normal 95% confidence intervals for the

aforementioned quantities (displayed in Figure 2 as shaded regions around the priming curves

and as error bars in the bar plot shown in the insets) (Efron and Tibshirani, 1986). In

computing confidence intervals for the 19 points of a priming curve, the critical value

corresponding to the chosen confidence level was adjusted for multiple comparisons using

Bonferroni correction.

Computational modeling

To interpret the results of our psychophysical tests, we simulated a decision neuron reading

out the activity of a simulated population of either component or pattern cells fed with either

grating or plaid stimuli. Three different simulations were performed to test the consistency of

our findings with three possible scenarios concerning the tuning of rat visual neurons for

motion direction and the choice of the decoding pool by the decision neuron. These

simulations have been already described in the Results and in the legends of Figures 3, 4 and

5. Below, we explain some additional technical details about how we modeled the

populations of simulated component and pattern cells and how we implemented the logistic

classifier used to simulate the decision neuron.

Simulations to check the consistency of the priming curves observed for the P group with a

pattern-based representation of global motion direction

To check whether the shape of the priming curves observed for the rats of the P group (Figure

2B) was more consistent with a component-like or a pattern-like representation, we trained a

logistic classifier to solve the same motion discrimination task that was administered to the

rats (i.e., 0º- vs. 180º-drifting plaids), based on the responses of a simulated population of

either component of pattern cells (Figure 3). We then tested how the trained classifier would

decode the full range of drifting plaids and gratings used to obtain, respectively, the identity-

and cross-priming curves in Figure 2B. To simulate the tuning for local and global motion

direction of the populations of component and pattern cells, we used 24 Von Mises functions

(Kohn and Movshon, 2004; Matteucci et al., 2019) centered at equispaced angles along the

circle (spaced by 15°). The von Mises functions, which are the circular analogs of Gaussian

functions, have a width that is controlled by the parameter k, which, in turn, is equivalent to

the inverse of the variance of a Gaussian distribution (Berens, 2009). The peak value of the

functions (and thus of the tuning curves) was set to 1 for all the simulated units.

For a pattern cell, the simulated tuning curve was the same regardless of whether the

input stimulus was a plaid or a grating stimulus (compare matching rows in the top and

bottom panels of Figure 3B). That is, by construction, each simulated pattern cell responded

to the global direction of the plaid stimuli. Conversely, for a component cell, the simulated

tuning curve was defined by a single Von Mises function in case of presentation of the

grating stimuli (Figure 3A, top), but was defined by the superimposition of two von Mises

functions (with the peaks being 120º apart) in case of presentation of the plaid stimuli (Figure

3A, bottom). That is, by construction, each simulated component cell responded to the local

directions of the constituent gratings of the plaid stimuli, rather than to the global direction of

the plaid. For both the simulated pattern and component cells, the Von Mises functions were

summed to a constant background activity. In addition, to emulate the variability of neuronal

firing in response to identical stimuli, in each simulated trial, the actual response to a given

stimulus was affected by Gaussian noise with zero mean and $\sigma = 0.25$. The k parameter of the

Von Mises functions of the simulated pattern and component cells was set to 7 in the mean

simulation shown in Figure 3, while it was allowed to range between 0.5 and 11 in the more

extended simulations of Figure 3–figure supplement 1. The ratio between the peak of the

tuning curves and the background rate was set to 10.

To model the training received by the rats in our experiment, we sampled for 500

times the responses of the whole population of either pattern or component cells to each of

the 0º- and 180º-drifting stimuli (either grating or plaids) used during the training phase. We

then trained a logistic regression classifier to discriminate the two sets of 0º and 180º

responses using gradient descent (implemented by the Matlab function "fminunc") and

including and L2 regularization term in the cost function (whose parameter lambda was set to

1). The weights resulting from such training are those shown in Figure 3C and E. Finally, we

fed to the classifier grating and plaid stimuli spanning the full range of possible global motion

directions (from 0º to 345º in steps of 15º) and we plotted the probability for each class of

stimuli to be classified as moving rightward as a function of its direction (Figure 3D and F).

Simulations to explore the mechanisms that could lead to a strong cross-priming only in case

of training with plaid stimuli – scenario #1

To obtain a proof of principle that the mechanisms illustrated in Figure 4A, B could lead to a

decoding pool enriched of pattern cells in case of training with plaids, we carried out a

second simulation. As before, we trained a logistic regression classifier in the same motion

direction discrimination task that was administered to our animals. This time, however, the

response properties of the simulated component and pattern cells, as well as their relative

proportion, better matched either the known properties derived from the rodent literature

(Juavinett and Callaway, 2015; Muir et al., 2015; Palagina et al., 2017) or those inferred

based on the monkey literature (Smith et al., 2005; Rust et al., 2006). First, the simulated

population of pattern and component cells was much larger (i.e., 1000 units). It was

composed of Poisson firing neurons with either "component" or "pattern" tuning curves, built

in the same way as described in the previous section. Second, their proportion was 80% and

20% respectively, so as to match the highest ratio of pattern to component cells observed in

rodent visual cortex (Juavinett and Callaway, 2015). As in the previous simulation, we used

Von Mises functions to define the tuning curves of component and pattern cells (see

description in the previous section). In this case however, the preferred directions of the

simulated neurons were not equispaced but drawn randomly from a uniform distribution over

the circle. Baseline value of tuning curves was set to 2 spikes/second. Von Mises function of

random amplitude (normally distributed around 8 spikes/s with standard deviation of 1 spike

per second) were added to this baseline. The firing rate of the units in each trial was drawn

from a Gaussian distribution with mean equal to the value set by the tuning curves and

standard deviation of 2 spikes/s. The width (i.e., the parameter k; see previous paragraph) of

the Von Mises functions was adjusted in such a way that the Orientation Selectivity Index

(OSI) of the resulting population of component cells was distributed according to a Gaussian

with a mean of 0.7 and a standard deviation of 0.1, while the Direction Selectivity Index

(DSI) was distributed according to a Gaussian with a mean of 0.6 and a standard deviation of

0.2. OSI and DSI are standard measures of orientation and direction tuning, whose definition

can be found, for instance, in (Niell and Stryker, 2008; Vermaercke et al., 2014; Matteucci et

al., 2019; Matteucci and Zoccolan, 2020).

To account for the imperfect cross-angle invariance documented by (Palagina et al.,

2017) we enlarged the width of Von Mises functions used to simulate the tuning of pattern

cells to plaids by dividing the k parameter previously used to define their tuning to gratings

(see previous paragraph) by a factor 4. This made the responses of our simulated pattern cells

better matched to the known properties of mouse pattern cells, reproducing their tendency to

have broader tuning curves when tested with plaids than with gratings, especially when the

plaids have a large cross-angle (such as the one used in our study, i.e. 120°).

Finally, we simulated cross-orientation suppression in component cells – that is, we

imposed that the peak responses of component cells to the plaids were half of their peak

responses to the gratings, while pattern cells were implemented as being fully unaffected by

cross-orientation suppression (i.e., same peak responses to plaids and gratings). This scenario

was simulated because our goal was to test the hypothesis that, if cross-orientation

suppression affected strongly component cells but not pattern cells, then the decision neuron

would learn to preferentially connect with the latter, in the case of training with the plaid

stimuli.

To model the training received by the rats in our experiment, we sampled for 200

times the responses of the mixed population of pattern or component cells to each of the 0°-

and 180°-drifting stimuli (either grating or plaids) used during the training phase. We then

trained a logistic regression classifier to discriminate the two sets of 0° and 180° responses

using gradient descent (implemented by the Matlab function "fminunc") and including and

L2 regularization term in the cost function (whose parameter lambda was set to 1). The

weights resulting from such training are those shown in Figure 4C and D (top). Finally, we

simulated a scenario where, as a result of this training, the decision neuron would maintain

only the strongest of its synaptic connections with sensory neurons encoding the drifting

stimuli. To incorporate this sparseness constraint on the connectivity of the decision neuron,

we progressively pruned the weakest synaptic weights by applying a quantile-based

thresholding to their magnitude after training (i.e., by setting to zero any weight below the

chosen quantile of the original weight magnitude distribution). We then quantified the

proportion of pattern and component cells that survived the pruning and remained in the

decoding pool of the decision neuron as a function of the quantile used as a threshold (Figure

4C and D, bottom). As a control, the whole simulation was also run without incorporating

cross-orientation suppression in the responses of component cells.

Simulations to explore the mechanisms that could lead to a strong cross-priming only in case

of training with plaid stimuli – scenario #2

Here, we simulated an alternative scenario where a decision neuron (modeled, again, as a

logistic classifier) is connected to a pool of sensory neurons made exclusively of pattern cells

that are strongly affected by cross-orientation suppression. As such, the parameters of the

simulations were the same as those described in the previous section, with only two notable

differences: 1) no component cells were included in the decoding pool, but only pattern cells;

2) cross-orientation suppression affected pattern cells instead of components cells, and its

impact was twice as strong as that of the previous stimulation (i.e., peak responses to plaids

were 25% of peak responses to gratings). The training of the classifier was the same as

described in the previous section. Note that the L2 regularization term in the cost function

played the role of a homeostatic constraint on the overall weights' magnitude. Again, we

compared the weights' distributions resulting from training with plaids and gratings (Figure

5B) and we also quantified the magnitude of weighted input (i.e., sensory "evidence") to the

decision neuron, depending on the training and prime stimuli (Figure 5D).

Acknowledgments. This work was supported by a European Research Council Consolidator

Grant (DZ, project n. 616803-LEARN2SEE). We thank Daniele Bertolini for his help in

developing and building the rigs for the behavioral experiments, and we thank Sofia Erica

Rossi and Marco Salluzzo for their help in training rats in the motion discrimination task.

Competing interests. None of the authors has financial and non-financial competing

interests of any nature.

References

Adelson EH, Movshon JA (1982) Phenomenal coherence of moving visual patterns. Nature
300:523–525.

Alemi-Neissi A, Rosselli FB, Zoccolan D (2013) Multifeatural Shape Processing in Rats
Engaged in Invariant Visual Object Recognition. J Neurosci 33:5939–5956.

Alwis DS, Richards KL, Price NSC (2016) Masking reduces orientation selectivity in rat
visual cortex. J Neurophysiol 116:2331–2341.

Bar M, Biederman I (1998) Subliminal Visual Priming. Psychol Sci 9:464–468.

Berens P (2009) CircStat: A MATLAB Toolbox for Circular Statistics. J Stat Softw 31.

Born RT, Bradley DC (2005) Structure and function of visual area MT. Annu Rev Neurosci
28:157–189.

Busse L, Wade AR, Carandini M (2009) Representation of Concurrent Stimuli by Population
Activity in Visual Cortex. Neuron 64:931–942.

Carandini M, Churchland AK (2013) Probing perceptual decisions in rodents. Nat Neurosci
16:824–831.

Chen T-W, Wardill TJ, Sun Y, Pulver SR, Renninger SL, Baohan A, Schreiter ER, Kerr RA,
Orger MB, Jayaraman V (2013) Ultrasensitive fluorescent proteins for imaging
neuronal activity. Nature 499:295.

DeAngelis GC, Robson JG, Ohzawa I, Freeman RD (1992) Organization of suppression in
receptive fields of neurons in cat visual cortex. J Neurophysiol 68:144–163.

Dell KL, Arabzadeh E, Price NSC (2018) Human-like perceptual masking is difficult to
observe in rats performing an orientation discrimination task. PLOS ONE
13:e0207179.

Dell KL, Arabzadeh E, Price NSC (2019) Differences in perceptual masking between
humans and rats. Brain Behav 9:e01368.

Djurdjevic V, Ansuini A, Bertolini D, Macke JH, Zoccolan D (2018) Accuracy of Rats in
Discriminating Visual Objects Is Explained by the Complexity of Their Perceptual
Strategy. Curr Biol 28:1005-1015.e5.

Douglas RM, Neve A, Quittenbaum JP, Alam NM, Prusky GT (2006) Perception of visual
motion coherence by rats and mice. Vision Res 46:2842–2847.

D'Souza RD, Bista P, Meier AM, Ji W, Burkhalter A (2019) Spatial Clustering of Inhibition in
Mouse Primary Visual Cortex. Neuron 104:588-600.e5.

Efron B, Tibshirani R (1986) Bootstrap Methods for Standard Errors, Confidence Intervals,
and Other Measures of Statistical Accuracy. Stat Sci 1:54–75.

Fennema CL, Thompson WB (1979) Velocity determination in scenes containing several
moving objects. Comput Graph Image Process 9:301–315.

Fenno L, Yizhar O, Deisseroth K (2011) The Development and Application of Optogenetics.
Annu Rev Neurosci 34:389–412.

Gao E, DeAngelis GC, Burkhalter A (2010) Parallel Input Channels to Mouse Primary Visual
Cortex. J Neurosci 30:5912–5926.

Ghodrati M, Alwis DS, Price NSC (2016) Orientation selectivity in rat primary visual cortex
emerges earlier with low-contrast and high-luminance stimuli. Eur J Neurosci
44:2759–2773.

Girman SV, Sauvé Y, Lund RD (1999) Receptive Field Properties of Single Neurons in Rat
Primary Visual Cortex. J Neurophysiol 82:301–311.

Glickfeld LL, Andermann ML, Bonin V, Reid RC (2013) Cortico-cortical projections in mouse
visual cortex are functionally target specific. Nat Neurosci 16:219–226.

Harvey RJ, De'sperati C, Strata P (1997) The Early Phase of Horizontal Optokinetic
969   Responses in the Pigmented Rat and the Effects of Lesions of the Visual Cortex.
970   Vision Res 37:1615–1625.

971   Hengen KB, Lambo ME, Van Hooser SD, Katz DB, Turrigiano GG (2013) Firing Rate
972   Homeostasis in Visual Cortex of Freely Behaving Rodents. Neuron 80:335–342.

973   Ji W, Gămănuţ R, Bista P, D'Souza RD, Wang Q, Burkhalter A (2015) Modularity in the
974   Organization of Mouse Primary Visual Cortex. Neuron 87:632–643.

975   Jin M, Glickfeld LL (2020) Mouse Higher Visual Areas Provide Both Distributed and
976   Specialized Contributions to Visually Guided Behaviors. Curr Biol 30:4682-4692.e7.

977   Juavinett AL, Callaway EM (2015) Pattern and Component Motion Responses in Mouse
978   Visual Cortical Areas. Curr Biol 25:1759–1764.

979   Kanai R, Verstraten FAJ (2005) Perceptual manifestations of fast neural plasticity: Motion
980   priming, rapid motion aftereffect and perceptual sensitization. Vision Res 45:3109–
981   3116.

982   Kohn A, Movshon JA (2004) Adaptation changes the direction tuning of macaque MT
983   neurons. Nat Neurosci 7:764–772.

984   Kravitz DJ, Kriegeskorte N, Baker CI (2010) High-Level Visual Object Representations Are
985   Constrained by Position. Cereb Cortex 20:2916–2925.

986   Leopold DA, O'Toole AJ, Vetter T, Blanz V (2001) Prototype-referenced shape encoding
987   revealed by high-level aftereffects. Nat Neurosci 4:89–94.

988   Li B, Thompson JK, Duong T, Peterson MR, Freeman RD (2006) Origins of Cross-
989   Orientation Suppression in the Visual Cortex. J Neurophysiol 96:1755–1764.

990   Licata AM, Kaufman MT, Raposo D, Ryan MB, Sheppard JP, Churchland AK (2017)
991   Posterior Parietal Cortex Guides Visual Decisions in Rats. J Neurosci 37:4954–4966.

992   Lisberger SG (2015) Visual Guidance of Smooth Pursuit Eye Movements. Annu Rev Vis Sci
993   1:447–468.

994   Luo L, Callaway EM, Svoboda K (2008) Genetic Dissection of Neural Circuits. Neuron
995   57:634–660.

996   Luo L, Callaway EM, Svoboda K (2018) Genetic Dissection of Neural Circuits: A Decade of
997   Progress. Neuron 98:256–281.

998   Madisen L, Garner AR, Shimaoka D, Chuong AS, Klapoetke NC, Li L, Van Der Bourg A,
999   Niino Y, Egolf L, Monetti C (2015) Transgenic mice for intersectional targeting of
1000   neural sensors and effectors with high specificity and performance. Neuron 85:942–
1001   958.

1002   Marques T, Summers MT, Fioreze G, Fridman M, Dias RF, Feller MB, Petreanu L (2018) A
1003   Role for Mouse Primary Visual Cortex in Motion Perception. Curr Biol CB 28:1703-
1004   1713.e6.

1005   Marr D (2010) Vision: a computational investigation into the human representation and
1006   processing of visual information. Cambridge, Mass: MIT Press.

Marshel JH, Garrett ME, Nauhaus I, Callaway EM (2011) Functional Specialization of Seven
Mouse Visual Cortical Areas. Neuron 72:1040–1054.

Matsui T, Ohki K (2013) Target dependence of orientation and direction selectivity of
corticocortical projection neurons in the mouse V1. Front Neural Circuits 7 Available
at: https://www.frontiersin.org/articles/10.3389/fncir.2013.00143/full [Accessed
January 26, 2018].

Matteucci G, Marotti RB, Riggi M, Rosselli FB, Zoccolan D (2019) Nonlinear Processing of
Shape Information in Rat Lateral Extrastriate Cortex. J Neurosci 39:1649–1670.

Matteucci G, Zoccolan D (2020) Unsupervised experience with temporal continuity of the
visual environment is causally involved in the development of V1 complex cells. Sci
Adv 6:eaba3742.

Muir DR, Roth MM, Helmchen F, Kampa BM (2015) Model-based analysis of pattern motion
processing in mouse primary visual cortex. Front Neural Circuits 9 Available at:
http://www.ncbi.nlm.nih.gov/pmc/articles/PMC4525018/ [Accessed April 24, 2017].

Newsome W, Pare E (1988) A selective impairment of motion perception following lesions of
the middle temporal visual area (MT). J Neurosci 8:2201–2211.

Niell CM, Stryker MP (2008) Highly selective receptive fields in mouse visual cortex. J
Neurosci 28:7520–36.

Nikbakht N, Tafreshiha A, Zoccolan D, Diamond ME (2018) Supralinear and Supramodal
Integration of Visual and Tactile Signals in Rats: Psychophysics and Neuronal
Mechanisms. Neuron 97:626-639.e8.

Nishida S, Kawabe T, Sawayama M, Fukiage T (2018) Motion Perception: From Detection to
Interpretation. Annu Rev Vis Sci 4:501–523.

Odoemene O, Pisupati S, Nguyen H, Churchland AK (2018) Visual Evidence Accumulation
Guides Decision-Making in Unrestrained Mice. J Neurosci 38:10143–10155.

Orban GA (2008) Higher Order Visual Processing in Macaque Extrastriate Cortex. Physiol
Rev 88:59.

Palagina G, Meyer JF, Smirnakis SM (2017) Complex Visual Motion Representation in
Mouse Area V1. J Neurosci 37:164–183.

Priebe NJ, Ferster D (2006) Mechanisms underlying cross-orientation suppression in cat
visual cortex. Nat Neurosci 9:552–561.

Raposo D, Kaufman MT, Churchland AK (2014) A category-free neural population supports
evolving demands during decision-making. Nat Neurosci 17:1784–1792.

Rasmussen R, Yonehara K (2017) Circuit Mechanisms Governing Local vs. Global Motion
Processing in Mouse Visual Cortex. Front Neural Circuits 11:109.

Roelfsema PR, Holtmaat A (2018) Control of synaptic plasticity in deep cortical networks.
Nat Rev Neurosci 19:166–180.

Rosselli FB, Alemi A, Ansuini A, Zoccolan D (2015) Object similarity affects the perceptual
strategy underlying invariant visual object recognition in rats. Front Neural Circuits
9:10.

Rust NC, Mante V, Simoncelli EP, Movshon JA (2006) How MT cells analyze the motion of
visual patterns. Nat Neurosci 9:1421–31.

Salzman CD, Britten KH, Newsome WT (1990) Cortical microstimulation influences
perceptual judgements of motion direction. Nature 346:174–177.

Samonds JM, Feese BD, Lee TS, Kuhlman SJ (2017) Nonuniform surround suppression of
visual responses in mouse V1. J Neurophysiol 118:3282–3292.

Samonds JM, Lieberman S, Priebe NJ (2018) Motion Discrimination and the Motion
Aftereffect in Mouse Vision. eNeuro 5.

Self MW, Lorteije JAM, Vangeneugden J, Beest EH van, Grigore ME, Levelt CN, Heimel JA,
Roelfsema PR (2014) Orientation-Tuned Surround Suppression in Mouse Visual
Cortex. J Neurosci 34:9290–9304.

Simoncelli EP, Heeger DJ (1998) A model of neuronal responses in visual area MT. Vision
Res 38:743–761.

Smith AT, Harris LR (1991) Use of plaid patterns to distinguish the corticofugal and direct
retinal inputs to the brainstem optokinetic nystagmus generator. Exp Brain Res 86
Available at: http://link.springer.com/10.1007/BF00228955 [Accessed November 23,
2019].

Smith MA, Majaj NJ, Movshon JA (2005) Dynamics of motion signaling by neurons in
macaque area MT. Nat Neurosci 8:220–228.

Solomon SS, Tailby C, Gharaei S, Camp AJ, Bourne JA, Solomon SG (2011) Visual motion
integration by neurons in the middle temporal area of a New World monkey, the
marmoset: Pattern motion sensitivity in middle temporal area of marmosets. J Physiol
589:5741–5758.

Stahl JS (2004) Using eye movements to assess brain function in mice. Vision Res
44:3401–3410.

Stirman JN, Townsend LB, Smith SL (2016) A touchscreen based global motion perception
task for mice. Vision Res 127:74–83.

Tafazoli S, Di Filippo A, Zoccolan D (2012) Transformation-Tolerant Object Recognition in
Rats Revealed by Visual Priming. J Neurosci 32:21–34.

Turrigiano GG (2008) The Self-Tuning Neuron: Synaptic Scaling of Excitatory Synapses.
Cell 135:422–435.

Turrigiano GG, Nelson SB (2000) Hebb and homeostasis in neuronal plasticity. Curr Opin
Neurobiol 10:358–364.

Vermaercke B, Gerich FJ, Ytebrouck E, Arckens L, Op de Beeck HP, Van den Bergh G
(2014) Functional specialization in rat occipital and temporal visual cortex. J
Neurophysiol 112:1963–1983.

Wuerger S, Shapley R, Rubin N (1996) "On the Visually Perceived Direction of Motion" by
Hans Wallach: 60 Years Later. Perception 25:1317–1367.

Zoccolan D (2015) Invariant visual object recognition and shape processing in rats. Behav
Brain Res 285:10–33.

Zoccolan D, Di Filippo A (2018) Chapter 5 - Methodological Approaches to the Behavioural
Investigation of Visual Perception in Rodents In A. Ennaceur & M. A. de Souza Silva,
eds. Handbook of Behavioral Neuroscience Handbook of Object Novelty
Recognition. Elsevier, p. 69–101.

Zoccolan D, Oertelt N, DiCarlo JJ, Cox DD (2009) A rodent model for the study of invariant
visual object recognition. Proc Natl Acad Sci U A 106:8748–53.

1093

1094

1095

1096

1097

1098

1099

1100

1101

1102

Figure supplements

1104
Figure 3 – figure supplement 1. Simulating the discrimination accuracy afforded by a component-
based representation with different degree of direction tuning. (**A**) Example tuning curves over the
direction axis, as obtained for increasingly large values of the parameter *k*, which controlled the width
of the Von Mises functions used to simulate the tuning of the component cells (see main text). For
each value of *k*, a population of 24 components cells was simulated, whose preferred directions
homogeneously spanned the [0° 360°] range (same as in Figure 3A). The figure shows how one of
such cells (the one with preferred direction = 160°) responded to drifting gratings (gray dashed curve)
and plaids (black solid curve) as a function of their direction. For *k* = 7, the curves report the same
values already shown in the rows of Figure 3A (top and bottom, respectively) corresponding to the

preferred direction of 160º. (**B**) Weights assigned to the 24 simulated component cells by a logistic classifier that was trained to discriminate a 0º- (rightward) from a 180º-drifting (leftward) plaid. The parameter k varies as indicated in A. (**C**) Fraction of times that the logistic classifier fed with the component-based representation (whose weights are shown in B) classified the test stimuli as drifting rightward (i.e., at 0º). The test stimuli were either plaids (solid black curve) or gratings (dashed gray curve) spanning the whole [0º 360º] range of directions. The parameter k varies as indicated in A. (**D**) Absolute difference between the pairs of curves shown in C for plaid and gratings (averaged across all tested directions) as a function of the parameter k.