Repeated subfunctionalization of a modular antimicrobial peptide gene for neural function

M.A. Hanson* and B. Lemaitre

Global Health Institute, School of Life Science, École Polytechnique Fédérale de Lausanne (EPFL), Lausanne, Switzerland.
* Corresponding author: M.A. Hanson (mark.hanson@epfl.ch)
ORCID IDs:
Hanson: https://orcid.org/0000-0002-6125-3672
Lemaitre: https://orcid.org/0000-0001-7970-1667

Abstract

Antimicrobial peptides (AMPs) are host-encoded antibiotics that combat invading pathogens. However recent studies have highlighted roles for AMPs in neurological contexts suggesting functions for these defence molecules beyond infection. Here we characterize the evolution of the *Drosophila Baramicin* (*Bara*) AMP gene family. During our immune study characterizing the *Baramicins*, we recovered multiple *Baramicin* paralogs in *Drosophila melanogaster* and other species, united by their N-terminal IM24 domain. Strikingly, some paralogs are no longer immune-induced. A careful dissection of the *Baramicin* family's evolutionary history indicates that these non-immune paralogs result from repeated events of duplication and subsequent truncation of the coding sequence from an immune-inducible ancestor. These truncations leave only the IM24 domain as the prominent gene product. Using mutation and targeted gene silencing, we demonstrate that two such genes are adapted for function in neural contexts in *D. melanogaster*, and show enrichment in the head for independent *Baramicin* genes in other species. The *Baramicin* evolutionary history reveals that the IM24 *Baramicin* domain is not strictly useful in an immune context. We thus provide a case study for how an AMP-encoding gene might play dual roles in both immune and non-immune processes via its multiple peptide products. We reflect on these findings to highlight a blind spot in the way researchers approach AMP research in in vivo contexts.

Introduction

Antimicrobial peptides (AMPs) are immune effectors best known for their

role in defence against infection. These antimicrobials are commonly encoded as a

polypeptide including both pro- and mature peptide domains (Zanetti 2005; Hanson

and Lemaitre 2020). AMP genes frequently experience events of duplication and

loss (Wang and Zhu 2011; Vilcinskas et al. 2013; Sackton et al. 2017; Hanson,

Lemaitre, et al. 2019) and undergo rapid evolution at the sequence level (Tennessen

2005; Jiggins and Kim 2007; Hellgren et al. 2010; Halldórsdóttir and Árnason 2015;

Hanson et al. 2016; Chapman et al. 2019). The selective pressures that drive these

evolutionary outcomes are likely the consequence of host-pathogen interactions

(Unckless et al. 2016). However AMPs and AMP-like genes in various species have

also recently been implicated in various non-immune roles in flies, nematodes, and

emerging evidence in humans. These new contexts suggest that the evolutionary

forces acting on AMP genes may not be driven strictly by trade-offs in host defence,

but rather by conflicts between roles in immunity and other non-immune functions.

For instance, Diptericins are membrane-disrupting antimicrobial peptides of

flies (Diptera) that are required for defence against infection by *Providencia* bacteria

(Unckless et al. 2016; Hanson, Dostálová, et al. 2019). It was therefore surprising

that the *D. melanogaster* gene *Diptericin B (DptB)* affects memory processes

(Barajas-azpeleta et al. 2018). In this study, *DptB* derived from the fly fat body

(analogous to the mammalian liver) regulated the ability of the fly to form long-term

memory associations (Barajas-azpeleta et al. 2018). Another AMP-like gene, nemuri,

regulates fly sleep and promotes survival upon infection (Toda et al. 2019). Studies

in nematodes have also shown that an immune-induced polypeptide (NLP-29) binds

to a G-protein coupled receptor (NPR-12) triggering neurodegeneration through

activation of the NPR-12-dependent autophagy pathway (Lezi et al. 2018), and

injury triggers epidermal AMPs including NLP-29 to promote sleep (Sinner et al.

2021). *Drosophila* AMPs have also recently been shown to regulate behaviours after

seeing parasitoid wasps (Ebrahim et al. 2021), during feeding with different bacteria

(Kobler et al. 2020), or following infection (Hanson et al. 2021). In humans, the

Cathelicidin gene encodes the AMP LL-37, which is implicated in glia-mediated

neuroinflammation and Alzheimer's disease (Lee et al. 2015; De Lorenzi et al. 2017),

alongside evidence of Alzheimer's being an infectious syndrome (Dominy et al.

2019); though the importance of this process is debated (Abbott 2020). Notably,

AMPs share a number of properties with classic neuropeptides (Brogden et al.

2005), further muddying the distinction between peptides of the immune and

nervous systems.

We recently described a novel antifungal peptide gene of *Drosophila*

melanogaster that we named *Baramicin A* (*BaraA*) (Hanson et al. 2021). A unique

aspect of *BaraA* is its precursor protein structure, which encodes a polypeptide
cleaved into multiple mature products by interspersed furin cleavage sites. The use
of furin cleavage sites to produce two mature peptides from a single polypeptide
precursor is widespread in animal AMP genes (Gerdol et al. 2020; Hanson and
Lemaitre 2020). However, *BaraA* represents an exceptional case as multiple tandem
repeat peptides are produced from the translation of a single coding sequence,
effectively resembling a "protein-based operon"; this tandem repeat structure has
also been found in two other AMPs of bees and flies (Casteels-Josson et al. 1993;
Hanson et al. 2016). The immature precursor protein of *D. melanogaster BaraA*
encodes three types of domains: an IM24 domain, three tandem repeats of IM10-like
domains, and an IM22 domain. *BaraA* mutants are susceptible to infection by fungi,
and *in vitro* experiments suggest the *BaraA* IM10-like peptides have antifungal
activity (Hanson et al. 2021). The other *Baramicin* domains encoding IM22 and
IM24 remain uncharacterized. Curiously, *BaraA* deficient flies also display an erect
wing behavioural phenotype upon immune stimulation even in the absence of
infection, suggesting that *BaraA* products could have non-microbial targets (Hanson
et al. 2021).
In this study, we describe the evolution of the Drosophilid *Baramicin* gene
family. Three unique *Baramicin* genes (*BaraA, B, and C*) are present in the genome of
D. melanogaster. Surprisingly, only *BaraA* is immune-induced, while *BaraB* and
BaraC are enriched in the nervous system. Both *BaraB* and *BaraC* have truncations
compared to the ancestral *Baramicin* gene, which focuses these genes towards
producing the *Baramicin* IM24 domain. We found similar truncations in other
species, and upon checking their patterns of expression, realized these overt gene
structure changes correlate with loss of immune expression and enrichment in the
nervous system. By resolving the genomic synteny of the various *Baramicin* genes in
different species, we confirmed that these repeated truncations focusing on IM24
production stem from independent events (convergent evolution). The exaggerated
'protein operon' polypeptide nature of *Baramicin* draws attention to the unique
roles that different mature peptides of AMP-encoding genes can play. Careful

attention paid to the multiple peptide products of AMP genes could explain how

these immune effectors contribute to both immune and neurological processes.

106   Results

107   *Baramicin is an ancestral immune effector*

The *Baramicin A* gene was only recently described as encoding antifungal

effectors by our group (Hanson et al. 2021), and another recent study also

confirmed *Baramicin*'s important contribution to Toll immune defence (Huang et al.

2020). These initial characterizations were done only in *D. melanogaster*, and only

focused on one *Baramicin* gene. We will therefore first provide a basic description of

the immune *Baramicins* of other species and also the larger *Baramicin* gene family of

D. melanogaster to establish that this is a classically immune gene family, and that

deviations from immune function are derived.

In *D. melanogaster, BaraA* is regulated by the Toll immune signalling pathway

(Huang et al. 2020; Hanson et al. 2021). Using BLAST, we recovered *BaraA*-like

genes encoding each Baramicin peptide (IM24, IM10-like, and IM22) across the

genus *Drosophila* and in the outgroup *Scaptodrosophila lebanonensis*. We performed

infection experiments to confirm that these *BaraA*-like genes were immune-

inducible by infecting the diverse species *D. melanogaster*, *D. pseudoobscura*, *D.*

willistoni, *D. virilis*, and *D. neotesteacea* (last common ancestor ~63mya (Tamura et

al. 2004)) with *Micrococcus luteus* and *Candida albicans*, two microbes that stimulate

the Toll pathway (**Fig.** 1A). In all five species, *BaraA*-like genes were immune-

induced (**Fig.** 1B-F). We therefore confirm the ancestral *Baramicin* was an immune-

induced gene.

128   *The four D. melanogaster Baramicins: BaraA1, BaraA2, BaraB and BaraC*

129   In *D. melanogaster*, we recovered four *Baramicin* genes. First, we realized

130   that a duplication of *BaraA* is actively segregating in wild flies (**Fig. 2**A). The

Figure 1: The ancestral *BaraA* gene was immune-induced. A) Cladogram of species used in B-F. B-F) Expression of *BomBc3* control genes (brown) or *BaraA-like* genes (orange) in diverse *Drosophila* species upon infection. In all cases, both *BomBc3* and *BaraA-like* genes are induced upon infection by either *C. albicans* yeast or *M. luteus* bacteria.

138
139 **Figure 2: The *D. melanogaster Baramicin* genes.** A) Schematic of the *BaraA* duplication. Using a PCR assay spanning the duplication-specific locus (PCR amplicon), we confirmed *BaraA* copy number is variable in various lab strains (Table S1) and wild-caught flies (Supplementary data file 1). B) *D. melanogaster* encodes two other *Baramicin* genes that we name *BaraB* and *BaraC*. These paralogs differ markedly in their precursor protein structure, which is truncated relative to *BaraA* including an actively segregating truncation event in the wild in *BaraB* (protein lengths implied by greyed out region).

D. melanogaster R6 genome assembly encodes two 100% identical *BaraA* genes

(*CG33470* and *CG18279*, *BaraA1* and *BaraA2* respectively). We screened 132 DGRP

lines for the *BaraA* duplication event, finding only ~14% (18/132) of strains were

PCR-positive for two *BaraA* copies (supplementary data file 1). Perhaps as a

consequence of the identical sequences of these two genes, this genome region is

poorly resolved in RNA sequencing studies and the Drosophila Genetic Reference

Panel (DGRP, see **Fig.** S1) (Mackay et al. 2012; Leader et al. 2018). Because this

region is poorly resolved, it is unclear if our PCR assay might be sensitive to cryptic

sequence variation. However our PCR screen nevertheless confirms that this region

is variable in the wild, and we additionally note that common fly strains seem to

differ in their *BaraA* copy number (Table S1), where extra gene copies correlated

with increased expression after infection (see (Hanson et al. 2021) S10 Fig).

We also recovered two paralogous *Baramicin* genes in *D. melanogaster*

through reciprocal BLAST searches: *CG13749* and *CG30285*, which we name *BaraB*

and *BaraC* respectively (**Fig.** 2B). The three *Baramicin* gene loci are scattered on the

right arm of chromosome II at cytological positions 44F9 (*BaraB*), 50A5 (*BaraA*),

and 57F8 (*BaraC*). These paralogous *Baramicins* are united by the presence of the

IM24 domain. In the case of *BaraB*, we additionally recovered a frameshift mutation

(2R_4821599_INS) causing a premature stop segregating in the DGRP leading to the

loss of IM13 and IM22 relative to the *BaraA* gene structure (**Fig. 2**B); this truncation

is present in the Dmel_R6 genome assembly, but many DGRP strains encode a CDS

with either a standard (e.g. DGRP38) or extended (e.g. DGRP101) IM22 domain (a

DGRP *BaraB* alignment is provided in supplementary data file 2). Moreover, in

contrast to BaraA, the initial IM10-like peptide of BaraB no longer follows a furin

cleavage site, and encodes a serine (R<u>S</u>XR) in its IM10-like motif instead of the

universal proline (R<u>P</u>XR) of *BaraA*-like IM10 peptides across the genus. Each of

these mutations prevents the secretion of classical IM10-like and IM22 peptides by

BaraB. Finally, *BaraC* encodes only IM24 tailed by a transmembrane domain at the

C terminus (TMHMM v2.0 (Krogh et al. 2001)), and thus lacks both the IM10-like

peptides and IM22 (**Fig. 2**B).

BaraB and BaraC are not immune-inducible

BaraA is strongly induced following microbial challenge (Fig. 1), being

predominantly regulated by the Toll pathway with a minor input from the Immune

Deficiency (Imd) pathway (Huang et al. 2020; Hanson et al. 2021). We therefore

assayed the expression of *BaraB* and *BaraC* in wild-type flies, and also flies with

defective Toll (*spz^{rm7}*) or Imd (*Rel^{E20}*) signalling to see if their basal expression relied

on these pathways. Surprisingly, neither gene was induced upon infection

regardless of microbial challenge (**Fig. 3**A and **Fig.** S2A-B). However *BaraC* levels

were consistently reduced in *spz^{rm7}* mutants regardless of treatment (cumulative

data in **Fig.** S2C, p = .005), suggesting *BaraC* basal expression is affected by Toll

signalling. We next generated a novel time course of development from egg to adult

to monitor the expression of the three *Baramicin* genes. We found that expression of

all genes increased over development and reached their highest level in young

adults (**Fig.** 3B). Of note, *BaraB* expression approached the lower limit of our assay's

detection sensitivity at early life stages. However *BaraB* was robustly detected

beginning at the pupal stage, indicating it is expressed during metamorphosis. *BaraC*

expression also increased markedly between the L3 larval stage and pupal stage.

Here we reveal that *BaraA* is part of a larger gene family. While the *BaraA*

gene was first described as an immune effector, the two *Baramicin* paralogs *BaraB*

and *BaraC* are not induced by infection in *D. melanogaster*. Both *BaraB* and *BaraC*

first see increased expression during pupation, and are ultimately expressed at their

highest levels in adults.

Dmel\BaraB is required in the nervous system over the course of development

A simple interpretation of the truncated gene structure and low levels of

BaraB expression is that this gene is undergoing pseudogenization. Indeed, AMP

gene pseudogenization is common in insects including *Drosophila* (Quesada et al.

2005; Rolff and Schmid-Hempel 2016; Hanson, Lemaitre, et al. 2019). To explore

Figure 3: *D. melanogaster* non-immune *Baramicins* have neural functions. A) Only *BaraA* is immune-induced. *BaraB* and *BaraC* do not respond to infection, though basal *BaraC* expression relies on Toll signalling (Fig. S3C). B) Time series of whole animal *Baramicin* expression over the course of development. Expression values are normalized within each gene, with expression in the adult set to 100% (left panel). For context, normalizing each gene to *BaraA* in adults shows that *BaraC* and especially *BaraB* expression is much lower at a whole fly level (right panel). C) *ΔBaraB* mutations generated in this study. *ΔBaraB^{LC1}* is an incidental hypomorph with reduced *BaraB* expression (Fig. S3A), while *ΔBaraB^{LC4}* encodes the intended genetic knock in of a DsRed cassette. Both express DsRed in the eyes, ocelli, and abdomen. D) Partial lethality of *ΔBaraB^{LC1}* hypomorphs or complete lethality of *ΔBaraB^{LC4}* null flies in varied genetic backgrounds. E) Example of a nubbin-like wing, which is phenocopied by *BaraB* gene silencing using the neural driver *elav-Gal4*. F) The *BaraC* gene almost perfectly matches the expression pattern of glia specific *Repo*-expressing cells in single cell RNA sequencing of the adult fly brain (FlyBrainAtlas (Davie et al. 2018)).

BaraB function, we used two mutations for *BaraB* (*ΔBaraB^{LC1}* and *ΔBaraB^{LC4}*,

generously gifted by S.A. Wasserman). These mutations were made using a CRISPR

double gRNA approach to replace the *BaraB* locus with sequence from the pHD-

DsRed vector. The *ΔBaraB^LC1* and *ΔBaraB^LC4* mutations differ in their ultimate effect,

as *ΔBaraB^LC1* is an incidental insertion of the DsRed cassette in the promoter of the

gene. This disruption reduces gene expression, resulting in a hypomorph state (**Fig.**

S3A). The *ΔBaraB^LC4* mutation however deletes the locus as intended, leading to

BaraB null flies (**Fig. 3**C).

We further introgressed both *ΔBaraB* mutations into the DrosDel isogenic

background (referred to as *iso*) for seven generations according to Ferreira et al.

(Ferreira et al. 2014). At the same time, we combined the original *ΔBaraB*

chromosomes with a CyO-GFP balancer chromosome in an arbitrary genetic

background to distinguish homozygous/heterozygous larvae. In all cases, *ΔBaraB^LC4*

homozygotes were entirely lethal during larval development, whereas the

hypomorphic *ΔBaraB^LC1* flies allowed for homozygous adults to emerge (**Fig. 3**D).

We further assessed *ΔBaraB^LC1* hypomorph viability using crosses between

ΔBaraB^LC1/CyO heterozygous females and *ΔBaraB^LC1* homozygous males, which

showed reduced viability and was exacerbated by rearing at 29°C (**Fig. S3**B). Using

our *CyO-GFP* reporter to track genotypes in larvae revealed that the major lethal

phase occurs primarily in the late larval and pupal stages (**Fig. S3**C-F), agreeing with

a role for *BaraB* in larvae/pupae previously suggested by increased expression at

this stage. Some emergent flies also exhibited locomotor defects, and/or a nubbin-

like wing phenotype (FlyBase: FBrf0220532 and e.g. in **Fig. 3**E) where the wings

were stuck in a shrivelled state for the remainder of the fly's lifespan. However, a

plurality of *ΔBaraB^LC1* homozygotes successfully emerged, and unlike their siblings,

had no immediate morphological or locomotory defects. The lifespan of

morphologically normal *iso ΔBaraB^LC1* adults is nevertheless significantly shorter

compared to wild-type flies and *iso ΔBaraB^LC1/CyO* siblings (**Fig. S4**G). We confirmed

these developmental defects using ubiquitous gene silencing with *Actin5C-Gal4 (Act-*

Gal4) to drive two *BaraB* RNAi constructs (*TRiP-IR* and *KK-IR*). Both constructs

resulted in significant lethality and occurrence of nubbin-like wings (**Table** S2).

Genomic deficiency crosses also confirmed significantly reduced numbers of

eclosing *BaraB*-deficient flies at 25°C (n = 114, χ^2 p < .001) and 29°C (n = 63, χ^2 p <

.001) (**Fig. S3**H). Thus full gene deletion is lethal in the larval/pupal transition stage,

and *BaraB* hypomorph flies suffer significant costs to fitness during development,

and have reduced lifespan even following successful eclosion.

These data demonstrate a significant cost of *BaraB* disruption. While whole-

fly *BaraB* expression is low, these results suggest that *BaraB* is not pseudogenized,

and instead performs an integral developmental role. The fact that there is a

bimodal outcome in hypomorph-like *ΔBaraB^{LC1}* adults (either severe defects or

generally healthy) suggests *BaraB* is involved in passing some checkpoint during

larval/pupal development. Flies deficient for *BaraB* may be more likely to fail at this

developmental checkpoint, resulting in either lethality or developmental defects.

The Baramicin paralogs BaraB and BaraC are expressed in the nervous system

We next sought to determine in which tissue(s) *BaraB* is required. A previous

screen using neural RNA interference highlighted *BaraB* for lethality effects (n = 15)

(Neely et al. 2010). Given this preliminary result, and alongside our observed *BaraB*

mutant locomotory defects, we started by silencing *BaraB* in the nervous system at

25°C or at 29°C for greater efficiency using the pan-neural *elav-Gal4* driver both the

TRiP-IR and *KK-IR BaraB-IR* lines. We additionally combined this approach with

UAS-Dicer2 (*Dcr2*) to further strengthen gene silencing as used previously (Neely et

al. 2010)*. In the event there was no lethality, it was expected that emerging

elav>TRiP-IR flies would follow simple mendelian inheritance. However both

elav>TRiP-IR and *elav>Dcr2, TRiP-IR* resulted in partial lethality and occasional

nubbin-like wings (χ^2 p < .02, **Table** S2). Crosses using *KK-IR* used homozygous

flies, and so we did not assess lethality using mendelian inheritance. However using

this construct, no adults emerged when *elav>Dcr2, KK-IR* flies were reared at 29°C.

Rare emergents (N = 11 after three experiments) occurred at 25°C, all of which bore

nubbin-like wings. Using *elav-Gal4* at 29°C without *Dcr2*, we observed greater

numbers of emerging adults, but 100% of flies had nubbin-like wings (**Fig. 3**E,

Table S2). Finally, *elav>KK-IR* flies at 25°C suffered both partial lethality and

nubbin-like wings, but normal-winged flies began emerging (χ^2 p < .001, **Table** S2).

This analysis indicates that *BaraB* is expressed in the nervous system, and this

expression readily explains both the lethality and nubbin-like wing phenotypes.

Moreover, we observed a consistent spectrum of developmental defects using *elav-*

Gal4>BaraB-IR wherein strength of gene silencing correlates with severity of

lethality and wing defect frequency. We additionally investigated the effect of *BaraB*

RNAi using Gal4 drivers in non-neural tissues including the fat body (*c564-Gal4*),

hemocytes (*hml-Gal4*), the gut (esg-Gal4), malpighian tubules (MyO-Gal4), the wing

disc (*nubbin-Gal4*), and in myocytes (*mef2-gal4*) to no effect. We also screened

neural drivers specific for glia (*Repo-Gal4*), motor neurons (*D42-*, *VGMN-*, and *OK6-*

Gal4), and a recently-made *BaraA-Gal4* driver that is expressed in the larval ventral

nervous system (Hanson et al. 2021). However all these *Gal4>BaraB-IR* flies were

viable and never exhibited overt morphological defects.

We also screened for effects of *BaraC* disruption using ubiquitous *Act-Gal4*

and neural *elav-Gal4>Dcr2* for developmental defects. However neither driver

produced overt phenotypes in morphology or locomotor activity (not shown).

Tissue-specific transcriptomic data indicate that *BaraC* is expressed in various

neural tissues including the eye, brain, and the thoracic abdominal ganglion (**Fig.**

S4A), but also the hindgut and rectal pads pointing to a complex expression pattern

(Hammonds et al. 2013; Leader et al. 2018). We next searched FlyBrainAtlas (Davie

et al. 2018) to narrow down which neural subtypes *BaraB and BaraC* were

expressed in. *BaraB* expressing cells were few and showed only low expression in

this dataset. However *BaraC* was robustly expressed in all glial cell types, fully

overlapping the glia marker *Repo* (**Fig. 3**F). To confirm the observation that *BaraC*

was expressed in glia, we compared the effects of *BaraC* RNA silencing (*BaraC-IR*)

using *Act-Gal4* (ubiquitous), *elav-Gal4* (neural) and *Repo-Gal4* (glia) drivers on

BaraC expression. *Act-Gal4* reduced *BaraC* expression to just ~14% that of control

flies (**Fig. S4**B). By comparison *elav-Gal4* reduced *BaraC* expression to ~63% that of

controls, while *Repo-Gal4* led to *BaraC* levels only 57% that of controls (overall

controls vs. neural/glia-IR, p = .002).

Collectively, our results support the notion that *BaraC* is expressed in the

nervous system, and are consistent with *BaraC* expression being most localized to

glial cells.

Extensive genomic turnover of the Baramicin gene family

Our results thus far show that *BaraA*-like genes are consistently immune-

induced in all *Drosophila* species (**Fig.** 1), however the two paralogs *Dmel\BaraB*

and *Dmel\BaraC* are not immune-induced, and are truncated in a fashion that

deletes some or all of the antifungal IM10-like peptides (**Fig. 2**B). These two

Baramicins are now enriched in the nervous system (**Fig. 3**E-F). In the case of *BaraB*,

a role in the nervous system is evidenced by severe defects recapitulated using pan-

neural RNA silencing. In the case of *BaraC*, nervous system expression is evidenced

by a clear overlap with *Repo*-expressing cells.

While *BaraA*-like genes are conserved throughout the genus *Drosophila*,

BaraB is conserved only in Melanogaster group flies, and *BaraC* is found only in

Melanogaster and Obscura group flies, indicating that both paralogs stem from

duplication events of a *BaraA*-like ancestor (**Fig. 4**). To determine the ancestry of

each *D. melanogaster Baramicin* gene, we traced their evolutionary history by

analyzing genomic synteny through hierarchical orthologous groups (Train et al.

2019). Ancestry tracing revealed that these three loci ultimately stem from a single-

locus ancestor encoding only one *Baramicin* gene that resembled *Dmel\BaraA* (**Fig.**

4A). This is evidenced by the presence of only a single *BaraA*-like gene in the

outgroup *S. lebanonensis*, and also in multiple lineages of the subgenus Drosophila

(**Fig. 4**B). Indeed, the general *BaraA* gene structure encoding IM24, tandem repeats

of IM10-like peptides, and IM22 is conserved in *S. lebanonensis* and all *Drosophila*

species (**Fig. 4**C). On the other hand, the *Dmel\BaraC* gene comes from an ancient

duplication restricted to the subgenus Sophophora, and *Dmel\BaraB* resulted from a

more recent duplication found only in the Melanogaster group (**Fig. 4**B).

We originally recovered outgroup *Baramicins* assayed for immune induction

(**Fig.** 1) through reciprocal BLAST searches. However following genomic synteny

analysis, we realized that the *D. willistoni BaraA*-like gene *Dwil\GK10648* is syntenic

with the *Dmel\BaraC* locus (**Fig. 4**A), yet this gene is immune-induced (**Fig.** 1D) and

retains a *BaraA*-like gene structure (**Fig. 4**C). On the other hand, *Dwil\GK10645* is

found at the locus syntenic with *BaraA*, but has undergone an independent

truncation to encode just an IM24 peptide (similar to *Dmel\BaraC*). Thus these two

D. willistoni genes have evolved similar to *D. melanogaster BaraA/BaraC*, but in a

vice versa fashion. This suggests a pattern of convergent evolution with two key

points: **i)** the duplication event producing *Dmel\BaraA* and *Dmel\BaraC* originally

copied a full-length *BaraA*-like gene to both loci, and **ii)** the derivation of an IM24-

specific gene structure has occurred more than once (*Dmel\BaraC* and

Dwil\GK10645). Indeed, another independent IM24-specific *Baramicin* gene is

present in *D. virilis* (*Dvir\GJ25897*), which is a direct sister of the *BaraA*-like gene

Dvir\GJ21309 (the signal peptides of these genes is identical at the nucleotide level,

and see **Fig. 4**C). Thus *Baramicins* in both *D. willistoni* and *D. virilis* have

convergently evolved towards an IM24-focused protein structure resembling

Dmel\BaraC. We checked the expression of these truncated *Baramicins* in each

species upon infection. As was the case for *Dmel\BaraC*, neither gene is immune-

induced (**Fig. S5**A-C). Given the glial expression of *Dmel\BaraC,* we reasoned that

the heads of adult flies (rich in nerve tissue) should be enriched in *BaraC* compared

to whole animals. Indeed we saw a significant enrichment of *BaraC* in the heads of *D.*

melanogaster males compared to whole flies, which was not the case for *BaraA* (**Fig.**

S5D). When we checked the heads of *D. willistoni* and *D. virilis*, we indeed saw a

consistent and significant enrichment in the head for the IM24-specific genes

Dwil\GK10645 and *Dvir\GJ25897*, while *BaraA*-like genes were more variable in

expression (**Fig. S5**E-F).

Thus, multiple independent IM24-specific *Baramicins* are not immune

induced and are more specifically enriched in the head. In the case of *Dmel\BaraC*,

this is likely due to expression in glia. Strikingly, we observe a parallel evolution of

expression pattern and gene structure in *Baramicins* of *D. willistoni* and *D. virilis*.

These expression data are summarized in **Fig. 4**C. Genomic synteny shows the gene

structure and immune expression of *BaraA* are the ancestral state, and *Dmel\BaraB*

and *Dmel\BaraC* are paralogs derived from independent duplication events.

Figure 4: *Baramicin* evolutionary history. A) Detailed map of genomic neighbourhoods in the outgroup drosophilids *S. lebanonensis, D. willistoni,* and *D. melanogaster*, detailing inferred duplication, inversion, and translocation events. Gene names are given as found in *D. melanogaster*. B) Cladogram and genomic loci further detailing the series of events leading to the extant *Baramicin* loci of *Drosophila* species. Loci in *S. lebanonensis* and flies in the subgenus *Drosophila* encode only one *Baramicin* gene, indicating the ancestral drosophilid likely encoded only one *Baramicin*. C) IM24-specific *Baramicins* arose from convergent evolution both in gene structure and expression profile. Genomic loci are described here as *ATP8A* or *stum* reflecting prominent genes neighbouring the *Baramicins* (see Fig. 4A). Expression dynamics relating to immune-induction or enrichment in the head (checked boxes) are shown in Fig. 2 and Fig. S5. The *Baramicin* loci in *D. willistoni* are syntenic with *D. melanogaster*, but evolved in a vice versa fashion (purple arrow). The *D. virilis Baramicins GJ21309* and *GJ25897* are direct sister genes (100% identity at N-terminus).

Residue 29 in the IM24 domain evolves in lineage-specific fashions

390
Multiple independent *Baramicin* genes have lost both IM10-like and IM22
peptides, converge on loss of immune induction, and are enriched in the head. Taken
together, these truncations and expression patterns suggest that the IM10-like
peptides and IM22 are strictly useful during the immune response, consistent with a
recently described antifungal role for IM10-like peptides (Hanson et al. 2021).
Inversely, non-immune *Baramicin* genes have repeatedly and independently
truncated to encode primarily IM24. We could not generate a reasonable model of
the IM24 peptide conformation using Phyre2 (Kelley et al. 2015), QUARK, or
TASSER protein modelling methodologies (Zhang et al. 2016). AlphaFold (Jumper et
al. 2021) also has only low confidence estimates for the mature structure of *D.*
melanogaster Baramicins. The IM24 domain unites the *Baramicin* gene family,
making its apparent non-immune functional roles in *BaraB* and *BaraC* intriguing.
Failing to model the protein, we next asked if we could highlight any residues in this
traditionally immune peptide that might correlate with immune or non-immune
gene lineages to gain insight into what governs the IM24-specific gene preference
for neural expression.
To do this, we screened for positive selection (elevated non-synonymous
mutation rate) in the IM24 domain using the HyPhy package implemented in
Datamonkey.org (Delport et al. 2010) using separate codon alignments of *Baramicin*
IM24 domains beginning at their conserved Q[1] starting residue. As is recommended
with the HyPhy package (Delport et al. 2010), we employed multiple statistical
approaches including Likelihood (FEL), Bayesian (FUBAR), and Count-based (SLAC)
analyses to ensure patterns in selection analyses were robust to different methods
of investigation. Specifically, we used locus-specific alignments (e.g. genes at the
stum locus in **Fig. 4**B were all analyzed together) independent of overall gene
structure to ensure IM24 evolution reflected locus-specific evolution. FEL, FUBAR,
and SLAC site-specific analyses each suggest strong purifying selection in many
residues of the IM24 domain (data in **supplementary data file 3**), agreeing with the
general protein structure of IM24 being broadly conserved (**Fig.** 5A). However one
residue (site 29) was consistently highlighted as evolving under positive selection

using each type of statistical approach for genes located at the Sophophora *ATP8A*

locus (*BaraA* genes and *Dwil\GK10645*: p-adj < .05; **Fig. 5**A). This site is universally

Proline in *Baramicin* genes located at the stum locus (*BaraC*-like) and in the

outgroup *S. lebanonensis*, but is variable in both the *BaraA* (commonly Threonine)

and *BaraB* (commonly Valine) lineages. Both the *S. lebanonensis* and the two *D.*

willistoni Baramicins encode Proline at site 29 independent of gene structure,

suggesting Proline is the ancestral state. We also note that two sites on either side of

site 29 (site 27 and site 31) similarly diverge by lineage in an otherwise highly

conserved region of the IM24 domain. FUBAR analysis (but not FEL or SLAC)

similarly found evidence of positive selection at site 31 in the *BaraA* locus genes (p-

adj = .026). Thus this neighbouring site could also be evolving in a non-random

fashion. Similar analyses of the *BaraB* and *stum* loci *Baramicins* did not find evidence

of site-specific positive selection.

While the structure of IM24 is unknown, HyPhy analysis highlights site 29 as

a key residue in IM24 that diverged in *Baramicin* lineage-specific fashions. This

ancestrally Proline residue has settled on a Threonine in most *BaraA*-like genes of

Obscura and Melanogaster group flies, and a Valine in most *BaraB* genes, which are

unique to the Melanogaster group.

Another IM24 domain in Baramicin lineages varies through relaxed selection

Visual inspection of aligned IM24 proteins makes it evident that the overall

IM24 domain is broadly conserved, except in sites 40-48 (**Fig. 5**A). This motif

uniquely encodes the residues [40]HHASSPAD[48] in *Dmel\BaraB*. Given the severe cost

of *BaraB* mutation, intriguingly the H[40] and D[48] residues are not found in any other

Baramicin genes. The three C-terminal residues of this motif are also diagnostic for

each gene lineage (*BaraA, BaraB,* and *BaraC* have RGE, PXE, or (S/N)GQ

respectively; **Fig. 5**A). However even with additional branch-site selection analyses

(aBSREL and BUSTED (Murrell et al. 2015)), we found no evidence of positive

selection at the [40]HHASSPAD[48] homologous domain (supplementary data file 3).

Thus while IM24 residues 40-48 are variable across lineages, this motif is not

evolving with elevated non-synonymous change. We suspect instead that this motif

is diversifying due to relaxed selection as six of nine sites in the *BaraA* locus analysis
failed to reach significance (p < .05) for purifying selection in e.g. SLAC analysis
(supplementary data file 3). It is nevertheless striking that this region is so variable
given the conservation of residues upstream and downstream of sites 40-48. This
pattern should have implications on the IM24 protein functional motifs, which
future protein folding investigations may decipher.
458
Overt IM24 structural change best explains loss of immune induction
Site 29 varies in lineage-specific fashions, encoding a derived Valine residue
in most species' BaraB IM24 domains. If the Valine at site 29 explains the *BaraB*
functional divergence relative to its sister *BaraA* lineage, this could suggest that
BaraB has long functioned in a neural role common to most Melanogaster group
flies. To this end, we performed infection experiments in diverse species across the
Melanogaster group to see if their *BaraB* genes had similarly lost immune induction
(see **Fig.** S6 for qPCR data). Surprisingly, we instead found that the non-immune
expression of *Dmel\BaraB* is extremely recent, as Melanogaster sister species like *D.*
sechellia and *D. mauritiana* nevertheless encode immune inducible *BaraB* loci
(summary in **Fig.** 5B). However, we also found that *D. simulans BaraB* lacked
immune induction, despite being most closely related to *D. sechellia*. This drew our
attention instead to the overall protein structure of the various *BaraB* genes. A
striking feature of the Dmel\BaraB protein is the absence of a signal peptide
structure (**Fig.** 2B). Signal peptide sequence is conserved in all *Baramicin* lineages,
except in *BaraB* of *D. melanogaster* and also *D. simulans* (last common ancestor
~3mya (Chakraborty et al. 2021)). Indeed despite *D. simulans* being more closely
related to *D. sechellia* and *D. mauritiana*, both *Dmel\BaraB* and *Dsim\BaraB* encode a
homologous N-terminus of parallel length (**Fig.** 5A). We also found that *D. yakuba*
BaraB is not immune-responsive, but note that *D. yakuba* has an insertion upstream
of residue 40 that elongates the IM24 domain (**Fig.** 5B black X boxes), and its sister
species *D. erecta* encodes multiple indels and premature stops suggesting *BaraB* is
pseudogenized in this lineage. While an insertion is also present in *D. sechellia*
BaraB in the signal peptide, this is still predicted to allow secretion (SignalP 5.0),
suggesting *Dsec\BaraB* is a functional immune protein.

Figure 5: *BaraB* and IM24 rapid evolution. A) Evolution of the signal peptide and core IM24 domain. Residue highlighting indicates agreement with *Dmel\BaraB*. Arrows indicate site 29 and the *D. melanogaster* [40]HHASSPAD[48] domain. Insertion events in *D. yakuba* and *D. sechellia BaraB* are denoted as XX to save space. The *D. sechellia* signal peptide is predicted to remain functional (SignalP 5.0). The cladogram on the left shows genomic relatedness (by speciation and locus) independent of sequence similarity. Background colouring is included to show birth of novel *Baramicin* loci/lineages. B) *BaraB* immune or non-immune expression by phylogeny. The *D. erecta BaraB* gene is pseudogenized by multiple premature stop codons, and the *D. yakuba* gene is not immune-induced and encodes a 9-residue insertion in the IM24 peptide bordering the HHASSPAD domain (see XX site in A). However the *BaraB* genes of *D. suzukii*, and both *D. mauritiana* and *D. sechellia* remain inducible by infection, while the *BaraB* genes of *D. simulans* and *D. melanogaster* are not and are expressed at very low levels (Fig. S6). This pattern suggests that *BaraB* of *D. melanogaster* (and *D. simulans*) acquired its non-immune role only recently, and is correlated with the loss of the *BaraB* signal peptide.

Loss of the BaraB signal peptide is therefore more specifically associated

with loss of immune expression in the Melanogaster species complex (*D. simulans, D.*

sechellia, D. mauritiana, and *D. melanogaster*). The last common ancestor of *D.*

simulans, D. sechellia, and *D. mauritiana* is estimated to be just ~250,000 years ago,

and these species diverged from *D. melanogaster* ~3 million years ago (Chakraborty

et al. 2021). The fact that *D. simulans* uniquely encodes this *Dmel\BaraB*-like

sequence suggests it was either introgressed from one species to the other prior to

the complete development of hybrid inviability, or reflects incomplete lineage

sorting of this locus in the Melanogaster species complex. In either case, this points

to an extremely young age for the novel function of *Dmel\BaraB* in the nervous

system. This loss of the signal peptide also occurs alongside a segregating allele that

truncates the mature *Dmel\BaraB* sequence (**Fig.** 2B), a pattern commonly found in

Baramicins derived for neural expression (**Fig.** 4C). This reinforces the fact that the

Dmel\BaraB gene had an immune function so recently that some wild flies still

produce the immune-relevant IM10-like and IM22 Baramicin peptides despite

neural expression of *BaraB*.

BaraB evolution therefore reinforces that the core driver of *Baramicin*

functional divergence is not based on minor sequence changes, but rather correlates

with overt protein structural change. *BaraB* has a mutation affecting secretion, while

BaraC now encodes a transmembrane domain, which should cause it to insert itself

into either its endogenous glial cell membrane or a neighbouring cell (e.g. a neuron).

In both cases IM24 is preferentially expressed and localized to the nervous system.

Discussion

We recently showed that *BaraA* deletion causes infected flies to display an

erect wing behavioural phenotype (Hanson et al. 2021). Notably, flies displayed

erect wing even when heat-killed bacteria were injected, indicating this behaviour

depends only on the triggering of the immune response in the absence of *BaraA*, and

not on active infection. Thus *BaraA* likely interacts with some host target(s) to

prevent this behaviour during the immune response.

Here we find the *Baramicin* IM24 domain has a predilection for interactions

with the nervous system. We speculate that the immune-mediated production of

IM24 by *BaraA* could protect the nervous system from autoimmune activity, which

occurs in the absence of *BaraA* to cause erect wing display. A notable aspect of this

hypothesis is it proposes that some peptides of AMP genes are responsible for

microbe killing, while others are co-secreted with the intent of preventing

autoimmune toxicity. For now this remains speculation, however it will be

interesting to clarify the mature structure of IM24 and determine what partner(s)

IM24 binds to. In this regard, we highlight site 29 as an important residue for IM24

function, and suggest that while residues 40-48 are variable, the sequence at this

motif does not experience the same sort of evolutionary selection. One possibility to

explain these evolutionary patterns is that site 29 is exposed in some way that an

IM24 binding partner uses. Meanwhile residues 40-48 could act as a sort of linker

between the two ends of the IM24 domain where the length of these residues is

important, but the exact sequence is malleable. Indeed, we found that *D. yakuba*

BaraB independently lost immune induction alongside an insertion at site 40.

Antimicrobial peptide genes have recently been implicated by a number of

studies in neural functions, regulating processes like memory, sleep, taste aversion,

behaviour, and neurodegeneration. The properties of these immune peptides share

many features with classic neuropeptides, including cationic charge and

amphipathicity (Brogden et al. 2005). Nevertheless it is unclear why AMPs can play

dual roles in either immunity or neural function. By characterizing the evolution of

the *Baramicin* gene family, we provide insight on how an ancestrally immune AMP

gene has adapted itself for neural function on a repeated basis. The mechanism

through which *Baramicin* achieves its neural effect is specific to the IM24 domain, as

the antifungal IM10-like peptides and IM10-related IM22 peptide are consistently

lost in *Baramicin* lineages now specific to neural expression. This realization is made

possible by the exaggerated polypeptide structure of *Baramicins,* which focuses the

interpretation on how different peptides of this modular protein structure may play

roles in either neurology or immunity. Other AMP genes similarly encode

polypeptides, but do not have so many tandem repeats of identical peptides and so

the polypeptide nature of their precursor protein is easily glossed over. The

polypeptide nature of these genes is lacking from the current conversation on AMP

involvements in the nervous system, despite AMP genes of fruit flies and other

animals encoding furin-cleaved polypeptides (Hanson and Lemaitre 2020).

One human AMP recently implicated in chronic neuroinflammatory disease is

the Cathelicidin LL-37 (Lee et al. 2015; De Lorenzi et al. 2017; Moir et al. 2018). Like

Baramicin, the *Cathelicidin* gene family is unified by its N-terminal "Cathelin"

domain. However to date no one has described antimicrobial activity of the Cathelin

domain in vitro (Zanetti 2005). Instead, *Cathelicidin* research has focused almost

exclusively on the mature peptide LL-37 at the C-terminus of mammalian

Cathelicidin genes. Reflecting on *Baramicin* evolution and the implication of

Cathelicidin in neurodegenerative diseases, what does the Cathelin domain do?

While this study was conducted in fruit flies, we hope we have emphasized the

importance of considering each peptide of AMP genes for in vivo function. This is

relevant to neural processes even if the gene is typically thought of for its role in

innate immunity. Indeed, recent studies of *Drosophila* AMPs have emphasized that in

vitro activity does not always predict the interactions that these genes can have in

vivo (Clemmons et al. 2015; Hanson, Dostálová, et al. 2019). Care should be taken

not to conflate in vitro activity with realized in vivo function. Most studies focus on

AMPs specifically in an immune role, but this is akin to 'looking for your keys under

the streetlight.' To understand AMP functions in vivo, genetic approaches will be

necessary that allow a more global view of gene function.

In summary, we characterize how an ancestral AMP-encoding gene has

repeatedly evolved for neural expression by truncating its protein sequence to

express just one peptide. It will be interesting to consider the functions of AMP

genes in neural processes not simply at the level of the gene, but at the level of the

mature peptides produced by that gene. Given the polypeptide of many AMP gene

structures and commonalities between AMPs and neuropeptides, these canonical

immune effectors may be adapted for neural function more often than appreciated.

Acknowledgements

We would like to thank Maria Litovchenko for advice, Ana Marija Jakšić for generously providing DGRP flies, Rob Unckless for stimulating discussion, Huang et al. (Huang et al. 2020) for collaborative cooperation, Brian McCabe for consultation, and Florent Masson, Hannah Westlake, and the anonymous reviewers and the editors at MBE for commentary on our initial manuscript. This research was supported by Sinergia grant CRSII5_186397 awarded to Bruno Lemaitre. The *BaraB^{LC1}* and *BaraB^{LC4}* mutations were graciously provided by Steven Wasserman and generated by Lianne Cohen, who we also thank for their critical involvement in characterizing *Baramicin A*.

Materials and Methods

DGRP population screening and bioinformatics analyses

Genomic sequence data were downloaded from GenBank default reference assemblies and Kim et al. (Kim et al. 2021), and DGRP sequence data from http://dgrp2.gnets.ncsu.edu/ (Mackay et al. 2012). Sequence comparisons and alignment figures were prepared using Geneious R10 (Kearse et al. 2012), Prism 7, and Inkscape. Alignments were performed using MUSCLE or MAFFT followed by manual curation, and phylogenetic analyses were performed to validate sequence patterns using the Neighbour Joining, PhyML, RaxML, and MrBayes plugins in Geneious. *BaraA* copy number screening was performed using primers specific to the duplication and *CG30059* control primers for DNA extraction (primers in supplementary data file 5). We found a significant correlation between *BaraA* PCR status and variant sites starting at 2R_9293471_SNP and extending to 2R_9293576_SNP (Pearson's correlation matrix: $0.0001 < \text{p-value} < 0.005$ at all nine sites), however the status of genetic variants at this site is poorly resolved and so we cannot be confident that our ~14% estimate for the *BaraA* duplication in the DGRP would hold true if long-read sequencing was employed. DGRP annotation of the *BaraA* locus in Fig. S1 was generated using the UCSC *D. melanogaster* DGRP2 genome browser. Selection analyses were performed using the HyPhy package implemented in datamonkey.org (Delport et al. 2010). Codon alignments of the IM24 domain used in **Fig. 5**A are included as a .fasta file in supplementary data file 3 alongside outputs from FEL, FUBAR, SLAC, and aBSREL selection analyses.

Fly genetics

The *BaraB^{LC1}* and *BaraB^{LC4}* mutations were generated using CRISPR with two gRNAs and an HDR vector by cloning 5' and 3' region-homologous arms into the pHD-DsRed vector, and consequently *ΔBaraB* flies express DsRed in their eyes, ocelli, and abdomen. The following PAM sites were used for CRISPR bordering the *BaraB* region. Slashes indicate the cut site: 5': GCGGGCAACAGATGTGTTCA/GGG 3': GTCCATTGCTTATTCAAAAA/TGG. These mutants were generated in the laboratory of Steve Wasserman by Lianne Cohen, who graciously allowed their use in this study. All fly stocks including Gal4 and RNAi lines are listed in **supplementary data file 4**. Experiments were performed at 25°C unless otherwise

indicated. When possible, genetic crosses of 6-8 males and 6-8 females were
performed in both directions to test for an effect of the X or Y chromosomes on
BaraB-mediated lethality; crosses in both directions yielded similar results in all
cases and reported data are pooled results. Fly diet consisted of a nutrient-rich lab
standard food: 3.72g agar, 35.28g cornmeal, 35.28g yeast, 36mL grape juice, 2.9mL
propionic acid, 15.9mL moldex, and H_2O to 600mL.
638
Infection experiments
Bacteria and yeast were grown to mid-log phase shaking at 200rpm in their
respective growth media (LB, BHI, or YPG) and temperature conditions, and then
pelleted by centrifugation to concentrate microbes. Resulting cultures were diluted
to OD = 200 at 600nm before infections to measure gene expression. The following
microbes were grown at 37°C: *Escherichia coli strain 1106* (LB) and *Candida albicans*
(YPG)*. Micrococcus luteus* was grown at 29°C in LB. For **Fig.** 1 and S2, pooled fly
samples were collected either 6 hours post-infection (*E. coli*) or 24 hours post-
infection (*C. albicans, M. luteus*) prior to RNA extraction on pools of 5 adult males.
These timepoints correspond to the maximal expression inputs of the Imd (6hpi) or
Toll (24hpi) NF-κB signalling pathways, which are most specifically induced by
Gram-negative bacteria (Imd) or Gram-positive bacteria or fungi (Toll) (Lemaitre et
al. 1997). Flies were pricked in the thorax as described in (Hanson, Dostálová, et al.
2019).
RNA extractions were performed using TRIzol™, Ambion DNAse treatment,
and PrimeScript RT according to manufacturer's protocols. RT-qPCR was performed
using PowerUP SYBR Green master mix with primers listed in supplementary data
file 5. Gene expression differences were analyzed using the PFAFFL method (Pfaffl
2001). For gene expression experiments requiring dissection of heads, pools of 20
males were used for either whole flies or heads dissected in ice-cold PBS and
transferred immediately to a tube kept on dry ice.
660
661   ### *Selection analysis using HyPhy package*
662
Codon aligned nexus tree files were generated using either the Neighbour-
joining (1000 bootstraps) or PhyML (100 bootstraps) methods including proteins
beyond those shown in **Fig. 5**. These tree files were analyzed using the HyPhy
package with only 174nt pertaining to just the IM24 domain codons included. The
cladogram in **Fig. 5**A is manually drawn from known species divergences (Kim et al.
2021). Use of either tree building method was chosen for convenience to best reflect
known lineage sorting, as use of just 174nt was too information poor to resolve
exact phylogenetic relatedness reliably. Tree files were qualitatively screened to
ensure topologies broadly matched known species sortings, and thus ensure only
relevant comparisons were made given the genomic synteny analysis in **Fig. 4** is
principally informative of true gene lineages. HyPhy analyses were run separately
for each *Baramicin* lineage within their clade, defined by genomic synteny; i.e. based
on locus (e.g. ATP8A locus), and not considering convergent gene structures. We
used three site-specific analyses (FEL, FUBAR, and SLAC) that use three independent

statistical approaches (Likelihood, Bayesian, and Count-based methods
respectively). We also employed both BUSTED and aBSREL branch-site analyses,
which are likelihood methods that differ in their approach of testing whole-
phylogeny selection or branch-specific comparisons respectively; an anology might
be performing analysis of variance (ANOVA) at the level of the entire ANOVA, or
comparing multiple groups against each other and subsequently using multiple test
correction. Each tree was rooted using the *Scaptodrosophila lebanonensis Baramicin*
as an outgroup with ancestral characteristics; we did not include *Baramicins* of the
subgenus Drosophila as including these resulted in long-branch attraction of the
Willistoni group *Baramicins* to subgenus Drosophila lineages, which would
confound relevant phylogenetic comparisons. When applicable, all internal branches
were assessed for potential selection. For *Baramicins* of the *ATP8A* locus, one site
(site 29) was highlighted as experiencing positive selection using FEL, FUBAR, and
SLAC analyses (p-adj = .011, .013, and .039 respectively). Additionally, site 31 was
also highlighted by FUBAR (p-adj = .026), but not FEL or SLAC analyses (p-adj > .05).
BUSTED analysis also supported diversifying selection in the *BaraA* lineage (ATP8A
locus, LRT p-adj = .008), indicating at least one site on at least one test branch has
experienced diversifying selection within the ATP8A lineage. The aBSREL branch-
site analysis specifically highlights the branch distinguishing the Willistoni group
Baramicins from the other Sophophora species (p-adj = .0045), suggesting variation
between these branches drives the signals of diversifying selection in the BUSTED
analysis. This result is intuitive, as we find a parallel but opposite evolution of
Baramicin protein structure in *Baramicins* of the *ATP8A* locus in *D. willistoni*
compared with *Baramicins* of other Sophophora species. Furthermore, in whole-
gene phylogenies, both *D. willistoni Baramicins* cluster together, supporting the
notion that these two daughter genes have evolved independent from the selection
that shaped the orthologues of *Dmel\BaraA* and *Dmel\BaraC*, also seen in qPCR data
that showed both genes were significantly enriched in the head (**Fig.** S6). This
phylogenetic clustering of the two *D. willistoni Baramicins* holds true when
additional *Baramicins* from recently sequenced genomes of the Willistoni group are
included (from (Kim et al. 2021) in supplementary data file 3), indicating this is
characteristic of the Willistoni group lineage and not specific to *D. willistoni*.

Supplementary figures and tables

Figure S1: The *BaraA* locus is poorly resolved in DGRP genome assemblies. The *BaraA1* and *BaraA2* gene regions are totally devoid of mapped variants (dashed boxes). We speculate this is due to an artefact during genomic assembly, where reads mapping equally to the two identical *BaraA* genes were discarded as non-specific. This would explain why *BaraA* is typically discarded in RNAseq datasets using such measures in their pipeline, but not in microarray data from De Gregorio et al. (De Gregorio et al. 2002) where it is called "IM10."

Genotype	Inferred copy #	Gtype	Junction PCR band	CG30059 band
Wild-type	2	*OR-R*	**YES**	**YES**
	2	*iso w1118*	**YES**	**YES**
	2	*Exelexis*	**YES**	**YES**
	1	*w VDRC*	NO	**YES**
	1	*yw*	NO	**YES**
	1	*Canton S*	NO	**YES**

Table S1: the *BaraA* duplication is variable in common lab stocks. Inferred *BaraA* copy numbers are given for six different lab stocks including *Oregon R* (*OR-R*), *iso DrosDel* (*iso w^{1118}*), *Exelexis*, a *w^{1118}* line from the VDRC, a *yellow white* (*yw*) line used previously by our group, and *Canton S*. All samples were additionally screened for *CG30059* as a DNA extraction positive control.

Figure S2: Additional assays of Baramicin expression upon infection. A) Neither *BaraB* nor *BaraC* are regulated by the Imd pathway, which is specifically stimulated by *E. coli* infection. **B)** Neither *BaraB* nor *BaraC* are induced after infection by *M. luteus*. **C)** *BaraC* levels were consistently depressed in *spz^{rm7}* flies in the unchallenged condition (UC) or upon infection with *C. albicans* (**Fig.** 3B) or *M. luteus* (**Fig.** S4B). Data here are pooled for *iso* wild type or *iso spz^{rm7}* flies without regard for infection treatment (student's t, p = .005).

738
Figure S3: ***BaraB*** **mutation is highly deleterious, even in** ***ΔBaraB^{LC1}***
hypomorphs. A) Diagram of *BaraB* mutant loci and qPCR showing the *ΔBaraB^{LC1}* is
a hypomorph mutation. Under our normal qPCR assay conditions, *BaraB* expression
is not detected in *ΔBaraB^{LC1}* homozygotes. However using highly concentrated cDNA
beyond our assay's valid range (100ng/10μL reaction), we could detect *BaraB*
transcript in *ΔBaraB^{LC1}* flies. Quantification shown here is intended only to show

that *BaraB* transcript can be recovered from *ΔBaraB^{LC1}* homozygotes, and to give a
sense of relative whole-fly expression levels. **B)** Emergent frequencies of *ΔBaraB^{LC1}*
flies at different temperatures. **C)** Aborted pupae (yellow arrows) are a common
occurrence in *ΔBaraB* vials, and sometimes contain fully-developed adults that
simply never eclosed. In **D-F:** ns = not significant, * = p < .05, *** p = < .001. **D)** The
ratio of *ΔBaraB^{LC1}*/CyO-GFP to *ΔBaraB^{LC1}* homozygous larvae drops between the S2
and S3 larval stages (χ^2, p = .515 and p = .012 respectively). **E)** Frequency of
successfully eclosing adults using *BaraB^{LC4}*/CyO-GFP flies. **F)** Frequency of
successfully eclosing adults using *BaraB^{LC1}*/CyO-GFP flies. **G)** *BaraB* mutation
negatively affects lifespan. *iso ΔBaraB^{LC1}* homozygotes suffer reduced lifespan even
relative to their *iso ΔBaraB^{LC1}/CyO* siblings. By comparison, *iso ΔBaraA* flies that
used the same vector for mutant generation live as wild-type. ATM[8] flies suffer
precocious neurodegeneration and are included as short-lived controls (Petersen et
al. 2013). **H)** *ΔBaraB^{LC1}* crossed to the genomic deficiency line (*Df(9063)*) supports a
partial-lethal effect of *BaraB* mutation.
760

Parents and temperature	Offspring	Sex	# eclosed	χ^2	p-value
TRiP					
y1 v1; P{TRiP.HMJ23624}attP40/CyO Act-Gal4/CyO-GFP ; + 25℃	Act-Gal4 / CyO	m	26	25.687	**p < .001**
		f	30		
	TRiP / CyO	m	26		
		f	29		
	Act-Gal4 > BaraB-IR{TRiP}	m	2		
		f	18		
y1 v1; P{TRiP.HMJ23624}attP40/CyO male elav>Dcr2 ;; female 25℃	elav>Dcr2 ; +/CyO	m	67	10.037	**p < .02**
		f	62		
	elav>Dcr2 ; +/BaraB-IR	m	38		
		f	47		
y1 v1; P{TRiP.HMJ23624}attP40/CyO male elav-Gal4> ;; female 25℃	elav-Gal4; +/CyO	m	29	13.593	**p < .01**
		f	46		
	elav-Gal4 ; +/BaraB-IR	m	21		
		f	22		
KK					
P{KK112854}VIE-260B Act-Gal4/CyO-GFP ; + 25℃	Act-Gal4 ; +/CyO	m	27	45.187	**p < .001**
		f	55		
	Act-Gal4 > BaraB-IR{KK}	m	14		
		f	11		
P{KK112854}VIE-260B male elav>Dcr2 ;; female 25℃	elav>Dcr2 ; +/BaraB-IR	m	0	11.000	**p < .001**
		f	11		
P{KK112854}VIE-260B male elav-Gal4> ;; female 29℃	normal wing	m	0	87.766	**p < .001**
		f	0		
	nubbin-like wing	m	28		
		f	57		
P{KK112854}VIE-260B male elav-Gal4> ;; female 25℃	normal wing	m	37	32.133	**p < .001**
		f	47		
	nubbin-like wing	m	31		
		f	5		

761
762 **Table S2: *BaraB* RNAi summary statistics.** Crosses used either the TRiP or KK
763 *BaraB-IR* lines, driven by either *Actin5C-Gal4* or *elav-Gal4,* sometimes including *UAS-*
764 *Dcr2*. Rearing at 29°C and inclusion of *UAS-Dcr2* increases the strength of RNA
765 silencing.
766

767
Figure S4: *BaraC* is expressed in the nervous system, but also the hindgut and rectal pads. A) FlyAtlas2 expression data for *BaraC*. **B)** RT-qPCR of *BaraC* in whole flies using different Gal4 drivers to express *BaraC* RNAi. *BaraC* is knocked down by both the *elav-Gal4* and *Repo-Gal4* nervous system drivers. Cumulatively, nervous system drivers significantly depress *BaraC* expression compared to *BaraC-IR* controls (student's t, p < .01). Ubiquitous knockdown using *Act>BaraC-IR* provides a comparative knockdown to better understand the strength of nervous system-specific knockdowns at the whole fly level.

777
Figure S5: RT-qPCR of *Baramicin* genes in diverse species. A-C) Independent
IM24-specific genes in *D. melanogaster* (A), *D. willistoni* (B), and *D. virilis* (C) are not
induced by infection. *BomBc3* is included as an immune-induced control. **D-F)** The
independent IM24-specific genes (blue) of *D. melanogaster* (D), *D. willistoni* (E), and
D. virilis (F) are each enriched in the head relative to whole flies. *BaraA-like* genes
(orange) were expressed more stochastically in the head, but also generally showed
an enrichment pattern relative to whole flies (not always significant). Each data
point represents an independent pooled sample from 20 male flies. Data were
analyzed using one-way ANOVA with Holm's-Sidak multiple test correction.
787

788
Figure S6: The *D. melanogaster BaraB* gene acquired its non-immune role only
recently. A) Cladogram of the Melanogaster species group. The presence of a
functional signal peptide (Fig. 5A), and the disruption of the *D. yakuba* IM24 peptide
by an in-frame insertion is noted. A summary of whether *BaraB* is an immune-
induced orthologue (B-G) is annotated. **B-G)** *Baramicin* expression data from
Melanogaster group flies either unchallenged or infected with *M. luteus*. *BaraB* is
immune-induced in *D. suzukii, D. sechellia,* and *D. mauritiana*, but not in *D. simulans*
and *D. melanogaster,* which both lack signal peptide structures. *Drosophila yakuba*
BaraB is not immune-induced (C), has an insertion event in its IM24 peptide (Fig.
5A), and its sister species *D. erecta* has pseudogenized its *BaraB* orthologue (Fig.
5B), suggesting pseudogenization may explain the lack of immune induction in *D.*
yakuba BaraB.

References:

Abbott A. 2020. Are infections seeding some cases of Alzheimer's disease? *Nature* 587:22–25.

Barajas-azpeleta R, Wu J, Gill J, Welte R. 2018. Antimicrobial peptides modulate long-term memory. *PLoS Genetics*:1–26.

Brogden KA, Guthmiller JM, Salzet M, Zasloff M. 2005. The nervous system and innate immunity: The neuropeptide connection.

Casteels-Josson K, Capaci T, Casteels P, Tempst P. 1993. Apidaecin multipeptide precursor structure: a putative mechanism for amplification of the insect antibacterial response. *The EMBO journal* 12:1569–1578.

Chakraborty M, Chang C-H, Khost DE, Vedanayagam J, Adrion JR, Liao Y, Montooth KL, Meiklejohn CD, Larracuente AM, Emerson JJ. 2021. Evolution of genome structure in the Drosophila simulans species complex. *Genome Res* 31:380–396.

Chapman JR, Hill T, Unckless RL. 2019. Balancing selection drives maintenance of genetic variation in Drosophila antimicrobial peptides. *Genome Biology and Evolution* 11:2691–2701.

Clemmons AW, Lindsay SA, Wasserman SA. 2015. An Effector Peptide Family Required for Drosophila Toll-Mediated Immunity. *PLoS Pathogens* 11.

Davie K, Janssens J, Koldere D, De Waegeneer M, Pech U, Kreft Ł, Aibar S, Makhzami S, Christiaens V, Bravo González-Blas C, et al. 2018. A Single-Cell Transcriptome Atlas of the Aging Drosophila Brain. *Cell* 174:982-998.e20.

De Gregorio E, Spellman PT, Tzou P, Rubin GM, Lemaitre B. 2002. The Toll and Imd pathways are the major regulators of the immune response in Drosophila. *EMBO Journal* 21:2568–2579.

De Lorenzi E, Chiari M, Colombo R, Cretich M, Sola L, Vanna R, Gagni P, Bisceglia F, Morasso C, Lin JS, et al. 2017. Evidence that the human innate immune peptide LL-37 may be a binding partner of amyloid-β and inhibitor of fibril assembly. *Journal of Alzheimer's Disease* 59:1213–1226.

Delport W, Poon AFY, Frost SDW, Kosakovsky Pond SL. 2010. Datamonkey 2010: A suite of phylogenetic analysis tools for evolutionary biology. *Bioinformatics* 26:2455–2457.

Dominy SS, Lynch C, Ermini F, Benedyk M, Marczyk A, Konradi A, Nguyen M, Haditsch U, Raha D, Griffin C, et al. 2019. Porphyromonas gingivalis in

Alzheimer's disease brains: Evidence for disease causation and treatment
with small-molecule inhibitors. *Science Advances* 5.

Ebrahim SAM, Talross GJS, Carlson JR. 2021. Sight of parasitoid wasps accelerates
sexual behavior and upregulates a micropeptide gene in Drosophila. *Nat*
Commun 12:2453.

Ferreira ÁG, Naylor H, Esteves SS, Pais IS, Martins NE, Teixeira L. 2014. The Toll-
dorsal pathway is required for resistance to viral oral infection in Drosophila.
PLoS Pathog. 10:e1004507.

Gerdol M, Schmitt P, Venier P, Rocha G, Rosa RD, Destoumieux-Garzón D. 2020.
Functional Insights From the Evolutionary Diversification of Big Defensins.
Front Immunol 11:758.

Halldórsdóttir K, Árnason E. 2015. Trans-species polymorphism at antimicrobial
innate immunity cathelicidin genes of Atlantic cod and related species. *PeerJ*
3:e976.

Hammonds AS, Bristow CA, Fisher WW, Weiszmann R, Wu S, Hartenstein V, Kellis M,
Yu B, Frise E, Celniker SE. 2013. Spatial expression of transcription factors in
Drosophila embryonic organ development. *Genome Biol.* 14:R140.

Hanson MA, Cohen LB, Marra A, Iatsenko I, Wasserman SA, Lemaitre B. 2021. The
Drosophila Baramicin polypeptide gene protects against fungal infection.
PLoS Pathog 17:e1009846.

Hanson MA, Dostálová A, Ceroni C, Poidevin M, Kondo S, Lemaitre B. 2019. Synergy
and remarkable specificity of antimicrobial peptides in vivo using a
systematic knockout approach. *eLife* 8.

Hanson MA, Hamilton PT, Perlman SJ. 2016. Immune genes and divergent
antimicrobial peptides in flies of the subgenus Drosophila. *BMC evolutionary*
biology 16:228.

Hanson MA, Lemaitre B. 2020. New insights on Drosophila antimicrobial peptide
function in host defense and beyond. *Curr. Opin. Immunol.* 62:22–30.

Hanson MA, Lemaitre B, Unckless RL. 2019. Dynamic Evolution of Antimicrobial
Peptides Underscores Trade-Offs Between Immunity and Ecological Fitness.
Frontiers in Immunology 10:2620.

Hellgren O, Sheldon BC, Buckling A. 2010. In vitro tests of natural allelic variation of
innate immune genes (avian ??-defensins) reveal functional differences in
microbial inhibition. *Journal of Evolutionary Biology* 23:2726–2730.

Huang J, Lou Y, Liu J, Bulet P, Jiao R, Hoffmann JA, Liegeois S, Li Z, Ferrandon D. 2020.
The BaramicinA gene is required at several steps of the host defense against
Enterococcus faecalis and Metarhizium robertsii in a septic wound infection
model in Drosophila melanogaster. bioRxiv Available from:
http://biorxiv.org/lookup/doi/10.1101/2020.11.23.394809

Jiggins FM, Kim KW. 2007. A screen for immunity genes evolving under positive
selection in Drosophila. *Journal of Evolutionary Biology* 20:965–970.

Jumper J, Evans R, Pritzel A, Green T, Figurnov M, Ronneberger O, Tunyasuvunakool
K, Bates R, Žídek A, Potapenko A, et al. 2021. Highly accurate protein
structure prediction with AlphaFold. *Nature*.

Kearse M, Moir R, Wilson A, Stones-Havas S, Cheung M, Sturrock S, Buxton S, Cooper
A, Markowitz S, Duran C, et al. 2012. Geneious Basic: An integrated and
extendable desktop software platform for the organization and analysis of
sequence data. *Bioinformatics*.

Kelley LA, Mezulis S, Yates CM, Wass MN, Sternberg MJE. 2015. The Phyre2 web
portal for protein modeling, prediction and analysis. *Nature protocols*
10:845–858.

Kim BY, Wang J, Miller DE, Barmina O, Delaney EK, Thompson A, Comeault AA,
Peede D, D'Agostino ERR, Pelaez J, et al. 2021. Highly contiguous assemblies
of 101 drosophilid genomes. *eLife* 10:e66405.

Kobler JM, Rodriguez Jimenez FJ, Petcu I, Grunwald Kadow IC. 2020. Immune
Receptor Signaling and the Mushroom Body Mediate Post-ingestion Pathogen
Avoidance. *Curr Biol* 30:4693-4709.e3.

Krogh A, Larsson B, von Heijne G, Sonnhammer EL. 2001. Predicting transmembrane
protein topology with a hidden Markov model: application to complete
genomes. *J Mol Biol* 305:567–580.

Leader DP, Krause SA, Pandit A, Davies SA, Dow JAT. 2018. FlyAtlas 2: A new version
of the Drosophila melanogaster expression atlas with RNA-Seq, miRNA-Seq
and sex-specific data. *Nucleic Acids Research* 46:D809–D815.

Lee M, Shi X, Barron AE, McGeer E, McGeer PL. 2015. Human antimicrobial peptide
LL-37 induces glial-mediated neuroinflammation. *Biochemical Pharmacology*
94:130–141.

Lemaitre B, Reichhart JM, Hoffmann JA. 1997. Drosophila host defense: differential
induction of antimicrobial peptide genes after infection by various classes of
microorganisms. *Proceedings of the National Academy of Sciences of the*
United States of America 94:14614–14619.

Lezi E, Zhou T, Koh S, Chuang M, Sharma R, Pujol N, Chisholm AD, Eroglu C,
Matsunami H, Yan D. 2018. An Antimicrobial Peptide and Its Neuronal
Receptor Regulate Dendrite Degeneration in Aging and Infection. *Neuron*
97:125-138.e5.

Mackay TFC, Richards S, Stone EA, Barbadilla A, Ayroles JF, Zhu D, Casillas S, Han Y,
Magwire MM, Cridland JM, et al. 2012. The Drosophila melanogaster Genetic
Reference Panel. *Nature* 482:173–178.

Moir RD, Lathe R, Tanzi RE. 2018. The antimicrobial protection hypothesis of
Alzheimer's disease. *Alzheimer's & Dementia* 14:1602–1614.

Murrell B, Weaver S, Smith MD, Wertheim JO, Murrell S, Aylward A, Eren K, Pollner
T, Martin DP, Smith DM, et al. 2015. Gene-wide identification of episodic
selection. *Mol Biol Evol* 32:1365–1371.

Neely GG, Hess A, Costigan M, Keene AC, Goulas S, Langeslag M, Griffin RS, Belfer I,
Dai F, Smith SB, et al. 2010. A Genome-wide Drosophila screen for heat
nociception identifies α2δ3 as an evolutionarily conserved pain gene. *Cell*
143:628–638.

Petersen AJ, Katzenberger RJ, Wassarman DA. 2013. The innate immune response
transcription factor relish is necessary for neurodegeneration in a Drosophila
model of ataxia-telangiectasia. *Genetics* 194:133–142.

Pfaffl MW. 2001. A new mathematical model for relative quantification in real-time
RT-PCR. *Nucleic Acids Res.* 29:e45.

Quesada H, Ramos-Onsins SE, Aguad?? M. 2005. Birth-and-death evolution of the
Cecropin multigene family in Drosophila. *Journal of Molecular Evolution* 60:1–
11.

Rolff J, Schmid-Hempel P. 2016. Perspectives on the evolutionary ecology of
arthropod antimicrobial peptides. *Philosophical Transactions of the Royal*
Society B: Biological Sciences 371.

Sackton TB, Lazzaro BP, Clark AG, Wittkopp P. 2017. Rapid expansion of immune-
related gene families in the house fly, musca domestica. *Molecular Biology*
and Evolution.

Sinner MP, Masurat F, Ewbank JJ, Pujol N, Bringmann H. 2021. Innate Immunity
Promotes Sleep through Epidermal Antimicrobial Peptides. *Current Biology*
31:564-577.e12.

Tamura K, Subramanian S, Kumar S. 2004. Temporal Patterns of Fruit Fly
(Drosophila) Evolution Revealed by Mutation Clocks. *Molecular Biology and*
Evolution 21:36–44.

Tennessen JA. 2005. Molecular evolution of animal antimicrobial peptides:
943   Widespread moderate positive selection.

944   Toda H, Williams JA, Gulledge M. 2019. A sleep-inducing gene, nemuri , links sleep
945   and immune function in Drosophila. 515:509–515.

946   Train C-M, Pignatelli M, Altenhoff A, Dessimoz C. 2019. iHam and pyHam: visualizing
947   and processing hierarchical orthologous groups. *Bioinformatics (Oxford,*
948   *England)* 35:2504–2506.

949   Unckless RL, Howick VM, Lazzaro BP. 2016. Convergent Balancing Selection on an
950   Antimicrobial Peptide in Drosophila. *Current Biology* 26:257–262.

951   Vilcinskas A, Mukherjee K, Vogel H. 2013. Expansion of the antimicrobial peptide
952   repertoire in the invasive ladybird Harmonia axyridis. *Proceedings of the*
953   *Royal Society B: Biological Sciences*.

954   Wang Y, Zhu S. 2011. The defensin gene family expansion in the tick Ixodes
955   scapularis. *Developmental and Comparative Immunology*.

956   Zanetti M. 2005. The role of cathelicidins in the innate host defenses of mammals.
957   *Curr Issues Mol Biol* 7:179–196.

958   Zhang W, Yang J, He B, Walker SE, Zhang H, Govindarajoo B, Virtanen J, Xue Z, Shen
959   H-B, Zhang Y. 2016. Integration of QUARK and I-TASSER for Ab Initio Protein
960   Structure Prediction in CASP11. *Proteins* 84 Suppl 1:76–86.