Natural Selection as the Sum over all Histories and a Differential Equation for Environmental Coevolution.

C E Neal-Sturgess
Emeritus Professor of Mechanical Engineering
University of Birmingham

Abstract: As evolution can be connected to the principle of least action, and if it is depicted in evolution-space versus time then it corresponds to the direction of ultimate causation. As an organism evolves and follows a path of proximate causation, if the vector is closely parallel to that of the Ultimate Causation then the changes will confer desirable attributes which will lead to further development. If, however, the variations do not occur in a direction close to that of the ultimate causation vector the evolved organism will quickly die out. This may be viewed as similar to Feynman's sum over all histories. Therefore, the principle of least action gives a direction, but not a purpose, to evolution. Taking the coevolution model of Lewontin, an equation of motion for environmental coevolution shows that it is the rate of evolutionary change that responds to changes in the environment, which is in line with some evidence from population studies.

Background:

Since the publication of "On the Origin of Species by Natural Selection" by Charles Darwin in 1859, there have been a number of attempts to link it to other scientific principles, notably the Principle of stationary Action, known popularly as "Least Action" [Nahin 2004]. In their paper Natural selection for least action (Kaila and Annila 2008) they depict evolution as a process conforming to the Principle of Least Action (PLA). This paper, although not giving any experimental evidence, shows that evolution, if conforming to the Second Law of Thermodynamics, will follow a trajectory of maximum Entropy production, which conforms to the PLA. To demonstrate this they rewrote the Gibbs-Duhem relationship in terms of possible states, to give a differential equation of evolution. This is a convincing argument as biology at root is a physical process, as expounded by Schrodinger 1944, and all physical processes are governed by the Second Law of thermodynamics.

The PLA first arose, in modern times in the 17[th] century, and was popularised by Maupertius although the major rigorous work was done by Euler [Kaila & Annila 2008]. was Feynman's guiding principle throughout this life [Feynman 1985]. As Wittgenstein 1922 said "people were using the Principle of Least Action before they knew it existed".

The Principle of Least (Stationary) Action is stated simply as:

$$S = \int_0^t U.dt \ \ and \ \ \frac{d}{dt}[S]=0 \tag{1}$$

Where U is the Gibb's Free Energy.

The second law can be stated as:

$$F = E - E_{nt} \tag{2}$$

Where: E is the total energy of the body, F is the "Gibbs Free Energy" and E_{nt} is the entropy.

Results:

Here it is hypothesised that an analogue exists to Feynman's "Sum Over All Histories" [Feynman 1985] . Richard Feynman, Nobel Laureate, was fascinated by the PLA for most of his life, and it led to one of his greatest breakthroughs in Quantum Electrodynamics or QED – the quantum theory of light [Feynman 1985]. Feynman considered that a Photon traversing a path from A to B in space-time was free to traverse any path. However, when the sum of all paths is taken, because of phase differences, the most likely path is that conforming to the PLA; this is known as the "Sum Over All Histories".

It is assumed here that there is a general overall direction of evolution. The vector of evolution is called here the Ultimate Causal Direction (green vector), in accordance with the terminology used in evolutionary biology(Mayr 2001), and this should conform to the PLA [Kaila & Annila 2008], in conformity with the external conditions, and could be shown graphically as shown in Fig.1.

Fig.1 Evolution-Space vs Time.

If during any genetic mutation leading to a change in the organism takes place (event A) this will in general be random, under constraints where the genome stays constant but the phenotype changes [Holmes 2013] and confers on the next generation of the organism's attributes which may or may not lead to a greater probability of survival in the external climate [Loewe 2010]. These changes to the phenotype are in evolutionary biological terms called "Proximate Causation". It is proposed that Proximate Causation will lead to greater

survivability if it is parallel, or closely parallel, to the direction of PLA, for the external conditions at the time (fitness). This is shown in Fig.1. as the red vector A-B-C. In this diagram, those variants not reasonably parallel to the trajectory of ultimate causality, the black lines ending in blobs, will die out. Further Epigenetic changes [Dupont 2006] could also more closely align the red vector with the Ultimate Causality. Although this diagram as an axis of Time, the interval between the changes is not known, as shown by the caption. The eventual unfolding of Proximate Causation towards Ultimate Causation as determined by both thermodynamics and the environment can be considered as analogous to Feynman's "Sum over all Histories", as the mutations leading to Proximate causation may be viewed as exploring all the possible states in evolution space, but only those approximately parallel to the environment vector will survive to generate new variants and lead to the Ultimate Causality; the other variants will die out [Loewe 2010]. Hence evolution towards the Ultimate Causality can be seen as the ultimate goal of the sum over all proximate evolutionary histories and corresponds to Natural Selection.

This is not to say the evolution to the PLA is teleological, there is no "purpose". It is simply that those evolutionary lines arising from proximate causality that are closely parallel to the line of PLA, are best adapted to whatever external environments that exist at the time, and so survive longer, and have a greater probability of reproduction (heritable); therefore, evolution has a direction but not a purpose.

There are also interesting developments in evolutionary thinking where some are beginning to posit that organisms make changes to their phenotype, as opposed to their genotype, by adjusting to their environment (Holmes 2010). This tendency here would be to add a rotation to the Proximate Causality, hence bringing the vector closer to the direction of the Ultimate Causality, in the direction of PLA.

Most investigators consider that natural selection is the cumulative effects of the genetic mutations, which is the position adopted here. Other investigators however classify natural selection as one of the mechanisms of mutation [Sanjuan & Domingo-Calap 2016], which is not adopted here.

It was difficult to reconcile the original Gaia hypothesis due to haemostasis [Lovelock 1979], however later variants of Gaia called "Coevolution" [Lewontin 1983] allows reconciliation with Gaia. Lewontin's description of Coevolution as two, coupled, differential equations to express the interaction between the organism (O) and the environment (E) as:

$$\frac{d}{dt}(O) = f(O, E) \tag{3}$$

and $\quad \frac{d}{dt}(E) = g(O, E) \tag{4}$

is an interesting development. It should be noted that in the literature coevolution can either be species coevolution and environmental evolution, this paper is solely concerned with environmental coevolution. Levin and Lewontin 1985 stated that solving these differential equations would be very difficult.

Taking the idea that the second law can be rewritten as an equation of motion [Kaila and Annila 2008[, and if Lagrangian's can be defined as shown below:

$$\mathcal{L}_O = f\{f(O, E)\} \tag{5}$$

$$\mathcal{L}_E = f\{g(O, E)\} \tag{6}$$

Then using Euler-Lagrange [Goldstein 1969] gives:

$$\frac{\partial}{\partial t} \mathcal{L}_O f\{f(O, E)\} - \mathcal{L}_E f\{g(O, E)\} = 0 \tag{7}$$

Hence:

$$\frac{\partial}{\partial t} \mathcal{L}_O f\{f(O, E)\} = \mathcal{L}_E f\{g(O, E)\} \tag{8}$$

The Lagangians will be difficult to define, but it is possible that they will contain a potential term proportional to the environment, and a kinetic term proportional to the organisms muatation rate.Therefore this equation gives an partial differential equation for environmental coevolution and shows that the rate of organism change, the rate of viral mutation or evolution rate , (the left hand side) is proportional to the magnitude of the change occurring in the environment, the right hand side, which is in agreement with some early evidence [Husby, Gross, Gianepp, Franks] dealt with below.

However, there is a measurement problem in evolution through the fossil record in that the rate of progress is inversely proportional to the timescale over which it is measured, leading to Gould's "Punctuated Equilibrium" ideas (Eldridge & Gould 1972, Gould 2002). Furthermore, another measurement problem is that of actual findings and observation. In general, the probability of finding a fossil will be proportional to the number of the species that existed, and so the time the fossil type lived. Therefore, the probability of finding a fossil type is dependent on how successful that fossil type was, and so most of the changes that have occurred in a species will probably never be found because they died out quickly. This is an exceptionally complicated problem, and as said by [Loewe and Hill] that it is "like looking for a needle in a haystack'; however, advances in Evolutionary Genetics may give answers in the future.

To look for empirical evidence of equation 8 the field can be broadly classified as 1) viral mutations, 2) bacterial mutations and 3) population level studies (considered by Levin and Lewontin as possibly the only level at which the equations may be solved).

Considering viral mutation rates first, there is a considerable number of investigations reported in the literature [Sanjuan et.al. 2010. Regoes et.al 2012, Peck et.al. 2015, Sanjua & Domingo-Calap 2015, Martınez-Padilla et.al. 2017. Peck & Lauring 2018, Duffy 2018]. However, a problem exists when trying to compare the reported "viral mutation rates" with the analysis conducted here. Equation 8 is a "rate of change" with respect to time, whereas typically the viral mutation rates are reported as substitution's/nucleotide/cell (s/n/c), which should not be confused with rates of evolutionary change [Peck & lauring 2018]. Although there is some evidence of a relationship between the two [Peck & Lauring 2018] see Fig3, it is likely that any such relationship will probably be a function of virus type and the mutation mechanism. Pleck et.al. 2015 state that Neutral theory posits that the evolutionary rate increases linearly with the mutation rate, which is interesting in that equation 8 could, with appropriate constants. apply to both viral rates and evolutionary rate. Although there is clearly a relationship between evolutionary rate and mutation rate there does not appear to be any systematic investigations of viral mutation rates or evolutionary rate and the environment. However, there seems to be a general conclusion that viral mutation rates do respond to selective environmental pressures from the environment, which could be in

agreement with the differential equation derived here. It is too early to comment on Covid-19, but it will be interesting to sees if the mutation frequency increases as the environment gets more difficult for the virus due yo the vaccine rollout.

Fig. 2. (A) Log-scale mean evolutionary rates against mutation rates for each Baltimore Class (B) Evolutionary rates against mutation rates for individual viruses. For both (A) and(B), the solid line represents the deleterious mutation model prediction, while the dashed line indicates the prediction from the within-host analytical model.
After Peck et.al 2015.

Dealing next with bacterial mutation rates, there is a considerable literature [Rosche et.al 2000, Zeyl & Devisser 2000, Denamur & Matic 2006, Pal et.al. 2007 Wielgossa et.al 2013, Chevallereau et.al 2018, Ferenci 2019, Ramiro RS et.al 2020] however the same type of problem in comparing equation 8 with the evidence exists as does that of viral mutation rates. One reference [Rosche et.al] discusses the problem with determining mutation rates in bacteria, and maintain that these methods are more accurate than time rate of change studies due to need to invoke theoretical models in the latter; this should not preclude such studies. Rosche et.al 2000 survey six different methods of calculation of mutation rates and conclude that the maximum likelihood rate is preferred, but more work is necessary. Ferenci 2019 from a review concludes that recent findings suggest a very uneven relationship between stress and mutations, and it remains to be investigated whether stress specific genetic variation impacts on evolvability differentially in distinct environments. Therefore, although again there do not appear to be any systematic investigations of bacterial mutation rates and environment, there seems to be a general conclusion that bacterial mutation rates do respond to selective pressures from the environment, which could again be in agreement with the differential equation derived here.

Research into evolution is extensive and has changed over time from qualitative to quantative measures, in some qualtative investigations it was difficult to separate phenotype plasticity from genetic evolvabilty. Following the landmark paper by Houle (1992) where he defined two quantative measures of genetic evolvablity namely the coefficient of evolvability V_A and its square I_A, with the comment that he considered I_A to be the most accurate; the field became mor quantative. Exploring genetic evolvability is complex and in the subsequent years there were many investigations which contained errors smmarised by Garcia-Gonzalez et.al. 2012 who reviewed 364 papers and found that there were errors in 47%.

In a major investgation of a 35 yerars longitudinal Husby (2011) showed that, in a wild population of great tits (Parus major), the strength of the directional selection gradients on timing of breeding increased with increasing spring temperatures, and that genotype-by-environment interactions also predicted an increase in additive genetic variance, and heritability, of timing of breeding with increasing spring temperature. There was a significant positive association between the annual selection differentials and the corresponding heritability.

Marrot (2013) Posited that evolutionary adaptation, as a response to climate change, is expected for fitness related traits affected by climate and exhibiting genetic variance. Their results indicated an increase in the strength of selection by 46% for every +1°C anomaly in warmer daily maximum temperatures in April. Such climate-driven influences on the strength of directional selection acting on laying date could favour an adaptive response in this trait, since it is heritable.

Franks (2013) Observed that as climate change progresses, there are widespread changes in phenotypes in many plant populations. Whether these phenotypic changes are directly caused by climate change, and whether they result from phenotypic plasticity or evolution, are active areas of investigation. Of the 38 studies that met their criteria for inclusion, all found plastic or evolutionary responses, with 26 studies showing both.

Wadgyner et.al (2019) conducted a major review (192 references) of the effects of climate change on evolution and concluded that climate change has the potential to directly and indirectly influence the direction and rate of evolution in natural populations. They also considered coevolution with both species and the environment.

In another major investgation Martınez-Padilla et.al (2017) they conducted a major bibliogaraphical review and found a final sample covering 20 populations of 12 species and asked the critical question: "do populations have the ability to evolve in response to these change?"; however, knowledge on how evolution works in wild conditions under different environmental circumstances is extremely limited. They used published data to collect or calculate 135 estimates of evolvability of morphological traits of European wild bird populations. The authors used a quantitative approach based on the two dimensionless ratios namely the coefficient of additive genetic variation (CV_A), and its square (I_A), as indexes of evolvability [Houle 1992]. If the time rate of change of evolution is proportianl to the environmental change in accordance with equation 8 the comparison with recent evidence of evolutionary change can be facilitated by rewriting equation 8 as follows: if a high rate of the coeficient of additive evolution (CV_A) is due to an increase in the the rate of change of mutations, then it can be inferred that:

$$\frac{\partial (CV_A)}{\partial E_F} \ or \ \frac{\partial (I_A)}{\partial E_F} \propto \mathcal{L}_E f\{g(O,E)\} \tag{9}$$

Where E_F is the environmental favourability, and the dimensionless coefiicents defined after Houle 1992 as:

$$CV_A = 100\frac{\sqrt{V_A}}{\bar{X}} \text{ and } I_A = CV_A{}^2 \qquad\qquad (10)$$

Where V_A ie the additive variation and $\bar{X}$ the population mean.
The environmental favourability was determined using Logistic Regression methods, after Real et.al 2006, to obtain Favourability functions.
Equation 9 makes it simpler to compare with the evidence presented below. The results see Fig 2 clearly show that the environmental favourability increases (fitness) the rate of evolvability (the slope of the curves) decreases; this is in accord with the differential equation of coevolution (equation 8).

Fig 2: Evolvability criteria versus environmental favourability after Martenaz et.al.

Figure 2a shows that as the environmental favourabilty approaches 1.0 the rate of evolvabilty deceases to approximately zero, with a reasonably large but unambiguous range. In Figure 2b the results for I_A are more definite and show that as the envirommental favourability approaches 1.0 the rate of evolvabity actually becomes negative, which is understandable for if the fitness to enviromental favourability is perfect why should the organism mutate? This is an important result as it as it shows that it is not evolution per.se which is driven by changes in the environment, but it is the rate of evolution (evlovability)which changes, which is in agrement with equation 8.
Curve fitting to the results above gives Fig 3a & 3b, from these figures it is obvious that the means are fitted very closely by quadratic functions as shown.

Fig.3 Coefficients against Environmental Favourability

Taking the analysis to the next step the rates of change for CA$_V$ and I$_A$ are:

$$\frac{\partial CA_V}{\partial E_F} = 0.02711 - 0.0325E_F \tag{11}$$

$$\frac{\partial I_A}{\partial E_F} = 0.001563 - 0.00219E_F \tag{12}$$

These equations, of the generic form A – BE$_F$, clearly show that the slope of the CA$_V$ and I$_A$ consistently reduce as the environmental favourability increases, and equations 10 and 11 are linear functions of the environmental favourability. This could be important in helping to untangle the importance of the environmental changes and assist the definition of the Lagrangian in equation 9. In the face of widely held views on mass extinctions because of climate change this gives some small hope [Husby]. However, this is potentially bad new for Covid-19 as. although the data is not yet available, as the stress on the environment for the virus increases due to the vaccination programme , then the mutation rate will increase and we are in a continual catch-up as with the flu virus.

Conclusions:
Evolution can be connected to the Principle of Least Action, and if it is depicted in "Evolution-Space vs Time" then it corresponds to the direction of Ultimate Causation. As an organism evolves and follows a path of Proximate Causation, if the vector is closely parallel to that of the Ultimate Causation then the changes will confer desirable heritable attributes which will lead to further development. If, however the variations do not occur in a direction close to the of the Ultimate Causation vector the evolved organism will quickly die out. This may be viewed as similar to Feynman's "sum over all histories. Therefore, the Principle of Least Action gives a direction, but not a purpose, to Evolution.
Taking the environmental Coevolution model of Lewontin, a differential equation for Coevolution shows that it is the rate of evolutionary change (evolvability) that responds to

changes in the environment, which agrees with a number of studies. Further analysis shows that the rate of change of CAv and IA with environmental favourability are linear, which could help in defining the Lagrangian of the environmental effects. However, this is potentially bad news for Covid-19 as. although the data is not yet available, as the stress on the environment for the virus increases due to the vaccination programme, then the mutation rate will increase and we are in a continual catch-up as with the flu virus.

References:

Bendall, D.S., ed.), from Molecules to Men (pp. 273–285, Cambridge

Chevallereau A, Meaden s, Edze S, Westra R, Rollie C. The effect of bacterial mutation rate on the evolution of CRISPR-Cas adaptive immunity, Phil. Trans. R. Soc. B 374: 20180094. http://dx.doi.org/10.1098/rstb.2018.0094

Darwin C, On the Origin of Species by Natural Selection, John Murray, London, 1859.

Denamur E, Matic I. Evolution of mutation rates in bacteria. Molecular Microbiology (2006) 60(4), 820–827

Duffy S (2018) Why are RNA virus mutation rates so damn high? PLoS Biol 16(8): e3000003. https://doi.org/10.1371/journal. pbio.3000003

Dupont C, Armant DR, Brenner CA (September 2009). "Epigenetics: definition, mechanisms and clinical perspective". Seminars in Reproductive Medicine. **27** (5): 351–7.

Ferenci T, Irregularities in genetic variation and mutation rates with environmental stresses. Environmental Microbiology (2019) 21(11), 3979–3988 doi:10.1111/1462-2920.14822

Feynman, R. P. (1985). QED: The Strange Theory of Light and Matter, Penguin.

Free A, and Barton NH, Do evolution and ecology need the Gaia hypothesis? TRENDS in Ecology and Evolution Vol.22 No.11

Franks S J, Weber J J and Aitken S N: Evolutionary and plastic responses to climate change in terrestrial plant populations Evolutionary Applications ISSN 1752-4571

Garcia-Gonzalez F, Simmons LW, Tomkins JL, Kotiaho JS and Evans JP. Comparing evolvabilities: common Errors surrounding the calculation And use of coefficients of additive Genetic variation. Evolution August 2012.

Gienapp P, Veronika N. Laine, A. Mateman C, van Oers K and Visser M E. Environment-Dependent Genotype-Phenotype Associations in Avian Breeding Time

Gould SJ & Eldridge N, Punctuated equilibria: an alternative to phyletic gradualism (1972) pp 82-115 in "Models in paleobiology", edited by Schopf, TJM Freeman, Cooper & Co, San Francisco.

Gould SJ, 2002 The Structure of Evolutionary Theory. Harvard University Press.

Goldstein H. Classical Mechanics. Wiley 1969.

Gross L (2011) Climate Change Could Change Rates of Evolution. PLoS Biol 9(2): e1001015. doi:10.1371/journal.pbio.1001015

Holmes , Life's Purpose, New Scientist, p33, 12 October 2013.

Houle D (1992) Comparing evolvability and variability of quantitative traits. Genetics 130: 195–204 1992,

Husby A, Visser ME, Kruuk LEB (2011) Speeding up Microevolution: The Effects of increasing Temperature on Selection and Genetic Variance in a Wild Bird Population. doi:10.1371/journal.pbio.1000585

Kaila, V. R. I. and A. Annila (2008). "Natural selection for least action." Proc.R. Soc A(464): 3055-3070.

Levin R, Lewontin ~R. The dialectical Biologist. 1985. Univeristy of Harvard.

Lewontin, R.C. (1983) Gene, organism and environment. In Evolution.

Loewe L and Hill WG, The population genetics of mutations: good, bad and indifferent; Phil. Trans. R. Soc. B (2010) 365, 1153–1167 doi:10.1098/rstb.2009.0317

Lovelock J, Gaia, 1979, OUP.

Martı´nez-Padilla J, Estrada A, Early R, Garcia-Gonzalez F. 2017 Evolvability meets biogeography: evolutionary potential decreases at high and low environmental favourability. Proc. R. Soc. B 284: 20170516. http://dx.doi.org/10.1098/rspb.2017.0516

Marrot P, Charmantier A , Blondel J : Garant D: Current spring warming as a driver of selection on reproductive timing in a wild Passerine: Journal of Animal Ecology, DOI: 10.1111/1365-2656.12794

Mayr, E. (2001). What Evolution Is, Basic Books.

Nahin, PJ, When Least is Best, Princeton UP. 2004.

Pal C, Macia MD, Oliver A, Schachar I, Buckling. A Coevolution with viruses drives the evolu,tion of bacteria mutation rates. NATURE| Vol 450| 13 December 2007.

Peck KM, Chan CHS. Tanaka MM. Connecting within-host dynamics to the rate of viral molecular evolution. Virus Evolution, 2015, 1(1): vev013.

Peck KM, Lauring AS. 2018. Complexities of viral mutation rates. J Virol 92:e01031-17. https://doi.org/10.1128/JVI .01031-17

Ramiro RS, Durão P, Bank C, Gordo I (2020) Low mutational load and high mutation rate variation in gut commensal bacteria. PLoS Biol 18 (3): e3000617. https://doi.org/10.1371/journal. pbio.3000617

Real R, Barbosa AM, Vargas JM. 2006 Obtaining environmental favourability functions from logistic regression. Environ. Ecol. Stat. 13, 237– 245.(doi:10.1007/s10651-005-0003-3).

Regoes RR, Hamblin S, Tanaka MM. 201 Viral mutation rates: modelling the roles of within-host viral dynamics and the trade-off between replication fidelity and speed. Proc R Soc B 280: 20122047. http://dx.doi.org/10.1098/rspb.2012.2047

Rosche WA and Foster PL, Determining Mutation Rates in Bacterial Populations, Methods. 2000 January ; 20(1): 4–17. doi:10.1006/meth.1999.0901.

Sanjua´ R, Domingo-Calap P. Mechanisms of viral mutation. Cell. Mol. Life Sci. (2016) 73:4433–4448 DOI 10.1007/s00018-016-2299-6.

Schrodinger, E. (1944). What is life?, Cambridge University Press 2007

Wittgenstein L, Tractatus Logicus-Philosophicus, Routledge 1922.

Wadgymar S M, MacTavish R, Anderson J T: Evolutionary consequences :of climate change: Ecosystem Consequences of Soil Warming , 2019 Elsevier Inc. https://doi.org/10.1016/B978-0-12-813493-1.00003-X

Wielgossa S, Barrickc JE, Tenaillone O, Wiserd, MJ, W. Dittmarh J, Cruveilleri S, Chane-Woon-Mingi B, Médiguei C, Lenskid RE,and Schneidera D, Mutation rate dynamics in a bacterial population reflect tension between adaptation and genetic load. PNAS | January 2, 2013 | vol. 110 |

Zeyl C, DeVisser JAGM. Estimates of the Rate and Distribution of Fitness Effects of Spontaneous Mutation in Saccharomyces cerevisiae. The Genetics Society of America, 2000.