Main Manuscript for

Elucidation of fibril helix structure responsible for swimming in
Spiroplasma using electron microscopy

Yuya Sasajima[a], Takayuki Kato[b], Tomoko Miyata[c], Akihiro Kawamoto[b], Keiichi Namba[c,d,e] and
Makoto Miyata[a,f]

[a]Graduate School of Science, Osaka City University, Osaka, 558-8585, Japan, [b]Institute for
Protein Research, Osaka University, 3-2 Yamadaoka, Suita, Osaka, 565-0871, Japan, [c]Graduate
School of Frontier Biosciences, Osaka University, 1-3 Yamadaoka, Suita, Osaka, 565-0871,
Japan, [d]RIKEN Center for Biosystems Dynamics Research and SPring-8 Center, 1-3
Yamadaoka, Suita, Osaka, 565-0871, Japan, [e]JEOL YOKOGUSHI Research Alliance
Laboratories, Osaka University, 1-3 Yamadaoka, Suita, Osaka, 565-0871, Japan, [f]The OCU
Advanced Research Institute for Natural Science and Technology (OCARINA), Osaka City
University, Osaka, 558-8585, Japan.

*Makoto Miyata

Email: miyata@osaka-cu.ac.jp

Author Contributions: YS and MM designed the research. YS performed the research. TK, TM,
AK, and KN provided techniques for structural analyses. YS, KN, and MM wrote the paper. All
authors discussed and checked the entire manuscript.

Competing Interest Statement: The authors have no competing interests to declare.

Classification: Major class: Biological Science. Minor class: Microbiology; Biophysics and
Computational Biology.

Keywords: Single particle analysis, Helical shape, Motility, Cytoskeleton, Filament.

This PDF file includes:

Main Text
Figures 1 to 6
Tables 1 to 2

Abstract

Spiroplasma, known pathogens and commensals of arthropods and plants, are helical-shaped
bacteria lacking the peptidoglycan layer. They swim by alternating between left- and right-handed
cell helicity, which is driven by an internal structure called the ribbon. This system is unrelated to
flagellar motility that is widespread in bacteria. The ribbon comprises the bacterial actin homolog
MreB and fibril, the protein specific to *Spiroplasma*. Here, we isolated the ribbon and its core, the
fibril filament, and using electron microscopy, found that the helicity of the ribbon and the cell is
linked to the helicity of the fibril. Single particle analysis using the negative-staining method
revealed that the three-dimensional structures of the fibril filament comprise a repeated ring
structure twisting along the filament axis. The handedness of these structures were verified by the

structures reconstituted through tomogram from quick-freeze deep-etch replica electron
microscopy. Based on these observations, we propose a scheme for the helicity-switching
mechanism in which the twists caused by the conformational changes in the fibril filament are
accumulated, transmitted to the ribbon, and then propel the cells by rotating the cell body like a
screw.

Significance Statement

Spiroplasma are commensal and pathogenic bacteria widespread globally in animals and plants.
A few species are also recognized as male-killing bacteria of insects. Their special swimming
mechanism is caused by helicity switching, which could be the simplest swimming mechanism,
because helicity shift can be caused by contraction and extension of paired filaments. This
mechanism has attracted research attention for many years because of the possible application
in the field of nano actuators; however, the details of this mechanism remain to be clarified. Here,
we propose a model for the swimming mechanism based on the structure of the core of the
Spiroplasma ribbon.
54
55 **Main Text**
56
57 **Introduction**
58
Mollicutes, which are parasitic or commensal bacteria, have evolved from the phylum Firmicutes
that includes *Bacillus* and *Clostridium* by reducing their genome sizes (1-4). In the course of
evolution, the cells have become softer and smaller owing to the loss of the peptidoglycan layer.
These changes allowed some species to transmit their internal housekeeping activities such as
the rotation of ATP synthesis to the outside, resulting in the acquisition of three unique motility
mechanisms (2, 5, 6). Two of the three, represented by *Mycoplasma mobile* and *Mycoplasma*
pneumoniae, exhibit gliding motilities on solid surfaces in which leg structures repeatedly catch
sialylated oligosaccharides on host cells, based on two different mechanisms (6-8). The other
motility system is the helicity-switching swimming of *Spiroplasma*, the subject of the present study
(Movie_S1) (7, 9-11). *Spiroplasma* species are parasitic to plants and arthropods, and are
characterized as polarized helical-shaped cells with a tapered end (12, 13). They also show
obvious chemotaxis despite the absence of genes for the two-component regulatory system in
the genome, which is generally responsible for bacterial chemotaxis (14). In general, swimming
bacteria such as *Escherichia coli*, *Salmonella*, and spirochetes can migrate by the rotational
motion of the flagellar motor fixed to the peptidoglycan layer, whereas *Spiroplasma* have a unique
swimming system in which kinks propagate along the cell body with a switch between left-handed
and right-handed cell helicity (Fig. 1A). This swimming is driven by an intracellular structure called
the "ribbon" which localizes along the innermost line of the helical cell structure, and structural
changes in the ribbon may switch the cell helicity (15, 16). Therefore, the detailed structure of the
ribbon should be elucidated to determine this swimming mechanism.

In previous studies using electron microscopy, two types of filaments in the ribbon were
visualized (15, 16). One comprises a protein "fibril," specific to *Spiroplasma*. The other is possibly
MreB, the bacterial actin homolog (14-16). Interestingly, all *Spiroplasma* species have as much
as five MreB classes (9, 17-19). As *Spiroplasma* MreBs are distantly located from other MreBs in
the phylogenetic tree, here, we use the term SMreB (17, 18). Fibril protein has been studied as a
linear motor protein which is thought to be responsible for the helicity-switching through
contraction and extension (16, 20, 21). The fibril filament is considered to function as a chain of
elliptical rings. However, the structure and function of the fibril protein remains unclear.

In the present study, we clarified the role of the fibril filament as the determinant of cell helicity,
using optical and electron microscopy (EM), and image analyses. Then, we proposed a scheme
for the helicity-switching swimming of *Spiroplasma.*

90
91
Results
Cell helicity is derived from the internal ribbon structure
To clarify which structure forms the helical cell morphology of *Spiroplasma*, we first measured the
helical pitches of the swimming cells using optical microscopy. The helical shape of the cells can
be observed as a series of density segments in the defocused image plane, relative to the cell
axis under phase contrast microscopy (Fig. 1B). The helical pitches between the left- and right-
handed segments along the cell axis were 696 ± 32 (n = 159) and 697 ± 37 nm (n = 146),
respectively. Next, we performed EM to analyze the internal ribbon structure to compare the
helical pitches of the cells and the ribbons. The cells were bound to EM grids non-specifically,
chemically fixed by glutaraldehyde, and then stained with uranyl acetate. Negative-staining EM
showed the images of helical-shaped cells with a narrow tip at one side (Fig. 1C). Next, we
exposed the internal ribbon structure by treating the cells with 0.1% Triton X-100 on the grid (Fig.
1D). The ribbon showed a "spiral" flat structure comprising protofilaments. However, generally, in
negative-staining EM, the specimens are placed in vacuum and dried and can result in
distortions, which is disadvantageous for helix observation. Therefore, we applied quick-freeze,
deep-etch (QFDE) EM to visualize the structure as close to the original as possible. In QFDE, a
sample is frozen in milliseconds, exposed by fracturing and etching, and then a platinum replica is
made by shadowing. The observation of the replica by transmission EM gives images with high
contrast and resolution much better than conventional scanning electron microscopy (SEM) (22,
23). The cells were non-specifically bound to mica flakes and fixed by quick freezing in a liquid
condition. Then, we prepared replicas by fracturing and platinum coating. QFDE-EM showed cell
morphology consistent with the results from negative-staining EM (Fig. 1E). Using QFDE-EM, we
also observed the ribbon exposed with 0.1% Triton X-100 treatment (Fig. 1F). The ribbon showed
the "helicoid" structure in which the twisted positions were aligned in a line. When the cells were
starved in phosphate-buffered saline (PBS) without glucose for 30 min, they all showed a left
helix with the same pitch. Therefore, we assumed that this structure is the default state of the cell.
Table 1 summarizes parameters that present the helicity of cells and ribbons obtained with the
two EM methods (Fig. 1G, H). The helical pitches of the cells and the ribbons were in good
agreement, indicating that the ribbon forms the cell helicity. As the helix diameter of the ribbon
was one-third that of the cell helix, the ribbon should be localized along the innermost line of the
cell helix.

Characterization of the isolated ribbon
We intended to use A22, an inhibitor of MreB polymerization to examine the role of MreBs in the
ribbon (Fig. 2A) (24), because the binding of A22 to SMreBs has been suggested from amino
acid sequences (17). First, we examined the effect of 1 mM A22 on swimming *Spiroplasma* cells
(Movie_S2). The cells shifted to a right-handed helix form and stopped moving in 2 min (Fig. 2B
a, b), suggesting that the functions of SMreBs are also inhibited by A22. The resulting helical
pitches of the cells were distributed around a peak at 426 ± 47 nm, shorter than the original (Fig.
2B c).

The cell suspension was treated with 1% Triton X-100 and subjected to stepwise gradient
centrifugation, consisting of 0%, 20%, 30%, 40%, 50%, and 60% sucrose layers. After

centrifugation, we found a dense layer of cell contents at the bottom of the 40% sucrose layer.
We recovered and observed the fraction under EM and found that the ribbon comprised
protofilaments with a width of 66 ± 12 nm and length longer than 2 μm (Fig. 2C a). The ribbons
were twisted with a pitch of 350 ± 17 nm (n = 47) (Fig. 2D a) consistent with the helical pitches of
the cells and the ribbons prepared on the grid (Fig. 1, Table 1) (P = 0.7 > 0.01). To analyze the
number and width of the protofilaments involved in the isolated ribbon, we traced a sectional
image profile of the ribbon (Fig. 2D b). Six to nine protofilaments were detected with widths
ranging between 4–16 nm (Fig. 2D c, d and Fig. S1). SDS-PAGE and peptide mass fingerprinting
analyses of this fraction showed five protein bands including six proteins (Table 2). The band (v)
was shown to contain SMreBs 2 and 4 (Table. S1). The whole ribbon fraction mainly comprised
fibril protein (band iii) and the protein mixture of SMreBs 2 and 4 (band v) with an intensity ratio of
47% and 37%, respectively (Fig. 2E).

Next, we examined the effects of A22 on the ribbon. We kept cells in 1 mM A22 for 2.5 h at 30 °C
and observed the ribbon. The ribbons were observed in a dispersed form with width distribution
characterized by three peaks, 9.5, 15.0, and 22.2 nm (Fig. 2C c, D d). SDS-PAGE analysis
clarified the content as 67% and 11% for fibril (band iii) and SMreB2 (band v) proteins,
respectively (Fig. 2E), suggesting that the protofilaments comprising fibril protein are bundled by
SMreBs in the ribbon structure.

Helical pitch of the isolated fibril filament
To analyze the detailed structure, we treated the ribbon fraction with cholic acid and isolated the
fibril protein using sucrose-gradient centrifugation. SDS-PAGE analysis showed that the fraction
contains only fibril protein (Fig. 3A). We examined the ATPase activity of fibril protein by
monitoring phosphate release because we expect ATP as the direct energy source for
Spiroplasma swimming. Fibril protein at 17 μM was mixed with 2 mM ATP in the presence of Mg
ions, but phosphate was not released even after 250 min at 30 °C. Then, we concluded that the
fibril protein does not have the obvious ATPase activity. Negative-staining EM showed that the
fibril protein forms filaments which include single- and double-stranded filaments, suggesting
various types of interactions between fibril protein molecules (Fig. 3B a). A single-stranded fibril
filament consisted of repeated ring units approximately 10 nm long (Fig. 3B b) and 7 nm wide as
observed in the side view (Fig. 3B c). A double-stranded fibril filament appeared to be formed by
aligning two single-stranded filaments to face each other at the side of the ring, resulting in a
thickness and width of 14 nm and 7 nm, respectively (Fig. 3B d), forming a 10 nm long repeating
structure. We analyzed the helical pitches for the double-stranded fibril filaments, because the
double-stranded fibril filament had enough persistent length to cover the helical pitch with a clear
twist of the ring pattern along the filament axis. The images of the fibril filament cropped from the
electron micrographs using the straightening selection tool of ImageJ software were subjected to
Fourier filtering to remove noise (Fig. 3C). To ensure that the FFT images did not include artificial
mistakes, we also generated the two-dimensional (2D) averaged images from 312 randomly
picked particles with a box size of 45 nm along the filament axis using RELION 3.0 software (Fig.
3D)(25). The three classes of images corresponded well to different positions of the filament
images generated by FFT, suggesting that the FFT images reflect the original structures. The
helical pitch estimated from the FFT images was 343 ± 22 nm (n = 158) (Fig. 3G a). However, we
could not conclude the handedness of the fibril filament, because the alignment of the filament on
the EM grid was not distinguishable with negative-staining EM. Therefore, we analyzed the
isolated fibril filament using QFDE-EM (Fig. 3F) because the replica is made by platinum, which is
not transparent to an electron beam. We succeeded in the determination of their handedness

(Fig. 3F) and concluded that the double-stranded fibril filament forms a left-handed helix with a
pitch distribution of 351 ± 33 nm (n = 50) (Fig. 3G b). The helix features were consistent with the
pitch of the cell in the default state and that of the isolated ribbon (Table 1), suggesting that the
fibril pitch is the determinant for the pitch of the ribbon and cell.

Three-dimensional reconstruction of fibril filaments
To detect the conformational changes in the fibril three dimensionally, we performed single
particle analysis based on negative-staining EM. The double-stranded fibril filament was not
suitable for image averaging owing to the positional variety in the binding of the two filaments
(Fig. 3 and Fig. S2). Therefore, we sonicated the purified fibril fraction to increase the proportion
of single-stranded forms and succeeded in the acquisition of single-stranded images (Fig. 4A).
From the selected 11 867 particles with good quality, the 2D-averaged images were summarized
into three types (i), (ii), and (iii) (Fig. 4A b). Then, the initial three-dimensional (3D) model was
reconstructed using the *ab-initio* 3D function of cisTEM software (26) and used as the reference
for the following 3D classification (Fig. 4A c). 3D structures of the fibril filament reconstructed from
the total of 11 867 particles by RELION 3.0 software showed three conformations, i.e., class 1:
left-handed mostly straight (49%), class 2: left-handed with curvature (24%), and class 3: right-
handed with curvature (27%) (Fig. 4A d and Fig. S3). Class 1 structure reconstituted with
rotational symmetry (C2) was not significantly different from that without symmetry (C1),
suggesting that fibril filament has rotational symmetry without filament directionality (Fig. S3).
Then, we calculated the structures of fibril filaments as C2 structures.

Two dimensional re-projections from these three structures corresponded well to the 2D class
averages, indicating the validity of the obtained 3D structures (Fig. S3). The 3D structure of the
fibril filament had repeating elliptical rings with a pitch of 8.7 nm along the filament axis, with long
and short axes of 11 and 6 nm, respectively. A short cylinder connects the ring units causing a
positive curvature (Fig. 4A d). These characteristics were common to all three classes. Cross-
sectional images perpendicular to the longitudinal axis of the filament showed a "boomerang-like
shape", which had a dent at one side and protrusion at the other (Fig. 4B). We evaluated the twist
of the fibril filament around the filament axis by measuring the directions of the Feret diameter,
the longest diagonal line of cross-sectional images. Fibril filaments twisted along the filament
axis. The twisting angles were estimated from the angle averages of the first and fourth units, as
5.9 (left-handed), 7.3 (left-handed), and 9.7 (right-handed) degrees for classes 1, 2, and 3,
respectively. Although superimposition of classes 2 and 3 showed diagonal shifts between these
two conformations, the positions responsible for the structural shift could not be identified owing
to the low resolution of structures (Fig. 4C). We constructed long filament models representing a
helical pitch by stacking the ring units (Fig. 4D). Class 1 formed a left-handed helix with a
diameter of 75 nm and a pitch of 290 nm. Class 2 formed the left-handed helix with a diameter of
80 nm and a pitch of 150 nm. Class 3 formed the right-handed helix with a diameter of 45 nm and
a pitch of 200 nm. We attempted to reconstruct the double stranded filaments through image
averaging and failed (Fig. S2). It is likely caused by the variation in binding positions in the
formation of the double strand. We reconstructed the double strand from the structure of the
single-stranded filament (Fig. 3E), which was consistent with the images obtained using the other
methods.

Although we found variations in curvatures and twists among the three classes, no variations
were detected in the lengths of periodic structures comprising rings and cylinders (Fig. 4A).

Handedness verified from tomography of QFDE replica

The 3D images reconstituted from negative-staining EM have common features, although they
showed variations in curvature and twist. They have rings and cylinders rising to the right along
the filament axis when they are viewed from front and back sides, respectively (Fig. 4A d),
showing that the three classes have the same handedness. As the images from negative-staining
EM are transparent, the structures reconstituted here may be mirror images of the real structures.
Then, we intended to verify the handedness of the reconstituted structures by tomography of
replica from QFDE EM (Fig. 5), because tomogram cannot be a mirror image (27, 28).

We made QFDE replicas from the fraction containing single-stranded fibril filaments, acquired
images every 1.5 degree to 50 degrees for both sides, reconstituted a tomogram（Movie_S4）
（Fig. 5A）, and then a structure was obtained by averaging 60 subtomograms（Fig. 5B）. The
resulting filament structure was featured by rings and cylinders, as expected. The rings and the
cylinders were tilted from the filament axis, rising to the right relative to the filament axis with 4–5
and 74–82 degrees viewed from front, and back, respectively (Fig. 5C, S4, S5), which are
consistent with the features of structures from negative-staining EM（Fig. S6）. These results
show that the classes of structures from negative-staining EM have the same handedness with
the real structures (Fig. 5C).

Discussion
242
Fibril and ribbon structures
Although the interconnected ring structure in the fibril filament has been observed (14, 20), the 3D
reconstruction of the fibril has not been achieved to date. In the present study, we clarified the 3D
structure of the fibril filament for the first time. The sonication in the isolation process was
effective in isolating the single stranded filament, whose uniform structure was advantageous for
image averaging (Fig. 4). The structure determined here showed a width of 10.5 nm, and was in
good agreement with the corresponding filament structure obtained through electron
cryotomography (16), suggesting that the filaments isolated here retained the original structure.
The left-handed conformation accounted for 73% of the fibril filament, suggesting that this
conformation is more stable than others, as an intrinsic character of fibril protein. This observation
may explain the fact that both the cells in a default state and the isolated ribbons were mostly left-
handed (29, 30). The fibril structure is likely more stable in the left-handed conformation than in
the right-handed one. The fibril filament did not show any polarity along the filament axis,
although *Spiroplasma* cells swim in a directed manner (Fig. 1) (9, 14). This directionality could be
caused by structures other than the fibril, for example, SMreB proteins and the dumbbell formed
at the cell front (14, 18, 31). We obtained three different filament conformations with different
curvatures, in other words, the helix diameter. This variety in curvature may function in directing
the ribbon formation, in which the filament at the ribbon edges requires a longer helix diameter
than those at the central positions (Fig. 6). Attractive models have been suggested to explain the
mechanism of helicity shift, where coupled length changes in fibril filaments cause torsion and
helicity shift (16, 20). However, the dimensions of periodical structure comprising ring and
cylinder did not show significant variations (Fig. 4), suggesting that the helicity shift is not resulted
from the conformational changes of fibril filaments.

Swimming mechanism

Based on these results, we can now suggest the core part of the helicity-switching mechanism (Fig. 6). The ribbon comprises 6–9 fibril filaments connected laterally and oriented along the innermost part of the helical cell (Fig. 6A). The fibril should support the cell membrane through their ring structures, because the fibril filament has a positive curvature toward the backbone (Fig. 6B a upper). Thus, the fibril twist forms the twist of the ribbon and the cell with the same handedness, because the fibril filament binds to the adjacent fibril filaments through their fixed positions (Fig. 6B a lower). If the fibril filaments in the ribbon have strong cooperativity along the ribbon axis and transmit the twist to the next levels, the helicity shift travels along the ribbon axis, with accumulation of the rotational angle (Fig. 6B b) (29). If we assume that such a ribbon is fixed at a tip on the front end, then, the shift initiated at the tip travels backward, along with rotation of the back portion to release the torsion, in a direction that pushes water backward (Fig. 6C) (Movie_S1). The ribbon structure drives the cell structure and propels the cell forward by pushing water backward like a screw.

Possible molecular mechanism for helicity-switching

The N-terminal region of the fibril protein has an amino acid sequence identity of as much as 25.6% with MTA/SAH nucleosidase from a bacterial species, *B. anthracis*. This protein is essential for bacterial growth because it recycles adenine and methionine through S-adenosylmethionine (SAM)-mediated methylation reactions and produces the universal quorum-sensing signal, autoinducer-2 (AI-2) (32). The fibril protein probably evolved from this protein, which is abundant in a cell, by acquiring a C-terminal region possessing polymerization activity. Generally, the energy for motility is supplied from ATP hydrolysis or membrane potential (2). Class Mollicutes lack the respiration pathway to generate membrane potential and produce ATP through metabolism such as glycolysis and arginine fermentation (33). In Mollicutes, the membrane potential is generated from ATP hydrolysis and not the primary energy source. Therefore, the energy for swimming should also be supplied by ATP rather than the membrane potential. In fact, the two motility mechanisms of Mollicutes genus, *Mycoplasma mobile*-type and *Mycoplasma pneumoniae*-type gliding mechanisms depend on the hydrolytic energy of ATP (34-37). The fibril protein derived from the MTA/SAH nucleosidase is unlikely to have ATPase activity, and no ATPase activity was detected in the fibril fraction. These facts suggest that other proteins may be involved in the helicity switching (16, 18, 38). The ribbon contains SMreB proteins, and interestingly, most genomes of the *Spiroplasma* code for five classes of SMreBs (17-19). Moreover, in the present study, we showed that SMreBs have roles to bundle fibril filaments (Fig. 2 C). Generally, MreBs exhibit polymerization dynamics based on ATP hydrolysis, and function to assign peptidoglycan synthesizing complexes (2, 22, 24, 39) and as a rail for the gliding motor of *Myxococcus xanthus* (40). SMreB5, a member in the five classes, is known to be essential for helix formation and swimming of *Spiroplasma citri* cells (18). These facts suggest that SMreBs support and drive the helices formed by the fibril. Perhaps, SMreB itself also forms filamentous structures along the cell axis (15, 16). In fact, MreB-like filaments have been observed along the cell axis by EM analyses of *Spiroplasma melliferum* (16).

Materials and Methods

Bacterial strains and culture conditions

The type strain, TDA-040725-5^T, of *Spiroplasma eriocheiris* was cultured in R2 medium (2.5%

312   [wt/vol] heart infusion broth, 8% sucrose, and 10% horse serum) at 30 °C to an optical density of

313   0.06 to 0.1 at 600 nm (14, 41).

Optical microscopy

Cultured cells were centrifuged at 11 000 × g, 10 °C for 10 min and suspended in PBS consisting

of 75 mM sodium phosphate [pH 7.3], 100 mM NaCl, containing 20 mM glucose and 0.6%

methylcellulose to be of a cell density 10-fold higher than that of the original (14, 41). The cells

were inserted into a tunnel chamber assembled by taping coverslips as previously described and

observed under an IX71 microscope (Olympus, Tokyo, Japan) (42). The video was captured

using a DMK33UX174 complementary metal–oxide–semiconductor (CMOS) camera (The

Imaging Source, Taipei, Taiwan) and analyzed using ImageJ v1.53a (https://imagej.nih.gov/ij/).

Electron microscopy

To observe the intact cells, the cell suspension was placed on a hydrophilized grid, fixed using

2% glutaraldehyde, washed with water, and stained with 2% uranyl acetate. To observe the

internal structure, the cell suspension on a grid was treated with PBS including 0.1 mg/mL DNase

and 1 mM $MgCl_2$ for 20 s, and washed and stained with 2% uranyl acetate. QFDE-EM was

performed as previously reported for specimens suspended with mica flakes (22). The images

were acquired using a JEM1010 EM (JEOL, Akishima, Japan) equipped with a FastScan-F214(T)

charged-coupled device (CCD) camera (TVIPS, Gauting, Germany) and analyzed using ImageJ

v1.53a. For tomography, the images were captured by Talos F200C EM (FEI, Eindhoven,

Netherlands) equipped by 4k × 4K Ceta CMOS camera (FEI). Single-axis tilt series were

collected covering an angular range from -50° to +50° with 1.5° steps and analyzed by IMOD (ver

4.11) and PEET (ver 1.15.0).

Isolation of ribbon and fibril

To isolate the internal structure, 10 mL of cell suspension in PBS was treated with 1% Triton X-

100, 0.1 mg/mL DNase, 1 mM $MgCl_2$, and 0.1 mM PMSF with shaking for 10 min at 4 °C. The

insoluble fraction was recovered by centrifugation at 20 000 × g, 4 °C for 30 min and suspended

in PBS to be 0.2 mL. The sample was laid at the top of sucrose solution layers of 0%, 20%, 30%,

40%, 50%, and 60%, and centrifugated at 20 000 × g, 4 °C for 20 min in a 1.5 mL tube with a

fixed angle. To isolate the fibril filament, the insoluble fraction was additionally treated with a

solution consisting of 2% choric acid, 20 mM Tris-Cl pH 8.0, 150 mM NaCl at 4 °C for 8 h and

subjected to stepwise density gradient centrifugation. SDS-PAGE and peptide mass fingerprinting

were performed as previously described (14, 43, 44).

ATPase assay

345   ATPase activity was assayed by a continuous spectrophotometric method using a 2-amino-6-

346   mercapto-7-methylpurine ribonucleoside–purine nucleoside phosphorylase reaction to detect the

347   released inorganic phosphate at 30 °C (EnzChek kit; Life Technologies, Carlsbad, CA, USA). The

348   reaction mixture was as follows: 17 µM purified fibril filament, 1 mM $MgCl_2$, 20 mM Tris-HCl (pH

349   7.5) in a total volume of 0.2 mL (45).

Reconstitution of 3D structure

The contrast transfer function (CTF) parameters for negative-staining EM images were estimated

using Gctf25 software (46). The images of fibril filaments were selected automatically by RELION

3.0 (25) as helical objects and segmented as squares of 200 × 200 pixels with 90% overlap.

These 14 543 images were 2D-classified and 11 867 images were selected for further analyses.

ab-initio reconstitution was performed by cisTEM (26) based on segmented images from 12
classes. The selected 11 867 particle images were 3D-classified using the 3D map in RELION 3.0
(25).

358
359

Acknowledgments
We thank Yuhei O Tahara, Daichi Takahashi, Hana Kiyama, and Ikuko Fujiwara at the Graduate
School of Science, Osaka City University, Japan for helpful discussions. This study was
supported by a Grants-in-Aid for Scientific Research A (MEXT KAKENHI, Grant Number
JP17H01544), JST CREST (Grant Number JPMJCR19S5), the Osaka City University (OCU)
Strategic Research Grant 2017 for top priority researches to MM, and JSPS KAKENHI (Grant
Number JP25000013), the Platform Project for Supporting Drug Discovery and Life Science
Research (BINDS) from AMED (Grant Number JP19am0101117 and support number 1282), the
Cyclic Innovation for Clinical Empowerment (CiCLE) from AMED (Grant Number
JP17pc0101020), and JEOL YOKOGUSHI Research Alliance Laboratories of Osaka University to
KN.

371
372
373

References

1.   H. Grosjean *et al.*, Predicting the minimal translation apparatus: lessons from the
reductive evolution of mollicutes. *PLoS Genet* **10**, e1004363 (2014).
2.   M. Miyata *et al.*, Tree of motility - A proposed history of motility systems in the tree of
life. *Genes Cells* **25**, 6-21 (2020).
3.   S. Razin, L. Hayflick, Highlights of mycoplasma research--an historical perspective.
Biologicals **38**, 183-190 (2010).
4.   S. Razin, D. Yogev, Y. Naot, Molecular biology and pathogenicity of mycoplasmas.
Microbiol Mol Biol Rev **62**, 1094-1156 (1998).
5.   T. Toyonaga *et al.*, Chained Structure of Dimeric F1-like ATPase in Mycoplasma mobile
Gliding Machinery. *mBio* 10.1128/mBio.01414-21, e0141421 (2021).
6.   M. Miyata, T. Hamaguchi, Prospects for the gliding mechanism of *Mycoplasma mobile*.
Curr Opin Microbiol **29**, 15-21 (2016).
7.   M. Miyata, T. Hamaguchi, Integrated information and prospects for gliding mechanism
of the pathogenic bacterium *Mycoplasma pneumoniae*. *Front Microbiol* **7**, 960 (2016).
8.   M. Miyata, Unique centipede mechanism of *Mycoplasma* gliding. *Annu Rev Microbiol* **64**,
519-537 (2010).
9.   S. Harne, P. Gayathri, L. Beven, Exploring *Spiroplasma* biology: opportunities and
challenges. *Front Microbiol* **11**, 589279 (2020).
10.   H. Wada, R. R. Netz, Hydrodynamics of helical-shaped bacterial motility. *Phys Rev E Stat*
Nonlin Soft Matter Phys **80**, 021921 (2009).
11.   J. W. Shaevitz, J. Y. Lee, D. A. Fletcher, *Spiroplasma* swim by a processive change in body
helicity. *Cell* **122**, 941-945 (2005).
12.   T. Harumoto, B. Lemaitre, Male-killing toxin in a bacterial symbiont of *Drosophila*.
Nature **557**, 252-255 (2018).
13.   G. E. Gasparich, *Spiroplasmas*: evolution, adaptation and diversity. *Front Biosci* **7**, d619-
640 (2002).
14.   P. Liu *et al.*, Chemotaxis without conventional two-component system, based on cell
polarity and aerobic conditions in helicity-switching swimming of *Spiroplasma*
eriocheiris. *Front Microbiol* **8**, 58 (2017).
15.   S. Trachtenberg *et al.*, Structure of the cytoskeleton of *Spiroplasma melliferum* BC3 and
its interactions with the cell membrane. *J Mol Biol* **378**, 778-789 (2008).
16.   J. Kürner, A. S. Frangakis, W. Baumeister, Cryo-electron tomography reveals the
cytoskeletal structure of *Spiroplasma melliferum*. *Science* **307**, 436-438 (2005).
17.   D. Takahashi, I. Fujiwara, M. Miyata, Phylogenetic origin and sequence features of MreB
from the wall-less swimming bacteria *Spiroplasma*. *Biochem Biophys Res Commun* **533**,
638-644 (2020).
18.   S. Harne *et al.*, MreB5 Is a determinant of rod-to-helical transition in the cell-wall-less
bacterium *Spiroplasma*. *Curr Biol* **30**, 4753-4762.e4757 (2020).
19.   C. Ku, W. S. Lo, C. H. Kuo, Molecular evolution of the actin-like MreB protein gene family
in wall-less bacteria. *Biochem Biophys Res Commun* **446**, 927-932 (2014).
20.   S. Cohen-Krausz, P. C. Cabahug, S. Trachtenberg, The monomeric, tetrameric, and
fibrillar organization of Fib: the dynamic building block of the bacterial linear motor of
Spiroplasma melliferum BC3. *J Mol Biol* **410**, 194-213 (2011).
21.   R. Townsend, D. B. Archer, K. A. Plaskitt, Purification and preliminary characterization of
Spiroplasma fibrils. *J Bacteriol* **142**, 694-700 (1980).

22.   I. Tulum, Y. O. Tahara, M. Miyata, Peptidoglycan layer and disruption processes in
Bacillus subtilis cells visualized using quick-freeze, deep-etch electron microscopy.
Microscopy (Oxf) **68**, 441-449 (2019).

23.   J. E. Heuser, The origins and evolution of freeze-etch electron microscopy. *J Electron*
Microsc (Tokyo) **60 Suppl 1**, S3-29 (2011).

24.   H. Shi, B. P. Bratton, Z. Gitai, K. C. Huang, How to build a bacterial cell: MreB as the
foreman of *E. coli* construction. *Cell* **172**, 1294-1305 (2018).

25.   J. Zivanov *et al.*, New tools for automated high-resolution cryo-EM structure
determination in RELION-3. *Elife* **7**, e42166 (2018).

26.   T. Grant, A. Rohou, N. Grigorieff, cisTEM, user-friendly software for single-particle image
processing. *Elife* **7**, e35383 (2018).

27.   A. Briegel, M. Pilhofer, D. N. Mastronarde, G. J. Jensen, The challenge of determining
handedness in electron tomography and the use of DNA origami gold nanoparticle
helices as molecular standards. *J Struct Biol* **183**, 95-98 (2013).

28.   G. J. Jensen (2015) Getting started in Cryo-EM with professor Grant Jensen. p
https://www.youtube.com/watch?v=gDgFbAqdM_c.

29.   D. Nakane, T. Ito, T. Nishizaka, Coexistence of two chiral helices produces kink
translation in *Spiroplasma* swimming. *J Bacteriol* **202**, e00735-00719 (2020).

30.   J. F. Boudet *et al.*, Large variability in the motility of spiroplasmas in media of different
viscosities. *Sci Rep* **8**, 17138 (2018).

31.   M. Garnier, M. Clerc, J. M. Bove, Growth and division of *Spiroplasma citri*: elongation of
elementary helices. *J Bacteriol* **158**, 23-28 (1984).

32.   N. Parveen, K. A. Cornell, Methylthioadenosine/S-adenosylhomocysteine nucleosidase, a
critical enzyme for bacterial metabolism. *Mol Microbiol* **79**, 7-20 (2011).

33.   C. M. Fraser *et al.*, The minimal gene complement of *Mycoplasma genitalium*. *Science*
270, 397-403 (1995).

34.   M. Mizutani, M. Miyata, Behaviors and energy source of *Mycoplasma gallisepticum*
gliding. *J Bacteriol* **201**, e00397-00319 (2019).

35.   Y. Kinosita, M. Miyata, T. Nishizaka, Linear motor driven-rotary motion of a membrane-
permeabilized ghost in *Mycoplasma mobile*. *Sci Rep* **8**, 11513 (2018).

36.   Y. Kinosita *et al.*, Unitary step of gliding machinery in *Mycoplasma mobile*. *Proc Natl*
Acad Sci U S A **111**, 8601-8606 (2014).

37.   A. Uenoyama, M. Miyata, Gliding ghosts of *Mycoplasma mobile*. *Proc Natl Acad Sci U S A*
102, 12754-12758 (2005).

38.   S. Trachtenberg, P. Schuck, T. M. Phillips, S. B. Andrews, R. D. Leapman, A structural
framework for a near-minimal form of life: mass and compositional analysis of the
helical mollicute *Spiroplasma melliferum* BC3. *PLoS One* **9**, e87921 (2014).

39.   A. J. F. Egan, J. Errington, W. Vollmer, Regulation of peptidoglycan synthesis and
remodelling. *Nat Rev Microbiol* **18**, 446-460 (2020).

40.   G. Fu *et al.*, MotAB-like machinery drives the movement of MreB filaments during
bacterial gliding motility. *Proc Natl Acad Sci U S A* **115**, 2484-2489 (2018).

41.   N. Terahara, I. Tulum, M. Miyata, Transformation of crustacean pathogenic bacterium
Spiroplasma eriocheiris and expression of yellow fluorescent protein. *Biochem Biophys*
Res Commun **487**, 488-493 (2017).

42.   A. Uenoyama, A. Kusumoto, M. Miyata, Identification of a 349-kilodalton protein
465   (Gli349) responsible for cytadherence and glass binding during gliding of *Mycoplasma*
466   *mobile*. *J. Bacteriol.* **186**, 1537-1545. (2004).
467   43.   Y. Kawakita *et al.*, Structural study of MPN387, an sssential protein for gliding motility of
468   a human-pathogenic bacterium, *Mycoplasma pneumoniae*. *J Bacteriol* **198**, 2352-2359
469   (2016).
470   44.   D. Nakane, M. Miyata, Cytoskeletal "jellyfish" structure of *Mycoplasma mobile*. *Proc Natl*
471   *Acad Sci U S A* **104**, 19518-19523 (2007).
472   45.   M. S. Nishikawa *et al.*, Refined mechanism of *Mycoplasma mobile* gliding based on
473   structure, ATPase activity, and sialic acid binding of machinery. *mBio* **10**, e02846-02819
474   (2019).
475   46.   K. Zhang, Gctf: Real-time CTF determination and correction. *J Struct Biol* **193**, 1-12
476   (2016).

479   **Table 1.** Dimensions of cell and ribbon

Parameters	Negatively stained electron microscopy	Quick-freeze, deep-etch electron microscopy	Optical microscopy
Cell thickness (a)	196 ± 26 nm	179 ± 23 nm	
Ribbon width (b)	147 ± 24 nm	118 ± 17 nm	
Cell helical pitch (c)	696 ± 73 nm	706 ± 43 nm (LH)	696 ± 32 nm (LH) 697 ± 37 nm (RH)
Ribbon 1/2 helical pitch (d)	348 ± 44 nm	350 ± 30 nm (LH)	
Ribbon helical pitch (e)	684 ± 60 nm	700 ± 60 nm (LH)	
Ribbon helical diameter (f)	137 ± 30 nm	118 ± 17 nm	
Cell helical diameter (g)	383 ± 45 nm	310 ± 26 nm	
Isolated ribbon pitch	350 ± 17 nm		
Isolated fibril pitch	343 ± 22 nm	351 ± 34 nm (LH)	

480   *, ** $p > 0.05$ (the agreements between cell pitch and ribbon pitch were supported by Student's t-test)
481
482

484 **Table 2.** Protein components of ribbon isolated from original cells[1].

Protein band	Gene ID	Annotation	Mascot Score[2]	Mass (kDa)[3]	Density ratio (%)	
					A22 untreated	A22 treated
(i)	SPE-1201	Hypothetical protein	72	85.8	4	5
(ii)	SPE-0013	FtsH	84	77.0	12	17
(iii)	SPE-0666	Fibril	206	58.7	47	67
(iv)	SPE-1231	MreB5	98	38.5	10	7
(v)	SPE-1224 SPE-1230	MreB2 MreB4	80	37.8 40.7	27	4

486 [1] From A22-treated cells, the proteins common with the original cells were identified for bands (i)–(iv). For band (v) only MreB2 was identified.

487 [2] Mascot Score is logarithm of probability that the observed match is a random event.

488 [3] Calculated from amino acid sequence as monoisotopic molecule.

Figures

3
Figure 1. Cell helicity derived from ribbon structure. (A) Schematic of helicity-switching
swimming. The swimming direction is indicated by an arrow. (B) Phase-contrast microscopy of
swimming cell. The blue and red segments indicate the left- and right-handed helicity,
respectively (upper). Histograms of both left- and right-handed helical pitches of swimming cells,
fitted by Gaussian curves with peak tops of 696 ± 32 and 697 ± 37 nm, respectively (lower). (C,
E) Field image of cells acquired by negative-staining and QFDE EM. (D, F) Intact cells (leftmost)

and ribbons prepared on grids (others). (G, H) Dimensions marked in cell schematics (a)-(g) (left)
are summarized in in histograms (right) and Table 1.
12

Figure 2. Isolation and characterization of ribbon. (A) Schematic for isolation and disassembly of
ribbon. (B) Effects of A22 on swimming cells. (a) Untreated. (b) A22-treated for 2 min. (c)
Histogram of the helical cell pitches after A22 treatment, fitted by a Gaussian curve with a peak at
426 ± 47 nm. (C) EM observation of isolated ribbon fractions. (a) The whole structure of the
isolated ribbon with helicity as shown by periodical wide positions (marked by arrows). (b) The
magnified image of the isolated ribbon and the helical pitch is indicated by a bidirectional arrow.
(c) Disassembled ribbon composed of protofilaments. (D) Numerical characterization. (a)
Histogram for the helical pitches of the isolated ribbon, fitted by a Gaussian curve with a peak at
351 ± 16 nm. (b) Sectional image profile of the area boxed in panel (C-b). The peaks correspond
to the center of protofilament. (c) Histogram for the number of protofilaments involved in a ribbon.
(d) Histogram for the protofilament width in ribbons. The distribution can be fitted by two
Gaussian curves marked (i) and (ii), with peaks around 7.0 and 10.5 nm, respectively. (Inset) The
images of reconstituted structures viewed from different angles (refer to Fig. 4). (e) Histogram for
the width of the protofilament in disassembled ribbons, fitted with three Gaussian curves with
peak tops of 9.5, 15, 22.2, and 30.1 nm. (E) SDS-10% PAGE of the ribbon fraction isolated from
original and A22-treated cells.
30
31

32
33
Figure 3. Helical pitches of isolated fibril filament. (A) Protein profiles of fractions in purification
process for fibril protein. (B) Purified fibril filaments observed by negative-staining EM. (a) Field
image. White and black arrows indicate single and double strands, respectively. (b, c) Front and
side views of the single-stranded fibril filaments. (d, e) Front and side views of the double-
stranded fibril filaments. (C) Double stranded filaments reconstituted through Fourier transform.
(D) Averaged images of double-stranded filaments reconstructed from 62, 11, and 23 images,
respectively for topmost, second and third panels. Repeated alignment of three images shown for
comparison with panel (C). (E) Model for the double-stranded fibril filament (Refer to Fig. 4). (F)
Fibril filaments observed by QFDE-EM. Field (a), single-stranded filament (b), and double-
stranded filaments (c, d) are presented. (G) Histograms of the helical pitches for the fibril
filaments measured for negative-staining (a) and QFDE (b) EM observations. Pitches were fitted
by Gaussian curves with peaks at 343 ± 22 and 351 ± 34 nm, respectively.
46
47

Figure 4. Three-dimensional reconstruction of fibril filaments. (A) Workflow of single particle
analysis by negative-staining EM. (a) Single-stranded fibril filaments prepared by sonication. (b)
Averaged images by "2D classification". (c) The initial 3D model generated by *ab-initio*
reconstruction of cisTEM software. (d) Three different conformations of the fibril filament
reconstituted by "3D classification". (B) Rotation of repeated units along the filament axis. Angles
of Feret diameter for the cross-sectional images were plotted along the filament is positions with a
Feret diameter longer than 80% of the filament maximum (right). (C) Superpose of class 1 (left
handed) and class 3 (right handed) structures. The fitting reference is indicated by the dashed
box. (D) Three types of fibril filament models showing 44 repeated units with dimensions in nm.

60
Figure 5. Comparison of 3D structures of fibril filament reconstructed from QFDE replica and
negative-staining EM methods. (A) Replica image of single-stranded fibril filaments. Left: A field
image is shown from a tilt series (Movie_S4). Right lower: Magnified images of single fibril
filament are shown as a raw image (a), a slice from tomogram (b), and a subtomogram (c). (B)
Structure averaged from 60 subtomograms. The leftmost and other images are presented under
different thresholds. (C) Superpose of 3D structures from single-particle analysis (grey) and
subtomogram averaging (magenta). Long axes of ring and cylinder are shown by broken red
arrows. The filament axes were detected by a function "relion_align_symmetry --sym d2" in
RELION-3.0.
70

71
Figure 6. Model schematic for swimming. (A) Ribbon alignment along innermost line of helical
cell body. (B) Relation between helicity of fibril filament and ribbon. (a) The ribbon is composed of
fibril filaments. Cross sections of ribbon at positions (i) and (ii) are shown in lower. The filaments
show positive curvature to the cytoplasmic side. The twists of individual fibril filaments result in
the twist of the whole ribbon, because the filaments bind to the neighboring filaments through
their fixed positions. (b) The twist of fibril filament accumulated through the stack of unit. The
horizontal positions of oval ends are shown for each unit. The positional difference of the end
points are shown by lines between red and blue dots. (c) Curvatures of the protofilament. For
ribbon formation, each protofilament should take different curvature. (C) Helix rotation caused by
accumulated twist (Movie_S3). The whole and small stack of ribbon are presented, respectively,
in left and right, for each of three conformations (i), (ii), and (iii). The ribbon is stabilized by a "Tip"
at the front end. The left-handed ribbon (i), the transition state of the ribbon from left-handed to
right-handed by helicity-switching, with a switch point traveling from front to back (ii), and the
resulted right-handed ribbon (iii) are presented. The areas of left-handed and right-handed are
marked by blue and red outlines, respectively. Rotation of small stacks from conformation (i) are
presented by traces with transition from light to deep colors. The twist of ribbon accumulates with
the switch traveling. The rotations caused by the accumulated twist pushes water backward,
resulting in the propelling force.
90