Isolation and structure of the fibril protein, a major component of the internal ribbon for *Spiroplasma* swimming

Yuya Sasajima[1], Takayuki Kato[2], Tomoko Miyata[3], Akihiro Kawamoto[3], Keiichi Namba[3,4,5] and Makoto Miyata[1,6*]

[1]Graduate School of Science, Osaka Metropolitan University, Osaka, Japan

[2]Institute for Protein Research, Osaka University, 3-2 Yamadaoka, Suita, Osaka, Japan

[3]Graduate School of Frontier Biosciences, Osaka University, 1-3 Yamadaoka, Suita, Osaka, Japan

[4]RIKEN Center for Biosystems Dynamics Research and Spring-8 Center, 1-3 Yamadaoka, Suita, Osaka, Japan

[5]JEOL YOKOGUSHI Research Alliance Laboratories, Osaka University, 1-3 Yamadaoka, Suita, Osaka, Japan

[6]The OCU Advanced Research Institute for Natural Science and Technology (OCARINA), Osaka Metropolitan University, Osaka, Japan.

*Correspondence:

Makoto Miyata

miyata@omu.ac.jp

Keywords: Helical shape, Motility, Cytoskeleton, Filament, Single particle analysis, Quick freeze replica electron microscopy, Electron tomography.

ABSTRACT

Spiroplasma, which are known pathogens and commensals of arthropods and plants, are helical-shaped bacteria that lack a peptidoglycan layer. *Spiroplasma* swim by alternating between left- and right-handed helicity. Of note, this system is not related to flagellar

motility, which is widespread in bacteria. A helical ribbon running along the inner side of

the helical cell should be responsible for cell helicity and comprises the bacterial actin

homolog, MreB, and a protein specific to *Spiroplasma*, fibril. Here, we isolated the

ribbon and its major component, fibril filament, for electron microscopy (EM) analysis.

Single-particle analysis of the fibril filaments using the negative-staining EM revealed a

three-dimensional chain structure composed of rings with a size of 11 nm wide and 6 nm

long, connected by a backbone cylinder with an 8.7 nm interval with a twist along the

filament axis. This structure was verified through EM tomography of quick-freeze deep-

etch replica sample, with a focus on its handedness. The handedness and pitch of the

helix for the isolated ribbon and fibril filament agreed with those of the cell in the resting

state. Structures corresponding to the alternative state were not identified. These results

suggest that the helical cell structure is supported by fibril filaments; however, the helical

switch is caused by the force generated by the MreB proteins. The isolation and structural

outline of the fibril filaments provide crucial information for an in-depth clarification of

the unique swimming mechanism of *Spiroplasma*.

1   Introduction

Mollicutes, which are parasitic or commensal bacteria, evolved from the phylum,

Firmicutes, including *Bacillus* and *Clostridium* by reducing their genome size (Razin et

al., 1998;Razin and Hayflick, 2010;Grosjean et al., 2014;Miyata et al., 2020). During the

course of evolution, the cells became softer and smaller owing to the loss of the

peptidoglycan layer. These changes may have allowed some species to transmit their

internal housekeeping activities, such as the rotation of ATP synthesis to the outside,

resulting in the acquisition of at least three unique motility mechanisms (Relich et al.,

2009;Miyata and Hamaguchi, 2016a;b;Distelhorst et al., 2017;Miyata et al.,

2020;Toyonaga et al., 2021). Two of the three well studied mechanisms are exerted by

Mycoplasma mobile and *Mycoplasma pneumoniae*. These species exhibit gliding

motilities on solid surfaces, in which leg structures repeatedly catch sialylated

oligosaccharides on host cells based on two mechanisms (Miyata, 2010;Miyata and

Hamaguchi, 2016a;b). Another motility system is the helicity-switching swimming of

Spiroplasma, which is the subject of the present study (Movie_S1) (Shaevitz et al.,

2005;Wada and Netz, 2009;Harne et al., 2020b;Sasajima and Miyata, 2021). *Spiroplasma*

species are parasitic to plants and arthropods and are characterized as polarized helical-

shaped cells with one tapered end (Gasparich, 2002;Harumoto and Lemaitre, 2018;Harne

et al., 2020b). These species exhibit obvious chemotaxis despite the absence of genes for

the two-component regulatory system in the genome, which is generally responsible for

bacterial chemotaxis (Liu et al., 2017). In general, swimming bacteria, including

spirochetes, can migrate through the rotational motion of the flagellar motor fixed to the

peptidoglycan layer, whereas *Spiroplasma* has a unique swimming system in which kinks

propagate along the cell body with a switch between left- and right-handed cell helicity

(Fig. 1A). The outline of this mechanism has been clarified as follows. The rotation of

helical cells linked to the helicity switch pushes the water back (Trachtenberg and Gilad,

2001;Trachtenberg et al., 2003b;Kürner et al., 2005;Shaevitz et al., 2005;Wada and Netz,

2009;Sasajima and Miyata, 2021). The helicity might be dominated by an intracellular

structure called the "ribbon," which localizes along the innermost line of the helical cell

structure and is composed of protofilaments. Based on structural studies, ribbons may

switch their helicity through changes in the protofilament length (Trachtenberg and

Gilad, 2001;Kürner et al., 2005;Cohen-Krausz et al., 2011). Ribbons are known to be

composed of fibril proteins specific for *Spiroplasma* species and some *Spiroplasma*

MreB (SMreB) proteins related to MreB that are common in rod-shaped bacteria.

Although fibril filaments are featured by repetitive ring structures, nanometer-order

three-dimensional structure has not been clarified (Trachtenberg and Gilad,

2001;Trachtenberg et al., 2003a;Kürner et al., 2005;Trachtenberg et al., 2008;Cohen-

Krausz et al., 2011;Liu et al., 2017).

In the present study, we isolated the filament of fibrils, the major component protein

of ribbons and clarified its nanometer-order three-dimensional structure using electron

microscopy (EM) and image analyses. The fibril filament has a repetitive structure

featuring a ring and a cylinder with a helical pitch similar to those of the ribbon and cell.

2   Materials and Methods

2.1 Bacterial strains and culture conditions

The type strain, TDA-040725-5[T], of *Spiroplasma eriocheiris* was cultured in R2 medium

(2.5% [wt/vol] heart infusion broth, 8% sucrose, and 10% horse serum) at 30 °C until an

optical density of 0.06 to 0.1 was achieved at 600 nm (Liu et al., 2017;Terahara et al.,

2017).

2.2 Optical microscopy

Cultured cells were centrifuged at 11 000 × g, 10 °C for 10 min and suspended in PBS

consisting of 75 mM sodium phosphate [pH 7.3], 100 mM NaCl containing 20 mM

glucose, and 0.6% methylcellulose, to achieve a cell density 10-fold higher than that of

the original (Liu et al., 2017;Terahara et al., 2017). Cells were inserted into a tunnel

chamber assembled by taping coverslips, as previously described, and observed under an

IX71 microscope (Olympus, Tokyo, Japan) (Uenoyama et al., 2004). A video was

captured using a DMK33UX174 complementary metal–oxide–semiconductor (CMOS)

camera (The Imaging Source, Taipei, Taiwan) and analyzed using ImageJ v1.53a

(https://imagej.nih.gov/ij/).

2.3 Electron microscopy

To observe the intact cells, the cell suspension was placed on a hydrophilized grid, fixed

using 2% glutaraldehyde, washed with water, and stained with 2% uranyl acetate. To

observe the internal structure, the cell suspension on a grid was treated with PBS

containing 0.1 mg/mL DNase and 1 mM $MgCl_2$ for 20 s, washed, and stained with 2%

uranyl acetate. QFDE-EM was performed as previously reported for specimens

suspended in a solution, 10 mM HEPES (pH 7.6), and 150 mM NaCl containing mica

flakes (Tulum et al., 2019). The Triton X-100 treatment was done on glass surface before

freezing, to observe the internal structure. Images were acquired using a JEM1010 EM

(JEOL, Akishima, Japan) equipped with a FastScan-F214(T) charge-coupled device

(CCD) camera (TVIPS, Gauting, Germany) and analyzed using ImageJ v1.53a. For

tomography, images were captured using a Talos F200C EM (FEI, Eindhoven,

Netherlands) equipped with a 4k × 4 K Ceta CMOS camera (FEI). Single-axis tilt series

were collected covering an angular range from -50° to +50° with 1.5° steps and analyzed

using IMOD (ver 4.11) and PEET (ver 1.15.0).

2.4 Isolation of the ribbon and fibril

To isolate the internal structure, 10 mL of cell suspension in PBS was treated with 1%

Triton X-100, 0.1 mg/mL DNase, 1 mM MgCl$_2$, and 0.1 mM PMSF, with shaking for 10

min at 4 °C. The insoluble fraction was recovered via centrifugation at 20 000 × *g* for 30

min at 4 °C, and suspended in PBS to obtain a final volume of 0.2 mL. The sample was

placed at the top of sucrose solution layers of 0%, 20%, 30%, 40%, 50%, and 60%, and

centrifuged at 20 000 × *g* for 20 min at 4 °C in a 1.5 mL tube at a fixed angle. To isolate

the fibril filament, the insoluble fraction was additionally treated with a solution

consisting of 2% cholic acid, 20 mM Tris-Cl pH 8.0, 150 mM NaCl at 4 °C for 8 h, and

subjected to stepwise density gradient centrifugation. SDS-PAGE and peptide mass

fingerprinting were performed as described previously (Nakane and Miyata,

2007;Kawakita et al., 2016;Liu et al., 2017). Band intensities were calculated using

ImageJ, from scanned gel images.

2.5 Preparation of the single-stranded fibril filament

The isolated fibril was adjusted to 1 mg/mL in 20 mM Tris-Cl pH 8.0 and 150 mM NaCl.

The fibril suspension (1 mL in a 1.5 mL test tube) was treated on ice for 5 s using a

sonicator (UR-21P, TOMY, Tokyo, Japan). The condition of the fibril filament was

checked via negative-staining electron microscopy (EM). The processes of sonication and

observation were repeated with the fibril suspension until more than 90% of the filaments

became single-stranded.

2.6 Reconstitution of the 3D structure

The contrast transfer function (CTF) parameters for negative-staining EM images were

estimated using the Gctf25 software (Zhang, 2016). The images of fibril filaments were

selected automatically using RELION 3.0 (Zivanov et al., 2018) as helical objects and

segmented as squares of 200 × 200 pixels with a 90% overlap. These 14,543 images were

2D-classified and 11,867 images were selected for further analyses. *Ab-initio*

reconstitution was performed using cisTEM (Grant et al., 2018) based on segmented

images from 12 classes. The selected 11,867 particle images were 3D-classified using the

3D map in RELION 3.0 (Zivanov et al., 2018).

3　　　Results

3.1 Cell helicity is derived from the internal ribbon structure

To clarify the relationship between the helical cell morphology and the internal ribbon structure, we first measured the helical pitches of the swimming cells using optical microscopy. Under phase-contrast microscopy, the helical shape of the cells can be observed as a series of dense segments in the defocused image plane relative to the cell axis (Fig. 1A). We measured the pitches along the cell axis for the segments of left and right handedness (Fig. 1F). The helical pitches were 709 ± 74 (n = 50) and 718 ± 65 nm (n = 50) for the left- and right-handed segments, respectively.

EM was subsequently employed to analyze the internal ribbon structure and compare the helical pitches of the cells and ribbons. Negative-staining EM revealed images of helical-shaped cells with a narrow tip on one side (Fig. 1B).

The internal ribbon structure was exposed by treating the cells with 0.1% Triton X-100 on the grid (Fig. 1C). The ribbon had a "helical" flat structure. These observations are consistent with those of previous studies (Trachtenberg and Gilad, 2001). Thereafter, the pitches of the cell and the exposed ribbon were measured (Fig. 1F). Generally, the specimens for negative-staining EM are placed in vacuum and dried, which can result in distortions and is disadvantageous for evaluating the helicity. We therefore applied quick-freeze, deep-etch (QFDE) EM to visualize the structure in a state as closely as possible to the original (Heuser, 2011). In QFDE, a sample is frozen in milliseconds and exposed by fracturing and etching. Thereafter, a platinum replica was created by shadowing. The observation of the replica by transmission EM provides images with high contrast and resolution, which is markedly better than that provided by conventional scanning electron microscopy (SEM) (Heuser, 2011;Tulum et al., 2019). Replicas were then prepared by fracturing and platinum coating. QFDE-EM revealed cell morphology consistent with that obtained using negative-staining EM (Fig. 1D). Using QFDE-EM, we observed the ribbons exposed to 0.1% Triton X-100 treatment (Fig. 1E). The ribbon had a structure in which the twisted positions were assembled in a line, showing that the images observed by negative-staining EM were flattened. Interestingly, all ribbons were left-handed (Fig. 1D, E). When the cells were starved in phosphate-buffered saline (PBS) without glucose

for 30 min, they showed a left-handed helix with the same pitch. Therefore, this structure

was assumed to be the default state of the cell, and the ribbon switched to the default

structure during the visualization process. The helical pitches of the cells and ribbons

aligned well with each other, indicating that the ribbon has a critical role for cell helicity

(Fig. 1F, Table 1).

3.2 Isolation and characterization of the ribbon

For further characterization, we isolated the internal ribbon structure. The cell suspension

was treated with 1% Triton X-100 and subjected to stepwise gradient centrifugation with

0%, 20%, 30%, 40%, 50%, and 60% sucrose layers. After centrifugation, we found a

dense layer of cell contents at the bottom of the 40% sucrose layer. The fraction was

recovered and then observed by EM. Based on the observation, the ribbon was found to

comprise protofilaments with a width of 66 ± 12 nm (n=20) and a length longer than 2

μm, which may correspond to the full length of the cell (Fig. 2A). To analyze the number

and width of the protofilaments involved in the isolated ribbon, we traced a cross

sectional image profile of the ribbon (Fig. 2D a). Six to nine protofilaments were

detected, with peak distances ranging between 4 and 16 nm (Fig. 2D b, c), consistent with

the findings of the previous studies (Trachtenberg and Gilad, 2001;Liu et al., 2017).

Ribbon twists are observed as periodic frays in the ribbons. The ribbon pitches were

measured from the frays as 350 ± 17 nm (n = 47) (Fig. 2D d), which is comparable to the

helical pitches of the cells and the ribbons exposed from cells on grids (Fig. 1, Table 1)

($P = 0.7 > 0.01$). SDS-PAGE and peptide mass fingerprinting analyses of this fraction

revealed five protein bands, including six proteins (Fig. 2B, Table 2, Table S1). Band (v)

contains SMreBs 2 and 4. The whole ribbon fraction mainly comprised the fibril protein

(band iii) and a protein mixture of SMreBs 2 and 4 (band v), with an intensity ratio of

47% and 37% of the total protein amount, respectively. Further studies are necessary to

conclude physical interactions of SPE-1201 and FtsH to fibril protein, because these

proteins are abundant in *S eriocheiris* cells (Liu et al., 2017).

We intended to use A22, an inhibitor of MreB polymerization, to examine the role of

SMreBs in ribbon formation (Shi et al., 2018); this is because the binding of A22 to

SMreBs has been suggested from amino acid sequences (Takahashi et al., 2020). First,

the effect of 1 mM A22 on the swimming *Spiroplasma* cells was determined. The cells

lost their original shape and stopped moving within 2 min (Fig. 2C), suggesting that the

functions of SMreBs were inhibited by A22. Thereafter, we isolated the ribbon from cells

maintained in 1 mM A22 for 2.5 h at 30 °C. The ribbons were found to be dispersed (Fig.

2A, c). SDS-PAGE analysis revealed contents of 67% and 11% for fibril (band iii) and

SMreB2 (band v) proteins, respectively (Fig. 2B), suggesting that the protofilaments

comprising fibril proteins are stabilized and modified by SMreBs in the ribbon structure.

3.3 Isolation and helical pitch of the fibril filament

To analyze the detailed structure of fibril filaments, we treated the ribbon fraction with

cholic acid and isolated fibril proteins using sucrose-gradient centrifugation. SDS-PAGE

analysis showed that the fraction only contained fibril protein (Fig. 3A). Negative-

staining EM revealed that the fibril protein formed filaments that included single-,

double-, and more-stranded filaments, suggesting various types of interactions between

the fibril protein molecules (Fig. 3B a). A single-stranded fibril filament consisted of

repeated ring units with approximately 9 nm intervals (Fig. 3B b c), consistent with

previous studies (Townsend et al., 1980;Williamson et al., 1991;Trachtenberg and Gilad,

2001;Liu et al., 2017). The ring units were connected by the backbone cylinder (Fig. 3B

c). The double-stranded fibril filament appeared to be formed via the alignment of two

single-stranded filaments contacting with each other at the ring side not the cylinder side,

resulting in a thickness of 14 nm, double that of the single-stranded filament (7 nm) (Fig.

3B d e). We analyzed the helical pitches for the double-stranded fibril filaments as the

double-stranded fibril filament had a sufficient length of stable helix to cover the pitch,

with a clear twist of the ring pattern along the filament axis. Images of the fibril filament

cropped from the electron micrographs using the straightening selection tool of the

ImageJ software were subjected to Fourier filtering to remove noise (Fig. 3C). However,

the handedness of the fibril filament could not be concluded as the negative-staining EM

images are projections of the object, and the alignment of the filament on the EM grid

was not distinguishable. Therefore, we analyzed the isolated fibril filament using QFDE-

EM (Fig. 3D) as the replica synthesized with platinum covers only one side of the object

surface. The structures shared features with those from the negative-staining EM (Fig.

3D, Fig. S1). We succeeded in determining their handedness (Fig. 3D a–f) and concluded

that the double-stranded fibril filament formed a left-handed helix. The half pitch was

distributed at 351 ± 33 nm (n = 50), which aligns with the results of negative-staining EM

(Fig. 3E). The agreement of helix pitches in the cell, isolated ribbon, and fibril filament

suggests that the fibril filament is a major component of ribbon formation and cell

helicity (Table 1).

3.4 Three-dimensional reconstruction of the fibril filaments

To clarify the fibril filament from a three-dimensional (3D) viewpoint, a single-particle

analysis was performed on negative-staining EM images. The double-stranded fibril

filament was not suitable for image averaging, which might be due to the positional

variation in the binding of the two filaments (Fig. 3 and Fig. S2). Therefore, we sonicated

the purified fibril fraction to increase the proportion of single-stranded filament and

successfully acquired single-stranded images (Fig. 4A). From the selected 11,867

particles with good quality, the 2D-averaged images were classified into three types: (i),

(ii), and (iii) (Fig. 4A b). The initial 3D image was reconstructed using the *ab-initio* 3D

function of cisTEM software (Grant et al., 2018), and used as the reference for the

subsequent 3D classification (Fig. 4A c). 3D structures of the fibril filament reconstructed

from 11,867 particles using RELION 3.0 revealed three different conformations (i.e.,

class 1, left-handed mainly straight (49%); class 2, left-handed with curvature (24%); and

class 3, right-handed with curvature (27%) (Fig. 4A d and Fig. S3). The class 1 structure

reconstituted with rotational symmetry (C2) was not significantly different from that

without symmetry (C1), suggesting that the fibril filament had rotational symmetry

without polarity (Fig. S3). We therefore reconstructed the structures of the fibril filaments

with C2 symmetry. The 2D reprojections from these three structures corresponded well

with the 2D class averages, indicating the validity of the obtained 3D structures (Fig. S3).

The 3D structure of the fibril filament had repeating elliptical rings with a pitch of 8.7 nm

along the filament axis, and the ring size was 11 wide and 6 nm long along the filament.

A short backbone cylinder tilted slightly to the right was found to connect the ring units,

resulting in a positive curvature (Fig. 4A d). These characteristics were common to all

three classes.

Although the superimposition of class 1 and others showed their structural

diffferences, the positions responsible for the differences could not be identified owing to

the low resolution of the structures (Fig. 4B). The fibril filaments of all classes were

twisted along the filament axis, but with different rotational angles (Fig. S4). The twisting

angles were estimated from the angle averages of the first and fourth units, as 5.9 (left-

handed), 7.3 (left-handed), and 9.7 (right-handed) degrees for classes 1, 2, and 3,

respectively. The twisting angles were estimated from the subunit numbers in the double-

stranded images (Fig. 3) for negative-staining and QFDE-EM as 4.9 °and 4.7 °,

respectively. These numbers slightly differed from those obtained from the reconstituted

3D structures, suggesting conformational differences between the curved and straight

filament forms. These structures can explain the peak distance observed in the density

profile of the isolated ribbon (Fig. 2D c, S5).

We proceeded to examine the variation in the ring interval (Fig. 4C, Fig. S6). 2D

averaged images were measured for 60 ring intervals. The intervals were 8.86 ± 0.24 nm

(n=60) and did not show group separation, suggesting that the intervals do not have clear

conformational change, despite some having an elasticity up to 2.7%.

3.5 Handedness verified based on the tomography of the QFDE replica

The 3D images reconstituted from negative-staining EM had common features, despite

variations in curvature and twist. The reconstructed structures all have rings and cylinders

tilted slightly to the right along the filament axis when viewed from the front and back

sides, respectively (Fig. 4A d), indicating that the three classes belong to the same side of

mirror images. As the images by negative-staining EM are projections of the objects, the

reconstituted structures may mirror images of the real structures. Thereafter, we intended

to verify the handedness of the reconstituted structures by EM tomography of the QFDE

replica sample (Fig. 5); this is because the tomogram cannot be a mirror image (Briegel et

al., 2013;Jensen, 2015).

We made QFDE replicas from the fraction containing single-stranded fibril

filaments, acquired images every 1.5° to 50° specimen tilt for both directions,

reconstituted tomograms (Movie_S4) (Fig. 5A), and then obtained a structure by

averaging 60 subtomograms (Fig. 5B). As expected, the resulting filament structure had

rings and cylinders. The rings and cylinders were tilted from the filament axis, rising to

the right from the horizontal axis by 4–5 ° and 74–82 ° when viewed from the front and

back, respectively (Fig. 5C and S7), which align well with the features of structures from

negative-staining EM. These results indicate that the classes of structures from negative-

staining EM had the same handedness as the real structures (Fig. 5C).

4   Discussion

4.1 Structures of the isolated fibril

The unique swimming of *Spiroplasma* is believed to be caused by the ribbon structure

(Kürner et al., 2005;Cohen-Krausz et al., 2011;Harne et al., 2020b;Sasajima and Miyata,

2021). In this study, we isolated filaments of fibrils, the major protein of the ribbon, and

revealed the 3D structure of the single-stranded filament at the nanometer scale using

EM. Fibril filaments have been isolated for a long time, and their EM images show a

characteristic ring repeat structure with a high contrast (Townsend et al.,

1980;Williamson et al., 1991;Trachtenberg and Gilad, 2001;Trachtenberg et al.,

2003a;Cohen-Krausz et al., 2011;Liu et al., 2017). However, a nanometer-order three-

dimensional fibril filament structure is yet to be revealed. Sonication during the isolation

process was effective in isolating the single-stranded filament, whose uniform structure

was advantageous for image averaging (Fig. 3). Negative-staining EM was used to

reconstruct the structure (Fig. 4). However, as this method produces projection images,

the handedness of the reconstructed structure may be incorrect. Therefore, we confirmed

the handedness of the structure by tomographic analysis of platinum replicas prepared by

QFDE-EM (Fig. 5). The final structure was a repeating structure of elliptical rings

connected by backbone cylinders aligned off-axis with a gentle left-handed helix, which

is consistent with that of previous studies. No polarity was observed in the filament

structure.

These results raise questions regarding the alignment of the 512 amino acid residues

of the fibril protein with the structure and the structure formed by the 1-228 amino acid

residues possessing obvious sequence similarity to methylthioadenosine/S-

adenosylhomocysteine (MTA/SAH) nucleosidase (Cohen-Krausz et al., 2011;Parveen

and Cornell, 2011;Sasajima and Miyata, 2021). These questions will be answered via

cryo-EM analysis of the single-stranded fibril filaments prepared in this study.

4.2 Ribbon structure in the cell

When *Spiroplasma* cells were lysed with a detergent, the ribbon structure appeared to run

along the entire length of the cell axis (Fig. 1) (Trachtenberg and Gilad, 2001). In this

study, we isolated ribbons with a length equivalent with the entire length of the cell (Fig.

2). These observations suggest that the ribbon is a relatively stable structure rather than a

highly dynamic one that disappears in a short time. Furthermore, as the extraction

procedure with cholic acid yielded a structure consisting only of fibril filaments (Fig. 3),

the stable properties of the ribbon are likely to be derived from the fibril filament. The

helix of the fibril filament was directly observed in the double-stranded filament (Fig. 3).

The constant helical pitch of a single strand could not be detected, which may be due to

its irregular attachment to the EM grid. The two strands of double-stranded filaments may

stabilize the inherent helical character of the filament by combining them. The

handedness and pitch observed in the duplexes were left-handed and 351±34 (702±68)

nm, respectively, aligning with the helical character of the cells at rest (Fig. 1, 3). As

previous observations revealed the presence of ribbons in the innermost portion of the

cell helix (Kürner et al., 2005;Trachtenberg et al., 2008), the helix of the resting cell

should directly reflect the characteristics of the fibril filament.

During swimming, the cell switches its helical form into a right-handed one with a

helical pitch similar to the left-handed one (Fig. 1). However, we could not find the

corresponding right-handed helical structures in the isolated fibril filaments or ribbons.

Only class 3 3D image reconstructed from 27% of the negative-staining EM images

suggested a right-handed helical structure (Fig. 4); however, a further investigation is

needed to conclude that this structure is stable one as the protein can be distorted by

sticking to the EM grid in this analysis. These observations suggest that the right-handed

helical structure observed in cells during swimming does not originate from another

stable fibril filament structure (Fig. 4). The helix switch can also be explained by

assuming that two types of filaments running parallel to the ribbon are alternately

extended and contracted (Kürner et al., 2005;Cohen-Krausz et al., 2011). To test this

notion, we examined the distribution of the fibril filament lengths and found that the
length distribution had a single peak at 8.86 ± 0.24 nm (Fig. 4). Such finding suggests
that the fibril filament has only one stable length and does not support a helical switch
caused by a length change in the fibril filament.

4.3 Role of fibril in the swimming mechanism

The fibril protein is conserved in most *Spiroplasma* species with high amino acid
sequence similarity (Ku et al., 2014). However, *Spiroplasma sabaudiense* and
Spiroplasma helicoide do not contain fibril proteins, despite exhibiting helicity-switching
swimming (Harne et al., 2020b). Recently, the expression of two SMreB proteins in the
non-swimming synthetic bacterium, syn3.0B, was demonstrated to reproduce cell helicity
and helicity-switching swimming (Hutchison et al., 2016; Kiyama et al., 2021).
Moreover, the expression of SMreB induced cell helicity and its switching in spherical
Mollicutes species (Lartigue et al., 2021), implying that the helix formation of the cell
and the force generation for switching are caused by SMreBs. Then, what is the role of
fibril filaments in most *Spiroplasma* species? Isolated MreB binds to fibril filaments
(Harne et al., 2020a). Further, our results (Fig. 2) support the binding of SMreB to the
fibril filaments. These observations suggest that MreB exerts a force on fibril filaments
for swimming. SMreBs might cause helicity-switching swimming, and fibril filaments
might be effective at obtaining high energy efficiency and chemotaxis; this is supported
by the observation that swimming reconstructed in syn3.0B by SMreBs lacks processivity
(Kiyama et al., 2021).

5   Conflict of Interest

The authors declare that the research was conducted in the absence of any commercial or
financial relationships that could be construed as a potential conflict of interest.

6   Author Contributions

YS and MM designed the experiments. YS performed the experiments. YS, TK, TM,
AK, and KN acquired and analyzed the images. YS and MM wrote the manuscript. All
authors discussed the data.

7 Funding

This study was supported by Grants-in-Aid for Scientific Research A (MEXT KAKENHI, Grant Number JP17H01544), JST CREST (Grant Number JPMJCR19S5), the Osaka City University (OCU) Strategic Research Grant 2017 for top priority research to MM, JSPS KAKENHI (Grant Number JP25000013), the Platform Project for Supporting Drug Discovery and Life Science Research (BINDS) from AMED (Grant Number JP19am0101117 and support number 1282), the Cyclic Innovation for Clinical Empowerment (CiCLE) from AMED (Grant Number JP17pc0101020), and JEOL YOKOGUSHI Research Alliance Laboratories of Osaka University to KN.

8 Acknowledgments

We thank Yuhei O Tahara, Daichi Takahashi, Hana Kiyama, and Ikuko Fujiwara at the Graduate School of Science, Osaka Metropolitan University, Japan, for their helpful discussions.

REFERENCES

Briegel, A., Pilhofer, M., Mastronarde, D.N., and Jensen, G.J. (2013). The challenge of determining handedness in electron tomography and the use of DNA origami gold nanoparticle helices as molecular standards. *J Struct Biol* 183**,** 95-98.

Cohen-Krausz, S., Cabahug, P.C., and Trachtenberg, S. (2011). The monomeric, tetrameric, and fibrillar organization of Fib: the dynamic building block of the bacterial linear motor of *Spiroplasma melliferum* BC3. *J Mol Biol* 410**,** 194-213.

Distelhorst, S.L., Jurkovic, D.A., Shi, J., Jensen, G.J., and Balish, M.F. (2017). The variable internal structure of the *Mycoplasma penetrans* attachment organelle revealed by biochemical and microscopic analyses: implications for attachment organelle mechanism and evolution. *J Bacteriol* 199.

Gasparich, G.E. (2002). *Spiroplasmas*: evolution, adaptation and diversity. *Front Biosci* 7**,** d619-640.

Grant, T., Rohou, A., and Grigorieff, N. (2018). cisTEM, user-friendly software for single-particle image processing. *Elife* 7**,** e35383.

Grosjean, H., Breton, M., Sirand-Pugnet, P., Tardy, F., Thiaucourt, F., Citti, C., Barre, A., Yoshizawa, S., Fourmy, D., De Crecy-Lagard, V., and Blanchard, A. (2014). Predicting the minimal translation apparatus: lessons from the reductive evolution of mollicutes. *PLoS Genet* 10**,** e1004363.

Harne, S., Duret, S., Pande, V., Bapat, M., Beven, L., and Gayathri, P. (2020a). MreB5 Is a determinant of rod-to-helical transition in the cell-wall-less bacterium *Spiroplasma*. *Curr Biol* 30**,** 4753-4762.e4757.

Harne, S., Gayathri, P., and Beven, L. (2020b). Exploring *Spiroplasma* biology: opportunities and challenges. *Front Microbiol* 11**,** 589279.

Harumoto, T., and Lemaitre, B. (2018). Male-killing toxin in a bacterial symbiont of *Drosophila*. *Nature* 557**,** 252-255.

Heuser, J.E. (2011). The origins and evolution of freeze-etch electron microscopy. *J Electron Microsc (Tokyo)* 60 Suppl 1**,** S3-29.

Hutchison, C.A., 3rd, Chuang, R.Y., Noskov, V.N., Assad-Garcia, N., Deerinck, T.J., Ellisman, M.H., Gill, J., Kannan, K., Karas, B.J., Ma, L., Pelletier, J.F., Qi, Z.Q., Richter, R.A., Strychalski, E.A., Sun, L., Suzuki, Y., Tsvetanova, B., Wise, K.S., Smith, H.O., Glass, J.I., Merryman, C., Gibson, D.G., and Venter, J.C. (2016). Design and synthesis of a minimal bacterial genome. *Science* 351**,** aad6253.

Jensen, G.J. (2015). "Getting started in Cryo-EM with professor Grant Jensen".).

Kürner, J., Frangakis, A.S., and Baumeister, W. (2005). Cryo-electron tomography reveals the cytoskeletal structure of *Spiroplasma melliferum*. *Science* 307**,** 436-438.

Kawakita, Y., Kinoshita, M., Furukawa, Y., Tulum, I., Tahara, Y.O., Katayama, E., Namba, K., and Miyata, M. (2016). Structural study of MPN387, an esssential protein for gliding motility of a human-pathogenic bacterium, *Mycoplasma pneumoniae*. *J Bacteriol* 198**,** 2352-2359.

Kiyama, H., Kakizawa, S., Sasajima, Y., Tahara, Y.O., and Miyata, M. (2021). Reconstitution of minimal motility system based on *Spiroplasma* swimming by expressing two bacterial actins in synthetic minimal bacterium. *bioRxiv*.

Ku, C., Lo, W.S., and Kuo, C.H. (2014). Molecular evolution of the actin-like MreB protein
gene family in wall-less bacteria. *Biochem Biophys Res Commun* 446**,** 927-932.
Lartigue, C., Lambert, B., Rideau1, F., Decossas, M., Hillion, M., Douliez, J.-P., Hardouin,
J., Lambert, O., Blanchard, A., and Béven, L. (2021). Tuning spherical cells into
kinking helices in wall-less bacteria. *bioRxiv*.
Liu, P., Zheng, H., Meng, Q., Terahara, N., Gu, W., Wang, S., Zhao, G., Nakane, D., Wang,
W., and Miyata, M. (2017). Chemotaxis without conventional two-component
system, based on cell polarity and aerobic conditions in helicity-switching
swimming of *Spiroplasma eriocheiris*. *Front Microbiol* 8**,** 58.
Miyata, M. (2010). Unique centipede mechanism of *Mycoplasma* gliding. *Annu Rev*
Microbiol 64**,** 519-537.
Miyata, M., and Hamaguchi, T. (2016a). Integrated information and prospects for gliding
mechanism of the pathogenic bacterium *Mycoplasma pneumoniae*. *Front*
Microbiol 7**,** 960.
Miyata, M., and Hamaguchi, T. (2016b). Prospects for the gliding mechanism of
Mycoplasma mobile. *Curr Opin Microbiol* 29**,** 15-21.
Miyata, M., Robinson, R.C., Uyeda, T.Q.P., Fukumori, Y., Fukushima, S.I., Haruta, S.,
Homma, M., Inaba, K., Ito, M., Kaito, C., Kato, K., Kenri, T., Kinosita, Y., Kojima, S.,
Minamino, T., Mori, H., Nakamura, S., Nakane, D., Nakayama, K., Nishiyama, M.,
Shibata, S., Shimabukuro, K., Tamakoshi, M., Taoka, A., Tashiro, Y., Tulum, I.,
Wada, H., and Wakabayashi, K.I. (2020). Tree of motility - A proposed history of
motility systems in the tree of life. *Genes Cells* 25**,** 6-21.
Nakane, D., and Miyata, M. (2007). Cytoskeletal "jellyfish" structure of *Mycoplasma*
mobile. *Proc Natl Acad Sci U S A* 104**,** 19518-19523.
Parveen, N., and Cornell, K.A. (2011). Methylthioadenosine/S-adenosylhomocysteine
nucleosidase, a critical enzyme for bacterial metabolism. *Mol Microbiol* 79**,** 7-20.
Razin, S., and Hayflick, L. (2010). Highlights of mycoplasma research--an historical
perspective. *Biologicals* 38**,** 183-190.
Razin, S., Yogev, D., and Naot, Y. (1998). Molecular biology and pathogenicity of
mycoplasmas. *Microbiol Mol Biol Rev* 62**,** 1094-1156.
Relich, R.F., Friedberg, A.J., and Balish, M.F. (2009). Novel cellular organization in a
gliding mycoplasma, Mycoplasma insons. *J Bacteriol* 191**,** 5312-5314.
Sasajima, Y., and Miyata, M. (2021). Prospects for the mechanism of *Spiroplasma*
swimming. *Front Microbiol* 12**,** 706426.
Shaevitz, J.W., Lee, J.Y., and Fletcher, D.A. (2005). *Spiroplasma* swim by a processive
change in body helicity. *Cell* 122**,** 941-945.
Shi, H., Bratton, B.P., Gitai, Z., and Huang, K.C. (2018). How to build a bacterial cell:
MreB as the foreman of *E. coli* construction. *Cell* 172**,** 1294-1305.
Takahashi, D., Fujiwara, I., and Miyata, M. (2020). Phylogenetic origin and sequence
features of MreB from the wall-less swimming bacteria *Spiroplasma*. *Biochem*
Biophys Res Commun 533**,** 638-644.

Terahara, N., Tulum, I., and Miyata, M. (2017). Transformation of crustacean pathogenic
bacterium *Spiroplasma eriocheiris* and expression of yellow fluorescent protein.
Biochem Biophys Res Commun 487**,** 488-493.
Townsend, R., Archer, D.B., and Plaskitt, K.A. (1980). Purification and preliminary
characterization of *Spiroplasma* fibrils. *J Bacteriol* 142**,** 694-700.
Toyonaga, T., Kato, T., Kawamoto, A., Kodera, N., Hamaguchi, T., Tahara, Y.O., Ando, T.,
Namba, K., and Miyata, M. (2021). Chained structure of dimeric F_1-like ATPase in
Mycoplasma mobile gliding machinery. *mBio***,** e0141421.
Trachtenberg, S., Andrews, S.B., and Leapman, R.D. (2003a). Mass distribution and
spatial organization of the linear bacterial motor of *Spiroplasma citri* R8A2. *J
Bacteriol* 185**,** 1987-1994.
Trachtenberg, S., Dorward, L.M., Speransky, V.V., Jaffe, H., Andrews, S.B., and Leapman,
R.D. (2008). Structure of the cytoskeleton of *Spiroplasma melliferum* BC3 and its
interactions with the cell membrane. *J Mol Biol* 378**,** 778-789.
Trachtenberg, S., and Gilad, R. (2001). A bacterial linear motor: cellular and molecular
organization of the contractile cytoskeleton of the helical bacterium *Spiroplasma
melliferum* BC3. *Mol Microbiol* 41**,** 827-848.
Trachtenberg, S., Gilad, R., and Geffen, N. (2003b). The bacterial linear motor of
Spiroplasma melliferum BC3: from single molecules to swimming cells. *Mol
Microbiol* 47**,** 671-697.
Tulum, I., Tahara, Y.O., and Miyata, M. (2019). Peptidoglycan layer and disruption
processes in *Bacillus subtilis* cells visualized using quick-freeze, deep-etch
electron microscopy. *Microscopy (Oxf)* 68**,** 441-449.
Uenoyama, A., Kusumoto, A., and Miyata, M. (2004). Identification of a 349-kilodalton
protein (Gli349) responsible for cytadherence and glass binding during gliding of
Mycoplasma mobile. *J. Bacteriol.* 186**,** 1537-1545.
Wada, H., and Netz, R.R. (2009). Hydrodynamics of helical-shaped bacterial motility.
Phys Rev E Stat Nonlin Soft Matter Phys 80**,** 021921.
Williamson, D.L., Renaudin, J., and Bove, J.M. (1991). Nucleotide sequence of the
Spiroplasma citri fibril protein gene. *J Bacteriol* 173**,** 4353-4362.
Zhang, K. (2016). Gctf: Real-time CTF determination and correction. *J Struct Biol* 193**,** 1-
12.
Zivanov, J., Nakane, T., Forsberg, B.O., Kimanius, D., Hagen, W.J., Lindahl, E., and
Scheres, S.H. (2018). New tools for automated high-resolution cryo-EM structure
determination in RELION-3. *Elife* 7**,** e42166.

1 Data Availability Statement

The datasets presented in this study can be found in the Supplementary Material.

Table 1. Dimensions of the cell and ribbon

Parameters	Negative-staining EM	QFDE-EM	Optical microscopy
Cell helical pitch	706 ± 74 nm 703 nm	711 ± 41 nm (LH) 711 nm (LH)	709 ± 74 nm (LH) 702 nm (LH) 718 ± 65 nm (RH) 711 nm (RH)
Ribbon helical pitch	691 ± 53 nm 700 nm	700 ± 60 nm (LH) 706 nm (LH)	
Isolated ribbon 1/2 helical pitch	350 ± 17 nm 352 nm		
Isolated fibril 1/2 helical pitch	341 ± 27 nm 335 nm	351 ± 34 nm (LH) 352 nm (LH)	

536   Handedness is represented by LH and RH.

537   The upper and lower rows show the mean, standard deviation, and median, respectively.

540   **Table 2.** Protein components of the ribbon isolated from original cells[1].

Protein Band[1]	Gene ID	Annotation	Mascot Score[2]	Mass (kDa)[3]	Density ratio (%)	
					Original	A22 treated
(i)	SPE-1201	Hypothetical protein	72	85.8	4	5
(ii)	SPE-0013	FtsH	84	77.0	12	17
(iii)	SPE-0666	Fibril	206	58.7	47	67
(iv)	SPE-1231	SMreB5	98	38.5	10	7
(v)	SPE-1224	SMreB2	80	37.8	27	4
	SPE-1230	SMreB4		40.7		

[1] From A22-treated cells, the proteins common to the original cells were identified for bands (i)–(iv). For band (v) only,

SMreB2 was identified.

[2] Mascot Score is the logarithm of probability that the observed match is a random event.

[3] Calculated from the amino acid sequence as a monoisotopic molecule.

Figure Legends

Figure 1. Helicity of the cell and ribbon structure. **(A)** Phase-contrast microscopy of

swimming cell. The blue and red segments, and broken line indicate the left- and right-

handed helicity, and cell axes, respectively. The pitch was measured as indicated by a

double headed arrow. **(B, C)** Cell and ribbon images acquired by negative-staining EM.

(D, E) Cell and ribbon images acquired by QFDE EM. (F) Helical pitches of cells and

ribbon measured by optical microscopy, negative-staining EM, and QFDE-EM.

Handedness was judged by optical microscopy and QFDE-EM. All cells analyzed by

QFDE-EM were left-handed as they were grown under a starved condition.

Figure 2. Isolation and characterization of the ribbon. **(A)** Isolated ribbon structure. (a)

The whole structure of the isolated ribbon with helicity as shown by periodical wide

positions (marked by arrows). (b) The magnified image of the isolated ribbon and the

helical pitch is indicated by a bidirectional arrow. (c) Ribbon fraction isolated from cells

treated with A22. **(B)** Protein profiles of the ribbon fraction isolated from cells untreated

and treated with A22. **(C)** Cell images before (a) and after (b) treatment with 1 mM A22

for 2 min. **(D)** Numerical characterization. (a) Sectional image profile of the area boxed

in panel (A b). The peaks correspond to the center of the protofilament. (b) Histogram for

the number of protofilaments involved in a ribbon. (c) Histogram for the protofilament

width in ribbons. The distribution can be fitted by two Gaussian curves marked (i) and

(ii), with peaks around 7.0 and 10.5 nm, respectively. (d) Histogram for the helical

pitches of the isolated ribbon, fitted by a Gaussian curve with a peak at 351 ± 16 nm

(n=47).

Figure 3. Structures of the isolated fibril filament. **(A)** Protein profiles of the fractions in

the purification process for fibril protein. (i) Whole cell lysate. (ii) Supernatant. (iii)

Isolated ribbon. The sample amount was adjusted to be delivered from the same cell

number. **(B)** Purified fibril filaments observed by negative-staining EM. (a) Field image.

White and black arrows indicate typical single and double strands, respectively. (b, c)

Front and side views of the single-stranded fibril filament. (d, e) Front and side views of

the double-stranded fibril filament. The ring intervals marked by an asterisk were 9 nm

for both single and double strands. **(C)** Double-stranded filaments reconstituted through

Fourier filtering. **(D)** Fibril filaments observed by QFDE-EM. Field (a) and single-

stranded filaments (b, c, d) are presented. The back (b), front (c), and side (d) views are

shown. (e) Single-stranded filament. (f, g) Double-stranded filament image (f) was

reduced for noise through Fourier filtering. (g). The helical pitch was measured as

depicted by a double headed broken arrow. The handedness was clearly observed at the

points marked by white arrows in the panel (a) and panel (f) image.

Figure 4. Three-dimensional reconstruction of the fibril filaments. **(A)** Workflow of

single particle analysis by negative-staining EM. (a) Field images of single-stranded fibril

filaments prepared by sonication. (b) Eight averaged images obtained by a function of 2D

classification in RELION software. (c) The initial 3D model generated by a function of

ab-initio reconstruction in cisTEM software. (d) Three different conformations of the

fibril filament reconstituted by a function of 3D refinement in RELION software. **(B)**

Superpose of class 1 (left-handed) and class 2 (left-handed) and 3 (right-handed)

structures. The fitting reference is indicated by a dashed box. **(C)** Distribution of the ring

intervals. Left: Ring intervals in an averaged image with complete rings. Right: Plotted

ring intervals.

Figure 5. Comparison of the 3D structures of the fibril filament reconstructed from

QFDE and negative-staining EM. **(A)** Replica image of mainly single-stranded fibril

filaments. Left: A field image is shown from a tilt series (Movie_S2). Right lower:

Magnified images of single fibril filament are shown as a raw image (a), a slice from the

tomogram (b), and a subtomogram (c). **(B)** Structure averaged from 60 subtomograms.

The leftmost image is presented under different thresholds from other three images. **(C)**

Superpose of the 3D structures from single-particle analysis (grey) and subtomogram

averaging (magenta). Long axes of ring and cylinder are depicted by broken red arrows.

The filament axes were detected by a function "relion_align_symmetry --sym d2" in

RELION-3.0.

Fig. 1

A

0 s | 0.03 s | 0.07 s | 0.10 s | 0.13 s

Kink

1 µm

B 1 µm | 200 nm

C 200 nm

Cell | Ribbon

D 1 µm | 200 nm

E 200 nm

Cell | Ribbon

F

Helical pitches (nm)

Opt_LH | Opt_RH | NS_Cell | NS_Ribbon | QF_Cell | QF_Ribbon

Fig. 2

Fig. 3

Fig. 4

A

(a)

100 nm

CTF estimation (Gctf_v1.06)

Particle picking (RELION-3.0)

Reference-free 2D classification (RELION-3.0)

(b)

(i) (ii) (ii) (iii) (iii)

(iii) (iii) (iii) (i) (iii)

20 nm

(c)

90°

10 nm

Initial model reconstruction Ab-initio 3D (cisTEM)

3D classification (RELION-3.0)

3D refinement, C2 (RELION-3.0)

11,867 particles

(d)

49%   24%   27%

Back   Front   Side

Ring   11 nm   8.7 nm

180°   45°   45°   Positive curvature

6 nm

Cylinder

10 nm

Class 1   Class 2   Class 3

B

Front   **Side**

Fitting ref.

45°   45°

■ Class 1   ■ Class 2

Fitting ref.

45°   45°

■ Class 1   ■ Class 3

10 nm

C

9.5

9.0

8.5

Ring intervals (nm)

Fig. 5

A

Step 1.5° - 50° ⟷ + 50°

Tilt series (Talos F200C)
Tomography software (FEI)

Generate tomogram (IMOD)

(a) 10 nm

(b)

(c) Front

100 nm

B

Front — Threshold: 190

Front — Threshold: 270 — Ring

Side — 90° — Cylinder

Back — 9 nm — 90° — Cylinder

10 nm

Subtomogram averaging 60 particles (PEET)

Superposing to SPA EM map

C

Tilt of the "ring" part

Tilt of the "cylinder" part

180°

10 nm