Evolution of thermal physiology alters the projected range of threespine

stickleback under climate change

Sara J.S. Wuitchik[1,2,3†,*], Stephanie Mogensen[1], Tegan N. Barry[1], Antoine

Paccard[4,‡], Heather A. Jamniczky[5], Rowan D.H. Barrett[4,§], and Sean M. Rogers[1,6,§]

[1]Department of Biological Sciences, University of Calgary, 2500 University Dr NW, Calgary,

AB, T2N 1N4, CANADA

[2]Informatics Group, Harvard University, 52 Oxford St, Cambridge, MA, 02138, USA

[3]Department of Biology, Boston University, 5 Cummington Mall, Boston, MA, 02215, USA

[4]Redpath Museum and Department of Biology, McGill University, 845 Sherbrooke St W,

Montreal, QC, H3A 0G4, CANADA

[5]Department of Cell Biology & Anatomy, Cumming School of Medicine, University of Calgary,

3330 Hospital Dr NW, Calgary, T2N 4N1, CANADA

[6]Bamfield Marine Sciences Centre, 100 Pachena Rd, Bamfield, BC, V0R 1B0, CANADA

[§]equal co-senior authors

Running title: Evolution alters projected species ranges

*Corresponding author: Sara J.S. Wuitchik

E: sjswuit@g.harvard.edu

T: 857-292-9977

[†]Present affiliations: Department of Biology, University of Victoria, 3800 Finnerty Rd, Victoria, BC, V8P 5C2, Canada & School of Environmental Science, Simon Fraser University, 8888 University Dr, Burnaby, BC, V5A 1S6, Canada
[‡]Present affiliation: McGill University Genome Center, 740 Dr Penfield Avenue, Montreal, QC, H3A 1A5, Canada

Abstract

Species distribution models (SDMs) are widely used to predict range shifts but could be

unreliable under climate change scenarios because they do not account for evolution. The

thermal physiology of a species is a key determinant of range and thus incorporating thermal trait

evolution into SDMs might be expected to alter projected ranges. We identified a genetic basis

for physiological and behavioural traits that evolve in response to temperature change in natural

populations of threespine stickleback (*Gasterosteus aculeatus*). Using these data, we created

geographic range projections using a mechanistic niche area approach under two climate change

scenarios. Under both scenarios, trait data was either static ('no evolution' models), allowed to

evolve at observed evolutionary rates ('evolution' models), or allowed to evolve at a rate of

evolution scaled by the trait variance that is explained by quantitative trait loci (QTL; 'scaled

evolution' models). We show that incorporating these traits and their evolution substantially

altered the projected ranges for a widespread panmictic marine population, with over 7-fold

increases in area under climate change projections when traits are allowed to evolve. Evolution-

informed SDMs should improve the precision of forecasting range dynamics under climate

change, and aid in their application to management and the protection of biodiversity.

Keywords: thermal biology, critical thermal limits, evolution, species distribution models,

climate change, threespine stickleback

Introduction

Temperature is a powerful driver of global biogeography and species distributions frequently reflect temperature gradients in both aquatic and terrestrial habitats (Hochachka & Somero, 2002). Many species adopt thermal strategies (such as thermoregulation or acclimation) that determine their thermal niche (Coutant, 1987; Huey & Kingsolver, 1989; Huey & Slatkin, 1976) and thermal traits can provide a target for directional selection if the environment changes to include temperatures outside the range encompassed by the thermal niche. Adaptation can thus permit species to persist at temperatures that would have previously led to extirpation (Hoffman & Sgrò, 2011; Sexton et al., 2009). Under moderate climate change scenarios, mean global oceanic temperature is projected to increase in excess of 2 °C by the end of the century (IPCC, 2014), with more extreme changes predicted in localized regions (Eyer et al., 2019; Walther et al., 2002) for both cooling and warming events (Hu et al., 2018). Accurately predicting species range patterns under climate change therefore requires data for temperature-associated adaptive trait evolution (Bay et al., 2017; Bush et al., 2016; Catullo et al., 2015).

The genetic basis of thermal traits has been explored in many terrestrial ectotherms (Rolandi et al., 2018; Angilletta et al., 2002; Bowler & Terblanche, 2008; Dillon et al., 2009) and there have been integrative frameworks proposed for how best to use this information to assess the vulnerability of organisms to environmental changes (Araújo et al., 2013; Bay et al., 2017; Huey et al., 2012). Given the differences in thermal biology between terrestrial and aquatic ectotherms (Sunday et al., 2011) and differential warming between terrestrial and aquatic habitats (IPCC, 2014, 2018), predicting how the thermal biology of aquatic organisms will respond to environmental changes requires an understanding of the genetic architecture underlying ecologically relevant thermal traits (Healy et al., 2018) in aquatic systems. The genetic basis of

thermal traits has been explored in a number of socioeconomically and culturally important fish

species (Everett & Seeb, 2014; Jackson et al., 1998; Jin et al., 2017; Larson et al., 2016; Muñoz

et al., 2014; Perry et al., 2001) and there has been an extensive body of work on the thermal

biology of the common killifish (*Fundulus heteroclitus*, Bryant et al., 2018; Fangue et al., 2006,

2009; Healy et al., 2018). These studies suggest that a significant proportion of phenotypic

variance in thermal traits can be explained by genetic variation (*e.g.*, 35 SNPs explained 51.9%

of the variation for upper thermal tolerance in killifish, Healy et al., 2018), highlighting the

potential for heritable responses to changes in thermal conditions.

72   While the climate and ocean temperatures are warming overall, thermal events

characterized by both cold and heat extremes are occurring with increasing frequency (Hu et al.,

2018; IPCC, 2014, 2018; Stott, 2016). Regionally downscaled environmental changes can be

highly spatially heterogenous relative to global temperature trends, and are likely to be the most

relevant for species-specific responses in the context of contemporary climate change (Walther et

al., 2002). For example, range expansions of marine species have been mediated by both ocean-

scale warming and regional cooling (Zeidberg & Robison, 2007). In such cases, population

persistence and adaptation will rely on both upper and lower critical thermal limits, forcing

organisms to respond independently to extreme heat and extreme cold in separate events

(Herring et al., 2015, 2018, 2020; Hu et al., 2018; Walther et al., 2002). Importantly, adaptation

in one thermal trait can shift the thresholds of correlated thermal traits (Buckley & Huey, 2016;

Denny & Dowd, 2012; Hoffman & Sgrò, 2011; Huey & Kingsolver, 2011). Associations

between traits influence the rate of trait adaptation and thus correlated traits must be considered

when attempting to predict future species distributions under climate change (Bestion et al.,

2015; Hoffman & Sgrò, 2011; Valladares et al., 2014). Recently there have been steps to

incorporate theoretical trait evolution into SDMs (*e.g.,* by including the breeder's equation for a

key phenotypic trait, Bush et al., 2016; Catullo et al., 2015; Kearney & Porter, 2009). However,

no model has used empirical estimates of evolutionary rates to inform projected species ranges

under climate change scenarios.

Threespine stickleback (*Gasterosteus aculeatus*, Fig. 1a) are a useful vertebrate species

for understanding the impact of adaptation on range dynamics under climate change. This

species exhibits widespread phenotypic variation (Hendry et al., 2013), there are publicly

available genomic resources (Jones, Grabherr, et al., 2012), and there is prior research on

temperature-associated evolutionary rates for this species (Barrett et al., 2011; Morris et al.,

2018). It is also feasible to artificially breed multiple hybrid generations in common garden lab

environments. Moreover, there is a regionally downscaled climate model for the Pacific

Northwest (Alexander et al., 2018; IPCC, 2014; Reynolds et al., 2007), which makes panmictic

marine populations of *G. aculeatus* in the eastern Pacific Ocean (Kirch et al., 2021; Morris et al.,

2018) an ideal system for testing how adaptive trait variation could affect projections of species

range under climate change.

The objective of this study is to incorporate ecologically and evolutionarily relevant data

on thermal traits into end-of-century projections of species distribution models. To accomplish

this, we assessed physiological and behavioural temperature-associated traits in two marine and

two freshwater populations of threespine stickleback, as well as within-population F1 families

and marine-freshwater hybrid F1 and F2 families. We used the hybrid families to construct

genetic linkage maps and identified multiple quantitative trait loci associated with thermal

preference, as well as upper and lower critical thermal limits. Using these genetically-based traits

and empirically derived evolutionary rate data for this species (Barrett et al., 2011), as well as

others (Morgan et al., 2020; Sanderson et al., 2021), we constructed species distribution models

using a mechanistic niche area approach that integrated the trait data reported in this study and

allowed those traits to evolve under two climate change scenarios, three distinct types of

evolutionary models, and three evolutionary rates.

Materials and Methods

Sample collection and husbandry

116   We collected adult *Gasterosteus aculeatus* from two marine populations (Bamfield, M1,

48°49'12.69"N 125° 8'57.90"W; Garden Bay Lagoon, M2, 49°37'52.84"N 124° 1'49.26"W) and

two freshwater populations (Hotel Lake, FW1, 49°38'26.94"N 124° 3'0.69"W; Klein Lake, FW2,

49°43'32.47"N 123°58'7.83"W) in southwestern British Columbia (Fig. 1b) at less than 0.5 m

depths. To ensure that any genetic variation identified would be relevant for adaptation within

this region, we confined our study to a single genetically panmictic cluster of populations

(Morris et al., 2018). Individuals were maintained in a flow-through system and under a

photoperiod that mimicked the natural source population environment during collection periods

before transport. We transported the fish to the Life and Environmental Sciences Animal

Resources Centre at the University of Calgary, where we separated the fish into population-

specific 113 L glass aquaria at a density of approximately 20 fish per aquarium. We acclimated

marine individuals to freshwater salinity over one week and maintained fish in a common

environment (salinity of 4-6 ppt, water temperature of 15 ± 2 °C, and a photoperiod of 16L:8D).

Individuals were allowed to acclimate for at least 2 weeks before experiments (1 week for stress

reduction post-transfer, 1 week for common garden environment acclimation and salinity ramp).

Each common garden aquarium was on a closed system with individual filters, air stones, and

water supply. We fed all adult fish *ad libitum* once per day with thawed bloodworms (Hikari

Bio-Pure Frozen Bloodworms). All collections and transfers were approved by the Department

of Fisheries and Oceans (marine collections and provincial transfers), the Ministry of Forests,

Lands, and Natural Resource Operations (freshwater collections), and the Huu-ay-aht First

Nations (marine collections).

Crossing design for marine and freshwater F1 families

We collected eggs from females and fertilized the eggs with extracted testes from

euthanized males. We transferred the fertilized egg mass to a mesh-bottomed egg incubator

suspended in a 37 L aquarium for hatching. Each hatching aquarium was maintained with a

single air stone and a filter. Once hatched, we reared the larval fish in 37 L hatching aquaria until

they reached a total length (TL) of approximately 1 cm, after which we split the families into

family-specific 113 L aquaria to maintain suitable densities. We fed the larval fish *ad libitum*

twice daily with live *Artemia spp.* nauplii, and then gradually transitioned the diet to chopped,

thawed bloodworms (Hikari Bio-Pure Frozen Bloodworms) *ad libitum* once daily as they

reached approximately 2 cm TL. The F1 families were maintained in a common garden

environment identical to that of the F0 populations. We produced one F1 family for each

population (M1_F1, M2_F1, FW1_F1, and FW2_F1).

Crossing design for hybrid mapping families

To generate genetically heterogeneous marine-freshwater F1 families from wild F0

parents, we collected eggs from marine females and fertilized the eggs with extracted testes from

euthanized freshwater males. Egg masses were hatched, and juveniles were reared, as detailed

above. We produced one F1 family of M1xFW1 hybrids (hereafter referred to as H1_F1) and

three F1 families of M1xFW2 hybrids (hereafter referred to as H2_F1). The hybrid F1 families

were maintained in a common garden environment identical to that of the F0 populations. To

generate F2 families for linkage map construction and quantitative trait loci (QTL) mapping, we

crossed siblings from the same F1 family using the same methodology that was used to generate

the F1 families. The amount of trait variation explained by these hybrid crosses is likely to be

higher than would be observed in marine-marine or freshwater-freshwater crosses, increasing our

likelihood of detecting QTL. Overall, we produced one F2 family of H1xH1 hybrids (referred to

as H1_F2) and three families of H2xH2 hybrids (referred to as H2_F2_1, H2_F2_2, and

H2_F2_3). All F2 individuals were raised as described above in a common garden environment

identical to that of the F0 and F1 individuals to ensure consistent thermal history.

Thermal tolerance and preference experiments

To assess the lower and upper limits of physiological thermal tolerance, we conducted

standard critical thermal minimum (CTmin) and maximum (CTmax) experiments on adult fish

(Barrett et al., 2011; Fangue et al., 2006; Hutchison, 1961). At these sublethal limits, the fish

experience a loss of equilibrium (LOE) at which they lose the ability to escape conditions that

would ultimately lead to their death (Beitinger et al., 2000). Before each experiment, individuals

were fasted for 24 hours. Our experimental tank held 1000 mL glass beakers aerated individually

to prevent thermal stratification. After a 15-minute acclimation to the experimental apparatus in

the individual beakers, we cooled or heated the water (for CTmin or CTmax, respectively) at a

rate of approximately 0.33 °Cmin^{-1}. We assessed wild F0 individuals ($n_{M1} = 32$, $n_{M2} = 14$, $n_{FW1} =$

15, $n_{FW2} = 16$, $N = 77$; Fig. S2), lab raised F1 ($n_{M1\_F1} = 13$, $n_{M2\_F1} = 15$, $n_{FW1\_F1} = 15$, $n_{FW2\_F1} =$

15, $N = 58$; Fig. S2), and F2 individuals ($n_{H1\_F2} = 28$, $n_{H2\_F2\_1} = 36$, $n_{H2\_F2\_2} = 21$, $n_{H2\_F2\_3} = 17$,

$N = 102$; Fig. S3). All individuals were assessed for CTmin, allowed to recover for at least three

days, then assessed for CTmax to keep thermal stress history consistent. The onset of erratic

behaviours associated with a behavioural stress response (i.e., 'agitation windows', Turko et al.,

2020) occurred below 5.0 °C and above 25.0 °C during CTmin and CTmax experiments,

respectively. Normal behaviour was observed between 5.0 °C and 25.0 °C, whereas outside of

those temperatures, individuals gradually exhibited more extreme stress responses (*e.g.,*

increased gilling rate, erratic movement, muscle spasms, listing, as outlined by the Canadian

Council of Animal Care guidelines) until reaching LOE and the inability of an individual to right

itself (the experimental endpoint, measured in 0.5 °C increments, Barrett et al., 2011; Fangue et

al., 2006; Hutchison, 1961).

186   To assess the range of behavioural thermoregulation, we conducted thermal preference

experiments in a temperature Shuttlebox (Loligo Systems, Viborg, Denmark). Experimental

pools were set with a static gradient of 10 °C in the cool side and 20 °C in the warm side,

connected with a water bridge at 15 °C (acclimation temperature). Temperature was continually

monitored and recorded by immersed temperature probes, connected to a computer-driven

temperature controller and data acquisition system (DAQ-M, Loligo Systems, Viborg,

Denmark). Movement in the Shuttlebox was tracked by an infrared-sensitive uEye USB 2.0

camera (IDS Imaging Development Systems GmbH, Obersulm, Germany) which connected to

Shuttlesoft (v2, Loligo Systems, Viborg, Denmark) and allowed monitoring of fish movement.

Following recovery from CTmax experiments (at least three days), individuals were acclimated

in the Shuttlebox for 15 minutes, with entry at 'cool' or 'warm' pool randomized, to allow for

exploration of the gradient. After acclimation to the experimental apparatus, the movement of

each individual was tracked in 1 second intervals for 30 minutes, recording the preferred ambient

water temperature (thermal preference). Individuals were assessed individually in the Shuttlebox

apparatus to avoid any confounding social behaviours (*e.g.,* as seen in shoaling behaviours,

Cooper et al., 2018). At the time of data collection for thermal trait experiments, all individuals

were adults.

Relationships between all thermal traits separated by population and generation were

assessed using the Pearson product-moment correlation with the *corrplot* (v0.84, Wei et al.,

2017) and *Hmisc* (v3.14-0, Harrell, 2014) packages in R (R Core Team, 2021).

Isolation and characterization of single nucleotide polymorphisms (SNPs)

Genomic DNA was extracted from caudal fin tissue using a phenol-chloroform-based

protocol. We digested tissues overnight in digestion buffer and proteinase K at 55 °C, then

performed multiple phenol-chloroform and ethanol washes to isolate the DNA. We assessed the

quantity of the extracted DNA using the Quant-iT PicoGreen dsDNA assay kit (ThermoFisher

Scientific, Waltham, MA, USA) and Synergy HT plate reader with the Gen5 associated software

(BioTek, Winooski, VT, USA). We prepared restriction site-associated DNA (RAD) libraries

(Peterson et al., 2012) using *MluCl* and *NlaIII* restriction enzymes (New England Biolabs,

Ipswich, MA, USA), ligation of individual barcodes, and pooling of 48 individuals per library at

equimolar concentrations. We performed a final PCR to amplify DNA and add library-specific

indices to allow for pooling of multiple libraries. We sequenced three libraries at the McGill

University and Génome Québec Innovation Center on one lane of Illumina HiSeq 4000 (Illumina

Inc., San Diego, CA, USA).

Assembly of the genetic linkage maps

After barcode demultiplexing and filtering out low quality reads in STACKS (Catchen et

al., 2013), we removed PCR duplicates from the raw sequences and aligned these cleaned

sequences to the *G. aculeatus* reference genome (Jones, Grabherr, et al., 2012) using the

Burrows-Wheeler transform (Li & Durbin, 2010). Individual libraries were concatenated and

filtered (Puritz et al., 2014) using *vcftools* (v3.0 , Danecek et al., 2011) and split into

chromosome-specific variant call format (VCF) files using *SnpSift* (Cingolani et al., 2021) to

assemble the linkage maps chromosome by chromosome. We assigned markers to a linkage

group with an initial logarithm of the odds (LOD) score of 3 after filtering out markers that

showed high levels of segregation distortion and missing observations (> 20% missing data) in

Lep-MAP3 (Rastas, 2017). Unassigned markers were subsequently added to the existing linkage

group at a LOD score of 3 and a minimum size limit of 5 markers per linkage group. We ordered

the markers using a minimum posterior value of 0.001 and collapsed multiple markers when the

probability difference between markers was < 0.01 (Rastas, 2017). The final linkage map was

phased in Lep-MAP3 (Rastas, 2017) and subset for use in R (R Core Team, 2021) with to

generate a list of informative SNPs to use in subsequent analyses with the *qtl* (v1.44-9, Broman

et al., 2003) package. The final linkage maps were similar across families in length and spacing

between markers, though the H2_F2 map had a higher density of markers (Table S1).

Quantitative trait loci (QTL) mapping

We analysed the four F2 families separately with the same methodology to assess the

presence of QTL associated with the thermal traits and avoid the confounding effects of unique

meiotic events. We calculated conditional genotype probabilities using a hidden Markov model,

allowing for possible genotyping errors at a level of 0.0001 using a Kosambi mapping function

with a fixed step width, prior to running genome scans with a single QTL model (Arends et al.,

2014; Broman & Sen, 2009). We determined the LOD score significance thresholds for each trait

through permutation tests for each family (5,000 permutations per chromosome). We pulled

significant QTL above the genome-wide significance threshold ($\alpha = 0.05$, Greenwood et al.,

2011), calculated confidence intervals of QTL location based on nearby markers, and estimated

the percent variance explained by each QTL peak marker ($PVE = 1 - 10^{\wedge}((-2*LOD)/n)$). We

identified two QTL associated with CTmin, two QTL associated with CTmax, and three QTL

associated with thermal preference (Table 1, Fig. 2).

Environmental variables used to construct species distribution models (SDMs)

We compiled environmental data widely used in the construction of SDMs to estimate

suitable habitat in both present day and end-of-century forecasts (Wiens et al., 2009), including

bathymetry, sea ice extent and concentration, salinity, and sea surface temperature. We used

2014 data as our baseline year to match the forecasting baseline of the Fifth Assessment Report

(IPCC, 2014). We assumed a suitable habitat range for this species in the Pacific Northwest to

consist of coastal areas (water depth less than 200 m) where sea ice is never present (*i.e.,* no sea

ice at the maximum extent). *G. aculeatus* has a broad salinity tolerance (Bayly, 2003; Divino et

al., 2016) and salinity is not limiting in any part of the range (Zweng et al., 2013), therefore

salinity was not included in the final present day or forecasted models. We obtained bathymetry

data from the General Bathymetric Chart of the Oceans (GEBCO) of the British Oceanographic

Data Centre (Weatherall et al., 2015), and maximum sea ice extent data from the Multisensory

Analyzed Sea Ice Extent – Northern Hemisphere (MASIE-NH) product (Fetterer et al., 2010).

We obtained maximum and minimum daily statistical mean sea surface temperatures (SST) from

Reynolds et al. (2007) to create rasters of the mean upper quartile, lower quartile, absolute

minimum, absolute maximum, and median temperatures for the baseline year (2014). We found

the minimum, maximum, and median temperature for each 1 km^2 grid cell across the range, then

reduced the area based on the constraints of sea ice extent and bathymetry. Stickleback thermal

trait data were used to set the limits of the ranges within the possible area delineated by sea ice

free water of a suitable depth. Thermal trait measurements were based on our experimental

findings reported in the present study.

In the end-of-century forecast for suitable habitat, we assumed bathymetry to be

consistent with the modern scenario. In contrast, because the Arctic Ocean is projected to be

predominantly free of sea ice in the summer by the end of the century (Johannessen et al., 2004),

with significant end-of-century reductions in winter/spring sea ice concentration (reduced to a

concentration of 0.1 at the Seward Peninsula, Johannessen et al., 2004), we conservatively set the

maximum northern extent of the suitable habitat to the western tip of the Seward Peninsula

(65°35' N). The water temperatures were increased based on projections for large marine

ecosystems of Northern Oceans from global climate models (Alexander et al., 2018; IPCC,

2014). Raster and maps were created in R (v4.1.2, R Core Team, 2021) using the packages *raster*

(v3.5.15, Hijmans et al., 2020) and *rgeos* (v0.5.9, Roger et al., 2020).

Trait inclusion in SDMs

We incorporated experimental data on the wild marine populations (Fig. 1c) to

understand how inclusion of trait data may affect range projections under climate change. These

trait-defined areas were overlain on the suitable habitat background to delineate projected

presence based on thermal traits in both current day and IPCC-projected Representative

Concentration Pathways (RCPs) 4.5 and 8.5. The trait values were kept constant from current

day to end-of-century (*i.e.,* not changed) in the 'no evolution' projections. In contrast, in the

'evolution' projections, we allowed CTmin and CTmax to change based on empirically derived

evolutionary rate estimates. We allowed CTmin to evolve at a rate of 0.63 *haldanes* estimated

from a selection experiment previously conducted on populations belonging to the same admixed

genetic cluster as the sampled populations (Barrett et al., 2011; Morris et al., 2018). At present,

there are no empirical estimates of CTmax evolution for threespine stickleback. Thus, as a proxy

we used a known estimate of CTmax evolution in fish, which was observed in zebrafish (0.19

haldanes; Morgan et al., 2020). Given that the evolutionary rates for CTmin and CTmax from

Barrett et al. (2011) and Morgan et al. (2020) are quite high relative to other recorded

evolutionary rates for phenotypic traits (particularly physiological traits; Sanderson et al., 2021),

we also explored the influence of evolution on projected ranges when evolution occurred at more

'typical' rates. To do so, we allowed both CTmin and CTmax to evolve at a mean rate from a

large meta-analysis of phenotypic trait evolution (0.14 *haldanes*; Sanderson et al. 2021).

Since the rates of evolution used here are estimates based on whole-organism tolerance

(Barrett et al., 2011; Morgan et al., 2020), and trait evolution is constrained by the underlying

genetic architecture of the trait (Barghi et al., 2020; Fisher, 1930; Orr, 1998; Ungerer &

Riesebero, 2003), we next explored how selection acting on discrete, identifiable loci could

impact evolutionary rates, and the subsequent effects on projected habitat range. To do so, we

constructed models that scaled the evolutionary rate of CTmin or CTmax based on the observed

genetic architecture from our QTL mapping (*i.e.*, we scaled by the percent of trait variation that

is explained by each locus; 'scaled evolution' projections). However, an important restriction in

the evolution of CTmin for all models incorporating trait evolution ('evolution' and 'scaled

evolution' projections) was a hard boundary at 0 °C, with the assumption that population

persistence in a sub-zero environment would require additional adaptations to CTmin

improvement (*e.g.,* extreme adaptations observed in Antarctic notothenioid fishes, Cheng &

Detrich, 2007; Detrich et al., 2000; Shin et al., 2014). Similarly, we defined a hard boundary for

CTmax evolution at 42 °C (Morgan et al., 2018), with the assumption that the strength of

selection for CTmax at this rate would result in decreased phenotypic variation and acclimatory

ability leading to an upper thermal limit plateau (Gilchrist & Huey, 1999; Hangartner &

Hoffmann, 2016; Morgan et al., 2020).

To quantify the differences in estimated suitable habitat under current day and end-of-

century conditions, we compared areas for each warming scenario to the equivalent scenario

under current conditions. Similarly, to compare the differences in evolutionary scenarios, the

area of each end-of-century evolutionary trajectory was compared to either the contrasting RCP

projection or scaled evolution projection. For these comparisons, we used North Pole Lambert

azimuthal equal area projection for all maps, and georeferenced to known landmarks in ArcGIS

v10.8 (Environmental Systems Research Institute, 2017) to calculate area from the maps

generated in R (v.4.1.2, R Core Team, 2021; conversion ratio of 7873.42).

Results

Genetic basis of thermal traits

Stickleback exhibited a wide thermal tolerance range bounded by a mean CTmin of 1.89

°C (+/- 1.04 °C SD) and a mean CTmax of 30.1 °C (+/- 1.78 °C SD)(Fig. 1c, S2), with these

physiological traits being highly correlated ($r = 0.79$, Fig. S1). These marine populations also

tolerated a wide range of temperatures within which there was no observable stress response (5.0

– 25.0 °C). Thermal preference values were slightly above the acclimation temperature, but with

more variance around the mean than is observed in the physiological traits (16.7 +/- 3.19 °C,

Figs. 1c, S2). To determine if these measured thermal traits have a genetic basis and could

therefore evolve in response to natural selection, we raised hybrid marine-freshwater F1 ($N = 2$)

and F2 ($N = 4$; Fig. S3) families under common garden conditions and used these fish for

genome-wide linkage map construction (Table S1) and quantitative trait loci (QTL) mapping.

Using 41,840 high-quality single nucleotide variants generated from RAD sequencing, we

identified two significant QTL for CTmin, two significant QTL for CTmax, and three significant

QTL for thermal preference (Table 1, Fig. 2). The trait variance that each QTL explained ranged

from 28.9% (CTmin QTL on linkage group III) to 87.1% (thermal preference QTL on linkage

group VII).

Geographic range of projected suitable habitat

Based on sea ice extent and bathymetry alone, our present-day suitable habitat model

suggests a marine range for this population from the southern Bering Sea to northern Washington

state, and along the southeast Alaskan Panhandle (combined shaded area in Fig. 3a,

approximately 654,122 km^2). End-of-century IPCC projections resulted in a substantial increase

in the overall suitable habitat area for stickleback, with an 873,059 km^2 increase in total area

(combined shaded area in Fig. 3b-e, approximately 1,527,181 km^2) in association with a

reduction in sea ice concentration at the northern end of the range.

'No evolution' SDM

We used our observed values of CTmin, CTmax, and thermal preference to first inform

the boundaries of three distinct environmental regions in species distribution models (SDMs): i)

a 'Normal Behaviour' (NB) area that included environmental temperatures associated with an

absence of an observable behavioural stress response (5.0 to 25.0 °C), ii) a 'Within Physiological

Limits' (WPL) area that included environmental temperatures within the range of their measured

physiological limits (1.8 to 30.1°C) , and iii) an 'Outside Physiological Limits' (OPL) area with

environmental temperatures that fall outside the measured physiological limits (below 1.8 and

above 30.1 °C). When we include thermal trait data from the wild marine populations, nearly the

entire range of suitable habitat was unaffected by thermal tolerance limits, apart from a slight

restriction at the northern end of the range (Fig. 3a). However, when restricted to the NB area,

the range becomes confined to the west of the northern tip of Kodiak Island (Fig. 3a), a limit

coinciding with the northern-most known marine population in the Pacific Northwest genetic

cluster (Morris et al., 2018). When temperature increases as projected by RCP 4.5, the 'no

evolution' model projects a 1,011,949 km^2 increase in the NB area within this newly suitable

habitat (Fig. 3b) when compared to the NB area in the current day model. Under RCP 8.5, the

entirety of suitable habitat area remains within tolerable limits in the 'no evolution' model (Fig.

3c), with a smaller proportion of the range (10.7%) falling outside of the NB area as compared to

the NB area under RCP 4.5 conditions.

'Evolution' SDMs

370   Incorporating the evolution of CTmin and CTmax into the SDMs ('evolution' models)

results in a large increase in the proportion of suitable habitat that falls within the NB area. At

the rate of 0.63 *haldanes* for CTmin (Barrett et al., 2011) and 0.19 *haldanes* for CTmax, the

entire range of the suitable habitat range falls within the NB area (Fig. 3d, 3e). These represent a

311,974 and 155,680 km^2 increase in the NB area relative to the 'no evolution' models under

RCP 4.5 and RCP 8.5, respectively, and an increase in 1,271,032 km^2 in the NB area when

compared to the current day model. If we only allow one of each of these traits to evolve at a

time, we find that only CTmin evolution results in a change to range distributions compared to

'no evolution' models (Fig. S4). Reducing the rate of evolution for these traits to the mean rate

of trait evolution reported in Sanderson et al. (2021) still results in all suitable habitat falling

within the NB area under both RCP 4.5 and RCP 8.5. Similarly, scaling CTmin to evolve at a

rate equivalent to the response expected if selection acted solely on the detected CTmin QTL on

either linkage group III (PVE = 28.9%) or XXI (PVE = 53.7%) resulted in identical range

distributions as the full 'evolution' models. Given that CTmax evolution had no impact on these

range projections, we did not construct 'scaled evolution' models for this trait.

Discussion

In this study, we have produced the first 'evolution-informed' species distribution models

that incorporate empirical data for temperature-associated traits and their evolution to understand

how end-of-century projections change with the inclusion of this data. We assessed physiological

(critical thermal minimum, CTmin, and maximum, CTmax), and behavioural (thermal

preference) thermal traits for threespine stickleback from wild marine and freshwater

populations, as well as hybrid F1 and F2 families. We provide novel characterization of

quantitative trait loci underlying these traits in this species, which are likely to be important

genetic targets of selection under climate change. Finally, we incorporated this trait data and

three empirical estimates of evolutionary rates into mechanistic niche area species distribution

models under two climate change scenarios. When trait data is included in projections but held

constant (*i.e.,* 'no evolution' models), there is an approximately 6-fold (RCP 4.5) and 7-fold

(RCP 8.5) increase in the geographic area of the NB area. The geographic ranges projected for

the end-of-century increased by over 7-fold when CTmin was allowed to evolve at observed

trait-specific rates in the 'evolution' models. Additionally, when a slower rate of CTmin was

used or the rate was scaled by the percent of the trait variance that is explained by individual loci

(reflective of a scenario where selection acted solely on those loci), this substantial increase in

NB area was still attained by the end of the century under either RCP scenario. These changes to

the projected species ranges under climate change underscore the importance of incorporating

behavioural as well as physiological data into SDMs (Sunday et al., 2012), as well as the key role

that thermal trait evolution could play in range shifts (Buckley et al., 2010; Evans et al., 2015;

Lyon et al., 2019).

Our models project a northward range expansion under climate change. This is

unsurprising because climate change opens newly available thermal niche space in waters north

of the current day geographic range (Alexander et al., 2018) (as is seen in the 'no evolution'

model northward expansion, Fig. 3b & c). Northward range expansion with climate change due

to increasing habitat availability has also been documented in birds (Melles et al., 2011; Rushing

et al., 2020; Tingley et al., 2009; Tombre et al., 2019), plants (D'Andrea et al., 2009), other

fishes (Fossheim et al., 2015; Spies et al., 2020; Yapıcı et al., 2016), and pest species (such as

ticks and mountain pine beetle, Clow et al., 2017; Kurz et al., 2008; Ogden et al., 2006; Sagurova

et al., 2019; Sambaraju et al., 2019), as well as in large scale analyses of diverse taxa assessing

the 'fingerprints' of climate change impacts (Parmesan & Yohe, 2003; Platts et al., 2019). What

is perhaps less expected is that our models reveal that evolution of cold tolerance, but not heat

tolerance, had a substantial impact on predicted ranges under climate change, despite end-of-

century climate change scenarios projecting an overall warmer, not cooler world. This result

occurs because even with climate change, waters within the marine range of this stickleback

population are not projected to reach temperatures that would surpass their current CTmax limit.

However, there is reason to think that the relevance of cold tolerance in climate change scenarios

is not specific to this system and could be generally important to consider. Climate change is

leading to an increase in the frequency of extreme temperature events (IPCC, 2014; Stott, 2016),

including both extreme heat and extreme cold (Herring et al., 2015, 2018, 2020), which could

drive selection on both CTmin and CTmax (Buckley & Huey, 2016; Denny & Dowd, 2012;

Hoffman & Sgrò, 2011; Kingsolver et al., 2011). Moreover, although northern waters are

warming on average, there has been considerable spatial heterogeneity in temperature change

(Walther et al., 2002). Importantly, northern waters are not predicted to be uniformly warmer

under climate change (Alexander et al., 2018; IPCC, 2014) relative to the southern waters in this

range, especially when seasonal variation is considered (Alexander et al., 2018). Seasonal

variation and more extreme thermal events under climate change scenarios may force organisms

to respond independently to both extreme heat and extreme cold in separate events (Herring et

al., 2015, 2018, 2020; Walther et al., 2002), and coupled with the correlation between CTmin

and CTmax (as observed here, Fig. S1), population persistence will rely on adaptation in both

upper and lower critical limits. Further investigations to test the empirical rate of evolution of

thermal behaviour, physiology, and the molecular underpinnings of these key traits would be

well served by assessing additional subpopulations along the latitudinal gradient inhabited by

stickleback to gain a more detailed understanding of these temperature-associated traits over a

wider environmental range.

In interpreting the projections we present here, it is important to note that using a

different methodology to assess thermal limits could result in different thresholds in the tolerance

and behaviour areas (Moyano et al., 2017). In addition, in the absence of an empirical estimate of

the rate of evolutionary change for CTmax in threespine stickleback, we are using a rate

observed for this trait in zebrafish as a proxy, which will necessarily be inaccurate for

stickleback. It is unclear whether the true rate of CTmax evolution in stickleback might be

slower or more rapid than the zebrafish rate. For instance, it is possible that in threespine

stickleback CTmax could evolve at similar rates as observed for CTmin, particularly given

similarities in genetic architecture and the correlation between these traits in this study. Although

incorporating evolution of CTmax had no impact on our projected ranges because temperatures

under RCP 4.5 and RCP 8.5 never exceeded current CTmax values, we suspect this trait will be

of strong importance in many other systems where populations reside in environments closer to

their current upper thermal limits (*e.g.,* Deutsch et al., 2008; Morgan et al., 2019; Nati et al.,

2021). Ultimately, given the dramatic temperature changes that are projected for marine systems

by the end-of-century (IPCC, 2014, 2018), there is a critical need for empirical measures of

evolutionary rates in thermal traits for predicting future species ranges.

While this set of mechanistic niche models builds upon current SDM work by

incorporating functional traits critical to population persistence as well as varied evolutionary

scenarios based on empirical measurements, there are limitations to these projections. We

present niche models of geographic species ranges, but future work in this area would benefit

from including a robust metric of species distribution and population genetic structure within

these mechanistic geographic ranges. Deriving distributional maps from statistical algorithms

that rely on species occurrence and bioclimatic variables would move beyond binary outcomes

of habitat suitability and provide a map of suitable habitat that is scored as an index and could

underlie the static and evolving trait areas. Species distributions that build upon these evolution-

informed species ranges could be an important tool for projecting population persistence under

climate change.

The efficiency of translating the selection acting on a trait into evolutionary response

across generations will depend on the genetic architecture of the trait and the extent of

environmental change (Dittmar et al., 2016; Orr, 2005; Rogers et al., 2012). Among-trait genetic

correlations can also either speed or slow adaptive evolution depending on whether correlations

are reinforcing or antagonistic to the direction of selection (Etterson & Shaw, 2001). In this case,

we observe a positive correlation between CTmin and CTmax that could facilitate selection for

increased temperature tolerance at both upper and lower limits. However, the empirical rates of

trait evolution that we incorporated into our SDMs were estimated using change in phenotypic

variation across generations (Barrett et al., 2011; Morgan et al., 2020), rather than evolution at

underlying loci. As such, it is likely that some proportion of the observed phenotypic change was

due to plastic responses. In our 'scaled evolution' models, we took a conservative approach by

restricting phenotypic evolution to only occur through heritable change via the loci shown here

to be associated with the trait. The large effect loci that we identified are consistent with

expectations from theory that suggest prolonged bouts of adaptation with gene flow (as expected

in this system, Jones, Chan, et al., 2012; Rogers et al., 2012; Schluter et al., 2010; Schluter et al.,

2021; Schluter & Conte, 2009) will favour architectures characterized by fewer, larger effect,

more tightly linked alleles (Griswold & Whitlock, 2003; Via et al., 2012; Yeaman & Otto, 2011;

Yeaman & Whitlock, 2011). However, the effects of the QTL identified from these populations

are likely overestimated, while other loci may have gone undetected (*sensu* the Beavis Effect,

Beavis, 1994). These loci may have been detected in some F2 families and not in others because

each mapping family has their own specific meiotic events that enable the identification of

certain loci. Additionally, there are many key functional traits that are relevant to population

persistence under a changing climate, and it is possible that the traits focused on here, or

unexplored traits related to thermal performance, could display antagonistic pleiotropy.

Collectively, the inclusion of thermal traits and their evolution alters the projected ranges

of marine threespine stickleback, with a substantial increase in the projected area that the species

may occupy under climate change forecasts. Many traits are evolving in response to climate

change (Eliason et al., 2011; Gómez-Ruiz & Lacher, 2019; Horton et al., 2020; Hovel et al.,

2016) and SDMs that do not take trait data (and trait evolution) into account could provide less

accurate predictions about future species distributions under climate change (Bush et al., 2016;

Guisan et al., 2013; Huey et al., 2012; Kearney & Porter, 2004) - an issue of particular concern

for species at risk and pest species undergoing range expansion (Bebber, 2015; Cullingham et al.,

2011; McLeod et al., 2012). Our results provide a framework for addressing this problem, which

will have critical implications for the application of these models in policy, sustainable resource

management, and the protection of biodiversity in a changing climate.

References

Alexander, M. A., Scott, J. D., Friedland, K. D., Mills, K. E., Nye, J. A., Pershing, A. J., &
Thomas, A. C. (2018). Projected sea surface temperatures over the 21st century: Changes in
the mean, variability and extremes for large marine ecosystem regions of Northern Oceans.
Elementa, *6*. https://doi.org/10.1525/elementa.191

Angilletta, M. J., Niewiarowski, P. H., & Navas, C. A. (2002). The evolution of thermal
physiology in ectotherms. *Journal of Thermal Biology*, *27*, 249–268.
https://doi.org/10.2741/E148

Araújo, M. B., Ferri-Yáñez, F., Bozinovic, F., Marquet, P. A., Valladares, F., & Chown, S. L.
(2013). Heat freezes niche evolution. *Ecology Letters*, *16*(9), 1206–1219.
https://doi.org/10.1111/ele.12155

Arends, D., Prins, P., Broman, K. W., & Jansen, R. C. (2014). Tutorial-Multiple-QTL Mapping
(MQM) Analysis for R/qtl. In *http://www.rqtl.org/tutorials/MQM-tour.pdf*.

Barghi, N., Hermisson, J., & Schlötterer, C. (2020). Polygenic adaptation: a unifying framework
to understand positive selection. *Nature Reviews Genetics*, *21*(12), 769–781.
https://doi.org/10.1038/s41576-020-0250-z

Barrett, R., Paccard, A., Healy, T., Bergek, S., Schulte, P., Schluter, D., & Rogers, S. (2011).
Rapid evolution of cold tolerance in stickleback. *Proceedings of Royal Society B*,
278(1703), 233–238. https://doi.org/10.1098/rspb.2010.0923

Bay, R. A., Rose, N., Barrett, R., Bernatchez, L., Ghalambor, C. K., Lasky, J. R., Brem, R. B.,
Palumbi, S. R., & Ralph, P. (2017). Predicting Responses to Contemporary Environmental
Change Using Evolutionary Response Architectures. *The American Naturalist*, *189*(5), 463–
473. https://doi.org/10.1086/691233

Bayly, I. A. E. (2003). Salinity Tolerance and Osmotic Behavior of Animals in Athalassic Saline
and Marine Hypersaline Waters. *Annual Review of Ecology and Systematics*, *3*(1), 233–268.

https://doi.org/10.1146/annurev.es.03.110172.001313

Beavis, W. (1994). The power and deceit of QTL experiments: lessons from comparative QTL
studies. *49th Annual Corn & Sorghum Research Conference*, 250–266.

Bebber, D. P. (2015). Range-Expanding Pests and Pathogens in a Warming World. *Annual*
Review of Phytopathology, *53*, 335–356. https://doi.org/10.1146/annurev-phyto-080614-
120207

Beitinger, T., Bennett, W., & McCauley, R. (2000). Temperature tolerances of North American
freshwater fishes exposed to dynamic changes in temperature. *Environmental Biology of*
Fishes, *58*, 237–275. https://doi.org/10.1023/A:1007676325825

Bestion, E., Clobert, J., & Cote, J. (2015). Dispersal response to climate change: Scaling down to
intraspecific variation. *Ecology Letters*, *18*(11), 1226–1233.
https://doi.org/10.1111/ele.12502

Bowler, K., & Terblanche, J. S. (2008). Insect thermal tolerance: What is the role of ontogeny,
ageing and senescence? *Biological Reviews*, *83*, 339–355. https://doi.org/10.1111/j.1469-
185X.2008.00046.x

Broman, K. W., & Sen, S. (2009). *A Guide to QTL Mapping with R/qtl* (Vol. 46). Springer.

Broman, K. W., Wu, H., Sen, Ś., & Churchill, G. A. (2003). R/qtl: QTL mapping in experimental
crosses. *Bioinformatics*, *19*(7), 889–890. https://doi.org/10.1093/bioinformatics/btg112

Bryant, H. J., Chung, D. J., & Schulte, P. M. (2018). Subspecies differences in thermal
acclimation of mitochondrial function and the role of uncoupling proteins in killifish. *The*
Journal of Experimental Biology, jeb.186320. https://doi.org/10.1242/jeb.186320

Buckley, L., & Huey, R. (2016). How extreme temperatures impact organisms and the evolution
of their thermal tolerance. *Integrative and Comparative Biology*, *56*(1), 98–109.
https://doi.org/10.1093/icb/icw004

Buckley, L., Urban, M., Angilletta, M., Crozier, L., Rissler, L., & MW, S. (2010). Can
mechanism inform species' distribution models? *Ecology Letters*, *13*, 1041–1054.
https://doi.org/10.1111/j.1461-0248.2010.01479.x

Bush, A., Mokany, K., Catullo, R., Hoffmann, A., Kellermann, V., Sgrò, C., McEvey, S., &
Ferrier, S. (2016). Incorporating evolutionary adaptation in species distribution modelling
reduces projected vulnerability to climate change. *Ecology Letters*, *19*(12), 1468–1478.
https://doi.org/10.1111/ele.12696

Catchen, J., Hohenlohe, P. A., Bassham, S., Amores, A., & Cresko, W. A. (2013). Stacks: An
analysis tool set for population genomics. *Molecular Ecology*, *22*(11), 3124–3140.
https://doi.org/10.1111/mec.12354

Catullo, R. A., Ferrier, S., & Hoffmann, A. A. (2015). Extending spatial modelling of climate
change responses beyond the realized niche: Estimating, and accommodating, physiological

limits and adaptive evolution. *Global Ecology and Biogeography*, *24*(10), 1192–1202.
https://doi.org/10.1111/geb.12344

Cheng, C. H. C., & Detrich, H. W. (2007). Molecular ecophysiology of Antarctic notothenioid
fishes. *Philosophical Transactions of the Royal Society B: Biological Sciences*, *362*(1488),
2215–2232. https://doi.org/10.1098/rstb.2006.1946

Cingolani, P., Patel, V.M., Coon, M., Nguyen, T., Land, S.J., Ruden, D.M., & Lu., X. (2012).
Using Drosophila melanogaster as a model for genotoxic chemical mutational studies with a
new program, SnpSift. *Frontiers in Genetics, 3*(1).

Clow, K. M., Leighton, P. A., Ogden, N. H., Lindsay, L. R., Michel, P., Pearl, D. L., & Jardine,
C. M. (2017). Northward range expansion of Ixodes scapularis evident over a short
timescale in Ontario, Canada. *PLoS ONE*, *12*(12), 1–15.
https://doi.org/10.1371/journal.pone.0189393

Cooper, B., Adriaenssens, B., & Killen, S. S. (2018). Individual variation in the compromise
between social group membership and exposure to preferred temperatures. *Proceedings of*
the Royal Society B: Biological Sciences, *285*(1880).
https://doi.org/10.1098/rspb.2018.0884

Coutant, C. C. (1987). Thermal preference: when does an asset become a liability?
Environmental Biology of Fishes, *18*(5), 161–172. https://doi.org/10.1007/BF00000356

Cullingham, C. I., Cooke, J. E. K., Dang, S., Davis, C. S., Cooke, B. J., & Coltman, D. W.
(2011). Mountain pine beetle host-range expansion threatens the boreal forest. *Molecular*
Ecology, *20*(10), 2157–2171. https://doi.org/10.1111/j.1365-294X.2011.05086.x

D'Andrea, L., Broennimann, O., Kozlowski, G., Guisan, A., Morin, X., Keller-Senften, J., &
Felber, F. (2009). Climate change, anthropogenic disturbance and the northward range
expansion of Lactuca serriola (Asteraceae). *Journal of Biogeography*, *36*(8), 1573–1587.
https://doi.org/10.1111/j.1365-2699.2008.02060.x

Danecek, P., Auton, A., Abecasis, G., Albers, C. A., Banks, E., DePristo, M. A., Handsaker, R.
E., Lunter, G., Marth, G. T., Sherry, S. T., McVean, G., Durbin, R., & Group, 1000
Genomes Project Analysis. (2011). The variant call format and VCFtools. *Bioinformatics*,
27(15), 2156–2158. https://doi.org/10.1093/bioinformatics/btr330

Denny, M. W., & Dowd, W. W. (2012). Biophysics, environmental stochasticity, and the
evolution of thermal safety margins in intertidal limpets. *Journal of Experimental Biology*,
215(6), 934–947. https://doi.org/10.1242/jeb.058958

Detrich, H. W., Parker, S. K., Williams, J., Nogales, E., & Downing, K. H. (2000). Cold
adaptation of microtubule assembly and dynamics. Structural interpretation of primary
sequence changes present in the α- and β-tubulins of antarctic fishes. *Journal of Biological*
Chemistry, *275*(47), 37038–37047. https://doi.org/10.1074/jbc.M005699200

Deutsch, C.A., Tewksbury, J.J., Huey, R.B., Sheldon, K.S., Ghalambor, C.K., Haak, D.C., &
Martin, P.R. (2008). Impacts of climate warming on terrestrial ectotherms across latitude.

Proceedings of the National Academic of Sciences, 105(18), 6668-6672.
www.pnas.orgcgidoi10.1073pnas.0709472105

Dillon, M. E., Wang, G., Garrity, P. A., & Huey, R. B. (2009). Thermal preference in
Drosophila. *Journal of Thermal Biology, 34*(3), 109–119.
https://doi.org/10.1016/j.jtherbio.2008.11.007.Review

Dittmar, E. L., Oakley, C. G., Conner, J. K., Gould, B. A., & Schemske, D. W. (2016). Factors
influencing the effect size distribution of adaptive substitutions. *Proceedings of the Royal*
Society B: Biological Sciences, *283*(1828), 20153065.
https://doi.org/10.1098/rspb.2015.3065

Divino, J. N., Monette, M. Y., McCormick, S. D., Yancey, P. H., Flannery, K. G., Bell, M. A.,
Rollins, J. L., Von Hippel, F. A., & Schultz, E. T. (2016). Osmoregulatory physiology and
rapid evolution of salinity tolerance in threespine stickleback recently introduced to fresh
water. *Evolutionary Ecology Research*, *17*(2), 179–201.

Eliason, E. J., Clark, T. D., Hague, M. J., Hanson, L. M., Gallagher, Z. S., Jeffries, K. M., Gale,
M. K., Patterson, D. A., Hinch, S. G., & Farrell, A. P. (2011). Differences in thermal
tolerance among sockeye salmon populations. *Science*, *332*(6025), 109–112.
https://doi.org/10.1126/science.1199158

Environmental Systems Research Institute. (2017). *ArcGIS Desktop: Release 10.8.* (10.8).

Etterson, J. R., & Shaw, R. G. (2001). Constraint to adaptive evolution in response to global
warming. *Science*, *294*(5540), 151–154. https://doi.org/10.1126/science.1063656

Evans, T. G., Diamond, S. E., & Kelly, M. W. (2015). Mechanistic species distribution
modelling as a link between physiology and conservation. *Conservation Physiology*, *3*(1),
1–16. https://doi.org/10.1093/conphys/cov056

Everett, M. V., & Seeb, J. E. (2014). Detection and mapping of QTL for temperature tolerance
and body size in Chinook salmon (Oncorhynchus tshawytscha) using genotyping by
sequencing. *Evolutionary Applications*, *7*(4), 480–492. https://doi.org/10.1111/eva.12147

Eyer, P. A., Blumenfeld, A. J., & Vargo, E. L. (2019). Sexually antagonistic selection promotes
genetic divergence between males and females in an ant. *Proceedings of the National*
Academy of Sciences of the United States of America, *116*(48), 24157–24163.
https://doi.org/10.1073/pnas.1906568116

Fangue, N. A., Hofmeister, M., & Schulte, P. M. (2006). Intraspecific variation in thermal
tolerance and heat shock protein gene expression in common killifish, Fundulus
heteroclitus. *The Journal of Experimental Biology*, *209*(15), 2859–2872.
https://doi.org/10.1242/jeb.02260

Fangue, N. A., Richards, J. G., & Schulte, P. M. (2009). Do mitochondrial properties explain
intraspecific variation in thermal tolerance? *The Journal of Experimental Biology*, *212*(4),
514–522. https://doi.org/10.1242/jeb.024034

Fetterer, F., Savoie, M., Helfrich, S., & Clemene-Colon, P. (2010). U.S. National Ice Center and
National Snow and Ice Data Center. In *Multisensor Analyzed Sea Ice Extent - Northern*
Hemisphere (MASIE-NH) (Version 1). NSIDC: National Snow and Ice Data Center.
https://doi.org/https://doi.org/10.7265/N5GT5K3K.

Fisher, R. (1930). *The Genetical Theory of Natural Selection*.

Fossheim, M., Primicerio, R., Johannesen, E., Ingvaldsen, R. B., Aschan, M. M., & Dolgov, A.
V. (2015). Recent warming leads to a rapid borealization of fish communities in the Arctic.
Nature Climate Change, *5*(7), 673–677. https://doi.org/10.1038/nclimate2647

Gilchrist, G. W., & Huey, R. B. (1999). The direct response of Drosophila melanogaster to
selection on knockdown temperature. *Heredity*, *83*(1), 15–29.
https://doi.org/10.1038/sj.hdy.6885330

Gómez-Ruiz, E. P., & Lacher, T. E. (2019). Climate change, range shifts, and the disruption of a
pollinator-plant complex. *Scientific Reports*, *9*(1), 1–10. https://doi.org/10.1038/s41598-
019-53670-9

Greenwood, A. K., Jones, F. C., Chan, Y. F., Brady, S. D., Absher, D. M., Grimwood, J.,
Schmutz, J., Myers, R. M., Kingsley, D. M., & Peichel, C. L. (2011). The genetic basis of
divergent pigment patterns in juvenile threespine sticklebacks. *Heredity*, *107*(2), 155–166.
https://doi.org/10.1038/hdy.2011.1

Griswold, C. K., & Whitlock, M. C. (2003). The Genetics of Adaptation: The Roles of
Pleiotropy, Stabilizing Selection and Drift in Shaping the Distribution of Bidirectional
Fixed Mutational Effects. *Genetics*, *165*(4), 2181–2192.
https://doi.org/10.1093/genetics/165.4.2181

Guisan, A., Tingley, R., Baumgartner, J. B., Naujokaitis-Lewis, I., Sutcliffe, P. R., Tulloch, A. I.
T., Regan, T. J., Brotons, L., Mcdonald-Madden, E., Mantyka-Pringle, C., Martin, T. G.,
Rhodes, J. R., Maggini, R., Setterfield, S. A., Elith, J., Schwartz, M. W., Wintle, B. A.,
Broennimann, O., Austin, M., … Buckley, Y. M. (2013). Predicting species distributions for
conservation decisions. *Ecology Letters*, *16*(12), 1424–1435.
https://doi.org/10.1111/ele.12189

Hangartner, S., & Hoffmann, A. A. (2016). Evolutionary potential of multiple measures of upper
thermal tolerance in Drosophila melanogaster. *Functional Ecology*, *30*(3), 442–452.
https://doi.org/10.1111/1365-2435.12499

Harrell, F. (2014). *Hmisc: Harrell Miscellaneous*. 2014.

Healy, T. M., Brennan, R. S., Whitehead, A., & Schulte, P. M. (2018). Tolerance traits related to
climate change resilience are independent and polygenic. *Global Change Biology*, *24*(11),
5348–5360. https://doi.org/10.1111/gcb.14386

Hendry, A. P., Peichel, C. L., Matthews, B., Boughman, J. W., & Nosil, P. (2013). Stickleback
research: The now and the next. *Evolutionary Ecology Research*, *15*(2), 111–141.

Herring, S., Christidis, N., Hoell, A., Hoerling, M., & Stott, P. (2020). Explaining extreme events
of 2018 from a climate perspective. *Bulletin of the American Meteorological Society*,
101(1), 1–146. https://doi.org/10.1175/1520-0477-95.9.s1.1

Herring, S., Christidis, N., Hoell, A., Kossin, J., Schreck, C., & Stott, P. (2018). Explaining
extreme events of 2016 from a climate perspective. *Bulletin of the American Meteorological*
Society, *99*(1), S1–S157. https://journals.ametsoc.org/view/journals/bams/99/1/bams-
explainingextremeevents2016.1.xml

Herring, S., Hoerling, M., Kossing, J., Peterson, T., & Stott, P. (2015). *Explaining extreme*
events of 2014 from a climate perspective. *96*(12), 1–180.

Hijmans, R. J., Etten, J. van, Sumner, M., Cheng, J., Bevan, A., Bevan, R., Busetto, L., Canty,
M., Forrest, D., Ghosh, A., Golicher, D., Gray, J., & Greenberg, J. A. (2020). *raster:*
Geographic data analysis and modelling. https://cran.r-
project.org/web/packages/raster/raster.pdf

Hochachka, P. W., & Somero, G. N. (2002). Mechanism and process in physiological evolution.
Biochemical Adaptation, *480*.

Hoffman, A., & Sgrò, C. (2011). Climate change and evolutionary adaptation. *Nature*, *470*, 479–
485.

Horton, K. G., La Sorte, F. A., Sheldon, D., Lin, T. Y., Winner, K., Bernstein, G., Maji, S.,
Hochachka, W. M., & Farnsworth, A. (2020). Phenology of nocturnal avian migration has
shifted at the continental scale. *Nature Climate Change*, *10*(1), 63–68.
https://doi.org/10.1038/s41558-019-0648-9

Hovel, R. A., Carlson, S. M., & Quinn, T. P. (2016). Climate change alters the reproductive
phenology and investment of a lacustrine fish, the three-spine stickleback. *Global Change*
Biology, 1–13. https://doi.org/10.1111/gcb.13531

Hu, D., Guan, Z., Tian, W., & Ren, R. (2018). Recent strengthening of the stratospheric Arctic
vortex response to warming in the central North Pacific. *Nature Communications*, *9*(1).
https://doi.org/10.1038/s41467-018-04138-3

Huey, R. B., Kearney, M. R., Krockenberger, A., Holtum, J. A. M., Jess, M., & Wiliams, S. E.
(2012). Predicting organismal vulnerability to climate warming: roles of behaviour,
physiology and adaptation. *Philosophical Transactions of the Royal Society B: Biological*
Sciences, *367*, 1665–1679. https://doi.org/10.1098/rstb.2012.0005

Huey, R. B., & Kingsolver, J. G. (1989). Evolution of thermal sensitivity of ectotherm
performance. *Trends in Ecology and Evolution*, *4*(5), 131–135.

Huey, R. B., & Kingsolver, J. G. (2011). Variation in universal temperature dependence of
biological rates. *Proceedings of the National Academy of Sciences of the United States of*
America, *108*(26), 10377–10378. https://doi.org/10.1073/pnas.1107430108

Huey, R. B., & Slatkin, M. (1976). Costs and benefits of lizard thermoregulation. *Quarterly*

Review of Biology, *51*(3), 363–384.

Hutchison, V. H. (1961). Comparative Biology Critical Thermal Maxima in Salamanders. *Physiological Zoology*, *34*(2), 92–125.

IPCC. (2014). Climate Change 2014: Synthesis Report. In *Fifth Assessment Report of the Intergovernmental Panel on Climate Change*. https://doi.org/10.1016/S0022-0248(00)00575-3

IPCC. (2018). *IPCC special report on the impacts of global warming of 1.5 °C - Summary for policy makers*. *October 2018*. http://www.ipcc.ch/report/sr15/

Jackson, T. R., Ferguson, M. M., Danzmann, R. G., Fishback, A. G., Ihssen, P. E., O'Connell, M., & Crease, T. J. (1998). Identification of two QTL influencing upper temperature tolerance in three rainbow trout (Oncorhynchus mykiss) half-sib families. *Heredity*, *80*(2), 143–151. https://doi.org/10.1038/sj.hdy.6882890

Jin, Y., Zhou, T., Geng, X., Liu, S., Chen, A., Yao, J., Jiang, C., Tan, S., Su, B., & Liu, Z. (2017). A genome-wide association study of heat stress-associated SNPs in catfish. *Animal Genetics*, *48*(2), 233–236. https://doi.org/10.1111/age.12482

Johannessen, O., Bengtsson, L., Miles, M., Nagurnyi, A., Zakharov, V., Bobylev, L., Pettersson, L., Hasselmann, K., & Cattle, H. (2004). Arctic climate change: observed and modelled temperature and sea-ice variability. *Tellus A: Dynamic Meteorology and Oceanography*, *56*(4), 328–341. https://doi.org/10.3402/tellusa.v56i4.14418

Jones, F. C., Chan, Y. F., Schmutz, J., Grimwood, J., Brady, S. D., Southwick, A. M., Absher, D. M., Myers, R. M., Reimchen, T. E., Deagle, B. E., Schluter, D., & Kingsley, D. M. (2012). A genome-wide SNP genotyping array reveals patterns of global and repeated species-pair divergence in sticklebacks. *Current Biology*, *22*(1), 83–90. https://doi.org/10.1016/j.cub.2011.11.045

Jones, F. C., Grabherr, M. G., Chan, Y. F., Russell, P., Mauceli, E., Johnson, J., Swofford, R., Pirun, M., Zody, M. C., White, S., Birney, E., Searle, S., Schmutz, J., Grimwood, J., Dickson, M. C., Myers, R. M., Miller, C. T., Summers, B. R., Knecht, A. K., … Kingsley, D. M. (2012). The genomic basis of adaptive evolution in threespine sticklebacks. *Nature*, *484*(7392), 55–61. https://doi.org/10.1038/nature10944

Kearney, M., & Porter, W. (2009). Mechanistic niche modelling: Combining physiological and spatial data to predict species' ranges. *Ecology Letters*, *12*(4), 334–350. https://doi.org/10.1111/j.1461-0248.2008.01277.x

Kearney, M., & Porter, W. P. (2004). Mapping the fundamental niche: Physiology, climate, and the distribution of a nocturnal lizard. *Ecology*, *85*(11), 3119–3131. https://doi.org/10.1890/03-0820

Kingsolver, J. G., Arthur Woods, H., Buckley, L. B., Potter, K. A., MacLean, H. J., & Higgins, J. K. (2011). Complex life cycles and the responses of insects to climate change. *Integrative and Comparative Biology*, *51*(5), 719–732. https://doi.org/10.1093/icb/icr015

Kirch, M., Romundset, A., Gilbert, M. T. P., Jones, F. C., & Foote, A. D. (2021). Ancient and
modern stickleback genomes reveal the demographic constraints on adaptation. *Current*
Biology, *31*(9), 2027-2036.e8. https://doi.org/10.1016/j.cub.2021.02.027

Kurz, W. A., Dymond, C. C., Stinson, G., Rampley, G. J., Neilson, E. T., Carroll, A. L., Ebata,
T., & Safranyik, L. (2008). Mountain pine beetle and forest carbon feedback to climate
change. *Nature*, *452*(7190), 987–990. https://doi.org/10.1038/nature06777

Larson, W. A., McKinney, G. J., Limborg, M. T., Everett, M. V., Seeb, L. W., & Seeb, J. E.
(2016). Identification of Multiple QTL Hotspots in Sockeye Salmon (Oncorhynchus nerka)
Using Genotyping-by-Sequencing and a Dense Linkage Map. *Journal of Heredity*, *107*(2),
122–133. https://doi.org/10.1093/jhered/esv099

Li, H., & Durbin, R. (2010). Fast and accurate long-read alignment with Burrows-Wheeler
transform. *Bioinformatics*, *26*(5), 589–595. https://doi.org/10.1093/bioinformatics/btp698

Lyon, N. J., Debinski, D. M., & Rangwala, I. (2019). Evaluating the Utility of Species
Distribution Models in Informing Climate Change-Resilient Grassland Restoration Strategy.
Frontiers in Ecology and Evolution, *7*(February), 1–8.
https://doi.org/10.3389/fevo.2019.00033

McLeod, D. J., Hallegraeff, G. M., Hosie, G. W., & Richardson, A. J. (2012). Climate-driven
range expansion of the red-tide dinoflagellate Noctiluca scintillans into the Southern Ocean.
Journal of Plankton Research, *34*(4), 332–337. https://doi.org/10.1093/plankt/fbr112

Melles, S. J., Fortin, M. J., Lindsay, K., & Badzinski, D. (2011). Expanding northward: Influence
of climate change, forest connectivity, and population processes on a threatened species'
range shift. *Global Change Biology*, *17*(1), 17–31. https://doi.org/10.1111/j.1365-
2486.2010.02214.x

Morgan, R., Finnøen, M. H., Jensen, H., Pélabon, C., & Jutfelt, F. (2020). Low potential for
evolutionary rescue from climate change in a tropical fish. *Proceedings of the National*
Academy of Sciences of the United States of America, *117*(52), 33365–33372.
https://doi.org/10.1073/PNAS.2011419117

Morgan, R., Finnøen, M. H., & Jutfelt, F. (2018). CTmax is repeatable and doesn't reduce
growth in zebrafish. *Scientific Reports*, *8*(1), 1–8. https://doi.org/10.1038/s41598-018-
25593-4

Morgan, R., Sundin, J., Finnøen, M.H., Dresler, G., Martínex Vendrell, M., Dey, A., Sarkar, K.,
& Jutfelt, F. (2019). Are model organisms representative for climate change research?
Testing thermal tolerance in wild and laboratory zebrafish populations. *Conservation*
Physiology, *7*(1), coz036. https://doi.org/10.1093/conphys/coz036

Morris, M. R. J., Bowles, E., Allen, B. E., Jamniczky, H. A., & Rogers, S. M. (2018).
Contemporary ancestor? Adaptive divergence from standing genetic variation in Pacific
marine threespine stickleback. *BMC Evolutionary Biology*, *18*(1), 1–21.
https://doi.org/10.1186/s12862-018-1228-8

Moyano, M., Candebat, C., Ruhbaum, Y., Álvarez-Fernández, S., Claireaux, G., Zambonino-
Infante, J. L., & Peck, M. A. (2017). Effects of warming rate, acclimation temperature and
ontogeny on the critical thermal maximum of temperate marine fish larvae. *PLoS ONE*,
12(7), 1–23. https://doi.org/10.1371/journal.pone.0179928

Muñoz, N. J., Farrell, A. P., Heath, J. W., & Neff, B. D. (2014). Adaptive potential of a Pacific
salmon challenged by climate change. *Nature Climate Change*, *5*(2), 163–166.
https://doi.org/10.1038/nclimate2473

Nati, J.J.H., Svendsen, M.B.S., Marras, S., Killen, S.S., Steffensen, J.F., McKenzie, D.J., &
Domenici, P. (2021). Intraspecific variation in thermal tolerance differs between tropical
and temperate fishes. *Scientific Reports, 11*(1), 1-8. https://doi.org/10.1038/s41598-021-
00695-8

Ogden, N. H., Maarouf, A., Barker, I. K., Bigras-Poulin, M., Lindsay, L. R., Morshed, M. G.,
O'Callaghan, C. J., Ramay, F., Waltner-Toews, D., & Charron, D. F. (2006). Climate
change and the potential for range expansion of the Lyme disease vector Ixodes scapularis
in Canada. *International Journal for Parasitology*, *36*(1), 63–70.
https://doi.org/10.1016/j.ijpara.2005.08.016

Orr, H. A. (1998). The population genetics of adaptation: The distribution of factors fixed during
adaptive evolution. *Evolution*, *52*(4), 935–949. https://doi.org/10.1111/j.1558-
5646.1998.tb01823.x

Orr, H. A. (2005). The genetic theory of adaptation: A brief history. *Nature Reviews Genetics*,
6(2), 119–127. https://doi.org/10.1038/nrg1523

Ouellette, L. A., Reid, R. W., Blanchard, S. G., & Brouwer, C. R. (2018). LinkageMapView-
rendering high-resolution linkage and QTL maps. *Bioinformatics*, *34*(2), 306–307.
https://doi.org/10.1093/bioinformatics/btx576

Parmesan, C., & Yohe, G. (2003). *A globally coherent fingerprint of climate change impacts*
across natural systems. *421*, 37–42. https://doi.org/10.1038/nature01286

Perry, G. M. L., Danzmann, R. G., Ferguson, M. M., & Gibson, J. P. (2001). Quantitative trait
loci for upper thermal tolerance in outbred strains of rainbow trout (Oncorhynchus mykiss).
Heredity, *86*(3), 333–341. https://doi.org/10.1046/j.1365-2540.2001.00838.x

Peterson, B. K., Weber, J. N., Kay, E. H., Fisher, H. S., & Hoekstra, H. E. (2012). Double digest
RADseq: An inexpensive method for de novo SNP discovery and genotyping in model and
non-model species. *PLoS ONE*, *7*(5). https://doi.org/10.1371/journal.pone.0037135

Platts, P. J., Mason, S. C., Palmer, G., Hill, J. K., Oliver, T. H., Powney, G. D., Fox, R., &
Thomas, C. D. (2019). Habitat availability explains variation in climate-driven range shifts
across multiple taxonomic groups. *Scientific Reports*, *9*(1), 1–10.
https://doi.org/10.1038/s41598-019-51582-2

Puritz, J. B., Hollenbeck, C. M., & Gold, J. R. (2014). *dDocent*□: a RADseq, variant-calling
pipeline designed for population genomics of non-model organisms. *PeerJ*, *2*, e431.

https://doi.org/10.7717/peerj.431

R Core Team. (2021). *R: A language and environment for statistical computing.*

Rastas, P. (2017). Lep-MAP3: Robust linkage mapping even for low-coverage whole genome
sequencing data. *Bioinformatics*, *33*(23), 3726–3732.
https://doi.org/10.1093/bioinformatics/btx494

Reynolds, R., Smith, T., Liu, C., Chelton, D., Casey, K., & Schlax, M. (2007). Daily High-
Resolution-Blended Analyses for Sea Surface Temperature. *Journal of Climate*, *20*(22),
5473–5496. https://doi.org/10.1175/2007jcli1824.1

Roger, A., Stuetz, R., Ove, K., Giraudoux, P., & Santilli, S. (2020). *rgeos: Interface to geometry*
engine - open.

Rogers, S. M., Tamkee, P., Summers, B., Balabahadra, S., Marks, M., Kingsley, D. M., &
Schluter, D. (2012). Genetic Signature of Adaptive Peak Shift in Threespine Stickleback.
Evolution, *Fisher 1930*, 2439–2451. https://doi.org/10.5061/dryad.6jj614kh

Rolandi, C., Lighton, J. R. B., de la Vega, G. J., Schilman, P. E., & Mensch, J. (2018). Genetic
variation for tolerance to high temperatures in a population of Drosophila melanogaster.
Ecology and Evolution, *8*(21), 10374–10383. https://doi.org/10.1002/ece3.4409

Rushing, C. S., Andrew Royle, J., Ziolkowski, D. J., & Pardieck, K. L. (2020). Migratory
behavior and winter geography drive differential range shifts of eastern birds in response to
recent climate change. *Proceedings of the National Academy of Sciences of the United*
States of America, *117*(23), 12897–12903. https://doi.org/10.1073/pnas.2000299117

Sagurova, I., Ludwig, A., Ogden, N. H., Pelcat, Y., Dueymes, G., & Gachon, P. (2019).
Predicted northward expansion of the geographic range of the tick vector amblyomma
americanum in North America under future climate conditions. *Environmental Health*
Perspectives, *127*(10), 1–14. https://doi.org/10.1289/EHP5668

Sambaraju, K. R., Carroll, A. L., & Aukema, B. H. (2019). Multiyear weather anomalies
associated with range shifts by the mountain pine beetle preceding large epidemics. *Forest*
Ecology and Management, *438*(September 2018), 86–95.
https://doi.org/10.1016/j.foreco.2019.02.011

Sanderson, S., Beausoleil, M., O'Dea, R. E., Wood, Z. T., Correa, C., Frankel, V., Gorné, L. D.,
Haines, G. E., Kinnison, M. T., Oke, K. B., Pelletier, F., Pérez☐Jvostov, F., Reyes☐Corral,
W. D., Ritchot, Y., Sorbara, F., Gotanda, K. M., & Hendry, A. P. (2021). The Pace of
Modern Life, Revisited. *Molecular Ecology*, *November*, 1–16.
https://doi.org/10.1111/mec.16299

Schluter, D, Marchinko, K., Barrett, R., & Rogers, S. (2010). Natural selection and the genetics
of adaptation in threespine stickleback. *Philosophical Transactions of the Royal Society B:*
Biological Sciences, *365*(1552), 2479–2486. https://doi.org/10.1098/rstb.2010.0036

Schluter, Dolph, & Conte, G. L. (2009). Genetics and ecological speciation. *Proceedings of the*

National Academy of Sciences, *3*, 47–64. https://doi.org/10.17226/12692

Schluter, Dolph, Marchinko, K. B., Arnegard, M. E., Zhang, H., Brady, S. D., Jones, F. C., Bell,
M. A., & Kingsley, D. M. (2021). Fitness maps to a large-effect locus in introduced
stickleback populations. *Proceedings of the National Academy of Sciences of the United
States of America*, *118*(3), 1–10. https://doi.org/10.1073/pnas.1914889118

Sexton, J. P., McIntyre, P. J., Angert, A. L., & Rice, K. J. (2009). Evolution and Ecology of
Species Range Limits. *Annual Review of Ecology, Evolution, and Systematics*, *40*(1), 415–
436. https://doi.org/10.1146/annurev.ecolsys.110308.120317

Shin, S. C. hu., Ahn, D. H. wa., Kim, S. J. i., Pyo, C. W. o., Lee, H., Kim, M. K., Lee, J., Lee, J.
E. u., Detrich, H. W., Postlethwait, J. H., Edwards, D., Lee, S. G. u., Lee, J. H. yuc., & Park,
H. (2014). The genome sequence of the Antarctic bullhead notothen reveals evolutionary
adaptations to a cold environment. *Genome Biology*, *15*(9), 468.
https://doi.org/10.1186/s13059-014-0468-1

Spies, I., Gruenthal, K. M., Drinan, D. P., Hollowed, A. B., Stevenson, D. E., Tarpey, C. M., &
Hauser, L. (2020). Genetic evidence of a northward range expansion in the eastern Bering
Sea stock of Pacific cod. *Evolutionary Applications*, *13*(2), 362–375.
https://doi.org/10.1111/eva.12874

Stott, P. (2016). How climate change affects extreme weather events. *Science*, *352*(6293), 1517–
1518. https://doi.org/10.1126/science.aaf7271

Sunday, J. M., Bates, A. E., & Dulvy, N. K. (2011). Global analysis of thermal tolerance and
latitude in ectotherms. *Proceedings of the Royal Society B: Biological Sciences*, *278*(1713),
1823–1830. https://doi.org/10.1098/rspb.2010.1295

Sunday, J. M., Bates, A. E., & Dulvy, N. K. (2012). Thermal tolerance and the global
redistribution of animals. *Nature Climate Change*, *2*(9), 686–690.
https://doi.org/10.1038/nclimate1539

Tingley, M. W., Monahan, W. B., Beissinger, S. R., & Moritz, C. (2009). Birds track their
Grinnellian niche through a century of climate change. *Proceedings of the National
Academy of Sciences of the United States of America*, *106*(SUPPL. 2), 19637–19643.
https://doi.org/10.1073/pnas.0901562106

Tombre, I. M., Oudman, T., Shimmings, P., Griffin, L., & Prop, J. (2019). Northward range
expansion in spring-staging barnacle geese is a response to climate change and population
growth, mediated by individual experience. *Global Change Biology*, *25*(11), 3680–3693.
https://doi.org/10.1111/gcb.14793

Turko, A. J., Nolan, C. B., Balshine, S., Scott, G. R., & Pitcher, T. E. (2020). Thermal tolerance
depends on season, age and body condition in imperilled redside dace Clinostomus
elongatus. *Conservation Physiology*, *8*(1), 1–15. https://doi.org/10.1093/conphys/coaa062

Ungerer, M. C., & Riesebero, L. H. (2003). Genetic architecture of a selection response in
Arabidopsis thaliana. *Evolution*, *57*(11), 2531–2539. https://doi.org/10.1111/j.0014-

3820.2003.tb01497.x

Valladares, F., Matesanz, S., Guilhaumon, F., Araújo, M. B., Balaguer, L., Benito-Garzón, M.,
Cornwell, W., Gianoli, E., van Kleunen, M., Naya, D. E., Nicotra, A. B., Poorter, H., &
Zavala, M. A. (2014). The effects of phenotypic plasticity and local adaptation on forecasts
of species range shifts under climate change. *Ecology Letters*, *17*(11), 1351–1364.
https://doi.org/10.1111/ele.12348

Via, S., Conte, G., Mason-Foley, C., & Mills, K. (2012). Localizing FST outliers on a QTL map
reveals evidence for large genomic regions of reduced gene exchange during speciation-
with-gene-flow. *Molecular Ecology*, *21*(22), 5546–5560. https://doi.org/10.1111/mec.12021

Walther, G. R., Post, E., Convey, P., Menzel, A., Parmesan, C., Beebee, T. J. C., Fromentin, J.
M., Hoegh-Guldberg, O., & Bairlein, F. (2002). Ecological responses to recent climate
change. *Nature*, *416*(6879), 389–395. https://doi.org/10.1038/416389a

Weatherall, P., Marks, K., Jakobsson, M., Schmitt, T., Tani, S., Arndt, J., Rovere, M., Chayes,
D., Ferrini, V., & Wigley, R. (2015). A new digital bathymetric model of the world's
oceans. *Earth and Space Science*, *2*(8), 331–345.
https://doi.org/10.1002/2015EA000107.Received

Wei, T., Simko, V., Levy, M., Xie, Y., Jin, Y., & Zemla, J. (2017). Package 'corrplot.'
Statistician, *56*, 316–324.

Wiens, J., Stralberg, D., Jongsomjit, D., Howell, C., & Snyder, M. (2009). Niches, models, and
climate change: Assessing the assumptions and uncertainties. *Proceedings of the National*
Academy of Sciences, *106*(Supplement_2), 19729–19736.
https://doi.org/10.1073/pnas.0901639106

Yapıcı, S., Bilge, G., & Filiz, H. (2016). Northwards range expansion of Sparisoma cretense
(Linnaeus, 1758) in the Turkish Aegean Sea. *Journal of Aquaculture Engineering and*
Fisheries Research, *August*, 201–207. https://doi.org/10.3153/jaefr16022

Yeaman, S., & Otto, S. P. (2011). Establishment and maintenance of adaptive genetic divergence
under migration, selection, and drift. *Evolution*, *65*(7), 2123–2129.
https://doi.org/10.1111/j.1558-5646.2011.01277.x

Yeaman, S., & Whitlock, M. C. (2011). The genetic architecture of adaptation under migration-
selection balance. *Evolution*, *65*(7), 1897–1911. https://doi.org/10.1111/j.1558-
5646.2011.01269.x

Zeidberg, L. D., & Robison, B. H. (2007). Invasive range expansion by the Humboldt squid,
Dosidicus gigas, in the eastern North Pacific. *Proceedings of the National Academy of*
Sciences, *104*(31), 12948–12950. https://doi.org/10.1073/pnas.0702043104

Zweng, M. M., Reagan, J. R., Antonov, J. I., Mishonov, A. V., Boyer, T. P., Garcia, H. E.,
Baranova, O. K., Johnson, D. R., Seidov, D., & Bidlle, M. M. (2013). World Ocean Atlas
2013, Volume 2: Salinity. In *NOAA Atlas NESDIS 74* (Vol. 2, Issue 1).
https://doi.org/10.1182/blood-2011-06-357442

Acknowledgments

The authors acknowledge that the species collections took place on Huu-ay-aht and Sechelt First

Nations traditional territories and are grateful for the opportunity to conduct their research in

protected and sacred areas. We would like to thank the Bamfield Marine Sciences Centre for the

resources required to conduct this research, Sam Owens for the photo in Figure 1a, Peter Peller

for creating Figure 1b, Carolyn Tepolt for providing the high-resolution sea surface temperature

dataset from Reynolds et al. (2007) during a government shutdown, as well as Kiyoko Gotanda

for calculating and providing mean *haldane* values from the Sanderson et al. (2021) database.

We would like to thank Daniel Wuitchik, Sam Yeaman, Jennifer Sunday, and Patrick Nosil for

feedback on the manuscript, and the anonymous peer reviewers for improving this work. This

research was funded by Discovery Grants from the Natural Sciences and Engineering Research

Council of Canada to RDHB and SMR, a Canada Research Chair to RDHB, and a New Faculty

Award from Alberta Innovates Technology Futures to SMR.

Author Contributions

This study was designed by SJSW, RDHB, and SMR; fish husbandry and breeding by SJSW and

TNB; experimental data collection by SJSW; DNA sequencing and initial processing by AP;

bioinformatic and QTL analyses by SJSW; species distribution modelling by SJSW and SM; the

manuscript was written by SJSW, RDHB, and SMR, with input from authors; the study was

funded by HAJ, RDHB, and SMR.

Data Accessibility

The data associated with this work is available as follows:

ddRAD DNA sequences: available on NCBI (BioProject PRJNA795470)

964   Experimental data associated with thermal trait data: available on Dryad

965   (https://doi.org/10.5061/dryad.70rxwdbz6)

966   Annotated code and example files: available on GitHub

967   (https://github.com/sjswuitchik/gasAcu_qtl_sdm)

968   **Figures and Tables**
969
970   **Figure 1.** a) Adult threespine stickleback (*Gasterosteus aculeatus*) from a single genetic cluster
971   were sampled from b) two marine and two freshwater populations in the Canadian Pacific
972   Northwest. These populations were assayed for c) thermal preference (green), critical thermal
973   minima (CTmin, blue) and maxima (CTmax, red). Thermal trait values for marine populations
974   (M1 and M2) were incorporated into the species distribution models.

975
976   **Figure 2.** Linkage maps constructed for quantitative trait loci (QTL) analyses for a) H1 F2
977   hybrids (M1 x FW1) and b) H2 F2 hybrids (M1 x FW2) families, rendered with
978   LinkageMapView (Ouellette et al., 2018), with locations of QTL indicated (Table 1).

979
Figure 3. Changes in the projected range of marine threespine stickleback (*Gasterosteus
aculeatus*) in the northeast Pacific Ocean as a result of incorporating thermal traits into models of
a) current day environmental conditions, and under IPCC end-of-century projections RCP 4.5
and 8.5, either without trait evolution (b & c; 'no evolution' model, orange box) and with trait
evolution at a rate of 0.63 *haldanes* for critical thermal minimum (CTmin) and a rate of 0.19
haldanes for critical thermal maximum (CTmax) (d & e; 'evolution' model, red box). Sampling
locations are in the southeast area of the range presented here.

987
988   **Table 1.** Significant quantitative trait loci (QTL) for critical thermal minimum (CTmin), critical
989   thermal maximum (CTmax), and thermal preference (Preference) traits with the percent of trait
990   variance explained (PVE) for each QTL, the logarithm of odds (LOD) threshold for significance,
991   and the LOD peak observed for each QTL.

992
993   **Figure S1.** Correlations between critical thermal minimum (CTmin), critical thermal maximum
994   (CTmax), and thermal preference for threespine stickleback (*Gasterosteus aculeatus*).

995
996   **Figure S2.** Thermal preference (green), critical thermal minimum (CTmin, blue) and maximum
997   (CTmax, red) measurements for threespine stickleback (*Gasterosteus aculeatus*) from wild
998   freshwater (FW) populations, pure F1 freshwater (FW*F1), wild marine (M), and pure F1 marine
999   (M*F1) families raised in a common garden under a constant thermal environment.

1000
1001   **Figure S3.** Thermal preference (green), critical thermal minimum (CTmin, blue) and maximum
1002   (CTmax, red) measurements for threespine stickleback (*Gasterosteus aculeatus*) from hybrid
1003   marine-freshwater F2 families raised in a common garden under a constant thermal environment.
1004

1005   **Figure S4.** Changes in the projected range of marine threespine stickleback (*Gasterosteus*
1006   *aculeatus*) in the northeast Pacific Ocean compared to a) current day range as a result of
1007   incorporating the evolution of only critical thermal minimum (CTmin, b & c; orange box) at a
1008   rate of 0.63 *haldanes* and only critical thermal maximum (CTmax, d & e; red box) at a rate of
1009   0.19 *haldanes* under IPCC end-of-century projections RCP 4.5 and RCP 8.5
1010
1011   **Table S1.** Summaries of linkage maps constructed for quantitative trait loci (QTL) analyses.
1012

a) Current day

■ Outside Physiological Limits
■ Within Physiological Limits
■ Normal Behaviour

b) RCP 4.5

c) RCP 8.5

d) RCP 4.5

e) RCP 8.5