**Evolution of thermal physiology alters predicted species distributions under**

**climate change**

Sara J.S. Wuitchik[1,†,*], Stephanie Mogensen[1], Tegan N. Barry[1], Antoine Paccard[2,‡],

Heather A. Jamniczky[3], Rowan D.H. Barrett[2,§], and Sean M. Rogers[1,4,§]

[1]Department of Biological Sciences, University of Calgary, 2500 University Dr NW, Calgary,

AB, T2N 1N4, CANADA

[2]Redpath Museum and Department of Biology, McGill University, 845 Sherbrooke St W,

Montreal, QC, H3A 0G4, CANADA

[3]Department of Cell Biology & Anatomy, Cumming School of Medicine, University of Calgary,

3330 Hospital Dr NW, Calgary, T2N 4N1, CANADA

[4]Bamfield Marine Sciences Centre, 100 Pachena Rd, Bamfield, BC, V0R 1B0, CANADA

[§]equal co-senior authors

19  **\*Corresponding author: Sara J.S. Wuitchik**

20  E: sjswuit@g.harvard.edu

21  T: 857-292-9977

[†]Present affiliations: Informatics Group, Harvard University, 38 Oxford St, Cambridge, MA, 02138, USA & Department of Biology, Boston University, 5 Cummington Mall, Boston, MA, 02215, USA
[‡]Present affiliation: McGill University Genome Center, 740 Dr Penfield Avenue, Montreal, QC, H3A 1A5, Canada

Species distribution models (SDMs) are widely adopted to predict range shifts but can be unreliable under climate change scenarios[1] because they do not account for evolution. The thermal physiology of a species is a key determinant of range[2,3] but the impact of thermal trait evolution on SDMs has not been addressed. We identified a genetic basis for physiological traits that evolve in response to temperature change in threespine stickleback. Using these data, we created geographic range projections under two climate change scenarios where trait data was either static ('no evolution' model), allowed to evolve in agreement with published evolutionary rates for the trait ('evolution' model)[4], or allowed to evolve with the rate of evolution scaled in association with the variance that is explained by QTL ('PVE' model). Here, we show that incorporating these traits and their evolution into SDMs substantially altered the predicted ranges for a widespread panmictic marine population, with increases in area of over 7-fold. Evolution-informed SDMs should therefore improve the precision of forecasting range dynamics under climate change, thereby aiding in their application to management and the protection of biodiversity[5–7].

Temperature is a powerful driver of global biogeography and organism distributions frequently reflect temperature gradients in both aquatic and terrestrial habitats[8]. Many species adopt thermal strategies (such as thermoregulation or acclimation) that determine their thermal niche[9–11] and thermal traits can provide a target for directional selection if the environment changes to include temperatures outside the range encompassed by the thermal niche. Adaptation may thus permit species to persist at temperatures that would have previously led to extirpation[12,13]. Under moderate climate change scenarios, mean global oceanic temperature is predicted to increase in excess of 2°C by the end of the century[14], with more extreme changes predicted in localized regions[15]. Predicting species distribution patterns under climate change

therefore requires data for temperature-associated adaptive trait evolution, which can vary by

species and population. While there have been recent steps to incorporate theoretical trait

evolution into SDMs[3], to date, no model has used empirical estimates of evolutionary rate to

inform predictions about future species distributions.

Due to widespread phenotypic variation[16], genomic resources[17], the availability of

temperature-associated ecological and evolutionary trait data[4], and the ability to artificially breed

multiple hybrid generations in a common garden lab environment, threespine stickleback fish

(*Gasterosteus aculeatus*, Fig. 1a) are a useful vertebrate species for understanding the impact of

adaptation on range dynamics under climate change. Here, we incorporate the specific capacity

of marine populations to adapt their physiology to rapidly changing climate conditions to

characterize how adaptive trait variation affects projections of species range distributions under

climate change[18].

**Results**

We collected and reared wild marine and freshwater stickleback from two marine and

two freshwater locations (Fig. 1b). These stickleback exhibited a wide thermal tolerance range

bounded by a mean CTmin of 2.09 °C (+/- 1.13 °C SD) and a mean CTmax of 30.4 °C (+/- 2.64

°C SD)(Fig. 1c) and also tolerated a wide range of temperatures within which there was no

observable stress response (5.0 – 25.0 °C). To determine if these measured thermal traits have a

genetic basis and could therefore be subject to adaptive evolution, we raised hybrid marine-

freshwater F1 (N=2) and F2 (N=4) families under common garden conditions and used these fish

for genome-wide linkage map construction (Table S1) and quantitative trait loci (QTL) mapping.

Using 25,001 high-quality single nucleotide variants generated from restriction site-associated

DNA (RAD) sequencing, we identified one significant QTL for each thermal tolerance trait (Fig.

2b) that explained a high percentage of trait variance (PVE; CTmin = 54%, CTmax = 64%).

We used these genetically based traits to inform the boundaries of three distinct

environmental regions in species distribution models (SDMs) for marine stickleback based on

varying levels of physiological performance: i) a 'normal behaviour' envelope with

environmental temperatures associated with an absence of an observable behavioural stress

response (5.0 to 25.0 °C), ii) a 'within physiological limits' envelope with environmental

temperatures that fall within the range of the measured physiological limits (0.85 to 31.9 °C) ,

and iii) an 'outside of physiological limits' envelope with environmental temperatures that fall

outside the measured physiological limits (below 0.85 and above 31.9 °C). Based on sea ice

extent and bathymetry alone, our present-day correlative SDM suggests a marine range

distribution for these stickleback from the southern Bering Sea to northern Washington state, and

along the southeast Alaskan Panhandle (combined shaded area in Fig. 3a). When we include

species-specific thermal trait data from the wild marine populations, nearly the entire range of

suitable habitat was unaffected by thermal tolerance limits, with the exception of a slight

restriction at the northern end of the range (Fig. 3a). However, when restricted to the Normal

Behaviour area, the range becomes confined to the west of the northern tip of Kodiak Island (Fig.

3a), a limit coinciding with the northern-most known marine population in the Pacific Northwest

genetic cluster[19].

We next generated SDMs based on predicted end-of-century environmental variables

according to the Intergovernmental Panel on Climate Change (IPCC) representative

concentration pathways (RCPs) 4.5 and 8.5 from the Fifth Assessment Report[14,20]. End-of-

century IPCC predictions resulted in a substantial increase in the overall suitable habitat area for

stickleback, with a 2.25-fold or 1,338,219 km$^2$ increase (combined shaded area in Fig. 3b-e) in

association with a reduction in sea ice concentration at the northern end of the range. When

temperature increases as predicted by RCP 4.5 in the 'no evolution' model, there is a 5.86-fold

(1,011,949 km$^2$) increase in the Normal Behaviour area within this newly suitable habitat (Fig.

3b) when compared to the current day model. Under RCP 8.5 in the 'no evolution' model, the

entirety of suitable habitat area remains within tolerable limits (Fig. 3c), with a smaller

proportion of the range falling outside of the Normal Behaviour area compared to RCP 4.5

(10.7%) 'no evolution' model.

Incorporating the evolution of CTmin into the SDMs ('evolution' model) results in a

large increase in the proportion of suitable habitat that falls within the Normal Behaviour area.

We allowed CTmin to evolve at a rate of 0.63 *haldanes*, which is equal to the rate observed for

CTmin in marine stickleback[4] (there are currently no empirical estimates of evolutionary rate for

CTmax). Under RCP 4.5, almost all (99.9%) of the suitable habitat range falls within the Normal

Behaviour area (Fig. 3d), while under the RCP 8.5 'evolution' model, the entire range of suitable

habitat is within the Normal Behaviour area (Fig. 3e). These represent a 7.45-fold increase

(1,336,123 km$^2$) for RCP 4.5 and 7.46-fold increase (1,338,219 km$^2$) for RCP 8.5 in the Normal

Behaviour area when compared to the current day SDM.

We next considered the effect of limiting the evolutionary rate of CTmin based on the

observed genetic architecture of a single major effect locus ('PVE' model). Under the RCP 4.5

'PVE' model, we observed a 1.08-fold reduction in the Normal Behaviour area (109,120 km$^2$,

Fig. S3) when compared to the 'evolution' model (Fig. 3d). Under RCP 8.5 projections with the

'PVE' model, we observed a 3,905 km$^2$ decrease in the Normal Behaviour area (Fig. S3) when

compared to the 'evolution' model (Fig. 3e). This relatively small reduction in Normal

Behaviour area with adjusted PVE for the thermal traits under RCP 8.5 still results in a

1,334,314 km$^2$ increase in area compared to the current day SDM.

**Discussion**

We assessed the critical thermal minimum (CTmin) and maximum (CTmax) for

threespine stickleback from wild marine and freshwater populations, as well as F1 and F2

families in order to determine the genetic basis underlying traits that will be important for

population persistence under climate change. We incorporated the empirical ecological and

evolutionary trait data from wild marine populations into mechanistic species distribution models

under two climate change scenarios. We estimated the species distribution in the Pacific

Northwest marine environment while these traits were held constant ('no evolution'), allowed

CTmin to evolve in accordance with evolutionary rate estimates ('evolution'), and allowed

CTmin to evolve at a constrained rate associated with the percent of trait variance explained by

the single QTL we detected ('PVE'). The geographic ranges predicted for the end-of-century

species distributions increased by over 7-fold (RCP 4.5: 1,336,123 km$^2$; RCP 8.5: 1,338,219

km$^2$) when CTmin was allowed to evolve, a substantial increase over the 'no evolution' model.

Additionally, when CTmin evolution was constrained in the PVE model, there remained a ~6-

fold increase (RCP 4.5: 1,227,002 km$^2$; RCP 8.5: 1,334,314 km$^2$) in the geographic range

compared to current day. These differences in the predicted distributions underline the

significance of incorporating empirical evolutionary data into SDMs[5,21,22], and in particular the

need to consider behaviour in addition to physiology when predicting range shifts[23].

While the results presented here showcase the importance of creating more robust and

informed SDMs, they also highlight a number of aspects that will benefit from additional

consideration when interpreting these models. The existing estimate of CTmin evolution in

sticklebacks considered the change in phenotypic variation across generations[4], rather than

evolution at underlying loci. As such, it is likely that some proportion of the observed phenotypic

change was due to plastic responses. In our PVE model, we take a conservative approach by

restricting phenotypic evolution of CTmin to only occur through heritable change via the locus

shown to be associated with the trait. The efficiency of translating the selection acting on a trait

into evolutionary response across generations can depend on the genetic architecture of the

trait[24]. The large effect loci that we identified here are consistent with expectations from theory

suggesting that prolonged bouts of adaptation with gene flow (as expected in this system[24–26])

should favour architectures characterized by fewer, larger effect, more tightly linked alleles[27–29].

However, it should be noted that the effects of the two QTL identified here are likely

overestimated and other loci might have gone undetected (*sensu* the Beavis Effect[30]). The joint

action of plastic effects and evolution at undetected loci might therefore result in range

distributions that are more similar to those predicted in our 'evolution' models.

Climate change is leading to an increase in the frequency of extreme temperature

events[14,31], including both extreme heat and extreme cold[32–34], which could drive selection on

both CTmin and CTmax[13,35–37]. Our models reveal that the evolution of cold tolerance can have a

significant impact on predicted range distributions despite most end-of-century climate change

scenarios involving an overall warmer, not cooler world. This counterintuitive result occurs

because climate change opens up newly available thermal niche space in waters north of the

current day geographic range[20], and the evolution of CTmin extends this range expansion further

still. Northward range expansion with climate change due to increasing habitat availability has

also been documented in birds[38–41], plants[42], other fishes[43–45], and pest species (such as ticks[46–48]

and mountain pine beetle[49,50]), as well as in large scale analyses of diverse taxa assessing the

'fingerprints' of climate change impacts[51,52]. However, it is likely that evolution of CTmax will

also play a role in responses to environmental change[53–55]. Although we have no empirical

estimates of CTmax evolution, it is interesting to explore how distributions would shift if we

observed the same rate of evolution in this trait as in CTmin. Using the same rate of haldanes and

incorporating our observed PVE for the locus associated with CTmax, we find that geographic

ranges predicted for the end-of-century species distributions also increased by over 7-fold (RCP

4.5: 1,227,002 km$^2$ increase; RCP 8.5: 1,334,314 km$^2$ increase; Fig. S4). Further investigations to

test the empirical rate of evolution of thermal behaviour, physiology and the molecular

underpinnings of these key traits would be well served by assessing additional samples along the

latitudinal gradient inhabited by stickleback to gain a more detailed understanding of these

temperature-associated traits over a wider environmental range.

Collectively, the inclusion of thermal traits and their evolution alters the projected ranges

of threespine stickleback, with a substantial increase in the predicted area that the species will

occupy under climate change forecasts. Many traits are evolving in response to climate change[56–

59] and SDMs that do not take trait data (and trait evolution) into account could provide inaccurate

predictions about future species distributions under climate change[3] - an issue of particular

concern for species at risk and pest species undergoing range expansion[60–62]. Our results provide

a framework for addressing this problem, which will have critical implications for the application

of these models in policy and resource management, and the protection of biodiversity in a

changing climate.

**Materials and Methods**

<u>Sample collection and husbandry</u>

We collected adult *Gasterosteus aculeatus* (Fig. 1a) from two marine populations

(Bamfield, M1, 48°49'12.69"N 125° 8'57.90"W; Garden Bay Lagoon, M2, 49°37'52.84"N 124°

1'49.26"W) and two freshwater populations (Hotel Lake, FW1, 49°38'26.94"N 124° 3'0.69"W;

Klein Lake, FW2, 49°43'32.47"N 123°58'7.83"W) in southwestern British Columbia (Fig. 1b).

Individuals were maintained in a flow-through system and photoperiod that mimicked the source

populations during collection periods before transport. We transported the fish to our aquatics

facility in the Life and Environmental Sciences Animal Resources Centre at the University of

Calgary, where we separated the fish into population-specific 113 L glass aquaria at a density of

approximately 20 fish per aquarium. We acclimated marine individuals to freshwater salinity

over one week and maintained fish in a common environment (salinity of 4-6 ppt, water

temperature of $15 \pm 2$ °C, and a photoperiod of 16L:8D). Individuals were allowed to acclimate

for at least 2 weeks before experiments (1 week for stress reduction post-transfer, 1 week for

common garden environment acclimation and salinity ramp). Each common garden aquarium

was on a closed system with individual filters, air stones, and water supply. We fed all adult fish

*ad libitum* once per day with thawed bloodworms (Hikari Bio-Pure Frozen Bloodworms). All

collections and transfers were approved by the Department of Fisheries and Oceans (marine

collections and transfers), the Ministry of Forests, Lands, and Natural Resource Operations

(freshwater collections), and the Huu-ay-aht First Nations (marine collections).

<u>Crossing design for marine and freshwater F1 families</u>

We collected eggs from females and fertilized the eggs with extracted testes from

euthanized males. We transferred the fertilized egg mass to a mesh-bottomed egg incubator

suspended in a 37 L aquarium for hatching. Each hatching aquarium was maintained with a

single air stone and a filter. Once hatched, we reared the larval fish in 37 L hatching aquaria until

they reached a total length (TL) of approximately 1 cm, after which we split the families into

family-specific 113 L aquaria to maintain suitable densities. We fed the larval fish *ad libitum*

twice daily with live *Artemia spp.* nauplii, and then gradually transitioned the diet to chopped,

thawed bloodworms (Hikari Bio-Pure Frozen Bloodworms) *ad libitum* once daily as they

reached approximately 2 cm TL. The F1 families were maintained in a common garden

environment identical to that of the F0 populations. We produced one F1 family for each

population (M1_F1, M2_F2, FW1_F1, and FW2_F1).

<u>Crossing design for hybrid mapping families</u>

To generate genetically heterogeneous marine-freshwater F1 families from wild F0

parents, we collected eggs from marine females and fertilized the eggs with extracted testes from

euthanized freshwater males. Egg masses were hatched, and juveniles were reared, as detailed

above. Overall, we produced one F1 family of M1xFW1 hybrids (hereafter referred to as H1_F1)

and three F1 families of M1xFW2 hybrids (hereafter referred to as H2_F1). The hybrid F1

families were maintained in a common garden environment identical to that of the F0

populations. To generate F2 families for linkage map construction, we crossed individuals from

the same F1 family with the same methodology used to generate the F1 families. Overall, we

produced one F2 family of H1xH1 hybrids (referred to as H1_F2) and three families of H2xH2

hybrids (referred to as H2_F2_1, H2_F2_2, and H2_F2_3). All F2 individuals were raised as

described above in a common garden environment identical to that of the F0 and F1 individuals

to ensure consistent history and use for QTL mapping.

Thermal tolerance experiments

To assess the lower and upper limits of physiological thermal tolerance, we conducted

standard critical thermal minimum (CTmin) and maximum (CTmax) experiments on adult

fish[4,63,64]. At these sublethal limits, the fish experiences a loss of equilibrium (LOE) at which

they lose the ability to escape conditions that would ultimately lead to their death in nature[65]. Our

experimental tank held 1000 mL glass beakers aerated individually to prevent thermal

stratification. Before each experiment, individuals were fasted for 24 hours. After a 15-minute

acclimation to the experimental apparatus in the individual beakers, we cooled or heated the

water (for CTmin or CTmax, respectively) at a rate of approximately 0.33 °C min$^{-1}$. We assessed

wild F0 individuals ($n_{M1}$ = 32, $n_{M2}$ = 14, $n_{FW1}$ = 15, $n_{FW2}$ = 16, $N$ = 77; Fig. 1c) and lab raised F1

($n_{M1\_F1}$ = 13, $n_{M2\_F1}$ = 15, $n_{FW1\_F1}$ = 15, $n_{FW2\_F1}$ = 15, $N$ = 58; Fig. S1) and F2 individuals ($n_{H1\_F2}$

= 28, $n_{H2\_F2\_1}$ = 36, $n_{H2\_F2\_2}$ = 21, $n_{H2\_F2\_3}$ = 17, $N$ = 102; Fig. S2). All individuals were

assessed for CTmin, allowed to recover for at least three days, then assessed for CTmax to keep

thermal stress history consistent. The onset of erratic behaviours associated with a behavioural

stress response occurred below 5.0 °C and above 25.0 °C during CTmin and CTmax

experiments, respectively. Normal behaviour was observed between 5.0 °C and 25.0 °C, whereas

outside of those temperatures, individuals gradually exhibited more extreme stress responses

(*e.g.,* increased gilling rate, erratic movement, muscle spasms, listing, as outlined by the

Canadian Council of Animal Care guidelines) until reaching LOE and the inability of an

individual to right itself (the experimental endpoint)[4,63,64]. At the time of data collection for

thermal trait experiments, all individuals were adults.

Isolation and characterization of single nucleotide polymorphisms (SNPs)

Genomic DNA was extracted from caudal fin tissue using a phenol-chloroform-based

protocol. We digested tissues overnight in digestion buffer and proteinase K at 55 °C, then

performed multiple phenol-chloroform and ethanol washes to isolate the DNA. We assessed the

quantity of the extracted DNA using the Quant-iT PicoGreen dsDNA assay kit (ThermoFisher

Scientific, Waltham, MA, USA) and Synergy HT plate reader with the Gen5 associated software

(BioTek, Winooski, VT, USA). We prepared restriction site-associated DNA (RAD)

libraries(Peterson et al. 2012) using *MluCl* and *NlaIII* restriction (New England Biolabs,

Ipswich, MA, USA), ligation of individual barcodes, and pooling of 48 individuals per library at

equimolar concentrations. We performed a final PCR to amplify DNA and add library-specific

indices to allow for pooling of multiple libraries. We sequenced three libraries at McGill

University and Génome Québec Innovation Center on one lane of Illumina HiSeq 4000 (Illumina

Inc., San Diego, CA, USA).

Assembly of genetic linkage map

After barcode demultiplexing and filtering out low quality reads in STACKS[66], we

removed PCR duplicates from the raw sequences and aligned to the *G. aculeatus* reference

genome[17] using the Burrows-Wheeler transform[67]. Individual libraries were concatenated and

filtered[68] using *vcftools* v3.0[69] and then split into chromosome-specific VCF files to assemble the

linkage maps chromosome by chromosome. We assigned markers to a linkage group with an

initial LOD score of 3 after filtering out markers that showed high levels of segregation

distortion and missing observations (> 20% missing data) in Lep-MAP3[70]. Unassigned markers

were subsequently added to the existing linkage group at a LOD score of 3 and a size limit of 5

markers per linkage group. We ordered the markers using a minimum posterior value of 0.001

and collapsed multiple markers when the probability difference between markers was $< 0.01$[70].

The final linkage map was subset for use in R[71] with a custom Python script to visualize the

linkage map and to generate a list of informative SNPs to use in subsequent analyses with the *qtl*

v1.44-9[72] and *qtlTools* v1.2.0[73] packages. Linkage maps were visualized using

*LinkageMapView*[74] in R[71]. The final linkage maps were similar across families in number of

markers, length, and spacing between markers, though the H1_F2 map did have a higher density

of markers (Table S1).

Quantitative trait loci (QTL) mapping

We analysed families separately with the same methodology to assess the presence of

QTL associated with the thermal traits. We calculated conditional genotype probabilities using

hidden Markov model technology and simulated genotypes based on the observed marker data

(allowing for possible genotyping errors at a level of 0.0001 using a Kosambi mapping function

with a fixed step width) prior to running genome scans with a single QTL model[75,76]. We

determined the logarithm of the odds (LOD) score significance thresholds for each trait through

permutation tests for each family (100,000 permutations) (Fig. 2a). We pulled significant QTL

above the genome-wide significance threshold ($\alpha = 0.05$[77]), calculated confidence intervals of

QTL location based on nearby markers, and estimated the percent variance explained by each

QTL peak marker. We identified two QTL on linkage group 4 (which corresponds to

chromosome 4 of the BROAD assembly[17]) associated with CTmin and CTmax (Fig. 2b).

Environmental variables and species distribution models (SDMs)

We compiled environmental data widely used in the construction of SDMs to estimate

suitable habitat in both present day and end-of-century forecasts[78], including bathymetry, sea ice

extent and concentration, salinity, and sea surface temperature. We used 2014 data as our

baseline year to match the forecasting baseline of the Fifth Assessment Report[14]. We assumed a

suitable habitat range for this species in the Pacific Northwest to consist of coastal areas (where

the water depth is less than 200 m) where sea ice is never present (*i.e.,* no sea ice at the

maximum extent). The salinity tolerance for *G. aculeatus* is very wide[79,80] and salinity was not

limiting in any of the habitat[81], therefore salinity was not included in the final present day or

forecasted models. We obtained bathymetry data from the General Bathymetric Chart of the

Oceans (GEBCO) of the British Oceanographic Data Centre[82], and maximum sea ice extent data

from the Multisensory Analyzed Sea Ice Extent – Northern Hemisphere (MASIE-NH) product[83].

We obtained maximum and minimum daily mean sea surface temperature (SST)[84]. Sea surface

temperature was used as a proxy for water temperature. Stickleback thermal trait data were used

to set the limits of the distribution within the possible area delineated by sea ice free water of a

suitable depth (Table S2). The thermal trait measurements were all based on our experimental

findings reported here.

In the end-of-century forecast for suitable habitat, we assumed bathymetry to be

consistent with the modern scenario. However, the Arctic Ocean is predicted to be

predominantly free of sea ice in the summer by the end of the century[85], with significant end-of-

century reductions in winter/spring sea ice concentration (reduced to a concentration of 0.1 at the

Seward Peninsula[85]), so we conservatively set the maximum northern extent of the suitable

habitat to be 65°35' N, which corresponds to the western tip of the Seward Peninsula (near

Wales, AK). The extent of sea ice was kept consistent between scenarios to control for area in

calculations of range expansion. The water temperatures were increased based on projections for

large marine ecosystems of Northern Oceans from global climate models[14,20]. Maps were created

in R[71] using the packages *raster* v. 3.3-13[86] and *rgeos* v. 0.5-3[87].

We incorporated the experimental data from the critical thermal minimum and maximum

trials on the wild marine populations (Fig. 1c) to understand how trait inclusion may affect range

projections under climate change. These trait-defined envelopes were overlain on the suitable

habitat background to delineate projected presence based on thermal traits in both current day

and IPCC predicted RCPs 4.5 and 8.5. The trait values were kept constant (*i.e.,* not changed) in

the 'no evolution' projections, but in the 'evolution' projections, we allowed CTmin to evolve an

improvement of 2.5 °C (*i.e.,* 2.5 °C lower than CTmin boundary in the 'no evolution' projection)

by the end of the century based on a rate of 0.63 *haldanes* from a selection experiment

previously conducted on populations from this same genetic cluster[4]. The 'evolution' model

assumes whole-organism tolerance evolution with selection acting on 100% of the loci affecting

CTmin evolution. Therefore, to account for the observed genetic architecture of a single, large

effect locus associated with CTmin, we next considered a 'PVE' model, where CTmin was

allowed to evolve to only 54% of the total estimated trait value from the 'evolution' model (*i.e.,*

2.5 °C * 0.54). However, a notable restriction in the evolution of CTmin for both models

('evolution' and 'PVE') was a hard boundary drawn at 0 °C under the assumption that

population persistence in a sub-zero environment would require many additional adaptations

alongside CTmin improvement (*e.g.,* extreme adaptations observed in Antarctic notothenioid

fishes[88–90]).

To quantify the differences in estimated suitable habitat under current day and end-of-

century conditions, we compared areas for each warming scenario to the equivalent scenario

under current conditions. Similarly, to compare the differences in evolutionary scenarios, the

area of each end-of-century evolutionary trajectory was compared to either the contrasting RCP

projection or adjusted PVE projection. For these comparisons, we used North Pole Lambert

azimuthal equal area projection for all maps, and georeferenced to known landmarks in ArcGIS

v10.8[91] to calculate area from the maps generate in R[71] (conversion ratio of 7873.42).

The datasets generated and analysed during the current study are available from the

corresponding author upon reasonable request. The annotated code, including all parameter

thresholds, for the above QTL analyses and SDM construction is publicly available on Github

(github.com/sjswuitchik/gasAcu_qtl_sdm).

**References**

1.      Kearney, M. R., Wintle, B. A. & Porter, W. P. Correlative and mechanistic models of
        species distribution provide congruent forecasts under climate change. *Conserv. Lett.* **3**,
        203–213 (2010).

2.      Kearney, M. & Porter, W. Mechanistic niche modelling: Combining physiological and
        spatial data to predict species' ranges. *Ecol. Lett.* **12**, 334–350 (2009).

3.      Bush, A. *et al.* Incorporating evolutionary adaptation in species distribution modelling
        reduces projected vulnerability to climate change. *Ecol. Lett.* **19**, 1468–1478 (2016).

4.      Barrett, R. D. H. *et al.* Rapid evolution of cold tolerance in stickleback. *Proc. R. Soc. B*
        **278**, 233–238 (2011).

5.      Evans, T. G., Diamond, S. E. & Kelly, M. W. Mechanistic species distribution modelling
        as a link between physiology and conservation. *Conserv. Physiol.* **3**, 1–16 (2015).

6.      Pearce, J. & Lindenmayer, D. Bioclimatic analysis to enhance reintroduction biology of
        the endangered helmeted honeyeater (Lichenostomus melanops cassidix) in southeastern

Australia. *Restor. Ecol.* **6**, 238–243 (1998).

7.    Araújo, M. B., Cabeza, M., Thuiller, W., Hannah, L. & Williams, P. H. Would climate

change drive species out of reserves? An assessment of existing reserve-selection

methods. *Glob. Chang. Biol.* **10**, 1618–1626 (2004).

8.    Hochachka, P. W. & Somero, G. N. Mechanism and process in physiological evolution.

*Biochem. Adapt.* **480**, (2002).

9.    Huey, R. B. & Slatkin, M. Costs and benefits of lizard thermoregulation. *Q. Rev. Biol.* **51**,

363–384 (1976).

10.    Coutant, C. C. Thermal preference: when does an asset become a liability? *Environ. Biol.*

*Fishes* **18**, 161–172 (1987).

11.    Huey, R. B. & Kingsolver, J. G. Evolution of thermal sensitivity of ectotherm

performance. *Trends Ecol. Evol.* **4**, 131–135 (1989).

12.    Sexton, J. P., McIntyre, P. J., Angert, A. L. & Rice, K. J. Evolution and Ecology of

Species Range Limits. *Annu. Rev. Ecol. Evol. Syst.* **40**, 415–436 (2009).

13.    Hoffman, A. & Sgrò, C. Climate change and evolutionary adaptation. *Nature* **470**, 479–

485 (2011).

14.    IPCC. *Climate Change 2014: Synthesis Report*. *Fifth Assessment Report of the*

*Intergovernmental Panel on Climate Change* (2014). doi:10.1016/S0022-0248(00)00575-

3

15.    Eyer, P. A., Blumenfeld, A. J. & Vargo, E. L. Sexually antagonistic selection promotes

genetic divergence between males and females in an ant. *Proc. Natl. Acad. Sci. U. S. A.* **116**, 24157–24163 (2019).

16. Hendry, A. P., Peichel, C. L., Matthews, B., Boughman, J. W. & Nosil, P. Stickleback research: The now and the next. *Evol. Ecol. Res.* **15**, 111–141 (2013).

17. Jones, F. C. *et al.* The genomic basis of adaptive evolution in threespine sticklebacks. *Nature* **484**, 55–61 (2012).

18. Barrett, R. D. H. & Hendry, A. P. Evolutionary rescue under environmental change. in *Behavioural responses to a changing world: mechanisms and consequences* 216–233 (Oxford University Press Oxford, UK, 2012).

19. Morris, M. R. J., Bowles, E., Allen, B. E., Jamniczky, H. A. & Rogers, S. M. Contemporary ancestor? Adaptive divergence from standing genetic variation in Pacific marine threespine stickleback. *BMC Evol. Biol.* **18**, 1–21 (2018).

20. Alexander, M. A. *et al.* Projected sea surface temperatures over the 21st century: Changes in the mean, variability and extremes for large marine ecosystem regions of Northern Oceans. *Elementa* **6**, (2018).

21. Buckley, L. *et al.* Can mechanism inform species' distribution models? *Ecol. Lett.* **13**, 1041–1054 (2010).

22. Lyon, N. J., Debinski, D. M. & Rangwala, I. Evaluating the Utility of Species Distribution Models in Informing Climate Change-Resilient Grassland Restoration Strategy. *Front. Ecol. Evol.* **7**, 1–8 (2019).

398    23.    Sunday, J. M., Bates, A. E. & Dulvy, N. K. Thermal tolerance and the global

399           redistribution of animals. *Nat. Clim. Chang.* **2**, 686–690 (2012).

400    24.    Rogers, S. M. *et al.* Genetic Signature of Adaptive Peak Shift in Threespine Stickleback.

401           *Evolution (N. Y)*. 2439–2451 (2012). doi:10.5061/dryad.6jj614kh

402    25.    Schluter, D., Marchinko, K. B., Barrett, R. D. H. & Rogers, S. M. Natural selection and

403           the genetics of adaptation in threespine stickleback. *Philos. Trans. R. Soc. B Biol. Sci.* **365**,

404           2479–2486 (2010).

405    26.    Jones, F. C. *et al.* A genome-wide SNP genotyping array reveals patterns of global and

406           repeated species-pair divergence in sticklebacks. *Curr. Biol.* **22**, 83–90 (2012).

407    27.    Yeaman, S. & Otto, S. P. Establishment and maintenance of adaptive genetic divergence

408           under migration, selection, and drift. *Evolution (N. Y)*. **65**, 2123–2129 (2011).

409    28.    Yeaman, S. & Whitlock, M. C. The genetic architecture of adaptation under migration-

410           selection balance. *Evolution (N. Y)*. **65**, 1897–1911 (2011).

411    29.    Via, S., Conte, G., Mason-Foley, C. & Mills, K. Localizing FST outliers on a QTL map

412           reveals evidence for large genomic regions of reduced gene exchange during speciation-

413           with-gene-flow. *Mol. Ecol.* **21**, 5546–5560 (2012).

414    30.    Beavis, W. The power and deceit of QTL experiments: lessons from comparative QTL

415           studies. in *49th Annual Corn & Sorghum Research Conference* 250–266 (1994).

416    31.    Stott, P. How climate change affects extreme weather events. *Science (80-. )*. **352**, 1517–

417           1518 (2016).

32.   Herring, S., Hoerling, M., Kossing, J., Peterson, T. & Stott, P. Explaining extreme events

of 2014 from a climate perspective. **96**, 1–180 (2015).

33.   Herring, S. C. *et al.* Explaining extreme events of 2016 from a climate perspective. *Bull.*

*Am. Meteorol. Soc.* **99**, S1–S157 (2018).

34.   Herring, S. C., Christidis, N., Hoell, A., Hoerling, M. P. & Stott, P. Explaining extreme

events of 2018 from a climate perspective. *Bull. Am. Meteorol. Soc.* **101**, 1–146 (2020).

35.   Kingsolver, J. G. *et al.* Complex life cycles and the responses of insects to climate change.

*Integr. Comp. Biol.* **51**, 719–732 (2011).

36.   Denny, M. W. & Dowd, W. W. Biophysics, environmental stochasticity, and the evolution

of thermal safety margins in intertidal limpets. *J. Exp. Biol.* **215**, 934–947 (2012).

37.   Buckley, L. B. & Huey, R. B. How extreme temperatures impact organisms and the

evolution of their thermal tolerance. *Integr. Comp. Biol.* **56**, 98–109 (2016).

38.   Tingley, M. W., Monahan, W. B., Beissinger, S. R. & Moritz, C. Birds track their

Grinnellian niche through a century of climate change. *Proc. Natl. Acad. Sci. U. S. A.* **106**,

19637–19643 (2009).

39.   Melles, S. J., Fortin, M. J., Lindsay, K. & Badzinski, D. Expanding northward: Influence

of climate change, forest connectivity, and population processes on a threatened species'

range shift. *Glob. Chang. Biol.* **17**, 17–31 (2011).

40.   Tombre, I. M., Oudman, T., Shimmings, P., Griffin, L. & Prop, J. Northward range

expansion in spring-staging barnacle geese is a response to climate change and population

growth, mediated by individual experience. *Glob. Chang. Biol.* **25**, 3680–3693 (2019).

41.   Rushing, C. S., Andrew Royle, J., Ziolkowski, D. J. & Pardieck, K. L. Migratory behavior

and winter geography drive differential range shifts of eastern birds in response to recent

climate change. *Proc. Natl. Acad. Sci. U. S. A.* **117**, 12897–12903 (2020).

42.   D'Andrea, L. *et al.* Climate change, anthropogenic disturbance and the northward range

expansion of Lactuca serriola (Asteraceae). *J. Biogeogr.* **36**, 1573–1587 (2009).

43.   Fossheim, M. *et al.* Recent warming leads to a rapid borealization of fish communities in

the Arctic. *Nat. Clim. Chang.* **5**, 673–677 (2015).

44.   Yapıcı, S., Bilge, G. & Filiz, H. Northwards range expansion of Sparisoma cretense

(Linnaeus, 1758) in the Turkish Aegean Sea. *J. Aquac. Eng. Fish. Res.* 201–207 (2016).

doi:10.3153/jaefr16022

45.   Spies, I. *et al.* Genetic evidence of a northward range expansion in the eastern Bering Sea

stock of Pacific cod. *Evol. Appl.* **13**, 362–375 (2020).

46.   Ogden, N. H. *et al.* Climate change and the potential for range expansion of the Lyme

disease vector Ixodes scapularis in Canada. *Int. J. Parasitol.* **36**, 63–70 (2006).

47.   Clow, K. M. *et al.* Northward range expansion of Ixodes scapularis evident over a short

timescale in Ontario, Canada. *PLoS One* **12**, 1–15 (2017).

48.   Sagurova, I. *et al.* Predicted northward expansion of the geographic range of the tick

vector amblyomma americanum in North America under future climate conditions.

*Environ. Health Perspect.* **127**, 1–14 (2019).

49.    Kurz, W. A. *et al.* Mountain pine beetle and forest carbon feedback to climate change.

459           *Nature* **452**, 987–990 (2008).

460    50.    Sambaraju, K. R., Carroll, A. L. & Aukema, B. H. Multiyear weather anomalies

461           associated with range shifts by the mountain pine beetle preceding large epidemics. *For.*

462           *Ecol. Manage.* **438**, 86–95 (2019).

463    51.    Parmesan, C. & Yohe, G. A globally coherent fingerprint of climate change impacts

464           across natural systems. **421**, 37–42 (2003).

465    52.    Platts, P. J. *et al.* Habitat availability explains variation in climate-driven range shifts

466           across multiple taxonomic groups. *Sci. Rep.* **9**, 1–10 (2019).

467    53.    Geerts, A. N. *et al.* Rapid evolution of thermal tolerance in the water flea Daphnia. *Nat.*

468           *Clim. Chang.* **5**, 665–668 (2015).

469    54.    Bozinovic, F., Medina, N. R., Alruiz, J. M., Cavieres, G. & Sabat, P. Thermal tolerance

470           and survival responses to scenarios of experimental climatic change: changing thermal

471           variability reduces the heat and cold tolerance in a fly. *J. Comp. Physiol. B Biochem. Syst.*

472           *Environ. Physiol.* **186**, 581–587 (2016).

473    55.    Cuenca Cambronero, M., Beasley, J., Kissane, S. & Orsini, L. Evolution of thermal

474           tolerance in multifarious environments. *Mol. Ecol.* **27**, 4529–4541 (2018).

475    56.    Eliason, E. J. *et al.* Differences in thermal tolerance among sockeye salmon populations.

476           *Science (80-. ).* **332**, 109–112 (2011).

477    57.    Hovel, R. A., Carlson, S. M. & Quinn, T. P. Climate change alters the reproductive

478              phenology and investment of a lacustrine fish, the three-spine stickleback. *Glob. Chang.*

479              *Biol.* 1–13 (2016). doi:10.1111/gcb.13531

480   58.   Gómez-Ruiz, E. P. & Lacher, T. E. Climate change, range shifts, and the disruption of a

481              pollinator-plant complex. *Sci. Rep.* **9**, 1–10 (2019).

482   59.   Horton, K. G. *et al.* Phenology of nocturnal avian migration has shifted at the continental

483              scale. *Nat. Clim. Chang.* **10**, 63–68 (2020).

484   60.   Cullingham, C. I. *et al.* Mountain pine beetle host-range expansion threatens the boreal

485              forest. *Mol. Ecol.* **20**, 2157–2171 (2011).

486   61.   McLeod, D. J., Hallegraeff, G. M., Hosie, G. W. & Richardson, A. J. Climate-driven

487              range expansion of the red-tide dinoflagellate Noctiluca scintillans into the Southern

488              Ocean. *J. Plankton Res.* **34**, 332–337 (2012).

489   62.   Bebber, D. P. Range-Expanding Pests and Pathogens in a Warming World. *Annu. Rev.*

490              *Phytopathol.* **53**, 335–356 (2015).

491   63.   Hutchison, V. H. Comparative Biology Critical Thermal Maxima in Salamanders. *Physiol.*

492              *Zool.* **34**, 92–125 (1961).

493   64.   Fangue, N. A., Hofmeister, M. & Schulte, P. M. Intraspecific variation in thermal

494              tolerance and heat shock protein gene expression in common killifish, Fundulus

495              heteroclitus. *J. Exp. Biol.* **209**, 2859–2872 (2006).

496   65.   Beitinger, T., Bennett, W. & McCauley, R. Temperature tolerances of North American

497              freshwater fishes exposed to dynamic changes in temperature. *Environ. Biol. Fishes* **58**,

498    237–275 (2000).

499    66.    Catchen, J., Hohenlohe, P. A., Bassham, S., Amores, A. & Cresko, W. A. Stacks: An

500           analysis tool set for population genomics. *Mol. Ecol.* **22**, 3124–3140 (2013).

501    67.    Li, H. & Durbin, R. Fast and accurate long-read alignment with Burrows-Wheeler

502           transform. *Bioinformatics* **26**, 589–595 (2010).

503    68.    Puritz, J. B., Hollenbeck, C. M. & Gold, J. R. *dDocent* : a RADseq, variant-calling

504           pipeline designed for population genomics of non-model organisms. *PeerJ* **2**, e431 (2014).

505    69.    Danecek, P. *et al.* The variant call format and VCFtools. *Bioinformatics* **27**, 2156–2158

506           (2011).

507    70.    Rastas, P. Lep-MAP3: Robust linkage mapping even for low-coverage whole genome

508           sequencing data. *Bioinformatics* **33**, 3726–3732 (2017).

509    71.    R Core Team. R: A language and environment for statistical computing. (2019).

510    72.    Broman, K. W., Wu, H., Sen, Ś. & Churchill, G. A. R/qtl: QTL mapping in experimental

511           crosses. *Bioinformatics* **19**, 889–890 (2003).

512    73.    Lovell, J. qtlTools. (2019).

513    74.    Ouellette, L. A., Reid, R. W., Blanchard, S. G. & Brouwer, C. R. LinkageMapView-

514           rendering high-resolution linkage and QTL maps. *Bioinformatics* **34**, 306–307 (2018).

515    75.    Broman, K. W. & Sen, S. *A Guide to QTL Mapping with R/qtl*. **46**, (Springer, 2009).

516    76.    Arends, D., Prins, P., Broman, K. W. & Jansen, R. C. *Tutorial-Multiple-QTL Mapping*

517    *(MQM) Analysis for R/qtl. http://www.rqtl.org/tutorials/MQM-tour.pdf* (2014).

518    77.    Greenwood, A. K. *et al.* The genetic basis of divergent pigment patterns in juvenile

519            threespine sticklebacks. *Heredity (Edinb).* **107**, 155–166 (2011).

520    78.    Wiens, J., Stralberg, D., Jongsomjit, D., Howell, C. & Snyder, M. Niches, models, and

521            climate change: Assessing the assumptions and uncertainties. *Proc. Natl. Acad. Sci.* **106**,

522            19729–19736 (2009).

523    79.    Bayly, I. A. E. Salinity Tolerance and Osmotic Behavior of Animals in Athalassic Saline

524            and Marine Hypersaline Waters. *Annu. Rev. Ecol. Syst.* **3**, 233–268 (2003).

525    80.    Divino, J. N. *et al.* Osmoregulatory physiology and rapid evolution of salinity tolerance in

526            threespine stickleback recently introduced to fresh water. *Evol. Ecol. Res.* **17**, 179–201

527            (2016).

528    81.    Zweng, M. M. *et al. World Ocean Atlas 2013, Volume 2: Salinity. NOAA Atlas NESDIS*

529            *74* **2**, (2013).

530    82.    Weatherall, P. *et al.* A new digital bathymetric model of the world's oceans. *Earth Sp. Sci.*

531            **2**, 331–345 (2015).

532    83.    Fetterer, F., Savoie, M., Helfrich, S. & Clemene-Colon, P. U.S. National Ice Center and

533            National Snow and Ice Data Center. *Multisensor Analyzed Sea Ice Extent - Northern*

534            *Hemisphere (MASIE-NH)* (2010). doi:https://doi.org/10.7265/N5GT5K3K.

535    84.    Reynolds, R. *et al.* Daily High-Resolution-Blended Analyses for Sea Surface

536            Temperature. *J. Clim.* **20**, 5473–5496 (2007).

537    85.    Johannessen, O. *et al.* Arctic climate change: observed and modelled temperature and sea-
538            ice variability. *Tellus A Dyn. Meteorol. Oceanogr.* **56**, 328–341 (2004).

539    86.    Hijmans, R. J. *et al.* raster: Geographic data analysis and modelling. (2020).

540    87.    Roger, A., Stuetz, R., Ove, K., Giraudoux, P. & Santilli, S. rgeos: Interface to geometry
541            engine - open. (2020).

542    88.    Detrich, H. W., Parker, S. K., Williams, J., Nogales, E. & Downing, K. H. Cold adaptation
543            of microtubule assembly and dynamics. Structural interpretation of primary sequence
544            changes present in the α- and β-tubulins of antarctic fishes. *J. Biol. Chem.* **275**, 37038–
545            37047 (2000).

546    89.    Cheng, C. H. C. & Detrich, H. W. Molecular ecophysiology of Antarctic notothenioid
547            fishes. *Philos. Trans. R. Soc. B Biol. Sci.* **362**, 2215–2232 (2007).

548    90.    Shin, S. C. hu. *et al.* The genome sequence of the Antarctic bullhead notothen reveals
549            evolutionary adaptations to a cold environment. *Genome Biol.* **15**, 468 (2014).

91.    Environmental Systems Research Institute. ArcGIS Desktop: Release 10.8. (2017).

Acknowledgments

The authors acknowledge that the species collections took place on Huu-ay-aht and Sechelt First

Nations traditional territories and are grateful for the opportunity to conduct their research in

protected and sacred areas. We would like to thank the Bamfield Marine Sciences Centre for the

resources required to conduct this research, Sam Owens for the photo in Figure 1a, Peter Peller

for creating Figure 1b, as well as Daniel Wuitchik, Sam Yeaman, Jennifer Sunday, and Patrick

Nosil for feedback on the manuscript.

**Author Contributions**

This study was designed by SJSW, RDHB, and SMR; fish husbandry and breeding by SJSW and

TNB; experimental data collection by SJSW; DNA sequencing and initial processing by AP;

bioinformatic and QTL analyses by SJSW; species distribution modelling by SJSW and SM; the

manuscript was written by SJSW, RDHB, and SMR, with input from authors; the study was

funded by HAJ, RDHB, and SMR.

Competing Interests: The authors declare no competing interests.

**Figures and Tables**

**Figure 1.** a) Adult threespine stickleback (*Gasterosteus aculeatus*) from a single genetic cluster
were sampled from b) two marine and two freshwater populations in the Canadian Pacific
Northwest. These populations were assayed for c) critical thermal minima and maxima. Thermal
trait values for marine populations (M1 and M2) were incorporated into the species distribution
models, while marine and freshwater populations (M1, FW1, and FW2) were used to generate F1
and F2 generations for linkage map construction and quantitative trait loci (QTL) analyses.

**Figure 2.** a) Quantitative trait loci (QTL) scan of linkage group 4 with trait-specific significance
thresholds for LOD scores, showing a significant LOD peak for CTmin (dark purple) and
CTmax (light blue) with b) an inset of linkage group 4 highlighting the position of the significant
QTL for upper and lower thermal tolerances (CTmax and CTmin, respectively).

**Figure 3.** Changes in the distribution of marine threespine stickleback (*Gasterosteus aculeatus*) as a result of incorporating thermal traits in a) current day environmental conditions and under IPCC end-of-century projections RCP 4.5 and 8.5 without trait evolution ('no evolution' model, b & c, orange box) and with trait evolution ('evolution' model, d & e, red box)**.**

**Supplementary Figures and Tables**

**Figure S1.** Critical thermal minimum (CTmin) and maximum (CTmax) measurements for adult threespine stickleback (*Gasterosteus aculeatus*) from pure F1 marine (M*F1) and freshwater (FW*F1) families raised in a common garden under a constant thermal environment.

**Figure S2.** Critical thermal minimum (CTmin) and maximum (CTmax) measurements for adult threespine stickleback (*Gasterosteus aculeatus*) from hybrid marine-freshwater F2 families raised in a common garden under a constant thermal environment.

**Figure S3.** Changes in the distribution of marine threespine stickleback (*Gasterosteus aculeatus*) as a result of incorporating thermal traits under IPCC end-of-century projections RCP 4.5 and 8.5 with trait evolution constrained by the underlying genetic architecture of critical thermal minimum (CTmin) as determined from hybrid F2 mapping families.

**Figure S4.** Changes in the distribution of marine threespine stickleback (*Gasterosteus aculeatus*) as a result of incorporating thermal traits under IPCC end-of-century projections RCP 4.5 and 8.5 with trait evolution constrained by the underlying genetic architecture of critical thermal minimum (CTmin) and critical thermal maximum (CTmax) as determined from hybrid F2 mapping families.

**Table S1.** Summaries of family-specific linkage maps constructed for quantitative trait loci (QTL) analyses.

| Map | $n_{markers}$ | Length (cM) | Avg. Max. Spacing |
|-----|---------|-------------|-------------------|
| KL1 | 2139 | 1370.4 | 7.6 |
| KL2 | 1558 | 1311 | 8.5 |
| KL3 | 1964 | 1359.5 | 7.9 |
| HL | 5247 | 1621.1 | 11.8 |

**Table S2.** Thermal trait data from *Gasterosteus aculeatus* used to inform the species distribution envelopes (rows) in the varied evolutionary scenarios (columns) projected for end-of-century conditions.

| | Current day | No evolution | Adjusted PVE | Evolution |
|---|---|---|---|---|
| Normal Behaviour | (5 - 25) | (5 - 25) | (3.6 - 25) | (2.5 - 25) |
| Within Physiological Limits | (0.9 - 31.9) | (0.9 - 31.9) | (0 - 31.9) | (0 - 31.9) |
| Outside Physiological Limits | $31.9 > x < 0.9$ | $31.9 > x < 0.9$ | $31.9 > x < 0$ | $31.9 > x < 0$ |