Versatile Cell-Based Assay for Measuring Base Excision Repair of DNA Alkylation Damage

Yong Li[1], Peng Mao[2,6], Evelina Y. Basenko[5,7], Zachary Lewis[3,4,5], Michael Smerdon[2], Wioletta Czaja[*][1,2,8]

[1] The Hormel Institute, University of Minnesota, Austin, Minnesota, 55912, USA. (current address)

[2] School of Molecular Biosciences, Washington State University, Pullman, Washington 99164, USA.

[3] Department of Microbiology, University of Georgia, Athens, Georgia, 30602, USA.

[4] Department of Plant Biology, University of Georgia, Athens, GA, 30602, USA.

[5] Department of Genetics, University of Georgia, Athens, GA, 30602, USA.

[6] Comprehensive Cancer Center, University of New Mexico, Albuquerque, NM, 87131, USA. (current address)

[7] Department of Biochemistry and Systems Biology, University of Liverpool, Liverpool L69 3BX, United Kingdom. (current address)

[8] Department of Biochemistry and Molecular Biology, University of Georgia, Athens, Georgia, 30602, USA.

[*]Address for correspondence

Wioletta Czaja
The Hormel Institute
University of Minnesota
Austin, Minnesota, 55912, USA
Email: wczaja@umn.edu

Abstract

DNA alkylation damage induced by environmental carcinogens, chemotherapy drugs, or endogenous metabolites plays a central role in mutagenesis, carcinogenesis, and cancer therapy. Base excision repair (BER) is a conserved, front line DNA repair pathway that removes alkylation damage from DNA. The capacity of BER to repair DNA alkylation varies markedly between different cell types and tissues, which correlates with cancer risk and cellular responses to alkylation chemotherapy. The ability to measure cellular rates of alkylation damage repair by the BER pathway is critically important for better understanding of the fundamental processes involved in carcinogenesis, and also to advance development of new therapeutic strategies. Methods for assessing the rates of alkylation damage and repair, especially in human cells, are limited, prone to significant variability due to the unstable nature of some of the alkyl adducts, and often rely on indirect measurements of BER activity. Here, we report a highly reproducible and quantitative, cell-based assay, named alk-BER (alkylation Base Excision Repair) for measuring rates of BER following alkylation DNA damage. The alk-BER assay involves specific detection of methyl DNA adducts (7-methyl guanine and 3-methyl adenine) directly in genomic DNA. The assay has been developed and adapted to measure the activity of BER in fungal model systems and human cell lines. Considering the specificity and conserved nature of BER enzymes, the assay can be adapted to virtually any type of cultured cells. Alk-BER offers a cost efficient and reliable method that can effectively complement existing approaches to advance integrative research on mechanisms of alkylation DNA damage and repair.

Key words: Alkylated DNA damage, alkylating agents, DNA methylation, base excision repair, DNA repair capacity, DNA damage quantitation

1. Introduction

DNA alkylation induced by methylating agents such as environmental carcinogens (e.g., smoke), by-products of cellular metabolism (e.g., methyl group donor S-adenosyl methionine), or chemotherapy drugs (e.g., temozolomide, procarbazine) represents one of the most abundant types of DNA base damage that forms in human cells. Monofunctional alkylating agents, like methyl methanesulfonate (MMS) or the anticancer drug temozolomide (TMZ) induce formation of N-methyl and O-methyl DNA adducts such as N7-methylguanine (7meG), N3-methyladenine (3meA) and O6-methylguanine (O6meG) [1, 2]. Methyl DNA adducts (MDAs) have cytotoxic and mutagenic properties because of their ability to block gene transcription and interfere with the fidelity of DNA replication. Persistent and inefficiently repaired methyl DNA adducts can induce microsatellite instability, frameshift mutations, and G→A transition mutations, that are commonly found in genes critical for malignant transformation, including the *H-ras* oncogene or *TP53* tumor suppressor gene [3-5]. Despite their carcinogenic properties, DNA alkylating agents, such as dacarbazine, temozolomide and streptozotin, have been used for decades in treating various cancers, including melanoma, glioma, and lymphoma [2, 6, 7]. Therapy with alkylating agents can be effective; however, these agents are extremely toxic and prolonged treatment often leads to chemoresistance and formation of secondary cancers [8, 9]. Human responses to alkylating agents vary considerably between individuals, which highlights the involvement of genetic and epigenetic mechanisms in the modulation of cellular toxicity to alkylating agents [2, 10, 11].

Base excision repair (BER) is the primary pathway involved in the removal of alkylation DNA damage induced by methylating agents [2, 12, 13]. Repair of methyl DNA adducts through the BER pathway is accomplished in four sequential steps, each carried out by a specific group of enzymes [14]. The first step is catalyzed by DNA glycosylases (e.g., AAG), which specifically recognize and bind to a damaged base, and subsequently catalyze cleavage of the glycosidic bond between the damaged base and DNA backbone [15, 16]. This reaction results in the release of the damaged base from DNA, and the formation of abasic AP (apurinic/apyridiminic) sites. The second step involves incision of the DNA backbone 5' upstream of the AP sites by AP endonucleases (e.g., APE1), which results in the formation of single strand DNA breaks (SSBs) [17, 18]. In mammalian cells, AP sites and SSBs are recognized by poly ADP-ribose polymerase 1 (PARP1) [19, 20]. Activated PARP1 catalyzes the formation of ADP-ribose chains, which serve as a docking platform that facilitates recruitment and assembly of the multiprotein BER complex (XRCC1-POLβ-LIG3). Breaks in DNA are filled in by DNA polymerases (primarily POLβ) using the undamaged complementary DNA strand as a template

[21]. Nicks in the damaged strand are sealed by ligases (e.g., LIG3), which finalizes repair of the

damaged DNA strand [22, 23].

Importantly, methylation-derived repair intermediates such as AP sites and SSBs are highly

cytotoxic and mutagenic. Therefore individual steps in the BER pathway need to be tightly

regulated and coordinated to prevent accumulation of those intermediates, cell death,

mutagenesis, and carcinogenesis [2, 24]. Genetic studies in yeast, mouse models, and human

cells have demonstrated that loss of the tight coordination between individual steps in the BER

pathway can trigger genome instability, increased mutagenesis, or cell death [2, 5, 24-27].

Levels and activities of BER proteins vary significantly between cells, tissues, and individuals

and correlate with cancer risk and response to alkylation chemotherapy [2, 10, 11, 25, 28-32].

Therefore, measuring and understanding differences in the rate of BER upon alkylation DNA

damage could contribute to the development of new approaches in personalized disease

prevention and treatment.

The BER pathway is dysregulated in many cancers and is often associated with cancer

heterogeneity, metastasis, and chemoresistance. Pharmacological inhibition of BER with PARP

inhibitors (e.g., olaparib) has shown enhanced cytotoxicity of various anticancer agents,

especially in tumors with defects in homologous recombination [33-37]. Identifying a pre-existing

BER imbalance within a tumor may be highly relevant for determining whether therapy involving

PARP inhibitors and alkylating agents can be beneficial.

Quantitation of DNA adduct formation and repair has greatly advanced our understanding of

DNA repair processes. A number of methods have been developed for quantitative analysis of

various enzymatic steps and the overall capacity of BER to repair alkylation DNA damage. The

most sensitive methods for detection and quantitation of alkyl DNA adducts include HPLC/^{32}P-

postlabeling, mass spectrometry-based adductomics, and radiolabeling [38-41]. These methods

offer high sensitivity; however, they require specialized equipment, expertise, and complex

sample preparation, which hinders the convenient use of those approaches to investigate

cellular BER mechanisms.

The most commonly used cell-based methods to investigate BER, include comet assays

and host cell reactivation (HCR) assays. The comet assay is a single-cell electrophoresis

technique that can be used to assess the capacity of BER to repair alkylation DNA damage

when performed under alkaline conditions [42]. This assay can be used to analyze total levels of

BER repair intermediates, such as alkali labile sites (e.g., abasic sites) and single strand breaks;

however, it does not quantitatively distinguish between these intermediates. The standard

comet assay may not reliably detect persistent and intact base modifications (e.g., 7meG or

3meA) that are not converted to AP sites or SSBs. In addition, the comet assay may not detect

lesions that form and persist within highly inaccessible heterochromatin fractions of the genome.

Furthermore, the standard comet assay workflow is laborious and prone to day-to-day

variability. It may also require extensive optimization of experimental conditions, including pH or

salts used during the alkaline electrophoresis steps, to achieve sensitivity and consistent

reproducibility [42, 43]. HCR is another method that has been used to measure the capacity of

BER to repair alkylation DNA damage in living cells [44, 45]. HCR relies on the transfection of

the non-replicating DNA plasmid with a reporter gene (e.g., *luciferase*) that contains chemically-

induced DNA base damage, which is subject to repair by the BER pathway. The presence of the

DNA base damage within the reporter gene inhibits its expression, whereas the repair of base

damage re-activates reporter expression. The HCR assay can be especially challenging to

assess repair of alkylation DNA damage, due to *in vitro* instability of some of the alkyl DNA

adducts (e.g., 7meG and 3meA), which can markedly affect assay reproducibility and

consistency [45]. Also, HCR involves a non-genomic DNA substrate that does not necessarily

reflect the complexity of the genomic chromatin environment.

More recently, high resolution, high throughput approaches such as LAF-seq (Lesion-

Adjoining Fragment Sequencing) or NMP-seq (N-methyl purine sequencing) utilizing next

generation DNA sequencing have been developed to enable precise mapping and quantitation

of methyl DNA adducts across the genome and at specific genomic loci [46, 47]. These

approaches offer unprecedented single base resolution, but can be laborious, involving

generation of DNA sequencing libraries and extensive bioinformatics analyses of the

sequencing data, especially when used with human cells. In addition, those methods may

require high (non-physiological) doses of DNA damaging agents and large amounts of input

DNA.

Here we report a reliable, gel-based method, called alk-BER, that offers a fast and

quantitative measure of BER capacity in living cells. Alk-BER was developed by an adaptation

of a previous method for DNA damage quantitation by alkaline gel electrophoresis, originally

developed by Sutherland et al. [48]. Alk-BER can be used to assess overall capacity of BER to

repair MMS-induced alkylation DNA damage within the genome of living cells. The assay can be

used to facilitate identification of new, conserved regulators of the BER pathway by using

complementary model eukaryotic systems, including fungal model organisms and human cells.

Application of the alk-BER assay could also facilitate identification of BER-deficient cancer sub-

types, which might represent a suitable targets for therapy with alkylating agents and/or PARP

inhibitors.

2. Materials and Methods

2.1. DNA damage and time course of repair in yeast cells. Yeast (*S. cerevisiae*) liquid cell cultures were inoculated from single colonies and grown in 10 ml of YPD (Yeast extract - Peptone-Dextrose) medium for ~ 16 h at 30°C in an orbital shaker. The following day, cells were sub-cultured in fresh media and grown until the cultures reached the logarithmic stage of growth, as determined by measuring the optical density of the cell culture (e.g., OD_{600} ~ 0.6). Next, MMS was added to the liquid cultures at a final concentration of 0.2% (vol/vol) and cells were incubated for 10 min at 30°C in an orbital shaker. Alternatively, cells were treated with 0.035% MMS for 1- 3 h, followed by removal of media containing MMS and repair time course in fresh media for 1-6 h. Cells were then harvested by centrifugation, and the supernatant fractions containing MMS were removed and disposed following DEHS guidelines. Cell pellets were washed with ice-cold phosphate buffered saline (1X PBS), re-suspended in a pre-warmed YPD media, and allowed to repair for a total of 3 h. Extended repair time points (longer than 4-5 hours) were avoided to ensure that restoration of the genome integrity was due to the activity of BER, and not due to lesion bypass and DNA replication. Cultures were incubated with continuous shaking and cell aliquots were collected at different repair time points [49, 50].

2.2. Yeast genomic DNA isolation. Yeast genomic DNA was extracted with the glass bead method following previously established protocols [51]. Briefly, cell pellets were mixed with 250 µL of DNA lysis buffer [2% (vol/vol) Triton X-100, 1% SDS, 100 mM NaCl, 10 mM Tris·HCl, pH 8.0, 1 mM EDTA, 250 µL of PCI (phenol: chloroform: isoamyl alcohol = 25:24:1), and 150 µL of acid-washed glass beads, and vortexed vigorously for 4 min. Next, 200 µL of 1xTE buffer (10 mM Tris·HCl, pH 7.5, 1 mM EDTA) were added and cell lysates were centrifuged at 14,000 rpm at 4°C. The supernatant fraction was transferred to a fresh Eppendorf tube and mixed with 1 mL of ice-cold 200 proof ethanol. Samples were incubated at -80°C for 15 min to facilitate formation of the DNA precipitate. Next, samples were centrifuged at 14,000 rpm at 4°C, and washed with 70% (vol/vol) ethanol. The DNA pellets were dissolved in 200 µL of 1xTE buffer and incubated with 2 µL of RNase A (Thermo Fisher Scientific, cat # EN0531) at 37°C for 1 h to remove RNA. DNA was subsequently ethanol-precipitated, dissolved in sterile deionized H_2O, and then stored at -80°C.

2.3. AAG and APE1 reactions. Purified genomic DNA was processed with (+) or without (-) an enzymatic cocktail composed of AAG (New England BioLabs, cat# M0313S) and APE1 (New

England BioLabs, cat# M0282), to convert MMS-induced 7meG and 3meA to SSBs. DNA
samples (0.6-1µg) were incubated with 1 µL of AAG and 1 µL of APE1 in the reaction buffer (70
mM MOPS, pH 7.5,1 mM dithiothreitol (DTT), 1 mM EDTA, 5% glycerol) at 37°C for 1 h in a
total reaction volume of 20 µL [52]. Methylated bases were cleaved by AAG glycosylase and the
resulting abasic sites were cleaved by APE1 endonuclease. The reactions were stopped by
adding DNA loading buffer (50 mM NaOH, 1 mM EDTA, 3% Ficoll, 0.025% Bromocresol green,
0.041% Xylene cyanol). Following enzymatic digestion of DNA cleavage products, single and
double strand breaks were resolved on 1.2% alkaline agarose gels and stained with SYBR Gold
[49, 50].

2.4. Alkaline agarose gel electrophoresis. Alkaline agarose gel electrophoresis was
performed following previously published protocols with modifications [48, 53]. The large gel
electrophoresis box and casting tray were used. The agarose solution was prepared by adding
4.3 g of agarose to 360 ml of H_2O in a 1 L Erlenmeyer flask and then heating in a microwave
oven until the agarose was dissolved. The solution was cooled to 55°C and then followed by
addition of a 0.1 volume (40ml) of 10X alkaline agarose gel electrophoresis buffer (500 mM
NaOH, 10 mM EDTA, pH 8.0). Addition of 10x alkaline buffer to a hot agarose solution should
be avoided because NaOH in the buffer may cause hydrolysis of the agar. The agarose solution
was poured into a large gel casting tray. After the gel was completely solidified, it was mounted
in the electrophoresis tank. Then the tank was filled with 1X alkaline electrophoresis buffer until
the gel was covered with the buffer at a depth of 3-5 mm above the gel. DNA samples after
AAG and APE1 digestion were collected and 6X alkaline gel-loading buffer (300 mM NaOH, 6
mM EDTA, 18% Ficoll, 0.15% Bromocresol green, 0.25% Xylene cyanol) was added to each
sample. Chelating all Mg^{2+} with EDTA (component of the 6X alkaline gel-loading buffer) is
important before loading the samples onto the alkaline agarose gel because in solutions with a
high pH, Mg^{2+} can form insoluble Mg $(OH)_2$ precipitates that entrap DNA and inhibit DNA
mobility through the gel. Samples were loaded and the gel was run at room temperature at 30V
for 19-24 h. Alternatively, the electrophoresis can be run at 4°C in a cold room. We found that
running the gel in a cold room helped improve the sharpness of the DNA bands. Note that after
the run is completed, the Bromocresol green dye may not be visible because of dye diffusion in
the gel. The gel was transferred to a large plexiglass tray, covered with neutralizing solution (1
M Tris-HCl, 1.5M NaCl), and incubated for 30 min with gentle shaking on an orbital shaker. Next
the gel was transferred to staining solution (1x TAE buffer with SYBR gold) and stained for 1 h
with gentle shaking. The container with the gel was covered with aluminum foil to protect the

staining solution from light. Following staining, the gel was rinsed and de-stained with H_2O for 30

min with gentle shaking. An image of the gel was acquired using the Storm phosphor imager.

2.5. Quantitative analysis of DNA damage and repair following MMS treatment.

Quantitation of methylated bases in genomic DNA was performed on phosphor image data by

using ImageQuant software (Molecular Dynamics) and number-average DNA length analysis.

The number-average length of genomic DNA (± AAG/APE treatment) was used to calculate the

average number of SSBs/kb, and the percentage of MDAs removed (% repair) was determined

as described previously [49, 51]. Briefly, using the ImageQuant software functions, each data

point on the gel image is marked with a box encompassing the entire length of the lane to give

the total area of each lane. The data point corresponding to 1/2 the total area, designated as

X_{med}, is then determined. The X_{med} value indicates the median migration distance of the DNA

fragments. X_{med} is converted to the median length L_{med} of DNA molecules by using a standard

curve generated from the migration of DNA-size markers. The number average molecular

length, L_n, is calculated from L_{med} by using the equation $L_n = 0.6\ L_{med}$ [54], assuming a Poisson

distribution of DNA fragments. Numbers of SSBs/kb is calculated using the following equation;

SSBs/kb = $1/L_n$ (+AAG+APE1) - $1/L_n$ (-AAG-APE1). Calculated numbers of SSBs per kb DNA

fragment indicate numbers of MDAs.

2.6. DNA damage and time course of repair in multicellular fungus *Neurospora crassa*.

Liquid cultures of *N. crassa* were grown for 11 h in VMM (Vogel's minimal medium) media in an

orbital shaker at 30°C. MMS was added to liquid cultures at a final concentration of 0.035%

(vol/vol). Cells were incubated in the presence of MMS for 3 h. Next, the cells were collected by

using a Buchnner funnel, and washed with 500 ml of VMM media to remove MMS. Washed

cells were transferred to pre-warmed VMM media and then allowed to grow at 30°C for 4 h to

enable repair of damaged DNA. Aliquots of cells were harvested and immediately frozen in

liquid nitrogen. Genomic DNA was isolated and 300 ng of genomic DNA was digested with

APE1 endonuclease (cat # M0282S, New England Biolabs), AAG glycosylase (cat # M0313S,

New England Biolabs), or both enzymes in MOPS reaction buffer (70 mM MOPS, pH 7.5,1 mM

dithiothreitol DTT, 1 mM EDTA, 5% glycerol) for 1 h and 15 min at 37°C. Reactions were

stopped by adding alkaline DNA loading buffer (50 mM NaOH, 1 mM EDTA, 3% Ficoll).

Samples were resolved on a 1.2% alkaline agarose gel. Agarose gels were run in a cold room

at 25V for 17 h and then incubated in neutralization buffer (1.5 M NaCl, 1 M Tris-Cl pH 7.6) for

45 min before being stained with SYBR Gold (cat # S-11494, Life Technologies) for 40 min and

de-stained for 30 min before imaging [55].

2.7. DNA damage and time course of repair in human cells. Human adrenal carcinoma cells

(SW13) were cultured in DMEM (Sigma, cat# D5796) supplemented with 10% fetal bovine

serum FBS (Gibco, cat# 26140-079). Human fibroblast CHON-002, leukemia HAP1, and

lymphoblastoid cells GM12878 were cultured according the ATCC guidelines. Cell cultures were

routinely tested for mycoplasma by using a mycoplasma detection kit (ATCC, cat# 30-1012K).

Cell doubling time was determined following ATCC guidelines and viability was routinely

monitored with trypan blue. Cells were seeded in T25 flasks (~500,000 cells per dish) and

grown for 16-24 h until cells reached 60-70% confluence. One T25 flask was set for each data

point to be collected. In DNA damage dose response experiments SW13 cells were treated with

(0- 20mM) MMS in 1xPBS for 10 min at RT. In DNA damage and repair experiments, SW13

cells were treated with 10 mM (0.1%) MMS for 10 min in 1xPBS at RT, or alternatively ice-cold,

serum-free media can be also used for MMS treatments. Ice-cold treatment is used to inhibit

endogenous background BER during the MMS treatment. MMS was removed and cells were

washed with 1XPBS. Next, fresh pre-warmed media were added and cells were allowed to

repair for 0, 3, 8, or 22 hours. GM 12878 cells were treated with 5mM (0.05%) and 10mM

(0.1%) MMS for 5 min. Total genomic DNA was purified from each time point using the PureLink

genomic DNA mini kit (K 182001, Thermo Fisher Scientific). DNA was processed with AAG and

APE1 enzymes as described above. DNA was resolved on a 1.2% alkaline agarose gel, run at

30V for 22 h, stained with SYBR gold, and quantified as described above.

3. Results

3.1. Alk-BER assay workflow. The conserved nature of the BER pathway enables easy adaptation of the assay to various cell types and model organisms from lower eukaryotes to human cells. We initially developed the alk-BER assay to measure BER in yeast cells (*S. cerevisiae*) and successfully adapted the assay to other fungal model organism (*N. crassa*) and human cells. The alk-BER assay is based on the enzymatic conversion of the MMS-induced damaged DNA bases to single strand breaks (SSBs), which are subsequently resolved on alkaline agarose gels and quantified. The rate of BER can be analyzed by performing a DNA damage and time course for repair and assessing the rate of removal of damaged bases from the total genomic DNA. The assay is performed in 5 simple steps (**Fig. 1**) and can be completed in 3 days. The first step involves exposing the cells to sub-lethal doses of MMS to induce formation of methyl DNA adducts (mainly 7meG and 3meA). In the second step, cell aliquots corresponding with DNA damage and repair time points are collected and total genomic DNA is isolated. The third step involves conversion of MDAs to SSBs with BER enzymes (AAG glycosylase and APE1 endonuclease) that specifically bind and cleave the DNA at sites of MDAs. The fourth step involves running the samples on alkaline agarose gels to separate DNA fragments containing SSBs from the bulk genomic DNA that does not contain damage. The final step involves staining of the separated DNA fragments in the gel, acquiring an image of the gel and performing quantitation of MDAs. The alk-BER method directly quantifies the numbers of MDAs in purified genomic DNA and analyzes the kinetics of DNA repair at the whole-genome level. In order to assess the capacity of BER using the alk-BER assay, cells should be exposed to sub-lethal doses of MMS, that induce detectable levels of MDAs, but do not induce substantial cell death. To determine sub-lethal doses for specific cell line it is recommended to perform MMS dose response experiment encompassing range of MMS doses from low to high. Cell death is induced when unrepaired lesions persist due to inability of BER to efficiently repair abnormally high levels of DNA damage or when BER activity is compromised (e.g., BER gene mutants). BER capacity should also be analyzed within a specific, experimentally determined window of time following DNA damage, typically 0-3 h for yeast cells, or 0-24 h for human cells, to avoid interference from lesion bypass and DNA replication.

Figure 1. Schematic outline of alk-BER assay. The assay involves exposing the cells to MMS (step 1), isolation of total genomic DNA (step 2), conversion of MMS-induced methylated bases to SSBs with damage specific enzymes AAG and APE1 (step 3), separation of DNA fragments containing SSBs by alkaline agarose gel electrophoresis (step 4), gel staining, imaging, and quantitation of MDAs (step 5).

3.2. Alk-BER in fungal cells. A DNA damage dose response assay was performed by exposing yeast cells (strain BY4741) to increasing doses of MMS (5, 10, 20 mM) for 10 min at 30°C. Total genomic DNA was isolated and treated with AAG and APE1 enzymes. DNA samples were resolved by alkaline agarose gel electrophoresis and numbers of MDAs were quantified as described above. As the gel is run at alkaline pH, hydrogen bonding between the two DNA strands is broken to facilitate separation of strands containing breaks from non-damaged genomic DNA. The denatured DNA is maintained in a single-stranded state and migrates through the alkaline gel as a function of its size, forming a distinct smear. Increased formation of MMS-induced DNA lesions in response to increased doses of MMS is shown as

increases in lower molecular weight DNA molecules, also visible as a smear (**Fig. 2A**). The
frequency of MDAs (7meG and 3meA) was calculated and plotted as the number of methyl A, G
per kb fragment as a function of increasing doses of MMS (**Fig. 2B**). The proportional
relationship between increasing MMS doses and numbers of MDAs indicates a high sensitivity
of the alk-BER method, ~1.5 MDAs per 10,000 bases, that is induced by 5mM MMS dose in
yeast cells.

DNA damage and time course of repair has been performed to evaluate the overall rate of
BER to repair alkylation DNA damage in the yeast wild type strain BY4741. DNA molecules
containing MMS-induced MDAs are converted to SSBs. Progressive shortening of the smear in
migration and restoration of the genome integrity indicated by formation of high molecular
weight DNA during the incubation time course indicates proficient DNA repair. DNA was
processed with AAG and APE1 and resolved on alkaline agarose gels as described previously.
A representative gel image showing DNA damage and repair in the BY4741 yeast strain is
shown (**Fig. 2C**) and corresponding quantitative analysis of the gel is also shown (**Fig. 2D**).
BER in the WT BY4741 yeast strain is proficient and nearly 90% of the total MDAs in the
genome are repaired after 3 h repair time at the dose used (**Fig. 2D**). The specificity of the alk-
BER method was validated using yeast and *Neurospora* mutant cells deficient in BER. The
yeast *mag1Δ* mutant has no Mag1 glycosylase (orthologue of human AAG) and is deficient in
cleaving MMS-induced 7meG and 3meA from the DNA. Cells from the WT and *mag1Δ* strains
were exposed to MMS for a total of 3 h followed by MMS removal and a 6 h-long repair time
course to allow cells to repair damaged DNA. During the 3 h-long MMS exposure, mutant cells
displayed higher levels of MDA formation as compared to WT because endogenous BER is
inactive in the mutant cells, which results in accumulation of MDAs under the conditions of
continuous MMS exposure. After removal of MMS, WT cells with proficient BER were able to
clear most of the lesions during the repair time course, whereas BER-deficient *mag1Δ* cells
contained high levels of unrepaired MDAs (**Fig. 3**). The alk-BER assay was successfully
adapted to multicellular fungal model *Neurospora crassa*. DNA damage and repair time course
was performed with *Neurospora* WT and *mag1Δ* cells [55]. *Neurospora* cells of the wild type
laboratory strain were exposed to 0.035% MMS continuously for 3 hours, followed by repair time
course in media without MMS for 5 hours. Genomic DNA was isolated and processed with
combinations of AAG and APE1 enzymes and resolved on alkaline agarose gel. BER capacity
to repair MMS-induced lesions in *Neurospora* cells is very efficient, with nearly complete
restoration of genome integrity following 2 h repair period (**Fig.S3**).

Figure 2. Alk-BER assay in yeast cells (*S. cerevisiae*). A) Representative alkaline agarose

gel image of MMS-induced DNA damage dose response in the BY4741 strain of *S. cerevisiae*.

C: control, DNA of cells not exposed to MMS and DNA of cells exposed to increasing doses of

MMS (5, 10, or 20 mM) were resolved on alkaline agarose gels. Each DNA sample was treated

with (+) and without (-) a cocktail of AAG and APE1 enzymes. **B)** Dose dependent increase in

the numbers of MMS-induced methyl G, A per 1 kb DNA fragment. Each data point denotes the

average value of 2 independent experiments. **C)** Representative gel image of DNA damage

and repair time course in the BY4741 strain of *S. cerevisiae*. M: DNA size standard

lambda/HindIII. C: control, cells not exposed to MMS, 0: cells collected after 10 min exposure to

20 mM MMS, 1-3 h: cells collected after 1, 2, 3 h of repair. **D)** Quantitative representation of

data displayed in panel C. Formation and repair of MMS-induced methyl G and A (7meG,

3meA), as a function of repair time. Each data point represents an average of 3 independent

experiments; error bars were calculated based on standard deviation.

Figure 3. Alk-BER assay validation with BER-deficient mutant cells, *mag1Δ*. The BER rate
was analyzed in *mag1Δ* mutant. WT and mutant cells were treated with 0.2% MMS for 1- 3 h,
followed by removal of media containing MMS and repair time course in fresh media for 1-6 h.

3.3. Alk-BER in human cells. The human adrenal carcinoma SW13 cell line was used to
adapt and optimize the alk-BER assay for assessing rates of BER in human cells. A series of
MMS-dose response experiments were initially performed to determine the appropriate range of
MMS concentration and time of the exposure for induction of detectable levels of MDAs at sub-
lethal MMS doses. Cell viability was routinely monitored with trypan blue, **Fig. S2C** [56].
Representative MMS dose response data are presented in **Fig. 4A,B**. Increases in the smear
length in response to increasing doses of MMS in the (+AAG&APE1) lanes indicate enhanced
formation of MDAs. The minimal smear in (-AAG&APE1) lanes reveals formation of MDA-
derived BER intermediate AP sites and SSBs that form in DNA as a result of the continuous
activity of endogenous BER during MMS exposure. AP sites are fragile in alkaline conditions
and can spontaneously convert to SSBs contributing to the smear in (-enzyme) lanes. The
proportional relationship between increasing MMS doses and numbers of MDAs indicates a
high sensitivity of the alk-BER method, ~0.7 MDAs per 10,000 bases with 5mM MMS dose in
human cells. The BER capacity in SW13 cells was analyzed by performing DNA damage and a
time course of repair as described in the methods section. Cells were exposed to 10 mM MMS
for 10 min in 1XPBS, at RT and allowed to repair for 22 h. Representative data showing the
image of the alkaline agarose gel and data quantitation are presented (**Fig. 4C,D**). Other human

cell lines were subject to alk-BER assay, including untransformed fibroblast cells CHON-002, and leukemia cancer cells HAP1. The BER capacity to remove MMS-induced adducts was quantitated over distinct repair time points. The repair rates vary significantly between different cell lines (**Fig.4D**). Additionally, alk-BER assay was performed with human lymphoblastoid cell line GM12878. Cells were exposed to two different doses of MMS (0.05% and 0.1%) for 5 min, and BER efficiency was analyzed at 2.5 and 5 h in the repair time in media without MMS. Clear dose-dependent DNA damage, followed by progressive and efficient repair are observed in the image of the alkaline agarose gel (**Fig. S4**). Specificity of alk-BER was analyzed by exposing SW13 cells to sub-lethal doses of temozolomide (TMZ), clinically used SN1 alkylator, known to induce 7meG, 3meA and O6meG adducts [57]. As expected, alk-BER can specifically detect TMZ-induced DNA methyl adducts (**Fig. S1A**), although the TMZ dose used in our experiment (10mM, 10min) appears less potent than MMS. We also analyzed performance of alk-BER assay with hydrogen peroxide-induced oxidative DNA damage. As expected, alk-BER does not detect oxidative DNA damage (**Fig.S1B**).

Figure 4. Alk-BER assay in human cells. A) MMS dose response in SW13 cells. Cells were treated with increasing doses of MMS for 10min at RT. Representative alkaline agarose gel image is shown. **B)** Quantification of methyl A, G per 1kb DNA fragment as a function of

increasing MMS dose. The graph represents quantification of the data in panel A. **C)** DNA damage and repair time course experiment was performed with several human cell lines; CHON-002, SW13, and HAP1. Cells (60-70% confluent) were exposed to 10mM (0.1%) MMS in 1xPBS for 10 min at RT, followed by DNA repair time course 0, 3, 8 and 22 hours at 37^{0}C. **D)** DNA repair rates were quantitated individually for each cell line and expressed as a percentage (%) of the removed methyl A,G, as compared to the 0hr time point.

4. Discussion

The efficiency of BER in repairing alkylation DNA damage varies substantially between different cells, tissues, and individuals, and has important implications in cancer development and treatment [1, 2, 58, 59]. BER efficiency is a result of a complex interplay between genetic and epigenetic factors influencing the abundance and activity of the BER enzymes, and the individual steps in the BER pathway. Ability to measure the formation and rates of repair of alkyl DNA adducts in genomic DNA provides a direct assessment of BER efficiency in a given cell type.

Here, we report a quantitative cell-based assay, alk-BER (<u>alk</u>ylation <u>B</u>ase <u>E</u>xcision <u>R</u>epair), adapted and optimized for measuring efficiency and rates of BER following alkylation DNA damage in fungal model organisms (*S.cerevisiae, N.crassa*) and human cell lines (SW13, CHON-002, HAP1 and GM12878). Alk-BER offers a simple, time- and cost-efficient, cell-based method for quantitative analysis of alkylation DNA damage and repair in the genomic DNA. The alk-BER assay can be used to determine BER efficiency in various cell types, by assessment of the rates of methyl DNA adducts removal from the DNA over time.

Yeast cells have been extensively used to study DNA repair processes in eukaryotic cells [60]. Yeast have a robust and conserved BER pathway to repair alkylation DNA damage. Using alk-BER assay, we found that in wild type strains of fungal model organisms (*S. cerevisiae* and *N. crassa*), BER proceeds quite rapidly, following removal of the MMS from the growth media. We showed that BER in the BY4741 strain of *S. cerevisiae* is nearly completed within 3-4 h post exposure since nearly 80-90% of the lesions were removed from the genome (**Fig. 2C, D**). Similarly, repair of alkylation DNA damage has been nearly completed within 2-4 h in the wild type laboratory strain of *N. crassa* [55] (**Fig.S3**). We also showed that BER-deficient *mag1Δ* yeast mutant cells were deficient to clear MMS-induced lesions and accumulated MDAs over time, which validates specificity of the alk-BER assay (**Fig. 3**). The human genome is much larger therefore repair is expected to take longer compared to lower eukaryotes. We found that the majority of the genome (~70% lesions removed) was restored following 22 h post MMS

exposure in SW13 cells (**Fig. 4C, S1A,D**), which is consistent with previous studies reporting

that that majority of DNA alkylation repair in mammalian cells can be completed within 24 h post

exposure, as revealed by results generated with various quantification methods [39, 61-65]. We

found that the rates of BER to remove MMS-induced DNA adducts vary significantly between

SW13 and CHON-002, and HAP1 cell lines (**Fig. 4C**). These rates can be influenced by the

endogenous levels and activities of BER enzymes and other regulators of DNA repair. Future

studies are needed to further investigate regulators of DNA alkylation repair in human cells.

In lower eukaryotes (*S.cerevisiae, N.crassa*), the rate of repair of MMS-induced methyl DNA

adducts is strongly dependent on the functional MAG1 glycosylase, where *mag1Δ* mutants

demonstrate abolished ability to repair MDAs. Interestingly, repair of MDAs in certain

mammalian cells does not appear to be exclusively dependent on AAG glycosylase. It has been

reported that the alkylated bases 3meA and 7meG, both AAG substrates generated from MMS

treatment, are removed from the genome of AAG-deficient embryonic stem cells, with slower

kinetics for 3meA but comparable kinetics for 7meG [66]. Other study revealed that similar

levels of 7meG were detected in livers of $AAG^{+/+}$ and $AAG^{-/-}$ mice 24hr after exposure to MNU

[67]. These studies suggest that methyl DNA adducts are excised in the absence of AAG

enzyme, in mammalian cells, perhaps by involvement of other glycosylases, or spontaneous

depurination. Future application of alk-BER could facilitate further understanding of the role of

AAG and other factors in regulation of human BER.

The alk-BER could serve as useful framework for number of approaches to study repair of

DNA alkylation. For example, alk-BER assay could be used to distinguish, in a quantitative way,

between the levels of MDAs from levels of AP sites and SSBs. Numbers of MDAs could be

revealed by treating the DNA sample with both AAG and APE1, whereas numbers of AP sites

could be detected by treating the aliquot of the same sample with APE1 enzyme only, and

numbers of SSBs would be revealed by running the same sample without enzymatic treatment.

Highly specific and sensitive detection of MDAs or AP sites could also be performed by

processing of the sample with the AAG enzyme only (converts MDA to AP sites) and

subsequent detection of AP sites using a highly sensitive AP site detection kit (e.g., Abcam, ab

211154). Alk-BER assay can also serve as a framework for quantification of gene-specific repair

when coupled with Southern blot and hybridization of gene-specific probes. Alk-BER could also

be useful in detection and quantification of MMS-induced methyl DNA adducts in preparation

and optimization of samples for the approaches involving next generation sequencing, such as

NMP-seq.

In summary, the alk-BER assay offers a versatile, reliable and affordable approach for

quantitative analysis of DNA damage formation and repair following exposure to DNA

methylating alkylating agents. The alk-BER assay can be easily optimized to be used in any

type of cultured cells, and integrated with the existing approaches to study mechanisms

regulating BER balance and capacity. The assay has the ability to detect imbalances in the

activity of the BER process, that is highly relevant to cancer development and treatment.

Quantitative analyses of DNA alkylation damage and repair using fungal genetic model

organisms and human cell lines offer unique opportunities to identify novel, conserved

regulators of BER.

Acknowledgements

We thank Dr. Ann Bode for help with proof reading the manuscript. We would like to

acknowledge Dr. John Wyrick for helpful discussions and intellectual contributions to this work.

This work was supported by grant R01ES002614 from the National Institute of Environmental

Health Sciences (NIEHS) (to M.J.S. and J.J.W.), R21ES028549 from NIEHS to W.C., and

Faculty Research Grant from University of Georgia to W.C. The contents are solely the

responsibility of the authors and do not necessarily represent the official views of the NIEHS,

NIH.

Figure S1. Specificity of alk-BER. A) Temozolomide (TMZ) induced DNA damage and repair time course. SW13 cells were exposed to sub-lethal dose of TMZ, 10mM for 10 min at RT, followed by TMZ removal and DNA repair for the indicated time. **B)** Hydrogen peroxide (H2O2) induced DNA damage and repair time course. SW13 cells were exposed to sub-lethal dose of H2O2, 5mM for 5min at RT, followed by H2O2 removal and DNA repair for the indicated time.

Figure S2. Alk-BER assay in SW13 cell line. A) Alkaline gel image representing DNA damage and repair time course. SW13 cells were exposed to 10mM MMS for 10min. Genomic DNA was isolated, and processed with double enzyme AAG&APE1 digest. **B)** Efficiency of double enzyme (AAG&APE1), and single enzymes: (AAG only), and (APE1 only), in converting methyl DNA adducts to SSBs. **C)** SW13 cell viability measured by trypan blue. **D)** Quantification of the data presented in panel A.

Time (hr)	0			1			2			3			4			5			6			7			8		
MMS	−			+			+			+			−			−			−			−			−		
APE1	+	−	+	+	−	+	+	−	+	+	−	+	+	−	+	+	−	+	+	−	+	+	−	+	+	−	+
AAG	−	+	+	−	+	+	−	+	+	−	+	+	−	+	+	−	+	+	−	+	+	−	+	+	−	+	+

Figure S3. Alk-BER assay in *Neurospora crassa*. Representative alkaline agarose gel image of MMS-induced DNA damage, followed by DNA repair in the wild type strain of *Neurospora*, 0: control DNA from cells not exposed to MMS; 1-3 hours: DNA from cells exposed to 0.035% MMS continuously for 1, 2 and 3 hours respectively; 4-8 hours: DNA from cells that were allowed to repair DNA in media without MMS for 1-4 hours respectively. Each DNA sample was treated with combination of human APE1 and AAG enzymes: APE1 (+) and without AAG (-), without APE1 (-) and with AAG (+), and with both enzymes APE (+) & AAG (+).

Figure S4. Alk-BER assay in human lymphoblastoid cell line GM12878. The representative alkaline agarose gel illustrates DNA damage and repair time course performed with GM12878 cells, using two different MMS doses (0.05% and 0.1%) respectively. Cells were treated with the indicated doses for 5 minutes, MMS was removed and cells were allowed to repair DNA for 2.5 and 5 hours.

References

1. Beranek, D.T., *Distribution of methyl and ethyl adducts following alkylation with monofunctional alkylating agents.* Mutat Res, 1990. **231**(1): p. 11-30.

2. Fu, D., J.A. Calvo, and L.D. Samson, *Balancing repair and tolerance of DNA damage caused by alkylating agents.* Nat Rev Cancer, 2012. **12**(2): p. 104-20.

3. Margison, G.P., M.F. Santibanez Koref, and A.C. Povey, *Mechanisms of carcinogenicity/chemotherapy by O6-methylguanine.* Mutagenesis, 2002. **17**(6): p. 483-7.

4. Saffhill, R., G.P. Margison, and P.J. O'Connor, *Mechanisms of carcinogenesis induced by alkylating agents.* Biochim Biophys Acta, 1985. **823**(2): p. 111-45.

5. Klapacz, J., et al., *Frameshift mutagenesis and microsatellite instability induced by human alkyladenine DNA glycosylase.* Mol Cell, 2010. **37**(6): p. 843-53.

6. Pfeifer, G.P., et al., *Tobacco smoke carcinogens, DNA damage and p53 mutations in smoking-associated cancers.* Oncogene, 2002. **21**(48): p. 7435-51.

7. Richardson, F.C. and K.K. Richardson, *Sequence-dependent formation of alkyl DNA adducts: a review of methods, results, and biological correlates.* Mutat Res, 1990. **233**(1-2): p. 127-38.

8. Head, R.J., et al., *Persistence of DNA adducts, hypermutation and acquisition of cellular resistance to alkylating agents in glioblastoma.* Cancer Biol Ther, 2017: p. 0.

9. McNerney, M.E., L.A. Godley, and M.M. Le Beau, *Therapy-related myeloid neoplasms: when genetics and environment collide.* Nat Rev Cancer, 2017. **17**(9): p. 513-527.

10. Chaim, I.A., et al., *In vivo measurements of interindividual differences in DNA glycosylases and APE1 activities.* Proc Natl Acad Sci U S A, 2017.

11. Nagel, Z.D., I.A. Chaim, and L.D. Samson, *Inter-individual variation in DNA repair capacity: a need for multi-pathway functional assays to promote translational DNA repair research.* DNA Repair (Amst), 2014. **19**: p. 199-213.

12. Chatterjee, N. and G.C. Walker, *Mechanisms of DNA damage, repair, and mutagenesis.* Environ Mol Mutagen, 2017. **58**(5): p. 235-263.

13. Soll, J.M., R.W. Sobol, and N. Mosammaparast, *Regulation of DNA Alkylation Damage Repair: Lessons and Therapeutic Opportunities.* Trends Biochem Sci, 2017. **42**(3): p. 206-218.

14. Kim, Y.J. and D.M. Wilson, 3rd, *Overview of base excision repair biochemistry.* Curr Mol Pharmacol, 2012. **5**(1): p. 3-13.

15. Samson, L., et al., *Cloning and characterization of a 3-methyladenine DNA glycosylase cDNA from human cells whose gene maps to chromosome 16.* Proc Natl Acad Sci U S A, 1991. **88**(20): p. 9127-31.

16. Samson, L.D., *The repair of DNA alkylation damage by methyltransferases and glycosylases.* Essays Biochem, 1992. **27**: p. 69-78.

17. Li, M. and D.M. Wilson, 3rd, *Human apurinic/apyrimidinic endonuclease 1.* Antioxid Redox Signal, 2014. **20**(4): p. 678-707.

18. Wilson, D.M., 3rd, M. Takeshita, and B. Demple, *Abasic site binding by the human apurinic endonuclease, Ape, and determination of the DNA contact sites.* Nucleic Acids Res, 1997. **25**(5): p. 933-9.

19. Ko, H.L. and E.C. Ren, *Functional Aspects of PARP1 in DNA Repair and Transcription.* Biomolecules, 2012. **2**(4): p. 524-48.

20. Wei, H. and X. Yu, *Functions of PARylation in DNA Damage Repair Pathways.* Genomics Proteomics Bioinformatics, 2016. **14**(3): p. 131-139.

21. Wilson, S.H., et al., *DNA polymerase beta and mammalian base excision repair.* Cold Spring Harb Symp Quant Biol, 2000. **65**: p. 143-55.

22.   Tomkinson, A.E. and Z.B. Mackey, *Structure and function of mammalian DNA ligases.*
Mutat Res, 1998. **407**(1): p. 1-9.
23.   Tomkinson, A.E. and A. Sallmyr, *Structure and function of the DNA ligases encoded by*
the mammalian LIG3 gene. Gene, 2013. **531**(2): p. 150-7.
24.   Ensminger, M., et al., *DNA breaks and chromosomal aberrations arise when replication*
meets base excision repair. J Cell Biol, 2014. **206**(1): p. 29-43.
25.   Calvo, J.A., et al., *Aag DNA glycosylase promotes alkylation-induced tissue damage*
mediated by Parp1. PLoS Genet, 2013. **9**(4): p. e1003413.
26.   Ebrahimkhani, M.R., et al., *Aag-initiated base excision repair promotes ischemia*
reperfusion injury in liver, brain, and kidney. Proc Natl Acad Sci U S A, 2014. **111**(45): p.
E4878-86.
27.   Meira, L.B., et al., *Aag-initiated base excision repair drives alkylation-induced retinal*
degeneration in mice. Proc Natl Acad Sci U S A, 2009. **106**(3): p. 888-93.
28.   Redaelli, A., et al., *AP endonuclease activity in humans: development of a simple assay*
and analysis of ten normal individuals. Teratog Carcinog Mutagen, 1998. **18**(1): p. 17-26.
29.   Brenerman, B.M., J.L. Illuzzi, and D.M. Wilson, 3rd, *Base excision repair capacity in*
informing healthspan. Carcinogenesis, 2014. **35**(12): p. 2643-52.
30.   Memisoglu, A.S., L., *Base excision repair in yeast and mammals.* Mutation Research,
2000. **451**: p. 39-51.
31.   Wallace, S.S., *Base excision repair: a critical player in many games.* DNA Repair (Amst),
2014. **19**: p. 14-26.
32.   Wallace, S.S., D.L. Murphy, and J.B. Sweasy, *Base excision repair and cancer.* Cancer
Lett, 2012. **327**(1-2): p. 73-89.
33.   Abbotts, R., N. Thompson, and S. Madhusudan, *DNA repair in cancer: emerging targets*
for personalized therapy. Cancer Manag Res, 2014. **6**: p. 77-92.
34.   Montaldi, A.P., P.R. Godoy, and E.T. Sakamoto-Hojo, *APE1/REF-1 down-regulation*
enhances the cytotoxic effects of temozolomide in a resistant glioblastoma cell line.
Mutat Res Genet Toxicol Environ Mutagen, 2015. **793**: p. 19-29.
35.   Montaldi, A.P. and E.T. Sakamoto-Hojo, *Methoxyamine sensitizes the resistant*
glioblastoma T98G cell line to the alkylating agent temozolomide. Clin Exp Med, 2013.
13(4): p. 279-88.
36.   Sultana, R., et al., *Synthetic lethal targeting of DNA double-strand break repair deficient*
cells by human apurinic/apyrimidinic endonuclease inhibitors. Int J Cancer, 2012.
131(10): p. 2433-44.
37.   Liu, L., Y. Nakatsuru, and S.L. Gerson, *Base excision repair as a therapeutic target in*
colon cancer. Clin Cancer Res, 2002. **8**(9): p. 2985-91.
38.   Guo, J. and R.J. Turesky, *Emerging Technologies in Mass Spectrometry-Based DNA*
Adductomics. High Throughput, 2019. **8**(2).
39.   Haque, K., et al., *Accurate and sensitive quantitation of N7-methyldeoxyguanosine-3'-*
monophosphate by 32P-postlabeling and storage-phosphor imaging. Chem Res Toxicol,
1997. **10**(6): p. 660-6.
40.   Liu, S. and Y. Wang, *Mass spectrometry for the assessment of the occurrence and*
biological consequences of DNA adducts. Chem Soc Rev, 2015. **44**(21): p. 7829-54.
41.   Thomas, B., et al., *A novel method for detecting 7-methyl guanine reveals aberrant*
methylation levels in Huntington disease. Anal Biochem, 2013. **436**(2): p. 112-20.
42.   Fortini, P., et al., *Analysis of DNA alkylation damage and repair in mammalian cells by*
the comet assay. Mutagenesis, 1996. **11**(2): p. 169-75.
43.   Hartmann, A., et al., *Recommendations for conducting the in vivo alkaline Comet assay.*
4th International Comet Assay Workshop. Mutagenesis, 2003. **18**(1): p. 45-51.

44. Alanazi, J.S. and J.J. Latimer, *Host Cell Reactivation: Assay for Actively Transcribed*
DNA (Nucleotide Excision) Repair Using Luciferase Family Expression Vectors. Methods
Mol Biol, 2020. **2102**: p. 509-528.
45. Wang, L., et al., *A modified host-cell reactivation assay to measure repair of alkylating*
DNA damage for assessing risk of lung adenocarcinoma. Carcinogenesis, 2007. **28**(7):
p. 1430-6.
46. Li, M., T. Ko, and S. Li, *High-resolution Digital Mapping of N-Methylpurines in Human*
Cells Reveals Modulation of Their Induction and Repair by Nearest-neighbor
Nucleotides. J Biol Chem, 2015. **290**(38): p. 23148-61.
47. Mao, P., et al., *Genome-wide maps of alkylation damage, repair, and mutagenesis in*
yeast reveal mechanisms of mutational heterogeneity. Genome Res, 2017. **27**(10): p.
1674-1684.
48. Sutherland, B.M., P.V. Bennett, and J.C. Sutherland, *DNA damage quantitation by*
alkaline gel electrophoresis. Methods Mol Biol, 2006. **314**: p. 251-73.
49. Czaja, W., et al., *Proficient repair in chromatin remodeling defective ino80 mutants of*
Saccharomyces cerevisiae highlights replication defects as the main contributor to DNA
damage sensitivity. DNA Repair (Amst), 2010. **9**(9): p. 976-84.
50. Czaja, W., P. Mao, and M.J. Smerdon, *Chromatin remodelling complex RSC promotes*
base excision repair in chromatin of Saccharomyces cerevisiae. DNA Repair (Amst),
2014. **16**: p. 35-43.
51. Bespalov, V.A., et al., *Improved method for measuring the ensemble average of strand*
breaks in genomic DNA. Environ Mol Mutagen, 2001. **38**(2-3): p. 166-74.
52. Alseth, I., et al., *Biochemical characterization and DNA repair pathway interactions of*
Mag1-mediated base excision repair in Schizosaccharomyces pombe. Nucleic Acids
Res, 2005. **33**(3): p. 1123-31.
53. Sambrook, J. and D.W. Russell, *Alkaline agarose gel electrophoresis.* CSH Protoc,
2006. **2006**(1).
54. Veatch, W. and S. Okada, *Radiation-induced breaks of DNA in cultured mammalian*
cells. Biophys J, 1969. **9**(3): p. 330-46.
55. Basenko, E.Y., et al., *The LSH/DDM1 Homolog MUS-30 Is Required for Genome*
Stability, but Not for DNA Methylation in Neurospora crassa. PLoS Genet, 2016. **12**(1):
p. e1005790.
56. Strober, W., *Trypan Blue Exclusion Test of Cell Viability.* Curr Protoc Immunol, 2015.
111: p. A3 B 1-A3 B 3.
57. Trivedi, R.N., et al., *The role of base excision repair in the sensitivity and resistance to*
temozolomide-mediated cell death. Cancer Res, 2005. **65**(14): p. 6394-400.
58. Dianov, G.L., *Base excision repair targets for cancer therapy.* Am J Cancer Res, 2011.
7(1): p. 845-851.
59. Dianov, G.L. and U. Hubscher, *Mammalian base excision repair: the forgotten*
archangel. Nucleic Acids Res, 2013. **41**(6): p. 3483-90.
60. Boiteux, S. and S. Jinks-Robertson, *DNA repair mechanisms and the bypass of DNA*
damage in Saccharomyces cerevisiae. Genetics, 2013. **193**(4): p. 1025-64.
61. Fortini, P., *The base excision repair: mechanisms and its relevance for cancer*
susceptibility. Biochimie, 2003. **85**(11): p. 1053-1071.
62. Lawley, P.D., et al., *Repair of chemical carcinogen-induced damage in DNA of human*
lymphocytes and lymphoid cell lines--studies of the kinetics of removal of O6-
methylguanine and 3-methyladenine. Chem Biol Interact, 1986. **57**(1): p. 107-21.
63. Scicchitano, D.A. and P.C. Hanawalt, *Repair of N-methylpurines in specific DNA*
sequences in Chinese hamster ovary cells: absence of strand specificity in the
dihydrofolate reductase gene. Proc Natl Acad Sci U S A, 1989. **86**(9): p. 3050-4.

64.   Wang, W., A. Sitaram, and D.A. Scicchitano, *3-Methyladenine and 7-methylguanine*
exhibit no preferential removal from the transcribed strand of the dihydrofolate reductase
gene in Chinese hamster ovary B11 cells. Biochemistry, 1995. **34**(5): p. 1798-804.
65.   Ye, N., G.P. Holmquist, and T.R. O'Connor, *Heterogeneous repair of N-methylpurines at*
the nucleotide level in normal human cells. J Mol Biol, 1998. **284**(2): p. 269-85.
66.   Smith, S.A. and B.P. Engelward, *In vivo repair of methylation damage in Aag 3-*
methyladenine DNA glycosylase null mouse cells. Nucleic Acids Res, 2000. **28**(17): p.
3294-300.
67.   Elder, R.H., et al., *Alkylpurine-DNA-N-glycosylase knockout mice show increased*
susceptibility to induction of mutations by methyl methanesulfonate. Mol Cell Biol, 1998.
18(10): p. 5828-37.