Physiological and transcriptomic variability indicative of differences in key functions within a single coral colony

Jeana L. Drake[1,+], Assaf Malik[1,+], Yotam Popovits[1,+], Oshra Yosef[1], Eli Shemesh[1], Jarosław Stolarski[2], Dan Tchernov[1,3], Daniel Sher[1*] and Tali Mass[1,3*]

[1]Department of Marine Biology, University of Haifa, 199 Aba Koushy Avenue, Haifa, Israel, 3498838
[2]Institute of Paleobiology, Polish Academy of Sciences, Twarda 51/55, Warsaw, Poland, 00-818
[3]Morris Kahn Marine Research Station, The Leon H. Charney School of Marine Sciences, University of Haifa, Sdot Yam, Israel

+These authors contributed equally

*Corresponding authors:
Jeana Drake (during review)
jeanadrake@g.ucla.edu
Tali Mass and Daniel Sher (for publication)
tmass@univ.haifa.ac.il, dsher@univ.haifa.ac.il

Keywords: Differential gene expression, biomineralization, toxins, host and symbiont

Abstract

Polyps in different locations on individual stony coral colonies experience variation in numerous environmental conditions including flow and light, potentially leading to transcriptional and physiological differences across the colony. Here, we describe high-resolution physiological measurements and differential gene expression from multiple locations within a single colony of *Stylophora pistillata*, aiming to relate these to environmental gradients across the coral colony. We observed broad transcriptional responses in both the host and photosymbiont in response to height above the substrate, cardinal direction, and, most strongly, location along the branch axis. Specifically, several key physiological processes in the host appear more active toward branch tips including several metabolic pathways, toxin production for prey capture or defense, and biomolecular mechanisms of biomineralization. Further, the increase in gene expression related to these processes toward branch tips is conserved between *S. pistillata* and *Acropora* spp. The photosymbiont appears to respond transcriptionally to relative light intensity along the branch and due to cardinal direction. These differential responses were observed across the colony despite its genetic homogeneity and likely inter-polyp communication. While not a classical division of labor, each part of the colony appears to have distinct functional roles related to polyps' differential exposure to environmental conditions.

1 Introduction

Coral reefs remain one of the most biodiverse ecosystems on Earth (Knowlton et al., 2010), despite facing a suite of anthropogenic stressors with both local and global causes (e.g., Hughes, 1994;Hoegh-Guldberg et al., 2019). Most stony corals (order Scleractinia), the sessile builders of the reef system, grow as colonies of thousands of genetically identical polyps formed by budding, with colonies ranging from centimeters to meters in diameter. The reef-building corals are also host to intracellular symbiotic dinoflagellates whose photosynthetic activity restricts the coral holobiont to the top ~100 m of the ocean and fulfills much of the host's carbon nutritional requirements (e.g., Falkowski et al.). Nevertheless, active feeding on particles from the water column plays a significant role as well (e.g., Martinez et al.), with active predation utilizing toxins produced and delivered by cnidocytes (recently reviewed by Schmidt et al.). These fixed carbon sources ultimately power the production of the external calcium carbonate skeleton that forms the reef and persists after the animal has died.

While the individual polyps in most colonial corals are genetically identical, each polyp inhabits a distinct location within the colony, which can be morphologically complex (e.g., in the case of branching corals). Thus, it has been suggested that individual polyps within a coral colony are differentially exposed to environmental conditions such as water flow and light availability (e.g., Carpenter and Patterson, 2007;Chang et al., 2009), which have distinct effects on coral physiology. These micro-environmental conditions may play a role in feeding capability (Sebens et al., 1997), reproductive success (Mass et al., 2011), photosynthesis (Dennison and Barnes;Mass et al., 2007), calcification rate (Dennison and Barnes, 1988), and waste removal (Mass et al., 2010). For instance, reductions in flow between the colony branches can trap food particles, making it easier for hunting tentacles of the polyps to efficiently pull them from the water column (Chang et al., 2009), and leading to downstream effects on metabolism (e.g., Gladfelter et al., 1989). Similarly, differences in light exposure due to the coral's branching leads to self-shading that can dramatically reduce light availability (Anthony et al., 2005), potentially leading to fine-scale spatial heterogeneity in photosynthetic parameters between individual polyps as well as between polyps and the coenosarc (Ralph et al., 2002). We hypothesize that such physiological differences will be underpinned by transcriptional differences. Plasticity in host gene expression has been observed at coral branch tips versus bases across multiple colonies (Hemond et al., 2014). However, it has not previously been linked, at high spatial resolution, to physiological and morphological measurements indicative of relevant environmental parameters and physiology within a single coral colony.

Here we combined high-resolution transcriptomic analysis with field measurements and physiological assays to describe intra-colony plasticity in the Indo-Pacific stony coral, *Stylophora pistillata*. We sought to answer two questions: first, are there differences in function by groups of polyps at different locations throughout the colony; and secondly, can we attribute those differences to specific location characteristics that are likely associated with varying microenvironments across the colony? Our whole-genome expression and physiological data are from a single coral colony at three locations along each branch taken from a different location within the colony. This high-resolution dataset enables us to resolve variability along three main axes – tip-junction-base, top to bottom, and the cardinal direction (Figure 1). Each of these axes is associated with changes in gene expression and physiology or morphology, which we suggest are due in part to light and current direction and strength. It appears that polyps in the interior of

the colony are exposed to relatively stagnant conditions, whereas the colony periphery likely
experiences higher flow, light exposure, chances for predation, and requirements for competition
and defense leading to the dynamic physiological and transcriptional responses observed here.
Our findings have implications for the interpretation of data from branches collected from
different locations across multiple coral colonies. Specifically, when fragments are taken from
multiple colonies for replication, considerations of the inherent differences across a colony
should be included in sampling design. Our findings also shed new light on the potential drivers
of plasticity of corals' responses to environmental stress and disease, and should be considered
when reconstructing paleophysiology and past microenvironmental conditions from well
preserved fossil coral colonies.
97
98 **2   Manuscript Type**
99 Original Research, Body text is 8048 words.
100
3   **Methods**
3.1 Field measurements
3.1.1 Sample collection
Ex situ measurements were conducted on a 15 cm diameter colony of *Stylophora pistillata*
collected during the morning in April 2016 at 8 m depth under a special permit from the Israel
Nature and Parks Authority in the coral nursery of the Inter-University Institute for Marine
Science (IUI), Eilat, Israel using SCUBA (Figure 1a). The colony was brought to the lab in a
tank filled with seawater and divided into individual branches and branch segments within 20
minutes of collection. Nine branches (numbered 1-9 in Figure 1b, c and supplementary
materials) were taken from three horizontal concentric rings (one 'top' branch, four 'middle'
branches, and four 'bottom' branches) and classified according to their relative height within the
coral colony (Figure 1b, c). The eight middle and bottom branches also represent duplicates from
each cardinal direction (N, S, E, W). All branches were further divided into sections according to
their position on the branch (tip, junction, base) and flash-frozen in TRI-Reagent (Sigma) (Figure
1 inset) for differential gene expression analysis. Five additional branches (numbered 20-24 in
Figure 1b, c and supplementary materials) distributed around the colony were chosen for
physiological examination, fragmented into tip, junction, and base segments, and then kept at –
80 ˚C.
119
3.1.2 Oxygen measurements
Immediately prior to removal of the colony intended for high spatial resolution examination from
the coral nursery, dissolved oxygen (DO) in seawater between branches of five *S. pistillata*
colonies located in the IUI coral nursery was measured in-situ. This nursery is a pair of raised
4x3 m frame structures (part of one can be seen behind the coral colony in Figure 1a) ranging in
depth from 5-10 m, so that conditions experienced by all well-spaced colonies are quite similar
across the nursery. Measurements were taken as up to five radial transects per colony, from
inside to just outside each colony (i.e., center, mid-way, and periphery), using Unisense oxygen
microsensors mounted on a Unisense UnderWater Meter System. The microsensors were moved
through each colony at a controlled pace in order to keep the intra-colony water as undisturbed as
possible and to prevent exchange of inter-branch water within the colony and with the
surrounding water. Each measurement was taken for a duration of two minutes to allow the
oxygen microsensor readings to stabilize. All measurements for each transect were normalized to

the outermost reading for that transect (Supplementary Table 1). DO concentrations across one
representative transect were calculated according to the following equation to establish that such
values were normoxic: DO (μmol/L) = (DO$_{sat}$(μmol/L)/(mV$_{sat}$-mV$_0$)) * in situ mV measurement.
136
3.1.3 Microbial counts
Microbial counts outside and within the coral colony were performed at the same time as DO
measurements on samples collected in-situ via radial transects on five different colonies growing
at the same site. We conducted preliminary experiments using small aliquots of concentrated
fluorescein dye injected to water between branches of *S. pistillata* skeletons, the same size as that
used in our physiological experiments, in a static laboratory aquarium to test the efficacy of
extracting water from between branches and at colony peripheries with minimal intra-colonial
water mixing. Briefly, we injected the dye on one side of a branch and then used a syringe to
carefully extract various volumes of water, from 0.4 to 12 ml in intervals of 0.4 ml, from the
other side of the branch, with five replicate experiments per extract volume tested. We then
measured the fluorescence of the extracted water, subtracted the fluorescence background of the
surrounding water, and represented the results as a percentage of the fluorescence from the
injected fluorescein dye aliquot. This showed that we could extract up to 1.6 ml of seawater from
between coral branches and from colony peripheries without pulling in water from the immediate
environment of the surrounding branches (Supplementary Figure 1). Water flow during our
sampling dive was noticeably very low. We therefore felt confident extracting 1 ml of seawater
from between the branches ('center' position) and the tip of the branch at the colony periphery
('periphery' position).of the live colonies. Upon completion of the dive, all water samples were
immediately carried to the lab and fixed in cryovials in a final concentration of 0.0625%
glutaraldehyde, incubated in the dark for 10 minutes, and then flash-frozen at − 80 °C until
analysis by flow cytometry.
158
Just prior to analysis, microbial count samples were thawed at room temperature and diluted 1:1
in ultra-pure water, and 2 μm-diameter fluorescent beads were added to each sample as an
internal standard (Polysciences, Warminster, PA, USA). The samples were analyzed on a
FACSCanto™ II Flow Cytometry Analyzer Systems (BD Biosciences). We first examined the
natural fluorescence of the cells (chlorophyll and phycoerythrin pigments). We then stained the
cells with SYBR Green I (Molecular Probes/ThermoFisher) according to the manufacturer's
instructions and counted the total microbial population as well as cyanobacterial sub-populations.
Data were acquired and processed with FlowJo software. Flow rates of the instrument were
determined several times during each run and the average value for a sample-free test run was
used for calculating cell per ml.
169
3.2 Physiological analysis
3.2.1 Tissue removal
Tissue was removed from five frozen coral fragment skeleton by airbrush with phosphate
buffered saline (PBS) and homogenized using an electrical homogenizer (MRC, HOG-160-1/2)
for 10 seconds. A sub-sample of each fragment's homogenate was stored for cell counts,
chlorophyll extraction, and total protein measurements and the rest of the homogenate was
centrifuged for 5 minutes at 5000 rpm. The supernatant was retained for host protein and
hemolysis assay. The pellet was observed under the microscope to quantify photosynthetic

symbionts. All quantities were adjusted to fragment extraction volume and normalized to cm^2 of
the fragment's surface area.
180
3.2.2 Chlorophyll extraction
Two ml of tissue homogenate was filtered on a Whatman GF/C filter and incubated with 1 ml
90% acetone for two hours at 4°C. After incubation, the filter was manually homogenized and
the solution was filtered through a 0.22 µm syringe filter into a glass cuvette. Spectrophotometric
measurements were conducted on a NanoDrop (Thermo-Fisher) and chlorophyll a concentrations
were calculated from light absorbance results based on the following equation (Ritchie, 2008):
chl-a [mg/ml]= $-0.3319(ABS630)-1.7485(ABS_{647nm})+11.9442(ABS_{664nm})-1.4306(ABS_{691nm})$.
188
3.2.3 Nematocysts and photosynthetic symbiont counts
100 µl of tissue homogenate was used for nematocyst (supernatant) and photosymbiont (pellet)
counts by hemocytometer (BOECO, Germany); cells were counted in five randomly selected
fields per fragment (1 mm^2 each) on a Nikon Eclipse Ti-S Inverted Microscope System.
Nematocysts were counted under white light with a contrast filter whereas photosynthetic
symbionts were viewed by fluorescence at 440 nm excitation and 590 nm emission. NIS
ELEMENTS (Nikon) software was used for the cell counts with automatic counting settings
limited to cell diameters smaller than 15 µm and circularity set to >0.5 but <1.
197
198   *3.2.4 Total protein and host protein*
199   We quantified homogenate and host total protein; sonication (Ultrasonic Atomizer Probe
200   Sonicator) was used for further extraction of symbiont protein. Protein concentration of each
201   sample was measured by bicinchoninic acid (BCA) assay (Pierce) against a bovine serum
202   albumin standard curve at an absorbance of 540 nm.
203
3.2.4 Hemolysis assay
Hemolytic assays were performed against O Rh positive human blood cells obtained from the
Rambam/Yoseftal Hospital Blood Bank, as described previously (Primor and Zlotkin, 1975).
Briefly, 2 ml of whole blood was diluted to a final volume of 15 ml in PBS (pH 7.4) and
centrifuged at 3000 g for 5 min. The supernatant was removed, and the process was repeated
with the pelleted erythrocytes until the supernatant was clear. Washed erythrocytes were then
resuspended in PBS to a final concentration of 20% (v/v). For each hemolysis assay, 160 µl of
homogenate supernatant was incubated with 40 µl of washed erythrocyte suspension (4%
erythrocytes v/v) at 37°C for 30 minutes in a water bath. At the end of the incubation, 400 µl of
PBS was added, and the assays were centrifuged at 3000 g for 3 min. The supernatant fluid
containing the hemoglobin released from lysed erythrocytes was transferred to 96-well
microplates and the absorbance at 540 nm was determined on a spectrophotometric microplate
reader (Perkin-Elmer). In addition, each experiment was normalized to a positive (100%
hemolysis) and negative control (0% hemolysis) by incubating erythrocytes with DDW and PBS
alone, respectively. HU_{50} was defined as the amount of homogenate required to cause 50%
hemolysis (Bartosz et al., 2008) in a dilution series of the coral extract.
220
3.2.5 Fragment surface area
Surface areas were estimated using the aluminum foil method (Marsh Jr, 1970). Briefly,
aluminum foil was molded over the skeleton of each fragment (without measuring skeleton

connective areas between the divided fragments), carefully removed, and weighed. This was
repeated three times on every fragment and the surface area was estimated using a standard curve
of the derived relationship between foil area and weight.
227
228 *3.2.6 Skeleton cleaning for assessments of surface area and corallite measurements*
229 After tissue extraction, all fragments were cleaned of organic residues overnight in 3% sodium
230 hypochlorite; the skeletons were then washed in deionized water and dried at 55°C.
231
3.2.7 Micromorphological analysis
The distance between neighboring polyps and the surface area of the polyp was measured using a
Nikon binocular microscope and analyzed using NIS-Elements (Nikon) software. For each coral
fragment, at least four corallites located in a position parallel to the focal plane were chosen.
Major and minor axes and circularity were measured for each corallite and the ratio of the two
axes (major divided by minor) was calculated; a ratio of 1 indicates a circle and the larger the
ratio, the more elliptical the shape. Surface area of each measured corallite was calculated from
the two-measured axes, using the following equation: $S = \pi * r \text{ (major axes)} * r \text{ (minor axes)}$.
Finally, distances between neighboring corallites were also measured to evaluate polyp density
in each area. Corallites were qualitatively analyzed by scanning electron microscopy.
242

3.3 Differential Gene Expression Analysis
3.3.1 RNA extraction, processing, and sequencing
Nine branches, each segmented into three portions along their axes, yielded 36 transcriptome
libraries across the three-dimensional space of the coral colony (Figure 1, colony locations in
Supplementary Table 3, read count statistics in Supplementary Table 4). Total RNA was
extracted from the holobiont in fragments stored in TRI-Reagent (Sigma) following the
manufacturer's protocol, with some modification at the homogenization step. Briefly, samples
frozen in TRI-Reagent were heated and centrifuged, and then bromochloropropane, at a ratio of
1:10, was added to the samples for separation. After incubation at room temperature for 10
minutes and centrifugation, RNA was purified from the clear phase using a Purelink RNA Mini
Kit (Ambion) according to the manufacturer's protocol. The RNA was washed in 70% ethanol
and then on-column DNase digestion was performed using a Qiagen RNase-free DNase Kit
according to the manufacturer's protocol. RNA-Seq libraries were prepared using an in-house
protocol at the Weizmann Institute of Science. Briefly, the polyA fraction (mRNA) was purified
from 500 ng of total RNA per sample followed by fragmentation and generation of double-
stranded cDNA. Then, end repair, A base addition, adapter ligation and PCR amplification steps
were performed. Libraries were evaluated by Qubit (Thermo Fisher) and TapeStation (Agilent).
Sequencing libraries were constructed with barcodes to allow multiplexing of all samples to be
run in each lane. Approximately 578 million total high-quality 125 bp paired end reads (15.45 ±
1.5 million paired reads per sample) were sequenced on an Illumina HiSeq 2500 across three
different lanes (i.e., each sample run in triplicate to remove batch effects).
264

265 *3.3.2 Host and symbiont RNA-Seq differential expression analysis*
266 Standard RNA-Seq quality filtering was conducted as described previously (Malik et al., 2021).
267 Briefly, RNA-Seq reads were adapter-trimmed using cutadapt 1.15
268 (https://cutadapt.readthedocs.io), and then low-quality regions were removed with Trimmomatic
269 0.3 (Love et al., 2014). In order to detect correct taxonomic classifications within our

transcriptome data, high-quality reads were mapped to all available NCBI and Reefgenomics (reefgenomics.org; (Liew et al., 2016)) genome-based proteome databases of Symbiodiniaceae species *Symbiodinium microadraticum, Cladocopium goreaui*, and *Fugacium kawagutii* (formerly *Symbiodinium* spp. clades A, C1, and F, respectively (LaJeunesse et al., 2018)), and Cnidaria as well as selected stramenopiles/alveolates/Rhizaria and Metazoa databases using Diamond (Buchfink et al., 2015). Top hits almost exclusively belonged to robust corals (mainly *S. pistillata*), and *S. microadriaticum* (formerly "clade A" (LaJeunesse et al., 2018)). These results are expected since *S. microadriaticum* was found to be a predominant symbiont in shallow water in Eilat (Malik et al., 2021). For symbiont RNA-Seq mapping, we further aligned the reads to the merged host genome assembly (NCBI GCA_002571385.1) and the symbiont genome assembly (NCBI GCA_001939145.1) using STAR (Dobin et al., 2013;Aranda et al., 2016;Voolstra et al., 2017). Across all branch fragments, 72-80% (~11-16 million reads) were concordantly mapped to the host genome, and 2%-10% (~0.5-1.5 million reads) were concordantly mapped to the symbiont genome. In total, 83-84% of the reads were aligned to either host or symbiont genomes. Differential expression (DE) analysis was conducted using Bioconductor DEseq2 (Love et al., 2014), separately for the host and the symbiont genes, using a DEseq2 generalized linear model. Specifically, we tested: (1) the additive effect of branch position, ring height and cardinal direction; (2) the additive and interaction effect of branch position and ring height (excluding cardinal direction factor); (Supplementary Tables 5-7). Note that the first model cannot include the four top branch samples (two tips, junction, base), since the single top branch is represented by an incomparable cardinal direction ("center"); as such, these samples were excluded from the analysis of expression of putative toxins and known biomineralization-related genes. For the symbiont analysis, only the first model was used. Although DESeq2 is designed to analyze samples of variable total read counts, in order to avoid read normalization problems, we only selected symbiont genes whose 25% quantile of read count was at least 12 reads (with 12,735 symbiont genes remaining); further, only symbiont samples with >300k total reads were retained. After filtering, the correct symbiont read normalization was indicated by a typical close-to-symmetric distribution of fold-changes around the DESeq2 MA plot horizontal axis (Supplementary Figure 2). NMDS analysis was conducted using metaDMS in the R Vegan package based on $\log_{10}FPM$ values of all expressed genes. For the symbiont NMDS, since read counts were relatively low, pairs of tips from the same branch are represented together based on the sum of read counts for both tips; however, symbiont DESeq2 data were not pooled for differential gene expression analysis.

3.3.3 Branch genetics and SNPs analysis

Single nucleotide polymorphisms (SNPs) analysis was conducted for each sample using the Broad Institute-recommended RNA-Seq SNPs practice (https://gatk.broadinstitute.org). The pipeline includes four steps: STAR reads mapping, a pre-processing step aimed at removing various alignment biases, variant calling using GATK version 3.5, and finally variant filtration and annotation (DePristo et al., 2011). In order to exclude variant-call biases due to changes in transcript abundance between samples, we considered only SNP loci with coverage greater than 30 RNA-Seq reads in all tested samples. Identity By Decent (IBD) analysis was conducted using SNPRealte in R.

3.3.4 Host and symbiont functional enrichment analysis

Biological terms were assigned to genes based on Uniprot *S. pistillata* (www.uniprot.org),
KEGG *S. pistillata* (www.kegg.jp)*, S. pistillata* Trinotate annotations (Bryant et al., 2017), and
Uniprot *S. microadriaticum* databases. Enrichment analysis was conducted in Bioconductor
GOSeq (Young et al., 2012), which corrects for enrichment biases associated with correlations
between gene size and DE significance. We also searched for functional enrichment using the
score-based tool GSEA (Subramanian et al., 2005;Sergushichev, 2016), which may be more
sensitive than p-value cutoff-based searches (such as GOSeq) when small non-significant
changes in relatively large groups of genes are expected. In GSEA, we used $\log_2$ fold changes as
scores. Because many enriched terms are functionally related, we hierarchically clustered the
biological terms based on pairwise distances between groups of genes: D =
$|xa\cap xb|/minimum(|a|,|b|)$, where a,b are two sets of gene ids, and xa and xa are DE genes ids
from a and b, respectively. Terms trees were constructed using Bioconductor-ggtree (Yu et al.,
2018). Terms enriched in ring height comparisons to the top of the colony should be viewed with
caution as only one branch was taken from that classification. Only significant enrichment results
are reported.

3.3.5 Focused S. pistillata *gene analyses*
Putative toxin genes were identified using reciprocal best-BLAST hits as described in
(Rachamim et al., 2015;Yosef et al., 2020) (Supplementary Table 14).Differential expression of
genes coding for known biomineralization-related proteins from *S. pistillata* skeleton or the
calicoblastic layer were examined (Puverel et al., 2005;Drake et al., 2013;Mass et al.,
2013;Zoccola et al., 2015;Peled et al., 2020) (Supplementary Table 16).

3.4 Statistical Analyses
Statistical tests of all skeletal, physiological, and environmental data described above were
performed in RStudio (RStudio Team). If a parameter displayed both homogeneity of variance
(Fligner-Killeen test) and normal distribution (Shapiro-Wilke test), a one-way ANOVA
(chlorophyll/total protein, chlorophyll/photosymbiont cell, photosymbiont/surface area, total
protein/surface area, host protein/surface area, nematocysts/total protein, nematocysts/surface
area, corallite diameter, corallite distanc) was performed, with TukeyHSD post hoc analysis for
parameters with significant differences between locations. Parameters with non-homogenous
variance and/or non-normal distribution were analyzed using Wilcoxon's rank sum test (water
column microbes, DO) or the Kruskal-Wallis rank sum test (chlorophyll/surface area,
photosymbiont/total protein, hemolytic units/total protein, hemolytic units/surface area, corallite
axis ratio), with Dunn's test for post hoc analysis for parameters with significant differences
between locations. Environmental parameters were pooled between all colonies and tested
between measured locations as described above. Coral and photosymbiont physiological and
morphological parameters were tested between branch axis locations across the colony (i.e., five
replicates each of junctions and bases, and 10 replicates of tips); however, there was not
sufficient replication at cardinal directions or ring height for statistical evaluation of parameters
between these locational classifications.

4   Results and Discussion
4.1 Differences in water DO and microbial counts across colony structure
Because branching corals' three-dimensional structure can lead to different micro-environments
within the colony, we first aimed to characterize indicators of the environmental conditions

experienced by polyps at different locations within the colony that could potentially affect their
gene expression. Rather than making direct measurements of flow within *S. pistillata* colonies at
the IUI reef, we instead measured several within-colony water parameters affected by flow. We
focused on dissolved oxygen because its concentration can vary within coral colonies, affecting
coral physiology (e.g., Mass et al., 2010). Likewise, microbial counts represent an easy to
quantify variable that can be used as a proxy of water stability. We measured higher oxygen
concentrations in water collected close to the core (center; ~250 μmol/L) of the corals, compared
to samples collected at colony peripheries (~230 μmol/L; $p<0.05$, Wilcoxon rank sum test;
Figure 2a; Supplementary Tables 1, 2). Although there was variability in DO signal between
colonies (Supplementary Figure 3), the 'center' versus 'periphery' differences held across all
colonies. We also observed higher microbe densities in water collected from the colony interiors
versus the immediately surrounding seawater ($p<0.05$, Wilcoxon rank sum test; Figure 2b;
Supplementary Tables 1, 2). Both of these observations may be due to lower rates of water
exchange at branch bases due to decreased water flow within the colony, as suggested by (Chang
et al., 2009). Lower mass transport has been observed within the less densely branched
Pocillopora eydouxi whose overall colony structure is similar to the *S. pistillata* studied here
(Figure 1a) compared to the more densely branched *P. meandrina* (Hossain and Staples). Flow
reduction within the colony would cause the accumulation of oxygen produced by
photosynthesis (Mass et al.) and we speculate might also support the retention of microbes
actively growing between branches (e.g., Schiller and Herndl;Ritchie, 2006), or those
chemotaxing toward the coral tissue (Garren et al., 2014). These differences further support the
notion that the environment differs along the coral branches from colony exterior to interior and
may affect the coral holobiont with some polyps inhabiting more or less favorable areas, which
we observe being reflected in transcriptomic and physiological differences across the colony.
385
4.2 Location-specific intracolonial variability of physiology and gene expression
To assess how the differences in environmental conditions resulted in different physiology and
gene expression across a colony, immediately after measuring DO and collecting microbial
samples we removed a single *S. pistillata* colony from the IUI reef, We then examined its
branches both at different cardinal directions and heights above the substrate (i.e., ring height),
with each branch further segmented along its axis from base to tip (Figure 1 inset;
Supplementary Table 3). The identified patterns in gene expression and physiology therefore
represent responses to microenvironments experienced by the colony at the specific time of
collection and depth at which the colony was growing. The *S. pistillata* colony examined here
was genetically homogenous (not chimeric, Supplementary Figure 4), and thus any changes in
physiology or gene expression between regions were likely a response of the polyps to their local
micro-environmental conditions rather than genotypic differences. While our study was limited
to a single colony to achieve high spatial resolution of data, many similar genetic patterns are
conserved between two major stony coral clades (see below). Additionally, differences were
observed in hemolytic potential between branch tips and bases, which were seen also in two
additional colonies from an independent study (Ben-Ari et al.). Taken together, it is clear that
there are transcriptional and physiological differences between various locations within the coral
colony, and that many of these differences are likely to be common across branching colonial
corals. The expression of >10,000 of the ~24,000 tested host genes was affected by at least one
of the tested factors: location at branch, ring height, and cardinal direction (Supplementary
Tables 4-6). We observed a clear differentiation in host gene expression between branch

positions and from the top to the bottom of the colony (Figure 2c, Figure 3, Supplementary Table 5), although comparisons to the lone top branch should be viewed with caution. Combined, we observed an interaction effect between factors (Supplementary Tables 4, 6), as, for instance, groups of genes involved in signaling pathways important in development that were significantly enriched in branch tips versus bases as compared to middle versus bottom interactions (Figure 3b, Supplementary Figure 5). Comparing colony interior with periphery, host gene expression patterns are more similar between junction and base branch positions compared to tips, although not the same at junctions and bases (Figure 2c). In contrast to the roughly ½ of all tested genes showing differential expression between location classifications, the only physiological parameters that differed along branch axis were hemolytic potential and skeleton morphology (described in detail below). Elucidating the microenvironmental drivers of these genetic and physiological differences across a colony may aid in, for example, understanding spatial differences of polyps' interactions with various infectious organisms that target the oral ectoderm (e.g., Gibbin et al.) leading to diseases such as skeletal band eroding disease (Winkler et al., 2004), which has been observed to commence at colony interiors. Other previously reported response differences across colonies that may be due to microenvironmental variations include bleaching that commences at colony perimeters in *Oculina* (Shenkar et al., 2005), greater storage of lipids and total fatty acids toward colony interiors in *Acropora* (Conlan et al., 2018), and differing relative abundance of various photosymbiont species in regions that later displayed differential bleaching susceptibility (Kemp et al., 2014).

In contrast to the host, of ~12,000 tested photosymbiont genes, only ~150 were differentially expressed (DE) (Supplementary Table 7). The host vs. symbiont differences are partly attributed to technical differences in statistical power due to lower symbiont read counts, and likely represent an underestimation of symbiont DE genes. DE gene patterns of the photosymbionts are not as clearly delineated as for the host (Figure 2d); nevertheless, the expression patterns differ between cardinal directions ($p<0.0007$, ADONIS test, Supplementary Table 8), likely reflecting the different light regimes experienced on different sides of the colony, visible as shading on the north (right) side of the colony and no shading on the south (left) side of the colony in Figure 1a, although we did not measure light levels around the colony. The ultimate drivers of the differences observed for groups of polyps in locations differing by ring height and cardinal direction remain to be resolved. Further, the observed differences across a single colony could have implications for studies using branches collected from different locations on separate colonies. While intracolonial differences that may diminish observable transcriptional and physiological differences between treatments - if chosen branches are from differing parts of selected colonies - may lessen over multi-year studies (e.g., Palumbi et al., 2014), epigenetic responses that differ across a colony may persist across generations (Liew et al., 2020).

4.3 Patterns in gene expression are conserved between coral taxa
A previous study, analyzing the differences in gene expression between branch tips and bases in *Acropora* spp. (Hemond et al., 2014), provided an opportunity to identify conserved intra-colonial patterns in gene expression between these two coral taxa, which likely diverged at least 200 million years ago (Park et al., 2012). We identified ~9,600 putative orthologous expressed genes between *S. pistillata* and two species of *Acropora* spp. (Hemond et al., 2014), based on reciprocal BLAST. Interestingly, from a total of 952 genes that were DE in *Acropora* spp. along the branch axis, ~70% were also DE in *S. pistillata* (672), and ~90% of these exhibited the same

fold change trend (p-value < 2.2e-16, Fisher's exact test; Figure 3a, Supplementary Table 9).
Overall, these results indicate that micro-environmental dependence of gene differential
expression at tips versus bases reflects an evolutionarily conserved pattern rather than a species-
specific pattern, being observed in members of both robust (here) and complex (Hemond et al.,
2014) scleractinian clades, as well as between taxa which possess physiologically unique axial
polyps (*Acropora*; (Wallace, 1978;1985) and a genus which does not (*Stylophora*). It also
supports that the occurrence of intracolonial differential responses to microenvironments in
general, and up-regulation of genes for specific processes such as protein production and energy
generation in particular (Figure 3b, green font), can likely be generalized across branching coral
taxa.
463
4.4 Gene expression patterns suggest differences in host versus photosymbiont metabolism
We next inferred potential broad-scale changes in cellular and metabolic functions by identifying
KEGG and Gene Ontology (GO) enrichment patterns associated with micro-environmental
changes in both the host (Figure 3b, Supplementary Tables 10, 11) and the photosymbiont
(Figure 3c, Supplementary Table 12). As expected, broad clusters of host KEGG terms and
photosymbiont GO terms were DE (Figure 3b, c), mirroring the pattern observed for all DE
genes within the host and symbiont (Figure 2c, d), with many more terms DE in the coral host
along the branch axis compared to the ring height or cardinal direction (Figure 3b), and more
terms differed along cardinal direction for the photosymbiont (Figure 3c).
473
Most DE host genes grouped by KEGG term tended to be up-regulated in branch tips relative to
the junctions or bases (Figure 3b, Supplementary Tables 10, 11). Specifically, many terms
related to cell metabolism were enriched in the tips versus the bases, including carbohydrate
metabolism (e.g., glycolysis and the TCA cycle, oxidative phosphorylation, mitochondrial
biosynthesis), amino acid biosynthesis and metabolism, and lipid metabolism (Figure 3b). We
further tested which functional groups were enriched with conserved tip vs. base DE gene
patterns between coral taxa (Figure 3a, b). As indicated by green KEGG term text in Figure 3b,
many fundamental processes such as metabolism show similar DE trends and enrichment
between these two genera (Supplementary Tables 9, 10), indicating a conserved response
between the taxa. "Zooming in" from the KEGG terms to the underlying genes, we note that
genes related to oxidative phosphorylation, such as ATP synthase, cytochrome c oxidase, co-
enzyme Q, and NADH dehydrogenase tend to be up-regulated at branch tips across the species.
In addition, multiple genes for succinate dehydrogenase as well as other individual genes
comprising the citric acid cycle, were also more highly expressed at branch tips than junctions or
bases in *S. pistillata* and many of these showed similar regulation patterns in the *Acropora* spp.
Finally, most of the >100 DE genes with KEGG terms for mitochondrial biogenesis were up-
regulated at tips compared with junctions and bases in *S. pistillata* and many of their likely
homologs were also up-regulated at branch tips in *Acropora* spp. The increased rates in
physiological processes involved in energy production at branch tips, as suggested by our gene
expression analysis, have been observed in the branching coral *A. palmata* as increased
respiration and mitotic index (Gladfelter et al., 1989).
495
Intriguingly, the patterns of gene expression observed in the host were not all recapitulated in the
photosymbiont. Although some symbiont GO terms associated with photosynthesis were
enriched in the tips vs bases in the symbionts (e.g., those involved in light-harvesting and the

thylakoid), likely in response to higher light intensities at tips compared to bases (e.g., Helmuth
et al., 1997;Kaniewska et al., 2008), others were not, including terms involved with carbon
fixation (Figure 3c). Additionally, no differing patterns in photophysiology were found along
branch axis, measured here as chlorophyll and symbiont density per both total protein and area
(Supplementary Tables 2, 13, Supplementary Figure 6), although we acknowledge that these are
limited measures of the photophysiological processes. Indeed, previous studies have shown that
the products of photosynthesis can be translocated from branch bases to the tips in branching
corals (e.g., Rinkevich and Loya). Rather than exhibiting tip-to-base functional differences,
many of the terms associated with symbiont photosynthesis (both light and dark reactions) were
found to differ along cardinal direction, being enriched in south-facing branches (Figure 3c),
possibly due to the direction of the sun. As such, we anticipate that future studies will find
differences in gene expression in the symbiont across a diel cycle, with, for example potentially
decreased differences between north- and south-facing branches during nighttime hours. As
noted above, these enrichment patterns are likely an underestimation. Furthermore, correlations
between mRNA abundance and physiological patterns may not be high due to translation level
effects, translocation of host/symbiont products, food particle capture by the host, etc., which
may lead to differences in symbiont transcriptional control.
516
4.5 Developmental pathways are DE along the branch axis
In addition to pathways related to metabolism, several KEGG terms associated with development
and morphological plasticity in colony architecture were DE between branch tips and junctions
or bases. For example, the Wnt signaling pathway was enriched at branch tips versus the bases in
both *S. pistillata* (Figure 3b) and *Acropora* spp. (Hemond et al., 2014). This pathway, which
transmits information from outside to inside cells, is associated with cell differentiation and axis
patterning in cnidarians (reviewed in Lee et al., 2006;e.g., Duffy et al., 2010) and bilaterians
(e.g., Siegfried and Perrimon). Similarly, the TGFβ signaling pathway was also enriched in *S.*
pistillata branch tips (Figure 3b), although this was not observed in *Acropora* spp. (Hemond et
al., 2014); this pathway has been implicated for a role in development and calcification in corals
(Gutner-Hoch et al., 2017). At the individual gene level, we observed that fibroblast growth
factors, which could aid in development of limbs (reviewed in Xu et al., 1999) and in production
of nerve cells (reviewed in Guillemot and Zimmer, 2011), were significantly up-regulated at
branch tips. Half of the genes involved in mitochondrial biosynthesis were also up-regulated at
branch tips. Further, two different bone morphogenetic proteins (BMPs), which may play a role
in axial patterning in cnidarian polyps (Reinhardt et al., 2004) and in budding (Surekha et al.,
2020), were up-regulated at branch tips here. Finally, several retinol dehydrogenases, proteins
associated with photoreceptors in higher organisms (Albalat, 2012), were up-regulated at branch
tips, where light levels tend to be higher, an observation shared with *Acropora* spp. (Hemond et
al., 2014).
537
In addition to the plasticity described above, gamete production has also been shown to vary
along branches axes across several coral taxa with gametogenesis typically occurring
preferentially deeper in the colony (e.g., Stimson). We observed the highest expression of several
testis-specific genes, such as testis-specific chromodomain protein Y 1-like and testis-specific
protein kinases at *S. pistillata* branch junctions (Supplementary Table 4). In humans, these genes
are associated with chromosome Y (Lahn and Page) and/or may be involved in sperm structure
(Walden and Cowan). Interestingly, we observed up-regulation of octopamine receptors at

branch tips, suggestive of higher maturation of ovaries and testes in that location (Chiu et al.,
2020).
547
4.6 Changes in expression of genes potentially related to predation along branch axis
As described above, many metabolic processes were enriched in the tips versus the bases,
whereas there is little evidence for higher photosynthesis by photosymbionts at branch tips.
Asking whether this could be due to more heterotrophic food input in the tips (e.g., through
predation), we analyzed the tissue toxicity (hemolytic activity) and gene expression patterns of
putative toxins. Although there was no difference in nematocyst abundance along branch axes
(Figure 4a, Supplementary Table 2), tissue hemolytic activity was highest at branch tips (Figure
4b, Supplementary Table 2), consistent with previous studies in *S. pistillata* that used branches
from two different colonies (Ben-Ari et al.). Thus, the observed branch axis patterning of
hemolysis potential is robust across at least three colonies from two independent studies. Using
reciprocal BLAST, we identified 22 putative toxin genes (Rachamim et al., 2015), encoding
cytolysins, phospholipase A2 toxins, SCRiPs (small cysteine rich peptides) and a potential
Kunitz/K channel toxin (Supplementary Table 14). We also compared these genes to the
reciprocal BLAST results against the *Acropora* spp. dataset (Hemond et al., 2014), finding only
two orthologous matches, a SCRiP and an uncharacterized protein. Like their *S. pistillata*
counterparts, they were not DE between branch tips and bases (Supplementary Table 15). In our
putative toxin dataset, we observed 13 and 11 DE genes between branch tips versus bases or
junctions, respectively (Figure 4c, d; Supplementary Table 15, Supplementary Figure 7). Up-
regulation of genes encoding potential toxins at branch tips has previously been observed in
colonial corals (Hemond et al.;Hemond and Vollmer). The two most abundantly expressed
putative toxin genes encode for a cytolysin (up-regulated at tips) and a second SCRiP (up-
regulated at bases), with the rest of the toxin genes having much lower overall expression levels
(Figure 4c, d; Supplementary Table 15). The highly expressed cytolysin is of particular interest
as it belongs to the actinoporin family, and a hemolytic protein from this family, likely the
protein or an isoform of the highly expressed gene here (Supplementary Figure 8), has
previously been isolated from *S. pistillata* (Ben-Ari et al., 2018). Actinoporins are ~20 kDa
water-soluble proteins which are lethal when injected into crustaceans (Giese et al., 1996), and
which insert themselves into membranes, particularly those containing sphingomyelins (Alvarez
et al., 2009). They have been immunolocalized to tentacle nematocysts as well as to mesenterial
filaments (Basulto et al., 2006). SCRiPs are ~5 kDa peptides which may be anthozoan-specific
(Sunagawa et al.;Jouiaei et al., 2015), with evidence that they have neurotoxic effects (Jouiaei et
al., 2015), although their function remains to be fully clarified.
580
The higher hemolytic activity and larger number of toxins more abundantly expressed at branch
tips suggest that the tips play the dominant role in venom-based competition, defense, and
predation, although more work is needed to identify the role of putative toxins such as SCRiPs,
which were more abundantly expressed at branch bases, and the potential contribution and roles
of photosymbiont toxin production (e.g., Nakamura et al., 1993). While it is difficult to
disentangle toxin effects on competition and defense from predation, based on higher flow at the
edges of colonies (Chang et al., 2009), colonies' edges are likely exposed to higher food loads
than are areas closer to the base of the branches. Consumption of these food sources could then
be reflected in increased respiration rates as the food is metabolized, as has been observed in
other species (Gladfelter et al., 1989) and may yield increased energy production which would

require up-regulation of genes involved in this process as we observed here. We also queried genes related to digestion and observed that many peptidases were DE, but not in a clear pattern along the branches with approximately half up-regulated at the tips. Similarly, multiple lipases were DE between branch locations, but no general trend was observed. However, pancreatic lipase expression, which has been observed in gastric cirri in a scyphozoan (Steinmetz, 2019), was found to be significantly up-regulated to a high degree at branch tips. We also observed up-regulation of three chitinases in branch tips versus bases, which could support extracellular digestion of arthropod prey (Steinmetz, 2019). In contrast, an amylase which converts starch and glycogen - biomolecules used by the photosymbiont and host, respectively, to store photosynthates (Kopp et al., 2015) - into simple sugars, was down-regulated at the tips versus junctions and bases. Thus, while our results suggest differences along the branch axis of functions related to prey capture and defense, including both toxicity and digestion, further work is needed to fully characterize the relative contributions of photosynthesized products and predation. Additionally, temporal differences in predation capabilities at both the transcriptional (i.e., genes related to digestion) and physiological (i.e., hemolytic potential) levels across individual colonies should be examined as zooplankton abundance on coral reefs changes on a diel cycle (Yahel et al., 2005). Future studies measuring water flow rates explicitly in conjunction with transcriptional changes across colonies with complex macromorphologies will also clarify precisely how particle capture and digestion are affected by differing flow patterns (e.g., Sebens et al., 1997).

4.7 Changes in skeleton structure and expression of known biomineralization genes along branch axis

Shallow water stony corals are most notable for their production of massive reef structures, growing in size by adding mineral and new polyps, particularly for branching corals, at colony tips or edges (e.g., D'A and Le Tissier, 1988;Meesters and Bak, 1995). We therefore examined in depth skeletal morphological patterns and related gene expression across the three branch axis locations. Unlike most other physiological parameters across the colony which were not significantly different between branch positions (Supplementary Tables 2, 13), those linked to differences in biomineralization were significantly different along branch axes. Qualitatively, coenosteal spines appear narrowest and thinnest at the tip while base spines exhibit a more distinct microtuberculate texture in their lower portions (Figure 5aii, iv, and vi). Such microtuberculate texture with better differentiation of mineral fiber bundles may suggest higher organic matrix content which separates the mineral phase packages to a greater extent (Mass et al., 2014;Coronado et al., 2019). Corallites were closer together and more elliptically shaped at the tips than at the branch junctions or bases (Figure 5b and c; Supplementary Tables 2, 13). Comparable trending toward ellipticity at branch tips has been reported in other Pocilloporids (Schmidt-Roach et al., 2014). Further, genetically identical *S. pistillata* branches have been shown to have more distantly spaced corallites with thicker theca and longer septa when grown under light spectra biased toward blue wavelengths (Rocha et al., 2014). While we did not measure light spectra in this study, such a spectral shift along a single branch, due to absorbance of red light by water, has been observed in *S. pistillata* tissue at the colony level on the sides versus at the top of ~30 cm diameter colonies (Kaniewska et al., 2011) and at the branch level along *Acropora millepora* branches toward the colony interior (Wangpraseurt et al., 2014).

We next examined differential expression of genes encoding proteins sequenced from *S. pistillata* skeleton and likely directly involved in the calcification mechanism (Puverel et al., 2005;Bertucci et al., 2011;Drake et al., 2013;Mass et al., 2014;Peled et al., 2020;Mummadisetti et al., 2021), as well as a bicarbonate anion transporter observed predominantly in the calicoblastic cell layer (Zoccola et al., 2015) (Figure 5e; Supplementary Table 16, Supplementary Figure 9). We did not include genes for known skeletal proteins from *Acropora millepora* and *A. digitifera* (Ramos-Silva et al.;Takeuchi et al.), as not all orthologs have been found in *S. pistillata* skeleton although they exist in the genome, or non-orthologous proteins within the same functional group occur in *S. pistillata* skeleton (Zaquin et al., 2021). Similar to the global gene expression pattern for the host, we observed an up-regulation of biomineralization genes biased toward the tips (Figure 5d, e; Supplementary Table 16). Of the 36 biomineralization-related genes that were DE between branch tips and bases, eight orthologous genes were also DE and in the same direction in *Acropora* spp. (Hemond et al., 2014) (Supplementary Table 17). In *S. pistillata*, the bicarbonate anion transporter SLCγ4, a coadhesin-like protein, uncharacterized skeletal organic matrix protein 8 (USOMP8), and a hypothetical protein were up-regulated at branch tips but were expressed in only low amounts. In contrast, a ZP-domain containing protein, carbonic anhydrase (STPCA2), mammalian ependymin-related protein, and coral acid rich protein 3 (CARP3) were moderately up-regulated at branch tips and were very highly expressed overall. These proteins exhibit very different functions with SLCγ4 and STPCA2 involved in regulating carbonate chemistry within the calcifying space (Zoccola et al., 2015;Zoccola et al., 2016), while CARP3, which is a highly acidic protein (35% Asp, 15%Glu (Mass et al., 2013)) can direct mineralogy toward formation of Mg-calcite and vaterite (Gavriel et al., 2018;Laipnik et al., 2020), and is up-regulated in newly-settled coral polyps (Akiva et al., 2018), in colonies growing at shallow depths (Malik et al., 2021), and in coral cell cultures reared at increased pCO_2 (Drake et al., 2017). Although the function of USOMP8 is not known, it is structurally similar to a BMP inhibitor (Drake et al., 2013). Of the remaining known highly acidic coral skeleton genes, CARPs 1 and 4 (CARP4 is sometimes also called skeletal aspartic acid rich protein 1; SAARP1 (Ramos-Silva et al., 2013)) were up-regulated at branch tips and were moderately expressed; both proteins are biased toward Asp residues (Mass et al., 2013) and are likely involved in nucleation initiation in early mineralization zones of growing septa (Mass et al., 2014). In contrast, CARP2 exhibited very low although significantly up-regulated expression at branch tips; while this protein is suggested to have a role in skeleton extension based on its localization to more soluble growth areas (Mass et al., 2014), its down-regulation upon settlement in pocilloporid larvae (Mass et al., 2016;Akiva et al., 2018), association with ACC phases pre-settlement (Akiva et al., 2018), and low overall expression in the present study may instead point to a role in inhibiting aragonite nucleation.

Previous and our present work provide strong genetic support for highest biomineralization activity at branch tips (e.g., Goreau, 1959). This faster calcification at the tips may result in the qualitatively thinner and narrower coenosteal spines we observed at branch tips (Figure 5) that would be thickened by polyps as they age. Energy for higher calcification rates at branch tips could come from two sources. First, there is evidence of translocation of ATP from tissue regions deeper in the colony with relatively high photosynthetic rates (Fang et al., 1989) to branch tips. Secondly, our and previous data suggest that the potential for predation is higher at branch tips, with higher toxin gene expression and hemolytic activity at that location (Figure 4) (Hemond et al., 2014;Ben-Ari et al., 2018), and up-regulation of genes related to metabolism and

682   energy production (Figure 3). This further suggests that biomineralization gene expression and
683   skeleton morphology patterns observed at the sub-polyp scale (i.e., centers of calcification in
684   growing tips of septa) are recapitulated at the branch scale (Malik et al., 2021).
685
5   Summary and Conclusions
Some colonial cnidarians, particularly hydrozoans, exhibit a clear division of labor inside the
colony, with functionally and morphologically distinct polyps across the colony (Cartwright et
al., 1999). While no obvious division of labor at the polyp scale is observed in corals, our results
show, from molecular to physiological and morphological scales on a single genetically-identical
colony, a functional specialization by clusters of polyps due to their locations on coral branches
and around the colony likely related to the microenvironments they encounter (Rueffler et al.,
2012) (Figure 6). In simplest terms, polyps at branch tips and toward the top of the colony are
more exposed to light and higher water flow than are those found lower in the branches and
closer to the substrate with consequent differential expression of a multitude of genes directing
many key functions and, at least along the branch axis, differences in physiology and skeleton
morphology. Our findings suggest that branch tips may also be sites of higher potential for
envenomation for colony defense and predation, as well as possibly for enhanced prey capture.
These plastic responses by genetically identical polyps can then be communicated through the
coenosarc to the rest of the colony to allow for enhanced resource sharing (Mackie, 1986). Such
plasticity may aid in colony recovery after partial mortality events, such as those due to
bleaching (Matsuda et al., 2020), and may be integral to adaptation to environmental change
(Kenkel and Matz, 2016). Further, the expression patterns underlying the tested micro-
environmental differences are found to be evolutionary conserved in the distantly-related stony
corals *S. pistillata* and *Acropora* spp. Future study of multiple colonies and species will clarify
the extent to which the general patterns we observe across the colony studied here are found
across differing colony morphologies, diurnal and seasonal cycles, and geographic location.
Further, the diversity of gene expression and physiology across a coral colony needs to be taken
into account when designing experiments with coral fragments from multiple colonies.
710
6   Figure Legends
Figure 1. **Micro-environments and responses in sections of an individual coral colony.**
Photograph of the *S. pistillata* colony analyzed for physiological and transcriptomic intracolonial
differences taken *in situ* (8 m depth) from the east side of the colony (a). Three-dimensional
reconstructions of the coral colony showing locations of branches based on cardinal direction
(top view, b) and ring height (oblique view, c). Inset in (c) shows branch axis sampling locations
at tips, junction, and base per branch.
718
719   Figure 2. In-situ water column concentrations of dissolved oxygen (a) and total microbial cells
720   (b) between branches inside and outside *S. pistillata* colonies. Non-metric multidimensional
721   scaling (NMDS) of all expressed genes of the coral host (c) and photosymbiont (d) with factor
722   effects color- and shape-coded. Host ring and branch position-based clusters are significant
723   (Supplementary Table 7) and marked by dashed and solid ellipses, respectively.
724
Figure 3. **Differential gene expression across coral species and terms clustering in host and**
symbiont. Venn diagram (a) representing the proportional counts of shared tip vs. base DE genes
in *S. pistillata* (here) and two species of *Acropora* (Hemond et al., 2014) (Supplementary Table

9), with shaded regions representing genes showing the same DE patterns and noting "up" and
"down" regulated genes. Enrichment patterns of clustered host KEGG (*S. pistillata*; b) and
photosymbiont GO terms (UniProt *S. microadriaticum*; c) across factor comparisons. Host terms
enriched in both *S. pistillata* (here) and two species of *Acropora* (Hemond et al., 2014) for tip vs
base comparisons, are shown as green text. Hierarchical tree clustering reflects functional
similarity between terms. Color-coded heatmap scores equal to $\log_{10}$(adjusted p-value) of
enriched terms multiplied by 1 and -1 for up- and down-regulated genes, respectively (for clarity,
values <-2 and >2 are shown as 2 and -2).

Figure 4. **Toxicity potential.** Nematocyst abundance (a) and hemolytic units per area (b), and
gene expression patterns of putative toxins as $\log_2$FC between branch tips (positive values) and
bases (negative values) (c), and $\log_{10}$FPKM along *S. pistillata* branch axis (d). Only significantly
DE toxin genes are shown (p<0.05). The two most abundant transcripts, a cytolysin annotated as
stichotoxin and an uncharacterized gene, for which reciprocal BLAST hits suggest it is a SCRiP
(Supplementary14 Table), are noted with a red star (d).

Figure 5. **Skeleton characteristics and biomineralization gene expression.** Microstructural
differences of corallites and coenosteal spines along the branch axis (a; tip in i, ii; junction in iii,
iv; base in v, vi). Corallite spacing (b) and ellipticity (c) along *S. pistillata* branch axis. Several
known biomineralization genes were DE, shown as $\log_2$FC between branch tips (positive values)
and bases (negative values) (d) and $\log_{10}$FPKM along the branch axis (e) to clarify genes with
high versus low expression. Only significantly DE genes are shown (p<0.05).

Figure 6. **Micro-environmental variation across a colony leads to physiological and**
transcriptional differences. Schematic of environmental parameters (e.g., flow and light
intensity, left) and resulting differential gene expression patterns and physiological outcomes for
key processes focusing on those enhanced at branch tips (right). We cannot assign gene
expression to specific tissue layers or cell types in most cases, except for the likely localization
of toxin and biomineralization genes to the oral ectoderm and calicoderm, respectively.

758   7   **Conflict of Interest**
759   *The authors declare that the research was conducted in the absence of any commercial or*
760   *financial relationships that could be construed as a potential conflict of interest.*

762   8   **Author Contribution Statement**
763   YP, OY, DT, DS, and TM conceived the study. YP, OY, ES prepared the samples. JLD, AM,
764   YP, OY, ES, and JS conducted formal analysis of the data. All authors wrote the manuscript and
765   approve of this submission.

767   9   **Funding**
This work was supported by grants 1239/13 and 312/15 from the Israel Science Foundation (to
DS and TM, respectively) and by the associated grant number 2236/16 from the Israeli National
Center for Personalized Medicine (to DS). This work has received funding to TM from the
European Research Council under the European Union's Horizon 2020 research and innovation
programme (grant agreement No 755876). JLD was supported by the Zuckerman STEM
Leadership Program. Computations presented in this work were performed on the Hive

computer cluster at the University of Haifa, which is partly funded by Israel Science Foundation
grant 2155/15.
776
777   **10 Acknowledgements**
778 We thank Yarden Ben-Tabou de-Leon for drawing the summary Figure 6 and Dikla
779 Aharonovich for guidance and laboratory assistance. We thank the Crown Genomics Institute of
780 the Nancy and Stephen Grand Israel National Center for Personalized Medicine, Weizmann
781 Institute of Science for transcriptome sequencing.
782

783   **11 Data Availability Statement**
784 The data underlying this article are available in the article, in its online Supplementary Material,
785 and on GitHub in the following repository:
786 https://github.com/jeanadrake/Single_coral_colony_transcriptomes_physiology.
787

12 References
Akiva, A., Neder, M., Kahil, K., Gavriel, R., Pinkas, I., Goobes, G., and Mass, T. (2018).
Minerals in the pre-settled coral *Stylophora pistillata* crystallize via protein and ion
changes. *Nature Communications* 9**,** 1880.
Albalat, R. (2012). Evolution of the genetic machinery of the visual cycle: a novelty of the
vertebrate eye? *Molecular Biology and Evolution* 29**,** 1461-1469.
Alvarez, C., Mancheno, J.M., Martínez, D., Tejuca, M., Pazos, F., and Lanio, M.E. (2009).
Sticholysins, two pore-forming toxins produced by the Caribbean sea anemone
Stichodactyla helianthus: their interaction with membranes. *Toxicon* 54**,** 1135-1147.
Anthony, K.R., Hoogenboom, M.O., and Connolly, S.R. (2005). Adaptive variation in coral
geometry and the optimization of internal colony light climates. *Functional Ecology* 19**,**
17-26.
Aranda, M., Li, Y., Liew, Y.J., Baumgarten, S., Simakov, O., Wilson, M.C., Piel, J., Ashoor, H.,
Bougouffa, S., and Bajic, V.B. (2016). Genomes of coral dinoflagellate symbionts
highlight evolutionary adaptations conducive to a symbiotic lifestyle. *Scientific Reports*
6**,** 1-15.
Bartosz, G., Finkelshtein, A., Przygodzki, T., Bsor, T., Nesher, N., Sher, D., and Zlotkin, E.
(2008). A pharmacological solution for a conspecific conflict: ROS-mediated territorial
aggression in sea anemones. *Toxicon* 51**,** 1038-1050.
Basulto, A., Pérez, V.M., Noa, Y., Varela, C., Otero, A.J., and Pico, M.C. (2006).
Immunohistochemical targeting of sea anemone cytolysins on tentacles, mesenteric
filaments and isolated nematocysts of *Stichodactyla helianthus*. *Journal of Experimental*
Zoology Part A: Comparative Experimental Biology 305**,** 253-258.
Ben-Ari, H., Paz, M., and Sher, D. (2018). The chemical armament of reef-building corals: inter-
and intra-specific variation and the identification of an unusual actinoporin in *Stylophora*
pistilata. *Scientific Reports* 8**,** 1-13.
Bertucci, A., Tambutté, S., Supuran, C., Allemand, D., and Zoccola, D. (2011). A new coral
carbonic anhydrase in *Stylophora pistillata*. *Marine Biotechnology* 13**,** 992-1002.
Bryant, D.M., Johnson, K., DiTommaso, T., Tickle, T., Couger, M.B., Payzin-Dogru, D., Lee,
T.J., Leigh, N.D., Kuo, T.-H., and Davis, F.G. (2017). A tissue-mapped axolotl de novo
transcriptome enables identification of limb regeneration factors. *Cell Reports* 18**,** 762-
776.

Buchfink, B., Xie, C., and Huson, D.H. (2015). Fast and sensitive protein alignment using
DIAMOND. *Nature Methods* 12**,** 59-60.
Carpenter, L.W., and Patterson, M.R. (2007). Water flow influences the distribution of
photosynthetic efficiency within colonies of the scleractinian coral *Montastrea annularis*
(Ellis and Solander, 1786); implications for coral bleaching. *Journal of Experimental*
Marine Biology and Ecology 351**,** 10-26.
Cartwright, P., Bowsher, J., and Buss, L.W. (1999). Expression of a Hox gene, Cnox-2, and the
division of labor in a colonial hydroid. *Proceedings of the National Academy of Sciences*
96**,** 2183-2186.
Chang, S., Elkins, C., Alley, M., Eaton, J., and Monismitha, S. (2009). Flow inside a coral
colony measured using magnetic resonance velocimetry. *Limnology and Oceanography*
54**,** 1819-1827.
Chiu, Y.-L., Shikina, S., Yoshioka, Y., Shinzato, C., and Chang, C.-F. (2020). De novo
transcriptome assembly from the gonads of a scleractinian coral, *Euphyllia ancora*:
molecular mechanisms underlying scleractinian gametogenesis. *BMC Genomics* 21**,** 1-20.
Conlan, J.A., Humphrey, C.A., Severati, A., and Francis, D.S. (2018). Intra-colonial diversity in
the scleractinian coral, *Acropora millepora*: identifying the nutritional gradients
underlying physiological integration and compartmentalised functioning. *PeerJ* 6**,** e4239.
Coronado, I., Fine, M., Bosellini, F.R., and Stolarski, J. (2019). Impact of ocean acidification on
crystallographic vital effect of the coral skeleton. *Nature Communications* 10**,** 1-9.
D'A, M., and Le Tissier, A. (1988). The growth and formation of branch tips of *Pocillopora*
damicornis (Linnaeus). *Journal of Experimental Marine Biology and Ecology* 124**,** 115-
131.
Dennison, W.C., and Barnes, D.J. (1988). Effect of water motion on coral photosynthesis and
calcification. *Journal of Experimental Marine Biology and Ecology* 115**,** 67-77.
DePristo, M.A., Banks, E., Poplin, R., Garimella, K.V., Maguire, J.R., Hartl, C., Philippakis,
A.A., Del Angel, G., Rivas, M.A., and Hanna, M. (2011). A framework for variation
discovery and genotyping using next-generation DNA sequencing data. *Nature Genetics*
43**,** 491.
Dobin, A., Davis, C.A., Schlesinger, F., Drenkow, J., Zaleski, C., Jha, S., Batut, P., Chaisson,
M., and Gingeras, T.R. (2013). STAR: ultrafast universal RNA-Seq aligner.
Bioinformatics 29**,** 15-21.
Drake, J.L., Mass, T., Haramaty, L., Zelzion, E., Bhattacharya, D., and Falkowski, P.G. (2013).
Proteomic analysis of skeletal organic matrix from the stony coral *Stylophora pistillata*.
Proceedings of the National Academy of Sciences 110**,** 3788-3793.
Drake, J.L., Schaller, M.F., Mass, T., Godfrey, L., Fu, A., Sherrell, R.M., Rosenthal, Y., and
Falkowski, P.G. (2017). Molecular and geochemical perspectives on the influence of CO_2
on calcification in coral cell cultures. *Limnology and Oceanography*.
Duffy, D.J., Plickert, G., Kuenzel, T., Tilmann, W., and Frank, U. (2010). Wnt signaling
promotes oral but suppresses aboral structures in *Hydractinia* metamorphosis and
regeneration. *Development* 137**,** 3057-3066.
Falkowski, P.G., Dubinsky, Z., Muscatine, L., and Porter, J.W. (1984). Light and the
bioenergetics of a symbiotic coral. *Bioscience* 34**,** 705-709.
Fang, L.-S., Chen, Y.-w.J., and Chen, C.-S. (1989). Why does the white tip of stony coral grow
so fast without zooxanthellae? *Marine Biology* 103**,** 359-363.

Garren, M., Son, K., Raina, J.-B., Rusconi, R., Menolascina, F., Shapiro, O.H., Tout, J., Bourne,
D.G., Seymour, J.R., and Stocker, R. (2014). A bacterial pathogen uses
dimethylsulfoniopropionate as a cue to target heat-stressed corals. *The ISME Journal* 8**,**
999-1007.
Gavriel, R., Nadav-Tsubery, M., Glick, Y., Yarmolenko, A., Kofman, R., Keinan-Adamsky, K.,
Berman, A., Mass, T., and Goobes, G. (2018). The coral protein CARP3 acts from a
disordered mineral surface film to divert aragonite crystallization in favor of Mg-calcite.
Advanced Functional Materials 28**,** 1707321.
Gibbin, E., Gavish, A., Domart-Coulon, I., Kramarsky-Winter, E., Shapiro, O., Meibom, A., and
Vardi, A. (2018). Using NanoSIMS coupled with microfluidics to visualize the early
stages of coral infection by *Vibrio coralliilyticus*. *BMC microbiology* 18**,** 1-10.
Giese, C., Mebs, D., and Werding, B. (1996). Resistance and vulnerability of crustaceans to
cytolytic sea anemone toxins. *Toxicon* 34**,** 955-958.
Gladfelter, E.H., Michel, G., and Sanfelici, A. (1989). Metabolic gradients along a branch of the
reef coral *Acropora palmata*. *Bulletin of Marine Science* 44**,** 1166-1173.
Goreau, T.F. (1959). The physiology of skeleton formation in corals. I. A method for measuring
the rate of calcium deposition by corals under different conditions. *Biological Bulletin*
116**,** 59-75.
Guillemot, F., and Zimmer, C. (2011). From cradle to grave: the multiple roles of fibroblast
growth factors in neural development. *Neuron* 71**,** 574-588.
Gutner-Hoch, E., Ben-Asher, H.W., Yam, R., Shemesh, A., and Levy, O. (2017). Identifying
genes and regulatory pathways associated with the scleractinian coral calcification
process. *PeerJ* 5**,** e3590.
Helmuth, B.S., Timmerman, B., and Sebens, K. (1997). Interplay of host morphology and
symbiont microhabitat in coral aggregations. *Marine Biology* 130**,** 1-10.
Hemond, E.M., Kaluziak, S.T., and Vollmer, S.V. (2014). The genetics of colony form and
function in Caribbean *Acropora* corals. *BMC Genomics* 15**,** 1133.
Hemond, E.M., and Vollmer, S.V. (2015). Diurnal and nocturnal transcriptomic variation in the
Caribbean staghorn coral, *Acropora cervicornis*. *Molecular Ecology* 24**,** 4460-4473.
Hoegh-Guldberg, O., Jacob, D., Taylor, M., Bolaños, T.G., Bindi, M., Brown, S., Camilloni, I.,
Diedhiou, A., Djalante, R., and Ebi, K. (2019). The human imperative of stabilizing
global climate change at 1.5° C. *Science* 365**,** eaaw6974.
Hossain, M.M., and Staples, A.E. (2020). Mass transport and turbulent statistics within two
branching coral colonies. *Fluids* 5**,** 153.
Hughes, T.P. (1994). Catastrophes, phase shifts, and large-scale degradation of a Caribbean coral
reef. *Science* 265**,** 1547-1551.
Jouiaei, M., Sunagar, K., Federman Gross, A., Scheib, H., Alewood, P.F., Moran, Y., and Fry,
B.G. (2015). Evolution of an ancient venom: recognition of a novel family of cnidarian
toxins and the common evolutionary origin of sodium and potassium neurotoxins in sea
anemone. *Molecular Biology and Evolution* 32**,** 1598-1610.
Kaniewska, P., Anthony, K.R., and Hoegh-Guldberg, O. (2008). Variation in colony geometry
modulates internal light levels in branching corals, *Acropora humilis* and *Stylophora*
pistillata. *Marine Biology* 155**,** 649-660.
Kaniewska, P., Magnusson, S.H., Anthony, K.R., Reef, R., Kühl, M., and Hoegh-Guldberg, O.
(2011). Importance of macro- versus microstructure in modulating light levels inside
coral colonies. *Journal of Phycology* 47**,** 846-860.

Kemp, D.W., Hernandez-Pech, X., Iglesias-Prieto, R., Fitt, W.K., and Schmidt, G.W. (2014).
Community dynamics and physiology of Symbiodinium spp. before, during, and after a
coral bleaching event. *Limnology and Oceanography* 59**,** 788-797.
Kenkel, C.D., and Matz, M.V. (2016). Gene expression plasticity as a mechanism of coral
adaptation to a variable environment. *Nature Ecology & Evolution* 1**,** 1-6.
Knowlton, N., Brainard, R.E., Fisher, R., Moews, M., Plaisance, L., and Caley, M.J. (2010).
Coral reef biodiversity. *Life in the world's oceans: diversity distribution and abundance***,**
65-74.
Kopp, C., Domart-Coulon, I., Escrig, S., Humbel, B.M., Hignette, M., and Meibom, A. (2015).
Subcellular investigation of photosynthesis-driven carbon assimilation in the symbiotic
reef coral *Pocillopora damicornis*. *MBio* 6.
Lahn, B.T., and Page, D.C. (1999). Retroposition of autosomal mRNA yielded testis-specific
gene family on human Y chromosome. *Nature Genetics* 21**,** 429-433.
Laipnik, R.a., Bissi, V., Sun, C.-Y., Falini, G., Gilbert, P.U., and Mass, T. (2020). Coral acid rich
protein selects vaterite polymorph in vitro. *Journal of Structural Biology* 209**,** 107431.
LaJeunesse, T.C., Parkinson, J.E., Gabrielson, P.W., Jeong, H.J., Reimer, J.D., Voolstra, C.R.,
and Santos, S.R. (2018). Systematic revision of Symbiodiniaceae highlights the antiquity
and diversity of coral endosymbionts. *Current Biology* 28**,** 2570-2580. e2576.
Lee, P.N., Pang, K., Matus, D.Q., and Martindale, M.Q. (2006). A WNT of things to come:
evolution of Wnt signaling and polarity in cnidarians. *Seminars in Cell & Developmental*
Biology 17**,** 157-167.
Liew, Y.J., Aranda, M., and Voolstra, C.R. (2016). Reefgenomics.org - a repository for marine
genomics data. *Database* 2016.
Liew, Y.J., Howells, E.J., Wang, X., Michell, C.T., Burt, J.A., Idaghdour, Y., and Aranda, M.
(2020). Intergenerational epigenetic inheritance in reef-building corals. *Nature Climate*
Change 10**,** 254-259.
Love, M.I., Huber, W., and Anders, S. (2014). Moderated estimation of fold change and
dispersion for RNA-Seq data with DESeq2. *Genome Biology* 15**,** 1-21.
Mackie, G. (1986). From aggregates to integrates: physiological aspects of modularity in
colonial animals. *Philosophical Transactions of the Royal Society of London. B,*
Biological Sciences 313**,** 175-196.
Malik, A., Einbinder, S., Martinez, S., Tchernov, D., Haviv, S., Almuly, R., Zaslansky, P.,
Polishchuk, I., Pokroy, B., and Stolarski, J. (2021). Molecular and skeletal fingerprints of
scleractinian coral biomineralization: from the sea surface to mesophotic depths. *Acta*
Biomaterialia 120**,** 263-276.
Marsh Jr, J.A. (1970). Primary productivity of reef-building calcareous red algae. *Ecology* 51**,**
255-263.
Martinez, S., Kolodny, Y., Shemesh, E., Scucchia, F., Nevo, R., Levin-Zaidman, S., Paltiel, Y.,
Keren, N., Tchernov, D., and Mass, T. (2020). Energy sources of the depth-generalist
mixotrophic coral *Stylophora pistillata*. *Frontiers in Marine Science* 7**,** 988.
Mass, T., Brickner, I., Hendy, E., and Genin, A. (2011). Enduring physiological and reproductive
benefits of enhanced flow for a stony coral. *Limnology and Oceanography* 56**,** 2176-
2188.
Mass, T., Drake, Jeana L., Haramaty, L., Kim, J.D., Zelzion, E., Bhattacharya, D., and
Falkowski, Paul G. (2013). Cloning and characterization of four novel coral acid-rich
proteins that precipitate carbonates in vitro. *Current Biology* 23**,** 1126-1131.

Mass, T., Drake, J.L., Peters, E.C., Jiang, W., and Falkowski, P.G. (2014). Immunolocalization
of skeletal matrix proteins in tissue and mineral of the coral *Stylophora pistillata*.
Proceedings of the National Academy of Sciences 111**,** 12728-12733.
Mass, T., Einbinder, S., Brokovich, E., Shashar, N., Vago, R., Erez, J., and Dubinsky, Z. (2007).
Photoacclimation of *Stylophora pistillata* to light extremes: metabolism and calcification.
Marine Ecology Progress Series 334**,** 93-102.
Mass, T., Genin, A., Shavit, U., Grinstein, M., and Tchernov, D. (2010). Flow enhances
photosynthesis in marine benthic autotrophs by increasing the efflux of oxygen from the
organism to the water. *Proceedings of the National Academy of Sciences* 107**,** 2527-2531.
Mass, T., Putnam, H.M., Drake, J.L., Zelzion, E., Gates, R.D., Bhattacharya, D., and Falkowski,
P.G. (2016). Temporal and spatial expression patterns of biomineralization proteins
during early development in the stony coral *Pocillopora damicornis*. *Proceedings of the*
Royal Society B: Biological Sciences 283**,** 20160322.
Matsuda, S.B., Huffmyer, A.S., Lenz, E.A., Davidson, J.M., Hancock, J.R., Przybylowski, A.,
Innis, T., Gates, R.D., and Barott, K.L. (2020). Coral bleaching susceptibility Is
predictive of subsequent mortality within but not between coral species. *Frontiers in*
Ecology and Evolution 8.
Meesters, E.H., and Bak, R.P. (1995). Age-related deterioration of a physiological function in the
branching coral *Acropora palmata*. *Marine Ecology Progress Series* 121**,** 203-209.
Mummadisetti, M.P., Drake, J.L., and Falkowski, P.G. (2021). The spatial network of skeletal
proteins in a stony coral. *Journal of The Royal Society Interface* 18**,** 20200859.
Nakamura, H., Asari, T., Ohizumi, Y., Kobayashi, J.i., Yamasu, T., and Murai, A. (1993).
Isolation of zooxanthellatoxins, novel vasoconstrictive substances from the zooxanthella
Symbiodinium sp. *Toxicon* 31**,** 371-376.
Palumbi, S.R., Barshis, D.J., Traylor-Knowles, N., and Bay, R.A. (2014). Mechanisms of reef
coral resistance to future climate change. *Science* 344**,** 895-898.
Park, E., Hwang, D.-S., Lee, J.-S., Song, J.-I., Seo, T.-K., and Won, Y.-J. (2012). Estimation of
divergence times in cnidarian evolution based on mitochondrial protein-coding genes and
the fossil record. *Molecular Phylogenetics and Evolution* 62**,** 329-345.
Peled, Y., Drake, J.L., Malik, A., Almuly, R., Lalzar, M., Morgenstern, D., and Mass, T. (2020).
Optimization of skeletal protein preparation for LC–MS/MS sequencing yields additional
coral skeletal proteins in *Stylophora pistillata*. *BMC Materials* 2**,** 8.
Primor, N., and Zlotkin, E. (1975). On the ichthyotoxic and hemolytic action of the skin
secretion of the flatfish *Pardachirus marmoratus* (Soleidae). *Toxicon* 13**,** 227-231.
Puverel, S., Tambutté, E., Pereira-Mouries, L., Zoccola, D., Allemand, D., and Tambutté, S.
(2005). Soluble organic matrix of two Scleractinian corals: Partial and comparative
analysis. *Comparative Biochemistry and Physiology Part B* 141**,** 480-487.
Rachamim, T., Morgenstern, D., Aharonovich, D., Brekhman, V., Lotan, T., and Sher, D. (2015).
The dynamically evolving nematocyst content of an anthozoan, a scyphozoan, and a
hydrozoan. *Molecular Biology and Evolution* 32**,** 740-753.
Ralph, P., Gademann, R., Larkum, A., and Kühl, M. (2002). Spatial heterogeneity in active
chlorophyll fluorescence and PSII activity of coral tissues. *Marine Biology* 141**,** 639-646.
Ramos-Silva, P., Kaandorp, J., Huisman, L., Marie, B., Zanella-Cleon, I., Guichard, N., Miller,
D.J., and Marin, F. (2013). The skeletal proteome of the coral *Acropora millepora*: The
evolution of calcification by cooption and domain shuffling. *Molecular Biology and*
Evolution.

Reinhardt, B., Broun, M., Blitz, I.L., and Bode, H.R. (2004). HyBMP5-8b, a BMP5-8 orthologue, acts during axial patterning and tentacle formation in *Hydra*. *Developmental Biology* 267**,** 43-59.

Rinkevich, B., and Loya, Y. (1983). Oriented translocation of energy in grafted reef corals. *Coral Reefs* 1**,** 243-247.

Ritchie, K.B. (2006). Regulation of microbial populations by coral surface mucus and mucus-associated bacteria. *Marine Ecology Progress Series* 322**,** 1-14.

Ritchie, R. (2008). Universal chlorophyll equations for estimating chlorophylls a, b, c, and d and total chlorophylls in natural assemblages of photosynthetic organisms using acetone, methanol, or ethanol solvents. *Photosynthetica* 46**,** 115-126.

Rocha, R.J., Silva, A.M., Fernandes, M.H.V., Cruz, I.C., Rosa, R., and Calado, R. (2014). Contrasting light spectra constrain the macro and microstructures of scleractinian corals. *PLOS ONE* 9.

Rueffler, C., Hermisson, J., and Wagner, G.P. (2012). Evolution of functional specialization and division of labor. *Proceedings of the National Academy of Sciences* 109**,** E326-E335.

Schiller, C., and Herndl, G.J. (1989). Evidence of enhanced microbial activity in the interstitial space of branched corals: possible implications for coral metabolism. *Coral Reefs* 7**,** 179-184.

Schmidt, C.A., Daly, N.L., and Wilson, D.T. (2019). Coral venom toxins. *Frontiers in Ecology and Evolution* 7**,** 320.

Schmidt-Roach, S., Miller, K.J., Lundgren, P., and Andreakis, N. (2014). With eyes wide open: a revision of species within and closely related to the *Pocillopora damicornis* species complex (Scleractinia; Pocilloporidae) using morphology and genetics. *Zoological Journal of the Linnean Society* 170**,** 1-33.

Sebens, K.P., Witting, J., and Helmuth, B. (1997). Effects of water flow and branch spacing on particle capture by the reef coral *Madracis mirabilis* (Duchassaing and Michelotti). *Journal of Experimental Marine Biology and Ecology* 211**,** 1-28.

Sergushichev, A.A. (2016). An algorithm for fast preranked gene set enrichment analysis using cumulative statistic calculation. *BioRxiv***,** 060012.

Shenkar, N., Fine, M., and Loya, Y. (2005). Size matters: bleaching dynamics of the coral *Oculina patagonica*. *Marine Ecology Progress Series* 294**,** 181-188.

Siegfried, E., and Perrimon, N. (1994). *Drosophila* wingless: a paradigm for the function and mechanism of Wnt signaling. *Bioessays* 16**,** 395-404.

Steinmetz, P.R. (2019). A non-bilaterian perspective on the development and evolution of animal digestive systems. *Cell and Tissue Research* 377**,** 321-339.

Stimson, J. (1978). Mode and timing of reproduction in some common hermatypic corals of Hawaii and Enewetak. *Marine Biology* 48**,** 173-184.

Subramanian, A., Tamayo, P., Mootha, V.K., Mukherjee, S., Ebert, B.L., Gillette, M.A., Paulovich, A., Pomeroy, S.L., Golub, T.R., and Lander, E.S. (2005). Gene set enrichment analysis: a knowledge-based approach for interpreting genome-wide expression profiles. *Proceedings of the National Academy of Sciences* 102**,** 15545-15550.

Sunagawa, S., DeSalvo, M.K., Voolstra, C.R., Reyes-Bermudez, A., and Medina, M. (2009). Identification and gene expression analysis of a taxonomically restricted cysteine-rich protein family in reef-building corals. *PLOS ONE* 4**,** e4865.

Surekha, K.L., Khade, S., Trimbake, D., Patwardhan, R., Nadimpalli, S.K., and Ghaskadbi, S.
(2020). Differential expression of BMP inhibitors gremlin and noggin in *Hydra* suggests
distinct roles during budding and patterning of tentacles. *Developmental Dynamics*.
Takeuchi, T., Yamada, L., Shinzato, C., Sawada, H., and Satoh, N. (2016). Stepwise evolution of
coral biomineralization revealed with genome-wide proteomics and transcriptomics.
PLOS ONE 11**,** e0156424.
RStudio Team. (2019). "RStudio: Integrated Development for R". (Boston, MA: RStudio, Inc.).
Voolstra, C.R., Li, Y., Liew, Y.J., Baumgarten, S., Zoccola, D., Flot, J.-F., Tambutté, S.,
Allemand, D., and Aranda, M. (2017). Comparative analysis of the genomes of
Stylophora pistillata and *Acropora digitifera* provides evidence for extensive differences
between species of corals. *Scientific Reports* 7**,** 17583.
Walden, P.D., and Cowan, N.J. (1993). A novel 205-kilodalton testis-specific serine/threonine
protein kinase associated with microtubules of the spermatid manchette. *Molecular and
Cellular Biology* 13**,** 7625-7635.
Wallace, C.C. (1978). The coral genus *Acropora* (scleractinia: Astrocoaniina: Acroporidae) in
the central and southern Great Barrier Reef province. *Memoires of the Queensland
Museum* 18**,** 273-319.
Wallace, C.C. (1985). Reproduction, recruitment and fragmentation in nine sympatric species of
the coral genus *Acropora*. *Marine Biology* 88**,** 217-233.
Wangpraseurt, D., Polerecky, L., Larkum, A.W., Ralph, P.J., Nielsen, D.A., Pernice, M., and
Kühl, M. (2014). The in situ light microenvironment of corals. *Limnology and
Oceanography* 59**,** 917-926.
Winkler, R., Antonius, A., and Abigail Renegar, D. (2004). The skeleton eroding band disease
on coral reefs of Aqaba, Red Sea. *Marine Ecology* 25**,** 129-144.
Xu, X., Weinstein, M., Li, C., and Deng, C.-X. (1999). Fibroblast growth factor receptors
(FGFRs) and their roles in limb development. *Cell and Tissue Research* 296**,** 33-43.
Yahel, R., Yahel, G., Berman, T., Jaffe, J.S., and Genin, A. (2005). Diel pattern with abrupt
crepuscular changes of zooplankton over a coral reef. *Limnology and Oceanography* 50**,**
930-944.
Yosef, O., Popovits, Y., Malik, A., Ofek-Lalzer, M., Mass, T., and Sher, D. (2020). A tentacle
for every occasion: comparing the hunting tentacles and sweeper tentacles, used for
territorial competition, in the coral *Galaxea fascicularis*. *BMC Genomics* 21**,** 1-16.
Young, M.D., Wakefield, M.J., Smyth, G.K., and Oshlack, A. (2012). goseq: Gene Ontology
testing for RNA-seq datasets. *R Bioconductor* 8**,** 1-25.
Yu, G., Lam, T.T.-Y., Zhu, H., and Guan, Y. (2018). Two methods for mapping and visualizing
associated data on phylogeny using ggtree. *Molecular Biology and Evolution* 35**,** 3041-
3043.
Zaquin, T., Malik, A., Drake, J.L., Putnam, H.M., and Mass, T. (2021). Evolution of protein-
mediated biomineralization in scleractinian corals. *Frontiers in Genetics* 12**,** 52.
Zoccola, D., Ganot, P., Bertucci, A., Caminiti-Segonds, N., Techer, N., Voolstra, C.R., Aranda,
M., Tambutté, E., Allemand, D., and Casey, J.R. (2015). Bicarbonate transporters in
corals point towards a key step in the evolution of cnidarian calcification. *Scientific
Reports* 5.
Zoccola, D., Innocenti, A., Bertucci, A., Tambutté, E., Supuran, C.T., and Tambutté, S. (2016).
Coral carbonic anhydrases: regulation by ocean acidification. *Marine Drugs* 14**,** 109.
1092

a

b

2, 5, 20

1, 8, 23

9   24

3, 7, 22

4, 6, 21

N

c

<u>Top</u> – 9, 24

<u>Middle</u>
5-8, 20, 21

<u>Bottom</u>
1-4, 22, 23

Tip1   Tip2

Junction

Base

Biomineralization

Hemolytic Toxins

Energy Production

Protein synthesis and folding

Fatty acid degradation

WNT signaling