Calprotectin-mediated zinc chelation inhibits *Pseudomonas aeruginosa* protease activity in cystic fibrosis sputum

Danielle M. Vermilyea[a], Alex W. Crocker[a], Alex H. Gifford[b,c], and Deborah A. Hogan[a,#]

[a] Geisel School of Medicine at Dartmouth, Hanover, New Hampshire

[b] Pulmonary and Critical Care Medicine, Dartmouth-Hitchcock Medical Center, Lebanon, New Hampshire

[c] The Dartmouth Institute for Health Policy and Clinical Practice, Lebanon, New Hampshire

Running title: Calprotectin inhibits *P. aeruginosa* proteases

[#] Address correspondence to:

Department of Microbiology and Immunology

Geisel School of Medicine at Dartmouth

Rm 208 Vail Building

Hanover, NH 03755

E-mail: dhogan@dartmouth.edu

Tel: (603) 650-1252

Abstract

Pseudomonas aeruginosa induces pathways indicative of low zinc availability in the cystic fibrosis (CF) lung environment. To learn more about *P. aeruginosa* zinc access in CF, we grew *P. aeruginosa* strain PAO1 directly in expectorated CF sputum. The *P. aeruginosa* Zur transcriptional repressor controls the response to low intracellular zinc, and we used the NanoString methodology to monitor levels of Zur-regulated transcripts including those encoding a zincophore system, a zinc importer, and paralogs of zinc containing proteins that do not require zinc for activity. Zur-controlled transcripts were induced in sputum-grown *P. aeruginosa* compared to control cultures, but not if the sputum was amended with zinc. Amendment of sputum with ferrous iron did not reduce expression of Zur-regulated genes. A reporter fusion to a Zur-regulated promoter had variable activity in *P. aeruginosa* grown in sputum from different donors, and this variation inversely correlated with sputum zinc concentrations. Recombinant human calprotectin (CP), a divalent-metal binding protein released by neutrophils, was sufficient to induce a zinc-starvation response in *P. aeruginosa* grown in laboratory medium or zinc-amended CF sputum indicating that CP is functional in the sputum environment. Zinc metalloproteases comprise a large fraction of secreted zinc-binding *P. aeruginosa* proteins. Here we show that recombinant CP inhibited both LasB-mediated casein degradation and LasA-mediated lysis of *Staphylococcus aureus*, which was reversible with added zinc. These studies reveal the potential for CP-mediated zinc chelation to post-translationally inhibit zinc metalloprotease activity and thereby impact the protease-dependent physiology and/or virulence of *P. aeruginosa* in the CF lung environment.

Importance

      The factors that contribute to worse outcomes in individuals with cystic fibrosis (CF) with chronic *Pseudomonas aeruginosa* infections are not well understood. Therefore, there is a need to understand environmental factors within the CF airway that contribute to *P. aeruginosa* colonization and infection. We demonstrate that growing bacteria in CF sputum induces a zinc-starvation response that inversely correlates with sputum zinc levels. Additionally, both calprotectin and a chemical zinc chelator inhibit the proteolytic activities of LasA and LasB proteases suggesting that extracellular zinc chelators can influence proteolytic activity and thus *P. aeruginosa* virulence and nutrient acquisition *in vivo*.

Introduction

In cystic fibrosis (CF), microbes such as *Pseudomonas aeruginosa* colonize airway mucus where they then compete with host cells and other microbes for nutrients, including metals. Divalent metal ions (e.g., Zn^{2+}, Fe^{2+}, Mn^{2+}, etc.) are essential micronutrients for host and microbe alike, in part, because they act as cofactors in enzymes important for a variety of cellular functions. While the concentration of metals, such as zinc, in CF sputum can vary, the concentration of zinc in expectorated sputum from CF patients is elevated, on average, compared to levels in samples from healthy controls [1-3]. However, studies investigating the transcriptional response of *P. aeruginosa* in CF sputum show that a common gene expression pattern is the increased expression of zinc uptake and transport genes [4-9], which are normally expressed when zinc is limited. The *P. aeruginosa* zinc-starvation response is regulated by the zinc uptake regulator (Zur), which is a transcriptional repressor [10]. When intracellular zinc is high, Zur monomers bind zinc, dimerize, and bind DNA to repress gene expression of zinc uptake pathways. When intracellular zinc becomes low, the dimeric, zinc-bound fraction of Zur decreases, which leads to derepression of genes involved in zinc uptake and the expression of zinc-free paralogs of essential proteins (zinc-sparing response). The *P. aeruginosa* Zur regulon [11, 12] includes the zinc transporter-encoding operon *znuABCD* [10, 13, 14], the zincophore-encoding operon *cntILMO* [15, 16], and zinc-free paralogs of ribosomal proteins (*PA3600* and *PA3601*) [13, 17] and transcription factors (*dksA2*) [18]. These responses not only reduce the requirement for zinc but liberate the zinc that was stored in the zinc-dependent forms of these proteins [19].

The host, on the other hand, utilizes nutritional immunity to sequester metal ions away from pathogens to reduce bacterial growth and control infection [20]. One of the most abundant zinc-binding host proteins in CF is calprotectin (CP), which was previously named "the cystic fibrosis antigen" because of its abundance in the serum, sputum, and bronchoalveolar lavage fluid (BALF) of individuals with CF [2, 21-24]. Neutrophils recruited to sites of inflammation release CP as S100A8/A9 heterodimers [25, 26], which then form tetramers in environments with

sufficient levels of calcium (27, 28). Each heterodimer has two divalent-metal binding sites: one

site has high affinity for zinc and low affinity for manganese while the other site is capable of

binding divalent manganese, iron, zinc, or nickel (29). CP is thought to induce zinc limitation as a

means to control infections caused by *Staphylococcus aureus*, *Acinetobacter baumannii* in

tissues, and *Salmonella enterica* serovar Typhimurium in the gastrointestinal tract (30-32).

However, little is known about the effect of CP-mediated zinc sequestration on *P. aeruginosa*

growth and physiology.

Additionally, CP has been shown to inhibit the activity of metalloproteases such as host

matrix metalloproteinases via zinc chelation (33). *P. aeruginosa* regulates expression of several

metalloenzymes, including zinc metalloproteases, by quorum sensing (QS), which is a

mechanism that regulates gene expression in accordance with cell density through the secretion

of signal molecules. The secretion of zinc metalloproteases LasB (PA3724), LasA (PA1871), AprA

(PA1249), ImpA (PA0572), PepB (PA2939), and Protease IV (PA4175) (**Table 1**) are regulated

by transcriptional regulators LasR and RhlR involved in QS (34, 35). This coordinated expression

may be of particular importance for optimal protease activity given recent findings showing that

LasB, Protease IV, and LasA are activated after being secreted by a QS-induced proteolytic

cascade in which LasB activates Protease IV and then Protease IV, in turn, activates LasA (36,

37). Expression of these zinc metalloproteases is important for *P. aeruginosa* colonization and

virulence because they play key roles in processes such as degrading host proteins (e.g., elastin)

(38), invading host cells (39), evading host immune responses (40-42), and lysing other bacteria

(e.g., *S. aureus*) (43, 44). While incubation of *P. aeruginosa* zinc metalloproteases with chemical

zinc chelators inhibits their activity (45, 46), the effect of physiologically relevant zinc chelators

such as CP on the activity of *P. aeruginosa* zinc metalloproteases remains unclear.

To test these hypotheses, we used a novel method in which *P. aeruginosa* strain PAO1

was grown directly in unamended expectorated CF sputum and matched sputum samples treated

with divalent metals (e.g., Zn^{2+} and Fe^{2+}) and zinc chelators (e.g., TPEN and CP). The effect of

zinc chelators on *P. aeruginosa* zinc metalloprotease activity was further assessed using

protease-specific assays. Overall, our findings support a model in which zinc chelation by CP in

the mucus of the CF lung may impact the ecology of colonizing *P. aeruginosa* by inhibiting the

activity of proteases involved in processes such as nutrient acquisition and interspecies

competition.

Results

***P. aeruginosa* exhibits a Zur-regulated zinc-starvation response when grown in CF sputum**

samples from different donors

Given that recent studies show that *P. aeruginosa* increases expression of Zur-regulated

genes in CF sputum (4-6), we first constructed a *lacZ* fusion to the promoter of *PA3600* on the

chromosome of *P. aeruginosa* strain PAO1 (PAO1 *att*::P$_{PA3600}$-*lacZ*) to act as a tool to explore

factors that influence the activation of the Zur regulon. *PA3600* encodes the Zur-regulated zinc-

independent isoform of the 50s ribosomal protein L36 (11-13, 17). Activation of the *PA3600*

promoter was first confirmed by measuring activity by *P. aeruginosa* grown in culture medium

(LB), medium containing TPEN (*N,N,N',N'*-tetrakis-2-pyridylmethyl-ethylenediamine), or medium

containing both TPEN and zinc (**Fig. 1a**). TPEN is a membrane permeable metal ion chelator with

a high affinity for zinc (47) and was therefore used to induce a zinc-starvation response in *P.*

aeruginosa. *P. aeruginosa* grown for 3 h in LB had little promoter activity (~23 Miller Units [MU]),

while growth in medium containing TPEN resulted in a seven-fold increase in promoter activity

(~150 MU) (**Fig. 1a**). The addition of TPEN and an excess of zinc (1 mM) did not stimulate

promoter activity (**Fig. 1a**). The ability of sputum to activate the *PA3600* promoter was then

determined by growing *P. aeruginosa* in M63 minimal medium containing 0.2% glucose (M63),

culture medium plus TPEN (positive control), or expectorated CF sputum from 10 different donors

(**Fig. 1b**). While *P. aeruginosa* grown for 3 h in M63 exhibited greater promoter activity (~85 MU)

than when grown in LB (~23 MU), growth in CF sputum resulted in a three-fold increase in

promoter activity (~281 MU). Average promoter activation in CF sputum was statistically the same

as promoter activity induced by TPEN (**Fig. 1b**).

To further assess the activity of Zur in CF sputum, we used a multiplex method to assess

expression of *PA3600* and three additional Zur-regulated genes. To do so, we used NanoString

technology, which is a hybridization-based method that is quantitative, not hindered by

contaminating DNA in sputum, and requires only a small amount of RNA. Consequently,

NanoString works well for the analysis of small clinical sample aliquots (e.g., sputum) as

previously demonstrated (48, 49). In this study, NanoString technology allowed for the analysis

of subset of Zur-regulated genes: *PA3600*, *cntO*, *znuA*, and *dksA2*. Analysis showed an induction

of these Zur-regulated genes in *P. aeruginosa* grown in sputum compared to M63 (**Fig. 1c**).

Amending samples with excess zinc (1 mM) was sufficient to reduce the expression of Zur-

regulated genes (**Fig. 1c**). Studies have shown regulatory crosstalk between iron and zinc as iron

starvation was previously shown to increase expression of Zur-regulated genes *cntO*, *cntM*, and

amiA, but not *znuA* (50). However, amending sputum samples with excess ferrous iron (1 mM)

did not reduce expression of Zur-regulated genes (**Fig. 1c**). Together these data support the

model that *P. aeruginosa* has limited access to zinc in sputum and that zinc and iron limitation are

separate signals.

Activation of the Zur-regulated *PA3600* promoter in CF sputum is inversely correlated with

concentration of zinc in sputum samples

While promoter activity of *P. aeruginosa* grown in CF sputum samples was overall higher

than medium controls, there was a range of promoter activity across sputum samples from

different subjects (**Fig. 1b**). We hypothesized that differences in promoter activities between

sputum samples from different CF patients were due to differences in sputum zinc concentrations.

To test this, inductively coupled plasma mass spectrometry (ICP-MS) was performed on

homogenized CF sputum samples to measure total metals (i.e., zinc, iron, and manganese)

concentrations. The ability of these same sputum samples to activate the *PA3600* promoter in

reporter strain PAO1 *att*::P*PA3600*-*lacZ* was tested in parallel. The data showed a significant inverse

correlation between sputum zinc concentration and induction of the *PA3600* promoter across

tested sputum samples (**Fig. 2**). There was no significant correlation between sputum iron or

manganese concentrations and induction of the *PA3600* promoter (**Fig. 2b**; **Fig. S1a**; **Fig. S1b**).

Induction of the *PA3600* promoter was also compared to clinical information, primarily lung

function (FEV1%) at the time of sputum collection, but there was no correlation found between

FEV1% and *PA3600* promoter activity (**Fig. S1c**). Therefore, the derepression of Zur-regulated

genes in *P. aeruginosa* grown in CF sputum inversely correlates with the total zinc concentration

in sputum samples.

Recombinant CP induces a *P. aeruginosa* zinc-starvation response *in vitro* and in

expectorated CF sputum

Studies report elevated levels of zinc in CF sputum (1, 2). Our ICP-MS data show that the

sputum sample in our study that elicited the strongest zinc-starvation response had a zinc

concentration of ~2 µg/g (~2000 µg/L, ~31 µM) (**Fig. 2a**). Given the concomitant high zinc

concentration in our CF sputum samples and the elevated zinc starvation response in *P.*

aeruginosa grown in these CF sputum samples, it is likely that the zinc in our CF sputum samples

is bound by zinc-sequestering proteins. CP is one such host zinc-sequestering protein that is

found in high concentrations in the sputum of CF patients (2, 22). CP has also been shown to

induce expression of Zur-regulated genes in *P. aeruginosa* strain PA14 (51). Therefore, we

hypothesized that CP binds zinc to induce a zinc starvation response in *P. aeruginosa* grown in

CF sputum. To test this, we first expressed and purified recombinant human CP as previously

described (52) and as illustrated in **Fig. S2**. The ability of our recombinant CP to induce a zinc-

starvation response was tested by growing *P. aeruginosa* strain PAO1 *att*::P*PA3600*-*lacZ* in culture

medium (LB), medium containing CP, or medium containing CP and zinc (**Fig. 3a**). CP

concentrations in the lung can reach 1 mg/ml (~40 μM) (29), therefore, 1 mg/ml CP was used for

all CP-based experiments. Growing *P. aeruginosa* in medium containing 1 mg/ml CP resulted in

a four-fold increase in promoter activation (~92 MU) compared to the control (~25 MU) (**Fig. 3a**).

The addition of excess zinc (1 mM) in the presence of CP prevented promoter activation (**Fig.**

3a). These results confirm that our purified recombinant human CP can induce a zinc-starvation

response in *P. aeruginosa* which is quenched with the addition of exogenous zinc.

Despite the reportedly high concentrations of CP in the serum, sputum, and BALF of CF

patients (2, 21-24), *P. aeruginosa* appears to be able to access enough zinc to persist. Various

environmental factors may influence CP zinc binding such as calcium concentrations (53), pH

(54), or the presence of oxidants (55, 56). Additionally, while CP in its tetrameric state is resistant

to proteolytic degradation, CP is susceptible to oxidation which in turn makes it susceptible to

proteolytic degradation by both host and bacterial proteases (55, 56). Because it was unclear if

CP in sputum would remain intact and/or active to bind zinc, we tested the ability of recombinant

human CP to bind zinc and thereby induce a zinc-starvation response in *P. aeruginosa* grown in

CF sputum. *P. aeruginosa* strain PAO1 *att*::P$_{PA3600}$-*lacZ* was grown in unamended CF sputum,

sputum supplemented with 1 mM zinc, and sputum supplemented with both 1 mM zinc and 1

mg/ml (~40 μM) CP (**Fig. 3b**). The addition of zinc lowered *PA3600* promoter activity in sputum

(**Fig. 3b**), supporting our NanoString data (**Fig. 1c**), while addition of CP to zinc-amended sputum

significantly prevented reduction of promoter activity (**Fig. 3b**). These data confirm that

recombinant CP added to CF sputum remains intact to bind zinc, which induces a zinc starvation

response in colonizing *P. aeruginosa*.

While recombinant CP added to zinc-amended sputum increased *P. aeruginosa PA3600*

promoter activity on average compared to zinc-amended sputum controls, the CF sputum

samples tested varied in their responses (**Fig. 3b, Fig. 1b**). The inverse correlation between

sputum zinc concentrations and induction of the *PA3600* promoter suggests that sputum samples

that result in high promoter activity have lower concentrations of zinc than samples that induce

low promoter activity, comparatively (**Fig. 2**). The high promoter activity by *P. aeruginosa* was

readily quenched by the addition of zinc but remained high when CP was also added (**Fig. 3b**;

green, lavender, lilac). Conversely, the low promoter activity by *P. aeruginosa* grown in sputum

samples with presumably high zinc is not affected greatly by the addition of zinc nor CP (**Fig. 3b**;

pink, light pink, gray). Overall, these data show that addition of recombinant CP to zinc-amended

sputum can induce a zinc-starvation response dependent on sputum zinc concentration.

Zinc metalloproteases are enriched amongst *P. aeruginosa*-secreted zinc-binding proteins

Since both TPEN and CP were confirmed to bind zinc and induce a zinc-starvation

response in *P. aeruginosa* in culture medium (**Fig. 1a**, **Fig. 3a**), we wanted to further measure the

effects of TPEN- and CP-mediated zinc sequestration on *P. aeruginosa* growth. Addition of TPEN

or CP to cultures grown in LB decreased the final OD_{600} of *P. aeruginosa* compared to control

conditions (**Fig. 3d**), but neither inhibited earlier growth stages (**Fig. 3c**). These data show that *P.*

aeruginosa grows in the presence of CP under the conditions tested.

While CP does not prevent the growth of *P. aeruginosa*, little is known about how CP-

mediated zinc starvation affects *P. aeruginosa* physiology. Unlike the chemical chelator TPEN,

CP is not membrane permeable and instead exerts its effects on pathogens by binding metals in

the extracellular environment. CF sputum has been reported to contain high concentrations of

both CP (2, 22, 23) and secreted *P. aeruginosa* proteases including zinc metalloproteases (57).

We performed a UniProt Knowledgebase (UniProtKB) analysis of the *P. aeruginosa* strain PAO1

proteome, which identified at least 72 zinc-binding proteins (**Table 2**). Of those 72, 64 were

described by Gene Ontology (GO) molecular function as having catalytic activity (**Table 2**), which

is consistent with the role of zinc as a cofactor. Of those 64 zinc-binding enzymes, 12 were further

described as proteases and 5 of those were secreted zinc metalloproteases LasB, LasA, AprA,

ImpA, and PepB (**Table 2**). We performed a second UniProtKB analysis of the *P. aeruginosa*

strain PAO1 proteome that identified at least 34 secreted proteins, of which 6 were proteases and

included the 5 aforementioned zinc metalloproteases in addition to Protease IV (PA4175).

UniProtKB does not show Protease IV as binding zinc, but Protease IV has been described as a

zinc metalloprotease and its enzymatic activity is reduced in a *P. aeruginosa* mutant lacking the

zinc importer-encoding gene *znuA* (14). These analyses suggest that 83-100% of secreted

proteases, important virulence factors, are zinc metalloproteases. Overall, previously published

studies and curated databases suggest that CP and *P. aeruginosa*-secreted zinc

metalloproteases are abundant in the extracellular milieu of the CF mucus environment.

Zinc chelation inhibits LasB-mediated proteolysis

Given the importance of zinc to the activity of zinc metalloenzymes, we hypothesized that

zinc chelation by TPEN and CP would inhibit the activity of secreted zinc metalloproteases. Our

initial studies suggested that LasB and LasA accounted for the majority of proteolytic activity by

P. aeruginosa strain PAO1 (WT) because filtered supernatants from Δ*lasAB* cultures spotted onto

milk plates cleared the milk plates substantially less than filtered WT supernatants (**Fig. 4a**, inset

i-ii). As a result, this study focuses on the effect of zinc chelation on LasB and LasA activity.

To test the above hypothesis, LasB activity was determined quantitatively using azocasein

as a substrate. The azocasein degradation assay was previously described to measure total

proteolytic activity (14). However, by comparing the ability of *P. aeruginosa* WT, Δ*lasA*, and

Δ*lasAB* supernatants to degrade azocasein, we found that azocasein degradation was LasB-

dependent under the conditions tested (**Fig. 4a**). As a result, we tested the effect of TPEN and

CP on LasB activity using the azocasein degradation assay. *P. aeruginosa* supernatants were

filtered and then left untreated, treated with TPEN or CP, or treated with both TPEN or CP and

zinc. Treatment with TPEN or CP inhibited LasB enzymatic activity while addition of excess zinc

(1 mM) in the presence of TPEN or CP restored LasB activity (**Fig. 4b**, **Fig. 4c**). Furthermore,

treatment of Δ*lasAB* supernatants with TPEN (**Fig. S3a**) or CP (**Fig. S3b**) without or with the

addition of excess zinc did not alter azocasein degradation. Therefore, treatment of *P. aeruginosa*

cell-free supernatants with zinc chelators TPEN and CP inhibits LasB-mediated caseinolytic

activity.

Zinc chelation inhibits LasA-mediated lysis of *S. aureus*

LasA activity was determined by monitoring the decrease in absorbance at 595 nm of a

heat-killed *S. aureus* suspension as previously described (14). Use of *P. aeruginosa* strain PAO1

(WT), Δ*lasA*, and Δ*lasA+lasA* (complemented mutant) supernatants confirmed that LasA is

necessary for the lysis of *S. aureus* and that this assay measures LasA-mediated lysis of *S.*

aureus under the conditions tested (**Fig. 5a-b**). This assay was then used to measure LasA

activity in *P. aeruginosa* cell-free supernatants left untreated, treated with TPEN or CP, or treated

with both TPEN or CP and zinc. Treatment of supernatants with TPEN or CP inhibited LasA

activity while treatment with TPEN or CP in the presence of excess zinc (500 µM and 160 µM,

respectively) restored LasA activity (**Fig. 5c-f**). Furthermore, treatment of Δ*lasA* supernatants with

zinc, TPEN, or CP had no effect on lysis of *S. aureus*, confirming that treatment of supernatants

did not have LasA-independent cytotoxic effects on *S. aureus* (**Fig. S5**). Therefore, treatment of

P. aeruginosa cell-free supernatants with zinc chelators TPEN and CP inhibits LasA-mediated

lysis of *S. aureus*.

Discussion

Here we show that *P. aeruginosa* strain PAO1 grown in aliquots of expectorated CF

sputum exhibits a zinc-starvation response despite relatively high concentrations of zinc in the

sputum samples. Treatment with recombinant host CP was sufficient to induce a zinc-starvation

response in *P. aeruginosa* grown in zinc-amended CF sputum samples from different subjects,

demonstrating that CP retains its function in sputum. Furthermore, treatment of *P. aeruginosa*

supernatants with CP inhibited the activity of secreted, extracellular zinc metalloproteases LasB

and LasA. The data presented in this study support a model in which CP released from recruited

neutrophils sequesters zinc from the environment to induce a zinc-starvation response in *P.*

aeruginosa and sequesters zinc from secreted virulence factors including zinc-dependent

metalloproteases LasA and LasB inhibiting *S. aureus* lysis, degradation of peptides, and/or

nutrient acquisition (**Fig. 6**).

A variety of strategies have been used to learn about the environment that *P. aeruginosa*

encounters in the CF lung including analysis of bacteria grown in buffered media supplemented

with CF sputum compared to bacteria grown in laboratory media (8, 9), and direct analysis of

gene expression by bacteria in expectorated CF sputum (4, 5, 58). While studies have varied in

their techniques, transcriptomic analyses have found that genes induced by low intracellular zinc

are elevated in sputum samples relative to controls (4-9). Our model differs from previous models

as it measures the transcriptional response of *P. aeruginosa* grown directly in expectorated

sputum from a variety of CF patients. Our study also found that *P. aeruginosa* activates its zinc-

starvation response in CF sputum on average but revealed differences across samples from

different CF donors (**Fig. 1b-c, Fig. 3b**). These findings taken together underscore the fact that

P. aeruginosa growth in laboratory media would not recapitulate the effect of low-zinc conditions

in the context of CF. To this end, our CF sputum model is one way to provide a low-zinc

environment and allows for investigation of the response of *P. aeruginosa* across sputum samples

from different donors which vary in levels of host factors like CP. This same approach would also

enable the investigation of different *P. aeruginosa* strains in sputum aliquots from a single donor.

CP concentrations during infections can reach 1 mg/ml or ~40 µM which is often posited

to be higher than or in excess of the bioavailable zinc concentration in most environments (29).

However, zinc concentrations in CF sputum are high relative to sputum from non-CF individuals

and other biological compartments. Smith et al. (1) found that the zinc concentration of 45 CF

sputum samples ranged from 678 µg/L (~10 µM) to 1181 µg/L (~18 µM) compared to 103 µg/L

(~2 µM) to 597 µg/L (~9 µM) in 8 non-CF sputum samples. Li et al. (3) reported that the zinc

concentration of 118 CF sputum samples ranged from ~5 µM to ~145 µM. In this study, the zinc

concentration of 8 CF sputum samples ranged from 1.002 µg/g (~15 µM) to 7.562 µg/g (~116 µM)

(**Fig. 2a**). Therefore, under certain conditions or in some microenvironments, CP may not be in

excess of environmental zinc.

There is mounting evidence that divalent-metal sequestration by CP affects *P. aeruginosa*.

Wakeman et al. (51) demonstrated that CP-mediated genetic responses in *P. aeruginosa* were

reversed upon treatment with zinc *in vitro* and that *P. aeruginosa* and CP colocalized at sites of

inflammation within a CF lung explant. D'Orazio et al. showed that CP-mediated growth inhibition

was enhanced in *P. aeruginosa* strain Δ*znuA*, which is a mutant lacking the gene encoding the

small zinc-binding protein of the ZnuABC zinc importer resulting in reduced intracellular zinc

accumulation (13, 14). Zygiel et al. (59) showed that treatment with CP significantly reduced

intracellular iron and manganese in *P. aeruginosa*, but did not significantly affect intracellular zinc,

though intracellular zinc trended downward (59). Our data show that CP induces a Zur-regulated

zinc-starvation response *in vitro* and in expectorated CF sputum which is repressed upon the

addition of excess zinc (**Fig. 3a-b**). We also observed CP-mediated growth defects *in vitro* (**Fig.**

3c) similar to those reported by Zygiel et al. (59) which were previously attributed to ferrous iron

chelation by CP. Taken together, the data show that *P. aeruginosa* and CP colocalize at sites of

inflammation in the CF lung and that CP is capable of inducing zinc- and/or iron-starvation

responses depending on test conditions.

Additionally, while Filkins et al. (60) showed that *in vitro* co-culture of *P. aeruginosa* and

S. aureus on CF bronchial epithelial cells reduced the viability of *S. aureus*, Wakeman et al. (51)

showed that zinc chelation by CP promotes *P. aeruginosa* and *S. aureus* co-culture in *in vitro*, *in*

vivo, and *ex vivo* models, in part, by downregulating genes encoding anti-staphylococcal factors

such as pyocyanin, hydrogen cyanide, and PQS/HQNO. Interestingly, treatment of *P. aeruginosa*

with CP did not reduce the expression of *lasA* though the functionality of LasA was not tested

(51). In this study, we show that CP-mediated zinc chelation inhibits LasA-mediated lysis of *S.*

aureus by *P. aeruginosa in vitro* (**Fig. 5e-f**). Therefore, while LasA may be expressed and

secreted by *P. aeruginosa* in the presence of CP, CP may post-translationally inhibit LasA activity

via zinc sequestration. Furthermore, colonization of the CF airways is usually described as a

pattern of succession where *S. aureus* is the predominant colonizer early on in younger patients

before being outcompeted by *P. aeruginosa* in older patients (60). However, Fischer et al. (61)

recently showed that *P. aeruginosa* and *S. aureus* chronically co-colonize the CF lung. Wakeman

et al. also showed that *P. aeruginosa*, *S. aureus*, and CP colocalize in CF lung explants (51).

Further studies are required to determine if CP modulates protease-dependent and/or protease-

independent co-colonization of *P. aeruginosa* and *S. aureus* in the CF lung.

Notably, *P. aeruginosa* strains chronically adapted to the CF lung, including *lasR* loss-of-

function (LasR-) mutants, have a reduced capacity to outcompete *S. aureus* (62). LasR is a QS

regulator that positively regulates the expression and secretion of several virulence factors

including zinc metalloproteases LasB, LasA, AprA, ImpA, PepB, and Protease IV (34, 35).

However, LasR- strains commonly arise during chronic CF infection and are associated with

worse lung function (63-68). While LasR- strains are common in CF infections, virulence factors

regulated by LasR such as zinc metalloproteases are still reported to be abundant in CF sputum

(57). Recent work by Mould et al. showed that when LasR+ and LasR- strains were cocultured,

the LasR+ strain increased production of RhlR-controlled virulence factors by the LasR- strain

(69). Interestingly, LasB and LasA are reportedly regulated by both the LasR and RhlR QS

regulators (35). Therefore, further investigation is needed to understand how intra- and

interspecies interactions within populations colonizing the CF airway impact the secretion and

function of virulence factors such as zinc metalloproteases LasB and LasA.

LasB is an abundant protease with broad substrate specificity that is implicated in amino

acid liberation and consumption (70). In addition to nutrient acquisition, LasB also plays a role in

the ability of *P. aeruginosa* to invade host epithelial cells (39) and to evade host immune

responses via processes such as degrading cytokines (40). Interestingly, degradation of pro-

inflammatory cytokines IL-8 and IL-6 by LasB reduces neutrophil recruitment and the overall IL-8

and IL-6 response (40). While LasB-mediated cytokine degradation has been reported to reduce

neutrophil recruitment, LasB can also induce neutrophil extracellular traps (NETs) (71, 72).

Neutrophils recruited to sites of inflammation can release CP through processes such as NET

formation (73) and in this study we show that CP-mediated zinc chelation inhibits the activity of

secreted LasB (**Fig. 4c**). Taken together, there appears to be a complex interplay between LasB,

neutrophils, and CP during the course of infection which may contribute to exacerbations in CF.

Furthermore, recent work suggests that secreted LasB activates Protease IV which then

predominantly processes and activates LasA (36, 37). Therefore, CP-mediated inhibition of

secreted LasB activity may have downstream effects on the processing and activity of other

secreted zinc metalloproteases.

In conclusion, the results of our study show that CP can induce a zinc-starvation response

in *P. aeruginosa* in CF sputum as well as chelate zinc to inhibit the activity of virulence-associated

zinc metalloproteases. Future studies will focus on how competition for zinc in a zinc-limited or

zinc-chelating environment such as CF mucus shapes polymicrobial infections and patient

outcomes, particularly considering the observed variability in zinc concentration and availability

across CF patients.

Materials and Methods

Strains and growth conditions

Bacterial strains and plasmids used in this study are listed in **Table S1**. *P. aeruginosa* and

Escherichia coli strains were maintained on lysogeny broth (LB) (1% tryptone, 0.5% yeast extract,

0.5% NaCl) with 1.5% agar and routinely grown in LB on a roller drum at 37°C LB. *P. aeruginosa*

plasmid strains were maintained by supplementing media with 300 µg/ml carbenicillin or 60 µg/ml

gentamicin. *E. coli* plasmid strains were maintained by supplementing media with 100 µg/ml

carbenicillin. *S. aureus* SH1000 was maintained on trypticase soy with 1.5% agar (TSA) or grown

in trypticase soy broth (TSB) on a roller drum at 37°C. *Saccharomyces cerevisiae* strains for

cloning were maintained on yeast-peptone-dextrose (YPD) medium with 2% agar.

Construction of plasmids

Primers used for plasmid construction are listed in **Table S2**. All plasmids were sequenced at the

Molecular Biology Core at the Geisel School of Medicine at Dartmouth. Plasmid GH121_P*PA3600*-

lacZ (DH3229) was constructed using a *S. cerevisiae* recombination technique as previously

described (74). Plasmid GH121_P*pqsA*-*lacZ* served as the vector backbone for this construct.

GH121_P*PA3600*-*lacZ* was purified from yeast using Zymoprep™ Yeast Plasmid Miniprep II

according to manufacturer's protocol and transformed into electrocompetent *E. coli* strain S17 by

electroporation. The plasmid was introduced into *P. aeruginosa* by conjugation and recombinants

were obtained using sucrose counter-selection and genotype screening by PCR.

Complementation plasmid pMQ70_*lasA* was generated using the NEBuilder HiFi DNA assembly

cloning kit (New England BioLabs). *P. aeruginosa* strain PAO1V Δ*lasA* was complemented *in*

trans by inserting a functional copy of *lasA* amplified from PAO1V genomic DNA into plasmid

pMQ70 under the control of the arabinose-inducible *BAD* promoter generating plasmid

pMQ70_*lasA*. Plasmid pMQ70_*lasA* was transformed into Δ*lasA* by electroporation.

Cystic Fibrosis (CF) sputum collection

Sputum samples were collected in accordance with protocols approved by the Committee for the Protection of Human Subjects at Dartmouth. Expectorated sputum samples used in this study were collected from adult subjects with CF during a routine office visit or upon admission for treatment of a disease exacerbation. Sputum samples were frozen upon collection and stored at −80°C until use.

Beta-galactosidase (β-Gal) assay

P. aeruginosa cells with a promoter fusion to *lacZ* integrated at the *att* locus were grown in 5 mL cultures of LB at 37°C for 16 h. Overnight cultures were diluted 1:50 in 50 ml culture medium (LB or M63) and then grown to an OD_{600} of 0.5. The cells were then centrifuged at 4,500 x g for 10 min, resuspended in culture medium, centrifuged at 10,000 x g for 2 min, and then resuspended in 500 µl culture medium. Ten µl of cell suspension were added per 100 µl culture medium or sputum sample in a 2 ml microcentrifuge tube. Samples were incubated at 37°C with shaking for 3 h. β-Gal activity was measured as described by Miller (75) using 50 µl of sample.

RNA isolation and NanoString analysis

Unamended sputum or sputum amended with 1 mM $ZnSO_4 \cdot 7 H_2O$ or $(NH_4)_2Fe(SO_4)_2 \cdot 6 H_2O$ (100 µL) was added to 2 ml microcentrifuge tubes. *P. aeruginosa* strain PAO1 was grown in 5 mL cultures of LB at 37°C for 16 h. Overnight cultures were diluted 1:50 in 50 ml M63 minimal medium with 0.2% glucose and then grown to an OD_{600} of 0.5. The cells were then centrifuged at 4,500 x g for 10 min, washed with water, centrifuged, and then resuspended in 500 µl water. Ten µl of cell suspension were added per 100 µl M63 minimal medium with 0.2% glucose (control) or sputum sample in a 2 ml microcentrifuge tube. Samples were then incubated at 37°C with shaking for 3 h. TriZol (900 µl) was added to 100 µl sputum containing 10 µl of PAO1 cell suspension. Samples

were stored overnight. RNA was prepared following DirectZol kit instructions and eluted in 50 µl

water.

For NanoString, 5 µl of a 1:10 dilution of RNA was used. Diluted RNA was applied to the codeset

PaV4 and processed as previously reported (49). Counts were normalized to the geometric mean

of spiked-in technical controls. Normalized counts were used for Z-score calculations and

heatmap construction.

Measurement of zinc in sputum samples

Sputum samples for zinc analysis were stored at −80°C until processed. Sputum zinc was

quantified by inductively coupled plasma-mass spectrometry (ICP-MS) following nitric acid

digestion of organic material according to the method of Heck et al. and is expressed as µg zinc

per g of sputum (76). ICP-MS was performed by the Dartmouth Trace Element Analysis (TEA)

Core.

Expression and Purification of recombinant calprotectin (CP)

Plasmid S100A8/A9 was obtained from Futami et al. (52) and recombinant CP was expressed

and purified as previously described with minor modification. Plasmid S100A8/A9 was first

confirmed by Sanger sequencing and then transformed into *E. coli* T7 Express cells. Transformed

T7 Express cells were then grown in LB containing 100 µg/ml carbenicillin at 37°C with shaking

and induced at about an OD_{600} of 0.5 with 0.5 mM β-D-1-thiogalactopyranoside (IPTG) for 3 h.

Cultures were centrifuged at 13,260 x g for 10 min at 4°C. Supernatant was discarded. Cell pellets

were resuspended in 30 ml wash solution (150 mM NaCl), transferred to a 50 ml conical tube,

and then centrifuged at 3,210 x g for 10 min at 4°C. Supernatant was discarded. Pellets were

weighed and then stored at -20°C.

Cell pellets were resuspended in 85 mL lysis buffer (50 mM Tris-HCl pH 7.5, 50 mM NaCl, 5 mM

MgCl$_2$) supplemented with Benzonase-HC to control viscosity of the sample. Cells were then

lysed using the microfluidizer with 3 passages at 18,000 psi. Final volume was about 100 ml. 15%

polyethylenimine (PEI) was added dropwise to a final concentration of 0.7% to precipitate nucleic

acids (about 5 ml). Samples were then centrifuged at 23,280 x g for 10 min at 4°C. Pellet

containing intact cells and precipitated nucleic acids was discarded. NH$_4$SO$_4$ (61.27 g) was added

slowly to clarified supernatant (about 115 ml) while stirring at 4°C until a saturation of 80%. The

sample become gradually turbid. Sample was stirred for an additional 30 min after complete

saturation. Sample was then centrifuged at 23,280 x g for 10 min at 4°C. Supernatant was

discarded and the pellet was dissolved in about 30 ml solubilization buffer (50 mM Tris-HCl pH

7.5, 30 mM dithiothreitol [DTT]) and incubated for 1 h at 37°C. Dissolved pellet was transferred to

dialysis cassettes and dialyzed overnight in 50 mM sodium phosphate pH 6.0 at 4°C using 3.5

kDa cut-off dialysis cassettes to change buffer. Sample was then centrifuged at 23,280 x g for 10

min at 4°C to remove any pellet.

CP was then purified using a HiTrap SP column (stored in 20% ethanol). The column was washed

with 5 column volumes (CV) of H$_2$O at about 5 ml/min. The column was then washed with 5 CV

of 100% SP Sepharose HP buffer B (50 mM sodium phosphate pH 6.0, 1 mM DTT, 1 M NaCl;

filtered/degassed) at about 5 ml/min. The column was equilibrated with 10 CV of SP Sepharose

HP buffer A (50 mM sodium phosphate pH 6.0, 1 mM DTT; filtered/degassed) at about 5 ml/min.

A superloop was assembled with the appropriate volume for sample application. Sample was then

loaded in the column using the superloop at 2.5 ml/min. The column was then washed with 10

CV of SP Sepharose HP buffer A at about 5 ml/min. The column was then washed with a step

gradient of SP Sepharose HP buffer B: 5 CV of 5% SP Sepharose HP buffer B, 10 CV at 30% SP

Sepharose HP buffer B and 5 CV at 100% SP Sepharose HP buffer B at about 5 ml/min.

Fractions were analyzed using SDS-PAGE (15% gel) and the appropriate fractions were then

pooled.

CP was then purified using a HiLoad 26/600 Sephadex S75 and CP buffer (50 mM Tris-HCl pH

7.5, 150 mM NaCl, 1 mM DTT; filtered/degassed). Sample (about 13 ml) was loaded in a 50 ml

superloop. Sample was then run on the HiLoad 26/600 Superdex 75p, a program composed of 2

CV equilibration, injection of 12 ml sample and elution with 1.2 CV at 2.6 ml/min. Flow rate is 2.6

ml/min and collection of 7 ml/tube. Tubes corresponding to three different fractions were pooled

to make fractions F1_I, F2_I, and F3_I. All other tubes containing calprotectin from both HiTrap

runs were concentrated using YM-10 Amicon centrifugal filters and re-loaded in the HiLoad

26/600 superdex 75 as before. Tubes corresponding to three different fractions were pooled to

make fractions F1_II, F2_II, and F3_II. Samples from all six fractions were analyzed using SDS-

PAGE (4-12% gel). Fractions F1_I and F1_II, F2_I and F2_II, and F3_I and F3_II were combined

to make fractions F1, F2, and F3, respectively. Fractions were concentrated with YM-10 Amicon

centrifugal filters. The final concentrations of the fractions were determined using a Bradford

protein assay.

Protease assays

P. aeruginosa culture supernatants were used for protease assays. 5 ml overnight cultures in LB

were centrifuged at 4,500 x g for 10 min. Supernatants were then filter sterilized using a 0.22 µm

syringe filter. For TPEN experiments, undiluted supernatants were used. For CP experiments,

stored aliquots of CP were first diluted to 3 mg/ml in CP buffer without DTT (50 mM Tris-HCl pH

7.5, 150 mM NaCl). Then 1 part 3 mg/ml CP was added to 2 parts supernatant for a final

concentration of 1 mg/ml.

Caseinolytic activity was determined qualitatively by spotting *P. aeruginosa* supernatants onto 1%

milk plates or quantitatively using azocasein as a substrate as previously described (14) with

modification. In brief, *P. aeruginosa* culture supernatants were treated overnight (16 h) with 50

μM TPEN or an equivalent volume of 100% EtOH, 1 mg/ml (~40 μM) CP or an equivalent volume

of CP buffer without DTT, and/or 1 mM $ZnSO_4 \cdot 7\ H_2O$ or an equivalent volume of diH_2O.

Treatment of WT supernatants with 50 μM to 2 mM $ZnSO_4 \cdot 7\ H_2O$ was found not to affect LasB

activity (**Fig. S3c**). The supernatants were then incubated at 37°C overnight (16 h). Supernatants

(25 μl) were mixed with 150 μl 2% azocasein in 10 mM Tris-HCl, 8 mM $CaCl_2$, pH 7.4. Samples

were incubated at 37°C for 15 min. 228 μl of 10% TCA were added to each sample, vortexed,

then incubated at room temperature for 15 min. Samples were then centrifuged for 10 min at

10,000 x g. Cleared supernatants (100 μl) were added to wells of a 96-well flat-bottom polystyrene

plate containing 200 μl 1 M NaOH. Absorbance was read at 440 nm.

Staphylolytic activity was determined by monitoring the decrease in absorbance at 595 nm of a

heat-killed *S. aureus* suspension as previously described (14) with modification. *S. aureus* strain

SH1000 (77) was cultured in TSB overnight (16 h) at 37°C with rolling. Cultures were centrifuged

at 4,500 x g for 10 min, resuspended in 20 mM Tris-HCl, pH 8.8 to a final OD_{600} of 1.0, and then

killed by heating at 100°C for 30 min. Heat-killed *S. aureus* suspensions were cooled to room

temperature before use. *P. aeruginosa* culture supernatants were treated overnight (16 h) with 50

μM TPEN or an equivalent volume of 100% EtOH, 1 mg/ml (~40 μM) CP or an equivalent volume

of CP buffer without DTT, and/or 160-500 μM $ZnSO_4 \cdot 7\ H_2O$ or an equivalent volume of diH_2O.

Because increasing concentrations of zinc were previously reported to inhibit LasA activity (46),

an appropriate concentration of zinc to use in add-back experiments was determined

experimentally. For undiluted WT supernatants, the addition of 500 μM zinc had no effect on LasA

activity, while increasing concentrations of zinc inhibited LasA-mediated lysis of *S. aureus* (**Fig.**

S4a-b). Therefore, we used 500 μM zinc for TPEN-based experiments. For CP-buffer diluted WT

supernatants, the addition of 50 µM zinc had no effect on LasA activity, while increasing

concentrations of zinc inhibited LasA-mediated lysis of *S. aureus* (**Fig. S4c-d**). However, a

tetramer of CP can potentially bind up to four zinc ions. Therefore, to ensure that zinc would be

in excess in CP-based experiments, we used 160 µM zinc which was four times the concentration

of CP but still less than 250 µM zinc which was the concentration tested that started to inhibit

LasA activity independent of CP. *P. aeruginosa* supernatants (20 µl) were added to 180 µl of heat-

killed *S. aureus* in wells of a 96-well flat-bottom polystyrene plate. Staphylolytic activity was

determined by monitoring the change in absorbance at 595 nm every 15 min for 3 h using a plate

reader. The plate was shaken before each read.

Statistical analysis

Statistical analysis was performed using GraphPad Prism 8 and results were expressed as the

mean values plus or minus standard deviations. Unless otherwise noted, one-way analysis of

variance (ANOVA) followed by Tukey's multiple-comparison test was performed to determine

statistical significance of the data. See the figure legends for other specific statistical tests used.

Acknowledgements

We would like to thank Pat Occhipinti for generating the promoter-*lacZ* fusion reporter strain used

in this study, Andreia Verissimo of the Institute for Biomolecular Targeting (bioMT) Molecular

Tools Core (MTC) for her assistance in expressing and purifying recombinant calprotectin used

in this study, and Nick Jacobs and Georgia Doing for their insightful comments and feedback

during manuscript preparation. Research reported in this publication was supported by the Cystic

Fibrosis Foundation (CFF) grants GIFFOR1610 and GIFFOR17Y5 awarded to A.H.G. and CFF

grant HOGAN19G0 awarded to D.A.H. Support for the project was also provided by bioMT

through NIH NIGMS grant P20 GM113132, the Dartmouth Cystic Fibrosis Research Center

(DartCF) through NIH NIDDK grant P30 DK117469, the CFF Research Development Program

(CF RDP) through CFF grant STANTO19R0, and the Dartmouth Trace Element Analysis (TEA)

Core through NIH NIEHS P42 ES007373. The content of this publication is solely the

responsibility of the authors and does not necessarily represent the official views of the funding

sources.

References

1.   Smith DJ, Anderson GJ, Bell SC, Reid DW. 2014. Elevated metal concentrations in the

CF airway correlate with cellular injury and disease severity. J Cyst Fibros 13:289-95.

2.   Gray RD, Duncan A, Noble D, Imrie M, O'Reilly DS, Innes JA, Porteous DJ, Greening AP,

Boyd AC. 2010. Sputum trace metals are biomarkers of inflammatory and suppurative

lung disease. Chest 137:635-41.

3.   Li K, Gifford AH, Hampton TH, O'Toole GA. 2020. Availability of zinc impacts interactions

between *Streptococcus sanguinis* and *Pseudomonas aeruginosa* in coculture. J Bacteriol

202.

4.   Cornforth DM, Dees JL, Ibberson CB, Huse HK, Mathiesen IH, Kirketerp-Moller K, Wolcott

RD, Rumbaugh KP, Bjarnsholt T, Whiteley M. 2018. *Pseudomonas aeruginosa*

transcriptome during human infection. Proc Natl Acad Sci U S A 115:E5125-E5134.

5.   Cornforth DM, Diggle FL, Melvin JA, Bomberger JM, Whiteley M. 2020. Quantitative

framework for model evaluation in microbiology research using *Pseudomonas aeruginosa*

and cystic fibrosis infection as a test case. mBio 11.

6.   Mastropasqua MC, Lamont I, Martin LW, Reid DW, D'Orazio M, Battistoni A. 2018.

Efficient zinc uptake is critical for the ability of *Pseudomonas aeruginosa* to express

virulence traits and colonize the human lung. J Trace Elem Med Biol 48:74-80.

7.   Tan J, Doing G, Lewis KA, Price CE, Chen KM, Cady KC, Perchuk B, Laub MT, Hogan

DA, Greene CS. 2017. Unsupervised extraction of stable expression signatures from

public compendia with an ensemble of neural networks. Cell Syst 5:63-71.e6.

8.   Palmer KL, Mashburn LM, Singh PK, Whiteley M. 2005. Cystic fibrosis sputum supports

growth and cues key aspects of *Pseudomonas aeruginosa* physiology. J Bacteriol

187:5267-77.

9.   Turner KH, Wessel AK, Palmer GC, Murray JL, Whiteley M. 2015. Essential genome of

Pseudomonas aeruginosa in cystic fibrosis sputum. Proc Natl Acad Sci U S A 112:4110-

5.

10.   Ellison ML, Farrow JM, Parrish W, Danell AS, Pesci EC. 2013. The transcriptional

regulator Np20 is the zinc uptake regulator in *Pseudomonas aeruginosa*. PLoS One

8:e75389.

11.   Mikhaylina A, Ksibe AZ, Scanlan DJ, Blindauer CA. 2018. Bacterial zinc uptake regulator

proteins and their regulons. Biochem Soc Trans 46:983-1001.

12.   Novichkov PS, Brettin TS, Novichkova ES, Dehal PS, Arkin AP, Dubchak I, Rodionov DA.

2012. RegPrecise web services interface: Programmatic access to the transcriptional

regulatory interactions in bacteria reconstructed by comparative genomics. Nucleic Acids

Res 40:W604-8.

13.   Pederick VG, Eijkelkamp BA, Begg SL, Ween MP, McAllister LJ, Paton JC, McDevitt CA.

2015. ZnuA and zinc homeostasis in *Pseudomonas aeruginosa*. Sci Rep 5:13139.

14.   D'Orazio M, Mastropasqua MC, Cerasi M, Pacello F, Consalvo A, Chirullo B, Mortensen

B, Skaar EP, Ciavardelli D, Pasquali P, Battistoni A. 2015. The capability of *Pseudomonas*

aeruginosa to recruit zinc under conditions of limited metal availability is affected by

inactivation of the ZnuABC transporter. Metallomics 7:1023-35.

15.   Mastropasqua MC, D'Orazio M, Cerasi M, Pacello F, Gismondi A, Canini A, Canuti L,

Consalvo A, Ciavardelli D, Chirullo B, Pasquali P, Battistoni A. 2017. Growth of

Pseudomonas aeruginosa in zinc poor environments is promoted by a nicotianamine-

related metallophore. Mol Microbiol 106:543-561.

16.   Lhospice S, Gomez NO, Ouerdane L, Brutesco C, Ghssein G, Hajjar C, Liratni A, Wang

S, Richaud P, Bleves S, Ball G, Borezée-Durant E, Lobinski R, Pignol D, Arnoux P,

Voulhoux R. 2017. *Pseudomonas aeruginosa* zinc uptake in chelating environment is

primarily mediated by the metallophore pseudopaline. Sci Rep 7:17132.

17.   Makarova KS, Ponomarev VA, Koonin EV. 2001. Two C or not two C: Recurrent disruption of Zn-ribbons, gene duplication, lineage-specific gene loss, and horizontal gene transfer in evolution of bacterial ribosomal proteins. Genome Biol 2:RESEARCH 0033.

18.   Blaby-Haas CE, Furman R, Rodionov DA, Artsimovitch I, de Crecy-Lagard V. 2011. Role of a Zn-independent DksA in Zn homeostasis and stringent response. Mol Microbiol 79:700-15.

19.   Gabriel SE, Helmann JD. 2009. Contributions of Zur-controlled ribosomal proteins to growth under zinc starvation conditions. J Bacteriol 191:6116-22.

20.   Hood MI, Skaar EP. 2012. Nutritional immunity: Transition metals at the pathogen-host interface. Nat Rev Microbiol 10:525-37.

21.   Barthe C, Figarella C, Carrere J, Guy-Crotte O. 1991. Identification of 'cystic fibrosis protein' as a complex of two calcium-binding proteins present in human cells of myeloid origin. Biochim Biophys Acta 1096:175-7.

22.   Gray RD, MacGregor G, Noble D, Imrie M, Dewar M, Boyd AC, Innes JA, Porteous DJ, Greening AP. 2008. Sputum proteomics in inflammatory and suppurative respiratory diseases. Am J Respir Crit Care Med 178:444-52.

23.   Gray RD, Imrie M, Boyd AC, Porteous D, Innes JA, Greening AP. 2010. Sputum and serum calprotectin are useful biomarkers during CF exacerbation. J Cyst Fibros 9:193-8.

24.   MacGregor G, Gray RD, Hilliard TN, Imrie M, Boyd AC, Alton EW, Bush A, Davies JC, Innes JA, Porteous DJ, Greening AP. 2008. Biomarkers for cystic fibrosis lung disease: application of SELDI-TOF mass spectrometry to BAL fluid. J Cyst Fibros 7:352-8.

25.   Edgeworth J, Gorman M, Bennett R, Freemont P, Hogg N. 1991. Identification of p8,14 as a highly abundant heterodimeric calcium binding protein complex of myeloid cells. J Biol Chem 266:7706-13.

26.   Gebhardt C, Németh J, Angel P, Hess J. 2006. S100A8 and S100A9 in inflammation and cancer. Biochem Pharmacol 72:1622-31.

27.   Strupat K, Rogniaux H, Van Dorsselaer A, Roth J, Vogl T. 2000. Calcium-induced
noncovalently linked tetramers of MRP8 and MRP14 are confirmed by electrospray
ionization-mass analysis. J Am Soc Mass Spectrom 11:780-8.

28.   Korndörfer IP, Brueckner F, Skerra A. 2007. The crystal structure of the human
(S100A8/S100A9)$_2$ heterotetramer, calprotectin, illustrates how conformational changes
of interacting alpha-helices can determine specific association of two EF-hand proteins. J
Mol Biol 370:887-98.

29.   Zygiel EM, Nolan EM. 2018. Transition metal sequestration by the host-defense protein
calprotectin. Annu Rev Biochem 87:621-643.

30.   Liu JZ, Jellbauer S, Poe AJ, Ton V, Pesciaroli M, Kehl-Fie TE, Restrepo NA, Hosking MP,
Edwards RA, Battistoni A, Pasquali P, Lane TE, Chazin WJ, Vogl T, Roth J, Skaar EP,
Raffatellu M. 2012. Zinc sequestration by the neutrophil protein calprotectin enhances
Salmonella growth in the inflamed gut. Cell Host Microbe 11:227-39.

31.   Grim KP, San Francisco B, Radin JN, Brazel EB, Kelliher JL, Parraga Solorzano PK, Kim
PC, McDevitt CA, Kehl-Fie TE. 2017. The Metallophore staphylopine enables
Staphylococcus aureus to compete with the host for zinc and overcome nutritional
immunity. mBio 8.

32.   Hood MI, Mortensen BL, Moore JL, Zhang Y, Kehl-Fie TE, Sugitani N, Chazin WJ, Caprioli
RM, Skaar EP. 2012. Identification of an *Acinetobacter baumannii* zinc acquisition system
that facilitates resistance to calprotectin-mediated zinc sequestration. PLoS Pathog
8:e1003068.

33.   Isaksen B, Fagerhol MK. 2001. Calprotectin inhibits matrix metalloproteinases by
sequestration of zinc. Mol Pathol 54:289-92.

34.   Schuster M, Lostroh CP, Ogi T, Greenberg EP. 2003. Identification, timing, and signal
specificity of *Pseudomonas aeruginosa* quorum-controlled genes: A transcriptome
analysis. J Bacteriol 185:2066-79.

35.   Nouwens AS, Beatson SA, Whitchurch CB, Walsh BJ, Schweizer HP, Mattick JS, Cordwell

SJ. 2003. Proteome analysis of extracellular proteins regulated by the *las* and *rhl* quorum

sensing systems in *Pseudomonas aeruginosa* PAO1. Microbiology (Reading) 149:1311-

1322.

36.   Li XH, Lee JH. 2019. Quorum sensing-dependent post-secretional activation of

extracellular proteases in *Pseudomonas aeruginosa*. J Biol Chem 294:19635-19644.

37.   Oh J, Li XH, Kim SK, Lee JH. 2017. Post-secretional activation of Protease IV by quorum

sensing in *Pseudomonas aeruginosa*. Sci Rep 7:4416.

38.   Bruce MC, Poncz L, Klinger JD, Stern RC, Tomashefski JF, Dearborn DG. 1985.

Biochemical and pathologic evidence for proteolytic destruction of lung connective tissue

in cystic fibrosis. Am Rev Respir Dis 132:529-35.

39.   Cowell BA, Twining SS, Hobden JA, Kwong MSF, Fleiszig SMJ. 2003. Mutation of *lasA*

and *lasB* reduces *Pseudomonas aeruginosa* invasion of epithelial cells. Microbiology

149:2291-2299.

40.   LaFayette SL, Houle D, Beaudoin T, Wojewodka G, Radzioch D, Hoffman LR, Burns JL,

Dandekar AA, Smalley NE, Chandler JR, Zlosnik JE, Speert DP, Bernier J, Matouk E,

Brochiero E, Rousseau S, Nguyen D. 2015. Cystic fibrosis–adapted *Pseudomonas*

aeruginosa quorum sensing *lasR* mutants cause hyperinflammatory responses. Sci Adv

1.

41.   Bardoel BW, Hartsink D, Vughs MM, de Haas CJ, van Strijp JA, van Kessel KP. 2012.

Identification of an immunomodulating metalloprotease of *Pseudomonas aeruginosa*

(IMPa). Cell Microbiol 14:902-13.

42.   Laarman AJ, Bardoel BW, Ruyken M, Fernie J, Milder FJ, van Strijp JA, Rooijakkers SH.

2012. *Pseudomonas aeruginosa* alkaline protease blocks complement activation via the

classical and lectin pathways. J Immunol 188:386-93.

43.   Kessler E, Safrin M, Olson JC, Ohman DE. 1993. Secreted LasA of *Pseudomonas*

670   *aeruginosa* is a staphylolytic protease. J Biol Chem 268:7503-8.

671   44.   Spencer J, Murphy LM, Conners R, Sessions RB, Gamblin SJ. 2010. Crystal structure of

672   the LasA virulence factor from *Pseudomonas aeruginosa*: Substrate specificity and

673   mechanism of M23 metallopeptidases. J Mol Biol 396:908-23.

674   45.   Cahan R, Axelrad I, Safrin M, Ohman DE, Kessler E. 2001. A secreted aminopeptidase of

675   *Pseudomonas aeruginosa*: Identification, primary structure, and relationship to other

676   aminopeptidases. J Biol Chem 276:43645-52.

677   46.   Kessler E, Safrin M, Abrams WR, Rosenbloom J, Ohman DE. 1997. Inhibitors and

678   specificity of *Pseudomonas aeruginosa* LasA. J Biol Chem 272:9884-9.

679   47.   Bertuchi FR, Papai R, Ujevic M, Gaubeur I, Cerchiaro G. 2014. General chelating action

680   of copper, zinc and iron in mammalian cells. Analytical Methods 6:8488-8493.

681   48.   Gifford AH, Willger SD, Dolben EL, Moulton LA, Dorman DB, Bean H, Hill JE, Hampton

682   TH, Ashare A, Hogan DA. 2016. Use of a multiplex transcript method for analysis of

683   *Pseudomonas aeruginosa* gene expression profiles in the cystic fibrosis lung. Infect

684   Immun 84:2995-3006.

685   49.   Grahl N, Dolben EL, Filkins LM, Crocker AW, Willger SD, Morrison HG, Sogin ML, Ashare

686   A, Gifford AH, Jacobs NJ, Schwartzman JD, Hogan DA. 2018. Profiling of bacterial and

687   fungal microbial communities in cystic fibrosis sputum using RNA. mSphere 3.

688   50.   Nelson CE, Huang W, Brewer LK, Nguyen AT, Kane MA, Wilks A, Oglesby-Sherrouse

689   AG. 2019. Proteomic analysis of the *Pseudomonas aeruginosa* iron starvation response

690   reveals PrrF small regulatory RNA-dependent iron regulation of twitching motility, amino

691   acid metabolism, and zinc homeostasis proteins. J Bacteriol 201.

692   51.   Wakeman CA, Moore JL, Noto MJ, Zhang Y, Singleton MD, Prentice BM, Gilston BA,

693   Doster RS, Gaddy JA, Chazin WJ, Caprioli RM, Skaar EP. 2016. The innate immune

protein calprotectin promotes *Pseudomonas aeruginosa* and *Staphylococcus aureus*

interaction. Nat Commun 7:11951.

52. Futami J, Atago Y, Azuma A, Putranto EW, Kinoshita R, Murata H, Sakaguchi M. 2016.

An efficient method for the preparation of preferentially heterodimerized recombinant

S100A8/A9 coexpressed in *Escherichia coli*. Biochem Biophys Rep 6:94-100.

53. Stephan JR, Nolan EM. 2016. Calcium-induced tetramerization and zinc chelation shield

human calprotectin from degradation by host and bacterial extracellular proteases. Chem

Sci 7:1962-1975.

54. Rosen T, Nolan EM. 2020. Metal sequestration and antimicrobial activity of human

calprotectin are pH-dependent. Biochemistry 59:2468-2478.

55. Hoskin TS, Crowther JM, Cheung J, Epton MJ, Sly PD, Elder PA, Dobson RCJ, Kettle AJ,

Dickerhof N. 2019. Oxidative cross-linking of calprotectin occurs in vivo, altering its

structure and susceptibility to proteolysis. Redox Biol 24:101202.

56. Stephan JR, Yu F, Costello RM, Bleier BS, Nolan EM. 2018. Oxidative post-translational

modifications accelerate proteolytic degradation of calprotectin. J Am Chem Soc

140:17444-17455.

57. Jaffar-Bandjee MC, Lazdunski A, Bally M, Carrère J, Chazalette JP, Galabert C. 1995.

Production of elastase, exotoxin A, and alkaline protease in sputa during pulmonary

exacerbation of cystic fibrosis in patients chronically infected by *Pseudomonas*

aeruginosa. J Clin Microbiol 33:924-9.

58. Rossi E, Falcone M, Molin S, Johansen HK. 2018. High-resolution *in situ* transcriptomics

of *Pseudomonas aeruginosa* unveils genotype independent patho-phenotypes in cystic

fibrosis lungs. Nat Commun 9:3459.

59. Zygiel EM, Nelson CE, Brewer LK, Oglesby-Sherrouse AG, Nolan EM. 2019. The human

innate immune protein calprotectin induces iron starvation responses in *Pseudomonas*

aeruginosa. J Biol Chem 294:3549-3562.

60.   Filkins LM, Graber JA, Olson DG, Dolben EL, Lynd LR, Bhuju S, O'Toole GA. 2015.
Coculture of Staphylococcus aureus with *Pseudomonas aeruginosa* drives *S. aureus*
towards fermentative metabolism and reduced viability in a cystic fibrosis model. J
Bacteriol 197:2252-64.

61.   Fischer AJ, Singh SB, LaMarche MM, Maakestad LJ, Kienenberger ZE, Peña TA, Stoltz
DA, Limoli DH. 2020. Sustained coinfections with *Staphylococcus aureus* and
Pseudomonas aeruginosa in cystic fibrosis. Am J Respir Crit Care Med.

62.   Baldan R, Cigana C, Testa F, Bianconi I, De Simone M, Pellin D, Di Serio C, Bragonzi A,
Cirillo DM. 2014. Adaptation of *Pseudomonas aeruginosa* in cystic fibrosis airways
influences virulence of *Staphylococcus aureus in vitro* and murine models of co-infection.
PLoS One 9:e89614.

63.   Marvig RL, Sommer LM, Molin S, Johansen HK. 2015. Convergent evolution and
adaptation of *Pseudomonas aeruginosa* within patients with cystic fibrosis. Nat Genet
47:57-64.

64.   Smith EE, Buckley DG, Wu Z, Saenphimmachak C, Hoffman LR, D'Argenio DA, Miller SI,
Ramsey BW, Speert DP, Moskowitz SM, Burns JL, Kaul R, Olson MV. 2006. Genetic
adaptation by *Pseudomonas aeruginosa* to the airways of cystic fibrosis patients. Proc
Natl Acad Sci U S A 103:8487-92.

65.   Hansen SK, Rau MH, Johansen HK, Ciofu O, Jelsbak L, Yang L, Folkesson A, Jarmer H,
Aanæs K, von Buchwald C, Høiby N, Molin S. 2012. Evolution and diversification of
Pseudomonas aeruginosa in the paranasal sinuses of cystic fibrosis children have
implications for chronic lung infection. ISME J 6:31-45.

66.   Köhler T, Buckling A, van Delden C. 2009. Cooperation and virulence of clinical
Pseudomonas aeruginosa populations. Proc Natl Acad Sci U S A 106:6339-44.

67.   Hoffman LR, Kulasekara HD, Emerson J, Houston LS, Burns JL, Ramsey BW, Miller SI.

2009. *Pseudomonas aeruginosa lasR* mutants are associated with cystic fibrosis lung

disease progression. J Cyst Fibros 8:66-70.

68.   Jiricny N, Molin S, Foster K, Diggle SP, Scanlan PD, Ghoul M, Johansen HK, Santorelli

LA, Popat R, West SA, Griffin AS. 2014. Loss of social behaviours in populations of

Pseudomonas aeruginosa infecting lungs of patients with cystic fibrosis. PLoS One

9:e83124.

69.   Mould DL, Botelho NJ, Hogan DA. 2020. Intraspecies signaling between common variants

of *Pseudomonas aeruginosa* increases production of quorum-sensing-controlled

virulence factors. mBio 11.

70.   Flynn JM, Phan C, Hunter RC. 2017. Genome-wide survey of *Pseudomonas aeruginosa*

PA14 reveals a role for the glyoxylate pathway and extracellular proteases in the utilization

of mucin. Infect Immun 85.

71.   Skopelja S, Hamilton BJ, Jones JD, Yang ML, Mamula M, Ashare A, Gifford AH, Rigby

WF. 2016. The role for neutrophil extracellular traps in cystic fibrosis autoimmunity. JCI

Insight 1:e88912.

72.   Skopelja-Gardner S, Theprungsirikul J, Lewis KA, Hammond JH, Carlson KM, Hazlett HF,

Nymon A, Nguyen D, Berwin BL, Hogan DA, Rigby WFC. 2019. Regulation of

Pseudomonas aeruginosa-mediated neutrophil extracellular traps. Front Immunol

10:1670.

73.   Urban CF, Ermert D, Schmid M, Abu-Abed U, Goosmann C, Nacken W, Brinkmann V,

Jungblut PR, Zychlinsky A. 2009. Neutrophil extracellular traps contain calprotectin, a

cytosolic protein complex involved in host defense against *Candida albicans*. PLoS

Pathog 5:e1000639.

74.   Shanks RM, Caiazza NC, Hinsa SM, Toutain CM, O'Toole GA. 2006. *Saccharomyces*

769   *cerevisiae*-based molecular tool kit for manipulation of genes from gram-negative bacteria.

770   Appl Environ Microbiol 72:5027-36.

771   75.   Miller JH. 1992. A short course in bacterial genetics. Cold Spring Harbor Press.

772   76.   Heck JE, Andrew AS, Onega T, Rigas JR, Jackson BP, Karagas MR, Duell EJ. 2009. Lung

773   cancer in a U.S. population with low to moderate arsenic exposure. Environ Health

774   Perspect 117:1718-23.

775   77.   Horsburgh MJ, Aish JL, White IJ, Shaw L, Lithgow JK, Foster SJ. 2002. SigmaB modulates

776   virulence determinant expression and stress resistance: Characterization of a functional

777   *rsbU* strain derived from *Staphylococcus aureus* 8325-4. J Bacteriol 184:5457-67.

Figure Legends

Fig. 1 *P. aeruginosa* inoculated into expectorated CF sputum from different donors exhibits a zinc-starvation response. (**a**) *P. aeruginosa* strain PAO1 P$_{PA3600}$-*lacZ* was grown in LB (Control), LB with 50 μM TPEN (TPEN), or LB with 50 μM TPEN and 1 mM ZnSO$_4$ • 7 H$_2$O (TPEN+Zn) for 3 h. The data shown represent the mean ± SD from three independent experiments. (**b**) *P. aeruginosa* strain PAO1 P$_{PA3600}$-*lacZ* was grown in M63 (Control), M63 with 50 μM TPEN (TPEN), or expectorated CF sputum (sputum) for 3 h. Each point in the sputum set indicates a separate sample from a different donor. The data were analyzed by Brown-Forsythe and Welch ANOVA with Dunnett's T3 multiple comparisons test. (**c**) *P. aeruginosa* strain PAO1 was inoculated into M63 (M63) or into sputum from two different donors (Sputum 1 and Sputum 2). The sputum was divided and left untreated (Sputum), treated with 1 mM ZnSO$_4$ • 7 H$_2$O (Sputum+Zn), or treated with 1 mM (NH$_4$)$_2$Fe(SO$_4$)$_2$ • 6 H$_2$O (Sputum+Fe). Each condition was analyzed in triplicate. The same lowercase letters indicate samples that are not significantly different and different lowercase letters indicate significant differences ($p<0.05$). *$p<0.05$, **$p<0.01$

Fig. 2 Activation of the *PA3600* promoter in CF sputum by *P. aeruginosa* is inversely correlated with total sputum zinc concentration. (**a**) *P. aeruginosa* strain PAO1 P$_{PA3600}$-*lacZ* was inoculated into 8 different CF sputum samples. Zinc concentration of the same 8 CF sputum samples was determined by ICP-MS. B-Gal activity on the left y-axis (Miller Units; gray bars) was then compared to sputum zinc concentration on the right y-axis (μg/g; red dots), (**b**) Pearson correlation matrix comparing B-Gal activity (Miller units), sputum zinc concentration, sputum iron concentration, and sputum manganese concentration. *$p<0.05$, **$p<0.01$

Fig. 3 Recombinant human CP added to CF sputum and culture medium induces a zinc-starvation response by *P. aeruginosa*. (**a**) *P. aeruginosa* strain PAO1 P$_{PA3600}$-*lacZ* was grown in culture medium (Control), medium with 40 μM CP (CP), or medium with 40 μM CP and 1 mM ZnSO$_4$ • 7

H_2O (CP+Zn) for 3 h. The data shown represent the mean ± SD from three independent

experiments. (**b**) *P. aeruginosa* strain PAO1 P*$_{PA3600}$*-*lacZ* was inoculated into CF sputum from 11

different donors. The sputum was divided and left untreated (Control), treated with 100 μM $ZnSO_4$

• 7 H_2O (Zn), or treated with 40 μM CP and 100 μM $ZnSO_4$ • 7 H_2O (CP+Zn) for 3 h. Different color

dots represent samples from different donors. The same color dots connected by a line are from

the same CF sputum donor. Data were analyzed by RM one-way ANOVA with Tukey's multiple

comparisons test. (**c**) Representative growth curves of *P. aeruginosa* strain PAO1 P*$_{PA3600}$*-*lacZ*

grown in LB, LB containing 50 μM TPEN, or LB containing 40 μM CP. Data shown represent the

mean ± SD of three technical replicates and are representative of three independent experiments.

(**d**) OD_{600} at 16 h of *P. aeruginosa* strain PAO1 P*$_{PA3600}$*-*lacZ* grown in LB, LB containing 50 μM

TPEN, or LB containing 40 μM CP. Data shown represent the mean ± SD of three independent

experiments. The same lowercase letters indicate samples that are not significantly different and

different lowercase letters indicate significant differences ($p<0.05$).

Fig. 4 Zinc chelation inhibits LasB enzymatic activity. (**a**) Filtered supernatants from 16 h cultures

of WT, Δ*lasA*, and Δ*lasAB* were incubated with 2% azocasein for 15 min. Inset are images

showing the ability of (**i**) WT and (**ii**) Δ*lasAB* cell-free supernatants to clear milk plates after 16 h.

(**b**) Filtered supernatants from WT 16 h cultures were left untreated (Control), treated with 50 μM

TPEN (TPEN), or treated with 50 μM TPEN and 1 mM $ZnSO_4$ • 7 H_2O (TPEN+Zn) for an additional

16 h. Supernatants were then incubated with 2% azocasein for 15 min. The data shown represent

the mean ± SD from three independent experiments. (**c**) Filtered supernatants from WT 16 h

cultures were left untreated (Control), treated with 40 μM CP (CP), or treated with 40 μM CP and

1 mM $ZnSO_4$ • 7 H_2O (CP+Zn) for an additional 16 h. Supernatants were then incubated with 2%

azocasein for 15 min. The same lowercase letters indicate samples that are not significantly

different and different lowercase letters indicate significant differences ($p<0.05$). An enzyme unit

(U) is defined as 1 μmol min^{-1}.

Fig. 5 Zinc chelation inhibits LasA enzymatic activity. (**a-b**) Lysis of heat-killed *S. aureus* strain SH1000 by cell-free supernatants from WT, Δ*lasA*, and Δ*lasA*+*lasA* (Δ*lasA* complemented *in trans* under the Control of arabinose-inducible P_{BAD}) 16 h cultures. (**c-d**) Lysis of heat-killed *S. aureus* strain SH1000 by WT and Δ*lasA* cell-free supernatants. WT supernatant was divided and left untreated (WT), treated with 50 μM TPEN (WT+TPEN), or treated with 50 μM TPEN and 500 μM $ZnSO_4 \cdot 7\ H_2O$ (WT+TPEN+Zn). (**e-f**) Lysis of heat-killed *S. aureus* strain SH1000 by WT and Δ*lasA* cell-free supernatants. WT supernatant was divided and left untreated (WT), treated with 40 μM CP (WT+CP), or treated with 40 μM CP and 160 μM $ZnSO_4 \cdot 7\ H_2O$ (WT+CP+Zn). (**a**), (**c**), (**e**) The data represent the mean from three independent experiments. Error bars have been omitted for clarity. (**b**), (**d**), (**f**) Quantification of data in (**a**), (**c**), and (**e**), respectively, using area under the curve (AUC). Data are the mean ± SD from three independent experiments. The same lowercase letters indicate samples that are not significantly different and different lowercase letters indicate significant differences ($p<0.05$).

Fig. 6 Model of the effects of CP-mediated zinc chelation in the CF lung on *P. aeruginosa*. *P. aeruginosa* colonizes the mucus in the airways of CF patients to high densities, which in part requires the uptake and utilization of zinc. At high densities, *P. aeruginosa* secretes a variety of quorum sensing-dependent virulence factors including zinc metalloproteases such as LasB and LasA. LasB is a protease that can degrade host proteins, such as elastin, as well as peptides. These degraded proteins/peptides can then be taken up and utilized as nutrients by *P. aeruginosa*. LasA is a protease that lyses *S. aureus* by cleaving pentaglycine bridges of peptidoglycan. LasA-mediated lysis of *S. aureus* allows *P. aeruginosa* to take up nutrients released from lysed *S. aureus* as well as to outcompete *S. aureus* in the CF lung. During infection, neutrophils are recruited to sites of infection/inflammation. Neutrophils may then release cellular contents such as CP. CP can then bind bioavailable zinc away from *P. aeruginosa* thus reducing

the overall abundance of *P. aeruginosa*, while also inducing a zinc-starvation response by *P.*

aeruginosa. Additionally, CP can bind zinc away from both LasB and LasA thereby inhibiting their

proteolytic activity. Furthermore, LasB and LasA activity have been shown to induce neutrophil

extracellular traps (NETs). Therefore, CP-mediated inhibition of LasB and LasA activity may lead

to less NET formation and, subsequently, less CP release. Black arrows indicate a positive

interaction. Red arrows indicate an inhibitory interaction.

Tables

Table 1 Zinc metalloproteases secreted by *P. aeruginosa*

Gene Number[a]	PDB Entry[b]	Protein Name and Description
PA0572	5KDW	ImpA, immunomodulating metalloprotease of *P. aeruginosa*
PA1249	1KAP	AprA, alkaline metalloprotease or aeruginolysin
PA1871	3IT5	LasA, staphylolytic protease
PA2939	N/A	PepB or PaAP, aminopeptidase
PA3724	1EZM	LasB, elastase or pseudolysin
PA4175	N/A	Protease IV, endoprotease

865 [a] From P. aeruginosa genome website, https://www.pseudomonas.com/.

866 [b] From Protein Data Bank (PDB) website, https://www.rcsb.org/.

868 **Table 2** Characteristics of zinc-binding proteins in *P. aeruginosa* as annotated by UniProtKB[a]

GO Molecular Function[b]	Number of Proteins	Subcellular Localization			
		Secreted	Inner Membrane	Cytoplasm	Not Listed
Zinc-Binding	72	8	5	21	38
Catalytic Activity: Non-peptidase	52	3	-	16	33
Catalytic Activity: Peptidase	12	5	3	1	3
Structural Binding Activity	2	-	-	-	2
Molecular Function Regulator	2	-	-	1	1
ATPase-Coupled Protein Transmembrane Transporter Activity	1	-	1	-	-

869 [a] From protein knowledgebase (UniProtKB) website, https://www.uniprot.org/uniprot/.

870 [b] Gene Ontology (GO)

Fig. 1 *P. aeruginosa* inoculated into expectorated CF sputum from different donors exhibits a zinc-starvation response. (**a**) *P. aeruginosa* strain PAO1 P$_{PA3600}$-*lacZ* was grown in LB (Control), LB with 50 μM TPEN (TPEN), or LB with 50 μM TPEN and 1 mM ZnSO$_4$ • 7 H$_2$O (TPEN+Zn) for 3 h. The data shown represent the mean ± SD from three independent experiments. (**b**) *P. aeruginosa* strain PAO1 P$_{PA3600}$-*lacZ* was grown in M63 (Control), M63 with 50 μM TPEN (TPEN), or expectorated CF sputum (sputum) for 3 h. Each point in the sputum set indicates a separate sample from a different donor. The data were analyzed by Brown-Forsythe and Welch ANOVA with Dunnett's T3 multiple comparisons test. (**c**) *P. aeruginosa* strain PAO1 was inoculated into M63 (M63) or into sputum from two different donors (Sputum 1 and Sputum 2). The sputum was divided and left untreated (Sputum), treated with 1 mM ZnSO$_4$ • 7 H$_2$O (Sputum+Zn), or treated with 1 mM (NH$_4$)$_2$Fe(SO$_4$)$_2$ • 6 H$_2$O (Sputum+Fe). Each condition was analyzed in triplicate. The same lowercase letters indicate samples that are not significantly different and different lowercase letters indicate significant differences (*p*<0.05). *p<0.05, **p<0.01

Fig. 2 Activation of the *PA3600* promoter in CF sputum by *P. aeruginosa* is inversely correlated with total sputum zinc concentration. (**a**) *P. aeruginosa* strain PAO1 P$_{PA3600}$-*lacZ* was inoculated into 8 different CF sputum samples. Zinc concentration of the same 8 CF sputum samples was determined by ICP-MS. B-Gal activity on the left y-axis (Miller Units; gray bars) was then compared to sputum zinc concentration on the right y-axis (µg/g; red dots), (**b**) Pearson correlation matrix comparing B-Gal activity (Miller units), sputum zinc concentration, sputum iron concentration, and sputum manganese concentration. *$p<0.05$, **$p<0.01$

Fig. 3 Recombinant human CP added to CF sputum and culture medium induces a zinc-starvation response by *P. aeruginosa*. (**a**) *P. aeruginosa* strain PAO1 P$_{PA3600}$-*lacZ* was grown in culture medium (Control), medium with 40 μM CP (CP), or medium with 40 μM CP and 1 mM ZnSO$_4$• 7 H$_2$O (CP+Zn) for 3 h. The data shown represent the mean ± SD from three independent experiments. (**b**) *P. aeruginosa* strain PAO1 P$_{PA3600}$-*lacZ* was inoculated into CF sputum from 11 different donors. The sputum was divided and left untreated (Control), treated with 100 μM ZnSO$_4$ • 7 H$_2$O (Zn), or treated with 40 μM CP and 100 μM ZnSO$_4$• 7 H$_2$O (CP+Zn) for 3 h. Different color dots represent samples from different donors. The same color dots connected by a line are from the same CF sputum donor. Data were analyzed by RM one-way ANOVA with Tukey's multiple comparisons test. (**c**) Representative growth curves of *P. aeruginosa* strain PAO1 P$_{PA3600}$-*lacZ* grown in LB, LB containing 50 μM TPEN, or LB containing 40 μM CP. Data shown represent the mean ± SD of three technical replicates and are representative of three independent experiments. (**d**) OD$_{600}$ at 16 h of *P. aeruginosa* strain PAO1 P$_{PA3600}$-*lacZ* grown in LB, LB containing 50 μM TPEN, or LB containing 40 μM CP. Data shown represent the mean ± SD of three independent experiments. The same lowercase letters indicate samples that are not significantly different and different lowercase letters indicate significant differences (*p*<0.05).

Fig. 4 Zinc chelation inhibits LasB enzymatic activity. (**a**) Filtered supernatants from 16 h cultures of WT, Δ*lasA*, and Δ*lasAB* were incubated with 2% azocasein for 15 min. Inset are images showing the ability of (**i**) WT and (**ii**) Δ*lasAB* cell-free supernatants to clear milk plates after 16 h. (**b**) Filtered supernatants from WT 16 h cultures were left untreated (Control), treated with 50 μM TPEN (TPEN), or treated with 50 μM TPEN and 1 mM $ZnSO_4 \cdot 7\ H_2O$ (TPEN+Zn) for an additional 16 h. Supernatants were then incubated with 2% azocasein for 15 min. The data shown represent the mean ± SD from three independent experiments. (**c**) Filtered supernatants from WT 16 h cultures were left untreated (Control), treated with 40 μM CP (CP), or treated with 40 μM CP and 1 mM $ZnSO_4 \cdot 7\ H_2O$ (CP+Zn) for an additional 16 h. Supernatants were then incubated with 2% azocasein for 15 min. The same lowercase letters indicate samples that are not significantly different and different lowercase letters indicate significant differences ($p < 0.05$). An enzyme unit (U) is defined as 1 μmol min^{-1}.

Fig. 5 Zinc chelation inhibits LasA enzymatic activity. (**a-b**) Lysis of heat-killed *S. aureus* strain SH1000 by cell-free supernatants from WT, Δ*lasA*, and Δ*lasA+lasA* (Δ*lasA* complemented *in trans* under the Control of arabinose-inducible P$_{BAD}$) 16 h cultures. (**c-d**) Lysis of heat-killed *S. aureus* strain SH1000 by WT and Δ*lasA* cell-free supernatants. WT supernatant was divided and left untreated (WT), treated with 50 µM TPEN (WT+TPEN), or treated with 50 µM TPEN and 500 µM ZnSO$_4$•7 H$_2$O (WT+TPEN+Zn). (**e-f**) Lysis of heat-killed *S. aureus* strain SH1000 by WT and Δ*lasA* cell-free supernatants. WT supernatant was divided and left untreated (WT), treated with 40 µM CP (WT+CP), or treated with 40 µM CP and 160 µM ZnSO$_4$•7 H$_2$O (WT+CP+Zn). (**a**), (**c**), (**e**) The data represent the mean from three independent experiments. Error bars have been omitted for clarity. (**b**), (**d**), (**f**) Quantification of data in (**a**), (**c**), and (**e**), respectively, using area under the curve (AUC). Data are the mean ± SD from three independent experiments. The same lowercase letters indicate samples that are not significantly different and different lowercase letters indicate significant differences (*p*<0.05).

Fig. 6 Model of the effects of CP-mediated zinc chelation in the CF lung on *P. aeruginosa*. *P. aeruginosa* colonizes the mucus in the airways of CF patients to high densities, which in part requires the uptake and utilization of zinc. At high densities, *P. aeruginosa* secretes a variety of quorum sensing-dependent virulence factors including zinc metalloproteases such as LasB and LasA. LasB is a protease that can degrade host proteins, such as elastin, as well as peptides. These degraded proteins/peptides can then be taken up and utilized as nutrients by *P. aeruginosa*. LasA is a protease that lyses *S. aureus* by cleaving pentaglycine bridges of peptidoglycan. LasA-mediated lysis of *S. aureus* allows *P. aeruginosa* to take up nutrients released from lysed *S. aureus* as well as to outcompete *S. aureus* in the CF lung. During infection, neutrophils are recruited to sites of infection/inflammation. Neutrophils may then release cellular contents such as CP. CP can then bind bioavailable zinc away from *P. aeruginosa* thus reducing the overall abundance of *P. aeruginosa*, while also inducing a zinc-starvation response by *P. aeruginosa*. Additionally, CP can bind zinc away from both LasB and LasA thereby inhibiting their proteolytic activity. Furthermore, LasB and LasA activity have been shown to induce neutrophil extracellular traps (NETs). Therefore, CP-mediated inhibition of LasB and LasA activity may lead to less NET formation and, subsequently, less CP release. Black arrows indicate a positive interaction. Red arrows indicate an inhibitory interaction.