1   Running Title: HGT and pH drive AMR in Salmonella infected chicks

2   Gene flux and acid-imposed selection are the main drivers of

3   antimicrobial resistance in broiler chicks infected with

4   *Salmonella enterica* serovar Heidelberg

Adelumola Oladeinde[1,2,10*], Zaid Abdo[2*], Maximilian O. Press[3¶], Kimberly Cook[4], Nelson

A. Cox[1], Benjamin Zwirzitz[5,6¶], Reed Woyda[2¶], Steven M. Lakin[2¶], Jesse C. Thomas IV[7],

Torey Looft[8], Douglas E. Cosby[1], Arthur Hinton Jr.[1], Jean Guard[1], Eric Line[1], Michael J.

Rothrock[1], Mark E. Berrang[1], Kyler Herrington[9], Gregory Zock[10], Jodie Plumblee

Lawrence[1], Denice Cudnik[1], Sandra House[1], Kimberly Ingram[1], Leah Lariscy[1], Martin

Wagner[5,6], Samuel E. Aggrey[10], Lilong Chai[10], and Casey Ritz[10]

Authors' Affiliation

[1]U.S. National Poultry Research Center, USDA-ARS, Athens, GA, USA. [2]Department of

Microbiology, Immunology and Pathology, Colorado State University, Fort Collins,

Colorado, USA. [3]Phase Genomics Inc, Seattle, WA, 98109, USA. [4]Office of National

Programs, USDA-ARS, Beltsville, Maryland, USA. [5]Institute of Food Safety, Food

Technology and Veterinary Public Health, University of Veterinary Medicine, Vienna,

Austria. [6]Austrian Competence Centre for Feed and Food Quality, Safety, and

Innovation FFoQSI GmbH, Tulln, Austria. [7]Division of STD Prevention, National Center

for HIV/AIDS, Viral Hepatitis, STD and TB Prevention, Center for Disease and Control,

Atlanta, Georgia, USA. [8]National Animal Disease Center, USDA-ARS, Ames, IA, USA,

[9]Medical College of Georgia, Augusta, GA, USA. [10] Poultry Science Dept, University of

Georgia, Athens, GA, USA

Corresponding authors* ade.oladeinde@usda.gov; zaid.abdo@colostate.edu

¶ These authors contributed equally to this work.

Abstract

Antimicrobial resistance (AR) spread is a worldwide health challenge, stemming in large

part, from the ability of microbes to share their genetic material through horizontal gene

transfer (HGT). Overuse and misuse of antibiotics in clinical settings and in food

production have been linked to this increased prevalence and spread of AR.

Consequently, public health and consumer concerns have resulted in a remarkable recent

reduction in antibiotics used for food animal production. This is driven by the assumption

that removing this selective pressure will favor the recovery of antibiotic susceptible taxa

and will limit AR sharing through HGT, allowing the currently available antibiotic arsenal

to be effective for a longer period. In this study we used broiler chicks raised antibiotic-

free and *Salmonella enterica* serovar Heidelberg (SH), as a model food pathogen, to test

this hypothesis. Our results show that neonatal broiler chicks challenged with an antibiotic

susceptible SH strain and raised without antibiotics carried susceptible and multidrug

resistance SH strains 14 days after challenge. SH infection perturbed the microbiota of

broiler chicks and gavaged chicks acquired antibiotic resistant SH at a higher rate. We

determined that the acquisition of a plasmid from commensal *Escherichia coli* population

conferred multidrug resistance phenotype to SH recipients and carriage of this plasmid

increased the fitness of SH under acidic selection pressure. These results suggest that

HGT of AR shaped the evolution of SH and that antibiotic use reduction alone is

insufficient to limit antibiotic resistance transfer from commensal bacteria to *Salmonella*.

Importance

The reported increase in antibiotic resistant bacteria in humans have resulted in a major shift away from antibiotics use in food animal production. This has been driven by the assumption that removing antibiotics will select for antibiotic susceptible bacterial taxa, and this in turn will allow the currently available antibiotic arsenal to be more effective. This shift in practice has highlighted new questions that need to be answered to assess the effectiveness of antibiotic removal in reducing the spread of antibiotic resistance bacteria. This research demonstrates that antibiotic susceptible *Salmonella* Heidelberg strains can acquire multidrug resistance from commensal bacteria present in the gut of neonatal broiler chicks, even in the absence of antibiotic selection. We demonstrate that exposure to acidic pH drove the horizontal transfer of antimicrobial resistance plasmids and suggests that simply removing antibiotics from food-animal production might not be sufficient to limit the spread of antimicrobial resistance.

Introduction

Antimicrobial resistance (AR) spread is a worldwide health challenge[1,2]. A major aspect of this challenge stems from the ability of microbes to share their genetic material through horizontal gene transfer (HGT)[3]. Exposure to antibiotics can select for a microbiota of commensals and pathogens that can withstand antibiotic selective pressure, creating a potential reservoir of resistance genes that are shared with susceptible taxa allowing for their survival and proliferation[3]. Overuse and misuse of antibiotics in a clinical setting[4] and in food production[5] have been shown to result in such selective environments contributing to the increased prevalence and spread of AR. These public health concerns have resulted in a major shift away from antibiotics use in food production. This was driven by the assumption that removing this selective pressure will favor the recovery of susceptible taxa, and that this in turn will allow the currently available antibiotic arsenal to be effective for a longer period[6-9]. This shift in food production practices has highlighted new questions that need to be answered to understand and assess the effectiveness of antibiotic removal in reducing the spread and prevalence of AR in food borne pathogens.

There has been a remarkable reduction in antibiotics sold or distributed in the United States (US) for use in food-producing animals since the publication of the US Food and Drug Administration's guidance for industry #209 (2012) and #213 (2013) and the Veterinary Feed Directive[10]. Chicken production accounted for the largest reduction in medically and nonmedically important antibiotics sold (~47% from 2016 to 2017), also driven by consumer concerns over food safety related to antibiotics use[11]. *Salmonella enterica* serovar Heidelberg (SH) strain is one of the prolific serovars causing salmonellosis in the US and poultry is by far the major vector for SH infections[12-14] . One

of the largest food-borne outbreaks in US history was associated with the consumption

of chicken contaminated with SH[15]. During this outbreak, SH was present in blood

samples of case-patients (15%) and resulted in an overall hospitalization rate of 38% (200

case-patients). Poultry and clinical strains of SH can be resistant to at least 1 antibiotic in

3 or more drug classes. Therefore, we focused on studying antibiotic-free broiler chicken

production, the most consumed meat in the US, with a goal to evaluate the prevalence of

AR and its potential transfer to SH, an important model pathogen.

Our results suggested that *Salmonella* can acquire multidrug resistance from the

bacteria of neonatal broiler chicks raised without antibiotics. We determined that *E. coli*

strains carrying IncI1 plasmids were the main reservoir of transferable AR in broiler chicks

and confirmed that *E. coli* was the donor of IncI1-ST26 (pST26) carrying antibiotic, metal,

and disinfectant genes to SH. The presence of pST26 did not impose a fitness cost to the

SH host and exposure to acidic pH posed a selection pressure on pST26 plasmid

population. Commensals like *E. coli* are a reservoir of mobile genetic elements including

plasmids and bacteriophages. Therefore, we conclude that a simple reduction in antibiotic

use might not be sufficient to prevent transfer of these mobile elements to a major food-

borne pathogen like *Salmonella.*

Results

S. Heidelberg abundance in ceca and litter is influenced by the route of challenge.
To investigate if the route of exposure to SH affected intestinal and litter colonization, we challenged 150 one day old broiler chicks with a nalidixic (nal) resistant strain of SH (SH-2813$_{nal}$R) either by oral gavage (OG), intracloacally (IC) or by the seeder (SB) method (i.e., a few chicks (n =5) were orally challenged and co-mingled with unchallenged chicks (n = 20)) (Fig. 1a). Chicks were grown for 14 days on fresh pine shavings in four separate floor pens (1.8 L ×1.16 W m or 5.9 L×3.8 W ft) including an unchallenged control (CTRL) group (n=25). Chicks were not administered any medication or antibiotics for the duration of the study. The challenge experiments, trial 1 and trial 2, were performed in September 2017 and April 2018, respectively and we have published the rate SH colonized the ceca[16]. SH concentration in the ceca of gavaged chicks (SB and OG) was higher than IC for trial 1 (H = 0.94, df =2, P = 0.62) and OG was higher than IC for trial 2 (H = 10.24, df =2, P = 0.006) (Fig. 1b). In the litter, SH levels were higher for IC pens compared to SB and OG in trial 1 and higher in OG pens compared to IC for trial 2 (Fig. 1c). The percentage of chicks that had SH detected in their ceca was 80% for OG and SB and 100% for IC during trial 1. For trial 2, 100% of the OG and IC were positive and 40% for SB[15]. The ceca and litter of CTRL were negative for SH by culture; however, one litter sample was positive after 24 h enrichment in buffered peptone water. Litter pH and moisture was higher for trial 1 (pH = 6.86 $\pm$ 0.27; moisture = 25$\pm$ 0.03 %) compared to trial 2 (6.54 $\pm$ 0.17; moisture = 21$\pm$ 0.03 %) (Extended data Fig. 1a and b).

S. Heidelberg perturbed the ceca and litter bacteria in a challenge route-specific manner. Next, we questioned if SH gut colonization perturbed changes in the ceca and

litter microbiota. We profiled the microbial community present in the ceca and litter using

16S ribosomal RNA gene amplicon sequencing. The bacterial alpha diversity was higher

for litter samples compared to the ceca for OG for all alpha diversity measures (Observed,

Chao1, Shannon and InvSimpson) examined ($P < 0.01$) (Extended data Fig. 2a). In

contrast, ceca from OG had lowest alpha diversity for all measures used compared to

CTRL ($P < 0.05$). Observed and Chao1 alpha diversity indices were higher for litter

compared to ceca for CTRL and IC ($P = 0.007$). Beta diversity differed between litter and

ceca and the route of SH challenge affected the beta diversity of the litter (F = 43.7, df =

11, $P = 0.001$) more than the ceca (F = 2.21, df = 29 $P = 0.007$). Cecal samples were

more dispersed than litter samples and OG chicks had the highest beta diversity (Fig. 2a).

Firmicutes, *Proteobacteria*, *Actinobacteria* and *Bacteriodetes* were the dominant

phyla present in litter (Fig. 2b). Litter from OG and SB carried higher abundance of

Bacteriodetes, whereas IC and CTRL carried higher abundance of *Firmicutes* than OG

and SB. *Firmicutes* dominated the cecal bacteria of broiler chicks and *Ruminococcaceae*

and *Lachnospiraceae* (class *Clostridia*) were the major families. *Proteobacteria* was

higher in the ceca of CTRL than OG or IC ($X^2= 0.6412$, $P=0.72$) (Fig.2b). Amplicon

sequence   variants   (ASV)   matching   *Subdolinagrum*,   *Lactobacillus*   and

Family_Lachnospiraceae were the most abundant ASVs in OG and IC compared to CTRL

in the ceca (Extended data Fig.2b). In contrast, *E. coli/Shigella* (*E. coli* and *Shigella* are

not distinguishable by 16S rRNA gene sequencing) and *Klebsiella* were the most

abundant ASVs in litter. *E. coli* was higher in IC and CTRL compared to OG and SB and

Klebsiella was higher in CTRL compared to OG, IC and SB (*E. coli*: $X^2= 9.9744$, $P=0.018$;

Klebsiella: $X^2=9.6667$, $P=0.021$). This result suggested that SH perturbed the bacterial

community present in the litter and ceca of broiler chicks especially the

Enterobacteriaceae.

S. Heidelberg developed multidrug resistance after IncI1 plasmid acquisition. We

investigated if SH isolates developed AR after 14 days of challenge. We used the national

antimicrobial resistance monitoring system sensititre ™ panel for Gram-negative bacteria

to conduct antibiotic susceptibility testing (AST) on 250 SH isolates recovered from either

the ceca or litter of challenged chicks. SH-2813$_{nal}$^{R} carries a *gyrA* mutation for nal

resistance, therefore all isolates recovered were resistant to nal. SH also carries a

chromosome-encoded *fosA7* gene that confers resistance to fosfomycin[17]. In trial 1, only

2 % of SH isolates (n = 92) developed resistance to an antibiotic. These two isolates were

resistant to gentamicin, streptomycin and sulfisoxazole. For trial 2, 40 % of the isolates

developed resistance to at least one antibiotic (Fig. 1d) and > 86 % of them were resistant

to 2 or more drug classes like the isolates from trial 1 (Extended data Fig. 3a). However,

trial 2 isolates developed AR to tetracyclines instead of sulfisoxazole. A disc diffusion

assay on a selected number of multidrug resistant (MDR) isolates (n = 4) revealed that

MDR isolates were also resistance to tobramycin, netilmicin and kanamycin (data not

shown). Additionally, OG chicks carried a higher percentage (46 %) of MDR isolates than

IC (24 %) or SB (8%) chicks (X^2 = 29.2, df = 2, *P* < 0.001) (Fig. 1e).

To find AR determinants responsible for the acquired AR phenotype, we performed

whole genome sequencing (WGS) using illumina short read sequencing technology on

sixty-nine SH isolates from trial 2 and two isolates from trial 1. We focused our resources

on trial 2, for which AR acquisition was higher than trial 1. The isolates sequenced were

either susceptible (n = 33) or antibiotic resistant (n = 38). The ancestral SH-2813$_{nal}$^{R}

(isolate used for challenge) harbors an IncX1 plasmid and carriage of this plasmid does

not confer AR in SH. The IncX1 *inc* region was not detected by plasmidFinder[18] in 42 of

the 71 isolates sequenced. Thirty-nine percent of SH isolates acquired Col plasmids, but

none carried a known antibiotic resistance gene (ARG). ARGs known to confer resistance

to aminoglycosides (*aadA1, aac(3)-Via*) and tetracyclines (*tetA*) were detected in all AR

isolates. The AR isolates from trial 1 harbored a sulphonamide resistance gene (*sul1*) in

addition to aminoglycoside genes but no *tet* gene. The ARGs were found on a plasmid

belonging to *inc* group IncI1 (Fig. 1f) and plasmid Multi Locus Sequence Type 26 (pST26).

A change in AR profile was observed in four SB isolates - for these isolates, AST

confirmed them to be MDR but after WGS, the AR determinants were not detected. Upon

AST re-test, these isolates were found to be susceptible. Acquired ARG was absent in

four isolates with resistance to either tetracycline or streptomycin.

HGT and chromosome rearrangement shaped the genome of *S*. Heidelberg. After

introduction of SH to the broiler gut microbiota, we hypothesized that successful

colonization will be dependent on gene flux (gain and loss of genes) and mutations. To

this end, we reconstructed a maximum-likelihood (ML) tree based on the pan genome

and mutations of SH strains recovered after 2 weeks in the gut or litter. The core genome

(genes present in ≥ 95% of the strains) and accessory genome (genes present in < 95%

of the strains) composed of 3729 and 1531 genes, respectively. A total of 91 new

mutations (SNPs and Indels) were on the chromosome of evolved strains and no single

mutation was present in every strain. Rather, mutations were unique to individual strains

or shared between 2-26 isolates (Supplementary data 1). The accessory genome tree

grouped the strains based on AR phenotype, plasmid and ARG carried (Fig. 3a).

Contrastingly, the core genome and SNP-based trees did not provide a clear division of

strains based on these characteristics and clades had low bootstrap support values

(Extended data Fig. 3b and c). Acquisition of IncI (IncI1 and IncI2) plasmid, Col plasmids

and the presence/absence of IncX1 plasmids defined the clades seen on the accessory

genome tree. Susceptible strains carrying only IncX1 plasmids represented Clade I and

MDR strains dominated Clade II. Clade II strains carried either IncI1 only or IncI1 plus a

Col plasmid or both were present with an IncX1 plasmid (Fig. 3a).

A few susceptible and strains with no plasmid (n = 12) were nested within clade II

and the genome of these strains exhibited signs of substantial gene loss (Fig. 3a;

Extended data Fig. 4a). Moreover, clade IIa strains had higher number of contigs, higher

misassemblies and lower genome size than the rest of the strains sequenced for this

study (H = 18.82, df =1, P <0.001; Extended data Fig. 4b-d). To determine if erroneous

assembly affected the clustering of this clade, we performed long read sequencing

(Pacific Biosciences Inc.) on one strain from this clade and used it in a hybrid approach

with Illumina short reads. This resulted in a partially complete chromosome (longest

contig - 4.7 Mbp) but 30 non- circular contigs were not scaffolded. An alignment of the

ordered contigs with the ancestor showed that the strain did not suffer significant gene

loss as suggested through short or long reads only assembly (Supplementary Table 1),

however a 3.2 Mbp chromosomal region was inverted (Fig. 3b). This strain also carried

an untypeable circular episome that carried uncharacterized proteins and two genes

encoding NADH-quinone oxidoreductase (*nuo*) (~22 kbp) (Extended data Fig.4e), and an

IncX1 plasmid with divergent protein sequences from the IncX1 present in the ancestor

(Extended data Fig.4f). Gene inversions can increase the diversity and virulence of

pathogens[19], hence we questioned if chromosomal inversion was present in other

evolved strains. To do this, we aligned protein sequences of two complete circular

chromosome from trial 1 and 2 with the ancestor. This analysis showed that the isolate

from trial 1 harbored a ~2 Mbp chromosomal inversion (Extended data Fig. 5a).

Gene inversion changes the leading or lagging strand sequence to its reverse

complement, thus altering the GC skew of the affected gene or genome from a positive

value to a negative value or negative to a positive[19]. Accordingly, we confirmed that there

was a GC skew inversion in these regions (Extended data Fig. 5b and c). Gene inversions

occur after rejoining of DNA breaks and inverted genes can introduce sequencing bias

and misassembly[20]. Here, deletions affecting the cytochrome c maturation gene cluster

(*ccm*) contributed to the inversion of genes in SH strain *og8-05a* (Fig. 3b; Extended data

Fig. 5b). Further, the *oriC* of *og8-05a* was reoriented when compared to the ancestor. On

the other hand, deletion of a putative YebC/PmpR transcriptional regulator and a deletion

between *cysK* and *cysZ* led to the reorientation of *oriC* in strain *ic9b* (Extended data Fig.

5a and c). The inverted region in these genomes encoded 34 - 60 % of the 168 virulence

genes present in the ancestor including *Salmonella* pathogenicity islands 1 and 2, fimbrial

and adherence genes, type 1 and 2 secretion system proteins and prophages

(Supplementary data 2; Extended data Fig. 6). Taken together, this result suggested that

gene flux and homologous recombination shaped the genome of SH strains after entry

into the broiler chick gut.

pST26 differed in their AR genetic context and genome architecture. To compare

pST26 plasmids, we used hybrid assemblies to achieve circular plasmids for two MDR

SH isolates recovered from trial 1 and trial 2. The pST26 plasmids from trial 1 (p1ST26)

and trial 2 (p2ST26) were determined to be ~ 112 kbp long and ~ 87 % identical (Fig. 1f).

Both carried the atypical IncI1 backbone including regions encoding replication, stability,

leading and conjugative transfer[21]. To determine the IncI1-complex group, we used the

classical *traY* and *excA* protein sequences of plasmid R64 (IncI1) and R621a (IncI1-

gamma). A reconstructed tree confirmed that both plasmids belong to the IncI1 group

(Fig. 4a). A 19.8 kb variable region encoding transposons and AR was found in both

plasmids. For p1ST26, this region encoded *aadA1, aac(3)-Via)*, *sul1* and quaternary

(quats) ammonium compound resistance *(qacEΔ1)* (Fig.1f; Extended data Fig. 7). In

contrast, this region carried *tetA* and *mer* operon in addition to *aadA1* and *aac(3)-Via* in

p2ST26 (Fig. 1f) .

To investigate whether gene flux influenced IncI1 genome architecture, we used

p1ST26 and p2ST26 as reference genomes for the rest of the MDR isolates carrying

IncI1. By aligning raw reads to complete pST26 genomes, we were able generate a

consensus IncI1 plasmid contig for each MDR isolate. A tree built with these plasmids

resulted in two clades represented by p1ST26 and p2ST26 plasmids, respectively (Fig.

4b). Multiple alignment of protein sequences revealed that pST26 plasmids differed in

number of genes carried and gene alleles. This was pronounced for regions encoding

AR, pilus/shufflon assembly proteins (*pil*) and IS66-family of transposases

(Supplementary data 3). A pangenome analysis revealed that the pST26 from this study

carried 70 core genes (genes present in ≥ 95% of the plasmids) and 90 accessory genes

(genes present in < 95% of the plasmids). A tree reconstructed with the core genes and

accessory genes divided the pST26 into 5 major groups (Fig. 4d and c). For both trees,

the p1ST26 formed one clade and the p2ST26 made up the other clades. No SNPs were

found between p2ST26 but p1ST26 differed from p2ST26 with 21 SNPs and 2 insertions

(Supplementary data 3; Supplementary Table 2). The insertions were present in only one

p1ST26 plasmid. One insertion was found on *tniA* (a DDE-type

integrase/transposase/recombinase protein) and the other was between tn3-like

transposon and *yadA*. Inserted DNA showed significant homology to *tniA* present in *E.*

coli plasmid EcPF5 (Genbank: CP054237) and IS5057 present in *E. coli* plasmid

pIOMTU792 (Genbank: C542972), respectively.

***E. coli* was the major IncI1 reservoir in broiler chicks**. The application of proximity-

ligation method (Hi-C) has improved the assembly of metagenomes and made it possible

to detect plasmid-host associations[22]. Therefore, we used Hi-C to determine the bacterial

species carrying IncI1 plasmids in the broiler chicken microbiota. Hi-C was done on two

cecal samples from chicks challenged with SH via the cloaca (hereafter referred to as Hi-

IC-FL1 and Hi-IC-FL2). We chose IC samples since all chicks in this group were

successfully colonized with SH compared to OG or SB. Our goal was to determine if Hi-

C could find the bacterial host of IncI1 plasmids and detect inoculated SH strains.

Salmonella enterica was found in Hi-IC-FL1 and Hi-IC-FL2 using metagenome sequence

classification bioinformatic tools[23,24] (Fig. 5a). However, serovar determination with

cluster contigs (~ 1.4 and 1.2 Mbp total contig size, respectively) was not possible with

Salmonella serovar prediction tool[25] or after alignment of contigs to SH reference

genome. We detected IncI1 plasmid contigs in Hi-IC-FL1 that were linked to *E. coli*

clusters (Fig. 5b and c; Supplementary Table 3). In addition, we found contigs (n = 151,

mean contig size = 17,607 bp, smallest contig = 1,027 bp, largest contig = 282,389 bp)

with protein sequences identical to pST26 in both Hi-IC-FL1 and Hi-IC-FL2

(Supplementary data 4). Other bacterial clusters belonging to the *Firmicutes* showed a

low level of linkage to IncI1 (~100X less in Hi-IC-FL1). These results suggested that *E.*

coli was the main reservoir of IncI1 plasmids.

Next, we investigated if SH challenge affected *E. coli* and IncI1 plasmid abundance.

To answer this, we used qPCR primers targeting the glyceraldehyde-3-phosphate

dehydrogenase (*gapA*) gene of *E. coli*, the *inc*RNAi region of IncI1 and selected genes

carried on p2ST26 (*intI1*, *aadA* and *tetA*) to determine their concentration in the ceca and

litter. Litter carried higher concentration of *E. coli*, IncI1 and p2ST26 genes than ceca on

a per gram basis (W = 1, *P* < 0.001). (Fig. 6a; Extended data Fig.8a). In the ceca, IncI1

was higher in OG compared to IC or CTRL (X^2 = 9.17, df =2, *P* = 0.01), while *aadA* (H =

8.25, df = 2, *P* = 0.016) was lower in OG compared to IC or CTRL (Fig. 6a). CTRL chicks

carried a higher concentration of *E. coli* (X^2 = 2.32, df =2, *P* = 0.31), *tetA* (X^2 = 5.74, df

=2, *P* = 0.05) and *intI1* (X^2 = 5.69, df =2, *P* = 0.05) genes than OG and IC. A principal

component analysis with these genes and *Salmonella* tetrathionate reductase (*ttr*)

revealed that samples within the control and challenge groups clustered with their

respective groups and that the challenge and control groups were separated. (Extended

data Fig. 8b). A correlation analysis between *Salmonella* and *E. coli* for challenged chicks

showed a significant positive correlation (S = 784, ρ = 0.55, *P* =0.007) (Fig. 6b). These

results suggested that the presence of SH affected *E. coli*, IncI1 and ARG concentrations.

Broiler chicks carried *E. coli* strains harboring pST26 IncI1 plasmids. To confirm if

broiler chicks from this study harbored *E. coli* populations that could serve as a reservoir

of pST26 plasmids, we retrospectively screened the cecal contents of the broiler chicks

used for Hi-C on CHROMagar™ supplemented with gentamicin and tetracycline (see

supplementary methods) and selected five colonies for whole genome sequencing. As

expected, we found pST26 in two strains (phylogroup F, MLST 6858), while the other

three strains (phylogroup D, MLST 69) carried an IncI1 that harbored no ARG, and an

unknown MLST (Table 1). Next, we compared the pST26 in SH to that found in *E. coli*

using complete plasmids. The p1ST26 and p2ST6 IncI1 plasmids of SH were determined

to be 86.8% and 99.4 % identical to pST26 present in *E. coli*. The main difference was is

in the region encoding AR and transposases in p1ST26 (Extended data Fig 9).

pST26 carriage did not pose a fitness cost under selection pressure. pST26 carried

accessory genes for AR, metal resistance or disinfectants (i.e., quaternary ammonium

compounds) Furthermore, SH strains carrying pST26 differed by route of exposure e.g.,

higher proportion in oral vs. cloacal. To this end, we questioned if any of these factors

could exert a selective pressure for pST26. We did not administer antibiotics to broiler

chicks throughout study, therefore we did not consider antibiotic selective pressure.

However, we detected metals including nickel, chromium, copper, and zinc in the starter

feed in parts per million and we found contaminants such as arsenic, lead and cadmium

in trace amounts (Supplementary Table 4). Disinfectants including quaternary ammonium

compounds were used for the cleaning of broiler houses and equipment, thus we

considered disinfectant as another source of selective pressure in this study. To

determine their effect on SH fitness, we used phenotype MicroArray™ (PM) 96-well plates

supplemented with selected metal chlorides and disinfectants at varying concentrations

to compare a pST26 carrying strain to a susceptible evolved strain. There was no

significant difference in metabolic activity for SH strain carrying pST26 compared to its

susceptible counterpart for metals and disinfectant but pST26 exhibited higher

metabolism for benzethonium chloride, chromate, tellurite, and zinc, whereas the

susceptible stain showed higher metabolism for dequalinum chloride, chromium, and

cesium (X^2= 1.83, df = 1, P = 0.1762) (Extended data Fig. 10a and b).

Cox and colleagues have suggested that the acidic pH of the upper GI tract was

associated with the lower number of chicks positive for *Salmonella* and *Campylobacter*

after oral gavage compared to cloacal inoculation[26-28]. Based on this premise, we

hypothesized that SH strains carrying pST26 will survive better under exposure to acidic

pH. To answer this question, we first compared the survival of the strains under different

pH levels (3.5 -10) with or without nitrogen sources and amino acids using PM plates. SH

strain carrying pST26 exhibited lower metabolic activity compared to the susceptible

strain at pH 3.5 – 4 (V = 3; P = 0.5), pH 4.5 with and without nitrogen sources (V = 599;

P = 3.426e-06) and at pH 9.5-10 with and without sources of nitrogen (V = 714; P =

6.547e-07) (Fig. 7a). Next, we compared the survival of the susceptible ancestor,

susceptible evolved strains without pST26 and antibiotic resistant evolved strains carrying

pST26 in pine shaving extract (PSE) adjusted to pH – 2.5. We acclimatized the bacterial

population to PSE (pH – 6.5) for 2 h before exposure to acidified PSE (Extended data

Fig. 10c). We screened for pST26 carrying populations using the gen^R marker on pST26.

After 2h of exposure to acidified PSE, the SH population carrying pST26 (resistant plus

susceptible cells) did not suffer a fitness cost, and cells carrying pST26 exhibited higher

fitness than their susceptible counterparts (X^2= 4.35, df = 2, P = 0.113) (Fig. 7b).

Notwithstanding, pST26 SH cells had a lower population size than the ancestor and

evolved susceptible strains (X^2 = 6.37, df =3, P = 0.095) (Fig. 7c). This result suggested

that pST26 carriage does not pose a fitness cost on SH host but benefits the host when

exposed to acidic pH.

Discussion

In this study, neonatal Cobb 500 broiler chicks challenged with a susceptible *S.*

Heidelberg (SH) strain carried susceptible and multidrug resistance SH population 2-

weeks after in their ceca and litter. This resistance phenotype was conferred by the

acquisition of IncI1-pST26 plasmids. pST26 plasmids are present in *S.* Heidelberg, *S.*

Typhimurium, *S.* Anatum, *S.* Derby, *S.* Schwarzengrund, *S.* Saintpaul and *E. coli* strains

isolated from animal and human sources (Extended data Fig. 11). The closest IncI1

plasmids to pST26 from this study are pST26 present in *E. coli* (CP018625) and *S.*

Typhimurium (CP027409). The plasmids carry aminoglycoside resistance genes with or

without tetracycline, sulphonamide, *qac* and *mer* resistance genes, making it a

multidrug/disinfectant/metal resistance plasmid. Additionally, pST26 transfer was higher

in chicks challenged orally compared to chicks inoculated via the cloaca. This route

specific rate of conjugation suggested that the selection for pST26 in our study was

associated with the differential pH in the upper GI versus lower GI tract. This was

corroborated by in vitro experiments, where acidic pH reduced the metabolism of SH cells

carrying pST26 compared to cells with no pST26. Furthermore, carriage of pST26

increased the fitness of pST26-carrying cells compared to the susceptible SH population

after exposure to acidic pine shaving extract. This type of acid-imposed selection for IncI1

alludes to the "negative frequency dependent selection hypothesis" - where the fitness of

the population is higher when a plasmid is present at a low frequency in the population[29].

This could also explain the higher levels of SH, but lower positivity rate found in the

ceca of broiler chicks gavaged compared to cloacally challenged chicks[16] (Fig. 1b). There

are multiple barriers that makes it challenging for SH to colonize the ceca of broiler chicks

gavaged including the acidity of the upper GI tract and the length of the GI tract to be

traveled. When we measured the distance of the cloaca and esophagus to the ceca of a

49-day old Cobb-500 broiler chicken (photo not shown), we estimated that the SH inocula

will have to travel ~20X farther to get to the ceca of gavaged chicks compared to cloacal

inoculated chicks. It is plausible that this longer "residence" time in the upper GI allows

more opportunities for SH to make contact and compete with other members of chicken

microbiota. In this study, such an event gave rise to SH cells that acquired pST26 from

E. coli. It is also noteworthy that AR acquisition occurred at a lower rate during trial 1

compared to trial 2. This could be due to several factors that warrants further study -

including the higher pH and moisture of the litter in trial 1 compared 2, seasonal

differences (trial 1 was done in the fall and trial 2 was in the spring) and the

antibiotic/disinfectant practices in place at the hatchery or breeder farms at time the

experiments were conducted.

The IncI1-complex are widespread in *Enterobacteriaceae* and have been reported to

be the major driver of AR in *Salmonella* and *E. coli*. Our understanding of their mode of

transfer and AR evolution have been limited to studies on their prevalence, mating

experiments and in vivo challenge with mice[21,30,31]. Fischer et al[32] showed that *E. coli*

populations in broiler chickens acquired β-lactam resistance after 4-days old broiler chicks

were gavaged with an *E. coli* strain carrying $bla_{CTX-M-1}$ on an IncI1 plasmid. In a twin pair

experiment, Hagbo et al[33] revealed that IncI1 carrying bla_{CMY-2}, ColE1 and P1

bacteriophage are frequently transferred between microbiota present in pre-term infant

gut. Recently, Nyirabahizi,et al[34] used likelihood-ratios to show that *E. coli* AR predicted

Salmonella resistance to β-lactams in retail meat, and to gentamicin, ceftriaxone, and

amoxicillin-clavulanic acid in cecal samples and suggested that the horizontal transfer of

bla$_{CMY-2}$ played a role. These studies have demonstrated that IncI1 is the most prevalent

plasmid carrying β-lactam in *Enterobacteriaceae* and that they are transferable from

Salmonella to *E. coli* and vice-versa. IncI1 plasmids can harbor transferable gene

cassettes with AR variants for β-lactam (*bla*$_{TEM}$, *bla*$_{CMY}$, *bla*$_{CTX}$), aminoglycoside (*aadA*,

aacA, *aph*), tetracycline (*tet*), sulphonamide (*sul*), fosfomycin (*fosA*), phenicol (*cmlA*, *floR*)

and trimethoprim (*dfrA*) resistance. Furthermore, mobile elements could carry AR genes

for disinfectants and metals including ammonium compounds (*qac, sug*), mercury (*mer*

operon) and arsenic (*ars* operon).

The nutrient-niche hypothesis suggests that SH will only be able to invade a new

niche if it can metabolize a growth-limiting resource or if it can outcompete a resident

species and a vacant niche arises due to a new component in diet or by eliminating the

competitor [35]. Competition for oxygen was shown to be the limiting factor that contributed

to the virulence of *S.* Enteritidis towards *E. coli* in neonatal chickens[36]. Competition within

and between SH and *E. coli* strains played a key role selecting for antibiotic resistant SH

in our study. SH challenge perturbed the bacterial community and ARGs of broiler chicks

and SH population was positively correlated with *E. coli*. Consequently, gene flux and

genome inversion shaped the evolution of SH in broiler chicks. For SH and *E. coli* to co-

evolve and persist in the same new niche, recombination of genes occurred and led to

the emergence of new SH strains. In our study, pST26 emerged after conjugation

between *E. coli* donors from broiler chicks raised on fresh litter and SH. As expected, SH

genomes were grouped into nine core genome MLST clusters with eight clusters

representing new strains (Extended data Fig. 12). These species/strains are expected to

be fitter in the new niche and persist longer.

Materials and Methods

SH inoculum preparation. We have previously described the characteristics of the SH

strain (SH-2813nal[R]) used for this study and how the inoculum was prepared[16]. Briefly,

the strain belongs to the multilocus sequence type 15, carries a 37-kb conjugative

plasmid, and is resistant to erythromycin, tylosin, and fosfomycin. The strain was

recovered from broiler chicken carcass in 2013 and made resistant to 200 ppm of nalidixic

acid (nal[R]) for selective enumeration. The nal[R] phenotype is conferred by a serine to

tyrosine substitution at position 83 of DNA gyrase subunit A protein (GyrA). The SH

inoculum was grown overnight in poultry litter extract, centrifuged, and resuspended in

1X phosphate buffered saline (PBS). The resuspended cells were used as inocula. The

genome of the ancestor to SH-2813nal[R] was sequenced using Illumina and PacBio

sequencing to achieve a complete circular chromosome and plasmid (Genbank). SH-

2813nal[R] carries six mutations (Supplementary data 5) including the mutation affecting

gyrA that are **not present in the ancestor.**

Challenging broiler chicks with SH. One-day-old Cobb 500 broiler chicks were

purchased from a commercial hatchery in Cleveland, GA, USA. Upon purchase, chicks

were placed in plastic crates lined with brown paper and transported to the University of

Georgia, Poultry Research Center (33.90693362620281, -83.37871698746522). One

hundred chicks were either uninoculated, gavaged or cloacally inoculated with 100 μl-

volume of SH inoculum (Fig.1a). We included a seeder-bird (SB) colonization method,

whereby five orally gavaged chicks were mingled with twenty uninoculated chicks. Each

inoculated chick received ~10^6 colony forming unites (CFU) of SH-2813nal^R. Afterwards,

chicks were placed in floor pens at a stocking density of 0.65 m^2/chick on fresh pine

shavings litter. Broiler chicks were given water and feed ad libitum and raised antibiotic-

free on starter diet for 2-weeks (starter feed was synthesized by the University of

Georgia's poultry research center's feed mill). The concentration of metals and priority

pollutants in feed was determined by the University of Georgia's feed and environmental

water laboratory (Athens, GA, USA) (Supplementary Table 4). Husbandry and

management followed commercial broiler chicken industry guidelines. After 2-weeks, ten

chicks from OG, IC and control groups and fifteen chickens from SB (five seeder chicks

and ten uninoculated pen mate) were sacrificed to determine the extent of SH colonization

in ceca. The experiments were done in two trials conducted in September 2017 (Trial 1)

and April 2018 (Trial 2). The study was approved by the University of Georgia Office of

Animal Care and Use under Animal Use Protocol: A2017 04-028-A2.

Cecal and litter bacteriological analyses. Ceca were aseptically removed from the

eviscera of 2-week-old broiler chicks, placed in a stomacher bag and transported on ice

to the US National Poultry Research Center for analysis. Ceca were weighed and

buffered peptone water (BPW) (BD Difco, MD, USA) was added 3X volume to the weight

(v/w) and stomached for 60s. Serial dilutions were made and plated onto Brilliant Green

Sulfa agar (BGS) (BD Difco, MD, USA) containing 200-ppm nal. In addition, a 10 µl

inoculating loop was used to streak cecal slurry (cecal contents in BPW) onto Xylose

Lysine Tergitol-4 agar (BD Difco, Sparks, MD). supplemented with 4 ppm tetracycline for

trial 1 and BGS supplemented 32 ppm ampicillin (amp) or 32 ppm streptomycin for trial

2. All antibiotics were purchased from Sigma-Aldrich (St. Louis, MO), unless otherwise

noted. Plates were incubated along with the cecal slurry for 24 h. All bacterial incubations

were carried out at 37°C, unless otherwise noted. After incubation, colonies were

manually counted and serial dilution plates with 2 – 100 colonies were used for CFU per

gram calculation. If no colonies appeared on serial dilution plates, cecal slurry was

streaked onto a new BGS plus nal plate and incubated overnight. After incubation, plates

were examined for the presence/absence of *Salmonella* colonies.

Broiler chicken litter was collected as grab samples from seven locations (4 corners

of the pen and under the waterer) in each pen after chicks were removed. The litter

samples were pooled and 30g was processed in duplicates from each pen as previously

described[37]. Serial dilutions of the litter slurry were made and plated onto BGS agar

containing 200 ppm nal. Plates were incubated overnight, and colonies were manually

counted and reported per gram litter dry weight. Litter pH and moisture was determined

as described previously[37]. We selected randomly 2 – 6 single colonies from BGS plates

supplemented with nal from each ceca and litter sample and archived them in 30% LB

glycerol at -80 °C. In addition, cecal slurry was saved at a 4:1 ratio in Luria Bertani broth

(BD Difco, MD, USA) containing 30% glycerol at –80°C, whereas litter samples were

stored in vacuum sealed whirl pak bags at -20°C.

Determining SH metabolism under selective pressure. To determine if exposure to

metals, disinfectants or acidic pH poses a selection on SH strains carrying IncI1 plasmids,

we used 96-well Phenotype Microarray (PM) ™ MicroPlates (PM10, PM12B, PM13B,

PM15B and PM16A) (Biolog, Inc., Hayward CA) to compare the metabolic profile of one

gentamicin resistant strain harboring Incl1 and one susceptible strain. PM™ plates uses

cell respiration via NADH production to determine cell metabolic activity. If the phenotype

is positive in a well, the cells respire actively, reducing a tetrazolium dye and forming a

strong color (Biolog, Inc.). Briefly, SH strains were sub-cultured twice in universal growth

agar (BUG+B) for 24 h at 37°C. Cells were removed with a sterile swab and transferred

to 16 ml IF-0 to achieve a final turbidimetric transmission of 42%T. Afterwards, 15 ml of

the 42%T cell suspension was transferred to 75 ml of IF-0 +dye to achieve a final

transmission of 85% T before 600 µl of the cell suspension was transferred to 120 ml IF-

10+ dye. The suspension was mixed and each well in a microplate was inoculated with

100 µl of the suspension. After inoculation, microplates were covered with a sterile plastic

film and monitored automatically for color development every 15 min for 48 h at 37 °C

using and OmniLog reader for 48 h. To identify phenotypes, the kinetic curves of the

gentamicin resistant and susceptible strains were compared using OmniLog PM software.

For each PM plate, the respiratory unit for each well at 22 h was extracted and $\log_{10}$

transformed. All supplies used were purchased from Biolog, Inc.

Exposing SH to acidic pH. To determine the fitness of evolved SH strains that acquired

antibiotic resistance compared to evolved susceptible strains after exposure to acidic pH,

we exposed gentamicin resistant (n = 3) and susceptible (n =3) strains recovered from

the litter of chicks gavaged with SH to acidified filter sterilized pine shaving extract. Pine

shaving extract (PSE) was prepared as described for poultry litter extract[37], using fresh

pine shavings collected from the University of Georgia, Poultry Research Center, Athens,

GA, USA. Three single colonies of each strain including the ancestral SH-2813nal[R] were

selected from overnight cultures grown on sheep blood agar and transferred to a

microcentrifuge tube containing 900 µl of PSE (pH = 6.52) i.e., one tube per strain. After

transfer, tubes were vortexed, covered with a gas permeable paper strip and incubated

at 41°C under microaerophilic conditions (5% O_2, 10% CO_2, and 85% N_2) for 2 h. After

incubation, tubes were vortexed and 100 µl of the suspension was transferred to a

microcentrifuge tube containing 900 µl of PSE (pH of PSE was adjusted to 2.54 using 1M

HCl (Spectrum Chemical Mfg. Corp., CA, USA) and 1M NaOH (Fisher Chemical, NJ,

USA) Afterwards, tubes were vortexed, and incubation continued for another 2 h as was

done for PSE at pH 6.5. To determine SH population in PSE, one replicate tube per strain

was removed and serially diluted in 1X PBS at timepoints 0 and 2 h for pH 6.5 and 0.5, 1

and 2 h for pH 2.5. Serial dilutions were plated onto BGS agar plates supplemented with

or without 16 ppm gentamicin. SH colonies were counted 18 - 24 h after incubation using

a calibrated automated colony counter.

The fitness of each evolved SH population relative to the SH-2813nal[R] population

was determined after 2h at pH 2.54 as described by San Millan et al[38]:

529
$$W_{strain} = \frac{log_e \frac{N_{final,evol}}{N_{initial,evol}}}{log_e \frac{N_{final,anc}}{N_{initial,anc}}}$$

where W_{strain} is the fitness of the evolved susceptible (total susceptible) or gentamicin

resistant populations (total resistant and gen[R] resistant), $N_{initial, evol}$ and $N_{final,evol}$ are the

numbers of cells (in CFU) of the evolved susceptible or gentamicin resistant population

at timepoint 0 and 2 h after exposure to PSE, and $N_{initial,anc}$ and $N_{final,anc}$ are the numbers

of cells of SH-2813nal[R] population at timepoint 0 and 2 h.

Antimicrobial susceptibility testing. We performed antimicrobial susceptibility testing

(AST) on two hundred and fifty SH isolates recovered from the ceca and litter of broiler

chicks following National Antimicrobial Resistance Monitoring System (NARMS) protocol

for Gram-Negative bacteria. Minimum inhibitory concentration for isolates were

determined by broth microdilution using the Sensititre semiautomated antimicrobial

susceptibility system (Thermo Fisher Scientific, Inc., MA, USA) and interpreted according

to clinical and laboratory standards institute guidelines when available; otherwise,

breakpoints established by NARMS were used. AST was also done on *E. coli* isolates (n

= 5) recovered from the ceca of two broiler chicks challenged intracloacally with SH. The

Kirby-Bauer disc diffusion assay for tobramycin, kanamycin, neomycin, netilmicin was

done on four gentamicin resistant SH isolates as previously described[37].

DNA extraction. DNA was extracted and purified from bacterial colonies using FastDNA

spin Kit (MP Biomedicals, LLC, CA, USA), whereas 250 mg of cecal and litter were

extracted with Qiagen DNeasy Power-Soil DNA kit (Hilden, Germany). The modifications

we made to the manufacturers protocol for DNA extraction have been reported [37,39].

Whole genome sequencing and processing. Illumina short read sequencing was

performed on DNA extracted from SH isolates recovered from ceca and litter. In addition,

five *E. coli* isolates recovered from two cecal samples were sequenced. Libraries were

prepared using either Nextera™ XT or Nextera™ DNA Flex library preparation kits

(Illumina, Inc., San Diego, CA) following the manufacturers protocol. Libraries were

sequenced on the Illumina MiSeq platform with 150 or 250-bp paired end reads.

Additionally, five SH and two *E. coli* isolates were selected for long read sequencing using

Sequel II System (PacBio Biosciences Inc.) or MinION (Oxford Nanopore technology).

Preparation and sequencing of long read libraries were done by next generation

sequencing core centers of University of Georgia and Colorado State University (see

supplementary methods for read quality control and demultiplexing).

Genome assembly, resistome characterization, and quality assessment of long reads

was done using Reads2Resistome pipeline v.1.1.1[40]. Reads2Resistome performed

hybrid assemblies using either Illumina reads and PacBio or Illumina and MinION reads,

using both Unicycler[41] and SPAdes[42]. In addition, hierarchical genome assembly process

(HGAP) assembly was performed using PacBio's SMRT Link v9.0 analysis software suite

using default settings. Assembly quality was assessed by QUAST v5.0.2[43], using default

settings) and genome annotation was done using Prokka[44], Rapid Annotation using

Subsystem Technology (RAST)[45] and BlastKOALA[46]. We confirmed that all SH isolates

were *Salmonella enterica* serovar Heidelberg using *Salmonella* In Silico Typing Resource

(SISTR)[25], and used ClermonTyping[47] and Multi-Locus Sequence Typing (MLST)[48] for *E.*

coli strains. For resistome characterization reads2resistome used ARG-ANNOT[49], the

Comprehensive Antibiotic Resistance Database[50], MEGARes[51], AMRFinderPlus[52],

PlasmidFinder[18], ResFinder[53] and VirulenceFinder database[54]. For plasmid typing and

IncI1 clonal complex determination, we used plasmid MLST[18]. PHAST[55] was used to

identify prophages present in chromosomal contigs and predicted prophage DNA

sequence was annotated with RAST and BlastKOALA. ProgressiveMAUVE v. 1.1.1[56] and

MAFFT v. 1.4.0[57] implemented in Geneious Prime® v 2020.0.1 were used for aligning

and comparing sequences. In addition, a pan genome analysis of annotated assemblies

was conducted with Roary[58]. Phylogenetic trees based on core genome and accessory

genome were reconstructed using the maximum likelihood (ML) method implemented in

RAxML-NG v 2.0.0[59]. When computationally possible, the best model of sequence

evolution predicted by jModelTest[60] was used for tree reconstruction, otherwise the GTR

+ GAMMMA model was implemented. Lastly, we used the bacterial genomes sequenced

in the study to create a BLAST database that can be searched for sequences of interest

in Geneious Prime®.

Illumina short reads were assembled de novo into contigs using Unicycler v.0.4.7 and

characterized with bioinformatic tools described for long reads. Illumina reads were used

to determine single nucleotide polymorphisms (SNPs) and indels present in SH isolates.

Alignment of raw FASTQ reads to the genome of SH-ancestor was done using Burrows-

Wheeler Aligner (BWA)[61] and SNPs/indels were called using Genome Analysis Toolkit[62]

as described previously[39]. Variant call format (VCF) files of identified SNPs/indels and the

Linux/Unix   shell   script   used   have   been   deposited   in   Dryad   Digital

Repository: https://doi.org/10.5061/dryad.4tmpg4f8d.   SNP   based   ML   tree   was

reconstructed by converting merged VCF files to PHYLogeny Inference Package format

using PGDSpider[63]. Afterwards, isolates with duplicated SNPs/indels were removed, and

a ML tree was drawn using the Jukes-Cantor model of nucleotide substitution and

GAMMA model of rate heterogeneity. For other bioinformatic analyses performed

including pST26-IncI1 consensus determination through FASTQ alignment, see

supplementary methods. All raw FASTQ reads for sequenced bacterial genomes are

publicly available under NCBI accession numbers:

Hi-C and cecal metagenome library preparation and analysis A shotgun and Hi-C

DNA library of two cecal samples was created using Nextera™ XT library preparation kit

and Phase Genomics (Seattle, WA) ProxiMeta Hi-C Microbiome Kit following the

manufacturer's instructions. Library sequencing was performed by Novogene corporation

(Sacramento, CA, USA) on the Illumina HiSeq platform using 150bp paired end reads.

Two libraries were sequenced per HiSeq flowcell lane resulting in a total of ~125 million

shotgun reads, and ~206 million Hi-C reads per sample. Metagenomic FASTQ files were

uploaded to the Phase Genomics cloud-based bioinformatics portal for subsequent

analysis. Shotgun reads were filtered and trimmed for quality using bbduk[64] and

normalized using bbtools[64] and then assembled with metaSPAdes[65] using default

options. Hi-C reads were then aligned to the assembly following the Hi-C kit

manufacturer's recommendations (https://phasegenomics.github.io/2019/09/19/hic-

alignment-and-qc.html). Briefly, reads were aligned using BWA-MEM with the -5SP and

-t 8 options specified, and all other options default. SAMBLASTER[66] was used to flag

PCR duplicates. Alignments were then filtered with samtools[67] using the -F 2304 filtering

flag to remove non-primary and secondary alignments. Lastly, Hi-C read alignments to

the assembly were filtered using Matlock (https://github.com/phasegenomics/matlock)

with default options (removing all alignments with MAPQ<20, edit distance greater than

5, and read duplicates). Metagenome deconvolution was performed with ProxiMeta[68,69],

resulting in the creation of putative genome and genome fragment clusters.

Clusters were assessed for quality using CheckM[70] and assigned preliminary

taxonomic classifications with Mash[71] and Kraken2 (v2.0.8-beta)[23]. To search the

metagenome for ARG and plasmids of interest, we used Geneious Prime® to create a

BLAST metagenome database. Metagenome contigs with significant homology (E value

< 0.05 and at least 1000 bp of aligned sequence) were considered contigs of the

respective plasmid or ARG. Hi-C data were then used to link identified sequences to host

genomes and genome fragments within ProxiMeta[22]. Phylogenetic visualizations of Hi-C

linkages used the placement of clusters in a large prokaryotic phylogeny as estimated by

CheckM[70]. Shotgun and Hi-C reads are publicly available under NCBI accession

numbers:

16S rRNA gene sequence processing and analysis. Cecal and litter DNA were used

for bacterial community analysis through the sequencing of the V4 hypervariable region

of the 16S rRNA gene of bacterial genomes. Sequencing was done using the paired-end

(250 × 2) method on the Illumina MiSeq platform. Raw 16S rRNA gene sequences were

processed with dada2 (version 1.14) in the R environment[72]. Briefly, the reads were

trimmed where the quality profile dropped below Q30 (250bp for forward and 240bp for

reverse reads) and all reads with more than 2 expected errors were removed entirely.

The remaining reads were used to learn the error rates and the samples were pooled

prior to sample inference. After merging of forward and reverse reads with the default

parameters, chimeric sequences were removed with the removeBimeraDenovo

command in the consensus method. High quality reads were then classified against the

SILVA database version 132[73]. To avoid bias introduced by spurious amplicon sequence

variants (ASVs) or from samples with low sequencing depth, any ASVs with less than 5

reads and samples with less than 5000 reads were removed from the dataset before

further analysis. Statistical analysis of microbial communities was performed in the R

environment using the packages "phyloseq", "Ampvis2", "vegan", and "MaAsLin2". Alpha

diversity indices were calculated with a dataset rarefied to the minimum sample size

(8740 sequences). After assessing normal distribution by qqplots, histograms, and the

Shapiro-Wilk normality test, the not normal distributed groups were compared using the

Wilcoxon Signed Rank test. Beta diversity was calculated with no initial data

transformation and ASVs that were not present in more than 0.1% relative abundance in

any sample have been removed. The principal coordinates analysis was based on Bray-

Curtis distances. The effect of the route of inoculation on individual members of the

bacterial community was computed with MaAsLin2 and a minimum prevalence threshold

of 0.1 and a minimum relative abundance threshold 0.01.Benjamini-Hochberg procedure

was applied as a correction method for computing the q-values. All raw FASTQ reads for

16S rRNA gene sequences have been deposited under NCBI accession number:

Real-time qPCR. Real-time qPCR amplification was performed on DNA extracted from

the ceca and litter of broiler chicks as described previously[37] using a CFX96 Touch Real-

Time PCR Detection System (Bio-Rad Inc., Hercules, CA). Reaction mixtures (20 µl) for

all assays contained 1X SsoAdvanced Universal SYBR Green Supermix (Bio-Rad Inc.,

Hercules, CA), 600 nM (each) primers (Supplementary Table 5), and 2 µl of cecal/litter

DNA. Calibration curves used for converting qPCR cycle threshold values to gene copies

per gram were determined using genomic DNA of *E. coli* or *Salmonella* strains harboring

the region or gene of interest

Statistical analyses. Continuous variables did not meet the assumption of a normal

distribution; therefore, non-parametric testing for direct comparisons was performed using

Wilcoxon rank sum and signed rank tests, and the Kruskal-Wallis rank sum test was used

for one-way analysis of variance tests. Furthermore, continuous variables were log-

transformed before any statistical tests were performed. Lastly, principal component

analysis was performed on litter and ceca samples using the copy number of genes

quantified by qPCR as features. The samples were then projected onto the first two

principal components and colored by route of SH challenge to visualize each route's

contribution to the data variance. Pearson correlation coefficients were calculated to

examine the correlation between *Salmonella ttr* and *E. coli gapA* gene copies present in

ceca and litter. Statistical analyses were performed using R (v 3.4.1).

Data availability. All raw FASTQ reads including short and long reads for sequenced

bacterial genomes are publicly available under NCBI accession numbers: PRJNA683658,

PRJNA684578 and PRJNA684580. Shotgun and Hi-C reads are publicly available under

NCBI accession number: PRJNA688069 and 16S rRNA gene sequences under NCBI

accession number: PRJNA669215. The whole genome assembly for *S*. Heidelberg

strains SH-2813-ancestor and ic9b harboring p1ST26 have been made available under

NCBI biosample number: SAMN12082795 and DDBJ/ENA/GenBank accession numbers

JAEMHU000000000, respectively. The whole genome assembly for *E. coli* strain Ec-FL1-

1X and Ec-FL1-2X carrying IncI1 are available under GenBank accession numbers

CP066836 and JAFCXR000000000, respectively. Variant call format (VCF) files of

identified SNPs/indels and the Linux/Unix shell script used have been deposited in Dryad

Digital Repository: https://doi.org/10.5061/dryad.4tmpg4f8d.

Ethics statement. All animal experiments were approved by the University of Georgia

Office of Animal Care and Use under Animal Use Protocol: A2017 04-028-A2.

Acknowledgements. We are grateful to Marlo Sommers, Jasmine Johnson, Carolina

Hall, Jeromey Jackson, and Latoya Wiggins for their logistical and technical assistance.

This work was supported by USDA Agricultural Research Service (Project Number: 6040-

32000-010-00-D), non-assistance cooperative agreement (58-6040-6-030) between

USDA Agricultural Research Service and University of Georgia, Research Foundation,

and research service agreement (58-6040-8-035) between USDA Agricultural Research

Service and Colorado State University. This work was also partially supported by

Colorado State University through startup funds to ZA. BZ was supported by the

competence center. The competence center (FFoQSI) is funded by the Austrian

ministries BMVIT, BMDW, and the Austrian provinces Niederoesterreich, Upper Austria

and Vienna within the scope of COMET-Competence Centers for Excellent Technologies.

The Austrian Research Promotion Agency FFG handle the program COMET. This study

was supported in part by resources and technical expertise from the Georgia Advanced

Computing Resource Center, a partnership between the University of Georgia's Office of

the Vice President for Research and Office of the Vice President for Information

Technology. Any opinions expressed in this paper are those of the authors and do not

necessarily reflect the official positions and policies of the USDA or the National Science

Foundation, and any mention of products or trade names does not constitute

recommendation for use. The authors declare no competing commercial interests in

relation to the submitted work. USDA is an equal opportunity provider and employer.

Author's contributions. A.O., Z.A., K.C. and N.A.C. designed the study. A.O., G.Z.,

D.E.C. and C.R. performed live-broiler chicken studies. A.O., J.L., G.Z., S.H. and D.C.

performed bacteriological analyses, antibiotic susceptibility testing, DNA extraction and

qPCR. J.L., G.Z. and A.O. performed Illumina whole genome sequencing. A.O. performed

cecal shotgun and Hi-C library preparation. T.L. made 16S rRNA gene libraries and

sequencing. B.Z. performed 16S rRNA bacterial community analysis and interpretation.

K.I., A.H., and A.O. performed phenotype microarray analyses with the help of J.G. and

E.L. A.O., Z.A., M.O.P., J.C.T., S.M.L., R.W., J.L., D.C. and L.L performed bioinformatic

analyses and data curation. K.H. and A.O performed *Salmonella* Heidelberg fitness

testing and analysis. A.O., J.L. and D.C. isolated multidrug resistant *E. coli* donors from

cecal contents with help from M.J.R. A.O., B.Z. and M.E.B. performed statistical analyses.

A.O., Z.A., N.A.C., B.Z., R.W., M.P., S.M.L., K.I., G.Z., J.L., K.H. and L.L. drafted the

manuscript, which was reviewed and edited by all authors. A.O., Z.A., S.A.A., L.C. and

M.W. supervised the study.

References

1   Frieri, M., Kumar, K. & Boutin, A. Antibiotic resistance. *J Infect Public Health* **10**,

369-378, doi:10.1016/j.jiph.2016.08.007 (2017).

2   Akova, M. Epidemiology of antimicrobial resistance in bloodstream infections.

Virulence **7**, 252-266, doi:10.1080/21505594.2016.1159366 (2016).

3   Von Wintersdorff, C. J. *et al.* Dissemination of antimicrobial resistance in microbial

ecosystems through horizontal gene transfer. *Frontiers in microbiology* **7**, 173

(2016).

4   Ventola CL. The Antibiotic Resistance Crisis. P T. 2015;40: 277–283.
735
5   Angulo FJ, Baker NL, Olsen SJ, Anderson A, Barrett TJ. Antimicrobial use in

agriculture: controlling the transfer of antimicrobial resistance to humans1 1This

article is based in part on an appendix written for an Institute of Medicine workshop

summary by Anderson AD, McClellan J, Rossiter S, Angulo FJ entitled Public

Health Consequences of Use of Antimicrobial Agents in Agriculture in "The

Resistance Phenomenon in Microbes and Infectious Disease Vectors: Implications

for Human Health and Strategies for Containment-Workshop Summary," Stacey

L. Knobler, Stanley M. Lemon, Marian Najafi, and Tom Burroughs (Eds.), Institute

of Medicine of the National Academies, Washington, DC, The National Academies

Press, 231–243, 2003. Seminars in Pediatric Infectious Diseases. 2004;15: 78–

85. doi:10.1053/j.spid.2004.01.0106

6   Ventola, C. L. The antibiotic resistance crisis: part 2: management strategies and

new agents. *Pharmacy and Therapeutics* **40**, 344 (2015).

7   Scott, A. M. *et al.* Is antimicrobial administration to food animals a direct threat to

human health? A rapid systematic review. *International journal of antimicrobial*

agents **52**, 316-323 (2018).

8   Aidara-Kane, A. *et al.* World Health Organization (WHO) guidelines on use of

medically important antimicrobials in food-producing animals. *Antimicrobial*

Resistance & Infection Control **7**, 1-8 (2018).

9   Tang, K. L. *et al.* Restricting the use of antibiotics in food-producing animals and

its associations with antibiotic resistance in food-producing animals and human

beings: a systematic review and meta-analysis. *The Lancet Planetary Health* **1**,

e316-e327 (2017).

10   Singer, R. & Porter, L. Estimates of on-farm antimicrobial usage in broiler chicken

and turkey production in the united states, 2013-2017. Retrieved from

https://www.uspoultry.org/poultry-antimicrobial-use-

report/docs/USPOULTRY_Antimicrobial-Report.pdf. (2019)

11   Karavolias, J., Salois, M. J., Baker, K. T. & Watkins, K. Raised without antibiotics:

impact on animal welfare and implications for food policy. *Translational Animal*

Science **2**, 337-348, doi:10.1093/tas/txy016 (2018).

12   Bekal, S. *et al.* Usefulness of High-Quality Core Genome Single-Nucleotide Variant

Analysis for Subtyping the Highly Clonal and the Most Prevalent Salmonella

enterica Serovar Heidelberg Clone in the Context of Outbreak Investigations. *J*

Clin Microbiol **54**, 289-295, doi:10.1128/Jcm.02200-15 (2016).

13   Liakopoulos, A. *et al.* Extended-Spectrum Cephalosporin-Resistant Salmonella

enterica serovar Heidelberg Strains, the Netherlands(1). *Emerg Infect Dis* **22**,

1257-1261, doi:10.3201/eid2207.151377 (2016).

14   Edirmanasinghe, R. *et al.* A Whole-Genome Sequencing Approach To Study

Cefoxitin-Resistant Salmonella enterica Serovar Heidelberg Isolates from Various

Sources. *Antimicrob Agents Chemother* **61**, doi:10.1128/AAC.01919-16 (2017).

15   Gieraltowski, L. *et al.* National Outbreak of Multidrug Resistant Salmonella

Heidelberg Infections Linked to a Single Poultry Company. *PLoS One* **11**,

e0162369, doi:10.1371/journal.pone.0162369 (2016).

16   Cox, N. A. *et al.* Research Note: Evaluation of several inoculation procedures for

colonization of day-old broiler chicks with Salmonella Heidelberg. *Poult Sci* **99**,

1615-1617, doi:10.1016/j.psj.2019.10.020 (2020).

17   Rehman, M. A., Yin, X., Persaud-Lachhman, M. G. & Diarra, M. S. First Detection

of a Fosfomycin Resistance Gene, fosA7, in Salmonella enterica Serovar

Heidelberg Isolated from Broiler Chickens. *Antimicrob Agents Chemother* **61**,

doi:10.1128/AAC.00410-17 (2017).

18   Carattoli, A. *et al.* In Silico Detection and Typing of Plasmids using PlasmidFinder

and Plasmid Multilocus Sequence Typing. *Antimicrob Agents Ch* **58**, 3895-3903,

doi:10.1128/Aac.02412-14 (2014).

19 Merrikh, C. N. & Merrikh, H. Gene inversion potentiates bacterial evolvability and
virulence. *Nat Commun* **9**, 4662, doi:10.1038/s41467-018-07110-3 (2018).

20 Lucas Lledo, J. I. & Caceres, M. On the power and the systematic biases of the
detection of chromosomal inversions by paired-end genome sequencing. *PLoS*
One **8**, e61292, doi:10.1371/journal.pone.0061292 (2013).

21 Carattoli, A., Villa, L., Fortini, D. & Garcia-Fernandez, A. Contemporary IncI1
plasmids involved in the transmission and spread of antimicrobial resistance in
Enterobacteriaceae. *Plasmid*, doi:10.1016/j.plasmid.2018.12.001 (2018).

22 Stalder, T., Press, M. O., Sullivan, S., Liachko, I. & Top, E. M. Linking the resistome
and plasmidome to the microbiome. *ISME J* **13**, 2437-2446, doi:10.1038/s41396-
019-0446-4 (2019).

23 Wood, D. E., Lu, J. & Langmead, B. Improved metagenomic analysis with Kraken
2. *Genome biology* **20**, 257 (2019).

24 Ondov, B. D. *et al.* Mash Screen: high-throughput sequence containment
estimation for genome discovery. *Genome Biol* **20**, 232, doi:10.1186/s13059-019-
1841-x (2019).

25 Yoshida, C. E. *et al.* The Salmonella In Silico Typing Resource (SISTR): An Open
Web-Accessible Tool for Rapidly Typing and Subtyping Draft Salmonella Genome
Assemblies. *PLoS One* **11**, e0147101, doi:10.1371/journal.pone.0147101 (2016).

26 Cox, N. A. *et al.* Fifty percent colonization dose for Salmonella typhimurium
administered orally and intracloacally to young broiler chicks. *Poult Sci* **69**, 1809-
1812, doi:10.3382/ps.0691809 (1990).

27 Bailey, J. S., Cox, N. A., Cosby, D. E. & Richardson, L. J. Movement and
persistence of Salmonella in broiler chickens following oral or intracloacal
inoculation. *J Food Prot* **68**, 2698-2701, doi:10.4315/0362-028x-68.12.2698
(2005).
28 Cox, N., Richardson, L., Cosby, D., Berrang, M. & Harrison, M. Recovery of
Campylobacter from External and Internal Spleen Samples from Baby Broiler
Chicks Following Various Routes of Inoculation. *Journal of Food Safety* **36**, 132-
135 (2016).
29 Dimitriu, T. *et al.* Negative frequency dependent selection on plasmid carriage and
low fitness costs maintain extended spectrum beta-lactamases in Escherichia coli.
Sci Rep **9**, 17211, doi:10.1038/s41598-019-53575-7 (2019).
30 Benz, F. *et al.* Plasmid- and strain-specific factors drive variation in ESBL-plasmid
spread in vitro and in vivo. *ISME J*, doi:10.1038/s41396-020-00819-4 (2020).
31 Wotzka, S. Y. *et al.* Escherichia coli limits Salmonella Typhimurium infections after
diet shifts and fat-mediated microbiota perturbation in mice. *Nat Microbiol* **4**, 2164-
2174, doi:10.1038/s41564-019-0568-5 (2019).
32 Fischer, E. A. J. *et al.* Competition between Escherichia coli Populations with and
without Plasmids Carrying a Gene Encoding Extended-Spectrum Beta-Lactamase
in the Broiler Chicken Gut. *Appl Environ Microbiol* **85**, doi:10.1128/AEM.00892-19
(2019).
33 Hagbo, M. *et al.* Experimental support for multidrug resistance transfer potential in
the preterm infant gut microbiota. *Pediatr Res* **88**, 57-65, doi:10.1038/s41390-019-
0491-8 (2020).

34   Nyirabahizi, E. *et al.* Evaluation of Escherichia coli as an indicator for antimicrobial

resistance in Salmonella recovered from the same food or animal ceca samples.

Food Control, 107280 (2020).

35   Pereira, F. C. & Berry, D. Microbial nutrient niches in the gut. *Environ Microbiol* **19**,

1366-1378, doi:10.1111/1462-2920.13659 (2017).

36   Litvak, Y. *et al.* Commensal Enterobacteriaceae Protect against Salmonella

Colonization through Oxygen Competition. *Cell Host Microbe* **25**, 128-139 e125,

doi:10.1016/j.chom.2018.12.003 (2019).

37   Oladeinde, A. *et al.* Hotspot mutations and ColE1 plasmids contribute to the fitness

of Salmonella Heidelberg in poultry litter. *PLoS One* **13**, e0202286,

doi:10.1371/journal.pone.0202286 (2018).

38   San Millan, A., Heilbron, K. & MacLean, R. C. Positive epistasis between co-

infecting plasmids promotes plasmid survival in bacterial populations. *ISME J* **8**,

601-612, doi:10.1038/ismej.2013.182 (2014).

39   Oladeinde, A. *et al.* Horizontal Gene Transfer and Acquired Antibiotic Resistance

in Salmonella enterica Serovar Heidelberg following In Vitro Incubation in Broiler

Ceca. *Appl Environ Microbiol* **85**, doi:10.1128/AEM.01903-19 (2019).

40   Woyda, R. R., Oladeinde, A. & Abdo, Z. Reads2Resistome: An adaptable and

high-throughput whole-genome sequencing pipeline for bacterial resistome

characterization. *bioRxiv*, 2020.2005.2018.102715,

doi:10.1101/2020.05.18.102715 (2020).

41   Wick, R. R., Judd, L. M., Gorrie, C. L. & Holt, K. E. Unicycler: Resolving bacterial
genome assemblies from short and long sequencing reads. *PLoS Comput Biol* **13**,
e1005595, doi:10.1371/journal.pcbi.1005595 (2017).

42   Antipov, D., Korobeynikov, A., McLean, J. S. & Pevzner, P. A. hybridSPAdes: an
algorithm for hybrid assembly of short and long reads. *Bioinformatics* **32**, 1009-
1015, doi:10.1093/bioinformatics/btv688 (2016).

43   Gurevich, A., Saveliev, V., Vyahhi, N. & Tesler, G. QUAST: quality assessment
tool   for   genome   assemblies.   *Bioinformatics*   **29**,   1072-1075,
doi:10.1093/bioinformatics/btt086 (2013).

44   Seemann, T. Prokka: rapid prokaryotic genome annotation. *Bioinformatics* **30**,
2068-2069, doi:10.1093/bioinformatics/btu153 (2014).

45   Brettin, T. *et al.* RASTtk: a modular and extensible implementation of the RAST
algorithm for building custom annotation pipelines and annotating batches of
genomes. *Sci Rep* **5**, 8365, doi:10.1038/srep08365 (2015).

46   Kanehisa, M., Sato, Y. & Morishima, K. BlastKOALA and GhostKOALA: KEGG
Tools for Functional Characterization of Genome and Metagenome Sequences. *J*
Mol Biol **428**, 726-731, doi:10.1016/j.jmb.2015.11.006 (2016).

47   Beghain, J., Bridier-Nahmias, A., Le Nagard, H., Denamur, E. & Clermont, O.
ClermonTyping: an easy-to-use and accurate in silico method for Escherichia
genus strain phylotyping. *Microb Genom* **4**, doi:10.1099/mgen.0.000192 (2018).

48   Larsen, M. V. *et al.* Multilocus sequence typing of total-genome-sequenced
bacteria. *J Clin Microbiol* **50**, 1355-1361, doi:10.1128/JCM.06094-11 (2012).

49   Gupta, S. K. *et al.* ARG-ANNOT, a New Bioinformatic Tool To Discover Antibiotic

878   Resistance Genes in Bacterial Genomes. *Antimicrob Agents Ch* **58**, 212-220,

879   doi:Doi 10.1128/Aac.01310-13 (2014).

880   50   Jia, B. *et al.* CARD 2017: expansion and model-centric curation of the

881   comprehensive antibiotic resistance database. *Nucleic Acids Res* **45**, D566-D573,

882   doi:10.1093/nar/gkw1004 (2017).

883   51   Lakin, S. M. *et al.* MEGARes: an antimicrobial resistance database for high

884   throughput   sequencing.   *Nucleic   Acids   Research*   **45**,   D574-D580,

885   doi:10.1093/nar/gkw1009 (2017).

886   52   Feldgarden, M. *et al.* Validating the AMRFinder Tool and Resistance Gene

887   Database by Using Antimicrobial Resistance Genotype-Phenotype Correlations in

888   a Collection of Isolates. *Antimicrob Agents Ch* **63**, doi:ARTN e00483-19

889   10.1128/AAC.00483-19 (2019).

890   53   Zankari, E. *et al.* Identification of acquired antimicrobial resistance genes. *J*

891   *Antimicrob Chemother* **67**, 2640-2644, doi:10.1093/jac/dks261 (2012).

892   54   Liu, B., Zheng, D., Jin, Q., Chen, L. & Yang, J. VFDB 2019: a comparative

893   pathogenomic platform with an interactive web interface. *Nucleic Acids Res* **47**,

894   D687-D692, doi:10.1093/nar/gky1080 (2019).

895   55   Zhou, Y., Liang, Y., Lynch, K. H., Dennis, J. J. & Wishart, D. S. PHAST: a fast

896   phage search tool. *Nucleic Acids Res* **39**, W347-352, doi:10.1093/nar/gkr485

897   (2011).

56   Darling, A. E., Mau, B. & Perna, N. T. progressiveMauve: multiple genome alignment with gene gain, loss and rearrangement. *PLoS One* **5**, e11147, doi:10.1371/journal.pone.0011147 (2010).

57   Katoh, K. & Standley, D. M. MAFFT multiple sequence alignment software version 7: improvements in performance and usability. *Mol Biol Evol* **30**, 772-780, doi:10.1093/molbev/mst010 (2013).

58   Page, A. J. *et al.* Roary: rapid large-scale prokaryote pan genome analysis. *Bioinformatics* **31**, 3691-3693, doi:10.1093/bioinformatics/btv421 (2015).

59   Kozlov, A., Darriba, D., Flouri, T., Morel, B. & Stamatakis, A. RAxML-NG: A fast, scalable, and user-friendly tool for maximum likelihood phylogenetic inference. *bioRxiv*, doi:10.1101/447110 (2018).

60   Darriba, D., Taboada, G. L., Doallo, R. & Posada, D. jModelTest 2: more models, new heuristics and parallel computing. *Nat Methods* **9**, 772, doi:10.1038/nmeth.2109 (2012).

61   Li, H. & Durbin, R. Fast and accurate short read alignment with Burrows-Wheeler transform. *Bioinformatics* **25**, 1754-1760, doi:10.1093/bioinformatics/btp324 (2009).

62   McKenna, A. *et al.* The Genome Analysis Toolkit: a MapReduce framework for analyzing next-generation DNA sequencing data. *Genome Res* **20**, 1297-1303, doi:10.1101/gr.107524.110 (2010).

63   Lischer, H. E. & Excoffier, L. PGDSpider: an automated data conversion tool for connecting population genetics and genomics programs. *Bioinformatics* **28**, 298-299, doi:10.1093/bioinformatics/btr642 (2012).

64   Bushnell, B., Rood, J. & Singer, E. BBMerge - Accurate paired shotgun read

merging via overlap. *PLoS One* **12**, e0185056, doi:10.1371/journal.pone.0185056

(2017).

65   Nurk, S., Meleshko, D., Korobeynikov, A. & Pevzner, P. A. metaSPAdes: a new

versatile   metagenomic   assembler.   *Genome   Res*   **27**,   824-834,

doi:10.1101/gr.213959.116 (2017).

66   Faust, G. G. & Hall, I. M. SAMBLASTER: fast duplicate marking and structural

variant   read   extraction.   *Bioinformatics*   **30**,   2503-2505,

doi:10.1093/bioinformatics/btu314 (2014).

67   Li, H. *et al.* The Sequence Alignment/Map format and SAMtools. *Bioinformatics*

25, 2078-2079, doi:10.1093/bioinformatics/btp352 (2009).

68   Press, M. O. *et al.* Hi-C deconvolution of a human gut microbiome yields high-

quality draft genomes and reveals plasmid-genome interactions. *biorxiv*, 198713

(2017).

69   Stewart, R. D. *et al.* Assembly of 913 microbial genomes from metagenomic

sequencing of the cow rumen. *Nat Commun* **9**, 870, doi:10.1038/s41467-018-

03317-6 (2018).

70   Parks, D. H., Imelfort, M., Skennerton, C. T., Hugenholtz, P. & Tyson, G. W.

CheckM: assessing the quality of microbial genomes recovered from isolates,

single   cells,   and   metagenomes.   *Genome   Res*   **25**,   1043-1055,

doi:10.1101/gr.186072.114 (2015).

71   Ondov, B. D. *et al.* Mash: fast genome and metagenome distance estimation using

MinHash. *Genome Biol* **17**, 132, doi:10.1186/s13059-016-0997-x (2016).

72　Callahan, B. J. *et al.* DADA2: High-resolution sample inference from Illumina

amplicon data. *Nat Methods* **13**, 581-583, doi:10.1038/nmeth.3869 (2016).

73　Quast, C. *et al.* The SILVA ribosomal RNA gene database project: improved data

processing and web-based tools. *Nucleic Acids Res* **41**, D590-596,

doi:10.1093/nar/gks1219 (2013).

968   **Table 1: Multidrug resistant *E. coli* genomes found in the ceca of broiler chicks**

Isolate ID	Resistance phenotype[†]	Antibiotic resistance genes[‡]	Plasmids[§]	Phylogroup	MLST
Ec-FL1-1X	**Amp Gen Str Fis Tet**	***aac(3)-lid aph(3')-la aph(6)-ld aph(3")-lb blaTEM-1B tet (B) sul2***	IncF:B:A-, IncI1	D	69
Ec-FL2-2X	**Amp Gen Str Fis Tet**	***aac(3)-lid aph(3')-la aph(6)-ld aph(3")-lb blaTEM-1B tet (B) sul2***	IncF:B:A-, IncI1	D	69
Ec-FL2-5X	**Amp Gen Str Fis Tet**	***aac(3)-lid aph(3')-la aph(6)-ld aph(3")-lb blaTEM-1B tet (B) sul2***	IncF:B:A-, IncI1	D	69
Ec-FL1-2X*	Gen Str Tet	*aadA1 aac(3)-Via tet(A)*	IncF:B:A- IncI1 IncI2 Col8282 ColRNAI	F	6858
Ec-FL1-5X*	Gen Str Tet	*aadA1 aac(3)-Via tet(A)*	IncF:B:A- IncI1 IncI2 Col8282 ColRNAI	F	6858

969
970
971
972
* *E. coli* isolates carrying IncI1-pST26 with >99% identical DNA sequence with p2ST26 acquired by SH.
† Amp – ampicillin, Gen – gentamicin, Str – streptomycin, Tet – tetracycline, Fis – sulfisoxazole.
‡ Boldness denotes antibiotic resistance genes encoded on the chromosome.
§ Illumina short reads and PacBio long reads were combined to assemble the plasmids of E. coli strains Ec-FL1-1X and Ec-FL1-2X.

Figure 1. **Broiler chicks challenged with an antibiotic susceptible strain of SH carried antibiotic resistance SH strains 2-weeks after challenge**. **a**, Experimental design for trial 1 and trial 2 conducted in

September 2017 and April 2018, respectively. **b,c** SH load in the ceca (n = 10 per trial) and litter (n = 2 per trial) 2 weeks after challenging 1-day broiler chicks (100 chicks/trial) with 10^6 colony forming units (CFU)

of SH by oral gavage, cloacal or through the seeder method (*P < 0.01 (Kruskal-Wallis test, multiple comparison correction; error bar = Standard deviation; limit of quantification = Log 1.3 CFU/g litter). **d**,

Percentage of SH isolates that acquired antibiotic resistance in trial 1 (n = 92) and trial 2 (n = 158) **e**, Percentage of SH isolates in trial 2 that acquired antibiotic resistance grouped by the route the broiler chicks

got challenged (n = 54 62, and 42 for oral, cloaca and seeder, respectively. **f,** BLASTn alignment of the Incl1-complex plasmid acquired during trial 1 (p1ST26) and trial 2 (p2ST26) – dashed black lines highlights

mobile region encoding antimicrobial resistance genes while blue dashed lines shows other regions that are different between p1ST26 and p2ST26.

Figure 2. SH infection perturbed the bacterial community present in ceca and litter of broiler chickens. a, Principal coordinates analysis of Bray-Curtis distances based on 16S rRNA gene libraries obtained
from ceca and litter samples. Each point represents values from individual libraries with colors expressing the route of inoculation of SH in for respective samples. b, Phylum-level classification of 16S rRNA gene
sequence reads in ceca and litter grouped by route of SH infection. Data represent average of amplicon sequence variant (ASV) counts from replicate libraries for each category.

Figure 3. Horizontal gene transfer and genome inversion shaped the fitness of SH in broiler chicks. a, Maximum likelihood tree constructed using accessory genes present in SH isolates (n=72) recovered

from the ceca and litter of broiler chicks colonized with SH. The GTR model of nucleotide substitution and the GAMMA model of rate heterogeneity were used for sequence evolution prediction. Numbers shown

next to the branches represent the percentage of replicate trees where associated isolates cluster together based on ~100 bootstrap replicates. Tree was rooted with the ancestral susceptible SH strain. Clade

numbers were arbitrary assigned to facilitate discussion on isolates that acquired antibiotic resistance. Blue rectangular box highlights the clade with susceptible strains nested within antibiotic resistance strains due

to sequencing bias and misassembly. (Legend: Nal – nalidixic acid, Gen – gentamicin, Str – streptomycin, Tet – tetracycline, Fis – sulfisoxazole, NA – Not applicable) **b,** Mauve visualization of the inverted genome

of SH strain og8-05a. The chromosomal contigs of og8-05a was aligned and ordered to the complete chromosome of SH ancestor. A colored similarity plot is shown for each genome, the height of which is

proportional to the level of sequence identity in that region. When the similarity plot points downward it indicates an alignment to the reverse strand of the SH ancestor genome i.e., inversion. Segment highlighted

with solid black rectangular box shows the inverted region in og8-05a, while dashed blue and red rectangular boxes denotes segments with relevant mutations.

1006

Figure 4. Gene flux contributed to the diversity of Incl1-pMLST26 plasmids. **a,** Maximum likelihood tree constructed using TraY and ExcA protein sequences from a representative Incl1 plasmid from this study, R64 (Incl1) and R621a (Incl1-gamma). The GTR model of nucleotide substitution and the GAMMA model of rate heterogeneity were used for sequence evolution prediction. Tree was rooted with traY sequence of R621a. **b - d,** Maximum likelihood tree of pST26 Incl1 plasmids from this study (n = 36) constructed using complete plasmid DNA sequences, core genes (**c**) and accessory genes (**d**). The GTR+I, GTR and JC+I model of nucleotide substitution and the GAMMA model of rate heterogeneity were used for sequence evolution prediction for b, c, and d, respectively (Tree was rooted with the closest relative (GenBank: CP016585) found through NCBI BLASTn)). Clade numbering were arbitrary assigned to show number of clades found and not order of evolution. Numbers shown next to the branches represent the percentage of replicate trees where associated isolates cluster together based on ~100 bootstrap replicates.

1018

Figure 5. ***Proteobacteria*** **was the reservoir of Incl1-plasmids in broiler chickens. a,** Percentage of metagenomic reads assigned to the top 20 bacterial genera of the phylum Proteobacteria present in samples
Hi-IC-FL1 and Hi-IC-FL2. Only reads assigned to phylum *Proteobacteria* (70.9 % of the metagenomic reads) were used and zoomed in to show 0-1% of total reads within phylum Proteobacteria (96.7% of reads
within phylum *Proteobacteria* were assigned to *Komagataeibacter*). **b and c,** Incl1 contigs hosts found by Hi-C contacts. Incl1-derived contigs (horizontal axis of heatmap) show specific Hi-C associations with
metagenome-assembled genomes (MAGs) present in samples Hi-IC-FL1 (**b**) and Hi-IC-FL2 (**c**) (vertical axis of heatmap). MAGs are derived from Hi-C deconvolution of the metagenome assembly and placed into
a bacterial phylogeny using CheckM. Cluster.10 in each sample is an *Escherichia coli* genome. In Hi-IC-FL1, one MAG (cluster.82) representing an extremely fragmented and partially contaminated *Escherichia
coli* genome is omitted for clarity (due to its contamination, this cluster was placed in *Clostridia* by CheckM). Heatmap values indicate transformed counts of Hi-C read contacts (indicating intracellular physical
proximity of Incl1 contigs to those genomes. Heatmap values were pseudocounted to facilitate plotting of log-transformed data including zeroes.

1026

1027

1028

1029 **Figure 6. SH challenge affected the relative abundance of antibiotic resistance genes present in the ceca and litter of broiler chickens.** a, Concentration of selected antibiotic resistance determinants,
1030 IncI1 and *E. coli* present in the ceca of broiler chickens. Gene copies were determined using qPCR. Primers targeting the housekeeping gene (*gapA*) and *inc*RNAI was used for *E. coli* and IncI1 quantification,
1031 respectively. b, Pearson correlation between *Salmonella ttr* and *E. coli gapA* gene copies present in ceca and litter (n =15).

1032

1033

1034

1035

1036

1037

1038

1039

1040

1041

1042

1043

Figure 7. SH exposure to acidic pH imposes negative frequency dependent selection for Incl1 (pST26). **a** Box plot comparing the metabolic activity of one evolved susceptible (no Incl1) and one multidrug (MDR carrying pST26) SH strain using phenotype microarray (PM) plates (Biolog Inc). Pre-configured PM plates are composed of microtiter plates with one negative control well and 95 wells pre-filled with or without nitrogen and at pH of 3.5 – 10 (n = 1 plate per strain; no of wells per plate for pH 3.5-4, 4.5, 5-8.5 and 9-10 was 2, 6, 37, and 38 respectively; ***$P < 0.001$ (Wilcoxon signed rank test). **b and c,** Fitness and abundance of evolved susceptible and MDR SH strains when exposed to pine shaving extract of pH 2.5. Each bar represents the average fitness or abundance (colony forming units per ml of pine shaving extract) of three individual population that was established from three single bacterial colonies from three different strains (Error bar = Standard deviation). The horizontal line in **b** represents the fitness of three SH-Ancestor population established from three single bacterial colonies. (Anc = Ancestor, Evol – Evolved i.e., SH isolate recovered during *in vivo* experiment, genR – population resistant to gentamicin).

1050

1051

1052

Gene flux and acid-imposed selection are the main drivers of antimicrobial resistance in broiler chicks infected with *Salmonella enterica* serovar Heidelberg

Adelumola Oladeinde[1,2,10*], Zaid Abdo[2*], Maximilian O. Press[3¶], Kimberly Cook[4], Nelson A. Cox[1], Benjamin Zwirzitz[5,6¶], Reed Woyda[2¶], Steven M. Lakin[2¶], Jesse C. Thomas IV[7], Torey Looft[8], Douglas E. Cosby[1], Arthur Hinton Jr.[1], Jean Guard[1], Eric Line[1], Michael J. Rothrock[1], Mark E. Berrang[1], Kyler Herrington[9], Gregory Zock[10], Jodie Plumblee Lawrence[1], Denice Cudnik[1], Sandra House[1], Kimberly Ingram[1], Leah Lariscy[1], Robert Wagner[5,6], Samuel E. Aggrey[10], Lilong Chai[10], and Casey Ritz[10]

Authors' Affiliation

[1]U.S. National Poultry Research Center, USDA-ARS, Athens, GA, USA. [2]Department of Microbiology, Immunology and Pathology, Colorado State University, Fort Collins, Colorado, USA. [3]Phase Genomics Inc, Seattle, WA, 98109, USA, [4]Office of National Programs, USDA-ARS, Beltsville, Maryland, USA. [5]Institute of Food Safety, Food Technology and Veterinary Public Health, University of Veterinary Medicine, Vienna, Austria. [6]Austrian Competence Centre for Feed and Food Quality, Safety, and Innovation FFoQSI GmbH, Tulln, Austria. [7]Division of STD Prevention, National Center for HIV/AIDS, Viral Hepatitis, STD and TB Prevention, Center for Disease and Control, Atlanta, Georgia, USA. [8]National Animal Disease Center, USDA-ARS, Ames, IA, USA, [9]Medical College of Georgia, Augusta, GA, USA. [10] Poultry Science Dept, University of Georgia, Athens, GA, USA

Corresponding authors* ade.oladeinde@usda.gov; zaid.abdo@colostate.edu

[¶] These authors contributed equally to this work.

Supplementary Methods

Isolation of *E. coli* from ceca of broiler chicks

To isolate *E. coli*, we retrospectively screened two cecal slurry from intracloacally challenged chicks on CHROMagar™ plates supplemented with gentamicin (8 ppm) and tetracycline (8 ppm). Frozen vials of cecal contents were thawed on ice and vortexed. For each sample, a 10 μl aliquot was struck for isolation, and a 100 μl aliquot was spread plated on CHROMagar™ with relevant antibiotics. Plates were incubated overnight at 37°C. Blue green and blue-cream colonies were counted as presumptive *E. coli* and colonies was restruck for isolation on CHROMagar™ supplemented with gentamicin and tetracycline. After incubation, 2- 3 colonies from each sample were selected to represent different forms of blue green/blue-cream colonies and struck in parallel onto sheep blood agar (SBA) and Eosin Methylene Blue (EMB) (Remel Inc., KS, USA) and incubated overnight. Growth on EMB was the characteristic green metallic sheen after incubation. Colonies from SBA were stored in 30% LB glycerol at -80°C and were used for antimicrobial susceptibility testing and whole genome sequencing.

Long read sequencing quality control. PacBio sequencing of samples was done using using the Sequel II system. MinION sequencing of samples was done using R9.5 chemistry on a 1D flowcell. Lima (https://github.com/PacificBiosciences/pbbioconda) was used to demultiplex barcoded PacBio samples and to convert the split BAM files into fastq format. BBMap reformat.sh (https://github.com/BioInfoTools/BBMap) was used to randomly subsample generated PacBio reads down to 200X coverage. Porechop v0.2.4 (https://github.com/rrwick/Porechop) was used to demultiplex barcoded MinION samples. Raw Illumina reads were trimmed using Trimmomatic v0.39[1] (command line parameters: PE ILLUMINACLIP:NexteraPE-PE.fa:2:30:10:3 LEADING:3 TRAILING:3 SLIDINGWINDOW:4:20 MINLEN:36).

Comparative genomics. To determine IncI1-pST26 consensus, first we used BWA[2] or Bowtie v. 7.2.1[3] (implemented in Geneious Prime®) to align raw FASTQ files against a complete circular p1ST26 or p2ST26 and used the binary alignment map file generated for consensus determination in Geneious Prime®. To identify identical IncI1-pST26 plasmids, we performed a BLAST search against the NCBI non-redundant database and selected the top 50 plasmids matching p2ST26 from this study. To construct a whole plasmid

based maximum likelihood tree, we downloaded the FASTA files for the top 50 plasmids from NCBI and used them for whole genome alignment.

Tools used for visualizing and exploring high-throughput sequence data. Gene presence/absence heatmap, with gene presence/absence data obtained from Roary, was generated using the pheatmap v1.0.12, tidyverse v1.3.0 and viridis v0.5.1 packages in R v4.0.2. BLAST Ring Image Generator[4] was used for genome comparison visualization including GC skew change and Phandango[5] was used for visualizing phylogenetic trees with their associated metadata. PHYLOViZ 2.0[6] was used to generate a minimum spanning tree of SH isolates from core genome information obtained through SISTR (i.e., wgMLST_330:complete-alleles, wgMLST_330:missing-alleles, wgMLST_330:partial-alleles,closest-public-genome-alleles-matching and cgMLST_cluster_level). SnapGene® was used for drawing linear maps of plasmid, prophages, and other regions of interest in bacterial genomes. Geneious Prime® and SnapGene® were used to view MAFFT alignments of DNA and amino-acid sequences.

References:

Bolger, A. M., Lohse, M. & Usadel, B. Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics* **30**, 2114-2120, doi:10.1093/bioinformatics/btu170 (2014).

Li, H. & Durbin, R. Fast and accurate short read alignment with Burrows-Wheeler transform. *Bioinformatics* **25**, 1754-1760, doi:10.1093/bioinformatics/btp324 (2009).

Langmead, B. & Salzberg, S. L. Fast gapped-read alignment with Bowtie 2. *Nat Methods* **9**, 357-359, doi:10.1038/nmeth.1923 (2012).

Alikhan, N. F., Petty, N. K., Ben Zakour, N. L. & Beatson, S. A. BLAST Ring Image Generator (BRIG): simple prokaryote genome comparisons. *BMC Genomics* **12**, 402, doi:10.1186/1471-2164-12-402 (2011).

Hadfield, J. *et al.* Phandango: an interactive viewer for bacterial population genomics. *Bioinformatics* **34**, 292-293, doi:10.1093/bioinformatics/btx610 (2018).

Nascimento, M. *et al.* PHYLOViZ 2.0: providing scalable data integration and visualization for multiple phylogenetic inference methods. *Bioinformatics* **33**, 128-129, doi:10.1093/bioinformatics/btw582 (2017).

Supplementary Table 1: Assembly report for a *S*. Heidelberg strain with sequencing bias

Assembly statistics[*]	Short read[†]	Long read [‡]	Hybrid[§]
Number of contigs	554	4	33
Total length (bp)	4314964	4703397	4865484
Reference length (bp)	4788855	4788855	4788855
Number of misassembled contigs	4	0	1
Misassembled contigs length (bp)	47430	0	4742328
Genome fraction of reference (%)	89.969	98.247	99.815
Largest alignment (bp)	60767	2669051	3233973
Total aligned length (bp)	4308655	4703397	4826756

[*] Assembly was done using Unicycler and SH-ancestor was used as the reference genome (4,788,855 bp)
[†] Illumina short technology was used for sequencing.
[‡] PacBio long read sequencing technology was used.
[§] Illumina short reads and PacBio long reads were combined.

Supplementary Table 2. Mutations found in p1ST26 but absent in p2ST26

Loci*	Location†	Product	Mutation‡
20998	Intergenic	Upstream replication protein	N/A
23599	CDS	*yacA*	Substitution
23637	CDS	*yacA*	Substitution
23661	CDS	*yacA*	Substitution
23695	CDS	*yacA*	Substitution
23698	CDS	*yacA*	Substitution
23704	CDS	*yacA*	Substitution
23731	CDS	*yacA*	Substitution
23734	CDS	*yacA*	Substitution
23826	CDS	*parE*	Substitution
25883	CDS	*tniA*	Insertion
35066	CDS	*aac3-Via*	Substitution
35608	CDS	*aadA1*	Substitution
35635	CDS	*aadA1*	Substitution
38685	CDS	*tn21*	Substitution
41737	Intergenic	downstream tn3; upstream hypothetical	N/A
42362	CDS	hypothetical	Substitution
49072	CDS	hypothetical	Substitution
70253	CDS	hypothetical	Substitution
70287	CDS	hypothetical	Substitution
70333	CDS	hypothetical	Substitution
70343	CDS	hypothetical	Substitution
92634	CDS	*icmG/dotF*	Substitution

*A circular and complete IncI1 plasmid from Trial 2 was used as the reference genome (111, 898 bp).
†CDS- Coding DNA sequence
‡N/A – Not applicable

Supplementary Table 3. Plasmids found in the cecal metagenome [†]

Hi-IC-FL1	Hi-IC-FL2
IncI1, IncI2, IncFIB, IncFII, IncFIC, IncHIB, p0111, Col8282, ColRNAI	p0111

[†] We searched the cecal metagenome of two cecal samples for plasmid *inc* groups that are commonly found in family *Enterobacteriaceae*

Supplementary Table 4: Concentration of metals and pollutants in starter feed.

Starter feed*	Parts per million (ppm)	Percent (%)	Pounds per ton
Total Acid Digestion			
Antimony (Sb)	<0.4	negligible	negligible
Arsenic (As)	0.999	negligible	negligible
Beryllium (Be)	<0.05	negligible	negligible
Cadmium (Cd)	0.109	negligible	negligible
Chromium (Cr)	2.74	0.0003	0.01
Copper (Cu)	7.96	0.0008	0.02
Lead (Pb)	0.326	negligible	negligible
Nickel (Ni)	1.67	0.0002	0
Selenium (Se)	0.985	negligible	negligible
Silver (Ag)	<0.05	negligible	negligible
Thallium (Tl)	<0.4	negligible	negligible
Zinc (Zn)	82.7	0.0083	0.17

*Unmedicated starter feed was synthesized by University of Georgia's poultry research center's feed mill.

Supplementary Table 5. Primers used for the qPCR analysis of ceca and litter.

Gene	Target description	Primer Sequences (5' - 3')	Amplification factor*	Reference
aadA	Aminoglycoside resistance	F: CAGCGCAATGACATTCTTGC R: GTCGGCAGCGACAYCCTTCG	2.02	73
gapA	D-glyceraldehyde-3-phosphate dehydrogenase (GAPDH) in *E. coli*	F: CCGTTGAAGTGAAAGACGGTC R: AACCACTTTCTTCGCACCAGC	1.98	74
IncI1	*inc*RNAi	F: CAGGAGAGATGGCATGTA R: GGGTTTTCCTTTTATGGC	1.94	This study
intI1	Class 1 integron	F: CCTCCCGCACGATGATC R: TCCACGCATCGTCAGGC	1.98	75
tetA	Tetracycline resistance	F: TGTCCACCAACTTATCAG R: TGCTCAGAATTACGATCA	1.81	This study
ttr	*Salmonella*	F: CTCACCAGGAGATTACAACATGG R: AGCTCAGACCAAAAGTGACCATC	1.86	76

*Genomic DNA extracted from an *E. coli* strain carrying *aadA1, aac(3)-Via, tetA* and *IntI1* on IncI1-pST26 plasmid was used for the calibration curves of *aadA, tetA, IntI1,* IncI1 and *gapA* whereas DNA from an SH-2813-ancestor was used for the calibration curve of *ttr*.

Extended Data Fig. 1. pH and moisture levels of pen litter. Litter was collected as grab samples (n = 7) and pooled into one whirl pak bag (n = 1 per pen). The pH of the litter was determined from the eluate made from a 1:5 dilution of litter in 1X PBS.

Extended Fig. 2 Challenging broiler chicks with SH affected the diversity of bacterial species in their ceca and litter. a, Average alpha diversity indices of ceca and litter samples grouped by route of SH infection. Boxes indicate the interquartile range (75th to 25th) of the data. Whiskers extend to the most extreme value within 1.5 * interquartile range and dots represent outliers beyond that range. d, Heatmap of the top 50 most abundant amplicon sequence variants (ASV) grouped by route of SH infection and split by sample type (Ceca, Litter). Data represents average of ASV counts from replicate 16S rRNA gene libraries for each category.

Extended Data Fig. 3. Acquisition of IncI1 plasmid conferred multidrug resistance on SH cells and influenced the contents of their genomes. Antimicrobial susceptibility and whole genome sequencing were performed on SH isolates recovered from the ceca and litter of broiler chicks 2-weeks after challenge. **a**, Percentage of SH isolates (n = 63) from Trial 2 that acquired resistance to two or more antibiotics compared to isolates resistance to only one antibiotic class. Core genome (**b**) and SNP (**c**) based maximum likelihood tree of SH isolates sequenced from Trial 1 (n =2) and Trial 2 (n=69). Isolates with duplicated SNPs/indels were removed before SNP based ML tree was reconstructed. GTR and JC model of nucleotide substitution was used for core and SNP tree, respectively and the GAMMA model of rate heterogeneity were used for sequence evolution prediction. Numbers shown next to the branches represent the percentage of replicate trees where associated isolates cluster together based on ~100 bootstrap replicates. Tree was rooted with the susceptible SH ancestor for core genome and nal resistant SH ancestor (SH-2813nalR) for SNP tree. (Legend: Nal – nalidixic acid, Gen – gentamicin, Str – streptomycin, Tet – tetracycline, Fis – sulfisoxazole, NA – Not applicable) .

Extended Data Fig. 4. Gene gain and loss contributed to SH diversity in the ceca and litter of broiler chicks a, Heatmap of genes present and absent in SH genomes determined by pangenome analysis. Thick rectangular box highlights SH strains that exhibited higher gene flux and strains with no plasmids but were grouped with SH strains carrying IncI1. **b,c,d** SH genome assembly statistics for SH strains with higher gene flux (n = 12; clade IIa of ML tree reconstructed using accessory genes (see Fig. 2a)) compared to other SH (n=59). (***P < 0.001 (Binomial logistic regression; Error bar = Standard deviation). **e**, Circular map of novel episome present in SH strain og8-05a. **f**, BLASTn alignment of IncX1 present in SH ancestor and IncX1 present in SH strain og8-05a.

Extended Data Fig. 5. Chromosomal inversion re-oriented the origin of replication of SH strains and positively skewed their GC. Complete chromosome contigs from three SH strains recovered from ceca during Trial 1 (ic9b) and 2 (ic4-3a and og8-05a) were aligned and compared to the ancestor. a, ProgressiveMauve alignment of SH strains ic4-3a and ic9b. A colored similarity plot is shown for each genome, the height of which is proportional to the level of sequence identity in that region. When the similarity plot points downward it indicates an alignment to the reverse strand of the genome of SH-ancestor i.e., inversion. Segment highlighted with solid black rectangular box denotes the inverted 2Mbp region in ic9b. Segment highlighted with horizontal blue box denotes regions with relevant mutations. b and c, GC skew in og8-05a and ic9b compared to SH-ancestor. Red arrow points towards reoriented oriC or ter, whereas red and blue arcs show the inverted region. The number of guanine and cytosine in inverted area was used for GC skew calculation. Dashed blue and red rectangular boxes denotes segments with mutations.

Extended Data Fig. 6 Linear maps of prophages found in SH ancestor.

Extended Data Fig. 7 Linear map of the variable region encoding antimicrobial resistance genes in IncI1. The ~19kb DNA region of the p1ST26 plasmid (~113kb) present in one multidrug resistant SH isolate from Trial 1 was used for drawing the map.

Extended Data Fig. 8. SH challenge affected the abundance of antimicrobial genes a, Concentration of *E. coli* and resistome (Incl1, Intl1 and antibiotic resistance genes) present in the litter of broiler chicks. Gene copies was determined using qPCR. b, Principal component analysis (PCA) characterizing the differences between samples grouped by route broiler chicks got challenged with SH. The PCA was constructed using the concentration of *E. coli, Salmonella* and resistome genes present in ceca and litter of broiler chicks challenged with SH and determined using qPCR. Each number and dot denote one sample, and data is from 32 samples. The farther apart two samples are from each other, the more different they are. Ellipses represents the mean/confidence interval if all the dots of the same color are represented by just one sample.

Extended Data Fig. 9. IncI1-pST26 acquired by SH are identical to the pST26 present in commensal *E. coli* strains. Complete pST26 contigs from two SH and one *E. coli* strain recovered from the ceca of broiler chicks were used for ProgressiveMauve alignment. A colored similarity plot is shown for each genome, the height of which is proportional to the level of sequence identity in that region. When the similarity plot points downward it indicates an alignment to the reverse strand of the genome of Ec-pST26 i.e., inversion. Segment highlighted with solid black rectangular box denotes the ~ 20kb region encoding antimicrobial resistance (AR), whereas blue horizontal box denotes region encoding transposons.

Extended Data Fig. 10. Exposing SH to metal chlorides, disinfecting compounds or neutral pH did not significantly change their metabolism. **a,b** Box plots comparing the metabolic activity of one evolved susceptible (no IncI1) and one multidrug (MDR carrying pST26) SH strain using phenotype microarray (PM) plates (Biolog Inc). Pre-configured PM plates are composed of microtiter plates with one negative control well and 95 wells pre-filled with or without nitrogen or amino acid nutrient source and at pH of 3.5 – 10 (n = 1 plate per strain; no of wells per plate per compound was 4) **c,** Concentration of evolved susceptible and MDR SH strains when exposed to pine shaving extract of pH 6.5. Each bar represents the average concentration (colony forming units per ml of pine shaving extract) of three individual population that was established from three single bacterial colonies from three different evolved strains. SH-Ancestor population was established from three single bacterial colonies of SH-2813nal[R]. (Anc = Ancestor, Evol – Evolved i.e., SH isolate recovered during in vivo experiment, genR – population resistant to gentamicin). (Error bars = Standard deviation; *P* > 0.05 (Wilcoxon signed rank test and Kruskal-Wallis rank sum test).

Extended Data Fig. 11. IncI1 plasmids of multilocus sequence type 26 (pST26) are carried by strains of *Salmonella enterica* and *E. coli* isolated from animal and human sources. Maximum likelihood tree reconstructed using the complete DNA sequence of the fifty closest plasmids to pST26 found using NCBI BLASTn search. After multiple alignment was performed, duplicated plasmids (n =4) before ML tree were built. JC model of nucleotide substitution and GAMMA model of rate heterogeneity was used for sequence evolution prediction. Numbers shown next to the branches represent the percentage of replicate trees where associated isolates cluster together based on ~100 bootstrap replicates. Tree was rooted with p2ST26 plasmid from this study. Taxa in blue text are pST26 plasmids from this study. (Note. ST/CC- Multi locus sequence/clonal complex; ARG – Antibiotic resistance gene)

Extended Data Fig. 12. Minimum spanning tree (MST) of SH genomes from this study. The core genome multilocus sequence type (cgMLST) data generated by SISTR for 63 genomes from this study including SH-ancestor were used for MST construction. Colors were attributed to nodes according to the number cgMLST cluster levels found (I), antibiotic resistance profile (II), isolation source (III) and plasmid carried. (i) The susceptible SH-ancestor is part of cgMLST cluster 1 (n = 18 isolates). All nodes with distances equal or above 0 were linked. All links from the MST with a distance value above 4 were deleted.

Supplementary data

Supplementary data 1: Mutations found in *S*. Heidelberg isolates recovered from ceca and litter.

Supplementary data 2: Virulence genes present in SH-ancestor and the inverted region of the genomes of SH strains ic9b and og8-05a. The number of virulence genes found in each genome is written in the column headers.

Supplementary data 3: Alignment of IncI1-pST26 plasmids from this study using ProgressiveMauve. Complete plasmid DNA consensus sequence (n= 36) was determined by aligning whole genome FASTQ reads to a complete circular pST26 plasmids from this study. IncI1-pST26 (GenBank: CP016585) present in a *S*. Heidelberg strain available on NCBI was used as a reference genome for the alignment. Segments highlighted with dashed horizontal blue box denotes regions with relevant mutations.

Supplementary data 4: Contigs matching p2ST26 found in the cecal metagenome.

Supplementary data 5: Assembly report for bacterial genomes sequenced using short and long reads technology.

Supplementary data 1: Mutations found in S. Heidelberg isolates recovered from ceca and litter

Loci[*]	Location[†]	Product	Mutation Change[‡]	Mutation type[§]	No of Isolates[‖]
95760	CDS	hypothetical protein	Pro23Thr	NS	1
130492	CDS	lipopolysaccharide 1,3-galactosyltransferase (WaaB)	Ile2Asn	NS	1
413914	CDS	putative surface-exposed virulence protein (BigA)	N/A	Ins	1
414022	CDS	putative surface-exposed virulence protein (BigA)	Val157Val	S	4
472700	CDS	16S ribosomal RNA (rRNA)	N/A	N/A	1
473535	Intergenic	upstream tRNA; downstream RNA	N/A	N/A	8
473536	Intergenic	upstream tRNA; downstream RNA	N/A	N/A	8
501858	CDS	hypothetical protein	N/A	Del	1
501860	CDS	hypothetical protein	N/A	Del	1
636934	CDS	hypothetical protein	N/A	Ins	2
693268	CDS	hypothetical protein (Kch)	Val58Met	NS	1
927855	CDS	putative regulatory protein (LysR)	Ile163Ile	S	1
1057273	CDS	N-acetylglucosaminyl transferase (MurG)	Ser322Ala	NS	1
1080335	CDS	aromatic amino acid transport protein (AroP)	Ser257Leu	NS	1
1080340	CDS	aromatic amino acid transport protein (AroP)	Leu255Leu	S	1
1208313	CDS	23S ribosomal RNA (rRNA)	N/A	N/A	21 and SH-2813nal[R]
1209197	CDS	23S ribosomal RNA (rRNA)	N/A	N/A	14
1209203	CDS	23S ribosomal RNA (rRNA)	N/A	N/A	13
1223192	Intergenic	upstream tRNA; downstream (TssA)	N/A	N/A	1
1339554	CDS	fimbriae usher protein (StbC)	N/A	Del	1
1351680	CDS	multidrug efflux system, outer membrane factor (OprM)	Thr120Pro	NS	1
1670308	CDS	altronate dehydratase (UxaA)	Leu21Leu	S	1
1744150	CDS	putative glycosyltransferase (WbbK)	Leu246Leu	S	1
1766719	Intergenic	upstream hypothetical protein; downstream hypothetical protein	N/A	N/A	1
2092586	Intergenic	upstream hypothetical protein; downstream hypothetical protein	N/A	N/A	1
2137198	CDS	glyoxylate/hydroxypyruvate reductase A (GhrA)	N/A	Del	1
2250686	CDS	putative outer membrane lipoprotein (RzoR)	Ser78Phe	NS	1
2260332	CDS	putative regulatory protein (CitB)	Arg110Trp	NS	1
2262081	Intergenic	upstream hypothetical protein; downstream hypothetical protein	N/A	N/A	1
2392728	Intergenic	upstream PykF; upstream SsaB	N/A	N/A	1
2415691	CDS	multidrug efflux transporter (MdtK/NorM)	Ala199Val	NS	1
2660396	Intergenic	downstream hypothetical protein; downstream YdbK	N/A	N/A	1
2961953	CDS	flagellin (FliC)	Gln129His	NS	70 and SH-2813nal[R]
3289224	CDS	DNA gyrase subunit A (GyrA)	Ser83Tyr	NS	70 and SH-2813nal[R]
3341410	CDS	NADH-ubiquinone oxidoreductase chain H (NuoH)	Val269Val	S	1
3356426	CDS	hexitol phosphatase (HxpA)	N/A	Del	1
3428428	Intergenic	upstream AlpA; downstream PgtE	N/A	Ins	1
3428439	Intergenic	upstream AlpA; downstream PgtE	N/A	Ins	1
3428448	Intergenic	upstream AlpA; downstream PgtE	N/A	Ins	1

3428454	Intergenic	upstream AlpA; downstream PgtE	N/A	N/A	1
3428456	Intergenic	upstream AlpA; downstream PgtE	N/A	Del	1
3428481	Intergenic	upstream AlpA; downstream PgtE	N/A	Del	19
3428490	Intergenic	upstream AlpA; downstream PgtE	N/A	N/A	19
3619800	CDS	transcriptional activator of cad operon (CadC)	N/A	Del	1
3672875	CDS	23S ribosomal RNA (rRNA)	N/A	N/A	2
3672877	CDS	23S ribosomal RNA (rRNA)	N/A	N/A	2
3672881	CDS	23S ribosomal RNA (rRNA)	N/A	Ins	2
3672883	CDS	23S ribosomal RNA (rRNA)	N/A	N/A	2
3672885	CDS	23S ribosomal RNA (rRNA)	N/A	Del	2
3672893	CDS	23S ribosomal RNA (rRNA)	N/A	N/A	2
3672899	CDS	23S ribosomal RNA (rRNA)	N/A	N/A	2
3672900	CDS	23S ribosomal RNA (rRNA)	N/A	Ins	2
3672905	CDS	23S ribosomal RNA (rRNA)	N/A	N/A	2
3672906	CDS	23S ribosomal RNA (rRNA)	N/A	N/A	2
3672912	CDS	23S ribosomal RNA (rRNA)	N/A	Del	2
3672922	CDS	23S ribosomal RNA (rRNA)	N/A	Del	2
3672935	CDS	23S ribosomal RNA (rRNA)	N/A	N/A	2
3672937	CDS	23S ribosomal RNA (rRNA)	N/A	N/A	2
3672939	CDS	23S ribosomal RNA (rRNA)	N/A	Del	2
3672948	CDS	23S ribosomal RNA (rRNA)	N/A	Del	2
3672953	CDS	23S ribosomal RNA (rRNA)	N/A	Ins	2
3672956	CDS	23S ribosomal RNA (rRNA)	N/A	N/A	2
3672958	CDS	23S ribosomal RNA (rRNA)	N/A	Ins	2
3672963	CDS	23S ribosomal RNA (rRNA)	N/A	N/A	1
3672966	CDS	23S ribosomal RNA (rRNA)	N/A	Ins	1
3672969	CDS	23S ribosomal RNA (rRNA)	N/A	N/A	1
3672970	CDS	23S ribosomal RNA (rRNA)	N/A	N/A	1
3672973	CDS	23S ribosomal RNA (rRNA)	N/A	N/A	1
3674946	CDS	16S ribosomal RNA (rRNA)	N/A	N/A	3
3674946	CDS	16S ribosomal RNA (rRNA)	N/A	N/A	4
3742216	CDS	putative inner membrane protein	Phe224Ser	NS	1
3744202	CDS	putative inner membrane protein	Glu267Glu	S	1
3930073	CDS	sulfite reductase [NADPH] hemoproteinbeta-component (CysI)	Ile352Ser	NS	70 and SH-2813nal^R
4002993	CDS	exodeoxyribonuclease V gamma chain (RecC)	Gln321Gln	S	1
4158109	CDS	hypothetical protein	Phe34Leu	NS	1
4251311	Intergenic	upstream YqjA; downstream SstT	N/A	Del	1
4369883	CDS	oxaloacetate decarboxylase Na (+) pump, alphachain (OadA)	Ala48Val	NS	1
4418343	CDS	23S ribosomal RNA (rRNA)	N/A	N/A	1
4418421	Intergenic	upstream 23S ribosomal RNA; downstream tRNA	N/A	N/A	2
4418422	Intergenic	upstream 23S ribosomal RNA; downstream tRNA	N/A	N/A	2

4696844	CDS	23S ribosomal RNA (rRNA)	N/A	N/A	38 and SH-2813nal^R
4698491	CDS	upstream 23S ribosomal RNA; downstream tRNA	N/A	N/A	68 and SH-2813nal^R
4699877	CDS	16S ribosomal RNA (rRNA)	N/A	N/A	2

* SH-ancestor was used as the reference genome (4,788,855 bp).

† CDS- Coding DNA sequence

‡ N/A – Not applicable

§ S – synonymous, NS – non-synonymous, Ins – insertion, Del – deletion, N/A – Not applicable

‖The number of evolved SH isolates carrying the same mutation. Six mutations are present in SH-2813nal^R but absent in SH-ancestor.

Virulence Factor Class	Virulence Factors	Related Genes	Ancestral=168	Ic9B=58	Og8-05a=102
Fimbrial Adherence Determinants	Agf/Csg	csgA			
		csgB			
		csgC			
		csgD			
		csgE			
		csgF			
		csgG			
	Bcf	bcfA			
		bcfB			
		bcfC			
		bcfD			
		bcfE			
		bcfF			
		bcfG			
	Fim	fimA			
		fimC			
		fimD			
		fimF			
		fimH			
		fimI			
		fimW			
		fimY			
		fimZ			
	Lpf	lpfA			
		lpfB			
		lpfC			
		lpfD			
		lpfE			
	Stb	stbA			
		stbB			
		stbC			
		stbD			
		stbE			
	Stc	stcA			
		stcB			
		stcC			
		stcD			
	Std	stdA			
		stdB			
		stdC			
	Ste	steA			
		steB			
		steC			
		steD			
		steE			
		steF			
	Stf	stfA			
		stfC			
		stfD			
		stfE			
		stfF			
		stfG			
	Sth	sthA			
		sthB			
		sthC			
		sthD			
		sthE			
	Sti	stiA			
		stiB			
		stiC			
		stiH			
	Stj	Undetermined			
		Undetermined			
		Undetermined			
		stjB			
		stjC			
	Stk	stkA			
		stkB			
		stkC			
		stkD			
		stkE			
		stkF			
		stkG			
	Tcf	tcfA			
		tcfB			
		tcfC			
		tcfD			
Macrophage Inducible	Mig-14	mig-14			
Magnesium Uptake	Mg2+ transport	mgtB			
		mgtC			
Nonfimbrial Adherence Determinants	MisL	misL			
	RatB	ratB			
	ShdA	shdA			
	SinH	sinH			
Regulation	PhoPQ	phoP			
		phoQ			
Secretion System	TTSS (SPI-1 encode)	hilA			
		hilC			
		hilD			
		iacP			
		iagB			
		invA			
		invB			
		invC			
		invE			
		invF			
		invG			
		invH			
		invI			
		invJ			
		orgA			
		orgB			
		orgC			
		prgH			
		prgI			
		prgJ			
		prgK			
		sicA			
		sicP			
		sipD			
		spaO			
		spaP			
		spaQ			
		spaR			
		spaS			
		sprB			
	TTSS (SPI-2 encode)	ssaC			
		ssaD			
		ssaE			
		ssaG			
		ssaH			
		ssaI			
		ssaJ			
		ssaK			
		ssaM			
		ssaN			
		ssaO			
		ssaP			
		ssaQ			
		ssaR			
		ssaS			
		ssaT			
		ssaU			
		ssaV			
		sscA			
		sscB			
		sseA			
		sseB			
		sseC			
		sseD			
		sseE			
		ssrA			
		ssrB			
	TTSS effectors translocated via both s	slrP			
	TTSS-1 translocated effectors	avrA			
		sipA			
		sipB			
		sipC			
		sopA			
		sopB/sigD			
		sopD			
		sopE2			
		sopE			
		sptP			
	TTSS-2 translocated effectors	pipB2			
		pipB			
		sifA			
		sifB			
		sopD2			
		spiC/ssaB			
		sseF			
		sseG			
		sseJ			
		sseK2			
		sseL			
		sspH2			
Stress Adaptation	SodCI	sodCI			

SH - CP016585

Pili/Shufflon

IS66 family of transposase

SH1 - Trial 2

SH2 - Trial 2

SH3 - Trial 2

SH4 - Trial 2

SH5 - Trial 2

SH6 - Trial 2

SH7 - Trial 2

SH8 - Trial 2

SH9 - Trial 2

SH10 - Trial 2

SH11 - Trial 2

SH12 - Trial 2

SH13 - Trial 2

SH14 - Trial 2

SH15 - Trial 2

SH16 - Trial 2

SH17 - Trial 2

SH18 - Trial 2

SH19 - Trial 2

SH20 - Trial 2

SH21 - Trial 2

SH22 - Trial 2

SH23 - Trial 2

SH24 - Trial 2

SH25 - Trial 2

SH26 - Trial 2

SH27 - Trial 2

SH28 - Trial 2

SH29 - Trial 2

SH30 - Trial 2

SH31 - Trial 2

SH32 - Trial 2

SH33 - Trial 2

SH34 - Trial 2

SH1 - Trial 1

tniA insertion

Inverted shufflon/pilV genes

SH2 - Trial 1

Supplementary data 4: Contigs matching p2ST26 found in the cecal metagenome

Protein ID	IC-FL2	IC-FL1	Number of contigs[*]
YhdJ	no	yes	1
AadA	no	yes	1
Aminoglycoside N(3)-acetyltransferase	no	yes	1
KlcA	no	yes	2
ParA	no	yes	1
Conjugal transfer protein	no	yes	3
Doc toxin	no	yes	3
Eaa	no	yes	1
Hypothetical	yes	yes	9
Hypothetical	no	yes	1
Hypothetical	no	yes	1
Hypothetical	no	yes	1
Hypothetical	yes	yes	6
Hypothetical	no	yes	3
Hypothetical	no	yes	1
Hypothetical	no	yes	2
Hypothetical	no	yes	1
DotA/TraY	no	yes	2
Replication initiation protein	yes	yes	2
DNA primase	no	yes	4
Hypothetical protein	no	yes	20
Pili assembly chaperone	no	yes	3
ParB/RepB/Spo0j	yes	yes	3
YubH	no	yes	1
Hypothetical	no	yes	2
Hypothetical	no	yes	2
Hypothetical	no	yes	1
Hypothetical	no	yes	1
MobC	no	yes	2
plasmid partitioning/stability family protein	no	yes	1
TraF	no	yes	2
TraE	no	yes	1
ArdA	no	yes	1
Rpn	yes	yes	4
3'-5' exonuclease	no	yes	1
ProQ/FINO	no	yes	1
Colicin	no	yes	1
Colicin transporter	no	yes	1
GroES	yes	yes	8
TraX	no	yes	1
TraQ	no	yes	1
TraP	no	yes	1
TraO	no	yes	1
TraN	no	yes	1
DotI/IcmL/TraM	no	yes	2
PilQ	yes	yes	2
PilR	no	yes	1
PilN	no	yes	2
TraB	no	yes	1
PilV	no	yes	2
PilS	no	yes	1
PilI	no	yes	2
PilK	no	yes	3
PilL	no	yes	1
PilM	no	yes	1
TraA	no	yes	1

Gene			
TraC	no	yes	2
TraG	no	yes	1
TraH	no	yes	1
TraI	no	yes	1
TraJ	no	yes	2
TraR	no	yes	1
TraS	no	yes	2
TraU	no	yes	2
TraW	no	yes	1
PilT	no	yes	1
PilP	no	yes	1
Integron integrase IntI1	no	yes	1
Mercuric ion reductase	yes	yes	2
MerR	no	yes	4
MerD	yes	yes	48
21PP	no	yes	1
MerT	no	yes	1
TnAs3	yes	yes	3
IS66 transposase	yes	yes	11
IS66 insertion sequence element	yes	yes	14
Helix-turn-helix domain-containing protein	yes	yes	3
Mobile element protein*	yes	no	2
MerP	yes	yes	2
DMT	no	yes	1
Plasmid conjugative transfer endonuclease	no	yes	1
PsiA	no	yes	2
PsiB	yes	yes	2
Phd	no	yes	2
StbA	no	yes	1
Relaxase/helicase	no	yes	1
Relaxase	no	yes	2
Relaxasome component	no	yes	1
Resolvase	yes	yes	3
Shufflon-specific DNA recombinase	no	yes	2
Single-stranded DNA-binding protein	yes	yes	4
Surface exclusion protein	no	yes	2
TetR	no	yes	1
TetA	no	yes	1
Urf2	no	yes	1
TniA	no	yes	2
InsC	yes	yes	3
InsD	yes	yes	4
Type-IV secretion leader peptidase/N-methyltransferase	no	yes	1
YkfF	yes	yes	2
YadA	no	yes	3
YDFB	no	yes	1

*Total number of contigs found encoding the gene in the cecal metagenome samples