Different underlying mechanisms for high and low arousal in probabilistic learning in humans

Luis F. Ciria[1,2*], Marta Suárez-Pinilla[2,3], Alex G. Williams[2], Sridhar R. Jagannathan[2], Daniel Sanabria[1], and Tristán A. Bekinschtein[2*]

[1] Mind, Brain & Behavior Research Center and Department of Experimental Psychology, University of Granada, Spain
[2] Consciousness and Cognition Lab, Department of Psychology, University of Cambridge, Downing Site, Cambridge, CB2 3EB, UK
[3] Office of the National Director for Dementia Research, Department of Neurodegenerative Disease, Institute of Neurology, University College of London, London, WC1N 3AX, UK

*Corresponding authors: lciria@ugr.es and tb419@cam.ac.uk

ABSTRACT
Humans are uniquely capable of adapting to highly changing environments by updating relevant information and adjusting ongoing behaviour accordingly. Here we show how this ability —termed cognitive flexibility— is differentially modulated by high and low arousal fluctuations. We implemented a probabilistic reversal learning paradigm in healthy participants as they transitioned towards sleep or physical extenuation. The results revealed, in line with our pre-registered hypotheses, that low arousal leads to diminished behavioural performance through increased decision volatility, while performance decline under high arousal was attributed to increased perseverative behaviour. These findings provide evidence for distinct patterns of maladaptive decision-making on each side of the arousal inverted u-shaped curve, differentially affecting participants' ability to generate stable evidence-based strategies, and introduces wake-sleep and physical exercise transitions as complementary experimental models for investigating neural and cognitive dynamics.

INTRODUCTION

Making mistakes is inherent to learning and the accomplishment of any task. We make mistakes every day, even when faced with the same task repeatedly. Our ability to learn from these errors and flexibly adapt ongoing behaviour according to changes in the environment is critical for our survival. This ability —termed cognitive flexibility— depends on our innate capacity to establish associations between stimuli (S), responses (R), and outcomes (O), as well as to integrate previously acquired knowledge and skills into effective strategies for coping with similar future demands.[1] Here, we implement a Probabilistic Reversal Learning (PRL) task to study the modulatory effect of low and high arousal on cognitive flexibility — participants continue to perform as they fall asleep or with increasing physical exercise— to map either side of the Yerkes-Dodson Curve (1908).[2]

Cognitive flexibility is often studied using PRL tasks, typically assigning probabilistic reinforcement contingencies to abstract S-R associations, that are later abruptly reversed, requiring participants to learn new S-R reinforcement contingencies by trial and error to overcome prepotent ones[3]. Efficient performance relies on learning from the reinforcement received[4], the estimation of the likelihood that a reversal may occur,[5,6] and the continuous integration of a history of choices and reinforcements.[7] Indeed, evidence from both human and animal studies suggests that different high- and low-order strategies or series of rules are adopted during reversal learning, leading to maladaptive response patterns when the external pressures change or when the internal milieu varies.[7,8] Parsing the microstructure of learning derived from trial-by-trial responses enables the dissociation of the cognitive processes and behavioural strategies that drive subjects' choices during reversal learning. Here we propose that arousal fluctuations may differentially modulate cognitive flexibility leading to distinct maladaptive behavioural patterns of performance.[9]

Fluctuations in arousal and alertness (hereafter described jointly as "arousal") occur constantly across the day but are exacerbated during transitions toward strained states such as sleep[10] or physical extenuation,[11] where arousal levels change drastically in a progressive and nonlinear manner.[12,13] These arousal fluctuations play a crucial role in modulating cognition, facilitating or hindering certain cognitive processes and performance to internal and external stimuli. [14,15,16,17,18]

The interaction between arousal and cognition has been traditionally approached from the perspective proposed by Yerkes and Dodson in 1908.[2] According to their famous inverted U-shaped law, the optimal level of cognitive performance in complex tasks is reached at moderate levels of arousal, whereas deviations from this optimal arousal point, below or beyond, result in cognitive performance impairments. Though reductionist, the inverted U-shaped law represents a useful minimal framework to characterize the neural and cognitive dynamics of many physiological states across the arousal spectrum. Among these physiological states, researchers have paid special attention to reduced arousal states, including sleep stages,[19] sedation,[20] sleep deprivation,[21] motivation[22] and fatigue.[23]

Sleep can be used as the gold standard model of transition toward low arousal.[10] This area looking at the interaction between homeostasis and cognitive function is understudied due to the complexity of capturing dynamically metastable states like mild sedation[24,25] and drowsiness.[17] When falling asleep, individuals manifest a wide range of changes, from physiological to phenomenological, that are categorized into several well-described sleep

stages.[26] One of these stages is drowsiness, a transitional stage of consciousness between
attentive wakefulness and light sleep, characterized by a progressive and sometimes nonlinear
loss of responsiveness to external stimuli which does not immediately imply
unconsciousness.[10,27,28,29] Drowsiness, as well as similar reduced arousal states, has been
repeatedly associated with an impairment of cognitive processing, and particularly the capacity
to deal with conflicting information,[18] attentional performance,[30] and perceptual decision-
making.[31] However, in drowsiness, and even during highly reduced arousal states, pre-attentive
and early bottom-up attentive processing can still be accomplished with and without conscious
awareness. [17,32,33]

The transition towards the other side of the arousal spectrum (i.e., heightened arousal
states) has received even less attention.[34] The absence of a theoretical model for progressive
physiological transitions towards high arousal states, has also contributed to a lack of advance
in the field. Here, we consider endurance physical exercise as a useful experimental model of
arousal transition upwards, with many commonalities with sleep transition. A single bout of
endurance physical exercise (e.g., running or cycling) up to physical extenuation involves a
complex transition encompassing a wide range of changes (e.g., neural, motor, endocrinal,
phenomenological, etc.), that are also categorized into several well-described stages, from
resting, through the aerobic and the anaerobic thresholds, up to the limit where the individual
has to stop.[35] This highly fluctuating transition has been also associated with changes in
cognitive processing to internal and external stimuli.[36,37,38] In particular, high-order top-down
processes that govern goal-directed behaviour in changing environments (i.e., cognitive
control) appear to benefit from increases in the level of arousal[39] up to a certain exercise
intensity. Further intensity increments approaching and exceeding the anaerobic threshold
seem to hinder cognitive performance,[36,37,38,40] in line with the Yerkes-Dodson law prediction.

Sleep and physical exercise provide complementary perspectives on the cognitive
dynamics, and experimental models, when the arousal level is altered. However, and despite
the fact that both sides of the arousal spectrum exhibit similar cognitive performance
impairments, they cannot be treated as mirroring states in terms of cognitive performance
without a fine-grained differentiation of the behavioural dynamics that lead to these global
impairments. Furthermore, the theoretical differences in the transitions towards sleep or
complete (physical) exhaustion have to be considered in the assumptions and interpretations of
this and future studies. Thus, it is crucial to ask when arousal is altered (increased or decreased),
which specific processes of cognitive flexibility and information processing are affected, and
whether low and high arousal states are characterized by different strategic behaviours
underlying decision-making. Here, we use a PRL task to disentangle the behavioural dynamics
of cognitive flexibility as they get modulated by ongoing fluctuations in arousal levels and to
further delineate the microstructure of learning derived from trial-by-trial responses to
conflicting evidence. In particular, we manipulated arousal level to facilitate natural transitions
to low alertness, from awake to asleep; or to elicit high arousal, instructing participants to
exercise during 60 minutes at the highest intensity and effort possible without reaching
premature exhaustion. During both arousal modulations, participants performed a PRL task,
requiring the adaptation of behaviour following changes in reinforcement and punishment, as
well as the maintenance of strategic response patterns in the face of misleading (probabilistic)
feedback.

Based on the premises that (1) drowsiness hinders the extraction of task-relevant
information from external stimuli and its integration, fragmenting specific aspects of cognition
while preserving crucial executive control processes;[18,31,33,41] (2) drowsiness has been
associated with more liberal decision-making;[17,30,31] (3) moderate-to-high intensity endurance
exercise leads to a selective enhancement of executive control processes while lower and higher
intensities result in an impairment or minimal effect;[40,42,43] and (4) high arousal promotes
habitual responding and reduced engagement of complex cognitive strategies;[44,45,46] we
predicted that behavioural performance would be enhanced in moderate-intensity physical
exercise, while drowsiness and high-intensity exercise would lead to diminished performance
in light of the inverted U-shaped Yerkes-Dodson Law. Specifically, we hypothesized that
reduced arousal states would be associated with an impairment of performance (compared to
baseline), which would be attributed to a tendency to apply a simple strategy (win-stay/lose-
shift) instead of using an integrated history of choices and outcomes to drive performance
(probabilistic switching behaviour). In contrast, while we also expected an impairment of
performance during heightened arousal states, we hypothesized it would be attributed to a
failure to disengage from ongoing behaviour (perseveration). In addition, we hypothesized that
altered arousal states might reduce the ability of participants to apply a proper higher order
strategy, resulting in wide periods of time-on-task in which participants would perform the task
simply responding to the tones (i.e., automatic rule) but without applying any strategy (i.e.,
higher order rule).[9]
141
RESULTS
To investigate the modulatory effect of arousal fluctuations on cognitive flexibility, a PRL task
was carried out with human participants (n=100) while they were transitioning toward deep
sleep or physical extenuation. Participants were instructed to associate an auditory stimulus (S)
—high pitch sound or low pitch sound— with a response (R) button —left or right. In this
auditory version of the PRL task, each S-R association leads to an auditory outcome (O) —
correct (ding sound) or incorrect (white noise)— which participants use to assess their choice,
and apply this knowledge to guide the next choices. Therefore, the task entails the use of, at
least, two rules to success, as participants have to press a button after each auditory stimulus
(i.e., automatic rule) and to use an integrated history of S-R-O associations to determine the
correct S-R association (i.e., high order rule). Once participants reach 90% accuracy or greater
on the latest 10 trials, the implicit abstract S-R association is reversed, and participants have to
infer the new association from the feedback received. The number of responses needed to attain
a reversal (RAR) of the abstract association is used as the main index of performance. We
hypothesized[9] that reduced arousal states would lead to reductions in behavioural performance
compared to baseline arousal state; while heightened arousal states would lead to improved
performance relative to baseline, but only to an optimal point (i.e., moderate arousal) after
which the performance will be deteriorated with further increases in arousal level (see figure
1A). These hypotheses were formulated in line with the famous psychology inverted u-shaped
law originally attributed to Yerkes and Dodson (1908)[2] relating arousal modulation
performance in complex tasks, but later more formally defined by Broadhurst (1958)[47] and
Brown (1961).[48]

Note that, as a probabilistic task, the feedback provided is not always truthful nor
reliable and misleads the participant 20% of the time (see figure 1B). Thus, the participant
could correctly apply the S-R association and press the correct button in response to the
auditory stimulus, and still receive negative feedback, thus indicating an incorrect choice. This
scenario of conflicting evidence can lead participants to two different maladaptive response
patterns (see figure 1C) while performing the task: 1) switching the pattern choice across trials
with little (i.e., one negative feedback against the choice) or no evidence (i.e., no feedback
against the choice) of an actual rule change (probabilistic switching); or 2) sticking with the
previous choice despite having strong evidence (i.e., two or more negative feedbacks against
the choice) of an actual rule change (perseveration). Relying on these response patterns lead to
poor performance,[7] as the optimal strategy in this task is to stick with the previous choice with
zero or one negative feedback against the choice, and to switch the pattern choice if two or
more consecutive negative feedbacks against the choice happen.

177
Figure 1. Experimental design and arousal level classification: A) Schematic representation of the
experimental design and main hypotheses. Arousal level was endogenously manipulated by facilitating the natural
transition of participants from awake to sleep, or instructing them to exercise during 60' at the highest intensity
and effort they could maintain without reaching premature extenuation. A probabilistic reversal learning task was
assessed continuously during the arousal modulation. Optimal performance of the task was expected at moderate
arousal state (exercising at moderate intensity), while lower (drowsiness) and higher (exercising at high-intensity)
arousal state were expected to result in task performance deterioration. B) In this auditory version of the
probabilistic reversal learning paradigm, an auditory stimulus was presented on each trial, and participants had to

associate the sound with a response button, left or right. After that, auditory feedback was provided according to
the ongoing implicit rule. Notice that the feedback provided was not always truthful nor reliable, and attempted
to mislead the participant 20% of the time. C) Task trials were grouped into sequences of trials following a
particular rule (trend) where a particular sound was implicitly associated with a response button (e.g., high pitch
sound with the left button, and low pitch sound with the right button). Participants were instructed to infer the rule
from the provided feedback to assess their previous choice and apply the knowledge of their accuracy to guide the
next choices, knowing that the rule might change after a certain time. Based on the feedback received, participants
could make probabilistic or perseverative errors in the following trials. D) Automatic classification of arousal
during a drowsy session (representative participant). The pink line depicts changes in the theta:alpha ratio
(occipital electrodes cluster) during the pre-trial period (2 seconds before the auditory stimulus onset). The
horizontal bars on top represent trials classified as baseline (grey) or low arousal (blue). The variability in the
reaction times (green circles) closely follows the changes in theta:alpha ratio. Notice that circles on the horizontal
axis (reaction time equal to zero) were non-responsive trials, usually during low arousal (drowsy) periods but also
observed during exercise periods. E) Automatic classification of arousal during a physical exercise session
(representative participant). The red line depicts changes in the heart rate during the pre-trial period (2 seconds
before sound onset), and the horizontal bars on top represent trials classified as baseline (grey), moderate (yellow)
or high arousal (red). Similar to the low arousal session, the reaction times (green circles) fluctuates with the
changes in heart rate.

Arousal modulates probabilistic information during a stream of conflicting evidence.

First, we calculate the average RAR per participant in each arousal state (low, baseline sitting, baseline cycling, moderate, high). To account for the dependencies potentially generated by any procedural differences between Experiments, we fitted RAR using hierarchical linear mixed-effects modelling, with arousal as fixed effect, and participant nested into Experiment as random effects. The model showed a strong effect of arousal on RAR, $F_{(3,113.02)} = 11.59$, $p < 0.001$, $\beta = 0.61$. indicating that the processing of probabilistic information that allows the detection of changing patterns in a stream of conflicting evidence was modulated by the arousal level. Next, we checked for non-linearity in the relationship between arousal and RAR, to test the famous u-shaped curve. As expected, we found that the quadratic ($AIC = 1243.6$; $BIC = 1262.3$) outperformed linear fitting ($AIC = 1264.8$; $BIC = 1280.4$), confirming a possible curvilinear pattern (U shaped) of the effect of arousal on RAR (see figure 2), with a reliable increase in the number of responses required by the participants to complete a trend reversal (i.e., decrease of performance) as the level of arousal progress towards the extremes of the defined arousal range, confirming, for reversal learning, convergence with the Yerkes-Dodson law, later reformulated by Broadhurst in 1958.[47]

Splitting the comparisons to its specific baselines per arousal condition (i.e., sitting baseline compared to low arousal in the drowsiness condition; cycling baseline compared to moderate and high arousal in the exercise condition) yielded a reliable increase of RAR in low arousal, $t_{(124.62)} = 5.67$, $p < 0.001$, $\beta = 1.02$, and high arousal state, $t_{(117.93)} = 2.57$, $p = 0.011$, $\beta = 0.45$, compared with their corresponding baselines. Notably, baseline performance did not differ across arousal conditions (see supplementary figure 1). Contrary to what we expected, moderate arousal state was not associated with a decrease of RAR (the expected peak in performance), relative to baseline ($t_{(114.85)} = 1.61$, $p = 0.11$, $\beta = 0.25$,). Moreover, that we did not find evidence for a potential dual-task confounding effect in the heightened arousal conditions (see supplementary material). In sum, these findings provide evidence for an impairment in the processing of probabilistic information when the arousal level is altered, regardless of the side of the arousal spectrum.

233
Figure 2. Number of responses needed to attain a trend reversal as a function of the arousal state. A) Violins
and overlaid box plots of mean responses to reverse across arousal states. In box plots, middle black mark indicates
the median, and bottom and top edges indicate 25th and 75th percentiles, respectively. The upper and lower
whiskers indicate the maximum value of the variable located within a distance of 1.5 times the interquartile range
above the 75th percentile and below the corresponding distance to the 25th percentile value. Surrounding the
boxes (shaded area) is a rotated kernel density plot, which is comparable to a histogram with infinitely small bin
sizes. Jittered dots represent the averaged response to reverse score for each participant in each arousal state.
Linear mixed-effects model analysis revealed a reliable quadratic fitting between arousal and task performance,
outlined by the dashed line. Low and high arousal states were associated with a worse task performance relative
to their own baseline arousal states. Moderate arousal state was not associated with the expected optimal
performance as no differences were found with the baseline arousal state. B) Baseline differences of each
participant across altered arousal states are represented by the bars (grey bars indicate that these participants
needed more trials to attain a trend reversal in the baseline compared with the altered arousal states; blue, yellow
and red bars depict that these participants needed more trials to attain a trend reversal when arousal level was
altered -increased or decreased- compared with baseline arousal state). Participants are sorted by performance
difference between baseline and the arousal state. Upper and bottom panels show a consistent impairment of task
performance across participants in low and high arousal states. Non-reliable differences were found between
moderate and baseline arousal.
252

Different underlying mechanisms explain decreased performance in low and high arousal
states
In the analysis above, performance under high and low arousal states was compared
irrespective of the strategy participants may have used to solve the task. To test for the
hypotheses of the differential mechanism driving changes in performance for each arousal side
of the u-shaped curve, we calculated: a) probabilistic switching, as the proportion of trials when
the participants change the pattern choice with little or no evidence (i.e., zero or one negative
feedback against the choice); and b) perseveration, the likelihood of sticking with the previous
choice despite strong evidence (i.e., receiving two or more negative feedbacks in a row) that
the pattern has changed. We hypothesized that the impairment of performance in low arousal

would be primarily attributed to an increase in probabilistic switching, relative to the baseline
arousal state; and in contrast, the observed impairment of performance in high arousal state
will be primarily due to an increase in perseverative behaviour. To test these hypotheses, we
fitted probabilistic switching and perseveration (separately for low and high arousal states)
using the hierarchical linear mixed-effects model structure defined previously. The analyses
revealed that, while the probabilistic switching increased consistently across subjects during
low arousal state compared with baseline arousal, $F(1,56) = 12.65$, $p < 0.001$, $\beta = 1.02$, no
reliable differences were observed in perseveration between these arousal states ($F < 1$). On the
other hand, high arousal states led to a reliable increase in perseverative behaviour compared
to the baseline state, $F(1,68) = 4.80$, $p = 0.031$, $\beta = 0.22$, with no reliable differences observed
in probabilistic switching ($F < 1$). These results suggest that altered arousal states lead to
distinct maladaptive decision-making patterns that affect participants' ability to generate stable
evidence-based strategies, although evidence-driven responses were present (see figure 3A).
To further prove that the impairment in performance in low and high arousal states
could be attributed to the different maladaptive behavioural patterns, we carried on a mediation
analysis separately for each arousal state (low, high). We first confirmed that probabilistic
switching and perseveration have an effect on the RAR, while controlling for the arousal state
(see figure 3B). These results, together with the previous analyses where we found an effect of
arousal state on probabilistic switching and perseveration, revealed a full mediation between
these variables. As figure 3B illustrates, the regression coefficient between arousal and RAR,
and the regression coefficient between probabilistic switching and RAR were statistically
reliable, showing a full mediation of probabilistic switching on the effect of low arousal on
RAR. The bootstrapped standardized indirect effect of low arousal on RAR, mediated by
probabilistic switching, was 0.65 ($p < 0.001$), and the 95% confidence interval ranged from
0.29 to 1.07. A similar fully mediation effect was observed in high arousal state, showing that
the effect of high arousal on behavioural performance was fully mediated via the perseverative
behaviour. The bootstrapped standardized indirect effect was 0.10 ($p = 0.014$), and the 95%
confidence interval ranged from 0.14 to 0.24. As predicted, participants showed an impairment
of performance during low arousal state, relative to baseline arousal, which was primarily
attributed to an increase of probabilistic switching (i.e., changing pattern choice with little or
no evidence of an actual rule change). In contrast, while participants also showed an
impairment of performance during high arousal state, relative to the baseline arousal, it was not
attributed to an increase in probabilistic switching, but to an increase in perseverative behaviour
(i.e., sticking with the previous choice despite consecutive negative feedbacks).
297

Figure 3. Maladaptive behavioural patterns across participants in low and high arousal states. A) Violins and overlaid box plots of the percentage of change from baseline to low (blue) and high (red) arousal states in probabilistic switching and perseveration. In box plots, middle black mark indicates the median, and bottom and top edges indicate 25th and 75th percentiles, respectively. The upper and lower whiskers indicate the maximum value of the variable located within a distance of 1.5 times the interquartile range above the 75th percentile and below the corresponding distance to the 25th percentile value. Surrounding the boxes (shaded area) is a rotated kernel density plot, which is comparable to a histogram with infinitely small bin sizes. Jittered dots represent the averaged response to reverse score for each participant in each arousal state. B) Mediation model diagram to illustrate that the general impairment in task performance found in low and high arousal states was mediated by different maladaptive behavioural patterns. Dashed lines (indirect effects) represent the effect of low (blue) and high (red) arousal on task performance (indexed by the averaged responses to attain a trend reversal) through probabilistic switching and perseveration, respectively. Solid lines depict direct effects between variables. Grey lines represent the absence of a direct effect of low arousal on perseveration and high arousal on probabilistic switching. Notice that a direct effect of an independent variable (arousal) onto the mediator (probabilistic switching, perseveration) is a prerequisite for mediation being possible. Standardized β regression coefficients are indicated in each effect (* depicts $p < 0.05$). Accordingly, the values of all effects are expressed as the number of standard deviations from the mean. For example, the direct effect of high arousal on RAR (β = 0.22) implies that a standard deviation change of 1 in the arousal variable would result in a standard deviation increase of 0.22 in RAR.

Arousal disrupt the reversal strategy

To maximise performance in the task, a good strategy is to not fall for the false feedback and stand your ground until the next feedback, as well as switch to the second consecutive feedback. The fact that participants sometimes needed an unreasonable high number of responses to attain a reversal in low and high arousal states suggests the existence of sections of time on task in which they responded to the tones but could not apply the strategy rules (see fig 4A). These sections without clear strategic behaviour, that we call breakdowns, have been often neglected in previous studies using PRL tasks as failures of compliances or "bad participant". The transient on/off nature of these breakdowns may provide valuable insight into the behavioural dynamics of participants in different states of arousal. We hypothesized that breakdowns sections would increase in low and high arousal states, relative to a baseline arousal state. First, we traced the sections of the task (more than 20 trials) in which participants did not attain a reversal. Second, we calculated the proportion of time these sections represented to the total time-on-task, and finally, we implemented a hierarchical linear mixed-effects model with the

structure defined in previous analyses, separately for each arousal state (low, high), with the number of breakdowns as the index of performance. As hypothesized, low and high arousal states lead to longer breakdown sections compared with baseline arousal state (t (127.99) = 3.40, $p < 0.001$, $\beta = 0.13$; t (121.69) = -2.97, $p = 0.003$, $\beta = 0.11$). Subject-by-subject results (fig 4C) show a consistent increase of breakdowns across participants in low arousal state. Although high arousal states also showed a reliable increase of breakdowns as a group, this effect was less systemic, with half of the participants showing the opposite effect, no difference or no breakdowns.

Figure 4. Behavioural strategy breaks as arousal changes. A) Automatic classification of a section of time where a representative participant responded without a clear behavioural strategy. The green circles show RTs and the blue line shows the ongoing accuracy of the task (10-points moving average). The grey shaded area flanked by the zigzagging vertical lines depicts the section of time classified as a breakdown. B) Violins and overlaid box plots of the averaged percentage of time-on-task without strategy across participants in low and high arousal states, compared with their respective baselines states. In box plots, the middle black mark indicates the median, and bottom and top edges indicate 25th and 75th percentiles, respectively. The upper and lower whiskers indicate the maximum value of the variable located within a distance of 1.5 times the interquartile range above the 75th percentile and below the corresponding distance to the 25th percentile value. Surrounding the boxes (shaded area) is a rotated kernel density plot, which is comparable to a histogram with infinitely small bin sizes. Jittered dots represent the averaged percentage of time-on-task without a strategy of each participant in each arousal state. Linear mixed-effects model analyses revealed that low and high arousal states lead to longer periods of breakdown relative to the baseline arousal state. Interestingly, violin plots show a considerable number of participants who had no breakdowns at baseline arousal states, something that completely disappears in low arousal state (all participants had breakdowns), and that is reduced in high arousal state. C) Baseline differences of each participant in low and high arousal states represented by horizontal bars (grey bars indicate that these participants spent more time performing the task without a particular strategy in the baseline arousal state compared with the altered

arousal states; blue and red bars depict that these participants were applying behavioural strategies less time when arousal level was altered (increased or decreased) than in baseline arousal state. Participants are sorted by performance difference between baseline and the arousal state. Both panels show a consistent impairment of task performance across participants in low and high arousal states.

DISCUSSION

In the present study, we facilitated natural transition of healthy participants towards the borders of non-pharmacological arousal states (drowsiness, physical exertion) to investigate the behavioural dynamics of cognitive flexibility. In line with our pre-registered hypotheses,[9] the findings revealed a quadratic-like pattern (inverted U-shape) of the effect of arousal fluctuations on cognitive performance. As the level of arousal progressed towards the extremes of the defined arousal range reversal learning performance decreased, in agreement with the predictions of the Yerkes-Dodson law (1908).[2] Although cognitive flexibility diminished in both under high and low arousal states, different maladaptive behavioural patterns drove this performance impairment. As predicted, the performance decline exhibited by our participants under drowsy states was primarily attributed to a more decision volatility (i.e., shifting pattern choice with little or no evidence of reinforcement contingencies change). In contrast, participants also showed a decline in performance during high arousal state but attributed to increased perseverative behaviour (i.e., sticking with a particular pattern choice despite having strong evidence that the contingencies have changed). Our findings also revealed that most participants undergo prolonged periods of time-on-task in which they seem unable to apply any specific higher order strategy. These breakdown periods, which can last for several minutes, are more frequent and sustained during high or low arousal. In short, our results provide solid evidence for distinct maladaptive decision-making patterns under altered arousal states, differentially affecting the participants' ability to generate stable evidence-based strategies.

Arousal fluctuations thus seem to elicit a distinctive behavioural distortion of cognitive flexibility as further indicated by the microstructure of learning derived from trial-by-trial responses to negative feedback. Healthy participants under high arousal exhibited normal acquisition of S-R reinforcement contingencies but perseverative response patterns when contingencies were reversed. This failure to disengage from ongoing behaviour is a translational phenomenon strongly linked to impulsivity and compulsivity,[49] and prevalent in numerous neuropsychiatric and medical conditions.[7,50,51] For instance, patients with lesions that include ventral prefrontal cortex and orbitofrontal cortex,[52] as well as chronic cocaine users[53] and patients with schizophrenia,[54] show normal acquisition of S-R contingencies but are severely impaired when those S-R reinforcement contingencies are abruptly reversed, exhibiting perseverative responding to the previously reinforced S-R contingency. Altogether, these findings suggest that high arousal undermines healthy individuals' capacity to engage in complex cognitive strategies driving them to rely on habitual response patterns, which, paradoxically, might also enhance behavioural control in terms of response inhibition.[46] Our findings not only further the understanding of the processes underlying automatized behaviour and habitual response tendencies, but high arousal may be used as a model to inform both impulsive and compulsive aspects of psychopathology.

In contrast, healthy participants under low arousal seemed unable to maintain the learned S-R reinforcement contingency and started to deviate from the evidence, revealing a

volatile pattern of behaviour. Since a crucial aspect of the PRL experimental design was the existence of a 20% of misleading feedback, to maximise performance, individuals should not fall for the false feedback and —ideally— stand their ground until the next feedback. Further and as part of a successful strategy, they should switch if two or more consecutive feedbacks are given against the previously reinforced choice pattern. Consequently, adaptive behaviour during the task requires a balance between both types of behaviour (stability and flexibility). Those participants under low arousal fell repeatedly for the misleading feedback, switching prematurely after negative feedback. Furthermore, they showed increased decision volatility by spontaneously switching even without any negative feedback. This volatile pattern of cognitive flexibility has been linked to serotonin[55] and dopamine systems,[56] and is observed in patients with major depression,[57,58,59] often linked to either an oversensitivity to punishment or an impaired control over negative feedback.[60,61] It is reasonable to speculate that low arousal levels render individuals more sensitive in updating S-R reinforcement contingencies, rather than increase sensitivity to punishment as in major depression. Moreover, low arousal may increase volatility by decreasing attentional resources, leading to spontaneous explorations, higher RT variability and periodic omissions (see supplementary figure 2).

The fragmentation of cognitive control due to changes in arousal has been primarily shown in sleep deprivation[62,63,64,65,66] and not in spontaneous fluctuations of alertness as we show in this study. The increased volatility in the PRL with low arousal suggests a decrease in cognitive control that is different from an increase in perseverative behaviour seen in high arousal. Indeed, we have previously shown that decreased levels of arousal can fragment or reconfigure specific aspects of cognition while preserving crucial executive control processes such as the capacity to detect and react to incongruity,[18] the efficiency in perceptual decision making,[31] and the precision of conscious access.[17] Here, we add further evidence showing that individuals under reduced arousal state, although struggling to maintain stable evidence-based decision-making patterns, are able to learn new S-R reinforcement contingencies, demonstrating flexibility of the human brain to adapt to increasing levels of endogenous (arousal) noise. The evidence of cognitive and —indirectly— neural reconfiguration of cognitive control networks suggests compensatory mechanisms elicited by the change in arousal.

Upon further examining the microstructure of learning derived from trial-by-trial performance of the PRL task, we uncovered the existence of prolonged periods of time-on-task in which participants did not seem to apply any particular high-order behavioural strategy. Although these breakdown periods emerged regardless of the arousal level, they were prevalent under low and high arousal states, lasting from few to several minutes. Remarkably, the transient on/off nature of these breakdowns suggests that extreme arousal levels alternate between different metastable cognitive states. The first state can be defined by a relatively successful application of the reinforcement information where participants can navigate the uncertainty of the PRL, while in the other metastable state they seem to only apply the simple auditory-motor S-R rule to respond to the auditory tones but are unable to use choice history to develop a successful strategy.

In the context of this study, arousal as a biological construct defined by the homeostatic regulatory capacity of the system and its responsiveness,[67] helps to link drowsiness and increased alertness during physical exercise in a common framework where the predictions of

the Yerkes-Dodson inverted U-shaped law can be experimentally tested. Despite the obvious
difference at the biological, neural and psychological level between both sides of the curve, the
common decrease in performance highlights the commonalities between the extremes in human
performance, adding to the fact that both —sleep and physical exertion— emerge as natural
transitions from a similar state (resting) traversing different stages, and exhibit nonlinear
dynamics and hysteresis processes in their transitions.[12] Thus, drowsiness and physical exertion
provide complementary perspectives on cognitive dynamics when the arousal level is altered.
The present findings point out their differences in the cognitive fragmentation leading to a
general decline in task performance.

Though the Yerkes-Dodson law was not initially formulated to be a general rule to
apply to all psychology subfields (learning, motivation, emotion, etc.), through the years, and
with the pressure to find common mechanisms in psychology, the findings initially defined for
learning were further extended and reinterpreted as a law about the relationship between
arousal and other physiological constructs to perceptual and cognitive performance.[68] Despite
this overgeneralization from its genuine formulation and its reductionist nature, our findings
rely on such inverted U-shaped law as a basic useful theoretical framework, providing an
attractive theoretical model to characterize the neural, cognitive and behavioural dynamics
involved in the impact of arousal fluctuations in a wide range of physiological states and
neuropsychiatric conditions.

Our findings bring some generalizations about the need to extend the traditional
framework of understanding the interplay between cognitive dynamics and arousal through the
prism of the homeostatic steady-state dynamics using pharmacological interventions[34] or
transient alterations of emotional state.[69] In addition to this classical approach, we believe that
drowsiness and physical exertion provide fruitful -naturally occurring- alterations of the arousal
level with a preserved capacity to behaviourally respond, which can be utilized to study the
modulation of neural function and cognitive processing. In the traditional steady-state
approach, such natural fluctuations of the arousal level may be undetected,[70] hindering or
distorting cognitive and neural markers of crucial aspects of information processing.[17]
Pharmacological and lesion perturbations of the brain are regarded as causal in cognitive
neuroscience and regarded as stronger in their explanatory power than conditions relying on
stimuli or psychological modulations. Arousal is an internally modulated change that can be
used to study cognition and may be regarded in the strong causality range due to its partial
independence from psychological processes.[18] The cases of drowsiness and physical exertion
as causal models to study the neural mechanism of cognitive flexibility may prove to be very
useful in the exploration of how cognition is fragmented or remain resilient under (reversible)
perturbations of arousal[17,33,71] Our findings highlight that further research should focus on the
rapidly changing dynamics of brain function and cognitive processing that appear to capture
key dynamics relevant to our behavioural and perhaps even phenomenological experience, as
we drift into strained physiological states.

METHODS
Participants
A total sample of 100 participants of an age range between 18 and 40 years old was included
in the present study. All participants reported normal binaural hearing, no visual impairment

and no history of cardiovascular, neurological or psychiatric disease. They were asked to get a
normal night rest on the day previous to testing, and not to consume stimulants like coffee or
tea on the day of the experiment.

The first experiment (herein Experiment#1) consisted of 35 participants (15 female; age
range 18-40). In addition to the general aforementioned inclusion criteria, only easy sleepers,
as assessed by the Epworth Sleepiness Scale (ESS),[72] were selected to increase the probability
that participants fell asleep. Recruited participants were considered healthy with relatively high
ESS scores but not corresponding to a condition of pathological sleep such as hypersomnia
(i.e., scores 7–14). They were recruited via the Cambridge Psychology SONA system.

The second experiment (herein Experiment#2) consisted of 15 participants (11 female;
age range 18-40). Inclusion criteria and recruitment processes were similar to Experiment#1.

The third experiment (herein Experiment#3) consisted of 50 participants (6 female; age
range 19-39). Additionally to the common inclusion criteria, only individuals who reported at
least 8 hours of cycling or triathlon per week were selected. Well-trained cyclists were selected
because they are used to maintaining the pedalling cadence at high intensity during long periods
of time. Furthermore, they are able to keep a fixed posture over time, which notably reduces
movement artefacts. They were recruited from the University of Granada (Spain) through
announcements on billboards and previous databases.

All participants from the three experiments gave written informed consent to participate
in the study and received a remuneration of 10€ per hour (i.e., approximately 30€ per
participant). The Cambridge Psychology Ethics Committee and the University of Granada
Ethics Committee approved the study (CPREC 2014.25; 287/CEIH/2017).

Experimental task
A modified version of the probabilistic reversal learning paradigm was used in all three
experiments, which was characterized by employing auditory stimuli and an abstract rule (see
figure 1B-C). In this task, participants learnt to choose one of two randomly presented tones
by receiving instructive auditory feedback tones after each response, indicating either a correct
or incorrect choice. When participants reached a 90% accuracy in the last 10 trials,
reinforcement/punishment contingencies were reversed so that the previously reinforced tone
was punished and vice versa. Within each reversal trend, a 20% probabilistic error trial was
included in which "wrong" feedback was given for correct choices, even though the
reinforcement contingencies had not changed. Participants were instructed to infer the rule
from the feedback received, knowing that sometimes it might be misleading and that the rule
might change after a certain time. The stimuli were binaurally presented at a random time
interval (between 1000 and 1500 ms) during 500 ms. They had to respond to both targets by
pressing a button with their right or left hand.

Procedure
In Experiment#1, participants were fitted with an EGI electrolyte 129-channel cap (Electrical
Geodesics, Inc. systems) after receiving the task instructions and subsequently signing the
informed consent. The whole session was completed in a comfortable adjustable chair with
closed eyes. Task instructions were to respond as fast and accurately as possible, reducing body
movements as possible and keeping the eyes closed. In the beginning, the back of the chair was

set up straight and the lights in the room were on. Participants were asked to remain awake with their eyes closed whilst performing the first block (awake block) of the task which consisted of 480 trials, lasting 30 min approximately. Then, the chair was reclined to a comfortable position, the lights were turned off and participants were offered a pillow and a blanket. They were explicitly told that they were allowed to fall asleep during this part of the task and that the experimenter would wake them up by making a sound (i.e. knocking on the wall) if they missed 5 consecutive trials. This block (drowsy block) also consisted of 480 trials. Then, the sequence of two blocks (awake-drowsy) was repeated. In total, participants completed 1920 trials divided into 4 blocks of 480 trials each one. The whole session lasted for 3 hours approximately.

In Experiment#2, the procedure was similar to the Experiment#1 except for the time to fall asleep that was increased to get a higher amount of low-arousal (i.e., drowsy) trials. Participants completed a total of 2120 trials, divided into 4 blocks. The order of the blocks was the same for all participants and followed the same sequence as in Experiment#1: awake-drowsy-awake-drowsy. Awake blocks had 100 trials each one, while drowsy blocks consisted of 960 trials each one. The session lasted for 3 hours approximately.

In Experiment#3, upon arrival to the laboratory, participants were seated in front of a computer in a dimly illuminated, sound-attenuated room with a Faraday cage. They received verbal and written instruction about the experiment and were prepared for electrophysiological measurement. They were fitted with a 64-channel high-density actiCHamp EEG system (Brain Products GmbH, Munich, Germany) and a Polar RS800CX heart rate (HR) monitor (Polar Electro Öy, Kempele, Finland). Notice that EEG data was acquired but was not used to test the hypotheses of this study, and will be reported elsewhere. The whole session consisted of 4 different blocks. The first one was an adaptation (non-exercise) block in which participants performed 100 trials while resting in a comfortable chair. Then, they got on a cycle-ergometer and completed 100 trials while warming-up at light intensity. Subsequently, they completed a self-paced 60' time-trial (i.e., high-intensity exercise) while performing the task, resulting in 850 trials approximately (the number of trials slightly varied as a function of the reaction time of participants). In line with previous experiments from our laboratory (Holgado et al., 2018; Holgado et al., 2019),[73,74,75] in the self-paced time-trial participants were instructed to achieve the highest average power (watts) during the 60' time-trial exercise, and were allowed to modify the power load during the exercise. They were encouraged to self-regulate effort in order to optimize physical performance without reaching premature exhaustion. That self-regulation yielded fluctuations of effort during the 60' exercise period, which allowed us to study the effect of arousal on the management of probabilistic information. Once the 60' time-trial block was finished, participants completed the last block while cooling down at light intensity, which was also composed of 100 trials. All participants completed the blocks in the same order, lasting around 3 hours.

Arousal classification

In Experiment#1 and Experiment#2, the arousal level was endogenously manipulated by facilitating the natural transition from awake to sleep. This transition reduces arousal and yields a considerable proportion of drowsy yet responsive trials as seen in previous experiments from our laboratory.[17,30,71] This way, we were able to study the effect of arousal (i.e. baseline arousal

[awake] trials vs. low-arousal [drowsy] trials) on the management of probabilistic information.
Given that awake-sleep transition is characterized by a decreasing alpha range activity, together
with an increasing theta range activity (Hori et al., 1994), progression of drowsiness was
quantified by the spectral power of respective EEG frequency bands[i]. We computed the spectral
power of EEG frequency oscillations for each trial from -2000 ms to 0 ms in respect to the
onset of a target tone using continuous wavelet transform, set from 3 cycles at 3 Hz to 8 cycles
at 40 Hz. Theta (4-6 Hz) and alpha (10-12 Hz) power were then averaged individually for each
trial across central (E36, E104) and occipital (E75, E70, E83) electrodes for theta and alpha
rhythms respectively. Finally, theta/alpha ratio was computed and smoothed with a 4-point
moving average resulting in a single "sleepiness" value per trial. Visual inspection of
theta/alpha ratio and RT dynamics of each participant confirmed the presence of clear
sleepiness-related fluctuations during the experimental session, especially during drowsy
blocks. Those participants who did not show clear fluctuations of the theta:alpha ratio were
removed from final analyses (5 subjects). Then, each trial for each participant was initially
categorized as drowsy (top 33% of lower theta-upper alpha ratio scores) or alert (lowest 33%).
Further, following the sleep hysteresis physiology criteria[77] isolated awake trials within
prolonged periods of drowsy (≥10 trials) were considered as drowsy to account for the gradual
homeostatic change during the sleep transition. In addition, the first 100 trials of each block
(awake and drowsy) were considered as awake trials.

In Experiment#3, the arousal level was endogenously manipulated by facilitating the
natural transition from a resting state to high-intensity physical exercise. This transition
increases the arousal level progressively, with continuous fluctuations that affect cognitive
performance as seen in previous studies from our laboratory.[40,75,78,79] We captured these arousal
fluctuations at a single trial level (moderate arousal trials, high arousal trials) by using the HR
response. To address the intersubject variability, HR data were transformed into differential
scores relative to the HRmax estimated using the equation of Tanaka et al., (2001)[80], a reliable
and well-established method to calculate HRmax in healthy individuals. Then, moderate and
high arousal trials were characterized based on percentage relative to HRmax. HR between
60% and 80% of HRmax were considered as moderate arousal, while HR higher than 80%
HRmax were considered as high arousal. Due to technical issues with HR monitoring, 4
subjects were removed for further analyses.
611

612   **Behavioural data analysis**
613   In probabilistic reversal learning paradigms, participants are instructed to infer an abstract rule
614   form the feedback they receive, knowing that sometimes it might be misleading and that the
615   rule might change. Since a reversal is triggered when a high-level accuracy is reached, the
616   number of responses needed to attain a reversal is considered one of the main indices of

---

[i] Deviation from pre-registration. Originally, we aimed to use the automated offline method developed by Jagannathan and collaborators based on frequency and sleep grapho elements to detect EEG micro variations in alertness and characterize awake and drowsy trials.[76] However, our PRL task design, especially the pretrial duration, which was limited to 2 seconds, did not fit the task features recommended by Jagannathan and collaborators (e.g., 4 seconds pretrial duration) for a reliable characterization of awake and drowsy trials. So we decided to classify awake/drowsy trials based on theta:alpha ratio, as seen in previous experiments from our laboratory.[17,30,71]

performance. To delineate the microstructure of learning derived from trial-by-trial responses we considered the likelihood of switching the pattern choice across trials as a function of the amount of consecutive negative feedback received. The likelihood of switching was considered the main index of strategic behaviour, and was divided into 2 different strategies: *i*) Probabilistic switching: the proportion of trials when the participants change the pattern choice with little (one negative feedback against the choice) or no evidence (no feedback against the choice) of an actual rule change; *ii*) Perseveration: likelihood that participants stay with the seemingly incorrect choice even after receiving two or more negative feedbacks in a row).

The number of breakdown sections was also used as an index of performance. We defined a breakdown as a section of time in which participants 'lose' the task, and do not follow any strategy, being unable to reach a change of trend during more than 20 consecutive trials. RT, accuracy, and omissions were also checked as secondary indices of behavioural performance.

Participants with overall accuracy under 60% during baseline period or a reduced amount of reversals attained were excluded (i.e., 4 subjects from Experiment 1; 2 subjects from Experiment 2; 6 subjects from Experiment 3).

Statistics

Single-subject analysis

In order to test the hypotheses, we took a set of strategies. We first captured the direction of effects for each of the key performance variables (i.e., RAR, RT, accuracy, omissions, and switching likelihood), and contrasted them for each participant, obtaining an indication of the direction and strength of the effects per participant. Descriptive and distribution measures, as well as single-subject statistics, were used as guidance of the variability of effect size in single variables, and for guiding the previously defined exploratory hypotheses. Per participant, effect sizes were calculated and depicted for each of the key performance variables to check the effect size of individual differences across arousal states.[ii]

Group analysis

To investigate the management of probabilistic information as a function of arousal, we conducted mixed-effects analyses including data from the three experiments collapsed into a single dataset with RAR as the main index of performance. In face of the diversity of samples' characteristics and experiment features, we fit RAR using hierarchical linear mixed-effects modelling, as implemented in the lme4 R package.[81] We treated RAR as obeying to a hierarchical data structure with arousal as fixed effect, and participant (level 2) nested into experiment (level 1) as random effects. This random part was common to all models. We tested the specific hypothesis by using the same approach based on multilevel linear mixed-effects modelling. Different variables (i.e., probabilistic switching, perseveration, breakdowns, RT variability and omissions) were analysed in a multilevel data structure, with the fixed (arousal) and random effects (experiment/participant) adjusted to the specific hypothesis tested.

[ii] Deviation from pre-registration. Spearman rank-order correlation tests and Bayes factors were finally not performed to estimate the degree of association between switch likelihood as a function of consecutive negative feedbacks and arousal states. We will check the slope and effect.

Models were compared using the Akaike Information Criterion (AIC), and a likelihood ratio test. Notice that AIC does not assume that the true model is among the set of candidates (and is just intended to select the one that is closest to the true one). In our case, fitting decisions were not about the truthiness of models, but to include or not a given factor. For model comparisons performed to identify the best-fitting model, a relatively lenient 0.010 p-value criterion was adopted.

Causal mediation analyses were conducted to estimate the proportional direct and indirect effects of arousal on task performance through probabilistic switching and perseveration strategies (mediators) using the "mediation" package in R[iii].[82] This method allowed us to assess a confidence interval of the mediation effect itself using rigorous sampling techniques with fewer assumptions of the data. The average causal mediation effect was determined using a nonparametric bootstrapping method (bias-corrected and accelerated; 1000 iterations) and reported as standardized β regression coefficients for direct comparison with each other. Confidence intervals were obtained using a quasi-Bayesian approximation.

Pre-registration
https://osf.io/tzw6d

Data and code
Data and codes used for the analyses presented here are available at the OSF repository (https://osf.io/xk379/), with unrestricted access.

Funding
This research was supported by a University of Granada Postdoctoral Fellowship (2019/P7/100) and the Junta de Andalucía (DOC_00225) to L.F.C, a research grant from the Ministerio de Economía y Competitividad (PSI2016-75956-P and PID2019- 105635GB-I00) to D.S., and Wellcome Trust Biomedical Research Fellowship (WT093811MA) awarded to T.A.B.

REFERENCES
1.   Salehi, B., Cordero, M. I. & Sandi, C. Learning under stress: The inverted-U-shape function revisited. *Learn. Mem.* **17**, 522–530 (2010).
2.   Yerkes, R. M. & Dodson, J. D. The relation of strength of stimulus to rapidity of habit-formation. *J. Comp. Neurol. Psychol.* **18**, 459–482 (1908).
3.   Shiu, L. & Chan, T. Unlearning a stimulus–response association. *Psychol. Res.* **70**, 193–199 (2006).
4.   Stalnaker, T. A., Cooch, N. K. & Schoenbaum, G. What the orbitofrontal cortex does not do. *Nat. Neurosci.* **18**, 620 (2015).
5.   Costa, V. D., Tran, V. L., Turchi, J. & Averbeck, B. B. Reversal Learning and Dopamine: A Bayesian Perspective. *J. Neurosci.* **35**, 2407–2416 (2015).
6.   Jang, A. I. *et al.* The role of frontal cortical and medial-temporal lobe brain areas in learning a

---

iii Deviation from pre-registration. The mediation analysis was not initially included in the pre-registration, however, we decided to run it in order to test whether the impairment in performance in low and high arousal states could be attributed to the different maladaptive behavioural patterns.

Bayesian prior belief on reversals. *J. Neurosci.* **35**, 11751–11760 (2015).

7.   Izquierdo, A., Brigman, J. L., Radke, A. K., Rudebeck, P. H. & Holmes, A. The neural basis of
reversal learning: An updated perspective. *Neuroscience* **345**, 12–26 (2017).

8.   Murray, E. A. & Gaffan, D. Prospective memory in the formation of learning sets by rhesus
monkeys (Macaca mulatta). *J. Exp. Psychol. Anim. Behav. Process.* **32**, 87 (2006).

9.   Ciria, L. F. *et al.* Probabilistic trend detection in different levels of arousal. OSF (2019).

10.   Goupil, L. & Bekinschtein, T. A. Cognitive processing during the transition to sleep. 15.

11.   Schmit, C. & Brisswalter, J. Executive functioning during prolonged exercise: a fatigue-based
neurocognitive perspective. *Int. Rev. Sport Exerc. Psychol.* **13**, 21–39 (2020).

12.   Ogilvie, R. D. & Wilkinson, R. T. The detection of sleep onset: behavioral and physiological
convergence. *Psychophysiology* **21**, 510–520 (1984).

13.   Bernaola-Galván, P. A., Gómez-Extremera, M., Romance, A. R. & Carpena, P. Correlations in
magnitude series to assess nonlinearities: Application to multifractal models and heartbeat fluctuations.
Phys. Rev. E **96**, 032218 (2017).

14.   Wickens, C. D., Hutchins, S. D., Laux, L. & Sebok, A. The impact of sleep disruption on
complex cognitive tasks: a meta-analysis. *Hum. Factors* **57**, 930–946 (2015).

15.   McMorris, T. & Hale, B. J. Differential effects of differing intensities of acute exercise on speed
and accuracy of cognition: A meta-analytical investigation. *Brain Cogn.* **80**, 338–351 (2012).

16.   McMorris, T., Sproule, J., Turner, A. & Hale, B. J. Acute, intermediate intensity exercise, and
speed and accuracy in working memory tasks: a meta-analytical comparison of effects. *Physiol. Behav.*
102, 421–428 (2011).

17.   Noreika, V. *et al.* Alertness fluctuations when performing a task modulate cortical evoked
responses to transcranial magnetic stimulation. *NeuroImage* **223**, 117305 (2020).

18.   Canales-Johnson, A. *et al.* Decreased alertness reconfigures cognitive control networks. *J.
Neurosci.* **40**, 7142–7154 (2020).

19.   Nir, Y., Massimini, M., Boly, M. & Tononi, G. Sleep and consciousness. in *Neuroimaging of
consciousness* 133–182 (Springer, 2013).

20.   Cimenser, A. *et al.* Tracking brain states under general anesthesia by using global coherence
analysis. *Proc. Natl. Acad. Sci.* **108**, 8832–8837 (2011).

21.   Hudson, A. N., Van Dongen, H. P. & Honn, K. A. Sleep deprivation, vigilant attention, and
brain function: a review. *Neuropsychopharmacology* **45**, 21–30 (2020).

22.   Wise, R. A. Dopamine, learning and motivation. *Nat. Rev. Neurosci.* **5**, 483–494 (2004).

23.   Tanaka, M., Ishii, A. & Watanabe, Y. Neural correlates of central inhibition during physical
fatigue. *PloS One* **8**, e70949 (2013).

24.   Chennu, S., O'Connor, S., Adapa, R., Menon, D. K. & Bekinschtein, T. A. Brain connectivity
dissociates responsiveness from drug exposure during propofol-induced transitions of consciousness.
PLoS Comput. Biol. **12**, e1004669 (2016).

25.   Lee, M. *et al.* Network properties in transitions of consciousness during propofol-induced
sedation. *Sci. Rep.* **7**, 1–13 (2017).

26.   Pace-Schott, E. F. & Hobson, J. A. The neurobiology of sleep: genetics, cellular physiology
and subcortical networks. *Nat. Rev. Neurosci.* **3**, 591–605 (2002).

27.   Bareham, C. A., Bekinschtein, T. A., Scott, S. K. & Manly, T. Does left-handedness confer
resistance to spatial bias? *Sci. Rep.* **5**, (2015).

28.   Overgaard, M. & Overgaard, R. Measurements of consciousness in the vegetative state. *The
Lancet* **378**, 2052–2054 (2011).

29.   Sanders, R. D., Tononi, G., Laureys, S. & Sleigh, J. Unconsciousness, not equal to
unresponsiveness. *Anesthesiology* **116**, 946–959 (2013).

30.   Bareham, C. A., Manly, T., Pustovaya, O. V., Scott, S. K. & Bekinschtein, T. A. Losing the left

side of the world: rightward shift in human spatial attention with sleep onset. *Sci. Rep.* **4**, 1–5 (2014).

31. Jagannathan, S. R., Bareham, C. A. & Bekinschtein, T. A. *Decreasing arousal modulates perceptual decision-making.* http://biorxiv.org/lookup/doi/10.1101/2020.07.23.218727 (2020) doi:10.1101/2020.07.23.218727.

32. Chennu, S. & Bekinschtein, T. A. Arousal Modulates Auditory Attention and Awareness: Insights from Sleep, Sedation, and Disorders of Consciousness. *Front. Psychol.* **3**, (2012).

33. Kouider, S., Andrillon, T., Barbosa, L. S., Goupil, L. & Bekinschtein, T. A. Inducing Task-Relevant Responses to Speech in the Sleeping Brain. *Curr. Biol.* **24**, 2208–2214 (2014).

34. Berridge, C. W. & Arnsten, A. F. Psychostimulants and motivated behavior: arousal and cognition. *Neurosci. Biobehav. Rev.* **37**, 1976–1984 (2013).

35. Kenney, W. L., Wilmore, J. H. & Costill, D. L. *Physiology of sport and exercise.* (Human kinetics, 2018).

36. Verburgh, L., Konigs, M., Scherder, E. J. A. & Oosterlaan, J. Physical exercise and executive functions in preadolescent children, adolescents and young adults: a meta-analysis. *Br. J. Sports Med.* **48**, 973–979 (2014).

37. Chang, Y. K., Labban, J. D., Gapin, J. I. & Etnier, J. L. The effects of acute exercise on cognitive performance: A meta-analysis. *Brain Res.* **1453**, 87–101 (2012).

38. Lambourne, K. & Tomporowski, P. The effect of exercise-induced arousal on cognitive task performance: A meta-regression analysis. *Brain Res.* **1341**, 12–24 (2010).

39. Erickson, K. I., Hillman, C. H. & Kramer, A. F. Physical activity, brain, and cognition. *Curr. Opin. Behav. Sci.* **4**, 27–32 (2015).

40. González-Fernández, F., Etnier, J. L., Zabala, M. & Sanabria, D. Vigilance performance during acute exercise. *Int. J. Sport Psychol.* **48**, 435–447 (2017).

41. Andrillon, T., Poulsen, A. T., Hansen, L. K., Léger, D. & Kouider, S. Neural markers of responsiveness to the environment in human sleep. *J. Neurosci.* **36**, 6583–6596 (2016).

42. Chmura, J., Krysztofiak, H., Ziemba, A. W., Nazar, K. & Kaciuba-Uścilko, H. Psychomotor performance during prolonged exercise above and below the blood lactate threshold. *Eur. J. Appl. Physiol.* **77**, 77–80 (1997).

43. Del Giorno, J. M., Hall, E. E., O'Leary, K. C., Bixby, W. R. & Miller, P. C. Cognitive function during acute exercise: a test of the transient hypofrontality theory. *J. Sport Exerc. Psychol.* **32**, 312–323 (2010).

44. Maran, T. *et al.* Lost in Time and Space: States of High Arousal Disrupt Implicit Acquisition of Spatial and Sequential Context Information. *Front. Behav. Neurosci.* **11**, (2017).

45. Packard, M. G. & Goodman, J. Emotional arousal and multiple memory systems in the mammalian brain. *Front. Behav. Neurosci.* **6**, 14 (2012).

46. Schwabe, L. & Wolf, O. T. Stress and multiple memory systems: from 'thinking' to 'doing'. *Trends Cogn. Sci.* **17**, 60–68 (2013).

47. Broadhurst, P. L. The interaction of task difficulty and motivation: The Yerkes Dodson law revived. *Acta Psychol. Amst.* (1959).

48. Brown, J. S. *Motivation of behavior.* (Prabhat Prakashan, 1961).

49. Izquierdo, A. & Jentsch, J. D. Reversal learning as a measure of impulsive and compulsive behavior in addictions. *Psychopharmacology (Berl.)* **219**, 607–620 (2012).

50. Swainson, R. *et al.* Probabilistic learning and reversal deficits in patients with Parkinson's disease or frontal or temporal lobe lesions: possible adverse effects of dopaminergic medication. *Neuropsychologia* **38**, 596–612 (2000).

51. Remijnse, P., Nielen, M., Uylings, H. & Veltman, D. Neural correlates of a reversal learning task with an affectively neutral baseline: An event-related fMRI study. *NeuroImage* **26**, 609–618 (2005).

52.   Clark, L., Cools, R. & Robbins, T. W. The neuropsychology of ventral prefrontal cortex:
Decision-making and reversal learning. *Brain Cogn.* **55**, 41–53 (2004).
53.   Ersche, K. D., Roiser, J. P., Robbins, T. W. & Sahakian, B. J. Chronic cocaine but not chronic
amphetamine use is associated with perseverative responding in humans. *Psychopharmacology (Berl.)*
197, 421–431 (2008).
54.   Leeson, V. C. *et al.* Discrimination learning, reversal, and set-shifting in first-episode
schizophrenia: stability over six years and specific associations with medication type and
disorganization syndrome. *Biol. Psychiatry* **66**, 586–593 (2009).
55.   Kanen, J. W. *et al.* Effect of lysergic acid diethylamide (LSD) on reinforcement learning in
humans. *bioRxiv* 2020–12 (2021).
56.   den Ouden, H. E. M. *et al.* Dissociable Effects of Dopamine and Serotonin on Reversal
Learning. *Neuron* **80**, 1090–1100 (2013).
57.   Murphy, F. C., Michael, A., Robbins, T. W. & Sahakian, B. J. Neuropsychological impairment
in patients with major depressive disorder: the effects of feedback on task performance. *Psychol. Med.*
33, 455 (2003).
58.   Dombrovski, A. Y., Szanto, K., Clark, L., Reynolds, C. F. & Siegle, G. J. Reward signals,
attempted suicide, and impulsivity in late-life depression. *JAMA Psychiatry* **70**, 1020–1030 (2013).
59.   Dombrovski, A. Y. *et al.* Corticostriatothalamic reward prediction error signals and executive
control in late-life depression. *Psychol. Med.* **45**, 1413 (2015).
60.   Tavares, J. V. T. *et al.* Neural basis of abnormal response to negative feedback in unmedicated
mood disorders. *Neuroimage* **42**, 1118–1126 (2008).
61.   Mueller, E. M., Pechtel, P., Cohen, A. L., Douglas, S. R. & Pizzagalli, D. A. POTENTIATED
PROCESSING OF NEGATIVE FEEDBACK IN DEPRESSION IS ATTENUATED BY
ANHEDONIA: Research Article: Anhedonia and Feedback Processing in MDD. *Depress. Anxiety* **32**,
296–305 (2015).
62.   Drummond, S. The Effects of Total Sleep Deprivation on Cerebral Responses to Cognitive
Performance. *Neuropsychopharmacology* **25**, S68–S73 (2001).
63.   Jackson, M. L. *et al.* Deconstructing and reconstructing cognitive performance in sleep
deprivation. *Sleep Med. Rev.* **17**, 215–225 (2013).
64.   Gevers, W., Deliens, G., Hoffmann, S., Notebaert, W. & Peigneux, P. Sleep deprivation
selectively disrupts top-down adaptation to cognitive conflict in the Stroop test. *J. Sleep Res.* **24**, 666–
672 (2015).
65.   Whitney, P. *et al.* Sleep deprivation diminishes attentional control effectiveness and impairs
flexible adaptation to changing conditions. *Sci. Rep.* **7**, 1–9 (2017).
66.   Whitney, P., Hinson, J. M. & Nusbaum, A. T. A dynamic attentional control framework for
understanding sleep deprivation effects on cognition. *Prog. Brain Res.* **246**, 111–126 (2019).
67.   Bekinschtein, T. A. *et al.* Neural signature of the conscious processing of auditory regularities.
Proc. Natl. Acad. Sci. **106**, 1672–1677 (2009).
68.   Teigen, K. H. Yerkes-Dodson: A Law for all Seasons. *Theory Psychol.* **4**, 525–547 (1994).
69.   Mauss, I. B. & Robinson, M. D. Measures of emotion: A review. *Cogn. Emot.* **23**, 209–237
(2009).
70.   Tagliazucchi, E. Decoding Wakefulness Levels from Typical fMRI Resting-State Data Reveals
Reliable Drifts between Wakefulness and Sleep. 14.
71.   Comsa, I. M., Bekinschtein, T. A. & Chennu, S. Transient Topographical Dynamics of the
Electroencephalogram Predict Brain Connectivity and Behavioural Responsiveness During
Drowsiness. *Brain Topogr.* **32**, 315–331 (2019).
72.   Johns, M. W. A new method for measuring daytime sleepiness: the Epworth sleepiness scale.
sleep **14**, 540–545 (1991).

73. Holgado, D. *et al.* Tramadol effects on physical performance and sustained attention during a
20-min indoor cycling time-trial: A randomised controlled trial. *J. Sci. Med. Sport* **21**, 654–660 (2018).
74. Holgado, D. *et al.* Transcranial direct current stimulation (tDCS) over the left prefrontal cortex
does not affect time-trial self-paced cycling performance: Evidence from oscillatory brain activity and
power output. *PloS One* **14**, e0210873 (2019).
75. Zandonai, T. *et al.* Novel evidence on the effect of tramadol on self-paced high-intensity
cycling. *J. Sports Sci.* 1–9 (2021) doi:10.1080/02640414.2021.1877440.
76. Jagannathan, S. R. *et al.* Tracking wakefulness as it fades: Micro-measures of alertness.
NeuroImage **176**, 138–151 (2018).
77. Saper, C. B., Fuller, P. M., Pedersen, N. P., Lu, J. & Scammell, T. E. Sleep State Switching.
Neuron **68**, 1023–1042 (2010).
78. Ciria, L. F., Perakakis, P., Luque-Casado, A. & Sanabria, D. Physical exercise increases overall
brain oscillatory activity but does not influence inhibitory control in young adults. *NeuroImage* **181**,
203–210 (2018).
79. Ciria, L. F. *et al.* Oscillatory brain activity during acute exercise: Tonic and transient neural
response to an oddball task. *Psychophysiology* e13326 (2019) doi:10.1111/psyp.13326.
80. Tanaka, H., Monahan, K. D. & Seals, D. R. Age-predicted maximal heart rate revisited. *J. Am.*
Coll. Cardiol. **37**, 153–156 (2001).
81. Bates, D. *et al.* Package 'lme4'. *Convergence* **12**, (2015).
82. Tingley, D., Yamamoto, T., Hirose, K., Keele, L. & Imai, K. **mediation** : *R* Package for Causal
Mediation Analysis. *J. Stat. Softw.* **59**, (2014).
863

SUPPLEMENTARY MATERIAL

Dual-tasking effect of physical exercise on baseline performance

Is the detrimental effect of heightened arousal on behavioural performance truly due to the increased arousal level, or does it simply reflect a dual-task confounding effect of the physical and the cognitive task occurring simultaneously? Although this question is partially tackled in main analyses as the baseline arousal state of the heightened arousal states was also a dual-task condition (i.e., warm-up), we specifically explored whether a dual-tasking arousal baseline might be associated with poorer performance (i.e., higher RAR and RT variability), relative to a non-exercise adaptation period that participants performed just before the warm-up. Contrary to what we expected, the mixed-effects model yielded no reliable performance differences between the adaptation period and the warm-up (RAR: $t (39) = 1.41$, $p = 0.167$, $\beta = 0.18$; RT variability: $t (39) = 1.50$, $p = 0.14$, $\beta = 0.19$). To further confirm that baseline performance was equal or similar for all experiments, we analyzed the behavioural performance during baselines of Experiments 1 and 2 (i.e., wakefulness periods), as well as during baseline of Experiment 3 (i.e., warm-up period). Neither the number of responses needed to attain a reversal (RAR) nor accuracy showed reliable differences in baseline performance between experiments ($F < 1$). Subject-by-subject results show a similar distribution of performance across subjects in each Experiment (see supplementary figure 1).

Supplementary figure 1: Subject-by-subject baseline performance. Individual behavioural measures during baseline across databases. Grey bars represent individual participants within each experiment. All subjects are arranged by performance, from best to worst in RAR, and from worst to best in accuracy. The analysis revealed no reliable differences in behavioural performance during baseline periods across experiments.

Low arousal deceleration in behavioural dynamics

The transition from wakefulness to sleep involves a progressive, and sometimes nonlinear loss of responsiveness to external stimuli and a progressive increase of RT variability.[10,18,30] To further characterise the behavioural pattern of this transition, and compared to previous falling asleep tasks, we investigated the responsiveness and RT dynamics of the participants in the low arousal condition. We fitted a mixed-effects model separately for RT variability and omissions as dependent variables. As predicted from other cognitive tasks, [17,18,30,33] low arousal led to higher RT variability ($t (27.99) = 4.59$, $p < 0.001$, $\beta = 0.54$), which was accompanied by

897   a drastic increase in omitted response to stimuli (t (27.99) = 5.11, p < 0.001, β = 0.67),
898   compared with the baseline arousal state (see supplementary figure 2). These findings confirm
899   the convergence to other tasks of our arousal manipulation in probabilistic reversal learning in
900   its basic effects.
901

Supplementary figure 2. RT dynamic in low arousal. A) RT distribution during low (blue bars) and baseline
(grey bars) arousal states. B) Violins and overlaid box plots of the averaged reaction time variability across
participants in low and baseline arousal states. C) Subject-by-subject baseline differences in RT variability in low
arousal. Grey bars represent participants with a higher RT variability in the baseline compared with low arousal
state. Blue bars depict participants with a higher RT variability when arousal level was reduced compared with
baseline arousal state. Participants are sorted by the RT variability difference between baseline and the arousal
state.
910
911
912