Temperature Specific Regulation of NDR Kinase Orb6 by MAP kinase Sty1 to Promote Heat Stress Resilience

Laura P. Doyle[1], Robert N. Tams[1], Chuan Chen[1], Illyce Nuñez[1], Patrick Roman Haller[1], Fulvia Verde[1]**

Author Affiliations: [1]Department of Molecular and Cellular Pharmacology, University of Miami, Miami, FL.

Keywords (max. 6): NDR kinase Orb6; MAP kinase Sty1; Cdc42 GTPase; Sts5 RNP granule assembly; heat stress; *Schizosaccharomyces pombe*

Abstract

Cellular response to environmental fluctuations, such as increased temperature, is
crucial in promoting cell survival and plays an increasingly recognized role in cancer
biology. Important cellular functions altered by heat stress are cell polarization and protein
translation. Previous studies have shown that heat stress alters the dynamics of Cdc42,
a key regulator of cell polarization in eukaryotes, and promotes RNP granule formation,
reprogramming protein translation. The biological mechanisms underlying these vast
changes are only partially known. Here, we report that conserved NDR kinase Orb6, a
homologue of mammalian STK38, responds to heat stress and regulates heat stress
resilience by modulating Cdc42 dynamics and promoting stress granule assembly. Also,
we discovered a finely tuned mechanism whereby stress-activated MAP kinase Sty1
negatively regulates Orb6 kinase and Orb6 C-terminal phosphorylation during heat
stress. Orb6 inhibition by Sty1 increases the sensitivity of the cell to heat stress in a
temperature-specific manner, fostering increased stress resilience and metabolic
adaptation. These observations highlight the role of NDR kinase in the process of heat
adaptation and thermotolerance during environmental cell exposure to elevated
temperatures.

Summary statement

Nuclear Dbf2-related kinase Orb6 inhibition by stress-activated protein kinase Sty1
promotes heat stress resilience in a temperature specific manner.

Introduction

The ability to respond to varying environmental conditions, such as elevated temperature, is a vital cellular defense mechanism that promotes adaptation and survival (Zayani *et al.*, 2025). Heat stress response activates adaptive mechanisms including antioxidant defenses, metabolic shifts, and heat shock proteins (HSPs) induction (Jung *et al.*, 2013; Gomez-Pastor, Burchfiel and Thiele, 2018; Ciocca, Arrigo and Calderwood, 2013; Hu *et al.*, 2022; Ding and Gao, 2025). Activation of the heat shock response usually provides cytoprotective effects, and HSPs have been found to promote survival of cancer cells (Wu *et al.*, 2017). An increase in HSP levels has also been found in chronic inflammatory conditions such as rheumatoid arthritis (Fouani *et al.*, 2022). In eukaryotes, severe heat stress (typically ≥ 42°C) rapidly disrupts cellular function, triggering a robust heat shock response, protein denaturation and aggregation, as well as a decrease in protein synthesis.

During heat stress, cells undergo distinct morphological changes. Mammalian cells often lose their spread morphology and become more rounded due to cytoskeletal reorganization (Velichko *et al.*, 2013; Dressler *et al.*, 2005; Gungor *et al.*, 2014). However, the mechanisms that regulate morphological changes during heat stress are poorly understood. In the fission yeast *Schizosaccharomyces pombe*, an amenable model system, mild, sublethal heat stress at 36°C alters cell polarization and the dynamic behavior of Cdc42 GTPase (Vjestica *et al.*, 2013), a key regulator of cell morphology in eukaryotes (Schwamborn and Püschel, 2004; Stankiewicz and Linseman, 2014). In unstressed conditions, Cdc42 displays oscillatory dynamics at cell tips (Das *et al.*, 2012). During heat stress (Vjestica *et al.*, 2013) as well as in response to different environmental

conditions such as pheromone exposure (Bendezú and Martin, 2013), oxidative stress

(Salat-Canela *et al.*, 2021), or nitrogen starvation (Chen *et al.*, 2019), an alternative

"exploratory" pattern of Cdc42 emerges, where dynamic patches of active Cdc42 appear

and disappear throughout the cell membrane. While the emergence of exploratory Cdc42

dynamic involves the activation of the conserved stress-activated MAP kinase Sty1/Spc1

(Salat-Canela *et al.*, 2021; Mutavchiev, Leda and Sawin, 2016; Doyle *et al.*, 2025), the

mechanism that regulates Cdc42 dynamics shift during thermal stress is insofar poorly

understood.

During heat stress, cells also rapidly suppress global protein synthesis to

conserve energy and prevent accumulation of misfolded protein by the inhibition of

translation initiation and mRNA sequestration into stress granules and P-bodies for

storage or decay (Kedersha *et al.*, 2005; Grousl *et al.*, 2009). Stress granules (SGs) are

membraneless cytoplasmic aggregates that form in response to various stress conditions,

including heat shock, oxidative stress, osmotic stress, and infections (Marcelo *et al.*,

2021)**.** Their primary role is to protect mRNA and regulate translation during stress,

helping cells survive adverse conditions. The formation process relies on liquid–liquid

phase separation (LLPS), driven by multivalent interactions between RNA-binding

proteins (RBPs) and untranslated mRNAs. Stress granules are critical adaptive structures

that integrate translational control with cellular stress signaling (Desai *et al.*, 2026).

Dysregulation of SG dynamics is increasingly recognized as a contributor to human

disease, and is strongly linked to neurodegeneration (ALS, FTD), cancer and viral

infections (Ramaswami, Taylor and Parker, 2013; Anderson, Kedersha and Ivanov, 2015;

Aguzzi and Altmeyer, 2016; Gilks *et al.*, 2004; Alberti and Dormann, 2019).

Nuclear Dbf2-related (NDR) kinases are highly conserved members of the AGC

protein kinase family which regulate essential cellular processes including

morphogenesis, growth and proliferation, mitosis, and apoptosis (Hergovich *et al.*, 2006;

Cornils *et al.*, 2011; Yang *et al.*, 2014; Vichalkovski *et al.*, 2008; Verde, Mata and Nurse,

1995; Chiba *et al.*, 2009; Hergovich *et al.*, 2007; Hergovich *et al.*, 2009; Verde, Wiley and

Nurse, 1998). Several studies have implicated dysregulation of NDR kinases in the

development of cancer, and NDR also plays a role in neuronal differentiation (Adeyinka

et al., 2002; Millward *et al.*, 1998; Hauschild *et al.*, 1999; Ross *et al.*, 2000; Cornils *et al.*,

2010; Zhang *et al.*, 2015; Napoletano *et al.*, 2011). The fission yeast *S. pombe* has a

NDR/LATS kinase known as Orb6, which is responsible for regulation of polarized cell

growth (Verde, Mata and Nurse, 1995). We previously reported that Orb6 kinase has

separate roles in the control of Cdc42 GTPase (Das *et al.*, 2015; Das *et al.*, 2009), and

mRNA translation (Nunez *et al.*, 2016; Chen *et al.*, 2019). Orb6 governs cell polarity by

promoting the Ras1-dependent Cdc42 control axis (Chen *et al.*, 2019) and by inhibiting a

stress-activated Cdc42 module (Doyle *et al.*, 2025; Salat-Canela *et al.*, 2021) composed

of the guanine nucleotide exchange factor Gef1 (Das *et al.*, 2015; Das *et al.*, 2012; Das

et al., 2009; Coll *et al.*, 2003) and the Cdc42 GAP (GTPase Activating Protein) Rga3

(Doyle *et al.*, 2025; Gallo Castro and Martin, 2018). Downregulation of Orb6 kinase, which

suppresses Ras1 GTPase and derepresses the stress-activated Cdc42 module, leads to

the emergence of Cdc42 exploratory dynamics (Chen *et al.*, 2019; Das *et al.*, 2009).

Orb6 kinase also promotes polarized cell growth by spatiotemporal regulation of

RNA-binding protein Sts5 (Nunez *et al.*, 2016; Chen *et al.*, 2019). Sts5 is an RNA binding

protein (RBP) homologous to human Dis3L2, which is associated with Perlman's

syndrome and Wilm's tumor in humans (Astuti *et al.*, 2012; Lv *et al.*, 2015; Robinson *et*

al., 2015; Malecki *et al.*, 2013; Toda *et al.*, 1996; Vaggi *et al.*, 2012; Jansen *et al.*, 2009;

Kurischko *et al.*, 2011). Sts5 functions to repress translation of specific mRNAs, many of

which encode proteins involved in polarized cell growth, via binding and sequestration

into ribonucleoprotein (RNP) granules (Nunez *et al.*, 2016), under conditions of nutritional

stress (Nunez *et al.*, 2016; Chen *et al.*, 2019; Magliozzi and Moseley, 2021).

Environmental stress such as nitrogen starvation, or crowded growth conditions leading

to stationary phase, inhibit Orb6 kinase activity, which results in the translational

repression of mRNAs involved in polarized growth via formation of Sts5 granules (Nunez

et al., 2016; Chen *et al.*, 2019).

In this paper, we report that heat stress downregulates Orb6 kinase activity,

promoting Cdc42 exploratory dynamics and Sts5 coalescence into stress granules.

Further, we find that Orb6 kinase inhibition increases stress granule formation and

promotes cell resilience to heat stress. Recently, we reported that stress activated MAP

kinase Sty1 negatively regulates Orb6 kinase activity during nitrogen starvation (Doyle *et*

al., 2025). In this paper, we show that MAP kinase Sty1 has also a role in the induction

of exploratory Cdc42 dynamics, Sts5 granule assembly, and the inhibition of Orb6 kinase

activity during mild heat stress. Furthermore, we find that the levels of phosphorylation of

Orb6-Thr456, the C-terminal hydrophobic motif, are decreased upon heat stress and that

Sty1 activation enhances the sensitivity of Thr456 phosphorylation and Orb6 kinase

function to heat stress. Thus, our observations identify a temperature specific regulation

of NDR kinase Orb6 by MAP kinase Sty1 to promote exploratory Cdc42 dynamics and

stress resilience.

Results

Orb6 kinase function and activity are downregulated by heat stress.

We have previously shown that Orb6 kinase is sensitive to nutritional stress, and that downregulation of Orb6 kinase activity alters cell morphology and protein translation during nitrogen deprivation (Chen *et al.*, 2019). During nutritional stress, Orb6 kinase activity decreases and therefore phosphorylation of the Cdc42 control factors and Orb6 substrates Gef1 (Cdc42 GEF) and Rga3 (Cdc42 GAP) declines, promoting exploratory Cdc42 dynamics (Das *et al.*, 2015; Chen *et al.*, 2019; Doyle *et al.*, 2025). The onset of exploratory Cdc42 dynamics and increased Gef1 membrane localization were previously reported to occur also during exposure to sublethal mild heat stress at 36°C (Vjestica *et al.*, 2013). Therefore, we investigated if Orb6 kinase has a role in the cellular response to heat stress. Consistent with previous findings, we found that Gef1-3YFP localizes to the cell membrane after 30 minutes at 36°C (Fig. 1A, B) as compared to the control which remained at 25°C. Further, we found that Gef1-3YFP also localizes to the cell membrane at the higher, severe heat stress temperature of 42°C (Fig. 1A, B).

As Gef1 is a substrate that is phosphorylated by Orb6 at serine 112, we sought to measure Orb6 activity using a previously described custom phospho-specific antibody for Gef1-S112 (Chen *et al.*, 2019). We found that Gef1-S112 phosphorylation decreases after 30 minutes at 36°C, a sublethal temperature that induces heat stress, and is further exacerbated at 42°C, a lethal temperature that induces a severe heat shock response (Fig. 1C,D). These results indicate the Orb6 activity decreases during both mild and severe heat stress exposure, fostering exploratory Cdc42 dynamics.

During heat exposure, cells reorganize their RNA and protein metabolism by

forming two major types of cytoplasmic granules: Stress Granules (SGs) (Guzikowski,

Chen and Zid, 2019; Mahboubi and Stochaj, 2017), and Processing Bodies (P-bodies)**.**

Both are membraneless organelles formed by liquid–liquid phase separation, but they

have distinct roles and compositions. We previously demonstrated that Orb6 kinase

controls polarized cell growth and protein translation by phosphorylating the highly

conserved RNA binding protein Sts5. Sts5, an Orb6 substrate, remains diffused in the

cytoplasm when Orb6 is active, whereas it partially co-localizes with P-(Processing)

Bodies following Orb6 inhibition, or in media lacking glucose or nitrogen (Nunez *et al.*,

2016; Magliozzi and Moseley, 2021).

Consistent with Orb6 kinase activity being downregulated by heat exposure, we

found that Sts5-3YFP remains diffused in the cytoplasm at 25°C, whereas it forms puncta

at both 36°C and 42°C (Fig. 1E, F). We found that Sts5-3YFP puncta readily associate

with stress granules (using Pabp1-dsRed stress granule marker) at 42°C (Figure 1E,g-i,

1G), but not at 36°C where we find no stress granule assembly (Figure 1E,d-f, 1G).

Conversely, we find that Sts5-3YFP colocalizes with P-body marker Dcp1-mCherry at

both 36°C and 42°C (Fig. S1).

We sought to further confirm Sts5 and stress granule co-localization, under

conditions of glucose limitation, which strongly induces stress granules formation (Nilsson

and Sunnerhagen, 2011) as well as Sts5 aggregation into RNP granules (Nunez *et al.*,

2016). We investigated the localization of *ssp1* mRNA, a Sts5-bound mRNA which is

translationally repressed upon formation of Sts5 puncta (Nunez *et al.*, 2016) by RNA-

FISH analysis. *ssp1* encodes a CamK kinase (Hanyu *et al.*, 2009; Matsusaka *et al.*, 1995).

We observed co-localization of Sts5 puncta, *ssp1* mRNA, and stress granules in cells

cultured in medium lacking glucose (Fig. 2A). We previously used *ssp1* mRNA to detect

co-localization of Sts5 with P-bodies (Nuñez *et al.*, 2016) .

Overall, these findings show that Orb6 kinase activity is downregulated upon heat

stress and heat shock, fostering Cdc42 exploratory dynamics and coalescence of RNA

binding protein Sts5 into stress granules and P bodies.

Orb6 inhibition and Sts5 Aggregation Promote Stress Granule Assembly.

We have previously demonstrated that Orb6 inhibition promotes the formation of

P-Bodies in a Sts5 dependent manner (Nunez *et al.*, 2016), suggesting that Sts5

aggregates may promote P-Body formation during glucose starvation. To test whether

Sts5 plays a role in stress granule nucleation, we followed the formation of stress granules

in the *sts5Δ* mutant using the stress granule marker Pabp1-DsRed. In this particular

experiment, we induced stress granules via glucose deprivation, which strongly induces

stress granules assembly, facilitating quantification. We found that the number of stress

granules decreased by 44% in *sts5Δ* mutant cells as compared to control cells (Fig. 2B,

b, d; Fig. 2D) cultured for 20 minutes in media lacking glucose. In media containing

glucose, Pabp1-DsRed is largely diffuse within the cytoplasm in both the *sts5Δ* and

control strains (Fig. 2B; a, c).

To determine if inhibition of Orb6 kinase drives stress granule formation, we

treated *orb6-as2 pabp1-DsRed* cells with 1-NA-PP1 inhibitor and found that inhibition of

Orb6-as2 at 25°C does not induce stress granule formation (Fig. 2C; a, c), These findings

indicate that Orb6 kinase inhibition alone is not sufficient to induce stress granule

formation, while it is sufficient to promote P-Body formation, as we previously reported

(Nunez *et al.*, 2016). However, when 1-NA-PP1 treated cells were subsequently heat

shocked for 20 minutes at 42ºC, we observed a remarkable increase in the number of

stress granules, as compared to the heat shocked DMSO control (Fig. 2C; b, d). The

number of stress granules formed was quantified for three independent experiments

(n=90 cells/condition), and the number of stress granules increased, on average, by 54%

for the 1-NA-PP1 treated cells as compared to the heat shocked DMSO control (Fig. 2E).

Overall, these data indicate that the presence of Sts5 is important for the formation of

stress granules, and that Orb6 kinase plays a role in this process by modulating the extent

of stress granule formation.

Sts5 aggregation Modulates Heat Resilience in *S. pombe.*

Stress granule formation provide a mechanism to protect cells from environmental stress

and promote cell survival (Mahboubi and Stochaj, 2017). These condensates can

modulate susceptibility to heat stress (Riback *et al.*, 2017; Kroschwald and Alberti, 2017).

Since Sts5 responds to heat by co-localizing with both stress granules and P-Bodies (Fig.

1E, Fig. S1) and is important for stress granule formation (Fig. 2B), we hypothesized that

modulating the assembly of Sts5 into RNP granules plays a role in cell protection from

heat stress. Sts5 contains an intrinsically disordered domain (IDD), which is

phosphorylated by Orb6 kinase at serine 86 (Chen *et al.*, 2019). Modification of Sts5

serine 86 to alanine promotes increased Sts5 puncta formation, increases cell resilience

during nutritional stress and promotes chronological lifespan extension (Chen *et al.*,

2019). Thus, we tested if loss of Sts5 (*sts5Δ*), or the presence of the *sts5-S86A* mutation

affect cell survival during extended heat stress or heat shock. To test the role of Sts5

during heat stress and in heat tolerance (exposure to heat for extended times), wild-type,

sts5Δ, *sts5-HA*, and *sts5-S86A-HA* log-phase cells were incubated for 6 hours at heat

stress temperature (36.5°C) or control temperature (25°C) in a shaking incubator and

plated to calculate colony forming units (CFUs) (see Methods). We found that the *sts5Δ*

strain exhibited decreased survival compared to wild-type (Fig. 3A). Conversely, the *sts5-*

S86A-HA strain displayed increased survival, as compared to the *sts5-HA* or wild-type

controls (Fig. 3A). The *sts5-HA* strain exhibited similar survival as compared to the wild

type, as expected. We previously established that addition of the *sts5-S86A* mutation

does not alter Sts5 protein levels (Chen *et al.*, 2019).

To test response upon acute heat shock, wild-type, *sts5Δ*, *sts5-HA*, and *sts5-*

S86A-HA strains were exposed to a 48°C heat shock for 15 minutes. We found that the

sts5Δ strain exhibited decreased survival as compared to the wild-type control (Fig. 3B).

Again, the *sts5-S86A-HA* mutant exhibited increased survival as compared either the

wild-type or *sts5-HA* controls (Fig. 3B).

Thus, our data indicate that Sts5 coalescence is important for survival and heat

stress resilience following various durations and intensities of exposure to elevated

temperatures. Therefore, we tested whether Sts5 granule assembly is modulated by

increased temperature and whether the *sts5-S86A* mutation increases formation of Sts5

puncta following exposure to heat. To do this, *sts5-3xYFP* and *sts5-S86A-3xYFP* strains

were cultured at 29°C, 32°C, 35.5°C, or 42°C for 30 minutes, while control *sts5-3xYFP*

and *sts5-S86A-3xYFP* cells were maintained at 25°C throughout the experiment.

Following heat stress, fluorescence microscopy was performed and the number of Sts5

puncta were counted at each temperature. The *sts5-3xYFP* control began to form visible

puncta at 35.5°C (Fig. 3C; a-d), whereas the *sts5-S86A-3xYFP* mutant began to form

puncta already at 25°C (Fig. 3C, f-j). These differences were statistically significant (Fig.

3D), and *sts5-S86A* predisposition to the formation of puncta was readily apparent until

the temperature reached 42°C at which point the control formed approximately the same

number of puncta (Fig. 3Ce, Cj, 3D).

Our findings indicate an important role for serine 86 phosphorylation in modulating

the extent of Sts5 puncta assembly in response to increased temperatures (Fig. 3C, D).

Since Orb6 kinase phosphorylates serine 86 (Chen *et al.*, 2019), we tested if

downregulation of Orb6 kinase also increases cell resilience to heat stress. To

transcriptionally repress Orb6, a thiamine-repressible o*rb6-as2* strain was used. Log

phase wild-type and *orb6-as2* cells were grown with or without 15 µM thiamine for 16

hours at 32°C. Samples were diluted to an equivalent optical density and were heat

shocked at 48°C for 15 minutes while control samples were maintained at 32°C. The

orb6-as2 strain, in the presence of thiamine, exhibited a striking increase in survival

following heat shock compared to the untreated control (Fig. 3E). This striking increase

in resilience was readily observed on Petri plates (Fig. S2) and was notably higher than

the relative increase in heat resilience observed for the *sts5-S86A* mutant (Fig. 3A-B).

Overall, these data indicate that the Orb6 kinase modulates stress granule assembly and

resilience to heat stress, in part, by phosphorylating mRNA binding protein Sts5 on serine

86.

Conserved MAP kinase Sty1 negatively regulates Orb6 activity in a temperature specific manner.

The mitogen-activated protein kinase (MAPK) Sty1/Spc1 is activated upon exposure to diverse stressors such as nitrogen starvation, osmotic changes, oxidative stress, and heat shock (Toone and Jones, 1998; Shiozaki and Russell, 1996; Millar, Buck and Wilkinson, 1995; Degols, Shiozaki and Russell, 1996). Recently, we have shown that Sty1 activation during nitrogen starvation negatively regulates Orb6 activity (Doyle *et al.*, 2025). Therefore, we tested if Sty1 has a role in Orb6 regulation during heat stress, by visualizing Sts5-3YFP aggregation during either mild or severe heat stress (36°C or 42°C) for 30 minutes in the presence or absence of *sty1.* We find that upon loss of *sty1,* Sts5-3YFP does not assemble into granules at 36°C but it still assembles into granules at 42°C (Fig. 4A). The number of Sts5 granules per cell at 42°C is similar in both the presence or absence of *sty1* (Fig. 4B). These results indicate that Sts5 granule assembly during heat stress is dependent on the presence of Sty1 at 36°C but not at 42°C.

Previously, we have shown that Orb6 phosphorylates two regulators of Cdc42, a GEF Gef1 (Das *et al.*, 2015) and a GTPase activating protein (GAP) Rga3 (Doyle *et al.*, 2025) to spatially regulate Cdc42 dynamics. When Orb6 is inhibited during nitrogen starvation, Gef1 and Rga3 move to the cell membrane and facilitate the onset of exploratory Cdc42 dynamics, where active Cdc42 forms dynamic patches along the cell membrane (Doyle *et al.*, 2025). These exploratory Cdc42 dynamics are also found upon different stress exposure, including heat stress at 36°C (Vjestica *et al.*, 2013). Thus, we sought to investigate if exploratory Cdc42 dynamics were dependent on Sty1 upon heat exposure. We visualized active Cdc42 using fluorescently tagged Cdc42-Rac Interactive

Binding domain bioreporter CRIB-GFP or the Cdc42 GEF Gef1 using Gef1-3YFP. We

find that exploratory Cdc42 dynamics and Gef1-3YFP membrane localization are induced

by heat exposure at 36°C and 42°C and that these effects are dependent on Sty1 at 36°C,

but not 42°C (Fig. 4C-F).

Further, we tested the effects of heat stress and heat shock on Gef1-S112

phosphorylation by Orb6 kinase, in the presence and absence of *sty1*. We find that Gef1-

S112 phosphorylation decreases upon exposure to 36°C and 42°C (Fig. 4G,H). In the

absence of *sty1,* we find that Gef1-S112 phosphorylation remains constant at 36°C but

decreases to similar levels as wild-type at 42°C (Fig. 4G, H). Sty1 activation in these

experimental conditions was validated by measuring levels of Atf1, a transcription factor

phosphorylated by Sty1 upon Sty1 activation (Shiozaki, Shiozaki and Russell, 1998;

Wilkinson *et al.,* 1996; Sun *et al.*, 2024). As expected, we find that Atf1 levels increase

upon cell exposure to 36°C or 42°C. (Fig. 4G, I), confirming that Sty1 is activated under

these experimental conditions.

Overall, these results indicate that Sty1 kinase activation during temperature stress

exerts a temperature specific effect on Orb6 kinase activity; Sty1 activation is necessary

to inhibit Orb6 activity and biological functions at 36°C while is not necessary at higher,

heat shock temperatures (42°C).

Phosphorylation of the C-terminal hydrophobic motif Orb6-T456 is temperature

sensitive.

To better study the molecular mechanisms of Orb6 kinase regulation, we

developed a phospho-specific antibody that recognizes Orb6 phosphorylation by an

upstream kinase at the C-terminal hydrophobic motif site, Threonine 456 (Fig. 5A).

Phosphorylation of the conserved phosphorylation site, Thr456, is essential for Orb6

function (Liu and Young, 2012). We validated the specificity of the phospho-Orb6-T456

antibody (hereafter referred to as pOrb6-T456) by performing a dot blot with 4 different

peptides of Orb6. The pOrb6-T456 antibody was specific only to phosphorylated

Threonine 456 and not to the corresponding unphosphorylated peptide or a similar

sequence at the C-terminal, Threonine 464 (Fig. 5B). Previous studies have identified

interactions between Nak1 kinase and Orb6 kinase suggesting that Nak1 functions

upstream of Orb6 (Liu and Young, 2012). However, direct phosphorylation of Orb6 by

Nak1 has yet to be reported. To test if Orb6-T456 phosphorylation is Nak1-dependent,

we exposed temperature-sensitive mutant of *nak1 (nak1-ts/orb3-167)* (Snell and Nurse,

1994; Leonhard and Nurse, 2005) to restrictive temperature (36°C) for 3 hours to

downregulate Nak1 activity and measured pOrb6-T456 levels in HA-tagged Orb6 (Fig.

5C). We find that pOrb6-T456 strongly decreases at 36°C in the *nak1-ts* mutant (Fig. 5C,

D). Further, we find that pOrb6-T456 also decreases to a lesser extent, at 36°C in the

control *nak1+* cells. (Fig. 5C, D). These results indicate the Orb6 phosphorylation at the

C-terminal hydrophobic motif T456 site is dependent on Nak1. Our observations also

suggest that Orb6-T456 phosphorylation is affected by heat stress.

To test if Sty1 kinase has a role in modulating Orb6-T456 phosphorylation, we

exposed cells to 36°C or 42°C for 30 minutes in the presence or absence of *sty1.* We

found that pOrb6-T456 phosphorylation decreases upon exposure to heat, in both control

sty1+ or *sty1Δ* deletion mutants (Fig. 5E-F). However, loss of *sty1* leads to overall

increased levels of pOrb6-T456, as compared to control cells (Fig. 5E-F). This increase

is particularly evident at 25°C and 36°C, indicating that Sty1 regulates the levels of

phosphorylation of the Orb6 protein at the T456 site. Our observations suggest that the

increased levels of Orb6 phosphorylation, in the absence of Sty1, play a role in

maintaining sustained Orb6 kinase activity during heat stress at 36°C, thereby

suppressing the onset of Cdc42 exploratory dynamics and the induction of Sts5 granule

formation. This idea is supported by the observation that the extent of Gef1-S112

phosphorylation (a readout of Orb6 kinase activity) is similar in *sty1Δ* cells at 36°C and

unstressed wild-type controls at 25°C. Since Orb6 T456 phosphorylation decreases in

response to heat in both control and *sty1Δ* deletion mutant cells, our data also point to a

Sty1-independent mechanism of downregulation of Orb6 activity in response to

increasing heat stress.

Discussion

Conserved NDR kinase Orb6 activity is thermo-sensitive.

Proper cellular response to environmental fluctuations, such as heat exposure, is critical

for survival and is increasingly recognized as a key factor in human disease. Heat stress

disrupts essential cellular processes, including cell polarization (Vjestica *et al.*, 2013;

Pawlik *et al.*, 2013; Singh *et al.*, 2016; Delley and Hall, 1999) and protein translation

(Kedersha *et al.*, 2005; Grousl *et al.*, 2009). However, the underlying mechanisms

mediating these effects are not completely understood.

Here, we demonstrate that the conserved fission yeast NDR kinase Orb6,

homologous to mammalian STK38, is inactivated by heat stress, a process that alters the

pattern of active Cdc42 distribution at the membrane and facilitates stress granule

assembly. Overall, our findings provide experimental evidence explaining, at least in part,

some of the physiological effects of heat stress. We also uncover a regulatory mechanism

in which the stress-activated MAP kinase Sty1 negatively regulates Orb6 activity and

Orb6 C-terminal phosphorylation during heat stress (Fig. 6). This inhibition sensitizes cells

to temperature-specific heat stress, promoting adaptive metabolic responses and

thermotolerance. These findings underscore the pivotal role of NDR kinase signaling in

cellular adaptation to elevated temperatures. Further, our observations reveal a novel role

for conserved stress activated MAP kinase Sty1 in RNP granule formation, mediated by

Orb6 kinase.

Orb6 downregulation fosters the emergence of exploratory Cdc42 dynamics during

heat stress.

Cdc42 GTPase is a key regulator of cell polarization and is a highly conserved

Rho-family GTPase that regulates cell polarity, actin cytoskeleton organization, and

vesicle trafficking**.** In mammalian myoblasts, heat stress reduces cell migration, at least

in part through *CDC42* downregulation (Lu *et al.*, 2023). In fission yeast, under normal

conditions, active Cdc42-GTP oscillates between cell tips to promote polarized growth

(Das *et al.*, 2012). Upon environmental stress (including heat), Cdc42 adopts

an exploratory activation pattern along the cell membrane, deviating from its usual tip-

localized oscillations (Vjestica *et al.*, 2013; Chen *et al.*, 2019; Salat-Canela *et al.*, 2021;

Haupt, Ershov and Minc, 2018). Here, we show that during both mild and severe heat

stress, Orb6 kinase is inhibited. We previously showed that Orb6 inhibition leads to

dephosphorylation of Cdc42 regulators Gef1 and Rga3, and their release from the grip of

14-3-3- protein Rad24 (Das *et al.*, 2015; Doyle *et al.*, 2025), thereby switching on the

stress-activated Cdc42 module, which Gef1 and Rga3 are part of (Salat-Canela *et al.*,

2021; Doyle *et al.*, 2025). Our findings provide a mechanistic explanation of the process

of exploratory Cdc42 dynamics activation during heat stress. The functional role of

exploratory Cdc42 dynamics activation during heat stress, or other stresses, however, is

still unclear. Induction of exploratory Cdc42 dynamics during nitrogen deprivation (Chen

et al., 2019) may have the function to facilitate the search for mating partners (Bendezú

and Martin, 2013). It is possible that the onset of Cdc42 exploratory dynamics in other

stresses is a byproduct of engaging the same MAP Kinase-Orb6 kinase regulatory axis

(Doyle *et al.*, 2025). Alternatively, temporary loss of cell polarity could signal a

physiological transition for the cell, providing the opportunity to activate and inactivate

alternative polarity sites upon recovery from stress. In fission yeast, recovery from a

disrupting event, such as actin depolymerization by Latrunculin A exposure, can alter the

pattern of Cdc42 distribution and ensuing cell polarity (Mutavchiev, Leda and Sawin,

2016). Finally, disrupting cell polarization, in combination with translational repression

(see below) may be part of a general physiological strategy to slow cell growth, in an effort

to preserve energy (Salat-Canela *et al.*, 2023) and extend cell lifespan (Chen *et al.*, 2019).

Sts5 assembles into RNP granules and promotes thermotolerance.

Upon exposure to a variety of external stressors, cells form stress granules that

help promote stress resilience and survival. The RNA binding protein Sts5 is highly

conserved in eukaryotes and is related to human Dis3L2, which is associated with

Perlman's syndrome and Wilm's tumor in humans. In fungi, Sts5 is closely related to

Saccharomyces cerevisiae and *Candida albicans* Ssd1, and *Neurospora crassa* GUL-1

(Astuti *et al.*, 2012; Lv *et al.*, 2015; Robinson *et al.*, 2015; Malecki *et al.*, 2013; Toda *et*

al., 1996; Vaggi *et al.*, 2012; Jansen *et al.*, 2009; Kurischko *et al.*, 2011). These RNA-

binding regulators fine-tune translation of cell wall and morphogenesis-related genes,

crucial for stress adaptation, survival, and pathogenicity. In the yeasts

Schizosaccharomyces pombe and *Saccharomyces cerevisiae*, Sts5 and Ssd1,

respectively, bind to 5′ UTRs of mRNAs encoding cell wall proteins, repressing their

translation to control cell wall biogenesis (Nuñez *et al.*, 2016; Hogan *et al.*, 2008). Fission

yeast Sts5 also alters the translational landscape of mRNAs involved in polarized cell

growth and Cdc42 activity, regulating the Ras1-Scd1-Cdc42 regulatory axis (Chen *et al.*,

2019; Nuñez *et al.*, 2016). By promoting translational repression during nutritional stress,

both Sts5 and Ssd1 influence entry into and recovery from quiescence (Chen *et al.*, 2019;

Miles *et al.*, 2019). Finally, both Sts5 (this paper) and Ssd1 (Mir, Fiedler and Cashikar,

2009) are essential for tolerance to heat. Thus, the conservation of structure and cellular

function through such enormous evolutionary distance highlights the crucial role of these

RNA binding proteins in cell adaptation and survival.

How are these effects exerted in the cell? These proteins all share the property to

undergo liquid-liquid phase separation upon exposure to environmental stress, in a

manner that engages their N-terminal Intrinsically Disordered Domains (IDRs). The N-

terminal domain is crucial to mediate liquid-liquid phase separation of Sts5, Ssd1 and

GUL-1 into RNP granules, and is phosphorylated by conserved NDR kinase Orb6 in

Schizosaccharomyces pombe (Chen *et al.*, 2019), Cbk1 in *Saccharomyces cerevisiae*

(Kurischko *et al.*, 2011; Jansen *et al.*, 2009), and COT-1 in *Neurospora crassa* (Stein et

al., 2020) and other kinases (such as *S. pombe* Pak1 (Magliozzi and Moseley, 2021)).

We previously showed that serine 86 in the N terminus of Sts5 has a crucial role in

regulating Sts5 assembly into RNP granules, by mediating N-terminal Sts5 interaction

with 14-3-3 protein Rad24 (Chen *et al.*, 2019). Association of Sts5 with Rad24 inhibits

Sts5 phase separation into granules and maintains active protein translation of Sts5-

associated RNAs (Nuñez *et al.*, 2016). Conversely, mutation of serine 86 to alanine (*sts5-*

S86A), which prevents phosphorylation at that site, inhibits Rad24 association with Sts5

and thereby promotes RNP granule assembly and decreased translation of Sts5-

assicated RNAs (Chen *et al.*, 2019). We found that by subtly promoting RNP assembly,

the *sts5-S86A* mutation seemingly enhances cell resilience to stress exposure, extending

chronological lifespan during nutritional stress (Chen *et al.*, 2019). Here, we show that the

sts5-S86A mutation increases cellular resilience to heat stress, while loss of Sts5

increases heat stress vulnerability. We previously showed that Sts5 alters the

translational landscape of the cell by decreasing specific mRNA availability for translation,

such as *ssp1* mRNA and *efc25* mRNA ((Nuñez *et al.*, 2016); Fig. 2C). Consistent with a

role for Sts5 in translational repression during heat exposure, Ssp1 protein levels

drastically increase upon heat stress, when *sts5* is deleted (Nuñez *et al.*, 2016).

Processing bodies and stress granules are types of cytoplasmic ribonucleoprotein

(RNP) granules involved in mRNA regulation. A defining function of P-bodies and stress

granules is their dynamism, as these granules quickly assemble upon stress exposure

and disperse when stress conditions ameliorate. The difference between P-bodies

(processing bodies) and stress granules lies in their composition, formation triggers, and

cellular roles. P-bodies are involved in mRNA decay, storage, and translational repression

and contain enzymes for mRNA degradation (e.g., decapping enzymes Dcp1/Dcp2)

(Sakuno et al., 2004; Ingelfinger et al., 2002; van Dijk et al., 2002; Wang et al., 2002). P-

bodies are present under normal conditions but increase in size and number during

cellular stress. Stress granules, conversely, form in response to cellular stress (e.g., heat

shock, oxidative stress) and are triggered by translation initiation inhibition. Stress

granules store untranslated mRNAs to prevent their degradation and help cells recover

by allowing mRNAs to re-enter translation once stress resolves (Marcelo *et al.*, 2021;

Panas, Ivanov and Anderson, 2016).

While it is known that P-bodies and stress granules initially assemble through a

process known as liquid-liquid phase separation (Aguzzi and Altmeyer, 2016), much

remains to be elucidated regarding the mechanisms and factors involved in their

formation. Coalescence of RNA binding proteins into puncta is thought to promote

nucleation of both P-Bodies and stress granules (Kroschwald et al., 2015; Lin et al., 2015;

Elbaum-Garfinkle et al., 2015; Patel et al., 2015; Kato et al., 2012; Hyman, Weber and

Julicher, 2014; Brangwynne et al., 2009; Lee et al., 2013; Brangwynne, 2013; Becker and

Gitler, 2015).

During nitrogen or glucose deprivation, Sts5 proteins assemble into puncta that

partially co-localize with the P-Body marker Dcp1-mCherry (Chen *et al.*, 2019; Nuñez *et*

al., 2016). In this paper, we find that also upon mild heat stress (36°C), Sts5 coalesces

into granules that colocalizes with P-body marker Dcp1. At the heat stress temperature

36°C, we do not observe formation of detectable Pabp1-containing stress granules. Upon

severe heat stress (42°C) when visible stress granules form, Sts5 also co-localize with

stress granules marker, Pabp1. Consistent with the fact that both P-Bodies and RNA

binding proteins are thought to promote the assembly of stress granules (Buchan,

Muhlrad and Parker, 2008; Wheeler et al., 2016), we find that loss of *sts5* strongly reduces

the number of stress granules and decreases cell survival during heat shock. Conversely,

Orb6 inhibition prior to heat exposure, which promotes Sts5 coalescence into granules,

(Nunez *et al.*, 2016; Chen *et al.*, 2019), substantially increases the amount of stress

granules formed in the cell during severe heat stress (42°C) and drastically increases

resilience to heat stress and heat shock. However, since increase in thermo-resilience is

much greater for Orb6 inhibition than for the *sts5-S86A* mutation, it is likely that Orb6 has

additional roles in thermotolerance which are independent of Sts5.

Furthermore, since Orb6 kinase inhibition alone is not sufficient to drive stress

granule formation, it is likely that additional factors and regulatory controls play a role in

driving Sts5-mediated stress granule formation. For example, other factors such as the

vigilin homologue Vgl1 (Wen *et al.*, 2010), signaling factors such as calcineurin (Higa *et*

al., 2015), and protein kinase C Pck2 (Kanda *et al.*, 2021) also associate with stress

granules in response to thermal stress, and stress granule formation is also stimulated in

response to heat stress by the RNA-binding protein Nrd1 (Satoh *et al.*, 2012).

The MAP kinase Sty1-NDR kinase Orb6 regulatory axis regulates Cdc42 dynamics

and RNP granule assembly during heat stress.

Conserved MAP kinase Sty1 is the stress-activated MAP kinase (SAPK) in fission

yeast. It is the central component of the Wis1–Sty1 MAPK pathway**,** analogous to

mammalian p38/JNK pathways. Heat stress transiently activates Sty1 through its dual

phosphorylation by Wis1 at two conserved residues, Threonine 171 and Tyrosine 173,

and leads to Sty1 nuclear accumulation (Degols, Shiozaki and Russell, 1996; Smith *et*

al., 2002). Sty1 phosphorylation during heat stress is regulated by threonine and tyrosine

phosphatases, where tyrosine phosphatases, Pyp1 and Pyp2, are inhibited by heat

contributing to Sty1 activation and threonine phosphatases, Ptc1 and Ptc3, work to

attenuate Sty1 activity (Nguyen and Shiozaki, 1999). *Sty1Δ* mutants (Sty1 deletion) show

severe thermo-sensitivity**,** losing viability rapidly compared to wild type (Yamada *et al.*,

1997). Sty1 activation is essential for survival under both severe heat stress conditions

(≥42°C) (Berlanga *et al.*, 2010; Yamada *et al.*, 1997; Degols, Shiozaki and Russell, 1996),

as well as mild heat stress temperature 37°C (Yamada *et al.*, 1997)**.**

Our results indicate that Orb6 activity decreases upon exposure to heat stress and

heat shock. We recently reported that MAP kinase Sty1 negatively regulates Orb6 during

nitrogen starvation (Doyle *et al.*, 2025) providing a mechanistic understanding as to why

Sty1 activation elicits exploratory Cdc42 dynamics, a behavior that may facilitate ensuing

cell mating (Bendezú and Martin, 2013; Merlini *et al.*, 2016). We showed that Sty1 inhibits

Orb6 kinase, thereby releasing the Orb6 substrates Cdc42 GEF Gef1 and Cdc42 GAP

Rga3 from their association with 14-3-3- protein Rad24 (Doyle *et al.*, 2025), Sty1 also

directly phosphorylates Gef1 and Rga3, likely activating their enzymatic activity (Salat-

Canela *et al.*, 2021). This "coherent feedforward" mechanism promotes exploratory

Cdc42 dynamics during stress response (Salat-Canela *et al.*, 2021; Doyle *et al.*, 2025).

In this paper, we show that this Sty1-NDR kinase Orb6 regulatory axis has a

function in other stress responses, namely heat stress. Further, our data indicate that,

through Orb6, Sty1 also modulates the phase separation properties of mRNA binding

protein Sts5, and therefore the coalescence of P-bodies and stress granules in response

to heat exposure.

Interestingly, our observations show that loss of Sty1 completely suppresses the

effects of heat dependent Orb6 inhibition (the induction of Cdc42 exploratory dynamics,

Gef1 dephosphorylation, Sts5 aggregation) during moderate heat stress at 36 °C, but not

under heat shock conditions at 42°C. Experimentally, these findings suggest a

"temperature-specific" control of Orb6 kinase by Sty1 kinase, namely that the Sty1

function in Orb6 inhibition is only effective at sublethal 36 °C temperatures, but not at

42°C heat stress temperatures.

The role of Orb6 T456 c-terminal hydrophobic motive in thermal sensation.

To explore the mechanism behind this effect, we tested the role of the C-terminal

hydrophobic motif of Orb6 kinase in thermal sensation. The highly conserved hydrophobic

motif (HM) is a critical regulatory element in the AGC family of protein kinases**.** It contains

a phosphorylatable serine or threonine residue (in Orb6, Threonine 456) flanked by

hydrophobic amino acids. Phosphorylation of the HM is essential for stabilizing the active

conformation of the kinase, facilitating proper alignment of catalytic residues for efficient

substrate phosphorylation, and strongly correlates with kinase activity (Millward, Hess

and Hemmings, 1999; Hergovich *et al.*, 2006; Hergovich, 2013; Jansen *et al.*, 2006;

Brace, Hsu and Weiss, 2011; Parker *et al.*, 2020).

We discovered that the level of HM T456 phosphorylation in Orb6 kinase is

temperature-dependent, in a manner that is inversely correlated with increasing

temperatures. We find that upon loss of Sty1, the overall levels of Orb6-T456

phosphorylation substantially increase, an effect that is particularly striking at 25°C and

36°C, indicating that Sty1 has a role in the negative control of Orb6-T456 phosphorylation.

Further, we find that Orb6-T456 phosphorylation decreases during mild heat stress and

severe heat stress in both control or *sty1Δ* strains, suggesting that a Sty1-independent

mechanism also mediates Orb6 HM dephosphorylation during heat stress and heat

shock. The combined effect is that in *sty1Δ* cells exposed to heat stress at 36°C, the level

of Orb6 T456 phosphorylation is comparable to wild-type control cells grown in

unstressed conditions at 25°C. Consistent with the idea that the activity of Orb6 remains

high in *sty1Δ* cells at 36°C, Gef1 phosphorylation, Cdc42 distribution and Sts5 localization

are unperturbed, and cells appear unresponsive to heat stress. Overall, these results

indicate that Orb6 kinase inhibition by Sty1 kinase increases heat sensitivity during mild

heat stress (36°C), promoting assembly of Sts5 into granules (Fig.1E, 4A), coalescence

of Sts5 granules with P-bodies (Fig. S1), and downregulation of Sts5-associated mRNA

translation (Nuñez *et al.*, 2016). Thus, this finely tuned regulatory network ensures the

inhibition of Orb6 during mild heat stress (36°C), when stress granules do not form, but

when reprogramming of the translational landscape, and morphogenetic changes at the

membrane may be necessary for survival.

It is unclear what could be the Sty1-independent mechanism of temperature-

sensing that leads to Orb6 inactivation. One possible mechanism is that Nak1, the kinase

upstream of Orb6, is directly inhibited by heat stress. Another possible mechanism is that

heat affects protein structure of the Orb6 regulatory complex, perhaps the association of

the co-activator Mob2 to Orb6, or the interaction of the Mor2 scaffold with Nak1 and Orb6.

Direct inhibition of the Mob2-Orb6 complex by an inhibitory protein is also possible, as

detailed for the *S. cerevisiae* protein Lre1, which does not however have homologues in

S. pombe or higher eukaryotes (Mancini Lombardi *et al.*, 2013; Versele and Thevelein,

2001). Future experiments will address these possibilities.

In conclusion, our study highlights the role of the MAP kinase Sty1-NDR kinase

Orb6 regulatory axis in promoting thermotolerance during environmental heat exposure.

Thermotolerance refers to the ability of an organism, tissue, or cell to withstand elevated

temperatures without sustaining irreversible damage. The acquisition of thermotolerance

has enormous implications for the survival of crops, the spread of pathogenic organisms

to different latitudes, or the success of disease treatments, such as during cancer

hyperthermia therapy to enhance chemotherapy or radiation sensitivity. Thus, these

findings may have implications in higher eukaryotes due to the highly conserved nature

of NDR kinases across evolution.

Materials and Methods:

Strains and Growth Medium

Schizosaccharomyces pombe strains used in this study were derived from wild-type

strains 972 or 975 and are displayed in Table S1. *S. pombe* was maintained in yeast

extract plus supplements (YES) or Edinburgh minimal medium supplemented with 0.5%

ammonium chloride (EMMN) and additional supplements as needed (Histidine, Leucine,

Adenine, or Uracil) at a concentration of 225 mg/L. All strains were cultured at 25°C unless

otherwise noted. Liquid cultures were incubated at 180 RPM in a shaking incubator, and

all cultures were diluted daily such that they were maintained in logarithmic growth for a

minimum of 8 generations prior to performing experiments.

Fluorescence Microscopy

For experiments utilizing microscopy, samples were imaged with an Olympus BX61

fluorescent microscope using appropriate filters. Exposure times range from 1000-

2500ms for each experiment and remain consistent among all samples for that

experiment. Images were acquired and processed using Intelligent Imaging Innovations

SlideBook image analysis software (Version 6.0.4; Denver, CO) and prepared with Fiji

(Schindelin *et al.*, 2012).

RNP Granule Induction

To determine localization of Sts5 puncta relative to stress granules, we introduced the

marker Pabp1-DsRed, constructing a *sts5-3xGFP pabp1-DsRed* strain (FV2361). We

tested the localization of Sts5 relative to stress granules upon heat stress at 36°C or 42°C

for 30 minutes. To test the effects of glucose limitation, cells were grown in minimal

medium (EMM) containing 2% glucose for at least 8 generations, and then were

centrifuged, washed with the appropriate medium, and resuspended either in EMM

lacking glucose or in EMM with 2% glucose. Images are representative of samples from

three independent experiments.

RNA Fluorescent In Situ Hybridization (RNA FISH)

Localization of *ssp1* mRNA was determined using RNA-FISH as previously described,

with modifications according to Nunez *et al.* (Nunez *et al.*, 2016; Nilsson and

Sunnerhagen, 2011; Heinrich *et al.*, 2013; Brengues and Parker, 2007). *sts5-3xGFP*

pabp1-DsRed cells (FV2361) were cultured in EMM lacking glucose for 20 minutes prior

to fixation and hybridization of RNA was performed with 20-mer DNA oligonucleotides

(Stellaris) labeled with Quasar 705 fluorochromes.

Glucose Starvation and Orb6 Inhibition Assays

To determine if Sts5 was required for the formation of stress granules, a glucose

starvation assay was performed with *sts5+ pabp1-DsRed* (FV1684) and *sts5Δ pabp1-*

DsRed (FV3192) mutants. Samples were cultured to log phase in minimal medium (EMM)

and were washed once with 5 mL minimal medium lacking glucose. Cells were then

resuspended in 5 mL minimal medium lacking glucose and were incubated at 25°C and

180 RPM in a shaking incubator for 20 minutes. Following incubation, 1 mL aliquots were

centrifuged at 4000 RPM for 1 minute and resuspended in a small amount of residual

medium. Microscopy was performed as previously described and the number of stress

granules was manually counted. This experiment was performed in triplicate, and n=90

cells per strain per strain per condition in total.

For Orb6 inhibition assays, *orb6-as2 pabp1-DsRed* (FV3226) and *pabp1-DsRed*

control (FV1684) strains were cultured to log phase in minimal medium containing

adenine. 1-NA-PP1 was added to samples at a final concentration of 50 µM. For assays

in which Orb6 was inhibited and then cells were heat shocked, 1-NA-PP1 was added to

cells 15 minutes prior to heat shock, whereas DMSO was added for control cells. Samples

were then shifted to a 42ºC shaking water bath incubator for 20 minutes. Additional

controls were maintained at 25 ºC. Microscopy was performed as previously described

and stress granules were manually counted. This experiment was performed in triplicate,

and n = 90 cells per strain per condition in total.

Heat Shock Survival Assay

Wild-type (FV2644), *sts5Δ* (FV2674), *sts5-HA* (FV2645), and *sts5-S86A-HA* (FV2649)

strains were cultured in EMM and maintained in log-phase for a minimum of eight

generations. Each strain was diluted to O.D.$_{595nm}$ = 0.2 (approximately 4×10^6 cells/mL) in

5 mL EMM and allowed to grow at this density for one hour at 25°C shaking at 180 RPM.

Samples were either maintained at 25°C or heat shocked at 48°C for 15 minutes.

Following heat shock, experimental and control samples were each serially diluted, plated

in triplicate on EMM plates, and incubated at 25°C for approximately 72 hours. Colony

forming units per mL (CFU's/mL) were calculated for each condition, and percent survival

was calculated by dividing CFU's/mL for heat shocked replicates by their respective

untreated control CFU's/mL. Percent survival of each technical replicate was normalized

to the average untreated control percent survival. The average percent survival for each

strain is as follows: *wt*, 1.8%; *sts5Δ*, 0.2%; *sts5-HA*, 1.3%; and *sts5-S86A-HA*, 4.0%. This

experiment was performed in biological triplicate, and data was normalized to the average

wild-type percent survival for each independent experiment and subsequently graphed

as the relative survival rate. To determine the impact of intermediate to long-term

restrictive temperature on these strains, this experiment was repeated with modifications.

In this case, samples were incubated at 25°C or 36.5°C for 6 hours. Following incubation,

cells were serially diluted and plated in triplicate, and incubated at 25°C for approximately

72 hours. Calculation of CFU's/mL and normalization of data was performed as previously

described, and this experiment was performed in biological triplicate. The average

percent survival for each strain (normalized to the control at 25°C) is as follows: *wt*,

125.6%; *sts5Δ*, 65.2%; *sts5-HA*, 120.8%; and *sts5-S86A-HA*, 179.9%.

For heat shock survival analysis of the *orb6-as2* strain (FV2527) and its respective

wild-type control (FV2530), cells were initially pre-cultured in minimal medium

supplemented with adenine at 180 RPM and 32ºC. In order to repress the expression of

orb6-as2 the culture was split into two sets, either with or without thiamine at a final

concentration of 15 μM and was allowed to grow for ~16 hours. Cells were diluted to an

equivalent O.D.$_{595nm}$ of 0.2 (4×10^6 cells/mL) in appropriate media, and cells were allowed

to grow for an additional 2 hours. Samples were then heat shocked in a 48ºC shaking

water bath for 15 minutes while controls were maintained at 32ºC. All samples were then

serially diluted and plated on minimal medium supplemented with adenine, but lacking

thiamine, for subsequent quantification of CFU's/mL as previously described. The

average percent survival for each strain is as follows: *wt* -Thiamine, 0.1%; *wt* +Thiamine,

0.1%; *orb6-as2* -Thiamine, 0.1%; and *orb6-as2* +Thiamine, 18.9%.

Temperature Dependent Analysis of Sts5 Puncta

sts5-3xYFP (FV2518) and *sts5-S86A-3xYFP* (FV2522) strains were used to determine

whether Sts5 puncta are induced at lower temperatures when Sts5 is hyperactive as

compared to a wild-type allele. Each strain was cultured to log-phase in EMM, diluted to

O.D.$_{595nm}$= 0.1, and cells were allowed to grow for a minimum of 1 hour at 25°C. Aliquots

of each strain were created, and samples were heat shocked at 29°C, 32°C, 35.5°C, or

42°C for 30 minutes. Control cells were maintained at 25°C throughout the experiment.

Following exposure to heat shock, samples were imaged using an Olympus BX61

microscope. To determine differences in number of puncta, 20 cells per strain were

manually scored for Sts5 (YFP) puncta at each temperature.

Development of anti-pOrb6-T456 phospho-specific antibody

The anti-pOrb6-T456 phospho-specific antibody was custom-made by 21st Century

Biochemicals against a synthetic peptide corresponding to the phosphorylated Thr-456

(amino acids 449-462, NLAFLGYT*YKKFNY) of the fission yeast Orb6 protein. To detect

pOrb6-T456, we used a strain expressing N-terminally tagged HA-Orb6 *(HA-orb6-as2)*,

integrated in single copy behind the *nmt1* promoter, since increased expression of Orb6

allows for better detection of pOrb6-T456, as compared to C-terminally tagged Orb6-HA

or endogenous Orb6 where the detection signal is too low.

Protein extraction and western blot analysis

Protein extraction was modified from previously described protocol(Matsuo *et al.*,

2006) and performed as previously described (Doyle *et al.*, 2025). ~7.5 × 10^7 total cells

growing exponentially were harvested and cell pellets were resuspended in 300µL of H$_2$O

with protease and phosphatase inhibitor cocktail (Halt™ Protease and Phosphatase

Inhibitor Cocktail (100X)) and transferred to 1.5mL microcentrifuge tube. 300µL of 0.6M

NaOH with inhibitor cocktail (see above) was added to the cells and resuspended. Cells

were left to incubate for 5 minutes at room temperature, inverting the tubes 2-3 times

halfway through incubation. Cells were then centrifuged at 2300 RCF for 2 minutes. The

supernatant was discarded, and the pellet was resuspended in 75µL modified SDS buffer

(60mM Tris-HCl [pH 6.8], 4% 2-Mercaptoethanol, 4% SDS, 5% glycerol) with inhibitor

cocktail (see above) and boiled at 98°C for 3 minutes. Cells were placed on ice and

centrifuged at 4°C at 3500 RCF for 1 minute. 65µL of the supernatant (protein extract)

was collected with 60µL stored at -80°C immediately, and 5µL used for downstream

protein quantification assay. To quantify protein, the RC DC protein assay kit was used

as it is compatible with the concentration of reducing agents and detergents present in

the modified SDS buffer.

Standard western blotting procedures were performed as follows: protein was

separated using SDS-PAGE and transferred onto a nitrocellulose membrane. The

membranes were probed with antibodies of interest and visualized through fluorescent

detection on Li-Cor Odyssey CLx. Gef1-3YFP strains were used in detecting total Gef1

protein with anti-GFP (Roche, CAT#: 11814460001; RRID:AB_390913; dilution factor

(DF): 1:1000) and phosphorylated Gef1-S112 was detected using a custom-made

previously described antibody(Chen *et al.*, 2019). HA-Orb6as2 strains were used in

detecting total Orb6 protein with anti-HA (Biolegend, CAT#901501; RRID:AB_2565006;

DF: 1:3000) and phosphorylated Orb6-T456 was detected using a custom-made antibody

(see subsection "Development of anti-pOrb6-T456 phosphospecific antibody" in Methods

section). Anti-β-Actin (Abcam, CAT#: AB8224; RRID:AB_449644; DF: 1:4000) was used

as a loading control for all experiments. Two-color detection was used with using LiCor

IRDye secondary antibodies anti-rabbit 800CW (Li-Cor, CAT#: 926-32211;

RRID:AB_621843; DF: 1:15000) for anti-pGef1-S112 or anti-pOrb6-T456 and anti-mouse

680RD (Li-Cor, CAT#: 102673-408; RRID:AB_10956588; DF: 1:15000) for anti-GFP

(Gef1-3YFP) or anti-HA (HA-Orb6as2). Quantification of the blots was performed using

Image Studio software. To quantify, a rectangle was drawn around the signal band of

interest and pixel intensity was recorded. To remove background signal, background

settings were set to median, segment: top and bottom, border width: 3. With the pixel

intensity measurements, a ratio was determined between pGef1-S112 or pOrb6-T456

(fluorescent channel 800) and total Gef1-3YFP or total HA-Orb6as2 (fluorescent channel

700) to measure how much total Gef1-3YFP protein is phosphorylated at S112 or total

HA-Orb6as2 protein is phosphorylated at T456. Data was normalized by creating a ratio

with all values with the average ratio of control strains and/or conditions. Similar

quantification was performed to measure ratio of Atf1 to β-Actin during heat stress in

control cells and *sty1* deletion mutants using anti-Atf1 (Abcam, CAT#: AB18123;

RRID:AB_444264; DF: 1:2000) and anti-β-Actin (Abcam).

Quantification and statistical analysis

Data are presented as described in legends. Experiments were completed in

independent biological triplicates. A two-tailed unpaired Student's t test was used to

assess statistical significance between two groups. One-way or two-way analysis of

variance (ANOVA) followed by appropriate post hoc test was applied to evaluate the

difference between more than two groups (one-way) or more than 2 groups and more

than one condition (two-way). Statistical analyses and visualization were performed with

GraphPad Prism 10 (GraphPad Software, San Diego, CA). p-value < 0.05 was set as

the threshold for statistical significance. Power analyses were performed using G*Power

to determine minimum sample size(Faul *et al.*, 2007; Kang, 2021).

Competing Interests

The authors declare no competing interests.

Funding

Research reported in this publication was supported by the National Institute of Health

(NIH) R01 GM129514 and by the Sylvester Comprehensive Cancer Center which

receives funding from the National Cancer Institute of the National Institutes of Health

under Award Number P30CA240139.

Data and Resource Availability

Data and resource availability: All relevant data and details of resources can be found

within the article and its supplementary information.

References

1. Adeyinka, A., Emberley, E., Niu, Y., Snell, L., Murphy, L. C., Sowter, H., Wykoff, C. C., Harris,

A. L. and Watson, P. H. (2002) 'Analysis of gene expression in ductal carcinoma in situ

of the breast', *Clin Cancer Res,* 8(12), pp. 3788-95.

2. Aguzzi, A. and Altmeyer, M. (2016) 'Phase Separation: Linking Cellular Compartmentalization

to Disease', *Trends Cell Biol,* 26(7), pp. 547-558.

3. Alberti, S. and Dormann, D. (2019) 'Liquid-Liquid Phase Separation in Disease', *Annu Rev*

Genet, 53, pp. 171-194.

4. Anderson, P., Kedersha, N. and Ivanov, P. (2015) 'Stress granules, P-bodies and cancer',

Biochim Biophys Acta, 1849(7), pp. 861-70.

5. Astuti, D., Morris, M. R., Cooper, W. N., Staals, R. H., Wake, N. C., Fews, G. A., Gill, H.,

Gentle, D., Shuib, S., Ricketts, C. J., Cole, T., van Essen, A. J., van Lingen, R. A., Neri,

G., Opitz, J. M., Rump, P., Stolte-Dijkstra, I., Muller, F., Pruijn, G. J., Latif, F. and Maher,

E. R. (2012) 'Germline mutations in DIS3L2 cause the Perlman syndrome of overgrowth

and Wilms tumor susceptibility', *Nat Genet,* 44(3), pp. 277-84.

6. Becker, L. A. and Gitler, A. D. (2015) 'It's all starting to come together', *Elife,* 4.

7. Bendezú, Felipe O. and Martin, Sophie G. (2013) 'Cdc42 Explores the Cell Periphery for

Mate Selection in Fission Yeast', *Current Biology,* 23(1), pp. 42-47.

8. Berlanga, J. J., Rivero, D., Martín, R., Herrero, S., Moreno, S. and de Haro, C. (2010) 'Role of

mitogen-activated protein kinase Sty1 in regulation of eukaryotic initiation factor 2alpha

kinases in response to environmental stress in Schizosaccharomyces pombe', *Eukaryot*

Cell, 9(1), pp. 194-207.

9. Brace, J., Hsu, J. and Weiss, E. L. (2011) 'Mitotic exit control of the Saccharomyces

cerevisiae Ndr/LATS kinase Cbk1 regulates daughter cell separation after cytokinesis',

Mol Cell Biol, 31(4), pp. 721-35.

10. Brangwynne, C. P. (2013) 'Phase transitions and size scaling of membrane-less organelles',

J Cell Biol, 203(6), pp. 875-81.

11. Brangwynne, C. P., Eckmann, C. R., Courson, D. S., Rybarska, A., Hoege, C., Gharakhani,

J., Julicher, F. and Hyman, A. A. (2009) 'Germline P granules are liquid droplets that

localize by controlled dissolution/condensation', *Science,* 324(5935), pp. 1729-32.

12. Brengues, M. and Parker, R. (2007) 'Accumulation of polyadenylated mRNA, Pab1p, eIF4E,

and eIF4G with P-bodies in Saccharomyces cerevisiae', *Mol Biol Cell,* 18(7), pp. 2592-

602.

13. Buchan, J. R., Muhlrad, D. and Parker, R. (2008) 'P bodies promote stress granule
assembly in Saccharomyces cerevisiae', *J Cell Biol,* 183(3), pp. 441-55.

14. Chen, C., Pino, M. R., Haller, P. R. and Verde, F. (2019) 'Conserved NDR/LATS kinase
controls RAS GTPase activity to regulate cell growth and chronological lifespan',
Molecular Biology of the Cell, 30(20), pp. 2598-2616.

15. Chiba, S., Ikeda, M., Katsunuma, K., Ohashi, K. and Mizuno, K. (2009) 'MST2- and Furry-
mediated activation of NDR1 kinase is critical for precise alignment of mitotic
chromosomes', *Curr Biol,* 19(8), pp. 675-81.

16. Ciocca, D. R., Arrigo, A. P. and Calderwood, S. K. (2013) 'Heat shock proteins and heat
shock factor 1 in carcinogenesis and tumor development: an update', *Arch Toxicol,*
87(1), pp. 19-48.

17. Coll, P. M., Trillo, Y., Ametzazurra, A. and Perez, P. (2003) 'Gef1p, a new guanine
nucleotide exchange factor for Cdc42p, regulates polarity in Schizosaccharomyces
pombe', *Mol Biol Cell,* 14(1), pp. 313-23.

18. Cornils, H., Kohler, R. S., Hergovich, A. and Hemmings, B. A. (2011) 'Downstream of human
NDR kinases: impacting on c-myc and p21 protein stability to control cell cycle
progression', *Cell Cycle,* 10(12), pp. 1897-904.

19. Cornils, H., Stegert, M. R., Hergovich, A., Hynx, D., Schmitz, D., Dirnhofer, S. and
Hemmings, B. A. (2010) 'Ablation of the kinase NDR1 predisposes mice to the
development of T cell lymphoma', *Sci Signal,* 3(126), pp. ra47.

20. Das, M., Drake, T., Wiley, D. J., Buchwald, P., Vavylonis, D. and Verde, F. (2012)
'Oscillatory Dynamics of Cdc42 GTPase in the Control of Polarized Growth', *Science,*
337(6091), pp. 239-243.

21. Das, M., Nuñez, I., Rodriguez, M., Wiley, D. J., Rodriguez, J., Sarkeshik, A., R.III, Y. J.,
Buchwald, P. and Verde, F. (2015) 'Phosphorylation-dependent inhibition of Cdc42 GEF
Gef1 by 14-3-3 protein Rad24 spatially regulates Cdc42 GTPase activity and oscillatory
dynamics during cell morphogenesis', *Molecular Biology of the Cell,* 26(19), pp. 3520-
3534.

22. Das, M., Wiley, D. J., Chen, X., Shah, K. and Verde, F. (2009) 'The conserved NDR kinase
Orb6 controls polarized cell growth by spatial regulation of the small GTPase Cdc42',
Curr Biol, 19(15), pp. 1314-9.

23. Degols, G., Shiozaki, K. and Russell, P. (1996) 'Activation and regulation of the Spc1 stress-
activated protein kinase in Schizosaccharomyces pombe', *Mol Cell Biol,* 16(6), pp. 2870-
7.

24. Delley, P. A. and Hall, M. N. (1999) 'Cell wall stress depolarizes cell growth via

hyperactivation of RHO1', *J Cell Biol,* 147(1), pp. 163-74.

25. Desai, M., Gulati, K., Agrawal, M., Ghumra, S. and Sahoo, P. K. (2026) 'Stress granules:

Guardians of cellular health and triggers of disease', *Neural Regen Res,* 21(2), pp. 588-

597.

26. Ding, X. and Gao, B. (2025) 'Heat Stress-Mediated Multi-Organ Injury: Pathophysiology and

Treatment Strategies', *Comprehensive Physiology,* 15(3), pp. e70012.

27. Doyle, L. P., Chen, J.-S., Gould, K. L., McCollum, D. and Verde, F. (2025) 'Control of stress-

activated Cdc42 dynamics by the MAP kinase Sty1—NDR kinase Orb6

regulatory axis', *iScience,* 28(9).

28. Dressler, C., Minet, O., Beuthan, J., Novkov, V. and Müller, G. (2005) 'Microscopical heat

stress investigations under application of quantum dots', *J Biomed Opt,* 10(4), pp.

41209.

29. Elbaum-Garfinkle, S., Kim, Y., Szczepaniak, K., Chen, C. C., Eckmann, C. R., Myong, S.

and Brangwynne, C. P. (2015) 'The disordered P granule protein LAF-1 drives phase

separation into droplets with tunable viscosity and dynamics', *Proc Natl Acad Sci U S A,*

112(23), pp. 7189-94.

30. Faul, F., Erdfelder, E., Lang, A. G. and Buchner, A. (2007) 'G*Power 3: a flexible statistical

power analysis program for the social, behavioral, and biomedical sciences', *Behav Res*

Methods, 39(2), pp. 175-91.

31. Fouani, M., Basset, C. A., Mangano, G. D., Leone, L. G., Lawand, N. B., Leone, A. and

Barone, R. (2022) 'Heat Shock Proteins Alterations in Rheumatoid Arthritis', *Int J Mol*

Sci, 23(5).

32. Gallo Castro, D. and Martin, S. G. (2018) 'Differential GAP requirement for Cdc42-GTP

polarization during proliferation and sexual reproduction', *Journal of Cell Biology,*

217(12), pp. 4215-4229.

33. Gilks, N., Kedersha, N., Ayodele, M., Shen, L., Stoecklin, G., Dember, L. M. and Anderson,

P. (2004) 'Stress granule assembly is mediated by prion-like aggregation of TIA-1', *Mol*

Biol Cell, 15(12), pp. 5383-98.

34. Gomez-Pastor, R., Burchfiel, E. T. and Thiele, D. J. (2018) 'Regulation of heat shock

transcription factors and their roles in physiology and disease', *Nat Rev Mol Cell Biol,*

19(1), pp. 4-19.

35. Grousl, T., Ivanov, P., Frýdlová, I., Vasicová, P., Janda, F., Vojtová, J., Malínská, K.,

Malcová, I., Nováková, L., Janosková, D., Valásek, L. and Hasek, J. (2009) 'Robust heat

shock induces eIF2alpha-phosphorylation-independent assembly of stress granules

containing eIF3 and 40S ribosomal subunits in budding yeast, Saccharomyces

cerevisiae', *J Cell Sci,* 122(Pt 12), pp. 2078-88.

36. Gungor, B., Gombos, I., Crul, T., Ayaydin, F., Szabó, L., Török, Z., Mátés, L., Vígh, L. and

Horváth, I. (2014) 'Rac1 Participates in Thermally Induced Alterations of the

Cytoskeleton, Cell Morphology and Lipid Rafts, and Regulates the Expression of Heat

Shock Proteins in B16F10 Melanoma Cells', *PLOS ONE,* 9(2), pp. e89136.

37. Guzikowski, A. R., Chen, Y. S. and Zid, B. M. (2019) 'Stress-induced mRNP granules: Form

and function of processing bodies and stress granules', *Wiley Interdiscip Rev RNA,*

10(3), pp. e1524.

38. Hanyu, Y., Imai, K. K., Kawasaki, Y., Nakamura, T., Nakaseko, Y., Nagao, K., Kokubu, A.,

Ebe, M., Fujisawa, A., Hayashi, T., Obuse, C. and Yanagida, M. (2009)

'Schizosaccharomyces pombe cell division cycle under limited glucose requires Ssp1

kinase, the putative CaMKK, and Sds23, a PP2A-related phosphatase inhibitor', *Genes*

Cells, 14(5), pp. 539-54.

39. Haupt, A., Ershov, D. and Minc, N. (2018) 'A Positive Feedback between Growth and

Polarity Provides Directional Persistency and Flexibility to the Process of Tip Growth',

Current Biology, 28(20), pp. 3342-3351.e3.

40. Hauschild, A., Engel, G., Brenner, W., Glaser, R., Monig, H., Henze, E. and Christophers, E.

(1999) 'S100B protein detection in serum is a significant prognostic factor in metastatic

melanoma', *Oncology,* 56(4), pp. 338-44.

41. Heinrich, S., Geissen, E. M., Kamenz, J., Trautmann, S., Widmer, C., Drewe, P., Knop, M.,

Radde, N., Hasenauer, J. and Hauf, S. (2013) 'Determinants of robustness in spindle

assembly checkpoint signalling', *Nat Cell Biol,* 15(11), pp. 1328-39.

42. Hergovich, A. (2013) 'Regulation and functions of mammalian LATS/NDR kinases: looking

beyond canonical Hippo signalling', *Cell & Bioscience,* 3(1), pp. 32.

43. Hergovich, A., Kohler, R. S., Schmitz, D., Vichalkovski, A., Cornils, H. and Hemmings, B. A.

(2009) 'The MST1 and hMOB1 tumor suppressors control human centrosome

duplication by regulating NDR kinase phosphorylation', *Curr Biol,* 19(20), pp. 1692-702.

44. Hergovich, A., Lamla, S., Nigg, E. A. and Hemmings, B. A. (2007) 'Centrosome-associated

NDR kinase regulates centrosome duplication', *Mol Cell,* 25(4), pp. 625-34.

45. Hergovich, A., Stegert, M. R., Schmitz, D. and Hemmings, B. A. (2006) 'NDR kinases

regulate essential cell processes from yeast to humans', *Nature Reviews Molecular Cell*

Biology, 7(4), pp. 253-264.

46. Higa, M., Kita, A., Hagihara, K., Kitai, Y., Doi, A., Nagasoko, R., Satoh, R. and Sugiura, R.
(2015) 'Spatial control of calcineurin in response to heat shock in fission yeast', *Genes*
Cells, 20(2), pp. 95-107.
47. Hogan, D. J., Riordan, D. P., Gerber, A. P., Herschlag, D. and Brown, P. O. (2008) 'Diverse
RNA-binding proteins interact with functionally related sets of RNAs, suggesting an
extensive regulatory system', *PLoS Biol,* 6(10), pp. e255.
48. Hu, C., Yang, J., Qi, Z., Wu, H., Wang, B., Zou, F., Mei, H., Liu, J., Wang, W. and Liu, Q.
(2022) 'Heat shock proteins: Biological functions, pathological roles, and therapeutic
opportunities', *MedComm (2020),* 3(3), pp. e161.
49. Hyman, A. A., Weber, C. A. and Julicher, F. (2014) 'Liquid-liquid phase separation in
biology', *Annu Rev Cell Dev Biol,* 30, pp. 39-58.
50. Ingelfinger, D., Arndt-Jovin, D. J., Luhrmann, R. and Achsel, T. (2002) 'The human LSm1-7
proteins colocalize with the mRNA-degrading enzymes Dcp1/2 and Xrnl in distinct
cytoplasmic foci', *Rna,* 8(12), pp. 1489-501.
51. Jansen, J. M., Barry, M. F., Yoo, C. K. and Weiss, E. L. (2006) 'Phosphoregulation of Cbk1
is critical for RAM network control of transcription and morphogenesis', *J Cell Biol,*
175(5), pp. 755-66.
52. Jansen, J. M., Wanless, A. G., Seidel, C. W. and Weiss, E. L. (2009) 'Cbk1 regulation of the
RNA-binding protein Ssd1 integrates cell fate with translational control', *Curr Biol,*
19(24), pp. 2114-20.
53. Jung, M. K., Kim, K. Y., Lee, N. Y., Kang, Y. S., Hwang, Y. J., Kim, Y., Sung, J. J., McKee,
A., Kowall, N., Lee, J. and Ryu, H. (2013) 'Expression of taurine transporter (TauT) is
modulated by heat shock factor 1 (HSF1) in motor neurons of ALS', *Mol Neurobiol,*
47(2), pp. 699-710.
54. Kanda, Y., Satoh, R., Takasaki, T., Tomimoto, N., Tsuchiya, K., Tsai, C. A., Tanaka, T.,
Kyomoto, S., Hamada, K., Fujiwara, T. and Sugiura, R. (2021) 'Sequestration of the PKC
ortholog Pck2 in stress granules as a feedback mechanism of MAPK signaling in fission
yeast', *J Cell Sci,* 134(2).
55. Kang, H. (2021) 'Sample size determination and power analysis using the G*Power
software', *J Educ Eval Health Prof,* 18, pp. 17.
56. Kato, M., Han, T. W., Xie, S., Shi, K., Du, X., Wu, L. C., Mirzaei, H., Goldsmith, E. J.,
Longgood, J., Pei, J., Grishin, N. V., Frantz, D. E., Schneider, J. W., Chen, S., Li, L.,
Sawaya, M. R., Eisenberg, D., Tycko, R. and McKnight, S. L. (2012) 'Cell-free formation

of RNA granules: low complexity sequence domains form dynamic fibers within

hydrogels', *Cell,* 149(4), pp. 753-67.

57. Kedersha, N., Stoecklin, G., Ayodele, M., Yacono, P., Lykke-Andersen, J., Fritzler, M. J.,

Scheuner, D., Kaufman, R. J., Golan, D. E. and Anderson, P. (2005) 'Stress granules

and processing bodies are dynamically linked sites of mRNP remodeling', *J Cell Biol,*

169(6), pp. 871-84.

58. Kroschwald, S. and Alberti, S. (2017) 'Gel or Die: Phase Separation as a Survival Strategy',

Cell, 168(6), pp. 947-948.

59. Kroschwald, S., Maharana, S., Mateju, D., Malinovska, L., Nuske, E., Poser, I., Richter, D.

and Alberti, S. (2015) 'Promiscuous interactions and protein disaggregases determine

the material state of stress-inducible RNP granules', *Elife,* 4, pp. e06807.

60. Kurischko, C., Kim, H. K., Kuravi, V. K., Pratzka, J. and Luca, F. C. (2011) 'The yeast Cbk1

kinase regulates mRNA localization via the mRNA-binding protein Ssd1', *J Cell Biol,*

192(4), pp. 583-98.

61. Lee, C. F., Brangwynne, C. P., Gharakhani, J., Hyman, A. A. and Julicher, F. (2013) 'Spatial

organization of the cell cytoplasm by position-dependent phase separation', *Phys Rev*

Lett, 111(8), pp. 088101.

62. Leonhard, K. and Nurse, P. (2005) 'Ste20/GCK kinase Nak1/Orb3 polarizes the actin

cytoskeleton in fission yeast during the cell cycle', *Journal of Cell Science,* 118(5), pp.

1033-1044.

63. Lin, Y., Protter, D. S., Rosen, M. K. and Parker, R. (2015) 'Formation and Maturation of

Phase-Separated Liquid Droplets by RNA-Binding Proteins', *Mol Cell,* 60(2), pp. 208-19.

64. Liu, G. and Young, D. (2012) 'Conserved Orb6 Phosphorylation Sites Are Essential for

Polarized Cell Growth in Schizosaccharomyces pombe', *PLOS ONE,* 7(5), pp. e37221.

65. Lu, J., Li, H., Yu, D., Zhao, P. and Liu, Y. (2023) 'Heat stress inhibits the proliferation and

differentiation of myoblasts and is associated with damage to mitochondria', *Front Cell*

Dev Biol, 11, pp. 1171506.

66. Lv, H., Zhu, Y., Qiu, Y., Niu, L., Teng, M. and Li, X. (2015) 'Structural analysis of Dis3l2, an

exosome-independent exonuclease from Schizosaccharomyces pombe', *Acta*

Crystallogr D Biol Crystallogr, 71(Pt 6), pp. 1284-94.

67. Magliozzi, J. O. and Moseley, J. B. (2021) 'Pak1 kinase controls cell shape through

ribonucleoprotein granules', *Elife,* 10.

68. Mahboubi, H. and Stochaj, U. (2017) 'Cytoplasmic stress granules: Dynamic modulators of

cell signaling and disease', *Biochim Biophys Acta Mol Basis Dis,* 1863(4), pp. 884-895.

69. Malecki, M., Viegas, S. C., Carneiro, T., Golik, P., Dressaire, C., Ferreira, M. G. and
Arraiano, C. M. (2013) 'The exoribonuclease Dis3L2 defines a novel eukaryotic RNA
degradation pathway', *EMBO J,* 32(13), pp. 1842-54.
70. Mancini Lombardi, I., Palani, S., Meitinger, F., Darieva, Z., Hofmann, A., Sharrocks, A. D.
and Pereira, G. (2013) 'Lre1 directly inhibits the NDR/Lats kinase Cbk1 at the cell
division site in a phosphorylation-dependent manner', *Curr Biol,* 23(18), pp. 1736-45.
71. Marcelo, A., Koppenol, R., de Almeida, L. P., Matos, C. A. and Nóbrega, C. (2021) 'Stress
granules, RNA-binding proteins and polyglutamine diseases: too much aggregation?',
Cell Death & Disease, 12(6), pp. 592.
72. Matsuo, Y., Asakawa, K., Toda, T. and Katayama, S. (2006) 'A Rapid Method for Protein
Extraction from Fission Yeast', *Bioscience, Biotechnology, and Biochemistry,* 70(8), pp.
1992-1994.
73. Matsusaka, T., Hirata, D., Yanagida, M. and Toda, T. (1995) 'A novel protein kinase gene
ssp1+ is required for alteration of growth polarity and actin localization in fission yeast',
EMBO J, 14(14), pp. 3325-38.
74. Merlini, L., Khalili, B., Bendezú, F. O., Hurwitz, D., Vincenzetti, V., Vavylonis, D. and Martin,
S. G. (2016) 'Local Pheromone Release from Dynamic Polarity Sites Underlies Cell-Cell
Pairing during Yeast Mating', *Curr Biol,* 26(8), pp. 1117-25.
75. Miles, S., Li, L. H., Melville, Z. and Breeden, L. L. (2019) 'Ssd1 and the cell wall integrity
pathway promote entry, maintenance, and recovery from quiescence in budding yeast',
Mol Biol Cell, 30(17), pp. 2205-2217.
76. Millar, J. B., Buck, V. and Wilkinson, M. G. (1995) 'Pyp1 and Pyp2 PTPases
dephosphorylate an osmosensing MAP kinase controlling cell size at division in fission
yeast', *Genes Dev,* 9(17), pp. 2117-30.
77. Millward, T. A., Heizmann, C. W., Schafer, B. W. and Hemmings, B. A. (1998) 'Calcium
regulation of Ndr protein kinase mediated by S100 calcium-binding proteins', *Embo j,*
17(20), pp. 5913-22.
78. Millward, T. A., Hess, D. and Hemmings, B. A. (1999) 'Ndr Protein Kinase Is Regulated by
Phosphorylation on Two Conserved Sequence Motifs *', *Journal of Biological Chemistry,*
274(48), pp. 33847-33850.
79. Mir, S. S., Fiedler, D. and Cashikar, A. G. (2009) 'Ssd1 is required for thermotolerance and
Hsp104-mediated protein disaggregation in Saccharomyces cerevisiae', *Mol Cell Biol,*
29(1), pp. 187-200.

80. Mutavchiev, Delyan R., Leda, M. and Sawin, Kenneth E. (2016) 'Remodeling of the Fission
Yeast Cdc42 Cell-Polarity Module via the Sty1 p38 Stress-Activated Protein Kinase
Pathway', *Current Biology,* 26(21), pp. 2921-2928.

81. Napoletano, F., Occhi, S., Calamita, P., Volpi, V., Blanc, E., Charroux, B., Royet, J. and
Fanto, M. (2011) 'Polyglutamine Atrophin provokes neurodegeneration in Drosophila by
repressing fat', *Embo j,* 30(5), pp. 945-58.

82. Nguyen, A. N. and Shiozaki, K. (1999) 'Heat-shock-induced activation of stress MAP kinase
is regulated by threonine- and tyrosine-specific phosphatases', *Genes Dev,* 13(13), pp.
1653-63.

83. Nilsson, D. and Sunnerhagen, P. (2011) 'Cellular stress induces cytoplasmic RNA granules
in fission yeast', *RNA,* 17(1), pp. 120-33.

84. Nunez, I., Rodriguez Pino, M., Wiley, D. J., Das, M. E., Chen, C., Goshima, T., Kume, K.,
Hirata, D., Toda, T. and Verde, F. (2016) 'Spatial control of translation repression and
polarized growth by conserved NDR kinase Orb6 and RNA-binding protein Sts5', *Elife,* 5.

85. Nuñez, I., Rodriguez Pino, M., Wiley, D. J., Das, M. E., Chen, C., Goshima, T., Kume, K.,
Hirata, D., Toda, T. and Verde, F. (2016) 'Spatial control of translation repression and
polarized growth by conserved NDR kinase Orb6 and RNA-binding protein Sts5', *eLife,*
5, pp. e14216.

86. Panas, M. D., Ivanov, P. and Anderson, P. (2016) 'Mechanistic insights into mammalian
stress granule dynamics', *Journal of Cell Biology,* 215(3), pp. 313-323.

87. Parker, B. W., Gogl, G., Bálint, M., Hetényi, C., Reményi, A. and Weiss, E. L. (2020)
'Ndr/Lats Kinases Bind Specific Mob-Family Coactivators through a Conserved and
Modular Interface', *Biochemistry,* 59(17), pp. 1688-1700.

88. Patel, A., Lee, H. O., Jawerth, L., Maharana, S., Jahnel, M., Hein, M. Y., Stoynov, S.,
Mahamid, J., Saha, S., Franzmann, T. M., Pozniakovski, A., Poser, I., Maghelli, N.,
Royer, L. A., Weigert, M., Myers, E. W., Grill, S., Drechsel, D., Hyman, A. A. and Alberti,
S. (2015) 'A Liquid-to-Solid Phase Transition of the ALS Protein FUS Accelerated by
Disease Mutation', *Cell,* 162(5), pp. 1066-77.

89. Pawlik, A., Nowak, J. M., Grzanka, D., Gackowska, L., Michalkiewicz, J. and Grzanka, A.
(2013) 'Hyperthermia induces cytoskeletal alterations and mitotic catastrophe in p53-
deficient H1299 lung cancer cells', *Acta Histochem,* 115(1), pp. 8-15.

90. Ramaswami, M., Taylor, J. P. and Parker, R. (2013) 'Altered ribostasis: RNA-protein
granules in degenerative disorders', *Cell,* 154(4), pp. 727-36.

91. Riback, J. A., Katanski, C. D., Kear-Scott, J. L., Pilipenko, E. V., Rojek, A. E., Sosnick, T. R.
and Drummond, D. A. (2017) 'Stress-Triggered Phase Separation Is an Adaptive,
Evolutionarily Tuned Response', *Cell,* 168(6), pp. 1028-1040 e19.
92. Robinson, S. R., Oliver, A. W., Chevassut, T. J. and Newbury, S. F. (2015) 'The 3' to 5'
Exoribonuclease DIS3: From Structure and Mechanisms to Biological Functions and
Role in Human Disease', *Biomolecules,* 5(3), pp. 1515-39.
93. Ross, D. T., Scherf, U., Eisen, M. B., Perou, C. M., Rees, C., Spellman, P., Iyer, V., Jeffrey,
S. S., Van de Rijn, M., Waltham, M., Pergamenschikov, A., Lee, J. C., Lashkari, D.,
Shalon, D., Myers, T. G., Weinstein, J. N., Botstein, D. and Brown, P. O. (2000)
'Systematic variation in gene expression patterns in human cancer cell lines', *Nat Genet,*
24(3), pp. 227-35.
94. Sakuno, T., Araki, Y., Ohya, Y., Kofuji, S., Takahashi, S., Hoshino, S. and Katada, T. (2004)
'Decapping reaction of mRNA requires Dcp1 in fission yeast: its characterization in
different species from yeast to human', *J Biochem,* 136(6), pp. 805-12.
95. Salat-Canela, C., Carmona, M., Martín-García, R., Pérez, P., Ayté, J. and Hidalgo, E. (2021)
'Stress-dependent inhibition of polarized cell growth through unbalancing the GEF/GAP
regulation of Cdc42', *Cell Reports,* 37(5), pp. 109951.
96. Salat-Canela, C., Pérez, P., Ayté, J. and Hidalgo, E. (2023) 'Stress-induced cell
depolarization through the MAP kinase–Cdc42 axis', *Trends in Cell Biology,* 33(2), pp.
124-137.
97. Satoh, R., Tanaka, A., Kita, A., Morita, T., Matsumura, Y., Umeda, N., Takada, M., Hayashi,
S., Tani, T., Shinmyozu, K. and Sugiura, R. (2012) 'Role of the RNA-binding protein
Nrd1 in stress granule formation and its implication in the stress response in fission
yeast', *PLoS One,* 7(1), pp. e29683.
98. Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T.,
Preibisch, S., Rueden, C., Saalfeld, S., Schmid, B., Tinevez, J.-Y., White, D. J.,
Hartenstein, V., Eliceiri, K., Tomancak, P. and Cardona, A. (2012) 'Fiji: an open-source
platform for biological-image analysis', *Nature Methods,* 9(7), pp. 676-682.
99. Schwamborn, J. C. and Püschel, A. W. (2004) 'The sequential activity of the GTPases
Rap1B and Cdc42 determines neuronal polarity', *Nature Neuroscience,* 7(9), pp. 923-
929.
100. Shiozaki, K. and Russell, P. (1996) 'Conjugation, meiosis, and the osmotic stress response
are regulated by Spc1 kinase through Atf1 transcription factor in fission yeast', *Genes
Dev,* 10(18), pp. 2276-88.

101. Shiozaki, K., Shiozaki, M. and Russell, P. (1998) 'Heat stress activates fission yeast
Spc1/Styl MAPK by a MEKK-independent mechanism', *Mol Biol Cell,* 9(6), pp. 1339-49.

102. Singh, D., Odedra, D., Lehmann, C. and Pohl, C. (2016) 'Acute heat shock leads to cortical
domain internalization and polarity loss in the C. elegans embryo', *Genesis,* 54(4), pp.
220-8.

103. Smith, D. A., Toone, W. M., Chen, D., Ba¨hler, J. r., Jones, N., Morgan, B. A. and Quinn, J.
(2002) 'The Srk1 Protein Kinase Is a Target for the Sty1 Stress-activated MAPK in
Fission Yeast*', *Journal of Biological Chemistry,* 277(36), pp. 33411-33421.

104. Snell, V. and Nurse, P. (1994) 'Genetic analysis of cell morphogenesis in fission yeast--a
role for casein kinase II in the establishment of polarized growth', *Embo j,* 13(9), pp.
2066-74.

105. Stankiewicz, T. R. and Linseman, D. A. (2014) 'Rho family GTPases: key players in
neuronal development, neuronal survival, and neurodegeneration', *Frontiers in Cellular*
Neuroscience, 8.

106. Stein, V., Blank-Landeshammer, B., Müntjes, K., Märker, R., Teichert, I., Feldbrügge, M.,
Sickmann, A. and Kück, U. (2020) 'The STRIPAK signaling complex regulates
dephosphorylation of GUL1, an RNA-binding protein that shuttles on endosomes', *PLoS*
Genet, 16(9), pp. e1008819.

107. Sun, L., Liu, L., Song, C., Wang, Y. and Jin, Q.-w. (2024) 'Heat stress-induced activation of
MAPK pathway attenuates Atf1-dependent epigenetic inheritance of heterochromatin in
fission yeast', *eLife,* 13, pp. e90525.

108. Toda, T., Niwa, H., Nemoto, T., Dhut, S., Eddison, M., Matsusaka, T., Yanagida, M. and
Hirata, D. (1996) 'The fission yeast sts5+ gene is required for maintenance of growth
polarity and functionally interacts with protein kinase C and an osmosensing MAP-kinase
pathway', *J Cell Sci,* 109 (Pt 9), pp. 2331-42.

109. Toone, W. M. and Jones, N. (1998) 'Stress-activated signalling pathways in yeast', *Genes*
Cells, 3(8), pp. 485-98.

110. Vaggi, F., Dodgson, J., Bajpai, A., Chessel, A., Jordan, F., Sato, M., Carazo-Salas, R. E.
and Csikasz-Nagy, A. (2012) 'Linkers of cell polarity and cell cycle regulation in the
fission yeast protein interaction network', *PLoS Comput Biol,* 8(10), pp. e1002732.

111. van Dijk, E., Cougot, N., Meyer, S., Babajko, S., Wahle, E. and Seraphin, B. (2002)
'Human Dcp2: a catalytically active mRNA decapping enzyme located in specific
cytoplasmic structures', *Embo j,* 21(24), pp. 6915-24.

112. Velichko, A. K., Markova, E. N., Petrova, N. V., Razin, S. V. and Kantidze, O. L. (2013)
'Mechanisms of heat shock response in mammals', *Cell Mol Life Sci,* 70(22), pp. 4229-
41.

113. Verde, F., Mata, J. and Nurse, P. (1995) 'Fission yeast cell morphogenesis: identification of
new genes and analysis of their role during the cell cycle', *J Cell Biol,* 131(6 Pt 1), pp.
1529-38.

114. Verde, F., Wiley, D. J. and Nurse, P. (1998) 'Fission yeast orb6, a ser/thr protein kinase
related to mammalian rho kinase and myotonic dystrophy kinase, is required for
maintenance of cell polarity and coordinates cell morphogenesis with the cell cycle', *Proc*
Natl Acad Sci U S A, 95(13), pp. 7526-31.

115. Versele, M. and Thevelein, J. M. (2001) 'Lre1 affects chitinase expression, trehalose
accumulation and heat resistance through inhibition of the Cbk1 protein kinase in
Saccharomyces cerevisiae', *Mol Microbiol,* 41(6), pp. 1311-26.

116. Vichalkovski, A., Gresko, E., Cornils, H., Hergovich, A., Schmitz, D. and Hemmings, B. A.
(2008) 'NDR kinase is activated by RASSF1A/MST1 in response to Fas receptor
stimulation and promotes apoptosis', *Curr Biol,* 18(23), pp. 1889-95.

117. Vjestica, A., Zhang, D., Liu, J. and Oliferenko, S. (2013) 'Hsp70-Hsp40 Chaperone
Complex Functions in Controlling Polarized Growth by Repressing Hsf1-Driven Heat
Stress-Associated Transcription', *PLOS Genetics,* 9(10), pp. e1003886.

118. Wang, Z., Jiao, X., Carr-Schmid, A. and Kiledjian, M. (2002) 'The hDcp2 protein is a
mammalian mRNA decapping enzyme', *Proc Natl Acad Sci U S A,* 99(20), pp. 12663-8.

119. Wen, W. L., Stevenson, A. L., Wang, C. Y., Chen, H. J., Kearsey, S. E., Norbury, C. J.,
Watt, S., Bahler, J. and Wang, S. W. (2010) 'Vgl1, a multi-KH domain protein, is a novel
component of the fission yeast stress granules required for cell survival under thermal
stress', *Nucleic Acids Res,* 38(19), pp. 6555-66.

120. Wheeler, J. R., Matheny, T., Jain, S., Abrisch, R. and Parker, R. (2016) 'Distinct stages in
stress granule assembly and disassembly', *Elife,* 5.

121. Wilkinson, M. G., Samuels, M., Takeda, T., Toone, W. M., Shieh, J. C., Toda, T., Millar, J.
B. and Jones, N. (1996) 'The Atf1 transcription factor is a target for the Sty1 stress-
activated MAP kinase pathway in fission yeast', *Genes Dev,* 10(18), pp. 2289-301.

122. Wu, J., Liu, T., Rios, Z., Mei, Q., Lin, X. and Cao, S. (2017) 'Heat Shock Proteins and
Cancer', *Trends in Pharmacological Sciences,* 38(3), pp. 226-256.

123. Yamada, H., Ohmiya, R., Yamamoto, E., Aiba, H. and Mizuno, T. (1997) 'Characterization
of multicopy suppressor genes that complement a defect in the Wis1-Sty1 MAP kinase

cascade involved in stress responses in Schizosaccharomyces pombe', *J Gen Appl*

Microbiol, 43(4), pp. 209-215.

124. Yang, R., Kong, E., Jin, J., Hergovich, A. and Puschel, A. W. (2014) 'Rassf5 and Ndr

kinases regulate neuronal polarity through Par3 phosphorylation in a novel pathway', *J*

Cell Sci, 127(Pt 16), pp. 3463-76.

125. Zayani, Z., Matinahmadi, A., Tavakolpournegari, A. and Bidooki, S. H. (2025) 'Exploring

Stressors: Impact on Cellular Organelles and Implications for Cellular Functions',

Stresses, 5(2). DOI: 10.3390/stresses5020026.

126. Zhang, L., Tang, F., Terracciano, L., Hynx, D., Kohler, R., Bichet, S., Hess, D., Cron, P.,

Hemmings, B. A., Hergovich, A. and Schmitz-Rohmer, D. (2015) 'NDR functions as a

physiological YAP1 kinase in the intestinal epithelium', *Curr Biol,* 25(3), pp. 296-305.

1140

1141

1127   **Main Figures and Legends**

1128

Figure 1

Figure 1: Orb6 activity is negatively regulated by heat stress.

(A) Gef1-3YFP localizes the cell membrane upon heat exposure at 36ºC or 42ºC for 30
minutes (arrows). Scale bar, 5 µm. (B) Quantification of percentage of cells that display
ectopic Gef1-3YFP localization from A based on three independent experiments. Data
presented as mean ± SD, *p* values are determined by one-way ANOVA with Tukey's HSD
test p ≤ 0.01, **. n = number of cells quantified. (C) Orb6 activity, as measured by Gef1-
S112 phosphorylation, decreases after 30 minutes at 36ºC or 42ºC. (D) Quantification of
pGef1-S112/tGef1 from C upon temperature stress exposure based on three independent
experiments. Data are presented as in B, p ≤ 0.05, *; p ≤ 0.001, ***. (E) *Sts5-3xGFP*
pabp1-DsRed cells were exposed to 36ºC or 42ºC for 30 minutes. At 36ºC, Sts5-3xGFP
formed cytoplasmic puncta (d, arrows) but no stress granule formation was visible (f). At
42ºC, Sts5 partially co-localized with the stress granule marker Pabp1-DsRed (g-i,
arrows). Scale bar, 5 µm. (F) Quantification of average number of Sts5 puncta present in
each cell from E based on three independent experiments. Data are presented as in B, p
≤ 0.01, **; p ≤ 0.001, ***, p ≤ 0.0001, ****. n = number of cells quantified. (G) Quantification
of average number of Sts5 puncta that colocalized by stress granule marker Pabp1 in
each cell from E based on three independent experiments. Data are presented as in B, p
≤ 0.0001, ****. n = number of cells quantified.

1147

1148

A

0% Glucose	ssp1 mRNA	Sts5-3xGFP	Pabp1-DsRed	Merge
	a	b	c	d

B

orb6-as2 pabp1-DsRed **C**

	Control	- Glucose
sts5⁺ pabp1-DsRed	a	b
sts5Δ pabp1-DsRed	c	d

	25°C	42°C
DMSO	a	b
1-NA-PP1	c	d

D

Stress Granules/Cell (-Glu)

***

sts5⁺   sts5Δ

E

Stress Granules/Cell (42°C)

***

DMSO   1-NA-PP1

Figure 2

Figure 2: Orb6 and Sts5 regulate stress granule formation.

(A) RNA FISH visualization of ssp1 mRNA in fixed cells cultured for 20 minutes in minimal
medium containing 0% glucose. Hybridization of RNA was performed with 20-mer DNA
oligonucleotides (Stellaris) labeled with Quasar 705 fluorochromes. ssp1 mRNA co-
localized with Sts5-3xGFP and the stress granule marker Pabp1-DsRed (arrows). Images
are deconvolved projections from Z-stacks (12 slices) separated by a step size of 0.3 μM
(Bar = 5 μM). (B) *pabp1-DsRed* and *sts5Δ pabp1-DsRed* cells were cultured in EMM
medium with (a, c) or without (b, d) glucose for 20 minutes. In untreated controls, the
stress granule marker Pabp1-DsRed remained diffuse throughout the cytoplasm.
Following glucose deprivation, *sts5Δ* cells formed less stress granules than the control
(b, d). Representative images are deconvolved projections from Z-stacks separated by a
step size of 0.3 μM (Bar = 5 μM). (C) Inhibition of Orb6 primes cells for the formation of
stress granules. *orb6-as2 pabp1-DsRed* cells were cultured in minimal medium
containing adenine, and 15 minutes prior to heat shock 1-NA-PP1 inhibitor was added,
or DMSO for controls. In cells maintained at 25ºC, Pabp1-DsRed remained diffuse within
the cytoplasm (a, c). However, in cells that were heat shocked for 20 minutes at 42ºC (15
minutes post addition of 1-NA-PP1), there was an increase in stress granule formation
compared to the DMSO control (b, d). Representative images are deconvolved
projections from Z-stacks separated by a step size of 0.3 μM (Bar = 5 μM). (D) The
number of stress granules were quantified for experiment represented in B (n=90
cells/strain/condition total, performed in biological triplicate), and there was a statistically
significant decrease in stress granule formation for the *sts5Δ* mutant as compared to the
control (***p<0.001; unpaired t-test). (E) The increase in stress granule formation from C
was statistically significant and performed in biological triplicates (n=90
cells/strain/condition total; ***p<0.001, unpaired t-test).
1174

1175

Figure 3

Figure 3: Orb6 and Sts5 modulate survival following heat stress.

(A) *Wild-type, sts5Δ, sts5-HA,* and *sts5-S86A-HA* cells were grown for 6 hours at either 25ºC or 36.5ºC. Following prolonged heat stress at 36.5ºC, s*ts5Δ* cells displayed decreased survival compared to wild type, whereas *sts5-S86A-HA* showed increased survival compared to the *sts5-HA* control. Data are presented as mean ± SD, p values determined by one-way ANOVA with Tukey's HSD test p ≤ 0.05, *; p ≤ 0.01, **; p ≤ 0.001, ***. (B) The experiment in A was repeated, but cells were instead incubated for 15 minutes at 48ºC temperature, while untreated controls were incubated at 25ºC. Again, *sts5Δ* cells exhibit decreased survival, whereas the *sts5-S86A-HA* mutant exhibited increased survival. Data are presented as mean ± SD, p values determined by one-way ANOVA with Tukey's HSD test p ≤ 0.05, *; p ≤ 0.001, ***. (C) In the *sts5-S86A-3xYFP* mutant, the number of Sts5 granules were increased at temperatures in which the *sts5-3xYFP* control did not form granules. (D) Quantification from results obtained in C. There is a statistically significant increase in average puncta per cell for the *sts5-S86A-3xYFP* strain as compared to the control for cells incubated between 25ºC and 35.5ºC Data are presented as mean ± SD, p values determined by two-way ANOVA with Tukey's HSD test p ≤ 0.01, **; p ≤ 0.001, ***. n=60 cells/strain performed in biological triplicates (E) *Wild-type* and *orb6-as2* strains were cultured in the presence (+Thiamine) or absence of thiamine (-Thiamine), heat shocked, and plated. Inhibition of Orb6 results in a 17-fold increase in survival as compared to the untreated control. Data are presented as mean ± SD, p values determined by two-way ANOVA with Tukey's HSD test p ≤ 0.0001, ****.

Figure 4

Figure 4: The distribution of Sts5, Gef1, and active Cdc42 is temperature sensitive and is modulated by Sty1 in a temperature specific manner.

(A) Sts5-3YFP forms puncta after 30 minutes at 36°C and 42°C (arrows). In the absence of *sty1,* Sts5-3YFP only clusters at 42°C. Images are a max projection Z-stack of 6 images separated by a step-size of 0.3 μm. Scale bar, 5 μm. (B) Quantification of average number of Sts5 puncta present in each cell from A based on three independent experiments. Data are presented as mean ± SD, p values determined by two-way ANOVA with Tukey's HSD test $p \leq 0.001$, ***; $p \leq 0.0001$, ****. n = number of cells quantified. (C) Active Cdc42 forms exploratory patches upon exposure to 36°C and 42°C when *sty1* is present but only forms patches at 42°C when *sty1* is absent (arrows). Active Cdc42 is localized at the cell tips in control cells at 25°C and in *sty1Δ* cells at 25°C and 36°C (brackets) Images are a max projection Z-stack of 6 images separated by a step-size of 0.3 μm. Scale bar, 5 μm. (D) Quantification of percentage of cells that display ectopic CRIB-GFP localization from C based on three independent experiments. Data presented as in D, $p \leq 0.0001$, ****. n = number of cells quantified. (E) Gef1-3YFP localizes to the cell membrane at 36°C and 42°C (arrows). Loss of *sty1* leads to Gef1-3YFP localization to the cell membrane only at 42°C. Scale bar, 5 μm. (F) Quantification of percentage of cells that display ectopic Gef1-3YFP localization from B based on three independent experiments. Data presented as in D, $p \leq 0.0001$, ****. n = number of cells quantified. (G) Gef1-S112 phosphorylation by Orb6 decreases after 30 minutes at 36°C or 42°C. In *sty1Δ* deletion cells, pGef1-S112 remains constant at 36°C but decreases at 42°C. Atf1 levels increase upon exposure to 36°C or 42°C for 30 minutes in control cells. β-Actin was used as a loading control. (H) Quantification of pGef1-S112/tGef1 from A upon temperature stress exposure in control or *sty1Δ* deletion mutant cells based on three independent experiments. Data are presented as mean ± SD, *p* values are determined by two-way ANOVA with Tukey's HSD test $p \leq 0.001$, ***; $p \leq 0.0001$, ****. (I) Quantification of Atf1/Actin from A upon temperature stress in control or *sty1Δ* deletion mutant cells based on three independent experiments. Data are presented as mean ± SD, *p* values are determined by two-way ANOVA with Tukey's HSD test $p \leq 0.001$, ***; $p \leq 0.0001$, ****.

Figure 5

Figure 5: The phosphorylation state of the Orb6 C-terminal hydrophobic motif is
temperature sensitive.

(A) Protein domain map of Orb6 kinase highlighting two phosphosites: S291 in the
activation segment and T456 in the hydrophobic motif. NTR:N-terminal region; KD: kinase
domain; AS: activation segment; HM: hydrophobic motif. (B) Dot plot showing specificity
of phosphor-Orb6-T456 antibody. (C) Orb6-456 phosphorylation decreases when
temperature sensitive mutant of *nak1* (*nak1-ts)* is exposed to restrictive temperature for
3 hours. (D) Quantification of pOrb6-T456/tOrb6 from C based on three independent
experiments. Data are presented as mean ± SD, *p* values are determined by two-way
ANOVA with Tukey's HSD test $p \leq 0.05$, *; $p \leq 0.01$, **; $p \leq 0.001$, ***. (E) Orb6-T456
phosphorylation decreases after 30 minutes at 36°C or 42°C in control or *sty1Δ* deletion
mutant cells. Orb6-T456 phosphorylation is higher in *sty1Δ* mutants. β-Actin was used as
a loading control. (F) Quantification of pOrb6-T456/tOrb6 from A based on three
independent experiments. Data are presented as mean ± SD, *p* values are determined
by two-way ANOVA with Tukey's HSD test $p \leq 0.01$, **; $p \leq 0.001$, ***.
1245

1246

A Optimal Temperature

B

Polarized Cdc42
Translation of Sts5-bound mRNAs
Polarized cell growth

Exploratory Cdc42 dynamics
Translational repression of Sts5-bound mRNAs
Increased heat stress resilience

● Cdc42-GTP
● Sts5 Granules

Figure 6

Figure 6: Regulation of NDR kinase Orb6 by temperature and MAP kinase Sty1.

(A) Orb6 kinase activity at optimal, not stressful temperatures (25°C) promotes the
canonical Cdc42 polarity module (via regulation of RNA Binding protein Sts5), and
polarized cell growth. (B) Upon heat stress, MAP kinase Sty1 is strongly activated and
inhibits Orb6 kinase activity, thereby promoting exploratory Cdc42 dynamics, Sts5
granule assembly, translational repression of Sts5-associated mRNAs, and increased
heat stress resilience. The induction of exploratory Cdc42 dynamics also involves direct
phosphorylation of Orb6 substrates by Sty1 Kinase (see Discussion). Inhibition of Orb6
kinase activity during heat stress also engages Sty1-independent regulatory
mechanisms, likely affecting other components of the MOR (Morphogenesis Orb6
network) pathway. The activity of MAP kinase Sty1 lowers the threshold temperature that
triggers Orb6 inhibition, thereby rendering Orb6 more temperature sensitive and
promoting thermotolerance.
1260
1261