1 **A fungal powdery mildew pathogen induces extensive local and marginal**

2 **systemic changes in the *Arabidopsis thaliana* microbiota**

4 **Paloma Durán [1,2], Anja Reinstädler [3], Anna Lisa Rajakrut [1], Masayoshi Hashimoto [4], Ruben**

5 **Garrido-Oter [1,2,5], Paul Schulze-Lefert [1,2,5], Ralph Panstruga [3,5]**

6 [1] Max Planck Institute for Plant Breeding Research, Department of Plant-Microbe

7 Interactions, Carl-von-Linné-Weg 55, 50829 Cologne, Germany

8 [2] Cluster of Excellence on Plant Sciences, 40225 Düsseldorf, Germany.

9 [3] RWTH Aachen University, Institute for Biology I, Unit of Plant Molecular Cell Biology,

10 Worringerweg 1, 52056 Aachen, Germany

11 [4] Graduate School of Agricultural and Life Sciences, The University of Tokyo, 113-8657 Tokyo,

12 Japan.

13 [5] co-corresponding authors

Summary

• Powdery mildew is a foliar disease caused by epiphytically growing obligate biotrophic

ascomycete fungi. How powdery mildew colonization affects host resident microbial

communities locally and systemically remains poorly explored.

• We performed powdery mildew (*Golovinomyces orontii*) infection experiments with

Arabidopsis thaliana grown in either natural soil or a gnotobiotic system and studied the

influence of pathogen invasion into standing natural multi-kingdom or synthetic bacterial

communities (SynComs).

• We found that after infection of soil-grown plants, *G. orontii* outcompetes numerous

resident leaf-associated fungi. We further detected a significant shift in foliar but not root-

associated bacterial communities in this setup. Pre-colonization of germ-free *A. thaliana*

leaves with a bacterial leaf-SynCom, followed by *G. orontii* invasion, induced an overall

similar shift in the foliar bacterial microbiota and minor changes in the root-associated

bacterial assemblage. However, a standing root SynCom in root samples remained robust

against foliar infection with *G. orontii*. Although pathogen growth was unaffected by the leaf

SynCom, fungal infection caused a more than two-fold increase in leaf bacterial load.

• Our findings indicate that *G. orontii* infection affects mainly microbial communities in local

plant tissue, possibly driven by pathogen-induced changes in source-sink relationships and

host immune status.

Keywords:

Arabidopsis thaliana, *Golovinomyces orontii*, gnotobiotic plant system, microbial multi-

kingdom interactions, plant microbiota, powdery mildew, synthetic community

Abbreviations

ASV   Amplificon sequence variant

ITS   Internal transcribed spacer

PCoA   Principal coordinates analysis

RA   Relative abundance

SynCom   Synthetic community

Introduction

Unlike plants grown under germ-free laboratory conditions, healthy plants in nature live in association and interact with a multitude of microorganisms belonging to several microbial classes, such as bacteria, fungi, oomycetes and protists, collectively called the plant microbiota (Bulgarelli *et al.*, 2013). Marker gene amplicon sequencing has served as an important tool for taxonomic profiling and quantitative surveys of microbial assemblages associated with different plant organs over a range of environmental and experimental conditions, revealing community composition and major factors explaining community structure (Hacquard & Schadt, 2015; Thiergart *et al.*, 2020). Root-associated bacterial assemblages, assessed by amplicon sequencing of the *16S* rRNA marker gene, are defined by a specific subset of bacteria that originate mainly from the highly diverse soil biota. These communities are characterized by a robust taxonomic pattern at the phylum rank, which comprises the dominant Proteobacteria as well as Actinobacteria, Bacteroidetes, and Firmicutes (Bulgarelli *et al.*, 2012; Lundberg *et al.*, 2012; Hacquard & Schadt, 2015). The main factors governing differences in root assemblages are, in order of importance, soil type (edaphic factors such as pH), plant species/genotype, and plant age (Lauber *et al.*, 2009; Hacquard *et al.*, 2015; Müller *et al.*, 2016; Finkel *et al.*, 2017). For leaf-associated bacterial assemblages, the seeding source is less defined because microbiota members can originate from aerosols, insects, and soil as well as upward microbial migration from the root (Vorholt, 2012; Müller *et al.*, 2016). Although root- and leaf-associated microbiota share a similar phylum-level taxonomic composition, the overall community structure in leaves is subject to larger fluctuations. Similar to bacterial communities, plant-associated fungi have been studied at the community level by amplicon sequencing of the internal transcribed spacer (ITS) region between the small- and large-subunit rRNA genes or the *18S* rRNA gene (Bazzicalupo *et al.*, 2013). Plant-associated fungal assemblages are dominated by members belonging to the Ascomycota and Basidiomycota phyla, and these communities display more stochastic variation compared with the bacterial microbiota, with biogeography as the strongest explanatory factor (Coleman-Derr *et al.*, 2016; Gao *et al.*, 2020; Thiergart *et al.*, 2020). In fact, taxonomically structured microbial communities with similar taxa on each plant individual have been described until now only for the bacterial root microbiota (Lundberg *et al.*, 2012).

Systematic establishment of plant-derived microbial culture collections of the plant

microbiota has enabled microbiota reconstitution experiments with germ-free plants and

taxonomically representative synthetic communities (SynComs) that can be used to address

principles underlying community assembly and proposed microbiota functions under

defined laboratory environments. This experimental approach has proven critical to the

advancement of microbiota research (Bai *et al.*, 2015; Lebeis *et al.*, 2015; Durán *et al.*, 2018;

Zhang *et al.*, 2019a). For example, more than 400 leaf- and root-derived bacterial

commensals have been isolated from healthy *A. thaliana* grown in natural soil, comprising

35 bacterial families belonging to the aforementioned four phyla (Bai *et al.*, 2015). This

microbiota culture collection represents the majority of bacterial taxa that are detectable by

culture-independent *16S* rRNA gene community profiling in the *A. thaliana* phyllo- and

rhizosphere. Microbiota reconstitution experiments using SynComs from this collection

revealed that the bacterial root microbiota provides indirect protection to its host against

soil-borne and root-associated harmful fungi and that this protection is essential for plant

survival (Duran et al., 2018). Similar reconstitution experiments have shown that bacterial

root commensals are necessary for iron nutrition of *A. thaliana* in naturally occurring

calcareous soils, where poor bioavailability of this soil mineral nutrient limits plant growth

(Harbort *et al.*, 2020).

Relatively little is known about how the plant microbiota responds to pathogen invasion. The

oomycete leaf pathogen *Albugo* has strong effects on epiphytic and endophytic bacterial

colonization in *A. thaliana*. Specifically, α-diversity decreased and β-diversity stabilized in the

presence of *Albugo* infection in leaves, whereas they otherwise varied between plants (Agler

et al., 2016). The effect of *Alb. laibachii* on leaf-associated fungal communities were less

consistent and not as clear. Upon foliar defense activation by the downy mildew oomycete

pathogen *Hyaloperonospora arabidopsidis*, the host *A. thaliana* specifically promotes three

bacterial species in the rhizosphere, namely *Xanthomonas*, *Microbacterium*, and

Stenotrophomonas sp., respectively (Berendsen *et al.*, 2018). Although separately these

bacteria did not affect the host significantly, together they induced systemic resistance

against downy mildew and promoted growth of the plant.

The analysis of powdery mildew-induced changes in plant leaf microbiota so far rests on field

studies with powdery mildew-infected leaf samples (diseased leaves) in comparison to

healthy/less infected leaves in Japanese spindle (*Euonymous japonicus*) (Zhang *et al.*, 2019b),

pumpkin (*Cucurbita moschata*) (Zhang *et al.*, 2018) and English oak (*Quercus robur*)

(Jakuschkin *et al.*, 2016) (reviewed in (Panstruga & Kuhn, 2019)). In the *E. japonicus* study,

the authors noticed a reduction in bacterial and fungal diversity, associated with a general

decrease in relative abundance (RA) at the genus level (Zhang *et al.*, 2019b). Similarly, the

richness and diversity of the fungal community was found to be reduced in pumpkin leaves

heavily infected by powdery mildew (*Podospharea* sp.) (Zhang *et al.*, 2018). Marked changes

in the composition of foliar fungal and bacterial communities were also observed in *Erysiphe*

alphitoides-colonized oak (Jakuschkin *et al.*, 2016). However, all these studies rely on field

samples and natural powdery mildew infections in fluctuating conditions, which complicates

deconvolution of microbiota changes caused by changes in environmental factors from

those driven by pathogen infection.

Here, we examined the effect of controlled powdery mildew (*G. orontii*) infection on the

structure of *A. thaliana* leaf and root microbiota in either soil-grown plants or a gnotobiotic

plant system pre-treated with defined root- or leaf SynComs. In both settings, we found

major powdery mildew-induced shifts in the composition of the local (foliar) assemblages of

fungal (natural soil) and bacterial (natural soil and sterile conditions) communities. In the

case of the leaf SynCom, this shift was also associated with a marked increase in bacterial

load. Apart from these major changes in the phyllosphere, we also observed a minor

systemic effect on the structure of the bacterial root microbiota in conjunction with the leaf

SynCom and powdery mildew challenge.

Materials and Methods

Plant material

A. thaliana Col-0 wild-type (Arabidopsis stock centre accession N60000) was used as model plant system. Seeds were surface-sterilized by treating them with 70% for 20 min and drying them under a sterile hood. Subsequently, the seeds were stratified overnight at 4 °C.

Microbial strains

The *At*-SPHERE bacterial strains used in this study have been previously reported (Bai *et al.*, 2015) and are summarized in Table S1. *G. orontii* (isolate MPIPZ) was used as powdery mildew infection agent. *G. orontii* was regularly propagated every week on 4- to 5-week-old supersuceptible *A. thaliana eds1* plants. Powdery mildew inoculation was conducted by leaf-to-leaf contact of healthy plants with rosette leaves of heavily infected *eds1* plants as reported previously (Acevedo-Garcia *et al.*, 2017).

Natural soil experiment

A.thaliana Col-0 seeds were sown into 7x7 greenhouse pots filled with Cologne Agricultural Soil (CAS) batch 12 and grown under short-day greenhouse conditions for 6.5 weeks (12 pots containing 5 seeds each). Then, half of the plants were inoculated with *G. orontii* and all pots were transferred to a growth chamber (day: 21 °C, 10 h light; night: 19 °C; 70% humidity). At 11 days post inoculation (dpi), bulk soil, rhizosphere, root and leaf samples were harvested as previously described (Figure 1A; (Bulgarelli *et al.*, 2012; Bai *et al.*, 2015)).

SynCom experiments in gnotobiotic system

Calcined clay was washed several times with tap water followed by MiliQ water. After removal of the liquid, the calcined clay was autoclaved following a liquid cycle (121 °C, 20 min), and oven-dried at 60 °C for 4 weeks. Of the washed, autoclaved and dried calcined clay, 100 g were transferred into a previously sterilized Magenta GA-7 plant culture box (Thermo Fisher Scientific, Schwerte, Germany), sealed and autoclaved again, and dried overnight prior to the experiment.

The bacterial strains used were pre-grown for 7 d in Tryptic Soy Broth 50 % (TSB 50 %, Sigma-Aldrich), and washed with 10 mM $MgCl_2$ (series of centrifugations and removal of supernatant, and final resuspension in $MgCl_2$) to remove any byproducts from the bacterial

cultures. A total of 88 and 103 bacterial strains were selected, which were differentiable

based on the their *16S* sequence (see Table S1), for the root and leaf SynComs, respectively,

and mixed in three separate inputs. For each of the three bacterial mixes, the OD_{600} was set

to 0.5 (root SynCom) and to 0.2 (leaf SynCom) with 10 mM $MgCl_2$. A sample of each of the

bacterial inputs was taken as a reference for bacterial community composition.

For the root SynCom, 70 mL ½ MS media (including vitamins without sucrose, pH 7, Sigma-

Aldrich GmbH, CatNo M5524-1L, Taufkirchen, Germany) were mixed with 1 mL of the SynCom

culture and used to inoculate one calcined-clay-containing Magenta box (n=12). As a control,

to a separate batch of magenta boxes only ½ MS media without SynCom culture was added

(n=12). Surface- sterilized *A. thaliana* Col-0 seeds were sown at the four corners of the boxes

and the lids closed. Plants were grown in a growth cabinet under short-day conditions (day:

22 °C, 10 h light; 70% humidity) for 4.5 weeks. Then, plants were inoculated with *G. orontii*

under sterile conditions (n=6, no SynCom culture, and n=6, previously treated with SynCom

culture). In addition, some boxes that were either mock- or SynCom-inoculated were opened,

but not inoculated with *G. orontii,* serving as a control. At 11 dpi, root and leaf tissues were

harvested (Figure 4a).

For the leaf SynCom, 70 mL ½ MS media (including vitamins without sucrose, pH 7, Sigma-

Aldrich GmbH, CatNo M5524-1L) were mixed into the calcined-clay-containing magenta

boxes, and surface-sterilized *A. thaliana* seeds sown at the four corners of the boxes. Plants

were grown in a growth cabinet under short-day conditions (day: 21 °C, 10 h light; night: 19

°C; 70% humidity) for 3.5 weeks. Then, a 10-fold dilution of the leaf SynCom in 10 mM $MgCl_2$

was used to spray five times with a Reagent Sprayer (CAMAG®Glass Reagent Spray, Muttenz,

Switzerland) onto the *A. thaliana* plants (see (Bai *et al.*, 2015); here*:* 1 spray volume = approx.

35 µL; n=18). At 14 d after the addition of the SynCom or respective mock treatments, plants

were inoculated with *G. orontii* under sterile conditions (n=5, no SynCom, and n=9, previously

treated with SynCom culture). In addition, some boxes that were either mock- or SynCom-

inoculated were opened, but not inoculated with *G. orontii,* serving as a control. At 7 dpi, root

and leaf tissues were harvested (Figure 4a).

Microbial profiling from plant tissues

Total DNA was extracted from the samples mentioned above using the FastDNA SPIN Kit for

Soil (MP Biomedicals, Solon, USA). Samples were homogenized in Lysis Matrix E tubes (MP

Biomedicals, Heidelberg, Germany) using the Precellys 24 tissue lyzer (Bertin Technologies,

Montigny-le-Bretonneux, France) at 6200 rpm for 30 s. DNA samples were then eluted in

80 µL nuclease-free water and used for bacterial and fungal community profiling (Durán *et*

al., 2018). Concentrations of DNA samples were fluorescently quantified, adjusted to

3.5 ng/µL, and samples used as templates in a two-step PCR amplification protocol. In the

first step, the V5–V7 region of bacterial *16S* rRNA (primers 799F-1192R), fungal *ITS1* (primers

ITS1F-ITS2) and *ITS2* (primers fITS7-ITS4) regions were amplified. Under a sterile hood, each

sample was amplified in triplicate in a 25 µL reaction volume containing 2 U DFS-Taq DNA

polymerase, 1x incomplete buffer (both Bioron GmbH, Ludwigshafen, Germany), 2 mM

$MgCl_2$, 0.3% BSA, 0.2 mM dNTPs (Life Technologies GmbH, Darmstadt, Germany) and 0.3 µM

forward and reverse primers. PCR was performed using the same parameters for all primer

pairs (94 °C/2 min, 94 °C/30 s, 55 °C/30 s, 72 °C/30 s, 72 °C/10 min for 25 cycles).

Afterwards, single-stranded DNA and proteins were digested by adding 1 µL of

Antarctic phosphatase, 1 µL xonuclease I and 2.44 µL Antarctic Phosphatase buffer (New

England BioLabs GmbH, Frankfurt, Germany) to 20 µL of the pooled PCR product. Samples

were incubated at 37 °C for 30 min and subsequently enzymes deactivated at 85 °C for

15 min. Samples were then centrifuged for 10 min at 4000 rpm, and 3 µL of this reaction were

used for a second PCR, prepared in the same way as described above, using the same

amplification protocol but with the number of cycles reduced to 10, and with primers

including barcodes with Illumina adaptors (Table S1). PCR product quality was controlled by

loading 5 µL of each reaction on an agarose gel and affirming that no band was detected in

the negative control. Afterwards, the replicated reactions were combined and purified as

follows: Bacterial amplicons were loaded on a 1.5 % agarose gel and run for 2 h at 80 V; bands

with the correct size of ~500 bp were cut out and purified using the QIAquick gel

extraction kit (Qiagen GmbH, Hilden, Germany). Fungal amplicons were purified using

Agencourt AMPure XP beads (Thermo Fisher Scientific). DNA concentration was again

fluorescently determined, and 30 ng DNA of each of the barcoded amplicons were pooled in

one library per microbial group. Each library was then purified and re-concentrated twice

with Agencourt AMPure XP beads, and 100 ng of each library were pooled together. Paired-

end Illumina sequencing was performed in-house using the MiSeq sequencer and custom

sequencing primers (Table S1).

Absolute quantification of microbial load in plant tissues

Genomic DNA from leaves inoculated with SynCom alone (n=9), *G. orontii* alone (n=9), both

SynCom and *G. orontii* (n=9), or mock-treated (n=9) was used for absolute quantification of

microbial load. Genomic DNA was fluorescently quantified and diluted to an equal

concentration of 4 µL. Each sample was subsequently used for absolute quantification *via* PCR

(qPCR) by adding SYBR green to monitor the PCR amplification in real time. Each sample was

amplified in duplicate: 4 µL of template were mixed with 7.5 µL of SYBR green (brand),

together with 1.2 µL of forward primer and 1.2 µL of reverse primer, to a final volume of 15

µL of reaction. For each organism, a specific primer pair was selected: for bacterial

assessment, the *16S* rRNA gene (primers 799F-1192R, (Wippel *et al.*, 2021)); for *G. orontii*, a

GDSL lipase-like gene (primers R263-R264, (Weßling & Panstruga, 2012)); and, for *A. thaliana*,

the *At4G26410* gene (primers L658-L659, (Hong *et al.*, 2010)). The following program was

used for amplification: pre-denaturation for 3 min at 94 °C, followed by 40 cycles of

denaturation for 15 s at 95 °C, annealing for 10 s at 62 °C and elongation for 10 s at 72 °C.

Melting curve analysis was performed from 55 °C to 95 °C, with a step-wise increase of 0.5

°C. The amount of bacterial and *G. orontii* genes was normalized to the reference plant gene

within each individual sample using the $2^{-\Delta\Delta Ct}$ equation (Pfaffl, 2001). Absolute abundances of

individual bacteria in SynCom experiments were obtained by multiplying the bacterial

gene/plant gene ratio to their individual relative abundances (RAs).

Processing of *16S* gene and *ITS* region amplicon data

Amplicon sequencing data from the natural soil experiment (plant tissues along with

unplanted controls) were demultiplexed according to their barcode sequence using the QIIME

pipeline (Caporaso *et al.*, 2010). Afterwards, DADA2 (Callahan *et al.*, 2016) was used to process

the raw sequencing reads of each sample. Unique amplicon sequence variants (ASVs) were

then inferred from error-corrected reads, followed by chimera filtering, also using the DADA2

pipeline. Next, ASVs were aligned to the SILVA database (Quast *et al.*, 2013) for taxonomic

assignment using the naïve Bayesian classifier implemented by DADA2. Next, raw reads were

mapped to the inferred ASVs to generate an abundance table, which was subsequently

employed for analyses of diversity (using the R package vegan, (Oksanen *et al.*, 2007)) and

differential abundance using the R package DESeq2 (Love *et al.*, 2014).

Sequencing data from SynCom experiments was processed using the Rbec tool (Zhang *et al.*,

2021). First, reads were de-replicated into unique tags and subsequently aligned to the

reference database, after which initial abundances were assigned to each strain according to

the copy number of each exactly aligned tag. Next, tags that were not exactly matched to any

sequence in the database were assigned a candidate error-producing reference based on *k*-

mer distances. Sequencing reads were then subsampled and an error matrix is calculated using

the mapping between subsampled reads and candidate error-producing sequences. The

parameters of the error model were recomputed iteratively until the number of re-

assignments fell below the set threshold. Strain abundances were then estimated from the

number of error-corrected reads mapped to each reference sequence. Next, we generated a

count table that was employed for downstream analyses of diversity with the R package vegan

(Oksanen *et al.*, 2007). Reads assigned to a given strain were normalized by its *16S* copy

number. Finally, amplicon data from all experimental systems were visualized using the

ggplot2 R package (Wickham, 2016).

Results

G. orontii-induced changes in root- and leaf-derived natural microbial communities

A. thaliana plants (accession Col-0) were grown in Cologne agricultural soil (CAS12; (Bulgarelli _et al._, 2012)) for 6.5 weeks and subsequently rosette leaves inoculated with _G. orontii_ conidiospores. Samples of leaf, root, and rhizosphere compartments as well as unplanted soil were harvested at 11 dpi (Figure 1A). Genomic DNA was prepared from these samples and used for PCR-based amplification of the bacterial _16S_ rRNA genes and the fungal ribosomal internal transcribed spacer (_ITS_) regions, _ITS1_ and _ITS2_. PCR amplicons were subjected to Illumina sequencing and the resulting sequencing data used for the analysis of the bacterial and fungal communities in the respective compartments. Information on the number and RA of amplicon sequence variants (ASVs) in each compartment was used to calculate α-diversity (Shannon index; within-sample diversity), β-diversity (Bray-Curtis dissimilarities; between-sample diversity), ASV enrichment, and taxonomic composition. Consistent with previous reports (Bulgarelli _et al._, 2012; Schlaeppi _et al._, 2014; Thiergart _et al._, 2020), we found the highest bacterial α-diversity in unplanted soil and the rhizosphere compartment, and lower α-diversity in roots and leaves. There was no significant difference in α-diversity between mock-treated and _G. orontii_-inoculated samples for any of the four compartments analyzed (Figure 1B). By contrast, we observed that in leaves, fungal α-diversity, determined _via ITS1_ sequences, was significantly (P=0.05) decreased in _G. orontii_-inoculated leaves (by ca. 70%; Figure 1B). This drop in α-diversity was retained upon _in silico_ depletion of _G. orontii_ reads, excluding the possibility that the decrease in species richness was due to an overrepresentation of powdery mildew reads (and thus underrepresentation of reads from other fungal taxa) in the samples. A similar outcome was obtained with _ITS2_ amplicons (Figure 1B). Together, these findings suggest that _G. orontii_ leaf colonization reduces the diversity of leaf-associated fungal communities whereas α-diversity of bacterial assemblages remains unaltered.

Analysis of β-diversity using principal-coordinates analysis (PCoA) of Bray-Curtis dissimilarities revealed distinctive community compositions in unplanted soil, rhizosphere, root and leaf compartments (30% of variance explained, P=0.001, Figure 2A). We noted that β-diversity of leaf-associated bacterial assemblages changes in response to _G. orontii_ inoculation (P=0.029), whereas bacterial communities remained indistinguishable in the

other compartments (Figure 2B and E). This suggests that *G. orontii* exerts a local effect on

bacterial community profiles, but not systemically on the bacterial root and/or rhizosphere

microbiota. Closer inspection of the *G. orontii*-induced bacterial community shifts in leaves

reveals a broad range of bacterial ASVs that show significantly altered RAs (differentially

abundant ASVs; *P*<0.05). This shift affects both abundant and low-abundant community

members with ASVs that are enriched or depleted (4.4% and 2.5% of differentially abundant

ASVs, respectively). The majority of taxa with differential fold changes are either

undetectable in the absence of *G. orontii* (enriched taxa) or undetectable following *G. orontii*

challenge (depleted taxa; Supplemental Figure 1). Similar to the leaf-associated bacterial

commensals, analysis of β-diversity of the leaf-associated fungal community reveals a

marked community shift (*ITS1* Figure 2C; *ITS2* Figure 2D). As with the changes in α-diversity

(Figure 1B), the differentiation in distinct leaf-associated fungal communities following

powdery mildew infection is not due to an overrepresentation of *G. orontii* reads, as

indicated by a similar outcome upon *in silico* depletion of the corresponding reads (Figure

2E).

Permutational multivariate analysis of variance revealed that *G. orontii* leaf colonization

explained ca. 15% and ca. 34-45% of the variation of the leaf-associated bacterial or fungal

communities, respectively (Figure 2E). To explore the changes driving this variation in

community structure, we calculated the proportion of bacterial ASVs with differential

abundance upon *G. orontii* infection at the order level. For the majority of bacterial orders

with multiple differentially abundant ASVs, both enriched and depleted ASVs were found,

although we observed more differentially enriched than differentially depleted ASVs. The

two orders with the greatest proportion of differentially abundant ASVs were

Burkholderiales and Rhizobiales (ca. 33% and 9.5%, respectively; Figure 3A), two abundant

bacterial taxa robustly found in *A. thaliana* leaf microbiota (Garrido-Oter *et al.*, 2018). We

also tested whether differentially enriched or depleted ASVs for a given bacterial order

affect its aggregated RA. Whereas the aggregated RA of Burkholderiales, Flavobacteriales

and Rhizobiales increases, the aggregated RA of Pseudonocardiales is reduced upon *G.*

orontii inoculation (*P*<0.05; Figure 3B). As these four bacterial orders belong to three phyla,

Proteobacteria, Actinobacteria and Bacteroidetes, *G. orontii* colonization influences the

abundance of phylogenetically distantly related bacterial leaf commensals.

Closer inspection of the *G. orontii*-induced fungal community shift showed that a great

proportion (15-34%) of taxa were depleted (Supplemental Figure 2), including abundant and

low abundant members of a variety of fungal classes. This result is consistent with the

aforementioned *G. orontii*-induced reduction in fungal α-diversity (Figure 1B). However, nine

fungal taxa are significantly enriched, with members of the Erysiphales and the Golubeviales

showing the highest fold change. Besides *G. orontii*, we noticed the presence of other

powdery mildew species (*Erysiphe* spp.) among the enriched fungal taxa. These could either

be contaminations in our inoculum or species introduced from the environment in the

course of the experiment and "hitchhiking" on the diseased plants.

In summary, we observed a major shift in the resident fungal community in *G. orontii*-

infected leaves, characterized by predominantly depleted taxa compared to non-infected

plants. In contrast, changes in the leaf-associated bacterial communities were more limited,

with both enriched and depleted taxa. No alterations were seen in the systemic root tissue.

G. orontii-induced changes in root- and leaf bacterial SynComs

We examined the impact of *G. orontii* leaf infection using a gnotobiotic *A. thaliana* system

and defined *A. thaliana* root and leaf bacterial consortia (root- and leaf-derived *At*-SPHERE

strains), which comprise representatives of the majority of taxa that are detectable by

culture-independent *16S* rRNA amplicon sequencing in association with plants grown in

natural soil (Bai *et al.*, 2015). In this gnotobiotic plant system, *A. thaliana* surface-sterilized

seeds were sown on a calcined clay matrix. In the first experiment, we inoculated prior to

sowing a defined bacterial consortium consisting of 88 root-derived bacterial commensals

(designated here 'root SynCom'), co-cultivated the consortium with the host for 4.5 weeks,

followed by either *G. orontii* conidiospore inoculation or mock treatment. At 11 dpi, leaf,

matrix and root samples were harvested and the corresponding DNA preparations subjected

to *16S* rRNA amplicon sequencing (Figure 4A, upper panel). PCoA of Bray-Curtis

dissimilarities of the bacterial consortia revealed their separation according to compartment

(12-22% of variance explained, *P*=0.001, Figure 4B-C). In addition, root-associated consortia

were found to be more similar to matrix-associated communities, while still separated from

the leaf-resident assemblages (Figure 4B; Supplementary Figure 4). The latter community is

likely the result of upward bacterial migration from roots to shoots during co-cultivation

(ectopic leaf colonization). In contrast to root samples, leaf samples collected from matrix-

inoculated root SynComs differentiated significantly upon *G. orontii* infection (*P*=0.02; Figure

4D and G; Supplementary Figure 4). This pattern is reminiscent of the *G. orontii*-induced

impact on *A. thaliana* plants grown in natural soil with an infection-induced change in the

bacterial leaf microbiota but non-responsiveness of the root-associated community (Figure

2F).

In the second experiment, *A. thaliana* plants were grown on sterilized calcined clay matrix

and at the age of 3.5 weeks we spray-inoculated leaves with a defined bacterial consortium

consisting of 103 leaf-derived bacterial commensals (designated here 'leaf SynCom').

Fourteen days after plant-bacteria co-cultivation, leaves were either inoculated with *G.*

orontii conidiospores or mock-treated. At 7 dpi, leaf and root samples were collected and

the corresponding microbial DNA preparations subjected to *16S* rRNA gene amplicon

sequencing (Figure 4A, lower panel). PCoA of Bray-Curtis dissimilarities of the bacterial

consortia shows their separation according to compartments (Figure 4C; Supplementary

Figure 5). Closer inspection revealed a significant separation of mock-treated and *G. orontii*-

inoculated samples in both roots and leaves (*P*=0.012 and *P*=0.002, respectively; Figure 4E-G;

Supplementary Figure 5). The *G. orontii*-induced shift in the leaf-associated bacterial

consortium is consistent with the shift of the leaf microbiota observed in *G. orontii*-

inoculated plants in natural soil (Figure 2E). However, unlike the absence of a systemic effect

on the bacterial root microbiota in natural soil upon leaf powdery mildew infection (Figure

2E), the ectopically located leaf SynCom commensals on roots were responsive to *G. orontii*

infection in the gnotobiotic plant system (see Discussion).

To assess potential changes in absolute abundance of leaf-associated bacterial communities

following *G. orontii* infection, we performed qPCR of the corresponding leaf samples with

PCR primers specific for bacterial *16S* rRNA and a *G. orontii*-specific gene and normalized

against an *A. thaliana*-specific amplicon (see Materials and Methods). Unexpectedly, we

found an approximately 2.1-fold increase in bacterial load in *G. orontii*-colonized leaf

samples, whereas *G. orontii* biomass on leaves in either the presence or absence of the

bacterial SynCom remained unaltered (Figure 5A). This was confirmed using PCoA of

absolute bacterial abundances in leaf samples spray-inoculated with the bacterial SynCom in

the presence or absence of *G. orontii*, explaining 56% of the observed variation (Figure 5B).

The increase in bacterial load in *G. orontii*-treated leaves could be seen across multiple

taxonomic classes, prominently in Actinobacteria, Alphaproteobacteria and Flavobacteria

(Figure 5C, Supplementary Figure 6 for data of individual SynCom strains). We further took

advantage of the *G. orontii*-infected plants in the gnotobiotic system to analyze the

composition of our inoculum. Although *G. orontii*, as expected, is the dominant taxum of the

inoculum, this revealed the presence of three additional fungal genera, including a known

hyperparasite of powdery mildews and another powdery mildew species (*Erysiphe* sp., >6%

average aggregated RA; Supplementary Figure 7).

Discussion

In this study, we analyzed in *A. thaliana* the influence of powdery mildew (*G. orontii*) infection on leaf- and root-associated natural and synthetic microbial communities in controlled environments (Figure 1A and 4A). For the experiments conducted in natural soil (Figure 1A), we observed no alteration in bacterial α-diversity in all tested compartments (soil, rhizosphere, root and leaf) following powdery mildew challenge, while α-diversity of the foliar fungal community was strongly reduced (Figure 1B). The more pronounced effect seen with the *ITS1* primers compared to the *ITS2* primer pair was probably a consequence of a greater molecular diversity recovered with the former primers (Bazzicalupo et al., 2013). The substantial reduction in α-diversity of the leaf-associated fungal community upon powdery mildew infection is an indication that the invasive pathogen outcompetes many resident fungal leaf endophytes (15-31% of ASVs), e.g. due to a powdery mildew pathogen-induced shift in sink-source relationships in infected compared to pathogen-free leaves. During a compatible powdery mildew interaction, photosynthetic activity of the plant host is progressively reduced both in cells directly below fungal colonies and in adjacent cells, and this process is associated with an increase in apoplastic invertase activity and an accumulation of hexoses thought to favour pathogen nutrition (Wright et al., 1995; Swarbrick et al., 2006; Eichmann & Hückelhoven, 2008). Alternative explanations may include fungus-specific antibiosis by *G. orontii* or activation of plant immune responses by the fungal invader. The former explanation appears unlikely because the obligate biotrophic pathogen has an unusually low genomic capacity for the biosynthesis of specialized metabolites (Spanu et al., 2010). *G. orontii* pathogenesis stimulates in leaves the accumulation of the defense hormone salicylic acid (SA) at later stages of infection (four dpi) and SA-dependent defense signaling limits hyphal growth and pathogen reproduction (Dewdney *et al.*, 2000; Stein *et al.*, 2008; Poraty-Gavra *et al.*, 2013). For this reason, we consider it plausible that the dramatic reduction in α-diversity of the resident community of asymptomatic leaf-associated fungi in response to *G. orontii* invasion is linked to pathogen-induced and SA-dependent immune responses and/or change in metabolic sink-source relationships. We can, however, not rule out that powdery mildew pathogens deploy secreted effector proteins to antagonize resident endophytic fungi that compete for the same ecological niche (Snelders *et al.*, 2018). Potentially different metabolic demands of leaf-associated commensal bacteria compared to fungi together with the ASV-level

compensatory changes seen within numerous bacterial orders of the leaf microbiota could

explain why the *G. orontii*-induced shift in nutrient availability and immune status does not

affect bacterial commensal diversity in the same way.

Consistent with a strong decrease in fungal species richness on *G. orontii*-infected leaves of

plants grown in natural soil, we found that the altered fungal community profile is

characterized by a reduction in RAs for most and an increase in RAs of a few taxa, while the

local bacterial community profile was only slightly affected (Figure 2; Supplementary Figure

3). Leaf-associated fungal endophytes whose RA increases upon *G. orontii* infection comprise

mainly one other powdery mildew species (*Erysiphe* sp.) or fungi typically associated with

them, such as *Golubevia* sp., which are basidiomycete hyperparasites of powdery mildew

pathogens (Russ *et al.*, 2021).

Closer inspection of the *G. orontii*-induced shift of bacterial community diversity in leaves

revealed 40 bacterial orders, containing several ASVs whose RA is depleted or enriched

(Figure 3A and Supplemental Figure 1). However, significant alterations in aggregated ASV-

level RAs were seen only in four bacterial orders, with three of them showing an increase

and one a decrease (Burkholderiales, Flavobacteriales, Rhizobiales and Pseudonocardiales,

respectively; Figure 3B). This pattern suggests widespread compensatory changes at the ASV

level within bacterial orders that may allow maintenance of the higher taxonomic structure

of the bacterial leaf microbiota during powdery mildew pathogenesis. The fungal pathogen

thus mediates local community shifts in the aggregated RAs limited to four bacterial orders

belonging to the three main phyla of the *A. thaliana* microbiota (Actinobacteria,

Proteobacteria and Bacteriodetes (Bulgarelli *et al.*, 2012)). In addition, in our SynCom

experiments, colonization by the biotrophic pathogen unexpectedly increased the load of

bacterial commensals in leaves approximately 2.1-fold, and this increase was seen roughly

equally proportional for all bacterial classes analyzed (Figure 5). The rise in bacterial

numbers could be an indirect consequence of the altered sink-source relationships upon

infection with the biotrophic pathogen, which turns infected leaves into a metabolic sink

associated with an altered availability of non-structural carbohydrates (Wright *et al.*, 1995;

Swarbrick *et al.*, 2006). Consequently, the bacterial commensals might benefit from an

accumulation and/or alteration in fluxes of apoplastic hexose sugars and reduction in export

of sucrose from the leaf.

In the case of bacteria, changes were only detectable locally (in the infected leaves) but not

systemically (in roots or rhizosphere compartments) and only applied to community

diversity, while species richness remains unaltered (Figure 1B and 2). The subtle shift in local

bacterial diversity could be an indirect consequence of the significant alterations in fungal

community richness and diversity upon *G. orontii* invasion. However, at the bacterial order

level, we noted a striking overall similarity of the *G. orontii*-induced local bacterial

community shifts between the natural soil and bacterial SynCom experiments (significant

increases in the RA of Burkholderiales, Flavobacteriales and Rhizobiales and a similar trend

for Caulobacterales; Supplementary Fig. 6B). This suggests that our gnotobiotic plant system

recapitulates features of the powdery mildew pathogen-induced bacterial community shifts

seen in plants grown in natural soil. It further rules out the possibility that these shifts are

the result of bacterial immigration into vacated leaf niches that had to be abandoned by the

resident fungal endophytes in the course of *G. orontii* pathogenesis.

The only systemic effect on root-associated microbes upon powdery mildew infection

reported so far involves nitrogen-fixing rhizobia, which engage in root symbiosis with

legumes. In pea (*Pisum sativum*), powdery mildew (*Erysiphe pisi*) colonization was found to

result in both a reduction of nodulation and reduced size of root nodules in the leaf-infected

plants (Singh & Mishra, 1992). The absence of a systemic effect on the bacterial root

microbiota following *G. orontii* challenge differs from *A. thaliana* plants infected with the

foliar downy mildew pathogen, the obligate biotrophic oomycete *H. arabidopsidis*.

Colonization of leaves with *H. arabidopsidis* promotes the specific enrichment of three

bacterial taxa (*Xanthomonas*, *Stenotrophomonas* and *Microbacterium* spp.) from soil to the

rhizosphere, where these three taxa synergistically induced systemic defence responses and

promoted plant growth, resulting in enhanced disease resistance against downy mildew

(Berendsen *et al.*, 2018). Although *G. orontii* and *H. aribidopsidis* are both obligate biotrophic

foliar pathogens, only the latter pathogen can infect leaf mesophyll cells and ramify inside

this organ, and this may be one reason why only *H. arabidopsidis* can systemically induce

changes in the bacterial root microbiota.

Our experiments using germ-free plants, pre-inoculated with a root- or a leaf-SynCom and

either mock-treated or challenged with *G. orontii* (Figure 4), recapitulated in parts the

pathogen-induced shift in bacterial community profiles of plants grown in natural soil.

However, in these experiments also the root samples showed a significant difference in β-

diversity of the bacterial root microbiota following *G. orontii* inoculation (Figure 4F and G).

As roots were not pre-inoculated in this experiment, the bacteria recovered from root

samples must have originated from the leaf inoculation and either reached the below-

ground roots in calcined clay accidentally (e.g. by wash-off) or by downward migration, e.g.

via the plant vasculature. Irrespective of the mechanism(s) underlying ectopic root

colonization of the leaf-inoculated bacterial SynCom, this result indicates a (subtle) systemic

effect of foliar inoculation with the powdery mildew pathogen on the root-associated

bacterial community, which was not seen in the experiment with plants grown in natural soil

(see above and Figure 2B). This difference is probably the result of different conditions in the

two experimental setups. The SynCom experiment involves a reduced complexity of the

bacterial community, allowing us to track the RA of many members with strain-specific

resolution rather than the RA of ASVs, which could represent an average of multiple

genetically polymorphic strains that share an identical *16S* rRNA gene. Therefore, the higher

complexity of the root microbiota in natural soil may mask putative systemic effects on

bacterial assemblages induced by *G. orontii* colonization.

It is surprising that the pre-inoculated leaf SynCom does not mediate detectable protective

activity against the fungal powdery mildew pathogen, e.g. through activation of defence-

associated gene expression in leaves or antagonistic bacteria-fungus interactions (Figure 5A;

(Vogel *et al.*, 2016; Durán *et al.*, 2018)). In roots, the presence of the bacterial microbiota is

essential for indirect protection and survival of *A. thaliana* against the otherwise detrimental

activity of diverse fungal root endophytes (Durán *et al.*, 2018). In *A. thaliana* leaves, several

members of the genus *Sphingomonas*, originally isolated from plants, conferred plant

protection against the foliar bacterial pathogen *P. syringae* DC3000 and *Xanthomonas*

campestris in a gnotobiotic system, whereas no protection was observed by colonization

with members of the genus *Methylobacterium* (Innerebner *et al.*, 2011). Thus, our results

suggest that the bulk of the resident bacterial leaf microbiota of *A. thaliana* does not have a

dedicated role in indirect protection against epiphytic *G. orontii* invasion in the tested

gnotobiotic system. Whether this is also true for fungal leaf pathogens that colonize

mesophyll cells in the leaf interior remains to be tested.

Part of the powdery mildew-induced changes in the *A. thaliana* microbiota could be due to

the suppression of plant immunity and reprogramming of host cells for parasitism by the

obligate biotrophic pathogen (Schulze-Lefert & Panstruga, 2003). The effect of defence

suppression in colonized and neighboring leaf epidermal cells is for example evident by the

phenomenon of "induced accessibility", which refers to enabled host cell entry of non-

adapted powdery mildews in the vicinity of established powdery mildew infection sites

(Yamaoka *et al.*, 1994; Lyngkjær & Carver, 1999b; Lyngkjær & Carver, 1999a; Lyngkjær *et al.*,

2001). Powdery mildew-induced alterations of the host transcriptome likely represent the

net outcome of counteracting activities of the plant immune system and the fungal intruder

(Fabro *et al.*, 2008). Although leaf-associated bacterial commensals also extensively

reprogram host transcriptomes, stimulating and/or repressing the activity of gene clusters

enriched in immunity-and metabolism-associated functions in leaves (Vogel *et al.*, 2016), the

additional pathogen-specific changes in the host transcriptional profile during infection likely

contribute to modulating the quantity and composition of the resident microbiota.

Based on the analysis of our *G. orontii* inoculum on gnotobiotic plants, we noted that

approximately 50% of the reads originate from different fungi, including a powdery mildew

hyperparasite and another powdery mildew species (*Erysiphe* sp.; Supplementary Figure 7).

Our finding highlights the difficulties of maintaining a plant pathogen with an obligate

biotrophic lifestyle that must be propagated on living host plants in pure culture. The co-

occurrence with these microbes may balance powdery mildew proliferation. These

observations might have implications not only for the interpretation of aspects of the data

obtained in this work, but also for studies that use other obligate biotrophic plant

pathogens.

Author contributions

PSL and RP conceived the project. RP, PSL and RGO designed the experiments. ALR, AR and HM performed the experiments with natural soil. ALR, AR and PD performed the SynCom experiments. PD and RGO analyzed the data. PD and RGO created the Figures. RP and PSL wrote the manuscript with support from AR, PD and RGO.

Data availability statement

Raw demultiplexed sequencing data and corresponding mapping files will be available at ENA accession number PRJEB43139.

Funding

This research was funded by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) under Germany's Excellence Strategy—EXC-number 2048/1—project 390686111 (R.G.-O. and P.S.-L.), the Priority Programme SPP 2125 DECRyPT (R.G.-O. and P.S.-L.), a European Research Council advanced grant (ROOTMICROBIOTA), a RIKEN grant (SYMBIOLOGY), and a cooperative research project with Dong-A University funded by the Republic of Korea to P.S.-L., as well as funds to P.S.-L. from the Max Planck Society. M.H. was supported by JSPS KAKENHI grants 20K05955, 19KT0033 and 15J04093.

References

Acevedo-Garcia, J, Gruner, K, Reinstädler, A, Kemen, A, Kemen, E, Cao, L, Takken, FLW, Reitz, MU, Schäfer, P, O'Connell, RJ, Kusch, S, Kuhn, H, Panstruga, R. 2017. The powdery mildew-resistant Arabidopsis *mlo2 mlo6 mlo12* triple mutant displays altered infection phenotypes with diverse types of phytopathogens. *Scientific Reports* **7**: 27.

Agler, MT, Ruhe, J, Kroll, S, Morhenn, C, Kim, S-T, Weigel, D, Kemen, EM. 2016. Microbial hub taxa link host and abiotic factors to plant microbiome variation. *PLoS Biology* **14**: e1002352.

Bai, Y, Müller, DB, Srinivas, G, Garrido-Oter, R, Potthoff, E, Rott, M, Dombrowski, N, Munch, PC, Spaepen, S, Remus-Emsermann, M, Huttel, B, McHardy, AC, Vorholt, JA, Schulze-Lefert, P. 2015. Functional overlap of the *Arabidopsis* leaf and root microbiota. *Nature* **528**: 364–369.

Bazzicalupo, AL, Bálint, M, Schmitt, I. 2013. Comparison of ITS1 and ITS2 rDNA in 454 sequencing of hyperdiverse fungal communities. *Fungal Ecology* **6**: 102–109.

Berendsen, RL, Vismans, G, Yu, K, Song, Y, Jonge, R de, Burgman, WP, Burmølle, M, Herschend, J, Bakker, PAHM, Pieterse, CMJ. 2018. Disease-induced assemblage of a plant-beneficial bacterial consortium. *ISME Journal*.

Bulgarelli, D, Rott, M, Schlaeppi, K, Ver Loren van Themaat, E, Ahmadinejad, N, Assenza, F, Rauf, P, Huettel, B, Reinhardt, R, Schmelzer, E, Peplies, J, Gloeckner, FO, Amann, R, Eickhorst, T, Schulze-Lefert, P. 2012. Revealing structure and assembly cues for *Arabidopsis* root-inhabiting bacterial microbiota. *Nature* **488**: 91–95.

Bulgarelli, D, Schlaeppi, K, Spaepen, S, Ver Loren Themaat, E, Schulze-Lefert, P. 2013. Structure and functions of the bacterial microbiota of plants. *Annual Review of Plant Biology* **64**: 807–838.

Callahan, BJ, McMurdie, PJ, Rosen, MJ, Han, AW, Johnson, AJA, Holmes, SP. 2016. DADA2: High-resolution sample inference from Illumina amplicon data. *Nature Methods* **13**: 581–583.

Caporaso, JG, Kuczynski, J, Stombaugh, J, Bittinger, K, Bushman, FD, Costello, EK, Fierer, N, Peña, AG, Goodrich, JK, Gordon, JI, Huttley, GA, Kelley, ST, Knights, D, Koenig, JE, Ley, RE, Lozupone, CA, McDonald, D, Muegge, BD, Pirrung, M, Reeder, J, Sevinsky, JR, Turnbaugh, PJ, Walters, WA, Widmann, J, Yatsunenko, T, Zaneveld, J, Knight, R. 2010. QIIME allows analysis of high-throughput community sequencing data. *Nature Methods* **7**: 335–336.

Coleman-Derr, D, Desgarennes, D, Fonseca-Garcia, C, Gross, S, Clingenpeel, S, Woyke, T, North, G, Visel, A, Partida-Martinez, LP, Tringe, SG. 2016. Plant compartment and biogeography affect microbiome composition in cultivated and native Agave species. *New Phytologist* **209**: 798–811.

Durán, P, Thiergart, T, Garrido-Oter, R, Agler, M, Kemen, E, Schulze-Lefert, P, Hacquard, S. 2018. Microbial interkingdom interactions in roots promote Arabidopsis survival. *Cell* **175**: 973-983.e14.

Fabro, G, Di Rienzo, JA, Voigt, CA, Savchenko, T, Dehesh, K, Somerville, S, Alvarez, ME. 2008. Genome-wide expression profiling Arabidopsis at the stage of *Golovinomyces cichoracearum* haustorium formation. *Plant Physiology* **146**: 1421–1439.

Finkel, OM, Castrillo, G, Herrera Paredes, S, Salas González, I, Dangl, JL. 2017.
Understanding and exploiting plant beneficial microbes. *Current Opinion in Plant Biology*
38: 155–163.
Gao, C, Montoya, L, Xu, L, Madera, M, Hollingsworth, J, Purdom, E, Singan, V, Vogel, J,
Hutmacher, RB, Dahlberg, JA, Coleman-Derr, D, Lemaux, PG, Taylor, JW. 2020. Fungal
community assembly in drought-stressed sorghum shows stochasticity, selection, and
universal ecological dynamics. *Nature Communications* **11**: 34.
Garrido-Oter, R, Nakano, RT, Dombrowski, N, Ma, K-W, McHardy, AC, Schulze-Lefert, P.
2018. Modular traits of the rhizobiales root microbiota and their evolutionary relationship
with symbiotic rhizobia. *Cell Host & Microbe* **24**: 155-167.e5.
Hacquard, S, Garrido-Oter, R, González, A, Spaepen, S, Ackermann, G, Lebeis, S, McHardy,
AC, Dangl, JL, Knight, R, Ley, R, Schulze-Lefert, P. 2015. Microbiota and host nutrition
across plant and animal kingdoms. *Cell Host & Microbe* **17**: 603–616.
Hacquard, S, Schadt, CW. 2015. Towards a holistic understanding of the beneficial
interactions across the *Populus* microbiome. *New Phytologist* **205**: 1424–1430.
Harbort, CJ, Hashimoto, M, Inoue, H, Niu, Y, Guan, R, Rombolà, AD, Kopriva, S, Voges,
MJEEE, Sattely, ES, Garrido-Oter, R, Schulze-Lefert, P. 2020. Root-secreted coumarins and
the microbiota interact to improve iron nutrition in *Arabidopsis*. *Cell Host & Microbe* **28**:
825–837.e6.
Hong, SM, Bahn, SC, Lyu, A, Jung, HS, Ahn, JH. 2010. Identification and testing of superior
reference genes for a starting pool of transcript normalization in Arabidopsis. *Plant and*
Cell Physiology **51**: 1694–1706.
Innerebner, G, Knief, C, Vorholt, JA. 2011. Protection of *Arabidopsis thaliana* against leaf-
pathogenic *Pseudomonas syringae* by *Sphingomonas* strains in a controlled model system.
Applied and Environmental Microbiology **77**: 3202–3210.
Jakuschkin, B, Fievet, V, Schwaller, L, Fort, T, Robin, C, Vacher, C. 2016. Deciphering the
pathobiome: Intra- and interkingdom interactions involving the pathogen *Erysiphe*
alphitoides. *Microbial Ecology* **72**: 870–880.
Lauber, CL, Hamady, M, Knight, R, Fierer, N. 2009. Pyrosequencing-based assessment of soil
pH as a predictor of soil bacterial community structure at the continental scale. *Applied*
and Environmental Microbiology **75**: 5111–5120.
Lebeis, SL, Paredes, SH, Lundberg, DS, Breakfield, N, Gehring, J, McDonald, M, Malfatti, S,
Glavina del Rio, T, Jones, CD, Tringe, SG, Dangl, JL. 2015. Salicylic acid modulates
colonization of the root microbiome by specific bacterial taxa. *Science* **349**: 860–864.
Love, MI, Huber, W, Anders, S. 2014. Moderated estimation of fold change and dispersion
for RNA-seq data with DESeq2. *Genome Biology* **15**: 550.
Lundberg, DS, Lebeis, SL, Paredes, SH, Yourstone, S, Gehring, J, Malfatti, S, Tremblay, J,
Engelbrektson, A, Kunin, V, Del Rio, TG, Edgar, RC, Eickhorst, T, Ley, RE, Hugenholtz, P,
Tringe, SG, Dangl, JL. 2012. Defining the core *Arabidopsis thaliana* root microbiome.
Nature **488**: 86–90.
Lyngkjær, HF, Carver, TLW. 1999a. Modification of *mlo5* resistance to *Blumeria graminis*
attack in barley as a consequence of induced accessibility and inaccessibility. *Physiological*
and Molecular Plant Pathology **55**: 163–174.

Lyngkjær, MF, Carver, TLW. 1999b. Induced accessibility and inaccessibility to *Blumeria*
graminis f.sp *hordei* in barley epidermal cells attacked by a compatible isolate.
Physiological and Molecular Plant Pathology **55**: 151–162.

Lyngkjær, MF, Carver, TLW, Zeyen, RJ. 2001. Virulent *Blumeria graminis* infection induces
penetration susceptibility and suppresses race-specific hypersensitive resistance against
avirulent attack in *Mla1*-barley. *Physiological and Molecular Plant Pathology* **59**: 243–256.

Müller, DB, Vogel, C, Bai, Y, Vorholt, JA. 2016. The plant microbiota: Systems-level insights
and perspectives. *Annual Review of Genetics* **50**: 211–234.

Oksanen, J, Kindt, R, Legendre, P, O'Hara, B, Stevens M.H.H., Oksanen, MJ. 2007. The
vegan package: community ecology package. *R package version*.

Panstruga, R, Kuhn, H. 2019. Mutual interplay between phytopathogenic powdery mildew
fungi and other microorganisms. *Molecular Plant Pathology* **20**: 463–470.

Pfaffl, MW. 2001. A new mathematical model for relative quantification in real-time RT-PCR.
Nucleic Acids Research **29**: e45.

Quast, C, Pruesse, E, Yilmaz, P, Gerken, J, Schweer, T, Yarza, P, Peplies, J, Glöckner, FO.
2013. The SILVA ribosomal RNA gene database project: Improved data processing and
web-based tools. *Nucleic Acids Research* **41**: D590-596.

Russ, L, Lombaers-van der Plas, C, Castillo-Russi, JD, Zijlstra, C, Köhl, J. 2021. Deciphering
the modes of action of *Golubevia sp.*, an antagonist against the causal agent of powdery
mildew in wheat, using an mRNA-based systems approach. *Biological Control* **152**: 104446.

Schlaeppi, K, Dombrowski, N, Oter, RG, van Loren Themaat, E ver, **Schulze-Lefert, P. 2014.**
Quantitative divergence of the bacterial root microbiota in *Arabidopsis thaliana* relatives.
Proceedings of the National Academy of Sciences of the United States of America **111**:
585–592.

Schulze-Lefert, P, Panstruga, R. 2003. Establishment of biotrophy by parasitic fungi and
reprogramming of host cells for disease resistance. *Annual Review of Phytopathology* **41**:
641–667.

Singh, UP, Mishra, GD. 1992. Effect of powdery mildew (*Erysiphe pisi*) on nodulation and
nitrogenase activity in pea (*Pisum sativum*). *Plant Pathology* **41**: 262–264.

Snelders, NC, Kettles, GJ, Rudd, JJ, Thomma, BPHJ. 2018. Plant pathogen effector proteins
as manipulators of host microbiomes? *Molecular Plant Pathology* **19**: 257–259.

Swarbrick, PJ, Schulze-Lefert, P, Scholes, JD. 2006. Metabolic consequences of susceptibility
and resistance (race-specific and broad-spectrum) in barley leaves challenged with
powdery mildew. *Plant, Cell & Environment* **29**: 1061–1076.

Thiergart, T, Durán, P, Ellis, T, Vannier, N, Garrido-Oter, R, Kemen, E, Roux, F, Alonso-
Blanco, C, Ågren, J, Schulze-Lefert, P, Hacquard, S. 2020. Root microbiota assembly and
adaptive differentiation among European Arabidopsis populations. *Nature Ecology &*
Evolution **4**: 122–131.

Vogel, C, Bodenhausen, N, Gruissem, W, Vorholt, JA. 2016. The Arabidopsis leaf
transcriptome reveals distinct but also overlapping responses to colonization by
phyllosphere commensals and pathogen infection with impact on plant health. *New*
Phytologist **212**: 192–207.

Vorholt, JA. 2012. Microbial life in the phyllosphere. *Nature Reviews Microbiology* **10**: 828–
840.

Weßling, R, Panstruga, R. 2012. Rapid quantification of plant-powdery mildew interactions
by qPCR and conidiospore counts. *Plant Methods* **8**: 35.

Wickham, H. 2016. ggplot2. Elegant graphics for data analysis. Cham: Springer International
Publishing.

Wippel, K, Tao, K, Niu, Y, Zgadzaj, R, Guan, R, Dahms, E, Zhang, P, Jensen, DB, Logemann, E,
Radutoiu, S, Schulze-Lefert, P, Garrido-Oter, R. 2021. Host preference and invasiveness of
commensals in the *Lotus* and *Arabidopsis* root microbiota. *bioRxiv*.

Wright, DP, Baldwin, BC, Shephard, MC, Scholes, JD. 1995. Source-sink relationships in
wheat leaves infected with powdery mildew. I. Alterations in carbohydrate metabolism.
Physiological and Molecular Plant Pathology **47**: 237–253.

Yamaoka, N, Toyoda, K, Kobayashi, I, Kunoh, H. 1994. Induced accessibility and enhanced
inaccessibility at the cellular level in barley coleoptiles .13. Significance of haustorium
formation by the pathogen *Erysiphe graminis* for induced accessibility to the non-
pathogen *E. pisi* as asessed by nutritional manipulations. *Physiological and Molecular Plant*
Pathology **44**: 217–225.

Zhang, J, Liu, Y-X, Zhang, N, Hu, B, Jin, T, Xu, H, Qin, Y, Yan, P, Zhang, X, Guo, X, Hui, J, Cao,
S, Wang, X, Wang, C, Wang, H, Qu, B, Fan, G, Yuan, L, Garrido-Oter, R, Chu, C, Bai, Y.
2019a. NRT1.1B is associated with root microbiota composition and nitrogen use in field-
grown rice. *Nature Biotechnology* **37**: 676–684.

Zhang, P, Spaepen, S, Bai, Y, Hacquard, S, Garrido-Oter, R. 2021. Reference-based error
correction of amplicon sequencing data from synthetic communities. *bioRxiv*.

Zhang, Z, Kong, X, Jin, D, Yu, H, Zhu, X, Su, X, Wang, P, Zhang, R, Jia, M, Deng, Y. 2019b.
Euonymus japonicus phyllosphere microbiome is significantly changed by powdery
mildew. *Archives of Microbiology* **201**: 1099–1109.

Zhang, Z, Luo, L, Tan, X, Kong, X, Yang, J, Wang, D, Zhang, D, Jin, D, Liu, Y. 2018. Pumpkin
powdery mildew disease severity influences the fungal diversity of the phyllosphere. *PeerJ*
6: e4559.

Figure 1. Analysis of α-diversity including all culture-independent samples (ASV-level

analysis) for both bacterial and fungal communities. **A)** Schematic experimental set-up and

representative pictures of mock-treated and *G. orontii*-inoculated plants. **B)** Boxplots of

within-sample diversity (Shannon index) for each compartment and condition. Significant

differences in bacterial community diversity within compartments are marked with an

asterisk (Student´s *t*-test, * *P*<0.05; n.s., not significant).

Figure 2. Analysis of β-diversity including all culture-independent samples (ASV-level

analysis), for both bacterial and fungal communities. A) PCoA plot of Bray-Curtis

dissimilarities between bacterial community samples, color-coded by compartment and

shaped based on treatment. **B)** Subset of leaf samples where separation between

treatments (mock- vs *G. orontii*-treated) can be observed. **C-D)** PCoA plots of fungal

communities in leaf samples for *ITS1* (**C**) and *ITS2* (**D**) profiles. **E)** Variance explained of

bacterial (left set of columns) and fungal community structure (right set of columns) upon *G.*

orontii infection. Note that for fungal communities, an *in silico* depletion of *G. orontii*-

assigned reads was performed.

Figure 3. Analysis of bacterial ASVs significantly affected by *G. orontii* infection on leaves.

A) Proportion of ASVs that significantly changed their relative abundance upon infection

with *G. orontii*, grouped by order level, in relation to the overall number of ASVs affected. **B)**

Log-transformed aggregated relative abundance of each bacterial order shown in **(A)**, which

has both enriched and depleted ASVs, compared between mock and *G. orontii*-treated

conditions.

Figure 4. Analysis of beta-diversity including all Root *At*-SPHERE SynCom samples and all

Leaf *At*-SPHERE SynCom samples based on relative abundances. A) Schematic experimental

set-up and representative pictures of mock-treated and *G. orontii*-inoculated plants, with

and without SynCom. **B)** PCoA of Bray-Curtis distances of all compartment inoculated with

Root *At*-SPHERE strains. **C)** PCoA of Bray-Curtis distances of all compartment inoculated with

Leaf *At*-SPHERE strains. **D-F)** Subset of leaf **(D and E)** and root samples **(F),** inoculated with

either Root *At*-SPHERE strains **(D)** or Leaf *At*-SPHERE strains **(E and F)**, showing the effect on

bacterial community structure upon *G. orontii* infection (different shapes). **G)** Variance

explained on bacterial community structure upon *G. orontii* infection (PERMANOVA analysis

on Bray-Curtis dissimilarities).

Figure 5. Absolute quantification of bacterial load on leaves inoculated with Leaf *At*-

SPHERE strains. A) Bacterial (left panel) and *G. orontii* (right panel) load calculated as the

relative quantification of each microbial gene to an *A. thaliana* reference gene. **B)** PCoA plot

of absolute abundances of bacterial communities in leaves inoculated with Leaf *At*-SPHERE

strains. **C)** Absolute abundance of each strain utilized in this experiment, color-coded by

their taxonomic assignment at the family level, in leaf SynCom-inoculated leaves mock- and

G. orontii-treated.

Supplementary Figure 1: Changes in RA of bacterial ASVs in *A. thaliana* leaves upon *G.*

***orontii* inoculation.** Each bacterial ASV that showed significant enrichment in leaf samples

treated with *G. orontii*, compared to mock-treated samples (*P*<0.05, DESeq package in R) is

shown. Each row represents a different ASV, where the size of each dot corresponds to the

log-transformed RA in a given sample (columns). The dots are color-coded based on the ASV

taxonomic assignment at the class level. The right panel depicts the fold change of each

significantly enriched ASV in *G. orontii*-treated leaves, compared to mock control.

Supplementary Figure 2: Changes in RA of fungal ASVs in *A. thaliana* leaves upon *G. orontii*

inoculation, for the ITS1 community profiles. Each fungal ASV that showed significant

enrichment in leaf samples treated with *G. orontii*, compared to mock-treated samples

(*P*<0.05, DESeq package in R) is shown here. Each row represents a different ASV, where the

size of each dot corresponds to the log-transformed RA in a given sample (columns). The dots

are color-coded based on the ASV taxonomic assignment at the class level. The right panel

depicts the fold change of each significantly enriched ASV in *G. orontii*-treated leaves,

compared to mock control.

Supplementary Figure 3: Changes in RA of fungal ASVs in *A. thaliana* leaves upon *G. orontii* inoculation, for the ITS2 community profiles. Each fungal ASV that showed significant enrichment in leaf samples treated with *G. orontii*, compared to mock-treated samples (*P*<0.05, DESeq package in R) is shown here. Each row represents a different ASV, where the size of each dot corresponds to the log-transformed RA in a given sample (columns). The dots are color-coded based on the ASV taxonomic assignment at the class level. The right panel depicts the fold change of each significantly enriched ASV in *G. orontii*-treated leaves, compared to mock control.

Supplementary Figure 4: Heatmap of bacterial strain-specific RA including SynCom samples (Root *At*-SPHERE strains). Warm colors correspond to abundant strains detected in at least one input sample (>0.1% RA). Samples (columns) are grouped by compartment and *G. orontii* treatment.

Supplementary Figure 5: Heatmap of bacterial strain-specific RAs including SynCom samples (Leaf *At*-SPHERE strains). Warm colors correspond to abundant strains detected in at least one input sample (>0.1% RA). Samples (columns) are grouped by compartment and *G. orontii* treatment.

Supplementary Figure 6: Comparison of absolute abundances of individual Leaf *At*-SPHERE bacteria to RAs in the culture-independent approach. A) Absolute abundance (shown as the product of the bacterial/plant gene ratio multiplied with the bacterial RA of each bacterial strain from the Leaf *At*-SPHERE inoculated in *A. thaliana* leaves without and with *G. orontii*. Boxplots are color-coded based on the taxonomic assignment of each strain at the class level. Significant differences in absolute abundances upon *G. orontii* inoculation are depicted with a red asterisk (Student's *t*-test, *P*< 0.05, FDR corrected). **B)** Aggregated relative (culture-independent approach) and absolute (culture-dependent approach) abundances of bacterial orders shared between the two experimental set-ups. Significant differences in abundances

upon *G. orontii* inoculation are depicted with a red asterisk (Student's *t*-test, *P*< 0.05, FDR

corrected).

Supplementary Figure 7: Profiling of *G. orontii* inoculum. A) Number of reads assigned to

either bacterial or fungal ASVs in *A. thaliana* leaves inoculated with *G. orontii* **B)** RAs of

bacterial ASVs color-coded based on their taxonomic assignment at the class level. **C)** RAs of

fungal ASVs from the ITS1 profiles, color-coded based on their taxonomic assignment at the

genus level. **D)** RAs of fungal ASVs from the ITS2 profiles, color-coded based on their

taxonomic assignment at the genus level.

Supplementary Table 1:

-   List of Root *At*-SPHERE strains used in this study and their taxonomic assignment

-   List of Leaf *At*-SPHERE strains used in this study and their taxonomic assignment

-   Primers used in library preparation for amplicon sequencing

-   Sequencing primers used for MiSeq sequencing

-   Natural soil experiment mapping file for the analysis of community profiles

-   SynCom experiments mapping file for the analysis of community profiles

-   Primers used in qPCR for microbial absolute quantification in plant tissues

A

1. Root-/soil-derived SynCom → Calcined clay → 2. Arabidopsis seeds → 4.5 weeks → +/- *G. orontii* conidiospores → 11 days → Leaves / Matrix / Roots

→ Calcined clay → 3.5 weeks → Leaf-derived SynCom → 14 days → +/- *G. orontii* conidiospores → 7 days → Leaves / Roots

Mock-treated | *G. o*-inoculated
SynCom-treated | SynCom+*G. o*-inoc.

B — All samples / Root-derived SynCom
PCoA 2 (16.71%) vs PCoA 1 (27.53%)

C — All samples / Leaf-derived SynCom
PCoA 2 (17.51%) vs PCoA 1 (33.53%)

Compartment
- SynCom Input
- Matrix
- Root
- Leaf

Treatment
- Mock
- *G. orontii*

D — Leaf samples / Root-derived SynCom
PCoA 2 (23.69%) vs PCoA 1 (35.47%)

E — Leaf samples / Leaf-derived SynCom
PCoA 2 (15.08%) vs PCoA 1 (37.16%)

F — Root samples / Leaf-derived SynCom
PCoA 2 (24.38%) vs PCoA 1 (29.22%)

G
Variance (%)

Compartment
- Root
- Leaf

16S SynCom (Root): n.s ; ** (p=0.02)
16S SynCom (Leaf): ** (p=0.012) ; *** (p=0.002)