Response of whitefly to the wild tomato *Solanum habrochaites*

Fengqi Li[1,*], Youssef Dewer[2], Jiahui Tian[1], Du Li[1], Cheng Qu[1], Zhen Yang[3] & Chen Luo[1]

[1] Beijing Key Laboratory of Environment Friendly Management on Fruit Diseases and Pests in North China, Institute of Plant and Environment Protection, Beijing Academy of Agriculture and Forestry Sciences, Beijing 100097, China

[2] Bioassay Research Department, Central Agricultural Pesticide Laboratory, Agricultural Research Center, Dokki, Giza 12618, Egypt; dewer72@yahoo.com

[3] Tianjin University of Traditional Chinese Medicine, Tianjin 300000, China

* Correspondence: pandit@163.com

Abstract

The whitefly Bemisia tabaci (Gennadius) causes severe damage to cultivated tomato in many regions of the world through direct feeding and indirectly through transmission of plant viruses. Field observations show that B. tabaci is rarely infested the non-host plants such as the wild tomato Solanum habrochaites; however, the molecular mechanism involved in the recognition of wild plant odors is still unclear. In this study, we assessed the effects of S. habrochaites on the survival, fecundity, and egg hatchability of the Mediterranean (MED) species of B. tabaci. Expression and splicing of stress-response genes in whitefly exposed to S. habrochaites was analyzed using RNA-sequencing data and alternative splicing analysis. These results indicated that the S. habrochaites treatment can induce the expression of environmental stress genes in B. tabaci. This study may help us to a better understanding of the molecular

mechanisms involved in the olfactory recognition of non-host volatiles particularly

wild tomato relative. Furthermore, the findings of this study may provide excellent

chances of finding a suitable antagonist of eco-friendly properties which can block the

perception of chemosensory signals. Thereby, the feeding behavior and food

preferences of B. tabaci can be manipulated and thus insect populations can

eventually be controlled.

Keywords: whitefly, *Solanum habrochaites*, transcriptome

1. Introduction

The Whitefly *Bemisia tabaci* (Gennadius) is one of the most important pests that

causes economic damage to crop plants (De et al., 1997). *B. tabaci* has a high

oviposition rate and rapid population growth (Erdogan et al., 2008). The Middle

East-Asia Minor1 (MEAM1) and Mediterranean (MED) subspecies are the most

invasive *B. tabaci* whiteflies worldwide (Chu et al., 2006). In China, the MED cryptic

species has now been replaced by MEAM1 in many regions   (Hu et al., 2011).

B. tabaci is one of the main tomato pests that reduces tomato growth through direct

feeding and indirectly through transmission tomato yellow leaf curl virus (TYLCV)

(Su et al., 2016). *B. tabaci* is the only vector of TYLCV, so whitefly control is one of

the key measures used to manage TYLCV (Wei et al., 2017). Chemical insecticides

remain the main tools in the management of whitefly, but the rapid development of

resistance to several chemical classes of insecticides has made it so difficult to control

(Elbert et al., 2000). Therefore, there are urgent demands for safe alternative strategies

in modern pest management, i.e. for controlling whitefly populations. One of these

novel approaches is based on the fact that chemical communication via the olfactory

system drives essential behaviors of insects (Ingham et al., 2020). Indeed, detecting

appropriate chemical signals is essential for insects to find their mating partners,

oviposition sites and food sources (Pelosi et al., 2017; Leal and Walter, 2013).

Accordingly, the olfactory system of many insects has evolved to chemo-detectors of

high sensitivity and accuracy, which allow the insect to detect plant volatile odors at

extremely low concentrations and to discriminate a large variety of odor cues (Suh et

al., 2014) . Thus, understanding of the mechanisms and signals that are involved in

non-host selection behavior of *B. tabaci* may lead to strategies that exploit the

odorous alert signals to enhance crop resistance.

B. tabaci is attracted to cultivated tomato, but it is repelled by the wild tomato

relative *S. habrochaites*. The repellent behavior response of *B. tabaci* exposed to *S.*

habroachaites as non-host plant was found to be associated with the emission of

7-epizingiberene and R-curcumene compounds (Bleeker et al., 2009). This two

terpene volatiles have shown strong repellency against *B. tabaci* and induced *S.*

habrochaites to develop enhanced resistance to insect infestation (Freitas et al., 2002;

Muigai et al., 2002). On the other hand, *B. tabaci* did not elicit behavioral responses

to the two corresponding chiral isomers: α-zingiberene and S-curcumene extracted

from ginger (Bleeker et al., 2011). Transformation of cultivated tomato with the gene

encoding *S. habrochaites* 7-epizingiberene synthase has become strongly repellent to

B. tabaci that reduced whitefly fecundity up to 87% (Bleeker et al., 2012). These

studies showed that *S. habrochaites* can exert strong repellent and toxic stresses

aganist *B. tabaci*, but it is unclear how it exerts its effects on the physiological and

molecular levels.

In order to achieve the goal, it was necessary to study the expression-related genes

that may be affected in the whitefly *B. tabaci* after exposure to the wild tomato

relative *S. habrochaites*.

2. Materials and methods

2.1 Insects rearing and plant materials

The MED cryptic species of whitefly was reared on tomato plants (Xianke 8) under

greenhouse conditions of 25 ± 2 °C, $65 \pm 5\%$ relative humidity (RH), and a 16:8 h

(L:D) photoperiod. The tomato plant was placed into cage measuring (60 cm × 60 cm

× 60 cm). The source of MED was collected from a greenhouse located in Beijing,

China and identified as MED by mitochondrial cytochrome oxidase I (mtCOI) gene

(Brown et al., 2005). One pair of newly emerged adults was used for colony initiation,

and the MED source after six generations was used for subsequent experiments.

2.2 Identification of wild tomato volatiles

S. habrochaites were placed into the cages measuring (60 cm × 60 cm × 60 cm) and

grown under greenhouse conditions of 25 ± 2°C, $65 \pm 5\%$ RH, and a 16:8 h (L:D)

photoperiod. Six-week-old plants were used for volatile analysis. We placed leaves of

a healthy tomato plant into a glass bottle (45 mL, cleman) and an SPME fiber

50/30-μm divinylbenzene/carboxen/polydimethylsiloxane (Supelco; Sigma-Aldrich,

http://www.sigmaaldrich.com) was used for analyte extraction. GC-MS analysis was

performed on an Agilent 7890 (Agilent Technologies, Tustin, CA, USA) gas

chromatography and Agilent 5975 mass selective detector.. Samples were separated

using both HP-5 and DB-WAX (both 30 m × 0.25 mm i.d., 0.25 mm film thickness,

Agilent Technologies, Tustin, CA, USA). Helium was used as carrier gas at a flow

rate of 1.7 mL/min, and the GC inlet was set in the split-less mode. The injector

temperature was 220 °C. The optimization of method parameters of SPME and

GC-MS were performed following to the method previously described (Li et al.,

2018). R-curcumene has been identified using GC-MS techniques by comparing its

retention time and mass spectrum with the same parameters from a reference

compound. As the 7-epizingiberene is thermally unstable (Bleeker et al., 2011),

therefore, the R-curcumene was used for conducting the behavioral and olfactory

studies. The synthesis of R-curcumene and S-curcumene was performed according to

the method previously reportedS (Song et al., 2012). The synthesized compounds

were confirmed by using various spectral technique like mass spectrometry and NMR

detection. The α-zingiberene was purchased from Sigma-Aldrich (St. Louis,

MO, USA).

2.3 RNA sequencing and bioinformatics analysis

To evaluate the effect of *S. habrochaites* on whiteflies at the molecular level,

RNA-sequencing was performed using MED females and males, which have been fed

on tomato Xianke 8 and the wild tomato *S. habrochaites* LA2175 for 6 and 24 h,

respectively. Sequencing libraries were constructed using strand-specific libraries.

FastQC (Andrews et al., 2010) was used to assess the quality of the sequencing data

while Fastp (Chen et al., 2018) was employed to remove adapters, contamination, and

duplicate sequences. The sequenced clean data were mapped to the MED whitefly

genome v1.0 (Xie et al., 2017) using HISAT2 (Kim et al., 2019)

ftp://www.whiteflygenomics.org/pub/whitefly/MED/v1.0. FeatureCounts (Liao et al.,

2014) was used to calculate the expression of each gene, edgeR (Roboison et al., 2010)

was used to analyze the difference in gene expression (we used Log2 Fold Change>1,

Adjusted P-value=0.05 as the threshold), rMATS (Shen et al., 2014) was used to

analyze the alternative splicing, and Sashimi plot was used to draw the alternative

splicing of genes. Finally, WGCNA (Langfelder and Horvath, 2008) was used to

calculate the correlation between the gene expression level (TPM values) and the

mortality phenotypes of whitefly.

3. Results and discussion

3.1 7-epizingiberene and R-curcumene are the main volatile components of *S.*

habrochaites

We synthesized a high-purity R-curcumene and confirmed its identity by mass

spectrometry and NMR (Supplementary Data 1). We found that 7-epizingiberene and

R-curcumene are the main volatile compounds of *S. habrochaites* using SPME and

GC-MS detection. GC-MS analysis revealed that these two volatile compounds were

not detected in the volatile blends of common tomato plants (Fig. 1). It was also found

that the peak areas ratio of 7-epizingiberene and R-curcumene in LA2175 was 2.37-

and 4.27-fold that of LA2329, respectively (Fig. 1). The data showed that LA2175

leaves had a relatively higher content of terpene volatiles than LA2329. This is

consistent with the biological analysis showing that LA2175 was more resistance to

whiteflies. Therefore, we chose and employed LA2175 for subsequent whitefly

high-throughput sequencing studies.

3.2 Expression of whitefly-related genes after stress exposure of *S. habrochaites*

A total of 24 samples of male and female whitefly adults (including two time points

and four treatments) were used for transcriptome sequencing (Fig. 2). About 21−24

million reads with an average length of 151 bp were aligned to the MED whitefly

reference genome by HISAT2 (Table 1). About 78% of the reads mapped to unique

loci (Table 1). After gene expression value analysis by featurecount and edgeR, we

found four significantly upregulated genes in wild tomato 6-h-treated female adults

(WF6) and wild tomato 24-h-treated female adults (WF24) (Fig. 2; Supplementary

Data 2). They were BTA016109 (Multicopper oxidase), BTA027707

(Sulfotransferase), BTA001651 (OHCU decarboxylase), and BTA011604 (Heat shock

protein). Wild tomato 6-h-treated male adults (WM6) and wild tomato 24-h-treated

female adults (WM24) also included four genes that are both significantly upregulated

(Fig. 2; Supplementary Data 2). They were BTA003886 (Heat shock protein),

BTA016667 (Unknown), BTA005900 (serine threonine-protein kinase SBK1-like),

and BTA011604 (Heat shock protein). One gene (BTA011604, Heat shock protein)

was significantly upregulated together in these four treatments. There were two genes

that were significantly downregulated in WF6 and WF24, BTA009028 (Unknown)

and BTA013802 (Farnesyl-diphosphate farnesyltransferase). Eighteen genes were

significantly downregulated in WM6 and WM24 (Fig. 2; Supplementary Data 2).

Among these, BTA022129 is solute carrier family 3, member 1, and the other genes

had unknown functions. These results indicated that the *S. habrochaites* treatment can

induce the expression of environmental stress genes such as heat shock protein in

whiteflies.

Alternative splicing is often related to the response of the organism to a stressful

environment (Sultan et al., 2008; Ling et al., 2015; Wang et al., 2008). We determined

if whiteflies were stressed by exposure to *S. habrochaites* and whether genes were

able to respond to the stress through alternative splicing. We performed alternative

splicing analysis on differently treated samples (comparison of male and female

samples treated with *S. habrochaites* and then treated with common tomato) through

rmats. There were five types of alternative splicing, including skipped exon (SE),

mutually exclusive exon (MXE), alternative to 5′splice site (A5SS), alternative to 3′

splice site (A3SS), and retained intron (RI). Among these, only SE and MXE had

significant differences between the two treatments (FDR <0.05), and the SE type

accounted for the main proportion (Table 2; Supplementary Data 3). GO enrichment

analysis showed that genes with alternative splicing of the SE type were significantly

enriched in three GO items including O-acetyltransferase activity, glycerol ether

metabolic process , and ether metabolic process (Supplementary Data 4). The

MXE-type alternative splicing genes were not significantly enriched.

Data availability conflict of interest

RNA-seq data have been deposited and are available in NCBI (BioProject ID is

PRJNA622388).

Conflict of interest: The authors declare no conflict of interest.

References

Andrews, S., FastQC: a quality control tool for high throughput sequence data. Babraham Bioinformatics, Babraham Institute, Cambridge, United Kingdom, 2010.

Bleeker, P. M., et al., 2009. The role of specific tomato volatiles in tomato-whitefly interaction. Plant Physiology. 151, 925-935.

Bleeker, P. M., et al., 2011. Tomato-produced 7-epizingiberene and R-curcumene act as repellents to whiteflies. Phytochemistry. 72, 68-73.

Bleeker, P. M., et al., 2012. Improved herbivore resistance in cultivated tomato with the sesquiterpene biosynthetic pathway from a wild relative. Proceedings of the National Academy of Sciences. 109, 20124-20129.

Chen, S., et al., 2018. fastp: an ultra-fast all-in-one FASTQ preprocessor. Bioinformatics. 34, i884-i890.

Chu, D., et al., 2006. The introduction of the exotic Q biotype of Bemisia tabaci from the Mediterranean region into China on ornamental crops. Florida Entomologist. 168-174.

De Barro, P., Driver, F., 1997. Use of RAPD PCR to distinguish the B biotype from other biotypes of Bemisia tabaci (Gennadius)(Hemiptera: Aleyrodidae). Australian Journal of Entomology. 36, 149-152.

Elbert, A., Nauen, R., 2000. Resistance of Bemisia tabaci (Homoptera: Aleyrodidae) to insecticides in southern Spain with special reference to neonicotinoids. Pest Management ence. 56.

Erdogan, C., et al., 2008. Insecticide resistance and biotype status of populations of the tobacco whitefly Bemisia tabaci (Hemiptera: Aleyrodidae) from Turkey. Crop Protection. 27, 600-605.

Freitas, J. A., et al., 2002. Inheritance of foliar zingiberene contents and their relationship to trichome densities and whitefly resistance in tomatoes. Euphytica. 127, 275-287.

Hu, J., et al., 2011. An extensive field survey combined with a phylogenetic analysis reveals rapid and widespread invasion of two alien whiteflies in China. PloS one. 6.

Ingham, V. A., et al., 2020. A sensory appendage protein protects malaria vectors from pyrethroids. Nature. 577, 1-5.

Kim, D., et al., 2019. Graph-based genome alignment and genotyping with HISAT2 and HISAT-genotype. Nature biotechnology. 37, 907-915.

Langfelder, P., Horvath, S., 2008. WGCNA: an R package for weighted correlation network analysis. BMC bioinformatics. 9, 559.

Leal, Walter, S., 2013. Odorant Reception in Insects: Roles of Receptors, Binding Proteins, and Degrading Enzymes. Annual Review of Entomology. 58, 373-391.

Liao, Y., et al., 2014. featureCounts: an efficient general purpose program for assigning sequence reads to genomic features. Bioinformatics. 30, 923-930.

Ling, Z., et al., 2015. Insect herbivory elicits genome□wide alternative splicing responses in N icotiana attenuata. The Plant Journal. 84, 228-243.

Muigai, S., et al., 2002. Mechanisms of resistance inLycopersicon germplasm to the whiteflyBemisia argentifolii. Phytoparasitica. 30, 347.

Pelosi, P., et al., 2017. Beyond chemoreception: Diverse tasks of soluble olfactory proteins in insects. Biological Reviews of the Cambridge Philosophical Society. 93.

Robinson, M. D., et al., 2010. edgeR: a Bioconductor package for differential expression analysis of digital gene expression data. Bioinformatics. 26, 139-140.

Shen, S., et al., 2014. rMATS: robust and flexible detection of differential alternative splicing from replicate RNA-Seq data. Proceedings of the National Academy of Sciences. 111, E5593-E5601.

Song, S., et al., 2012. Enantioselective iridium☐catalyzed hydrogenation of β, γ☐unsaturated carboxylic acids: An efficient approach to chiral 4☐alkyl☐4☐aryl butanoic acids. Angewandte Chemie International Edition. 51, 2708-2711.

Su, Q., et al., 2016. Tomato yellow leaf curl virus differentially influences plant defence responses to a vector and a non☐vector herbivore. Plant, cell & environment. 39, 597-607.

Suh, E., et al., 2014. Peripheral olfactory signaling in insects. Curr Opin Insect Sci. 6, 86-92.

Sultan, M., et al., 2008. A global view of gene activity and alternative splicing by deep sequencing of the human transcriptome. Science. 321, 956-960.

Wang, E. T., et al., 2008. Alternative isoform regulation in human tissue transcriptomes. Nature. 456, 470-476.

Wei, J., et al., 2017. Vector development and vitellogenin determine the transovarial transmission of begomoviruses. Proceedings of the National Academy of Sciences. 114, 6746-6751.

Xie, W., et al., 2017. Genome sequencing of the sweetpotato whitefly Bemisia tabaci MED/Q. GigaScience. 6, gix018.

Figure legends

Fig.1: GC−MS analysis of the volatile profiles of wild tomato *Solanum habrochaites* LA2175, LA2329, and common tomato Xianke 8. 1. R-curcumene, 2. 7-epizingiberene, 3 . *p*-cymene, 4. β-phellandrene, 5. elemene, 6. α-copaene, 7. unknow, 8.β-caryophyllene, 9. α-humulene.

Fig.2: Analysis of the RNA-Seq data of whiteflies treated by common tomato and wild tomato. (A) Principal component analysis (PCA) plot showing clustering of RNA-Seq of whiteflies. (B) Bar graph showing the number of upregulated (upper bars) and downregulated (lower bars) genes in whitefly male and female adults. (C) Venn diagram showing the number of induced differently expressed genes unique to or common within the four conditions. (D) Venn diagram showing the number of suppressed differently expressed genes unique to or common within the four conditions. WF6: wild tomato (*Solanum habrochaites*) 6 hours treated female adults; WM6: wild tomato 6 hours treated male adults; WF24: wild tomato 24 hours treated female adults; WM24: wild tomato 24 hours treated male adults.

Fig.1

Fig.2

Table legends

Table 1. Summary of the RNA-Seq Data. CKF6_1-3, whitefly females grown on common tomato for 6 h replicates 1–3. CKM6_1-3, whitefly males grown on common tomato for 6 h replicates 1–3. CKF24_1-3, whitefly females grown on common tomato for 24 h replicates 1–3. CKM24_1-3, whitefly males grown on common tomato for 24 h replicates 1–3. WF6_1-3, whitefly females grown on wild tomato LA2175 for 6 h replicates 1–3. WM6_1-3, whitefly females grown on wild tomato LA2175 for 6 h replicates 1–3. WF24_1-3, whitefly females grown on wild tomato LA2175 for 24 h replicates 1–3. WM24_1-3, whitefly males grown on wild tomato LA2175 for 24 h replicates 1–3.

Table 2. Genome-wide effects of *Solanum habrochaites* stress on alternative splicing of whitefly. WF6: wild tomato (*Solanum habrochaites*) 6-h-treated female adults; WM6: wild tomato 6-h-treated male adults; WF24: wild tomato 24-h-treated female adults; WM24: wild tomato 24-h-treated male adults. CKF6: common tomato 6-h-treated female adults; CKM6: common tomato 6-h-treated male adults; CKF24: common tomato 24-h-treated female adults; CKM24: common tomato 24-h-treated male adults.

Table 1

	No. of paired-end reads	No. of mapped paired-end reads	Proportion of mapped paired-end reads
CKF6_1	22729871	19170373	84.34%
CKF6_2	24263436	20446798	84.27%
CKF6_3	24587112	20852330	84.81%
CKM6_1	22886414	18199276	79.52%
CKM6_2	22949620	18146265	79.07%
CKM6_3	22716792	18100740	79.68%
CKF24_1	22159172	18759955	84.66%
CKF24_2	22190660	18620183	83.91%
CKF24_3	23055250	19278800	83.62%
CKM24_1	22912092	18384663	80.24%
CKM24_2	22576192	17744887	78.60%
CKM24_3	22106168	17342289	78.45%
WF6_1	21695473	17994225	82.94%
WF6_2	25184704	21089671	83.74%
WF6_3	24590350	20636222	83.92%
WM6_1	23754024	19112488	80.46%
WM6_2	24277496	19358875	79.74%
WM6_3	22011538	17624639	80.07%
WF24_1	22934225	19340432	84.33%
WF24_2	22669251	18994565	83.79%
WF24_3	24285715	20487429	84.36%
WM24_1	23292401	19060172	81.83%
WM24_2	21426439	17460405	81.49%
WM24_3	24205186	19790160	81.76%

Table2

AS Events Summary	WF6 vs CKF6	WM6 vs CKM6	WF24 vs CKF24	WM24 vs CKM24
SE	7	13	11	14
MXE	0	1	2	5
RI	0	0	0	0
A5SS	0	0	0	0
A3SS	0	0	0	0