The Epstein-Barr virus ubiquitin deconjugase BPLF1 regulates the activity of Topoisomerase II during virus replication

Jinlin Li[1,4] Noemi Nagy[1], Jiangnan Liu[1], Soham Gupta[1,5], Teresa Frisan[2] Thomas Hennig[3], Donald P. Cameron[1], Laura Baranello[1], and Maria G. Masucci[1§]

[1]Department of Cell and Molecular Biology, Karolinska Institutet, S-17165, Stockholm, Sweden

[2]Department of Molecular Biology, Umeå Center for Microbial Research, Umeå University, S-90187 Umeå, Sweden

[3]Institute for Virology and Immunobiology, University of Würzburg, 97078 Würzburg, Germany

[4]Current address: Institute of Medical Biochemistry and Microbiology, Uppsala University, S-75121, Uppsala, Sweden

[5]Current address: Division of Clinical Microbiology, Department of Laboratory Medicine Karolinska Institutet, S-14152, Huddinge, Sweden.

Short title: Regulation of Topoisomerase-II by BPLF1

§ Corresponding author: Maria G. Masucci, CMB, Biomedicum A6, Karolinska Institutet, S-17165, Stockholm, Sweden; e-post: maria.masucci@ki.se

Abstract

Topoisomerases are essential for the replication of herpesviruses but the mechanisms by which

the viruses hijack the cellular enzymes are largely unknown. We found that topoisomerase-II

(TOP2) is a substrate of the Epstein-Barr virus (EBV) ubiquitin deconjugase BPLF1. BPLF1

selectively inhibited the ubiquitination of TOP2 following treatment with topoisomerase

poisons, interacted with TOP2α and TOP2β in co-immunoprecipitation and *in vitro* pull-down,

stabilized Etoposide-trapped TOP2 cleavage complexes (TOP2cc) and promoted TOP2

SUMOylation, which halted the DNA-damage response and reduced Etoposide toxicity.

Induction of the productive virus cycle promoted the accumulation of TOP2βcc, enhanced

TOP2β SUMOylation, and reduced Etoposide toxicity in lymphoblastoid cell lines carrying

recombinant EBV encoding the active enzyme. Attenuation of this phenotype upon expression

of a catalytic mutant BPLF1-C61A impaired viral DNA synthesis and virus release. These

findings highlight a previously unrecognized function of BPLF1 in promoting non-proteolytic

pathways for TOP2cc debulking that favor cell survival and virus production.

Introduction

Epstein–Barr virus (EBV) is a human gamma-herpesvirus that establishes life-long persistent infections in most adults worldwide. The virus has been implicated in the pathogenesis of a broad spectrum of diseases ranging from infectious mononucleosis (IM) to a variety of lymphoid and epithelial cell malignancies including both Hodgkin and non-Hodgkin lymphomas, undifferentiated nasopharyngeal carcinoma, and gastric cancer(Shannon-Lowe & Rickinson, 2019).

Like other herpesviruses, EBV establishes latent or productive infections in different cell types. In latency, few viral genes are expressed resulting in the production of proteins and non-coding RNAs that drive virus persistence and cell proliferation(Babcock et al., 1998). In contrast, productive infection requires the coordinated expression of a large number of immediate early, early and late viral genes, which leads to the assembly of progeny virus and death of the infected cells(Hammerschmidt & Sugden, 2013). Although much of the EBV-induced pathology has been attributed to viral latency, the importance of lytic products in the induction of chronic inflammation and malignant transformation is increasingly recognized(Munz, 2019), pointing to inhibition of virus replication as a useful strategy for preventing EBV associated diseases.

EBV replication is triggered by the expression of immediate early genes, which transcriptionally activates a variety of viral and host cell factors required for subsequent phases of the productive cycle(Countryman & Miller, 1985; Feederle et al., 2000; Murata, 2014). Among the cellular factors, DNA topoisomerase-I and -II (TOP1 and TOP2) were shown to be essential for herpesvirus DNA replication(Hammarsten et al., 1996; M Kawanishi, 1993; Wang et al., 2008), raising the possibility that topoisomerase inhibitors may serve as antivirals. Indeed, non-toxic concentrations of TOP1 and TOP2 inhibitors were shown to suppress EBV-DNA replication(M Kawanishi, 1993), and different TOP1 inhibitors reduced the transcriptional

activity of the EBV immediate-early protein BZLF1 and the assembly of viral replication

complexes(Wang et al., 2009). However, the mechanisms by which the virus harnesses the

activity of these essential cellular enzymes remain largely unknown.

Topoisomerases sustain DNA replication, recombination and transcription by inducing

transient single or double-strand DNA breaks that allow the resolution of topological problems

arising from strand separation(Champoux, 2001; Wang, 2002). TOP2 homodimers mediate

DNA disentanglement by inducing transient double strand-breaks (DSBs) through the

formation of enzyme-DNA adducts, known as TOP2 cleavage complexes (TOP2ccs), between

catalytic tyrosine residues and the 5'ends of the DSBs(Nitiss, 2009). Following the passage of

the second DNA strand, TOP2 rejoins the DNA ends via reversion of the trans-esterification

reaction. While TOP2-indued DSBs are relatively frequent in genomic DNA(Morimoto et al.,

2019), failure to resolve TOP2ccs, as may occur upon endogenous or chemical stress that

inhibits TOP2 activity, results in the formation of stable TOP2-DNA adducts that hinder DNA

replication and transcription and trigger apoptotic cell death(Kaufmann, 1998). Thus, cellular

defense mechanisms attempt to resolve the TOP2ccs via proteolytic or non-proteolytic

mechanisms(Sun, Saha, et al., 2020). These may involve the displacement of TOP2 via

ubiquitin(Mao et al., 2001) or SUMO and ubiquitin-dependent(Sun, Miller Jenkins, et al., 2020)

proteasomal degradation, which, following the removal of residual peptide-DNA adducts by

the Tyrosyl-DNA phosphodiesterase-2 (TDP2) resolving enzyme(Gao et al., 2014; Pommier et

al., 2014), unmasks the DNA breaks and promotes activation of the DNA damage response

(DDR)(Pommier et al., 2014). Alternatively, SUMOylation may induce conformational

changes in the TOP2 dimer the expose the covalent bonds to the direct action of

TDP2(Schellenberg et al., 2017). Two TOP2 isozymes expressed in mammalian cells share

~70% sequence identity and have similar catalytic activities and structural features but are

differentially regulated and play distinct roles in biological processes(Nitiss, 2009). While

TOP2α is preferentially expressed in dividing cells and is essential for decatenating intertwined

sister chromatids during mitosis(Chen et al., 2015), TOP2β is the only topoisomerase expressed

in non-proliferating cells and is indispensable for transcription(Madabhushi, 2018)·(McKinnon,

2016).

Ubiquitin-specific proteases, or deubiquitinating enzymes (DUBs), regulate protein turnover by

disassembling poly-ubiquitin chains that target the substrate for proteasomal

degradation(Komander, 2009). Several human and animal viruses encode DUB homologs that

play important roles in the virus life cycle by promoting viral genome replication and inhibiting

the host antiviral response(Bailey-Elkin et al., 2017; Gastaldello et al., 2010; Kattenhorn et al.,

2005). In this study, we report that TOP2 interacts with and is a substrate of the DUB encoded

in the N-terminal domain of the EBV large tegument protein BPLF1 and provide evidence for

the capacity of BPLF1 to promote non-proteolytic pathways for the resolution of TOP2ccs,

which enhances cell survival and virus replication.

Results

BPLF1 selectively inhibits the degradation of TOP2 in cells treated with topoisomerase poisons

To investigate whether the EBV encoded DUB regulates the proteasomal degradation of topoisomerases, FLAG-tagged versions of the 325 amino acid long N-terminal catalytic domain of BPLF1 and an inactive mutant where the catalytic Cys61 was substituted with Ala (BPLF1-C61A) were stably expressed by lentivirus transduction in HEK-293T cells under the control of a Tet-on regulated promoter (HEK-rtTA-BPLF1/BPLF1-C61A cell lines). Inducible expression was monitored by probing immunoblots of cells treated for 24 h with increasing concentration of doxycycline (Dox) with antibodies to the FLAG or V5 tags (Fig. S1A). Although the steady-state levels of BPLF1-C61A appeared to be lower, which may be due to rapid turnover, both polypeptides were readily detected by anti-FLAG immunofluorescence in approximately 50% of the induced cells, while the fluorescence was weak or below detection in the remaining cells (Fig. S1B).

To monitor ubiquitin-dependent proteasomal degradation, HEK-rtTA-BPLF1/BPLF1-C61A cells cultured overnight in the presence or absence of Dox were treated with the TOP1 poison Campthothecine or the TOP2 poison Etoposide in the presence or absence of the proteasome inhibitor MG132, and topoisomerase levels were assessed by western blot. Campthothecine and Etoposide trap TOP1-DNA and TOP2-DNA covalent adducts, respectively(Pommier, 2013), while MG132 prevents the proteasomal degradation of stalled topoisomerase-DNA intermediates(Mao et al., 2001). As expected, TOP1 was efficiently degraded in Campthothecine treated cells (Fig. 1A and Fig. 1C upper panels), while treatment with Etoposide promoted the degradation of both TOP2α and TOP2β (Fig. 1B and 1C middle and

lower panels). The degradation was inhibited by treatment with MG132, confirming the

involvement of the proteasome in the clearance of poisoned topoisomerases. Expression of wild

type or mutant BPLF1 did not affect the Campthothecine-induced degradation of TOP1. In

contrast, expression of BPLF1 was accompanied by stabilization of both TOP2α and TPO2β in

Etoposide-treated cells, while the mutant BPLF1-C61A had no effect. Thus, the viral DUB

selectively inhibits the degradation of TOP2 isozymes by the proteasome.

Figure 1. BPLF1 selectively binds to TOP2 and inhibits the degradation of TOP2 in cells
treated with topoisomerase poisons. *HEK-293T cell expressing inducible FLAG-BPLF1 or*
FLAG-BPLF1-C61A were seeded into 6 well plates and treated with 1.5 µg/ml doxycycline
(Dox) for 24 h. After treatment for 3 h with 5 µM of the TOP1 poison Campthothecine (Cpt) or
6 h with 40 µM of the TOP2 poison Etoposide (Eto) with or without the addition of 10 µM
MG132, protein expression was analyzed in western blots probed with the indicated antibodies.
*GAPDH was used as the loading control. (**A**) Representative western blots illustrating the*
expression of TOP1 in control and Campthothecine treated cells. The treatment induced
degradation of TOP1 by the proteasome, which was not affected by the expression of BPLF1
*or BPLF1-C61A in Dox treated cells. (**B**) Representative western blots illustrating the*

expression of TOP2α and TOP2β in Etoposide treated cells. Expression of BPLF1 reduced the

Etoposide-induced degradation of both TOP2α and TOP2β while BPLF1-C61A had no

appreciable effect. (C) The percentage degradation of TOP1, TOP2α and TOP2β in

Campthothecine or Etoposide treated cells versus untreated controls was calculated from the

intensity of the specific bands recorded in two (TOP1) or three (TOP2α and TOP2β)

independent experiments using the ImageJ software. Data from HEK-rtTA-BPLF1/BPLF1-

C61A cultured in the absence of Dox were pulled. Statistical analysis was performed using

Student's t-test. **P< 0.01; ***P < 0.001; ns, not significant. (D) HEK293T cells transfected

with FLAG-BPLF1, FLAG-BPLF1-C61A, or FLAG-empty vector were treated with 40 μM

Etoposide for 30 min and cell lysates were either immunoprecipitated with anti-FLAG

conjugated agarose beads or incubated for 3 h with anti-TOP2α or TOPO2β antibodies

followed by the capture of immunocomplexes with protein-G coated beads. Catalytically active

and inactive BPLF1 binds to both TOP2α and TOP2β in both untreated and Etoposide treated

cells (upper panels). Conversely, TOP2α (middle panels) and TOP2β (lower panels) interacts

with both catalytically active and inactive BPLF1. Representative western blots from one of

two independent experiments where all conditions were tested in parallel are shown.

TOP2 is a BPLF1 substrate

To assess whether topoisomerases are direct substrates of BPLF1, we first investigated whether

they interact in cells and in pull-down assays performed with recombinant proteins. Lysates of

HEK-293T cells transiently transfected with FLAG-BPLF1 or FLAG-BPLF-C61A were

immunoprecipitated with antibodies recognizing FLAG, TOP1, TOP2α or TOP2β. In line with

the failure to rescue Campthothecine-induced degradation, BPLF1 did not interact with TOP1

(Fig. S2A), whereas both TOP2α and TOP2β were readily detected in western blots of the

FLAG immunoprecipitates and, conversely, BPLF1 was strongly enriched in the TOP2α and

TOP2β immunoprecipitates indicating that the proteins interact in cells (Fig. 1D). Of note, co-

immunoprecipitation was more efficient when BPLF1-C61A was the bite, suggesting that

TOP2 may be a substrate of the viral enzyme. To gain insight on the nature of the interaction,

equimolar concentration of yeast expressed FLAG-TOP2α, or a TOP2α mutant lacking the

unique C-terminal domain that is not conserved in the TOP2β isozyme, FLAG-TOP2α-ΔCTD,

were mixed with bacterially expressed His-BPLF1 and reciprocal pull-downs were performed

with anti-FLAG (Fig. S2B) or Ni-NTA coated beads (Fig. S2C). A weak BPLF1 band was

reproducibly detected in western blots of the FLAG-TOP2α pull-downs probed with a His-

specific antibody and, conversely, a weak FLAG-TOP2α band was detected in the His pull-

downs. The binding of BPLF1 to TOP2α was not affected by deletion of the TOP2α C-terminal

domain (Fig. S2D), pointing to a TOP2α and TOP2β shared domain shared as the likely site of

interaction. Notably, comparison of the efficiency of *in vitro* pull-down versus co-

immunoprecipitation suggests that binding may be stabilized by factors or TOP2 modifications

that are only present in cell lysates. To further investigate whether the viral DUB

deubiquitinates TOP2, BPLF1, TOP2α and TOP2β were immunoprecipitated from lysates of

control and Etoposide-treated HEK-293T cells transiently transfected with BPLF1 or BPLF1-

C61A and western blots were probed with a ubiquitin-specific antibody. The cell lysates were

prepared under denaturing conditions to exclude non-covalent protein interactions and working

concentrations of NEM and iodoacetamide were added to all buffers to inhibit DUB activity.

In line with the capacity of Etoposide to promote the proteasomal degradation of TOP2, smears

of high molecular weight species corresponding to ubiquitinated TOP2α and TOP2β were

detected in the immunoprecipitates of Etoposide-treated cells (Fig. 2A). The intensity of the

smears was strongly decreased in cells expressing catalytically active BPLF1, while the mutant

BPLF1-C61A had no appreciable effect, confirming that TOP2 is a bona-fide BPLF1 substrate.

Figure 2. BPLF1 deubiquitinates TOP2 and stabilizes TOP2cc. (**A**) HEK293T cells were transiently transfected with plasmids expressing FLAG-BPLF1, FLAG-BPLF1-C61A, or the empty FLAG vector, and aliquots were treated with 40 μM Etoposide for 30 min. TOP2α and TOP2β were immunoprecipitated from cell lysates prepared under denaturing conditions in the presence of DUB inhibitors and western blots were probed with antibodies to TOP2α, TOP2β and ubiquitin. The expression of catalytically active BPLF1 inhibits the ubiquitination of TOP2α and TOP2β in Etoposide treated cells. Western blots from one representative experiment out of three are shown in the figure. (**B**) HEK-rtTA-BPLF1 cells were treated with 1.5 μg/ml Dox for 24 h followed by treatment with 80 μM Etoposide for the indicated time with or without the addition of 10 μM MG132. RADAR assays were performed as described in Materials and Methods and TOP2 trapped in 10 μg DNA was detected in western blots using antibodies to TOP2α or TOP2β. Trapped TOP2 appeared as a major band of the expected size and a smear of higher molecular weight species. The intensity of the trapped TOP2α and TOP2β bands decrease over time in control untreated cells due to proteasomal degradation, while the decrease was significantly reduced upon expression of BPLF1 in Dox treated cells. Western blots from one representative experiment out of two are shown in the figure. (**C**) The percentage of Etoposide-induced TOP2 degradation was calculated from the intensity of the specific bands measured with the ImageJ software. MG132 prevented the degradation of

TOP2α and TOP2β trapped into TOP2cc in control BPLF1 negative cells whereas TOP2
degradation was significantly reduced in BPLF1 expressing cells. The mean ± SD of two
independent experiments is shown in the figure.

Degradation of TOP2 by the proteasome plays an important role in the debulking of persistent

TOP2ccs generated by topoisomerase poisons. To investigate whether the viral DUB may

interfere with this process, HEK-rtTA-BPLF1 cells cultured for 24 h in the presence or absence

of Dox were treated for 30 min or 4 h with Etoposide with or without addition of MG132, and

DNA-trapped TOP2 was detected by RADAR (rapid approach to DNA adduct recovery)

assays(Anand et al., 2018). Neither TOP2α nor TOP2β were detected in control DMSO treated

cells confirming that only covalently DNA-bound species are isolated by this method (Fig. 2B).

In Etoposide treated cells, TOP2α and TOP2β appeared as major bands of the expected size

and smears of high molecular weight species that are likely to correspond to different types of

post-translational modifications. Comparable amounts of trapped TOP2α and TOP2β were

detected in cells treated with Etoposide for 30 min, independently of BPLF1 expression or

MG132 treatment, indicating that neither treatment, either alone or in combination, affects the

formation of TOP2ccs. As expected, the intensity of the TOP2 band decreased after Etoposide

treatment for 4 h, which was inhibited by MG132, confirming the involvement of proteasome-

dependent degradation in the debulking of Etoposide-induced TOP2ccs. The degradation of

both TOP2α and TOP2β was significantly decreased at the 4 h time point in Dox treated cells,

resulting in levels of stabilization comparable to those achieved by treatment with MG132 (Fig.

2C). Thus, BPLF1 deubiquitinates and stabilizes TOP2 trapped in covalent DNA adducts. The

finding was independently confirmed in experiments where TOP2ccs were stabilized by

alkaline lysis(Ban et al., 2013) (Fig. S3A). Smears of high molecular weight species were

readily detected above the main TOP2β band in Dox-induced Etoposide-treated HEK-rtTA-

BPLF1 cells, whereas high molecular weight species were not detected when the blots were

probed with a TOP1 specific antibody, confirming that the high molecular weight species

correspond to DNA-trapped TOP2 (Fig. S3A). As expected, the intensity of the high molecular

weight species decreased with time in BPLF1 negative cells, and the decrease was inhibited by

MG132 confirming the involvement of proteasomal degradation. In cells expressing

catalytically active BPLF1, the intensity of the high molecular weight species remained

virtually constant over the observation time, resulting in significantly higher amounts of

residual TOP2ccs (Fig. S3B). Similar results were obtained when the blots were probed with

antibodies to TOP2α.

BPLF1 inhibits the detection of Etoposide-induced DNA damage and promotes

TOP2 SUMOylation and cell survival

The removal of TOP2 trapped in TOP2ccs induces a DNA damage response (DDR) that, while

limiting Etoposide toxicity, may also promote genomic instability and apoptosis(Lee et al.,

2016; Mao et al., 2001; Sciascia et al., 2020). To test whether the capacity of BPLF1 to stabilize

TOP2ccs interferes with DDR activation, HEK-rtTA-BPLF1/BPLF-C61A cells cultured in the

presence or absence of Dox for 24 h and then treatment with Etoposide for 1 h. The

accumulation of phosphorylated histone H2AX (γH2AX), a validated DDR marker(Mah et al.,

2010), was monitored by immunofluorescence in BPLF1 positive and negative cells. As

illustrated by representative fluorescence micrographs (Fig. 3A, upper panels) and plots of

fluorescence intensity in BPLF1 positive and negative cells (Fig. 3B, upper panels), a diffuse

γH2AX fluorescence was readily detected in Etoposide-treated BPLF1 negative cells and in

cells expressing the mutant BPLF-C61A.

Figure 3. BPLF1 selectively inhibits the detection of TOP2-induced DNA damage. *HEK-*
rtTA-BPLF1/BPLF1-C61A cells grown on cover-slides were treated with 1.5 μg/ml Dox for 24
h to induce the expression of BPLF1 followed by treatment for 1 h with 40 μM Etoposide or
0.5 μg/ml of the radiomimetic Neocarzinostatin (NCS) before staining with the indicated
*antibodies. (**A**) The cells were co-stained with antibodies against FLAG (red) and antibodies*
to γH2AX or 53BP1 (green) and the nuclei were stained with DAPI (blue). Expression of the
catalytically active BPLF1 was associated with a significant decrease of nuclear γH2AX
fluorescence and decreased formation of 53BP1 foci while the BPLF1-C61A mutant had no
effect. Neither the catalytically active nor the inactive BPLF1 affected the induction of γH2AX
in cells treated with NCS. Representative micrographs from one out of two experiments where
*all conditions were tested in parallel are shown. Scale bar = 10 μm. (**B**) Quantification of*
γH2AX fluorescence intensity and 53BP1 foci in BPLF1/BPLF1-C61A positive and negative
cells from the same image. The Mean ± SD of fluorescence intensity in at least 50 BPLF1-
positive and 50 BPLF1-negative cells recorded in each condition is shown. Statistical analysis
was performed using Student's t-test. ***P <0.001; ns, not significant.

Cells expressing active BPLF1 showed significantly weaker γH2AX fluorescence, suggesting

that the viral enzyme counteracts DDR activation. Accordingly, DNA repair was not triggered

as assessed by the impaired formation of 53BP1 foci in BPLF1 positive compared to negative

cells (Fig. 3A and 3B, middle panels). A comparable BPLF1-dependent decrease of γH2AX

fluorescence and formation of 53BP1 and BRCA1 foci was observed upon Etoposide treatment

in HeLa cells transiently transfected with BPLF1/BPLF1-C61A (Fig. S4), confirming that the

effect is not cell-type specific. To assess whether the failure to activate the DDR may be due to

the capacity of BPLF1 to target events downstream of the formation of DSBs, cells expressing

BPLF1/BPLF1-C61A were treated with the radiomimetic agent Neocarzinostain (NCS)(Povirk,

1996). Neither BPLF1 nor BPLF1-C61A altered the induction of γH2AX in NCS treated cells

(Fig. 3A and 3B, lower panels). Thus, BPLF1 selectively inhibits the DDR and DNA repair

responses triggered by Etoposide-induced DSBs.

In the absence of TOP2 degradation, TOP2cc may be resolved via a non-proteolytic process

whereby SUMOylation-dependent conformational changes expose the tyrosyl-DNA bond to

the activity of Tyrosyl-DNA-phosphodiesterase-2 (TDP2), which enables DSBs repair without

the need of nuclease activity(Schellenberg et al., 2017). To assess whether this pathway may

be engaged in BPLF1 expressing cells, Dox-treated HEK-rtTA-BPLF1 cells were exposed to

Etoposide for 30 min and western blots of TOP2ccs isolated by RADAR were probed with

antibodies to ubiquitin and SUMO2/3. As expected, smears of high molecular weight species

reacting with both ubiquitin- and SUMO2/3-specific antibodies were highly enriched in

Etoposide treated cells (Fig. 4A).

Figure 4. BPLF1 promotes TOP2 SUMOylation and cell viability following Etoposide treatment. (A) HEK-rtTA-BPLF1 cells were cultured for 24 h in the presence or absence of 1.5 μg/ml Dox and then treated with 80 μM Etoposide for 30 min followed by detection of DNA trapped TOP2 by RADAR assay. Western blots of proteins bound to 10 μg DNA were probed with antibodies to ubiquitin, SUMO2/3 and TOP2. The expression of BPLF1 was associated with strongly decreased ubiquitination of the TOP2ccs while SUMOylation was only marginally affected. (B) The intensity of the ubiquitin, SUMO2/3 and TOP2bcc specific bands was quantified by densitometry using the ImageJ software. Relative intensity was calculated as the % intensity in Dox-treated versus untreated cells after normalization to TOP2cc. Mean ± SD of two independent experiments. (C) HEK-rtTA-BPLF1/BPLF1-C61A cells were cultured for 24 h in the presence or absence of 1.5 μg/ml Dox and then treated overnight with the indicated concentration of Etoposide before assessing cell viability by MTT assays. The expression of catalytically active BPLF1 decreased the toxic effect of Etoposide over a wide range of concentrations while BPLF1-C61A had no appreciable effect. The mean ± SD of two independent experiments is shown.

Although the formation of TOP2ccs was not affected (Fig. 4A, lower panel), expression of the viral DUB was accompanied by a dramatic decrease of ubiquitinated species, while the intensity of the SUMO2/3 smear was largely unaffected (Fig 4A upper and middle panels and 4B). Thus, in addition to preventing the detection of TOP2-induced DNA damage, by inhibiting TOP2

ubiquitination BPLF1 may shift the processing of TOP2ccs towards non-proteolytic pathways,

which could counteract the toxic effect of Etoposide. To test this possibility, cell viability was

assessed by Thiazolyl blue tetrazolium bromide (MTT) assays in untreated and Dox-induced

HEK-rtTA-BPLF1/BPLF1-C61A cells following overnight exposure to increasing

concentrations of Etoposide. Similar levels of cell viability were observed in cells treated with

Etoposide in the absence of Dox whereas, in line with the hypothesized protective effect of

BPLF1, cell viability was consistently improved in cells expressing the active enzyme over a

wide range of Etoposide concentrations and the BPLF1-C61A mutant had no appreciable effect

(Fig. 4C).

BPLF1 regulates the activity of TOP2β during productive infection

In the final set of experiments, we asked whether physiological levels of BPLF1 regulate the

activity of TOP2 during the productive virus cycle in EBV infected cells. To this end, infectious

virus rescued from HEK-293-EBV cells carrying recombinant EBV expressing wild type or

mutant BPLF1(Gupta et al., 2018) was used to transform normal B-lymphocytes into

immortalized lymphoblastoid cell lines (LCL-EBV-BPLF1/BPLF1-C61A). To optimize the

induction of the productive virus cycle, the LCLs were stably transduced with a recombinant

lentivirus expressing the viral transactivator BZLF1 under the control of a tetracycline-

regulated promoter. Treatment with Dox induced the expression of early (BMRF1) and late

(BFRF3) viral antigens detected in western blots probed with specific antibodies (Fig. S5A),

and BPLF1 mRNA detected by qPCR (Fig. S5B). Of note, the induction of late antigens was

consistently weaker in LCL-EBV-BPLF1-C61A (Fig. S5A), pointing to impairment of the late

phase of the virus cycle.

Consistent with the establishment of a pseudo-S-phase where progression to G2/M is blocked

and cellular DNA synthesis is inhibited(Kudoh et al., 2003), induction of the productive cycle

was associated with a strong decrease of TOP2α mRNA (Fig. S5C) and protein levels (Fig. 5A)

while TOP2β protein and mRNA showed either no change or a small increase of protein levels

in cells expressing catalytically active BPLF1 (Fig. 5A and S5C).

Figure 5. BPLF1 regulates the expression and activity of TOP2β during productive infection.

The productive virus cycle was induced by treatment with 1.5 mg/ml Dox in LCL cells carrying

recombinant EBV encoding wild type or catalytic mutant BPLF1 and expressing a tetracycline

*regulated BZLF1 transactivator. (**A**) The expression of TOP2α and TOP2β was assessed by*

western blot and the intensity of the specific bands was quantified using the ImageJ software.

Induction of the productive cycle was associated with a highly reproducible downregulation of

TOP2α while TOP2β was either unchanged or slightly increased. The effect was stronger in

cells expressing wild type BPLF1. Representative western blots and quantification of specific

*bands in three to five independent experiments are shown. (**B**) The formation of TOP2bcc was*

investigated by RADAR assays in untreated and induced LCLs. Representative western blot

illustrating the significant increase of TOP2bcc upon induction of the productive virus cycle in

*LCL cells expressing catalytically active BPLF1. BPLF1-C61A had no appreciable effect. One representative western blots and quantification of the intensity of the TOP2b smears in three independent experiments are shown. Fold increase was calculated as the ratio between the smear intensity in control versus induced cells. *P<0.05. (**C**) TOP2β was immunoprecipitated from total cell lysates of control and induced LCLs and western blots were probed with antibodies to TOP2β, ubiquitin and SUMO2/3. Western blots illustrating the decreased ubiquitination and increased SUMOylation of TOP2β in cells expressing catalytically active BPLF1. One representative experiment out of three is shown. (**D**) The intensity of the bands corresponding to immunoprecipitated TOP2β, ubiquitinated and SUMOylated species was quantified using the ImageJ software and the SUMO/Ub ratio was calculated after normalization to the intensity of immunoprecipitated TOP2b. The mean ± SD of three independent experiments is shown. *P<0.05. (**E**) The productive cycle was induced in LCL-EBV-BPLF1/BPLF1-C61A by culture for 72 h in the presence 1.5 μg/ml Dox. After washing and counting, $5x10^4$ live cells were seeded in triplicate wells of 96 well plates and treated overnight with the indicated concentration Etoposide before assessing cell viability by MTT assays. The expression of catalytically active BPLF1 enhanced cell viability over a wide range of Etoposide concentration with BPLF1-C61A had no appreciable effect. The mean ± SD of cell viability in three independent experiments is shown. (**F**) The amount of cell associated and release EBV DNA was measures in the cell pellets ad supernatants after induction for 72 h. Fold induction was calculated relative to uninduced cells. Mean ± DS of 3 experiments.*

This was associated with a significant increase of TOP2βccs relative to uninduced cells or cells expressing the mutant BPLF1-C61A (Fig. 5B), and with decreased TOP2β ubiquitination (Fig. 5C top panel). As previously reported, higher molecular weight species detected by the SUMO2/3 specific antibody were increased in induced cells due to viral micro-RNA-dependent downregulation of RNF4(Li et al., 2017). SUMOylated species were also increased in immunoprecipitated TOP2β (Fig. 5C lower panel), resulting in a significant shift of the SUMO/ubiquitin ratio towards TOP2β SUMOylation in cells expressing catalytically active BPLF1 (Fig. 5D). As observed with the inducible HEK-rtTA-BPLF1 cell line, expression of

the active viral DUB counteracted the toxic effect of Etoposide (Fig. 5E). Furthermore, the

BPLF1-mediated regulation of TOP2β expression and ubiquitination was associated with

higher levels of viral DNA replication and efficient release of infectious virus particles as

measured by qPCR in cell pellets and culture supernatants (Fig. 5F).

Discussion

Although compelling evidence points to a pivotal role of topoisomerases in the replication of

herpesviruses and other DNA viruses(M. Kawanishi, 1993; Wang et al., 2009; Wang et al.,

2008), very little is known about the mechanisms by which the viruses harness the activity of

these cellular enzymes. In this study, we have shown that the ubiquitin deconjugases encoded

in the N-terminal domain of the EBV large tegument protein BPLF1 regulates the activity of

TOP2β during productive EBV infection by promoting the proteasome-independent debulking

of TOP2-DNA adducts, which favors cell survival and the faithful replication and transcription

of viral DNA. The findings highlight a previously unrecognized function of the viral enzyme

in hijacking cellular functions that enable efficient virus production. Our proposed model for

the activity of BPLF1 is shown in Fig. 6.

We found that the viral DUB that is physiologically released during productive infection via

caspase-1-mediated cleavage of the EBV large tegument protein BPLF1(Gastaldello et al.,

2013) selectively binds to TOP2α and TOP2β, and effectively counteracts their ubiquitination

and proteasomal degradation in cells treated with Etoposide (Fig. 1, Fig 2A). In the absence of

proteasomal degradation, TOP2ccs were stabilized (Fig 2B, 2C), which prevented the

unmasking of TOP2-induced DSBs (Fig 3) and promoted resistance to the toxic effect of

Etoposide (Fig 4C). Several lines of evidence suggest that the potent DDR triggered by the

proteolytic debulking of TOP2ccs may be detrimental to cell survival and genomic integrity.

Following the degradation of TOP2, protein-free DSBs engage multiple pathways for error-free

or error-prone repair, including MRE11 nuclease-dependent homologous recombination

(HR)(Hoa et al., 2016) and non-homologous end joining (NHRJ)(Gomez-Herreros et al., 2013;

Gomez-Herreros et al., 2017).   Recent findings suggest that a substantial fraction of the

Etoposide induced DSBs undergo extensive DNA end-resection(Sciascia et al., 2020), which

favors mispairing and the occurrence of chromosomal rearrangements that compromise cell

viability or promote genomic instability. Of note, these genotoxic effects were efficiently

counteracted by inhibition of the proteasome prior or during Etoposide treatment, supporting

the notion that the non-proteolytic resolution of TOP2ccs can minimize DSB misrepair and

promote genomic integrity(Sciascia et al., 2020).

We found that expression of the catalytically active viral deubiquitinase closely mimicked the

stabilization of TOP2ccs (Fig 2B) and inhibition of both DDR activation (Fig. 3) and Etoposide

toxicity (Fig. 4C and 5E) observed upon inhibition of the proteasome. While in line with the

notion that ubiquitination is strictly required for the targeting of substrates to the proteasome,

this finding points to the capacity of BPLF1 to shift the cellular strategy for TOP2cc debulking

towards proteasome-independent pathways that may ensure higher fidelity of DNA repair and

reduce toxicity. In this context, it is important to notice that, while inhibiting ubiquitination,

catalytic active BPLF1 did not affect the SUMOylation of TOP2cc in Etoposide treated cells

(Fig. 4A) and promoted the preferential SUMOylation of TOP2β during productive EBV

infection (Fig. 5C and 5D). SUMOylation plays multiple roles in the debulking of TOP2ccs. It

may mediate TOP2 proteolysis by serving as a recognition signal for ubiquitination mediated

by SUMO-targeted ubiquitin ligases such as RNF4(Sun, Miller Jenkins, et al., 2020) or may

recruit SprT-family metalloproteases, such as SPRTN(Lopez-Mosqueda et al., 2016) and

ARC/GCNA(Borgermann et al., 2019), that are involved in the proteasome-independent

proteolytic debulking of DNA-protein adducts. Interestingly, the activity of SPRTN is inhibited

by mono-ubiquitination(Stingele et al., 2016) and yet unpublished findings suggest that failure

to deubiquitinate SPRTN upon depletion of the cellular deubiquitinase USP11 leads to the

accumulation of unrepaired DNA-protein adducts(Perry et al., 2020). Thus, BPLF1 could

mimic the activity of the cellular DUB. In addition, SUMOylation of TOP2 by the ZNF451

ligase was shown to promote the non-proteolytic resolution of TOP2-DNA cross-links via

direct recruitment of TDP2 through a "split-SIM" SUMO2 engagement platform(Schellenberg

et al., 2017). SUMOylation was shown to alter the conformation of the trapped TOP2 dimers,

thereby facilitating the access of TDP2 to the tyrosyl-DNA covalent bond and promoting error-

free rejoining of the DSBs. This may be accomplished by the T4 DNA ligase(Gomez-Herreros

et al., 2013) or, upon removal of Etoposide, by TOP2 (Sciascia et al., 2020).

While the pivotal role of topoisomerases in both the latent and lytic replication of herpesviruses

is firmly established(Benson & Huang, 1988; Ebert et al., 1990; M Kawanishi, 1993), the roles

of the individual enzymes are not well understood. The torsion-relieving activity of TOP1 was

shown to be essential for reconstitution of the HSV replication machinery using purified viral

proteins(Nimonkar & Boehmer, 2004), and its recruitment to the viral replication complex was

required for the lytic origin (OriLyt)-driven replication of EBV(Wang et al., 2009) and

KSHV(Wang et al., 2008). Less is known about the function of the TOP2 isozymes, although

the importance of TOP2 is underscored by its upregulation during the productive cycle of

HCMV(Benson & Huang, 1990) and KSHV(Wang et al., 2008) despite a general host proteins

shutoff during virus replication. We have found that TOP2α mRNA is strongly downregulated

upon induction of the productive virus cycle in EBV infected cells (Fig. S5C) and the protein

becomes virtually undetectable in cells expressing catalytically active BPLF1 (Fig. 5A). While possibly related to the virus-induced arrest of the cell cycle in G1/S, the precise mechanism of this downregulation remains unknown. Nevertheless, our findings exclude a major role of TOP2α in the replication of the viral genome. In contrast, the expression of TOP2β was either not affected or slightly upregulated during productive infection, which is in line with the exclusive expression of this topoisomerase in resting cells and its essential role in transcription. Most importantly, we found that catalytically active BPLF1 was required for the accumulation of TOP2βccs in cell entering the productive cycle (Fig 5B), which, in the absence of topoisomerase poisons, is likely to indicate a significant increase of TOP2β activity driven by viral DNA replication and/or transcription. Conceivably, the capacity of BPLF1 to stabilize TOP2ccs and their resolution by non-proteolytic pathways that favor error-free repair and cell survival may be instrumental to ensure faithful and proficient replication and transcription of the viral genome. Of note, the BPLF1-mediated salvage of TOP2β from proteolytic disruption is likely to be reinforced by the concomitant downregulation of RNF4(Li et al., 2017), which may ensure that sufficient levels of the protein remain available throughout the productive cycle to sustain efficient virus production.

Aberrant expression of BPLF1 in the context of abortive lytic cycle reactivation has been reported in EBV associated malignancies such as undifferentiated nasopharyngeal carcinoma, NK-T cell lymphomas, and a subset of gastric cancers[(Peng et al., 2019),(Borozan et al., 2018)]. Etoposide and other topoisomerase poisons are used clinically as therapeutic anticancer agents against these malignancies(Delgado et al., 2018). Our data suggest that the expression of BPLF1 could be potentially used as a biomarker to predict the effectiveness of chemotherapeutic regimens that incorporate topoisomerase poison.

Figure 6. Model of TOP2 regulation by the BPLF1. *TOP2 (violet) trapped in TOP2ccs (yellow) is targeted for proteasomal degradation via SUMOylation (light blue) and ubiquitination (red) mediated by the SUMO ligase ZNF451, the SUMO-targeting ubiquitin ligase RNF4 and other cellular ubiquitin ligases, leading to the display of partially digested 5'-phosphotyrosyl-DNA adducts. Processing by the TDP2 resolvase generates protein-free DSBs that trigger the DDR and error-free or error-prone DNA repair. Imprecise repair leads to apoptosis and genomic instability. BPLF1 inhibits the degradation of Etoposide-poisoned TOP2, which inhibits activation of the DDR. In the absence of proteasomal degradation, SUMOylation may alter the conformation of the TOP2 dimer allowing direct access of TDP2 to the 5'-phosphotyrosyl-DNA bonds, which promotes error-free repair. During productive EBV infection, the concomitant expression of BPLF1 and downregulation of RNF4 favors the accumulation of SUMOylated TOP2β and the activation of non-proteolytic pathways for TOP2ccs debulking, which, in the absence of TOP2 poisons, may be mediated by TOP2 itself. This ensures the fidelity of virus replication and transcription and enhances cell survival and virus production.*

Acknowledgements

We are deeply grateful to Dr Arne Lindquist and Prof Nico Dantuma (CMB, Karolinska Institutet) for generously sharing knowledge and reagents and to Ms. Lisa Wohlgemuth (Ulm University, Germany) for technical assistance. This investigation was supported by grants awarded by the Swedish Cancer Society and The Medical Research Council. The work of SG and TH was partially supported by a grant awarded to the European ERA-NET eDEVILLI consortium.

Materials and Methods

Chemicals

IGEPAL CA-630 (NP40, I3021), Sodium dodecyl sulphate (SDS, L3771), N-Ethylmaleimide (NEM, E1271), Iodoacetamide (I1149), Sodium deoxycholate monohydrate (D5670), Triton X-100 (T9284), Bovine serum albumin (BSA, A7906), Tween-20 (P9416), Trizma base (Tris, 93349), Ethylenediaminetetraacetic acid disodium salt dehydrate (EDTA-E4884), Doxycycline cyclate (D9891), MG132 (M7449), Etoposide (E1383), Neocarzinostain (N9162) and Imidazole (I5513), were purchased from Sigma-Aldrich (St. Louis, MO, USA). RNase A (12091-021) and DNAzol (10503027) were purchased from Invitrogen (Carlsbad, CA, USA). Micrococcal nuclease (88216) was from Thermo Fisher Scientific (Rockford, lL, USA). Complete protease inhibitors cocktail tablets (04693116001), and phosphatase inhibitor cocktail (04906837001) were from Roche Diagnostic (Mannheim, Germany). Campthothecin was purchased from Selleckchem (Munich, Germany).

Antibodies

The antibodies that were used in this study: mouse anti-βactin (AC-15, 1:20000) and mouse anti-FLAG (F-3165, 1:7000; IF 1:300) from Sigma-Aldrich; rabbit anti-TOP1 (A302-589A,1:5000), TOP2α (A300-054A, 1:4000), TOP2β (A300-950A, 1:2000) and 53BP1(A300-272A,1:150) were from Bethyl Laboratories (Montgomery, Texas, USA); rabbit anti phospho-histone H2A.X clone 20E3 (#9718, IF:1:100) from Cell Signaling Technology (Danvers, Massachusetts); mouse monoclonal anti-ubiquitin (P4D1, sc-8017 1:1000), mouse anti-EBV-BZLF1 (sc-53904, 1:1000) and mouse anti-BRCA1 (D-9, sc-6954, IF 1:100) from Santa Cruz Biotechnology (Dallas, Texas, USA); mouse anti-SUMO2/3 (8A2, ab81371, 1:2000) from Abcam (Cambridge, MA, USA); monoclonal rat anti-EBV-BPLF1 (1:1500) (van Gent et al., 2014) from the MAB core facility, Helmholtz Center, Munich, Germany; mouse anti-EBV-BMRF1(1:10000) and rat anti-EBV-BFRF3 (1:1000) from Dr. Jaap M. Middeldorp (VU University Medical Center, Amsterdam, Netherlands). Alexa Fluor anti-rabbit-488 (A31570, 1:1000) and anti-mouse-555 (A315721, 1:1000) conjugated secondary antibodies raised in donkey were from Thermo Fisher (Waltham, Massachusetts, USA).

Plasmids and recombinant lentivirus vectors

Eukaryotic expression vectors encoding the N-terminal domain of the EBV large tegument protein 3xFLAG-BPLF1 (amino acid 1-235) and the catalytic mutant BPLF1-C61A(Ascherio & Munger, 2015) and the bacterial expression vector His-BPLF1(Gupta et al., 2019) were described previously. Lentiviral vectors encoding N-terminal 3xFLAG and V5 tandem tagged versions of BPLF1 aa 1-325 and the corresponding catalytic mutant BPLF1-C61A under control of the doxycycline-inducible pTight promoter were produced by cloning the corresponding open reading frames(Ascherio & Munger, 2015) into ta modified version of the pCW57.1 plasmid (gift from David Root, Addgene plasmid #41393). The Gal1/10 His6 TEV

Ura S. cerevisiae expression vector (12URA-B) was a gift from Scott Gradia (Addgene plasmid

#48304) a plasmid expressing human TOP2α was kindly provided by the James Berger (John

Hopkins School of Medicine, Baltimore, USA). The FLAG-TOP2α construct was created by

in-frame cloning the 3xFLAG coding sequence (amino acids

DYKDHDGDYKDHDIDYKDDDDKL) at the N-terminus of the TOP2α open reading frame.

All cloning was performed using the ligation independent cloning protocol from the QB3

Macrolab at Berkeley (macrolab.qb3.berkeley.edu). A recombinant lentivirus vector expressing

the coding sequence of the EBV transactivator BZLF1 under control of a tetracycline-regulated

promoter was constructed by cloning the open reading frame amplified with the primers 5'-

CGACCGGTATGATGGACCCAAACTCGAC-3' and 5'- CGACGCGTTTAGAAATTTAA

GAGATCCTCGTGT-3' into the Age I and Mlu I sites of the pTRIPZ lentiviral vector (Thermo

Fisher Scientific, USA). For virus production, HEK293FT cells were co-transfected with the

pTRIPZ-BZLF1, psPAX and pMD2G plasmids (Addgene, Cambridge, MA) using JetPEI

(Polyplus, Illkirch, France) according to the manufacture's protocol and cultured overnight in

complete medium. After refreshing the medium, the cells were cultured for additional 48 h to

allow virus production. Virus containing culture supernatant was briefly centrifuged and passed

through a 0.45 μm filter to removed cell debris before aliquoting and storing at -80°C for future

use.

Cell lines and transfection

HeLa cells (ATCC RR-B51S) and HEK293T (ATCC CRL3216) cell lines were cultured in

Dulbecco's minimal essential medium (DMEM, Sigma-Aldrich), supplemented with 10% FBS

(Gibco-Invitrogen) and 10 μg/ml ciprofloxacin (17850, Sigma-Aldrich) and grown in a 37°C

incubator with 5% CO2. Stable HEK-rtTA-BPLF1/BPLF1-C61A cell lines were produced by

lentiviral transduction followed by selection in medium containing 2μg/puromycin for 2 weeks.

Expression of FLAG-BPLF1/BPLF1-C61A was induced by treatment with 1.5 μg/ml

doxycycline and confirmed by anti-FLAG immunofluorescence and Western blot analysis.

Clones expressing high levels of the transduced proteins were selected by limiting dilution.

HeLa cells were transiently transfected with plasmids expressing FLAG-tagged version of

BPLF1/BPLF1-C61A using the lipofectamine 2000 (Invitrogen, California, USA) or jetPEI®

(Polyplus transfection, Illkirch FR) DNA transfection reagent according to the protocols

recommended by the manufacturer.

Production of EBV immortalized lymphoblastoid cell lines (LCLs)

Peripheral blood mononuclear cells were purified from Buffy coats (Blood Bank, Karolinska

University Hospital, Stockholm, Sweden) by Ficoll-Paque (Lymphoprep, Axis-shield PoC AS,

Oslo, Norway) density gradient centrifugation, and B-cells were affinity-purified using CD19

microbeads (MACS MicroBeads, Miltenyi Biotec, Bergisch Gladbach, Germany) resulting in

>95% pure B-cell populations. Infectious EBV encoding wild type or catalytic mutant BPLF1

were rescued for HEK293-EBV cells as previously described(Gupta et al., 2018). One million

B-cells were incubated in 1 ml virus preparation for 1.5 h at 37°C, followed by the addition of

fresh complete medium and incubation at 37°C in a 5% CO2 incubator until immortalized LCLs

were established. Sublines expressing a doxycycline-inducible BZLF1 transactivator were

produced by culturing 10^6 LCL cells with the recombinant lentivirus in presence of 8 μg/ml

polybrene (TR-1003-G, Sigma-Aldrich) for 24 hours followed by replacement of the infection

medium with fresh complete medium. The transduced cells were selected in medium containing

0.8μg/ml (LCL-BPLF1) or 0.25μg/ml (LCL-BPLF1-C61A) puromycin for one or two weeks.

Immunofluorescence

Transfected HeLa and HEK-Tta-BPLF1/BPLF1-C61A cells were grown on coverslips and

induced with 1.5 µg/ml doxycycline for 24 h. For immunofluorescence analysis, the cells were

fixed with 4% formaldehyde for 20 min, followed by permeabilization with 0.05% Triton X-

100 in PBS for 5 min and blocking in PBS containing 4% bovine serum albumin for 40 min.

After incubation for 1 h with primary antibodies and washing 3x5 min in PBS, the cells were

incubated for 1 h with the appropriate Alexa Fluor-conjugated secondary antibodies, followed

by washing and mounting in Vectashield-containing DAPI (Vector Laboratories, Inc.

Burlingame, CA, USA). Images were acquired using a fluorescence microscope (Leica DM

RA2, Leica Microsystems, Wetzlar, Germany) equipped with a CCD camera (C4742-95,

Hamamatsu, Japan). Fluorescence intensity was quantified using the ImageJ® software.

Western blots

Cells were lysed in RIPA buffer (25 mM Tris•HCl pH 7.6, 150 mM NaCl, 1% Igepal, 1%

sodium deoxycholate, 2% SDS) supplemented with protease inhibitor cocktail. Loading buffer

(Invitrogen) was added to each sample followed by boiling for 10 min at 100°C. The lysates

were fractionated in acrylamide Bis-Tris 4-12% gradient gel (Life Technologies Corporation,

Carlsbad, USA). After transfer to PVDF membranes (Millipore Corporation, Billerica, MA,

USA), the blots were blocked in TBS (VWR, Radnor, Pennsylvania, USA) containing 0.1%

Tween-20 and 5% non-fat milk, and the membranes were incubated with the primary antibodies

diluted in blocking buffer for 1 h at room temperature or over-night at 4°C followed by washing

and incubation for 1 h with the appropriate horseradish peroxidase-conjugated secondary

antibodies. The immunocomplexes were visualized by enhanced chemiluminescence (GE

Healthcare AB, Uppsala, SE). For detecting topoisomerase-DNA adducts after treatment with

topoisomerase poisons, the cells were lysed in alkaline buffer(Ban et al., 2013). Briefly, cells

treated for the indicated time with 5 µM Campthothecine or 80 µM Etoposide were lysed in

100 µl in buffer containing 200 mM NaOH, 2 mM EDTA, followed by the addition of 100 µl

of 1M HEPES buffer (pH 7.4). Nucleic acids were removed by addition of 10 µl 100 mM

CaCl2, 2 µl 1M DTT, and 200U of micrococcal nuclease followed by incubation at 37°C for

20 min. Seventy µl of 4xLDS loading buffer (Invitrogen) were added to each sample followed

by boiling for 10 min at 100°C before SDS-PAGE fractionation and western blot analysis.

Immunoprecipitation and pull-down assays

Cells were harvested 48h after transfection and lysed in NP40 lysis buffer (150 mM NaCl, 50

mM Tris-HCl pH7.6, 5mM MgCl2, 1mM EDTA, 1% Igepal, 1 mM DTT, 10% glycerol)

supplemented with protease/phosphatase inhibitor cocktail, 20 mM NEM and 20 mM

Iodoacetamide for 30 min on ice. For immunoprecipitations under denaturing condition the

lysis buffer was supplemented with 1% SDS followed by dilution to 0.1% SDS. For

BPLF1/BPLF1-C61A co-immunoprecipitation, the lysates were incubated for 3 h with 50 µl

anti-FLAG packed agarose affinity gel (A-2220; Sigma) at 4 °C with rotation. After washing

4 times with lysis buffer, the immunocomplexes were eluted with FLAG peptide (F4799;

Sigma). For TOP2α and TOP2β immunoprecipitation, specific antibodies were added to cell

lysates and incubated at 4°C for 3 h with rotation. The protein-antibody complexes were

captured with protein-G coupled Sepharose beads (GE Healthcare) by incubation at 4 °C for 1

h. The beads were washed 4 times with lysis buffer followed by boiling in 2xSDS-PAGE

loading for 10 min at 100°C. The production of 6xHis-BPLF1 in bacteria and purification of

the recombinant protein were done as previously described(Gupta et al., 2019). Recombinant

human TOP2α was expressed and purified according to a previously published protocol with

slight modifications(Lee et al., 2017). Briefly, URA-deficient yeast (kindly provided by Lena

Ström, CMB Karolinska Institutet) were transformed with the TOP2α expression plasmid,

grown initially in uracil-deficient media, then in YPLG (1% yeast extract, 2% peptone, 2%

sodium DL-lactate, 1.5% glycerol) before induction of expression by addition of 2% galactose.

The yeast was harvested by centrifugation and snap-frozen in liquid nitrogen. Proteins were

extracted using a cryo-mill, and the filtered lysate was passed sequentially through HisTrap

Excel nickel and HiTrap CP cation exchange columns (GE Healthcare) to purify the tagged

TOP2α protein before incubation overnight with His-tagged TEV protease. The following day,

the protein was passed through a HisTrap column to remove the cleaved His-tag and the TEV

protease. The TOP2α protein was further purified on a Superdex 200 16/60 column,

concentrated, and stored at -80°C. Relaxation and decatenation assays along with western

blotting were performed to confirm protein purity and activity. Equimolar concentration of

purified His-BPLF1 (0.35 μg) and FLAG-TOP2α (2 μg) were incubated in binding buffer (100

mM NaCl, 50 mM Tris-HCl, 1mM DTT, 0.5% Igepal) for 20 min at 4°C. Anti-FLAG agarose

affinity gel (A-2220; Sigma) or Ni-NTA beads (Qiagen) were added followed by incubation

for 60 min or 20 min at 4°C with rotation. The beads were intensively washed, and bound

proteins were eluted with FLAG peptide or 300 mM imidazole in buffer containing 50 mM

Tris-HCl pH 7.6, 50 mM NaCl and 1 mM DTT.

Rapid approach to DNA adduct recovery (RADAR) assay

TOP2ccs were isolated by RADAR assays as described(Kiianitsa & Maizels, 2013). Briefly,

cells cultured in 6 well plates were treated with 80 μM etoposide for 30 min or 4 h and then

lysed in 800 μl DNAzol. Following the addition of 400 μl absolute ethanol, the lysates were

cooled at -20°C and then centrifuged at 14000 rpm for 20 min at 4°C. After repeated washing

in 75% ethanol the nucleic acid pellets were dissolved in 100 μl H_2O at 37°C for 15 minutes,

followed by treatment with 100 μg/ml RNaseA. The concentration of DNA was measured and

10 μg DNA from each sample were treated with 250 U micrococcal nuclease supplemented

with 5 mM CaCl₂ before the addition of loading buffer and detection of trapped protein by

western blot.

Reverse transcription and real-time PCR

Total RNA was isolated using the Quick-RNA MiniPrep kit (Zymo Research, Irvine, CA, USA)

with in-column DNase treatment according to the instructions of the manufacturer. One

microgram of total RNA was reverse transcribed using SuperScript VILO cDNA Synthesis kit

(Invitrogen). PCR amplification was performed with the LC FastStart DNA master SYBR green

I kit in a LightCycler 1.2 instrument (Roche Diagnostic) using the following specific primers:

TOP1 5'-AGTGGAAAGAAGTCCGGCATGA-3', 5'-GCCAGTCCTTCTCACCCT TGAT-

3'; TOP2α 5'-AAGCCCAGCAAAAGGTTCCA-3', 5'-TGGCTTCAACAGCCTCCA AT-3';

TOP2β 5'-GGTTCGTGTAGAGGGGTCAA-3', 5'-CCCAGTTTCATCCAATTTGT C-3';

BPLF1 5'-CATACACCGTGCGAAAAGAA-3', 5'-GATGGCGGGTAATACATGCT-3'; and

MLN51 (Metastatic Lymph Node 51) 5'-CAAGGAAGGTCGTGCTGGTT-3', 5-AC

CAGACCGGCCACCAT-3'as endogenous control gene. The PCR reactions were denatured at

95°C for 10 min, followed by 40 cycles at 95°C for 8 sec, 60°C for 5 sed, 72°C for 8 sec. The

relative levels of mRNA were determined from the standard curve using MLN51 as reference.

MTT assay

For assay of cell viability, $2x10^4$ HEK-rtTABPLF1/BPLF1-C61A or $5x10^4$ LCLEBV-

BPLF1/BPLF1-C61A were plated in 150 μl medium in triplicate wells of a 96 well plate

without or with the addition of the indicated concentrations of Etoposide. After incubation for

20 h at 37°C in a 5% CO2 incubator, 50 μl culture medium containing 1 mg/ml

Methylthiazolyldiphenyl-tetrazolium bromide (MTT, M5655, Sigma-Aldrich) were added to

the wells followed by incubation for additional 4 h. The MTT formazan crystals produced by

mitochondrial dehydrogenases in living cells were solubilized by the addition of 50 μl 10%

SDS and O.D. was measured at 540 nm in a plate reader. Relative viability was calculated after

subtraction of the background O.D. of media alone.

EBV DNA replication and release of infectious virus

Virus replication and the release of infectious virus were monitored after induction with 1.5

μg/ml Doxycycline for 3 days in cell pellets and culture supernatants by quantitative PCR.

Briefly, DNA was isolated from cell pellets and culture supernatants cleared of cell debris by

centrifugation of 5 min at 14000 rpm and treated with 20 U/ml DNase I (Promega, Madison,

WI, USA) to remove free viral DNA, using the DNeasy Blood & Tissue Kit (Qiagen, Hilden,

Germany). Quantitative PCR was performed Quantitative PCR was performed as described

above with primers specific for a unique sequence in EBNA1 5'- GGCAGTGGACCTCAAAG

AAG-3', 5'-CTATGTCTTGGCCCTGATCC-3' and the cellular EF1α (Elongation

factor 1α) 5'-CTGAACCATCCAGGCCAAAT-3', 5'-GCCGTGTGGCAATCCAAT-3' as

reference. Virus replication was calculated as the ratio between the amount of viral DNA in

induced versus untreated cells.

REFERENCES

Anand, J., Sun, Y., Zhao, Y., Nitiss, K. C., & Nitiss, J. L. (2018). Detection of Topoisomerase
Covalent Complexes in Eukaryotic Cells. *Methods Mol Biol, 1703*, 283-299.
https://doi.org/10.1007/978-1-4939-7459-7_20
715

716   Ascherio, A., & Munger, K. L. (2015). EBV and Autoimmunity. *Curr Top Microbiol Immunol,*
717   *390*(Pt 1), 365-385. https://doi.org/10.1007/978-3-319-22822-8_15
718

719   Babcock, G. J., Decker, L. L., Volk, M., & Thorley-Lawson, D. A. (1998, Sep). EBV persistence in
720   memory B cells in vivo. *Immunity, 9*(3), 395-404. https://doi.org/10.1016/s1074-
721   7613(00)80622-6
722

723   Bailey-Elkin, B. A., Knaap, R. C. M., Kikkert, M., & Mark, B. L. (2017, Nov 10). Structure and
724   Function of Viral Deubiquitinating Enzymes. *J Mol Biol, 429*(22), 3441-3470.
725   https://doi.org/10.1016/j.jmb.2017.06.010
726

727   Ban, Y., Ho, C. W., Lin, R. K., Lyu, Y. L., & Liu, L. F. (2013, Oct). Activation of a novel ubiquitin-
728   independent proteasome pathway when RNA polymerase II encounters a protein
729   roadblock. *Mol Cell Biol, 33*(20), 4008-4016. https://doi.org/10.1128/MCB.00403-13
730

731   Benson, J. D., & Huang, E. S. (1988, Dec). Two specific topoisomerase II inhibitors prevent
732   replication of human cytomegalovirus DNA: an implied role in replication of the viral
733   genome. *J Virol, 62*(12), 4797-4800. https://doi.org/10.1128/JVI.62.12.4797-
734   4800.1988
735

736   Benson, J. D., & Huang, E. S. (1990, Jan). Human cytomegalovirus induces expression of cellular
737   topoisomerase II. *J Virol, 64*(1), 9-15. https://doi.org/10.1128/JVI.64.1.9-15.1990
738

739   Borgermann, N., Ackermann, L., Schwertman, P., Hendriks, I. A., Thijssen, K., Liu, J. C., Lans, H.,
740   Nielsen, M. L., & Mailand, N. (2019, Apr 15). SUMOylation promotes protective
741   responses to DNA-protein crosslinks. *EMBO J, 38*(8).
742   https://doi.org/10.15252/embj.2019101496
743

744   Borozan, I., Zapatka, M., Frappier, L., & Ferretti, V. (2018, Jan 15). Analysis of Epstein-Barr
745   Virus Genomes and Expression Profiles in Gastric Adenocarcinoma. *J Virol, 92*(2).
746   https://doi.org/10.1128/JVI.01239-17
747

748   Champoux, J. J. (2001). DNA topoisomerases: structure, function, and mechanism. *Annu Rev*
749   *Biochem, 70*, 369-413. https://doi.org/10.1146/annurev.biochem.70.1.369
750

751   Chen, T., Sun, Y., Ji, P., Kopetz, S., & Zhang, W. (2015, Jul 30). Topoisomerase IIalpha in
752   chromosome instability and personalized cancer therapy. *Oncogene, 34*(31), 4019-
753   4031. https://doi.org/10.1038/onc.2014.332
754

755 Countryman, J., & Miller, G. (1985, Jun). Activation of expression of latent Epstein-Barr
756   herpesvirus after gene transfer with a small cloned subfragment of heterogeneous
757   viral DNA. *Proc Natl Acad Sci U S A, 82*(12), 4085-4089.
758   https://doi.org/10.1073/pnas.82.12.4085
759

760 Delgado, J. L., Hsieh, C. M., Chan, N. L., & Hiasa, H. (2018, Jan 23). Topoisomerases as
761   anticancer targets. *Biochem J, 475*(2), 373-398. https://doi.org/10.1042/BCJ20160583
762

763 Ebert, S. N., Shtrom, S. S., & Muller, M. T. (1990, Sep). Topoisomerase II cleavage of herpes
764   simplex virus type 1 DNA in vivo is replication dependent. *J Virol, 64*(9), 4059-4066.
765   https://doi.org/10.1128/JVI.64.9.4059-4066.1990
766

767 Feederle, R., Kost, M., Baumann, M., Janz, A., Drouet, E., Hammerschmidt, W., & Delecluse, H.
768   J. (2000, Jun 15). The Epstein-Barr virus lytic program is controlled by the co-operative
769   functions of two transactivators. *EMBO J, 19*(12), 3080-3089.
770   https://doi.org/10.1093/emboj/19.12.3080
771

772 Gao, R., Schellenberg, M. J., Huang, S. Y., Abdelmalak, M., Marchand, C., Nitiss, K. C., Nitiss, J.
773   L., Williams, R. S., & Pommier, Y. (2014, Jun 27). Proteolytic degradation of
774   topoisomerase II (Top2) enables the processing of Top2.DNA and Top2.RNA covalent
775   complexes by tyrosyl-DNA-phosphodiesterase 2 (TDP2). *J Biol Chem, 289*(26), 17960-
776   17969. https://doi.org/10.1074/jbc.M114.565374
777

778 Gastaldello, S., Chen, X., Callegari, S., & Masucci, M. G. (2013). Caspase-1 promotes Epstein-
779   Barr virus replication by targeting the large tegument protein deneddylase to the
780   nucleus of productively infected cells. *PLoS Pathog, 9*(10), e1003664.
781   https://doi.org/10.1371/journal.ppat.1003664
782

783 Gastaldello, S., Hildebrand, S., Faridani, O., Callegari, S., Palmkvist, M., Di Guglielmo, C., &
784   Masucci, M. G. (2010, Apr). A deneddylase encoded by Epstein-Barr virus promotes
785   viral DNA replication by regulating the activity of cullin-RING ligases. *Nat Cell Biol,*
786   *12*(4), 351-361. https://doi.org/10.1038/ncb2035
787

788 Gomez-Herreros, F., Romero-Granados, R., Zeng, Z., Alvarez-Quilon, A., Quintero, C., Ju, L.,
789   Umans, L., Vermeire, L., Huylebroeck, D., Caldecott, K. W., & Cortes-Ledesma, F.
790   (2013). TDP2-dependent non-homologous end-joining protects against topoisomerase
791   II-induced DNA breaks and genome instability in cells and in vivo. *PLoS Genet, 9*(3),
792   e1003226. https://doi.org/10.1371/journal.pgen.1003226
793

794 Gomez-Herreros, F., Zagnoli-Vieira, G., Ntai, I., Martinez-Macias, M. I., Anderson, R. M.,
795   Herrero-Ruiz, A., & Caldecott, K. W. (2017, Aug 10). TDP2 suppresses chromosomal
796   translocations induced by DNA topoisomerase II during gene transcription. *Nat*
797   *Commun, 8*(1), 233. https://doi.org/10.1038/s41467-017-00307-y
798

799 Gupta, S., Yla-Anttila, P., Callegari, S., Tsai, M. H., Delecluse, H. J., & Masucci, M. G. (2018, Jan).
800   Herpesvirus deconjugases inhibit the IFN response by promoting TRIM25

autoubiquitination and functional inactivation of the RIG-I signalosome. *PLoS Pathog, 14*(1), e1006852. https://doi.org/10.1371/journal.ppat.1006852

Gupta, S., Yla-Anttila, P., Sandalova, T., Sun, R., Achour, A., & Masucci, M. G. (2019, Nov). 14-3-3 scaffold proteins mediate the inactivation of trim25 and inhibition of the type I interferon response by herpesvirus deconjugases. *PLoS Pathog, 15*(11), e1008146. https://doi.org/10.1371/journal.ppat.1008146

Hammarsten, O., Yao, X., & Elias, P. (1996, Jul). Inhibition of topoisomerase II by ICRF-193 prevents efficient replication of herpes simplex virus type 1. *J Virol, 70*(7), 4523-4529. https://www.ncbi.nlm.nih.gov/pubmed/8676478

Hammerschmidt, W., & Sugden, B. (2013, Jan 1). Replication of Epstein-Barr viral DNA. *Cold Spring Harb Perspect Biol, 5*(1), a013029. https://doi.org/10.1101/cshperspect.a013029

Hoa, N. N., Shimizu, T., Zhou, Z. W., Wang, Z. Q., Deshpande, R. A., Paull, T. T., Akter, S., Tsuda, M., Furuta, R., Tsutsui, K., Takeda, S., & Sasanuma, H. (2016, Nov 3). Mre11 Is Essential for the Removal of Lethal Topoisomerase 2 Covalent Cleavage Complexes. *Mol Cell, 64*(3), 580-592. https://doi.org/10.1016/j.molcel.2016.10.011

Kattenhorn, L. M., Korbel, G. A., Kessler, B. M., Spooner, E., & Ploegh, H. L. (2005, Aug 19). A deubiquitinating enzyme encoded by HSV-1 belongs to a family of cysteine proteases that is conserved across the family Herpesviridae. *Mol Cell, 19*(4), 547-557. https://doi.org/10.1016/j.molcel.2005.07.003

Kaufmann, S. H. (1998, Oct 1). Cell death induced by topoisomerase-targeted drugs: more questions than answers. *Biochim Biophys Acta, 1400*(1-3), 195-211. https://doi.org/10.1016/s0167-4781(98)00136-5

Kawanishi, M. (1993, Oct). Topoisomerase I and II activities are required for Epstein-Barr virus replication. *J Gen Virol, 74 (Pt 10)*, 2263-2268. https://doi.org/10.1099/0022-1317-74-10-2263

Kawanishi, M. (1993). Topoisomerase I and II are required for Epstein-Barr virus replication. *J Gen Virol, 74*, 2263-2269. https://doi.org/10.1099/0022-1317-74-10-2263

Kiianitsa, K., & Maizels, N. (2013, May). A rapid and sensitive assay for DNA-protein covalent complexes in living cells. *Nucleic Acids Res, 41*(9), e104. https://doi.org/10.1093/nar/gkt171

Komander, D. (2009, Oct). The emerging complexity of protein ubiquitination. *Biochem Soc Trans, 37*(Pt 5), 937-953. https://doi.org/10.1042/BST0370937

Kudoh, A., Fujita, M., Kiyono, T., Kuzushima, K., Sugaya, Y., Izuta, S., Nishiyama, Y., & Tsurumi, T. (2003, Jan). Reactivation of lytic replication from B cells latently infected with Epstein-Barr virus occurs with high S-phase cyclin-dependent kinase activity while

848   inhibiting cellular DNA replication. *J Virol, 77*(2), 851-861.
849   https://doi.org/10.1128/jvi.77.2.851-861.2003

851   Lee, J. H., Wendorff, T. J., & Berger, J. M. (2017, Dec 22). Resveratrol: A novel type of
852   topoisomerase II inhibitor. *Journal of Biological Chemistry, 292*(51), 21011-21022.
853   https://doi.org/10.1074/jbc.M117.810580

855   Lee, K. C., Bramley, R. L., Cowell, I. G., Jackson, G. H., & Austin, C. A. (2016, Mar 1). Proteasomal
856   inhibition potentiates drugs targeting DNA topoisomerase II. *Biochem Pharmacol, 103*,
857   29-39. https://doi.org/10.1016/j.bcp.2015.12.015

859   Li, J., Callegari, S., & Masucci, M. G. (2017, Apr). The Epstein-Barr virus miR-BHRF1-1 targets
860   RNF4 during productive infection to promote the accumulation of SUMO conjugates
861   and the release of infectious virus. *PLoS Pathog, 13*(4), e1006338.
862   https://doi.org/10.1371/journal.ppat.1006338

864   Lopez-Mosqueda, J., Maddi, K., Prgomet, S., Kalayil, S., Marinovic-Terzic, I., Terzic, J., & Dikic,
865   I. (2016, Nov 17). SPRTN is a mammalian DNA-binding metalloprotease that resolves
866   DNA-protein crosslinks. *Elife, 5*. https://doi.org/10.7554/eLife.21491

868   Madabhushi, R. (2018, Jun 29). The Roles of DNA Topoisomerase IIbeta in Transcription. *Int J*
869   *Mol Sci, 19*(7). https://doi.org/10.3390/ijms19071917

871   Mah, L. J., El-Osta, A., & Karagiannis, T. C. (2010, Apr). gammaH2AX: a sensitive molecular
872   marker of DNA damage and repair. *Leukemia, 24*(4), 679-686.
873   https://doi.org/10.1038/leu.2010.6

875   Mao, Y., Desai, S. D., Ting, C. Y., Hwang, J., & Liu, L. F. (2001, Nov 2). 26 S proteasome-mediated
876   degradation of topoisomerase II cleavable complexes. *J Biol Chem, 276*(44), 40652-
877   40658. https://doi.org/10.1074/jbc.M104009200

879   McKinnon, P. J. (2016, Nov). Topoisomerases and the regulation of neural function. *Nat Rev*
880   *Neurosci, 17*(11), 673-679. https://doi.org/10.1038/nrn.2016.101

882   Morimoto, S., Tsuda, M., Bunch, H., Sasanuma, H., Austin, C., & Takeda, S. (2019, Oct 30). Type
883   II DNA Topoisomerases Cause Spontaneous Double-Strand Breaks in Genomic DNA.
884   *Genes (Basel), 10*(11). https://doi.org/10.3390/genes10110868

886   Munz, C. (2019, Nov). Latency and lytic replication in Epstein-Barr virus-associated
887   oncogenesis. *Nat Rev Microbiol, 17*(11), 691-700. https://doi.org/10.1038/s41579-
888   019-0249-7

890   Murata, T. (2014, Jun). Regulation of Epstein-Barr virus reactivation from latency. *Microbiol*
891   *Immunol, 58*(6), 307-317. https://doi.org/10.1111/1348-0421.12155

893 Nimonkar, A. V., & Boehmer, P. E. (2004, May 21). Role of protein-protein interactions during
894 herpes simplex virus type 1 recombination-dependent replication. *J Biol Chem,*
895 *279*(21), 21957-21965. https://doi.org/10.1074/jbc.M400832200
896
897 Nitiss, J. L. (2009, May). DNA topoisomerase II and its growing repertoire of biological
898 functions. *Nat Rev Cancer, 9*(5), 327-337. https://doi.org/10.1038/nrc2608
899
900 Peng, R. J., Han, B. W., Cai, Q. Q., Zuo, X. Y., Xia, T., Chen, J. R., Feng, L. N., Lim, J. Q., Chen, S.
901 W., Zeng, M. S., Guo, Y. M., Li, B., Xia, X. J., Xia, Y., Laurensia, Y., Chia, B. K. H., Huang,
902 H. Q., Young, K. H., Lim, S. T., Ong, C. K., Zeng, Y. X., & Bei, J. X. (2019, Jun). Genomic
903 and transcriptomic landscapes of Epstein-Barr virus in extranodal natural killer T-cell
904 lymphoma. *Leukemia, 33*(6), 1451-1462. https://doi.org/10.1038/s41375-018-0324-5
905
906 Perry, M., Sundeep Kollala, S., Beigert, M., Su, G., Kodavati, M., Mallard, H., Kreling, N.,
907 Holfrook, A., & Ghosal, G. (2020). USP11 deubiquitinates monoubiquitinated SPRTN to
908 repair DNA-protein crosslinks. *bioRxiv, https://doi.org/10.1101/2020.06.30.180471*.
909
910 Pommier, Y. (2013, Jan 18). Drugging topoisomerases: lessons and challenges. *ACS Chem Biol,*
911 *8*(1), 82-95. https://doi.org/10.1021/cb300648v
912
913 Pommier, Y., Huang, S. Y., Gao, R., Das, B. B., Murai, J., & Marchand, C. (2014, Jul). Tyrosyl-
914 DNA-phosphodiesterases (TDP1 and TDP2). *DNA Repair (Amst), 19*, 114-129.
915 https://doi.org/10.1016/j.dnarep.2014.03.020
916
917 Povirk, L. F. (1996, Aug 17). DNA damage and mutagenesis by radiomimetic DNA-cleaving
918 agents: bleomycin, neocarzinostatin and other enediynes. *Mutat Res, 355*(1-2), 71-89.
919 https://doi.org/10.1016/0027-5107(96)00023-1
920
921 Schellenberg, M. J., Lieberman, J. A., Herrero-Ruiz, A., Butler, L. R., Williams, J. G., Munoz-
922 Cabello, A. M., Mueller, G. A., London, R. E., Cortes-Ledesma, F., & Williams, R. S. (2017,
923 Sep 29). ZATT (ZNF451)-mediated resolution of topoisomerase 2 DNA-protein cross-
924 links. *Science, 357*(6358), 1412-1416. https://doi.org/10.1126/science.aam6468
925
926 Sciascia, N., Wu, W., Zong, D., Sun, Y., Wong, N., John, S., Wangsa, D., Ried, T., Bunting, S. F.,
927 Pommier, Y., & Nussenzweig, A. (2020, Feb 14). Suppressing proteasome mediated
928 processing of topoisomerase II DNA-protein complexes preserves genome integrity.
929 *Elife, 9*. https://doi.org/10.7554/eLife.53447
930
931 Shannon-Lowe, C., & Rickinson, A. (2019). The Global Landscape of EBV-Associated Tumors.
932 *Front Oncol, 9*, 713. https://doi.org/10.3389/fonc.2019.00713
933
934 Stingele, J., Bellelli, R., Alte, F., Hewitt, G., Sarek, G., Maslen, S. L., Tsutakawa, S. E., Borg, A.,
935 Kjaer, S., Tainer, J. A., Skehel, J. M., Groll, M., & Boulton, S. J. (2016, Nov 17).
936 Mechanism and Regulation of DNA-Protein Crosslink Repair by the DNA-Dependent
937 Metalloprotease SPRTN. *Mol Cell, 64*(4), 688-703.
938 https://doi.org/10.1016/j.molcel.2016.09.031
939

940 Sun, Y., Miller Jenkins, L. M., Su, Y. P., Nitiss, K. C., Nitiss, J. L., & Pommier, Y. (2020, Nov). A
941   conserved SUMO pathway repairs topoisomerase DNA-protein cross-links by engaging
942   ubiquitin-mediated proteasomal degradation. *Sci Adv, 6*(46).
943   https://doi.org/10.1126/sciadv.aba6290
944
945 Sun, Y., Saha, L. K., Saha, S., Jo, U., & Pommier, Y. (2020, Oct). Debulking of topoisomerase
946   DNA-protein crosslinks (TOP-DPC) by the proteasome, non-proteasomal and non-
947   proteolytic pathways. *DNA Repair (Amst), 94*, 102926.
948   https://doi.org/10.1016/j.dnarep.2020.102926
949
950 van Gent, M., Braem, S. G., de Jong, A., Delagic, N., Peeters, J. G., Boer, I. G., Moynagh, P. N.,
951   Kremmer, E., Wiertz, E. J., Ovaa, H., Griffin, B. D., & Ressing, M. E. (2014, Feb). Epstein-
952   Barr virus large tegument protein BPLF1 contributes to innate immune evasion
953   through interference with toll-like receptor signaling. *PLoS Pathog, 10*(2), e1003960.
954   https://doi.org/10.1371/journal.ppat.1003960
955
956 Wang, J. C. (2002, Jun). Cellular roles of DNA topoisomerases: a molecular perspective. *Nat
957   Rev Mol Cell Biol, 3*(6), 430-440. https://doi.org/10.1038/nrm831
958
959 Wang, P., Rennekamp, A. J., Yuan, Y., & Lieberman, P. M. (2009, Aug). Topoisomerase I and
960   RecQL1 function in Epstein-Barr virus lytic reactivation. *J Virol, 83*(16), 8090-8098.
961   https://doi.org/10.1128/JVI.02379-08
962
963 Wang, Y., Li, H., Tang, Q., Maul, G. G., & Yuan, Y. (2008, Mar). Kaposi's sarcoma-associated
964   herpesvirus ori-Lyt-dependent DNA replication: involvement of host cellular factors. *J
965   Virol, 82*(6), 2867-2882. https://doi.org/10.1128/JVI.01319-07
966
967