Insights into biotic and abiotic modulation of ocean mesopelagic communities

Janaina Rigonato[1,17][*][#], Marko Budinich[2,3][*], Alejandro A. Murillo[4], Manoela C. Brandão[5], Juan J. Pierella Karlusich[6], Yawouvi Dodji Soviadan[5], Ann C. Gregory[7], Hisashi Endo[8], Florian Kokoszka[6,9], Dean Vik[7], Nicolas Henry[2], Paul Frémont[1], Karine Labadie[1], Ahmed A. Zayed[7], Céline Dimier[2,6,17], Marc Picheral[5,17], Sarah Searson[5], Julie Poulain[1,17], Stefanie Kandels[4,10], Stéphane Pesant[11,12], Eric Karsenti[6,10], The Tara Oceans coordinators[§], Peer Bork[6,13], Chris Bowler[6,17], Samuel Chaffron[3,17] , Colomban de Vargas[2,17], Damien Eveillard[3,17], Marion Gehlen[14], Daniele Iudicone[9], Fabien Lombard[5,17], Hiroyuki Ogata[8], Lars Stemmann[5,17], Matthew B. Sullivan[7,15], Shinichi Sunagawa[4,16], Patrick Wincker[1,17] and Olivier Jaillon[1,17][#]

Génomique Métabolique, Genoscope, Institut de Biologie François Jacob, Commissariat à l'Energie Atomique (CEA), CNRS, Université Evry, Université Paris-Saclay, 91000 Evry, France
Sorbonne Université, CNRS, Station Biologique de Roscoff, AD2M, UMR 7144, 29680 Roscoff, France
Université de Nantes, CNRS UMR 6004, LS2N, F-44000 Nantes, France
Structural and Computational Biology, European Molecular Biology Laboratory, Meyerhofstr. 1, 69117 Heidelberg, Germany
Sorbonne Université, CNRS, Laboratoire d'Océanographie de Villefranche (LOV), 06230 Villefranche-sur-Mer, France
Institut de Biologie de l'ENS (IBENS), Département de biologie, Ecole normale supérieure, CNRS, INSERM, Université PSL, 75005 Paris, France
Department of Microbiology, The Ohio State University, Columbus, OH 43214, USA
Bioinformatics Center, Institute for Chemical Research Kyoto University, Gokasho, Uji, Kyoto, 611-0011, Japan
Stazione Zoologica Anton Dohrn, Villa Comunale, 80121 Naples, Italy
Directors' Research European Molecular Biology Laboratory Meyerhofstr. 1 69117 Heidelberg Germany
MARUM, Center for Marine Environmental Sciences, University of Bremen, Bremen, Germany
PANGAEA, Data Publisher for Earth and Environmental Science, University of Bremen, Bremen, Germany
Department of Bioinformatics, Biocenter, University of Würzburg, Würzburg, Germany
Institut Pierre Simon Laplace, Laboratoire des Sciences du Climat et de l'Environnement, CEA, CNRS, Université Paris-Saclay, 91191 Gif-sur-Yvette cedex, France
Department of Civil, Environmental and Geodetic Engineering, The Ohio State University, Columbus OH 43214 USA
Department of Biology; Institute of Microbiology and Swiss Institute of Bioinformatics, ETH Zurich; Zurich, 8093; Switzerland
Research Federation for the study of Global Ocean Systems Ecology and Evolution, FR2022/Tara Oceans GOSEE, 3 rue Michel-Ange,75016 Paris, France

[*]These authors share the first authorship
Corresponding authors: jana_rigonato@protonmail.com (JR);
                         ojaillon@genoscope.cns.fr (OJ)

Running title: Mesopelagic community ecology

Abstract

Marine plankton mitigate anthropogenic greenhouse gases, modulate biogeochemical cycles, and provide fishery resources. Plankton is distributed across a stratified ecosystem of sunlit surface waters and a vast, though understudied, mesopelagic 'dark ocean'. In this study, we mapped viruses, prokaryotes, and pico-eukaryotes across 32 globally-distributed cross-depth samples collected during the *Tara* Oceans Expedition, and assessed their ecologies. Based on depth and O_2 measurements, we divided the marine habitat into epipelagic, oxic mesopelagic, and oxygen minimum zone (OMZ) eco-regions. We identified specific communities associated with each marine habitat, and pinpoint environmental drivers of dark ocean communities. Our results indicate that water masses primarily control mesopelagic community composition. Through co-occurrence network inference and analysis, we identified signature communities strongly associated with OMZ eco-regions. Mesopelagic communities appear to be constrained by a combination of factors compared to epipelagic communities. Thus, variations in a given abiotic factor may cause different responses in sunlit and dark ocean communities. This study expands our knowledge about the ecology of planktonic organisms inhabiting the mesopelagic zone.

Keywords: mesopelagic community, metabarcoding, plankton, pan-oceanic expedition, oxygen minimum zone

Introduction

Below the ocean's sunlit layer lies the mesopelagic zone that occupies around 20% of the global ocean volume [1]. The mesopelagic zone is biologically defined as starting where photosynthesis no longer occurs (<1% irradiance around 200m depth), down to its lower boundary where there is no detectable sunlight (around 1000m depth) [2]. This twilight ecosystem cannot rely on photoautotrophy, but sustains its energetic requirements by the combination of heterotrophic, chemoautotrophic, and chemo-mixotrophic metabolisms, together with physicochemical processes. Among the latter, the fraction of upper ocean productivity that escapes epipelagic recycling and sinks by gravity or is delivered by the daily migration of zooplankton constitutes an essential energy source in deep waters and is a vector for attached organisms [3].

The biodiversity and biomass in mesopelagic communities have been underestimated in the past [2, 4], and previous work showed these communities hold an enormous unexploited biological resource [5, 6, 1]. Mesopelagic organisms are considered to be a vast source of fat and protein, potentially becoming the primary source of global bioeconomy [7]. So far, efforts have been made to increase knowledge of mesopelagic mega/macrofauna by studying the abundance and diversity of nekton. These efforts are of great importance, given the rapid increase in the exploitation of this zone by nutraceutical and fisheries industries [6]. However, less attention has been devoted to the mesopelagic community's microscopic fraction, despite the pivotal role of the marine microbiome in biogeochemical cycles. The marine microbiome makes crucial links in the food web between primary production and dark ocean specialized consumers. Previous reports have shown stratification of planktonic communities with depth. In this regard, the mesopelagic zone displays a distinct assemblage of dsDNA viruses [8], giant viruses [9], prokaryotes [10, 11], and eukaryotes [12]. However, unlike the epipelagic layer, mesopelagic plankton diversity does not show the latitudinal diversity gradient trends from pole-to-pole, peaking at lower latitudes [13].

Among the studies conducted in mesopelagic zones, we highlight the efforts to explore regions of extreme conditions, such as oxygen minimum zones (OMZs). These zones are formed by

relatively old slowly upwelling waters, often lying below highly productive surface zones [14], and are currently increasing in volume in the oceans [15]. OMZ prokaryotic communities are well documented and predominated by taxa such as Nitrospira, Marinimicrobia, and anammox bacteria from the phylum Planctomycetes, while Thaumarchaeota abundance is frequently lower in these zones [16, 17, 18, 19, 20]. In contrast, knowledge of eukaryotic diversity in OMZs is still rudimentary, but a prevalence of specific taxa such as Ciliophora, Dinoflagellata, MALV, and Acantharia has been reported, together with a higher metabolic activity of these taxa [21, 22, 23]. Understanding plankton community structure and dynamics is fundamental to anticipate the impacts of global warming and acidification in these regions.

The last decades have seen a significant increase in large-scale oceanic surveys [24, 25, 26]. However, most mesopelagic community studies have been limited to geographically or ecologically fragmented regions, or to specific taxonomic groups, mainly because of the inherent difficulties of accessing this zone on a global scale [5]. Hence, these studies have given us a limited picture of community composition. Moreover, the factors influencing community structure, presumably a combination of biotic and abiotic factors [27, 28], have been little explored in the mesopelagic zone.

The present study takes advantage of the *Tara* Oceans large-scale survey conducted in different water layers using a systematic sampling protocol, spanning viruses to small eukaryote size fractions, to investigate the mesopelagic biome [29]. We capitalized on genomic data together with extensive contextual data and ocean geography to explore the particularities of mesopelagic communities compared to communities found in the euphotic zone. We also investigated potential water deoxygenation effects on these communities by comparing OMZ communities with those from well-oxygenated waters. This work expands our knowledge of the web of relationships underpinning mesopelagic plankton ecosystems on a broad geographic scale.

Materials and Methods

- **Sample collection and pre-processing**

The environmental and biological data were obtained during the *Tara* Oceans expedition

(2009- 2012) in 32 oceanographic stations located in the Indian Ocean (IO - 037, 038, 039),

Pacific Ocean (PO - 097, 098, 100, 102, 106, 109, 110, 111, 112, 122, 131, 132, 133, 135,

137, 138), South Atlantic Ocean (SAO - 068, 070, 072, 076, 078) and North Atlantic Ocean

(NAO - 142, 143, 144, 145, 146, 148, 149, 152) comprising tropical and subtropical regions

(Figure 1). Physico-chemical environmental data were obtained along a vertical profile at each

station. Temperature, salinity, and oxygen were measured using a CTD-rosette system with a

coupled dissolved oxygen sensor. Chlorophyll-a concentrations were measured using high-

performance liquid chromatography. Nutrient concentrations were determined using

segmented flow analysis. Metadata are available at PANGAEA [30, 31, 32, 33, 34, 35] -

(https://doi.pangaea.de/10.1594/PANGAEA.875582).

The vertical distribution of marine particles was investigated with an Underwater Vision Profiler

(UVP, [36, 37]) mounted on the CTD-Rosette. The UVP acquires images in a coherent water

volume (1 L) delimited by a light sheet issued from red light-emitting diodes. Automatic

identification of objects was made using Ecotaxa, based on a learning set of visually identified,

manually classified objects and associated features. Images were classified to distinguish

mesozooplankton from non-living objects and artifacts (e.g., detritic particles, fibers, and out-

of-focus objects).

Water vertical profiles of temperature and salinity generated from the CTD were used to identify

the water masses by plotting a temperature x salinity (T/S) diagram using the Ocean Data View

V 5.0 (ODV) software package [38].

Three different water layers were sampled: surface (SRF, 3-7 m), deep chlorophyll maximum

(DCM - depth identified according to the peak of chlorophyll-a fluorescence obtained *in situ*),

and mesopelagic (ranging from 200-1000 m) [39]. The planktonic community was sampled by

partitioning the seawater by filtering each sampled depth with different filter sizes [34]. Among

the mesopelagic zones, 13 of them were identified as deficient in oxygen and classified as

oxygen minimum zone (OMZ, stations IO - 037, 038, 039 / PO - 100, 102, 106, 109, 110, 111,

133, 135, 137, 138). The OMZ were categorized as suboxic: <10 µM O_2/kg seawater and

anoxic: (<0.003 µM/kg seawater or undetectable with most sensitive techniques, e.g., STOX

sensors) [Units of O_2 concentration: 1 mL.L^{-1}=1.43 mg. L^{-1}=44.64 µM] [22].

Our dataset comprises different organismal size-fractions from viruses (two dsDNA-virus

families *Podoviridae* and *Myoviridae* - hereafter named as phages and NCDLV giant viruses -

hereafter named as giruses) to pico-eukaryotes. Phage libraries were constructed from

seawater samples filtered at 0.22 µm, concentrated using iron chloride flocculation, and treated

with deoxyribonuclease (DNase). Girus *polB* and prokaryotic 16S rDNA sequences were

extracted from plankton metagenomes sequenced from 0.22–1.6 or 0.22-3 µm filters, and the

pico-eukaryote dataset was obtained by V9-18S rDNA marker amplification from 0.8-3 or 0.8-

5 µm filters. Details of sample preparation and sequencing procedures are described in Alberti

et al. [40].

Phage relative abundance was accessed through the search for the marker genes *gp23*

(*Myoviridae*) and *polA* (*Podoviridae*) in the protein collection GOV2.0 derived from

metagenomic sequencing described in Gregory et al. [8]. The girus abundance profile was

obtained from *polB* marker gene gathered from the OM-RGC.v2 catalog [11] as described in

Endo et al. [9]. The Prokaryote 16S rDNA marker derived from the metagenome assembly,

named 16S Mitag, is described in Sunagawa et al. [10]. Sequences matching "Eukaryota",

"chloroplasts", and "mitochondria" were removed from the final table. Clustering and annotation

of pico-eukaryote V9-18S rDNA amplicons are described in de Vargas et al. [41], and functional

annotation of taxonomically assigned V9-18S rDNA metabarcodes was improved afterward; in

this case, we conserved in the final data only sequences assigned to the "Eukaryota" domain.

We concatenated SRF and DCM samples for each taxonomic group to obtain an epipelagic

dataset (EPI). Counts of OTUs shared in SRF and DCM samples were summed. OTU

abundance was normalized by the total counts for each taxonomic group within each sample.

- **Ecological Analysis**

Epipelagic and Mesopelagic Community and Environmental differences

We applied an NMDS analysis based on the Bray-Curtis dissimilarity matrix on relative abundances using the 'metaNMDS' function from the vegan R package [42] to confirm community differences between epipelagic and mesopelagic layers. Homogeneity of the sampled environmental parameters was checked using the 'betadisper' function (Homogeneity of multivariate dispersions in the vegan package). The analysis was conducted using the Euclidean distance matrix of the environmental variables using the depths (epipelagic, mesopelagic) as group factor. A permutation test statistically confirmed the results.

Ecological model selection

We used the Species Abundance (SAD) and Rank Abundance Distribution curves (RAD: log abundances vs. rank order) to fit some of the most popular ecological models assessing how communities are assembled [43, 44]. Niche/deterministic models presume that the community is under selection due to biotic and abiotic interactions, while neutral/stochastic models assume that random processes structure the community such as drift, migration, birth, and death [45, 46]. SAD is essential for describing and understanding community assemblage and its management [44]. Further, by showing logarithmic species abundance against a rank order, RAD is commonly used to investigate a community's structure from observations made at one point in space and time [43, 44]. All the abundance distributions were fitted using 'fitsad' (sads R package - [47]), and 'radfit' (vegan) functions using maximum likelihood estimation. A set of candidate models was selected *a priori* to be tested by SAD: Log-series, Poisson-lognormal, Broken Stick, Power Law, and the neutral ecological model Zero-sum multinomial distribution. The neutral ecological model describes the SAD of a sample taken from a neutral metacommunity under random drift. The models selected to be tested by RAD were niche pre-emption models (geometric series or Motomura model), lognormal, Zipf, and Zipf–Mandelbrot,

and the null model that infers that the individuals are randomly distributed among observed

species. For a complete description of each model see Magurran and McGill [48].

The Akaike's Information Criterion (AIC) was used to evaluate the fitted model's quality based

on log-likelihood penalized by the number of estimated parameters. AIC estimates the loss of

information if the model is assumed for a given dataset. In this manner, models with lower AIC

values are selected as better fit conditions [49]. The AIC values produced for each model were

compared using the delta AIC (Δ_i). A Δ_i value <2 indicates equally likely models, values 3 <Δ_i

<9 indicate less likely models, and Δ_i > 10 for no likely models [49].

Ecological inferences and statistics

Ecological patterns were inferred using environmental variables to constrain the variation

observed in biological data for planktonic samples using Canonical Correspondence Analysis

(CCA) in the vegan R package. A set of physico-chemical variables for the discrete depths

were selected for the ecological inferences, such as nitrate (NO_3^-), oxygen, temperature,

salinity, density, and particles using particle flux UVP data. In order to avoid collinearity among

factors, the selected variables were checked for variance inflation factor using the vif.cca

function and tested for significance by 'anova' implemented in vegan with 999 permutations.

Each variable effect significance was tested individually using all the others as covariables

(independently from their order in the model) by applying the option 'margin' to the 'anova'

function in vegan.

Permutational multivariate analysis of variance (PERMANOVA) was performed with the

function 'adonis' in vegan to determine the relationship between mesopelagic community

composition and predefined water masses based on 999 permutations.

Organism Eco-region classification

In order to detect organisms specific to epipelagic (EPI), oxic mesopelagic (Oxic MES), and

OMZ eco-regions, only data containing both epipelagic and mesopelagic information were

considered: in total, 25 stations for giruses, prokaryotes, and pico-eukaryotes, and 13 for

phages. First, we discarded OTUs with fewer than 100 reads to remove biases due to rare

species' presence, and then recalculated the relative abundances for each dataset. Next, we

ran a Kruskal-Wallis test ('kruskal.test' from stats R package [50]) to detect differential OTU

relative abundances between eco-regions, followed by a Bonferroni correction to avoid Type I

error. Organisms with a p-value <0.05, indicating a difference within groups, were subject to a

post-hoc Dunn test ('dunn.test' from dunn.test R package [51]) to identify preferential Eco-

regions for each OTU. From these results, OTUs statistically equally abundant in all Eco-

regions or non-significant Kruskal-Wallis tests were assigned to the "ubiquitous" group. In

contrast, those with significant p-values were classified as EPI, Oxic MES, or OMZ if only the

corresponding Eco-region was elected according to the Dunn test. Organisms with no

significant differences between Oxic MES and OMZ were assigned to Core MES.

Co-occurrence Network

For investigation of associations between organisms across Eco-regions, a co-occurrence

network was inferred. In this analysis, phage samples were not included due to the lower

number of stations sampled. Therefore, samples for giruses, prokaryotes and pico-eukaryotes

from stations 038, 039, 068, 070, 072, 076, 078, 098, 100, 102, 109, 110, 111, 112, 122, 132,

133, 137, 138, 142, 145, 146, 148, 149 and 152 were retained. OTUs with a relative abundance

$<10^{-4}$ and counting fewer than 5 observations were discarded. Next, each sample was

normalized by applying a centered-log-ratio (CLR) transformation. Network inferences were

performed using Flashweave version 0.18 developed in Julia version 1.2 [52], using the

sensitive and heterogeneous mode.

We analyzed this global co-occurrence network by delineating communities (or modules) using

the Clauset-Newman-Moore algorithm [53]. These modules are subsets of OTUs, obtained by

maximizing the co-occurrences within the module and minimizing connections between them.

Next, we investigated modules enriched in OTUs from specific Eco-regions using Fisher's

exact test using the "fisher.test" function from the stats R package.

Results and Discussion

Leveraging the resources produced by the *Tara* Oceans project, we deciphered differences

between epipelagic and mesopelagic beta-diversity stratification, with a particular emphasis

on the role of environmental variables such as temperature, oxygen, salinity, NO_3^-, chlorophyll-

a, and particle flux (see Methods). As previously reported, we observed a stratification by depth

of the epipelagic and the mesopelagic communities (i.e., phages, giruses, prokaryotes, and

pico-eukaryotes) (Supplementary Figure 1). Consequently, we investigated differences among

epipelagic and mesopelagic sampling sites based on Euclidean distance of physicochemical

characteristics from each site. We observed a high dissimilarity gradient among sites for both

layers (Supplementary Figure 2a, b). Mesopelagic samples were spread in the plot, with most

of the points placed distant from the group centroid (located in the center of the cloud of points

identified for each group) (Supplementary Figure 2a). In contrast, epipelagic points displayed

a large variance due to a few samples positioned apart from the main cluster (Supplementary

Figure 2a). These results probably underlie the heterogeneity of environmental conditions

encountered in both sampled layers, and this environmental variation may be an important

factor that can directly influence community composition.

Community composition variations can also be shaped by four main eco-evolutionary

processes: selection, dispersal, drift, and speciation [54]. Mathematical models based on these

processes and applied to species abundance (SAD) and rank abundance (RAD) distributions

are historically used to infer the ecological or evolutionary mechanisms that structure a given

community [55]. Here, we addressed these biological processes to infer the abundance

distributions as observed in such natural communities [56, 57] in the most diverse

environments (terrestrial and aquatic) [45, 46, 43]. We used both SAD and RAD as evidence

of ecological processes related to variations in plankton community composition. When applied

at the local scale, we excluded random/neutral evolutive effects because all epipelagic and

mesopelagic communities fit the niche/deterministic ecological models. In fact, in all cases, we

found a Δ_i value <2 for both methods evaluated, which allows us to select one best-fitting model

without a subjective judgment (Supplementary Material Table 1 and 2) [49]. In these cases,

the community assemblages are mediated by a combination of environmental conditions,

interspecies interactions (competition, predation, or mutualisms), and species traits (for

instance, phototrophy, parasitism). Specifically, 93% and 65% of epipelagic and mesopelagic

samples better fit the Lognormal model (Poisson Lognormal SAD [58] and Lognormal RAD

[59]), respectively). These results suggest that EPI and MESO plankton populations are

affected by the combination of many independent variables, including competitive biotic

interactions and abiotic factors [43]. According to this model, the community has a broad and

elementary form of organization [57]. Ser-Giacomi et al. [60] stated that the eukaryotic rare-

biosphere (non-dominant OTU's) composed of "transient" or occasional taxa in the ocean sunlit

layer are mainly governed by dispersal/neutral events. In contrast, and in the case of our

findings, the dominant fraction of the assemblage follows idiosyncratic environmental

conditions.

Next, to quantify how much of the differences in the assemblages' variance can be explained

by environmental conditions, we employed canonical correspondence analysis (CCA) using

the environmental variables measured at discrete depths as constraint variables. The results

showed that the environment explains only a small fraction of community variance for both

layers (32% on average) (Figure 2). The phage assemblage was the exception, for which about

55% of the epipelagic variation and 65% of the mesopelagic variation could be explained by

the variables investigated (Figure 2). We tested the variance explained by single explanatory

variables individually. In contrast with epipelagic communities mainly governed by temperature

and oxygen, as observed here and elsewhere [10, 61, 8, 13, 12, 62], we could not identify a

single environmental predictor structuring entire mesopelagic assemblages. However, a few

different variables appeared to be significant for each group (Table 1, complete analysis in

Supplementary Material Table 3). Notably, our analysis identified oxygen as the main

mesopelagic driver for phage and prokaryote assemblages, confirming previous reports [20,

63, 64].

Even though we observed in the ordination-plots the distinction of OMZ and oxic mesopelagic

(Oxic MES) stations for giruses and pico-eukaryote assemblages (Figure 2, diamond and down

triangle), we could not disentangle the effect of oxygen from the other variables included in the

analyses (Table 1, complete analysis in Supplementary Material Table 3). This result shows

that these assemblages are probably affected by a combination of the predictors evaluated,

reflecting their need to cope with a broader environmental gradient that maximizes their niche-

space partitioning. Previous studies have identified oxygen as one of the main drivers of the

eukaryotic community structure in OMZ regions [65, 22, 23]. These studies mainly compared

community composition along the oxygen gradient within the water column depth, from the

surface downwards. However, depth stratification of plankton communities is evident even in

regions with high oxygen concentrations, so distinct parameters co-varying with depth must be

taken into account in addition to the oxygen gradients [66].

In addition to the physicochemical parameters, our results show that particle flux derived from

UVP measurements was also a significant variable structuring the phage assemblages in both

epipelagic and mesopelagic layers (Table 1, complete analysis in Supplementary Material

Table 3). This data supports previous reports about the high correlation of this environmental

factor with phages, finding possible relevance for the carbon pump's functioning in epipelagic

layers [67]. This observation may also reflect the association with virus inputs from overlaying

water layers via sinking particles [68].

In situ physico-chemical measurements have revealed the dynamics and fluctuating nature of

the ocean, even at short time scale [69]. The heterogeneity in mesopelagic layers given by

deep currents, and by the impact of the surface production, together with the low mixing levels,

may favor a diversification in the mesopelagic community living in different water masses,

leading to species adaptation-acclimation. The *Tara* Oceans expedition route included

samples from common or distinct water masses defined by temperature/salinity profiles - T/S, comprising regionally connected or unconnected stations. We identified nine different water masses in the mesopelagic sampled locations (Figure 3). We could confirm significant differences in mesopelagic communities sampled in different water masses based on the PERMANOVA test (Table 2). This result indicates that the oceanic patchiness created by distinct water masses can favor beta-diversity diversification, indicating it to be a critical component for mesopelagic community variation for all the assemblages studied (phages, giruses, prokaryotes, and pico-eukaryotes). Thus, we hypothesize that this result may be explained by two non-exclusive causes related to water masses: (i) past common origin among water masses that have drifted or (ii) constant connectivity by ocean circulation between sampled sites belonging to the same water mass.

Following the biotic control suggested by the deterministic model previously identified, we addressed another lingering question, resolving planktonic community signatures of Oxic MES and OMZ regions for epipelagic communities. For this, we classified OTUs based on their relative abundance into three eco-regions: 1) EPI, 2) Oxic MES, and 3) OMZ. OTUs were classified as Core MES when commonly present in Oxic MES and OMZ samples. The taxa that were either equally abundant in all three eco-regions or not statistically confirmed to a single eco-region were classified as ubiquitous (Supplementary Figure 3). Using this approach, we could identify ubiquitous taxa, that are likely to thrive in a wide range of environmental conditions, or that may be detected in mesopelagic samples due to the simple vertical movement of sinking particles. This classification should help avoid putative biases inherent of the metabarcoding methodology.

More specifically, we were able to identify Oxic MES and OMZ signatures mainly at the infra-taxonomic level (OTU-species) for all biotic groups investigated (Figure 4, Supplementary Figure 3, 4, 5, 6, 7). This reflected the wide ecological niche occupied by the different species at a higher taxonomic level (i.e. family). At the species level, we observed large taxonomic plasticity of OTUs that occurred equally in both Oxic MES and OMZ samples, called Core MES,

principally for girus assemblages. However, most OTUs are not yet classified at the infra-

taxonomic level (data not shown). This observation reflects the knowledge gap about the

biodiversity and functional plasticity of species thriving in this ecosystem.

The great majority of phage taxa occurred at similar abundance in all regions (ubiquitous)

(Supplementary Figure 3, 4). Surprisingly, this ubiquity is vertically linked at each independent

station (Supplementary Figure 4), supporting the seed-bank hypothesis raised by Brum et al.

[60], and the correlation to the sinking particles observed here. We observed taxa specific to

the mesopelagic layer in second place, mostly related to the OMZ eco-region (Figure 4a,

Supplementary Figures 3, 8). This mesopelagic specificity agrees with the sharp increase in

marine phage microdiversity following depth, as previously shown by Gregory et al. [8]. Our

results emphasize that one cause for phage stratification in the water column might be the

adaptation to the mesopelagic environment. Two hypotheses arise here, 1) the environment

acts as a strong driver, directly selecting phages independently of their hosts, and 2) there is

higher phage-host specificity in the mesopelagic layer, promoting phage selection. Following

the first hypothesis, we can posit that the environment can directly impact phage assemblage

composition. The direct contact with the environment of free phage particles (released from

their hosts) may reduce infectivity, degrade, or remove virus particles, and adversely affect

adsorption to the host [70]. This direct environmental effect over marine phages was reported

for different ionic gradients [71], daylight conditions, and temperature [72]. However, the

enrichment of prokaryotic OTUs specific to mesopelagic regions (Supplementary Figures 3,

8), especially in OMZs, does not exclude the phage-host indirect selection relationship.

We found fewer but abundant mesopelagic-specific girus OTUs in both Oxic and OMZ eco-

regions, indicating that giruses can be less diverse in the mesopelagic layer (Figure 4b,

Supplementary Figures 5, 3, 8). Also, giruses can encode genes such as transporters for

ammonium, magnesium, and phosphate that are important in marine oligotrophic areas [73].

This characteristic can improve the host's fitness in the short-term but ultimately favor girus

fecundity and endurance. This property is named NCLDV-mediated host reprogramming [73].

The great majority of mesopelagic girus OTUs were assigned to the Core MES group,

indicating that these entities may infect a wide range of hosts adapted to diverse environmental

conditions.

We could better distinguish the prokaryotic mesopelagic signatures between Oxic MES and

OMZ, confirming the influence of oxygen reported here and in previous studies [18, 19, 20]

(Figure 4c, Supplementary Figures 6, 3, 8). Among the planktonic microorganisms,

prokaryotes have been, so far, the most investigated group in OMZ regions, especially in the

Pacific Ocean [60, 19, 18]. We observed similar occurrence and abundance for the OMZ

signature taxa in the Indian Ocean stations (IO - 037, 038, 039) and in stations PO - 100, 137,

and 138 from the Pacific Ocean (Figure 4c). These Pacific stations are located in the open

ocean (PO - 137 and 138 located in the Equatorial upwelling zone and station PO - 100 in the

South Pacific Subtropical Gyre). They present a strong upwelling signature, disclosing an

intense decrease in oxygen concentration almost reaching shallow waters. Likewise, the

sampling stations in the Indian Ocean are located in well-stratified waters, markedly

characterized by the abrupt decrease of oxygen concentration below the thermohaline at 100-

120m depth. In all these stations, the oxygen concentration ranges from 0.83 to 3 µmol/kg,

characterizing functionally anoxic waters since this oxygen level cannot sustain aerobic

metabolisms [74]. The other OMZ stations in the Pacific Ocean (PO - 102, 109, 110, 111) are

located in coastal areas. Although they are also under the influence of upwellings presenting

an oxygen depletion, the oxygen level does not achieve anoxic conditions, and thus are

classified as suboxic waters. This microoxic environment is enough to completely change the

microbial metabolism delineating the community composition in those sites. Also, differences

in offshore and coastal upwelling formation, for instance, or the influence of river runoffs,

transporting anthropogenic nutrient enrichment from the continent to coastal zones [74], could

be crucial to support the differences in OMZ communities we observed.

The same combination of OMZ anoxic and suboxic samples was observed for the pico-

eukaryotic groups MALV-II and Diplonemida, suggesting these OTUs as the true OMZ

eukaryotic signatures (Figure 4d). Some OTUs of these groups exhibited similar occurrences

in the anoxic Indian and Pacific Oceans but not in suboxic samples from the Pacific Ocean.

However, we observed a lower number of pico-eukaryotic taxa in the OMZ eco-region, the

prevailing OTUs being specific to Oxic MES locations in most cases. Although the CCA

analysis did not disentangle the oxygen from the other variables to explain the pico-eukaryote

assemblage variations, here we can verify that OMZ conditions do act negatively on selection

of pico-eukaryotes in marine environments.

Another step to better understand mesopelagic community dynamics is to dissect the

ecological relationships among species that thrive in this layer. Co-occurrence networks can

indicate how the environment may structure the community acting as a filter for resident

species [76]. They can also give us glimpses of organisms' ecological interactions based on

species connectivity [76, 77]. Combining the girus, prokaryote, and pico-eukaryote data, we

inferred a network containing 6,154 nodes and 12,935 edges (Figure 5a, Table 3). Due to the

lower number of stations sampled for phages, we excluded this group from the analysis. We

found mainly positive relationships (94%), suggesting a predominance of putative biotic

interactions (e.g. competition, symbiosis) rather than taxa avoidance or exclusion. This

dominance of positive relations was also reported for epipelagic plankton communities [27,78].

The global network had a modularity value greater than 0.4 (Table 3), indicating that the

network has a modular structure [74]. Using a module detection algorithm, we were able to

identify 36 distinct modules in the global network. Three of them were mainly composed of

OTUs significantly enriched in mesopelagic OTUs (Oxic MES enriched module 1 and OMZ

enriched modules 4 and 17; Figure 5). Together, these three modules cover almost the total

richness found in the mesopelagic zone (Figure 5b), and present similar values for the average

degree, clustering coefficient, and average path length (Table 3). These parameters indicate

a network complexity [76], hinting at distinct ecological niches within the mesopelagic at the

level of investigated organismal fractions.

In more detail, the OMZ signature modules were composed of a few connected nodes (323

and 175 nodes for modules 4 and 17, respectively), potentially indicating two distinct OMZ

community niches. The Oxic MES module 1 counted more nodes composing the network

associations (731 nodes), and both modules presented a variation in taxonomic composition

and proportions. OMZ module 4 contained mainly prokaryotic (23%) and girus (55%) OTUs

(Figure 5c, d). Among the prokaryotes, we detected taxa previously determined as OMZ

signatures (Nitrospinae, Marinimicrobia SAR 406, and Planctomycetes). Module 17 is mainly

comprised of pico-eukaryotes (82%), notably MALV-II (14%) and Diplonemida (17%)

previously indicated as OMZ signatures. Module 1 is taxonomically more diverse but is mainly

comprised of giruses and pico-eukaryotes. These groups accounted for 36% and 55%,

respectively, of OTUs in this module (Figure 5c, d). Giruses contributed to 598 associations

(edges) in mesopelagic module 1, of which 177 occurred between giruses and pico-

eukaryotes. Giruses from the *Mimiviridae* family are the most numerous taxa in all three

mesopelagic modules. *Mimiviridae* is a very abundant family in the ocean, present in various

size ranges from piconanoplankton (0.8-5 µm) up to mesoplankton (180- 2,000 µm) [9, 79].

This observation supports our findings that giruses are a prosperous group in mesopelagic

waters, undertaking different strategies to endure in such environmental conditions. In all three

modules, we observed the presence of Foraminifera, of which some species can use nitrate

over oxygen as an electron acceptor, favoring their survival in OMZ regions [80].

Sugihara [57] affirms that a hierarchical niche structure can explain the lognormal abundance

pattern in communities, and this assertion is valid for small assemblages and for large

ensembles. Consistently, the observation of three distinct community modules supports the

evidence of the lognormal ecological model empirically defined locally for each assemblage.

Our results converge and suggest that the mesopelagic presents at least three well-defined

ecological niches (Oxic MES, OMZ-4 and OMZ-17), with established conditions and resources

(abiotic and biotic) that allow the survival of a given species in these environments. Differences

between OMZ and Oxic MES networks suggest a potential loss of connections and interactions

among mesopelagic community members, directly affecting ecosystem stability due to habitat

change.

Conclusions

In this study, we explored mesopelagic pico-plankton ecological structuring, and concluded

that this component of oceanic plankton is heterogeneous regarding its environmental

parameters. The ecological parameters drive mesopelagic community assemblages, as they

fit niche ecological models. Although we could not identify a single driver of community

composition (such as temperature for sunlit ocean) for all organisms, we could pinpoint the

relevance of oxygen for phages and prokaryotic fractions and the relation of the former with

particle flux. Also, we show that water masses defined by their T/S profiles can explain the

differences in the observed pico-plankton structure, pointing to the role of a set of

environmental parameters rather than single drivers for community composition.

By establishing Eco-regions (Epipelagic, Oxic MES, and OMZ), we were able to discriminate

specific OTUs for all fractions studied. While we recovered known markers for Oxic MES and

OMZ regions at high taxonomic levels, we also found that most of these OTU signatures are

observed at low taxonomic levels, which sometimes cannot be resolved using current

databases. Crossing these specific OTUs with co-occurrence networks, we identified three

niches with biotic and abiotic conditions that characterize mesopelagic waters.

The limiting access to data is usually the bottleneck for knowledge about mesopelagic

dynamics. Our study benefits from a more significant number of organism samples and distinct

oceanic provinces than previous ones, allowing us to combine data to derive an expanded

vision of mesopelagic composition. Our results emphasize the need for better understanding

of mesopelagic life, in particular by improving our knowledge about oxic and oxygen-depleted

mesopelagic-dwelling communities, especially as climate change can be expected to expand

marine OMZs shortly.

Acknowledgments

This study is part of the "Ocean Plankton, Climate and Development" project conducted by the Tara Ocean Foundation with the support of the French Facility for the Global Environment (FFEM). Rigonato J., Budinich M., Murillo A.A., Brandão M.C., Pierella Karlusich J.J. and Soviadan Y.D. received financial support from FFEM to execute the project. Brandão, M.C. also received financial support from Coordination for the Improvement of Higher Education Personnel of Brazil (CAPES 99999.000487/2016-03). Tara Oceans (which includes both the Tara Oceans and Tara Oceans Polar Circle expeditions) would not exist without the leadership of the *Tara* Ocean Foundation and the continuous support of 23 institutes (https://fondationtaraocean.org). We further thank the commitment of the following sponsors: CNRS (in particular Groupement de Recherche GDR3280 and the Research Federation for the study of Global Ocean Systems Ecology and Evolution, FR2022/Tara Oceans-GOSEE), European Molecular Biology Laboratory (EMBL), Genoscope/CEA, The French Ministry of Research, and the French Government 'Investissements d'Avenir' programs OCEANOMICS (ANR- 11-BTBR-0008), FRANCE GENOMIQUE (ANR-10-INBS-09-08), MEMO LIFE (ANR-10-LABX-54), and PSL* Research University (ANR-11-IDEX-0001-02). We also thank the support and commitment of agnès b. and Etienne Bourgois, the Prince Albert II de Monaco Foundation, the Veolia Foundation, Region Bretagne, Lorient Agglomeration, Serge Ferrari, World Courier. The global sampling effort was enabled by countless scientists and crew who sampled aboard the schooner *Tara* from 2009-2013, and we thank MERCATOR-CORIOLIS and ACRIST for providing daily satellite data during the expeditions. We are also grateful to the countries who graciously granted sampling permission. The authors declare that all data reported herein are fully and freely available from the date of publication, with no restrictions, and that all of the analyses, publications, and ownership of data are free from legal entanglement or restriction by the various nations whose waters the Tara Oceans expeditions sampled in. This article is contribution number XX of Tara Oceans.

Tara Oceans Coordinators (alphabetical order) Silvia G. Acinas, Marcel Babin, Peer Bork, Emmanuel Boss, Chris Bowler, Guy Cochrane, Colomban de Vargas, Michael Follows, Gabriel Gorsky, Nigel Grimsley, Lionel Guidi, Pascal Hingamp, Daniele Iudicone, Olivier Jaillon, Stefanie Kandels, Lee Karp-Boss, Eric Karsenti, Fabrice Not, Hiroyuki Ogata, Stéphane Pesant, Nicole Poulton, Jeroen Raes, Christian Sardet, Sabrina Speich, Lars Stemmann, Matthew B. Sullivan, Shinichi Sunagawa and Patrick Wincker.

Conflict of Interest

The authors declare that they have no conflict of interest.

References

1. Proud R, Cox MJ, and Brierley AS. Biogeography of the Global Ocean's Mesopelagic Zone. en. Current Biology 2017 Jan; 27:113–9. doi: 10.1016/j.cub.2016.11.003

2. Robinson C, Steinberg DK, Anderson TR, Ar´ıstegui J, Carlson CA, Frost JR, et al. Mesopelagic zone ecology and biogeochemistry – a synthesis. en. Deep Sea Research Part II: Topical Studies in Oceanography 2010 Aug; 57:1504–18. doi: 10.1016/j.dsr2.2010.02.018

3. Mestre M, Ruiz-Gonz´alez C, Logares R, Duarte CM, Gasol JM, and Sala MM. Sinking particles promote vertical connectivity in the ocean microbiome. en. Proceedings of the National Academy of Sciences 2018 Jul; 115:E6799–E6807. doi: 10.1073/pnas.1802470115

4. Whitman WB, Coleman DC, and Wiebe WJ. Prokaryotes: The unseen majority. en. Proceedings of the National Academy of Sciences 1998 Jun; 95:6578–83. doi: 10.1073/pnas.95.12.6578

5. Hidalgo M and Browman HI. Developing the knowledge base needed to sustainably manage mesopelagic resources. en. ICES Journal of Marine Science 2019 May; 76:609–15. doi: 10.1093/icesjms/fsz067

6. St. John MA, Borja A, Chust G, Heath M, Grigorov I, Mariani P, et al. A Dark Hole in Our Under- standing of Marine Ecosystems and Their Services: Perspectives from the Mesopelagic Community. en. Frontiers in Marine Science 2016 Mar; 3. doi: 10.3389/fmars.2016.00031

7. Glover AG, Wiklund H, Chen C, and Dahlgren TG. Managing a sustainable deep-sea 'blue economy' requires knowledge of what actually lives there. eLife 2018 Nov; 7. Ed. by Groll H and Rodgers PA:e41319. doi: 10.7554/eLife.41319

8. GregoryAC, ZayedAA, Conceição-NetoN, TempertonB, BolducB, AlbertiA,et al. MarineDNA Viral Macro- and Microdiversity from Pole to Pole. en. Cell 2019 Apr :S0092867419303411. doi: 10.1016/j.cell.2019.03.040

9. Endo H, Blanc-Mathieu R, Li Y, Salazar G, Henry N, Labadie K, et al. Biogeography of marine giant viruses reveals their interplay with eukaryotes and ecological functions. en. Nature Ecology & Evolution 2020 Sep. doi: 10.1038/s41559-020-01288-w

10. Sunagawa S, Coelho LP, Chaffron S, Kultima JR, Labadie K, Salazar G, et al. Structure and function of the global ocean microbiome. en. Science 2015 May; 348:1261359–1261359. doi: 10.1126/science. 1261359

11. Salazar G, Paoli L, Alberti A, Huerta-Cepas J, Ruscheweyh HJ, Cuenca M, et al. Gene Expression Changes and Community Turnover Differentially Shape the Global Ocean Metatranscriptome. en. Cell 2019 Nov; 179:1068–1083.e21. doi: 10.1016/j.cell.2019.10.014

12. Giner CR, Pernice MC, Balague ́ V, Duarte CM, Gasol JM, Logares R, et al. Marked changes in diversity and relative activity of picoeukaryotes with depth in the world ocean. en. The ISME Journal 2020 Feb; 14:437–49. doi: 10.1038/s41396-019-0506-9

13. Ibarbalz FM, Henry N, Brand ̃ao MC, Martini S, Busseni G, Byrne H, et al. Global Trends in Marine Plankton Diversity across Kingdoms of Life. en. Cell 2019 Nov; 179:1084–1097.e21. doi: 10.1016/j. cell.2019.10.008

14. Capone DG and Hutchins DA. Microbial biogeochemistry of coastal upwelling regimes in a changing ocean. en. Nature Geoscience 2013 Sep; 6:711–7. doi: 10.1038/ngeo1916

15. Breitburg D, Levin LA, Oschlies A, Gr ́egoire M, Chavez FP, Conley DJ, et al. Declining oxygen in the global ocean and coastal waters. en. 2018 :13

16. Stevens H and Ulloa O. Bacterial diversity in the oxygen minimum zone of the eastern tropical South Pacific. en. Environmental Microbiology 2008 May; 10:1244–59. doi: 10.1111/j.1462-2920.2007. 01539.x

17. Stewart FJ, Ulloa O, and DeLong EF. Microbial metatranscriptomics in a permanent marine oxygen minimum zone: OMZ community gene expression. en. Environmental Microbiology 2012 Jan; 14:23– 40. doi: 10.1111/j.1462-2920.2010.02400.x

18. Ulloa O, Wright JJ, Belmar L, and Hallam SJ. Pelagic Oxygen Minimum Zone Microbial Communities. en. The Prokaryotes. Ed. by Rosenberg E, DeLong EF, Lory S, Stackebrandt E, and Thompson F. Berlin, Heidelberg: Springer Berlin Heidelberg, 2013 :113–22. doi: 10.1007/978-3-642-30123-0_45

19. Ulloa O and Pantoja S. The oxygen minimum zone of the eastern South Pacific. en. Deep Sea Research Part II: Topical Studies in Oceanography 2009 Jul; 56:987–91. doi: 10.1016/j.dsr2.2008.12.004

20. Wright JJ, Konwar KM, and Hallam SJ. Microbial ecology of expanding oxygen minimum zones. en. Nature Reviews Microbiology 2012 Jun; 10:381–94. doi: 10.1038/nrmicro2778

21. Duret MT, Pachiadaki MG, Stewart FJ, Sarode N, Christaki U, Monchy S, et al. Size-fractionated diversity of eukaryotic microbial communities in the Eastern Tropical North Pacific oxygen minimum zone. en. FEMS Microbiology Ecology 2015 May; 91. doi: 10.1093/femsec/fiv037

22. Orsi W, Song YC, Hallam S, and Edgcomb V. Effect of oxygen minimum zone formation on communities of marine protists. en. The ISME Journal 2012 Aug; 6:1586–601. doi: 10.1038/ismej.2012.7

23. Parris DJ, Ganesh S, Edgcomb VP, DeLong EF, and Stewart FJ. Microbial eukaryote diversity in the marine oxygen minimum zone off northern Chile. en. Frontiers in Microbiology 2014 Oct; 5. doi: 10.3389/fmicb.2014.00543

24. Rusch DB, Halpern AL, Sutton G, Heidelberg KB, Williamson S, Yooseph S, et al. The
Sorcerer II Global Ocean Sampling Expedition: Northwest Atlantic through Eastern
Tropical Pacific. en. PLOS Biology 2007 Mar; 5:e77. doi: 10.1371/journal.pbio.0050077

25. Karsenti E, Acinas SG, Bork P, Bowler C, De Vargas C, Raes J, et al. A Holistic
Approach to Marine Eco-Systems Biology. en. PLoS Biology 2011 Oct; 9:e1001177. doi:
10.1371/journal.pbio.1001177

26. Pernice MC, Forn I, Gomes A, Lara E, Alonso-Sáez L, Arrieta JM, et al. Global
abundance of planktonic heterotrophic protists in the deep ocean. en. The ISME Journal
2015 Mar; 9:782–92. doi: 10.1038/ismej.2014.168

27. Lima-Mendez G, Faust K, Henry N, Decelle J, Colin S, Carcillo F, et al. Determinants of
community structure in the global plankton interactome. en. Science 2015 May;
348:1262073–1262073. doi: 10. 1126/science.1262073

28. Louca S, Hawley AK, Katsev S, Torres-Beltran M, Bhatia MP, Kheirandish S, et al.
Integrating bio- geochemistry with multiomic sequence information in a model oxygen
minimum zone. en. Proceedings of the National Academy of Sciences 2016 Oct;
113:E5925–E5933. doi: 10.1073/pnas.1602897113

29. Ardyna M, Tara Oceans Consortium C, and Tara Oceans Expedition P. Environmental
context of all station from the Tara Oceans Expedition (2009-2013), about the annual
cycle of key parameters estimated daily from remote sensing products at a spatial
resolution of 9km. en. type: dataset. 2017. doi: 10.1594/PANGAEA.883613

30. Sunagawa, S., Acinas, S. G., Bork, P., Bowler, C., Acinas, S. G., Babin, M., … de
Vargas, C. (2020). Tara Oceans: towards global ocean ecosystems biology. *Nature
Reviews Microbiology*, *18*(8), 428–445. doi: 10.1038/s41579-020-0364-5

31. Guidi L, Morin P, Coppola L, Tremblay JE, Pesant S, Tara Oceans Consortium C, et al.
Environmental context of all samples from the Tara Oceans Expedition (2009-2013),
about nutrients in the targeted environmental feature. en. type: dataset. 2017. doi:
10.1594/PANGAEA.875575

32. Guidi L, Picheral M, Pesant S, Tara Oceans Consortium C, and Tara Oceans Expedition
P. Environ- mental context of all samples from the Tara Oceans Expedition (2009-2013),
about sensor data in the targeted environmental feature. en. type: dataset. 2017. doi:
10.1594/PANGAEA.875576

33. Guidi L, Ras J, Claustre H, Pesant S, Tara Oceans Consortium C, and Tara Oceans
Expedition P. Environmental context of all samples from the Tara Oceans Expedition
(2009-2013), about pigment concentrations (HPLC) in the targeted environmental
feature. en. type: dataset. 2017. doi: 10.1594/PANGAEA.875569

34. Pesant S, Tara Oceans Consortium C, and Tara Oceans Expedition P. Methodology
used on board to prepare samples from the Tara Oceans Expedition (2009-2013). en.
type: dataset. 2017. doi: 10.1594/ PANGAEA.875580

35. Speich S, Chaffron S, Ardyna M, Pesant S, Tara Oceans Consortium C, and Tara
Oceans Expedition P. Environmental context of all samples from the Tara Oceans
Expedition (2009-2013), about the water column features at the sampling location. en.
type: dataset. 2017. doi: 10.1594/PANGAEA.875579

36. Picheral M, Guidi L, Stemmann L, Karl DM, Iddaoud G, and Gorsky G. The Underwater
Vision Profiler 5: An advanced instrument for high spatial resolution studies of particle
size spectra and zooplankton. Limnology and Oceanography: Methods 2010; 8:462–73

37. Picheral M, Searson S, Taillandier V, Bricaud A, Boss E, Ras J, et al. Vertical profiles of
environmental parameters measured on discrete water samples collected with Niskin
bottles during the Tara Oceans expedition 2009-2013. en. type: dataset. 2014. doi:
10.1594/PANGAEA.836319

38. Schlitzer R. Interactive analysis and visualization of geoscience data with Ocean Data
View. en. Computers & Geosciences 2002 Dec; 28:1211–8. doi: 10.1016/S0098-
3004(02)00040-7

39. Pesant, S., Not, F., Picheral, M., Kandels-Lewis, S., Le Bescot, N., Gorsky, G., …
Searson, S. (2015). Open science resources for the discovery and analysis of Tara
Oceans data. *Scientific Data*, *2*(Lmd), 1–16. doi: 10.1038/sdata.2015.23

40. Alberti A, Poulain J, Engelen S, Labadie K, Romac S, Ferrera I, et al. Viral to metazoan
marine plankton nucleotide sequences from the Tara Oceans expedition. en. Scientific
Data 2017 Aug; 4:170093. doi: 10.1038/sdata.2017.93

41. Vargas C de, Audic S, Henry N, Decelle J, Mahe F, Logares R, et al. Eukaryotic plankton
diversity in the sunlit ocean. en. Science 2015 May; 348:1261605–1261605. doi:
10.1126/science.1261605

42. Jari Oksanen FGB, Friendly M, Kindt R, Legendre P, McGlinn D, Minchin PR, et al.
vegan: Community Ecology Package. R package version 2.5-6. 2019

43. Wilson JB. Methods for fitting dominance/diversity curves. en. Journal of Vegetation
Science 1991 Feb; 2:35–46. doi: 10.2307/3235896

44. Matthews TJ and Whittaker RJ. Fitting and comparing competing models of the species
abundance distribution: assessment and prospect. en. Frontiers of Biogeography 2014
Jun; 6. doi: 10.21425/ F56220607

45. Chave J. Neutral theory and community ecology: Neutral theory and community
ecology. en. Ecology Letters 2004 Feb; 7:241–53. doi: 10.1111/j.1461-
0248.2003.00566.x

46. McGill BJ, Maurer BA, and Weiser MD. EMPIRICAL EVALUATION OF NEUTRAL
THEORY. en. Ecology 2006 Jun; 87:1411–23. doi: 10.1890/0012-
9658(2006)87[1411:EEONT]2.0.CO;2

47. Paulo I. Prado MDM and Chalom A. sads:Maximum Likelihood Models for Species
Abundance Distributions. R package version 0.4.2. 2018

48. Magurran AE and McGill BJ. Biological Diversity frontiers in measurement and
assessment. en. 24. New York: Oxford University press, 2011

49. Burnham, K. P., & Anderson, D. R. (2004). Multimodel inference: Understanding AIC
and BIC in model selection. *Sociological Methods and Research*, *33*(2), 261–304. doi:
10.1177/0049124104268644

50. R Core Team. R: A Language and Environment for Statistical Computing. ISBN 3-
900051-07-0. R Foundation for Statistical Computing. Vienna, Austria, 2013

51. Dinno A. Dunn's Test of Multiple Comparisons Using Rank Sums. R package version
1.3.5. 2017

52. Tackmann J, Rodrigues JFM, and Mering C von. Rapid inference of direct interactions
in large-scale ecological networks from heterogeneous microbial sequencing data. Cell
Systems 2019. doi: 10.1016/ j.cels.2019.08.002

53. Clauset, Newman, and Moore, "Finding Community Structure in Very Large Networks."
Physical Review E. 2004; 70:066111 doi: 10.1103/PhysRevE.70.066111

54. Vellend M. Conceptual Synthesis in Community Ecology. en. The Quarterly Review of
Biology 2010 Jun; 85:183–206. doi: 10.1086/652373

55. Filho RC, Martins FR, and Gneri MA. Fitting abundance distribution models in tropical
arboreal communities of SE Brazil. en. Community Ecology 2002; 3:169–80

56. Magurran AE and Henderson PA. Explaining the excess of rare species in natural
species abundance distributions. en. 2003; 422:3

57. Sugihara G. Minimal Community Structure: An Explanation of Species Abundance
Patterns. en. The American Naturalist 1980; 116:770–87

58. Bulmer MG. On Fitting the Poisson Lognormal Distribution to Species-Abundance Data.
en. Biometrics 1974 Mar; 30:101. doi: 10.2307/2529621

59. Preston FW. The Commonness, And Rarity, of Species. en. Ecology 1948 Jul; 29:254–
83. doi: 10. 2307/1930989

60. Ser-Giacomi, E., Zinger, L., Malviya, S., De Vargas, C., Karsenti, E., Bowler, C., & De
Monte, S. (2018). Ubiquitous abundance distribution of non-dominant plankton across
the global ocean. *Nature Ecology and Evolution*, *2*(8), 1243–1249. doi: 10.1038/s41559-
018-0587-2

61. Brum JR, Ignacio-Espinoza JC, Roux S, Doulcier G, Acinas SG, Alberti A, et al. Patterns
and ecological drivers of ocean viral communities. en. Science 2015 May;
348:1261498–1261498. doi: 10 . 1126 / science.1261498

62. Ghiglione JF, Galand PE, Pommier T, Pedrós-Alió C, Maas EW, Bakker K, et al. Pole-
to-pole bio- geography of surface and deep marine bacterial communities. en.
Proceedings of the National Academy of Sciences 2012 Oct; 109:17633–8. doi:
10.1073/pnas.1208160109

63. Ulloa O, Canfield DE, DeLong EF, Letelier RM, and Stewart FJ. Microbial oceanography
of anoxic oxygen minimum zones. en. Proceedings of the National Academy of Sciences
2012 Oct; 109:15996– 6003. doi: 10.1073/pnas.1205009109

64. Aldunate M, De la Iglesia R, Bertagnolli AD, and Ulloa O. Oxygen modulates bacterial
community composition in the coastal upwelling waters off central Chile. en. Deep Sea
Research Part II: Topical Studies in Oceanography 2018 Oct; 156:68–79. doi:
10.1016/j.dsr2.2018.02.001

65. De la Iglesia R, Echenique-Subiabre I, Rodríguez-Marconi S, Espinoza JP, Dassow P
von, Ulloa O, et al. Distinct oxygen environments shape picoeukaryote assemblages
thriving oxygen minimum zone waters off central Chile. en. Journal of Plankton
Research 2020 Sep; 42. Ed. by Dolan J:514–29. doi: 10.1093/plankt/fbaa036

66. Schnetzer A, Moorthi SD, Countway PD, Gast RJ, Gilg IC, and Caron DA. Depth
matters: Microbial eukaryote diversity and community structure in the eastern North
Pacific revealed through environmen- tal gene libraries. en. Deep Sea Research Part I:
Oceanographic Research Papers 2011 Jan; 58:16–26. doi: 10.1016/j.dsr.2010.10.003

67. Guidi L, Chaffron S, Bittner L, Eveillard D, Larhlimi A, Roux S, et al. Plankton networks
driving carbon export in the oligotrophic ocean. en. Nature 2016 Feb; 532:465–70. doi:
10.1038/nature16942

68. Parada V, Sintes E, Aken HM van, Weinbauer MG, and Herndl GJ. Viral Abundance,
Decay, and Diversity in the Meso- and Bathypelagic Waters of the North Atlantic. en.
Applied and Environmental Microbiology 2007 Jul; 73:4429–38. doi:
10.1128/AEM.00029-07

69. Bates AE, Helmuth B, Burrows MT, Duncan MI, Garrabou J, Guy-Haim T, et al.
Biologists ignore ocean weather at their peril. en. Nature 2018 Aug; 560:299–301

70. Mojica KDA and Brussaard CPD. Factors affecting virus dynamics and microbial host–
virus interactions in marine environments. en. FEMS Microbiol Ecol 2014 :21

71. Kukkaro P and Bamford DH. Virus–host interactions in environments with a wide range
of ionic trengths. en. Environmental Microbiology Reports 2009 :7

72. Bettarel Y, Bouvier T, and Bouvy M. Viral persistence in water as evaluated from a
tropical/temperate cross-incubation. en. Journal of Plankton Research 2009 Aug;
31:909–16. doi: 10.1093 / plankt / fbp041

73. Schulz F. Giant virus diversity and host interactions through global metagenomics. en. Nature 2020; 578:432–6. doi: https://doi.org/10.1038/s41586-020-1957-x

74. Thamdrup B, Dalsgaard T, and Revsbech NP. Widespread functional anoxia in the oxygen minimum zone of the Eastern South Pacific. en. Deep Sea Research Part I: Oceanographic Research Papers 2012 Jul; 65:36–45. doi: 10.1016/j.dsr.2012.03.001

75. Helly JJ and Levin LA. Global distribution of naturally occurring marine hypoxia on continental margins. en. Deep Sea Research Part I: Oceanographic Research Papers 2004 Sep; 51:1159–68. doi: 10.1016/j.dsr.2004.03.009

76. Berry D and Widder S. Deciphering microbial interactions and detecting keystone species with co- occurrence networks. en. Frontiers in Microbiology 2014 May; 5. doi: 10.3389/fmicb.2014.00219

77. Zhu W, Qin C, Ma H, Xi S, Zuo T, Pan W, et al. Response of protist community dynamics and co- occurrence patterns to the construction of artificial reefs: A case study in Daya Bay, China. en. Science of The Total Environment 2020 Nov; 742:140575. doi: 10.1016/j.scitotenv.2020.140575

78. Chaffron, S., Delage, E., Budinich, M., Vintache, D., Henry, N., Nef, C., … Eveillard, D. (2020). Environmental vulnerability of the global ocean plankton community interactome. *BioRxiv*. doi: 10.1101/2020.11.09.375295

79. Mihara T, Koyano H, Hingamp P, Grimsley N, Goto S, and Ogata H. Taxon Richness of "Megaviridae" Exceeds those of Bacteria and Archaea in the Ocean. en. Microbes and Environments 2018; 33:162–71. doi: 10.1264/jsme2.ME17203

80. Glock N, Roy AS, Romero D, Wein T, Weissenbach J, Revsbech NP, et al. Metabolic preference of nitrate over oxygen as an electron acceptor in foraminifera from the Peruvian oxygen minimum zone. en. Proceedings of the National Academy of Sciences 2019 Feb; 116:2860–5. doi: 10.1073/pnas. 1813887116

Figure legends

Figure 1: Geographical locations of *Tara* Oceans epipelagic and mesopelagic sampling sites included in this study. Symbols: ▲ refers to epipelagic, ▼ Oxic MES and ♦ OMZ eco-regions. Symbol colors represent organism groups evaluated in the present study.

Figure 2: Ordination plot of epipelagic (left) and mesopelagic (right) communities based on OTU composition based on canonical correspondence analysis (CCA). Percentages in parentheses are the amount of variation constrained, in titles represent the total in each analysis, and in the axis the correspondent value for each dimension. Arrows represent environmental quantitative explanatory variables with arrowheads indicating their direction of increase. Shapes represent sampling sites. Shape formats represent eco-regions, epi: epipelagic, Oxic MES: oxic mesopelagic, OMZ: oxygen minimum zone mesopelagic. IO: Indian Ocean, NAO: North Atlantic Ocean, NPO: North Pacific Ocean, SAO: South Atlantic Ocean, SPO: South Pacific Ocean.

Figure 3: Temperature and salinity plot indicating water mass designation for all mesopelagic samples. Formats represent the different oceanic basins (■ - North Atlantic Ocean, ● - South Atlantic Ocean, ▲ - Pacific Ocean, ★ - Indian Ocean). Colors indicate the oxygen concentration at the sampling depth. LSW - Labrador Sea Water; AAIW - Antarctic Intermediate Water; tNPIW transitional North Pacific Intermediate Water; SAMW - Subantarctic Mode Water; SPSTMW - South Pacific Subtropical Mode Water; modAAIW - modified Antarctic Intermediate Water; PGW - Persian Gulf Water mass; RSW - Red Sea Water mass; NASTMW - North Atlantic Subtropical Mode Water.

Figure 4: Relative abundance of OTUs assigned to mesopelagic eco-regions. A) Phages, B)
Giruses, C) Prokaryotes and D) pico-Eukaryotes.

Figure 5: Co-occurrence network in epipelagic and mesopelagic communities. A) Global
network, with connected modules for OMZ (purple and orange) and MES (green) highlighted.
B) Relative taxa abundance in each module in each station and depth. C) Relative number of
OTUs classified in taxonomic groups. D) Network representation of modules.

Supplementary Figure 1: Non-metric multidimensional scaling (NMDS) showing epipelagic
and mesopelagic community stratification for each organism group.

Supplementary Figure 2: Epipelagic and mesopelagic group dispersion based on physical-
chemical oceanic properties (Euclidian method). A) First two axes of PCoA. B) Dispersion of
distances from samples to centroids.

Supplementary Figure 3: Relative abundance of OTUs classified into different eco-regions
by ocean layers.

Supplementary Figure 4: Normalized relative abundance of phages and their preferred eco-
region.

Supplementary Figure 5: Normalized relative abundance of giruses and their preferred eco-
region.

Supplementary Figure 6: Normalized relative abundance of prokaryotes and their preferred
eco-region.

Supplementary Figure 7: Normalized relative abundance of pico-eukaryotes and their
preferred eco-region.

Supplementary Figure 8: Relative abundance of OTUs from taxonomic groups for epipelagic
and mesopelagic (Oxic MES and OMZ) samples enriched in each eco-region (UBI: ubiquitous,
EPI: epipelagic, Core MES: core mesopelagic, Oxic MES: oxic mesopelagic, OMZ: oxygen
minimum zone. A) Phages B) Giruses C) Prokaryotes D) pico-eukaryotes.

 # Figures:

Figure 1: Geographical locations of Tara Oceans epipelagic and mesopelagic sampling sites included in this study. Symbols up-triangle refers to epipelagic, down-triangles Oxic MES and diamond OMZ eco-regions. Symbols colors represent organism groups evaluated in the present study

Figure 2: Ordination plot of epipelagic (left) and mesopelagic (right) communities based on OTU's composition based on canonical correspondence analysis (CCA). Percentages in parenthesis are the amount of variation constrained, in titles represents the total in each analysis, and in the axis the correspondent value for each dimension. Arrows represent environmental quantitative explanatory variables with arrowheads indicating their direction of increase. Shapes represent sampling sites. Shape formats represent eco-regions, epi: epipelagic, Oxic_mes: oxic mesopelagic, OMZ: OMZ mesopelagic. IO: Indian Ocean, NAO: North Atlantic Ocean, NPO: North Pacific Ocean, SAO: South Atlantic Ocean, SPO: South Pacific Ocean

Figure 3: Temperature and salinity plot indicating water masses designation for all mesopelagic samples. Formats represent the different oceanic basins (□- North Atlantic Ocean, ◯ - South Atlantic Ocean, △ - Pacific Ocean, ★ - Indian Ocean). Colours indicate the oxygen concentration at the sampling depth. LSW - Labrador Sea Water; AAIW - Antarctic Intermediate Water; tNPIW ? transitional North Pacific Intermediate Water; SAMW - Subantarctic Mode Water; SPSTMW - South Pacific Subtropical Mode Water; modAAIW - modified Antarctic Intermediate Water; PGW - Persian Gulf Water mass; RSW - Red Sea Water mass; NASTMW - North Atlantic Subtropical Mode Water.

Figure 4: Relative abundance of OTUs assigned to Mesopelagic Eco-regions. A) Phages, B) Giruses, C) Prokaryotes and D) Eukaryotes

Figure 5: Co-Ocurrence Network in Epipelagic and Mesopelagic communities. A) Total Network, with connected modules for OMZ (purple and orange) and MES (green) highlighted. B) Relative taxa abundance in each Module in each station and depth. C) Relative number of OTUs classified in taxonomic groups. D) Network representation of modules

 # Tables

Table 1: ANOVA p_value for each environmental factor used as explanatory variable. Global correspond to the p-value for the whole set of variables, the following columns to p-values for each of the environmental variables considering the others as covariable.

assemblage	Depth	Global	Temp. °C	Salinity	O_2 [μmol/kg]	[NO3]$^-$ [μmol/l]	Chl-a [mg/m^3]	Particle flux
Phages	EPI	0.001	0.057	0.144	0.043	0.031	0.420	0.017
Pahges	MES	0.001	0.059	0.004	0.027	0.124	0.110	0.002
Giruses	EPI	0.001	0.023	0.074	0.051	0.152	0.053	0.221
Giruses	MES	0.002	0.032	0.211	0.145	0.134	0.008	0.211
Prokaryotes	EPI	0.035	0.048	0.141	0.002	0.006	0.044	0.568
Prokaryotes	MES	0.006	0.501	0.304	0.039	0.444	0.966	0.486
pico-Eukaryotes	EPI	0.025	0.027	0.166	0.412	0.292	0.216	0.659
pico-Eukaryotes	MES	0.001	0.606	0.191	0.243	0.468	0.271	0.477

Table 2: Proportion of the variation in community composition that explained by water masses using the Permutation multivariated analysis of variance (PERMANOVA)

assemblage	Df	Sum Of Squares	Mean Squares	F. Model	R^2	Pr(¿F)
Phages	4	1.87	0.47	2.32	0.51	0.002
Giruses	8	3.19	0.40	1.81	0.46	0.001
Prokaryotes	8	1.30	0.16	3.29	0.60	0.001
pico-Eukaryotes	8	2.60	0.32	1.62	0.36	0.001

Table 3: Network topological features derived from global analysis including giruses, prokaryotes and pico-eukaryotes samples in epipelagic and mesopelagic depths

Name	Global	Mod 1	Mod 4	Mod 17
Nodes	6154	731	323	175
Positive Edges	12193	1236	480	223
Negative Edges	742	70	49	9
Avg. degree	4.20	3.57	3.28	2.65
Clustering	0.03	0.03	0.09	0.05
Density	0.00	0.00	0.01	0.02
Average.path.length	7.28	6.01	6.27	6.30
Betweenness	0.01	0.05	0.10	0.22
Degree Centralization	0.00	0.01	0.02	0.04
Modularity	0.47	0.60	0.67	0.66

 # Supplementary Figures:

Supplementary Figure 1: Non-metric multidimensional scaling (NMDS) showing epipelagic and mesopelagic communities stratification for each organism group

Supplementary Figure 2: Epipelagic and mesopelagic group dispersion based on physical-chemical oceanic properties (Euclidian method). A) First two axes of PCoA. B) Dispersion of distances from samples to centroids.

Supplementary Figure 3: Relative abundance of OTUs classified into different eco-regions in to ocean layers

Supplementary Figure 4: Normalized Relative abundance of Phages and their preferred eco-region

Supplementary Figure 5: Normalized Relative abundance of giruses and their preferred eco-region

Supplementary Figure 6: Normalized Relative abundance of Prokaryotes and their preferred eco-region

Supplementary Figure 7: Normalized Relative abundance of pico-Eukaryotes and their preferred eco-region

Supplementary Figure 8: Relative abundances of OTUs from taxonomic groups for epipelagic and mesopelagic (Oxic MES and OMZ) samples enriched in each eco-region (UBI: ubiquitous, EPI: epipelagic, Core MES: core mesopelagic, Oxic MES: oxic mesopelagic, OMZ: oxygen minimum zone. A) Phages B) Giruses C) Prokaryotes D) pico-Eukaryotes