Improved correspondence of fMRI visual field localizer data after

cortex-based macroanatomical alignment

Running Title: Visual field localizer & CBA

Mishal Qubad[1], Catherine V. Barnes-Scheufler[1], Michael Schaum[2], Eva Raspor[1],

Lara Rösler[1,3], Benjamin Peters[4,5], Carmen Schiweck[1], Rainer Goebel[3,6], Andreas

Reif[1], Robert A. Bittner[1,7]*

[1] Department of Psychiatry, Psychosomatic Medicine and Psychotherapy and Brain

Imaging Center, University Hospital, Goethe University, Frankfurt, Germany

[2] Leibniz Institute for Resilience Research, Mainz, Germany

[3] Netherlands Institute for Neuroscience, Amsterdam, The Netherlands

[4] Institute of Medical Psychology, University Hospital, Goethe University, Frankfurt,

Germany

[5] Zuckerman Mind Brain Behavior Institute, Columbia University, New York, NY, USA

[6] Department of Cognitive Neuroscience, Faculty of Psychology and Neuroscience,

Maastricht University, Maastricht, The Netherlands

[7] Ernst Strüngmann Institute for Neuroscience (ESI) in Cooperation with Max Planck

Society, Frankfurt am Main, Germany

*Corresponding author. Department of Psychiatry, Psychosomatic Medicine and Psy-

chotherapy, University Hospital Frankfurt, Goethe University, Heinrich-Hoffmann-Str.

10, D-60528 Frankfurt am Main, Germany.

Phone: +49 69 6301 84713.

Fax: +49 69 6301 81775.

Email: bittner@med.uni-frankfurt.de

Total number of words in abstract: 196/200

Total number of words in main text: 4279/4500

ORCID IDS

M.Q.: orcid.org/0000-0002-9291-5739

C.V.B.-S.: orcid.org/0000-0002-4730-1981

M.S.: orcid.org/0000-0002-6589-4530

E.R.: orcid.org/0000-0003-4418-7084

L.R.: orcid.org/0000-0003-1046-1073

B.P.: orcid.org/0000-0002-0948-8976

C.S.: orcid.org/0000-0003-4215-0448

R.G.: orcid.org/0000-0003-1780-2467

A.R.: orcid.org/0000-0002-0992-634X

R.A.B.: orcid.org/0000-0003-2021-0358

ABSTRACT

Studying the visual system with fMRI often requires using localizer paradigms to de-

fine regions of interest (ROIs). However, the considerable interindividual variability of

the cerebral cortex represents a crucial confound for group-level analyses. Cortex-

based alignment (CBA) techniques reliably reduce interindividual macroanatomical

variability. Yet, their utility has not been assessed for visual field localizer paradigms,

which map specific parts of the visual field within retinotopically organized visual ar-

eas. We evaluated CBA for an attention-enhanced visual field localizer mapping ho-

mologous parts of each visual quadrant in 50 participants. We compared CBA with

volume-based alignment and a surface-based analysis, which did not include macro-

anatomical alignment. CBA led to the strongest increase in the probability of activa-

tion overlap (up to 40 percent). On the group level, CBA led to the most consistent in-

crease in ROI size while preserving vertical ROI symmetry. Overall, our results indi-

cate, that in addition to the increased signal-to-noise ratio of a surface-based analysis

macroanatomical alignment considerably improves statistical power. These findings

confirm and extend the utility of CBA for the study of the visual system in the context

of group analyses. CBA should be particularly relevant when studying neuropsychiat-

ric disorders with abnormally increased interindividual macroanatomical variability.

KEYWORDS

INTRODUCTION

The visual system includes a multitude of topographical representations of varying

resolution across increasingly specialized visual areas[1]. Functional magnetic reso-

nance imaging (fMRI) offers a variety of methods either to map these topographical

representations in full, or to localize specific visual areas or retinotopic positions

within their topography. These approaches are essential not only for the fine-grained

study of fundamental properties of the visual system[1], but also for investigating the

role of these areas for higher-order cognitive processes such as visual attention and

working memory[2-6]. This also extends to translational studies of visual dysfunction

and its cognitive consequences in neuropsychiatric disorders[7,8].

Methods for fMRI-based visual mapping, i.e., techniques to define regions of interest

in the visual system based on specific functional properties, fall in in three broad cat-

egories: retinotopic mapping, visual field localizer and functional localizer paradigms.

Retinotopic mapping and the more advanced population receptive field mapping al-

low the complete delineation of early visual areas[1,9,10]. Conversely, visual field local-

izer paradigms can map a circumscribed region within a retinotopically organized vis-

ual area[11,12]. Finally, functional localizers can detect higher-order visual areas such

as the fusiform face area, parahippocampal place area (PPA), extrastriate body area

(EBA) and lateral occipital complex (LOC), which are clustered and show specializa-

tion for the processing of specific categories of complex visual information[1,13,14]. In

most fMRI studies, high interindividual anatomical variability of cortical areas in terms

of both size and location constitutes an important challenge[15-23]. For instance, it has

been shown that primary visual cortex (V1) can differ in size by about 2-fold between

individuals[17]. Furthermore, anatomical variability in terms of location has been shown

to be particularly pronounced in extra striate visual areas[24]. This crucial confound re-

duces the power to reliably map visual areas at the group level.

One way to mitigate this problem for the visual system is to pool single-subject re-

gions of interest (ROIs), while simultaneously using the overall group-based probabil-

ity for that ROI at each point in a Cartesian coordinate system as a constraint[25-27].

While such a single-subject-based analysis improves sensitivity and functional reso-

lution compared to a standard group-based approach, it does not actually reduce

macroanatomical variability. Additionally, studying the interplay between visual areas

and other cortical areas more directly involved in higher-order cognitive processes

with whole-brain methods such as functional connectomics network analyses[28] might

preclude a single-subject based strategy.

Such group-based analyses typically require spatial normalization of structural and

functional imaging data to a common Cartesian coordinate system such as Talair-

ach[29] or MNI[30] space. In its most basic form, volume-based spatial normalization em-

ploys a linear transformation that matches the overall extent of the brains to a stand-

ard brain template. While transformation into Talairach space relies on anatomical

landmarks, transformation into MNI space utilizes fully data driven registration of

structural images to an average template brain[30]. While these spatial normalization

approaches inherently lead to an alignment of brains, the underlying algorithms are

not geared specifically toward aligning homologous brain structures. Conversely,

more refined methods employ non-linear warping algorithms guided by intensity dif-

ferences to improve macroanatomical alignment[31]. Thus, all of these methods can be

categorized as volume-based alignment (VBA) techniques. However, both linear and

nonlinear VBA mostly disregard the topological properties of the cerebral cortex and

its geometric features such as sulci and gyri. Consequently, VBA methods result in a

considerable amount of residual interindividual anatomical variability[32,33].

Surface-based procedures constitute an important alternative approach. Surface-

based spatial normalization typically uses a geodesic coordinate system, which al-

lows for a two-dimensional representation of the cerebral cortex and respects the cor-

tical topography to a much larger degree than traditional Cartesian coordinate sys-

tems[18,34]. This approach offers two main advantages over VBA. First, surface-based

spatial normalization allows to constrain data readout and data preprocessing such

as spatial smoothing to cortical tissue, greatly reducing signal contamination by white

matter and cerebrospinal fluid. It also avoids contamination from cortical areas proxi-

mal in volume space but considerably more distant in surface space. Consequently,

spatial smoothing in surface space is superior to spatial smoothing in volume

space[19,35,36]. The second advantage of surface-based spatial normalization is the

possibility to use individual cortical folding patterns for an additional, fully data-driven

macroanatomical alignment of the cerebral cortex[34]. Compared to VBA techniques,

these cortex-based alignment (CBA) methods considerably improve anatomical cor-

respondence of cortical structures while respecting cytoarchitectonic boundaries[37].

Thus, CBA leads to a notable reduction of inter-individual anatomical variability[18,34,38-

40].

Importantly, previous studies have often exclusively compared surface-based data

before and after macroanatomical alignment[19,41], essentially using the former ap-

proach as a proxy for VBA. Yet, this comparison only reflects the second advantage

of CBA, namely the use of macroanatomical alignment instead of VBA. However, in

this case both data sets benefit equally from reduced signal contamination, likely un-

derestimating the full effects of CBA. Assessing the impact of this first advantage of

surface-based analyses in isolation requires a comparison of VBA with a surface-

based analysis without macroanatomical alignment. We refer to this intermediate ap-

proach as a "surface-based analysis using VBA" (SBAV). Thus, studying both ad-

vantages of CBA requires the comparison of three approaches: VBA, SBAV and

CBA.

Due to the advantageous properties outlined above, CBA methods have been pro-

posed as an alternative approach to VBA specifically for the visual system[26]. Several

studies have compared the impact of VBA and CBA methods on specific visual map-

ping techniques. For retinotopic mapping an improvement of functional overlap in

both V1 and V2 after CBA has been demonstrated[34,42]. Furthermore, for functional lo-

calizer data CBA substantially increases the overlap of object processing areas LOC,

FFA and PPA across subjects[19,43-45]. Conversely, the effects of CBA on visual field

localizer paradigms mapping specific retinotopic positions have not been studied.

Thus, the utility of CBA has been demonstrated for two of the three main categories

of visual mapping methods, i.e., those methods, which map whole areas, either de-

fined by cytoarchitectonic (e.g. V1) or functional (e.g. FFA) properties. Conversely, it

remains unclear, to which degree CBA can improve the alignment of ROIs mapped

by visual field localizer paradigms. Such paradigms are required for the detailed

study of the local processing of simple visual stimuli in early visual areas[11,12,46-48].

Flashing checkerboards covering the exact area of interest within the visual field are

primarily used for this purpose. Checkerboards lead to a particularly strong BOLD-

signal increase in early visual areas (V1-V3)[49]. To maximize the fidelity of the result-

ing localizer maps, visual field localizer paradigms typically utilize the fact that atten-

tional modulation induced by task demands significantly enhances response reliability

across visual areas. This can be accomplished by adding a simple target detection

task[50].

We used such an attention-enhanced visual field localizer paradigm to map a circum-

scribed location in each visual quadrant across early visual areas aiming to define

ROIs to be used for the study of higher cognitive processes. We chose a CBA

method using a dynamic group average as the target brain[19], thus eliminating the

possible confound of a static CBA target based on an individual brain, whose cortical

folding pattern might by chance deviate considerably from the group average.

Our primary goal was to examine the effects of CBA for a visual field localizer para-

digm. More specifically, we aimed to determine, whether macroanatomical alignment

improves the reliability of mapping subregions within retinotopically organized visual

areas delineated by such a paradigm at the group level. To this end, in addition to the

analysis of the full single-subject ROIs, we also examined the correspondence of sin-

gle-subject ROI peak vertices, i.e., single vertices showing the strongest level of acti-

vation in each subject for each visual quadrant. We conducted this analysis, because

peak vertices are a good approximation of the center of a ROI and thus allow for a

more precise assessment and visualization of the effects of CBA. Based on previous

findings for other localizer paradigm classes and the relatively good structural-func-

tional correspondence in posterior occipital cortex, we expected to observe a benefit

of CBA compared to SBAV when aligning subregions within early visual cortex for

both full ROIs and peak vertices. Our second goal was to examine the effects of

SBAV. More specifically, we aimed to assess the impact of surface-based functional

data readout and pre-processing without macroanatomical alignment. Here, we ex-

pected a general improvement of the signal-to-noise ratio for SBAV compared to VBA

and a corresponding global increase in ROI size for all visual quadrants.

Notably, several studies have shown differential response properties such as recep-

tive field size by visual quadrant or hemifield for homologous early visual areas. For

instance, improved behavioral performance and higher BOLD-signal amplitudes were

observed in the lower visual hemifield[51-54]. We were therefore also interested,

whether we could observe differences between upper and lower visual hemifields in

our group analysis after CBA. Overall, the aim of the study was to close an important

gap in the evaluation of CBA for the study of the visual system. Since visual field lo-

calizers are crucial for investigating contributions of the visual system to higher-order

cognitive processes, our results should have implications for the study of visual cog-

nition in both basic and translational research.

RESULTS

Visual quadrant ROIs (group level)

Group-level mapping of the four visual quadrants revealed notable differences for the

three alignment techniques (VBA, SBAV, CBA) (Figure 4, Table 1). For the lower

right visual quadrant, ROI size increased considerably from VBA to SBAV, but de-

creased for CBA. For the lower left visual quadrant, ROI size decreased slightly from

VBA to SBAV, but increased considerably for CBA. For the upper left visual quadrant,

ROI size increased considerably from VBA to SBAV and increased further for CBA.

For the upper right visual quadrant ROI size increased slightly from VBA to SBAV

and increased considerably for CBA. Thus, three out of four visual quadrant ROIs ex-

hibited a pattern of continuously increasing cluster size reflecting an increasing extent

of significant position selectivity across alignment techniques. Conversely, the ROI of

the lower right visual quadrant showed a cluster size decrease after CBA. For SBAV

this ROI also showed by far the greatest extent of any ROI even encompassing pos-

terior parts of temporal cortex.

Within group ROIs, average time courses showed clear position selectivity, which

was not affected by alignment technique as indicated by the negative results of our

linear mixed models (Table 2). Notably, asymmetry indices (AIs) revealed markedly

greater vertical symmetry of both upper and lower hemifield ROIs for VBA and CBA

compared to SBAV (Table 3). After CBA, ROI sizes for the lower visual hemifield

were considerably larger size than for the upper visual hemifield (Table 1).

Probability maps

For all three data sets, the maximum probability of activation overlap was consistently

located at the center of each ROI as defined in our previous group analysis (Figure

4). For VBA data, probability maps (PMs) showed a relatively wide spread of func-

tional activation around the core ROIs (Figure 5a, Table 4). Maximum probability of

activation overlap was approximately 50 percent. For SBAV data, PMs showed an

even wider spread of functional activation around the core ROIs (Figure 5b, Table 4).

Maximum probability of activation overlap was approximately 45 percent. For CBA

data, PMs showed a noticeable decrease in the spread of functional activation

around the core ROIs with a corresponding increase in the maximum probability of

overlap at the center of the core ROIs (Figure 5c, Table 4). Maximum probability of

activation overlap was approximately 75 percent.

Probability difference maps

Probability difference maps (PDMs) revealed a differential impact of the individual

methodological elements of our overall macroanatomical alignment approach.

For the impact of surface-based functional data readout and pre-processing com-

pared to standard volume-based alignment, the corresponding PDM (SBAV minus

VBA) showed an increase in the probability of activation overlap of up to 15 percent

around the central ROIs. Conversely, at the location of the central ROIs we mostly

observed a decrease in the probability of activation overlap of up to 10 percent (Fig-

ure 6a, Table 5). Notably, changes were widespread, partly extending into posterior

temporal and parietal cortex. For the additional impact of macroanatomical alignment,

the corresponding PDM (CBA minus SBAV) showed an increase in the probability of

activation overlap of up to 35 percent in the central ROIs (Figure 6b, Table 5). Con-

versely, more peripheral occipital regions showed a decrease in the probability of ac-

tivation overlap of up to 35 percent. Overall, changes were considerably less wide-

spread than for the SBAV minus VBA comparison. For the additive impact of both

methodological elements, the corresponding PDM (CBA minus VBA) showed an in-

crease in the probability of activation overlap of up to 30 percent in the central ROIs

(Figure 6c, Table 5). Conversely, more peripheral occipital regions as well as poste-

rior temporal and parietal cortex showed a decrease in the probability of activation

overlap of up to 20 percent. Overall, the spatial extent of these effects fell in between

that of the other two comparisons.

Spatial variability of ROI peak vertex distribution (single-subject level)

The rate of success for detecting a ROI for each subject was as follows: lower left

visual quadrant 96 percent (48 out of 50 subjects), lower right visual quadrant 100

percent (50 out of 50 subjects), upper left visual quadrant 100 percent (50 out of 50

subjects), upper right visual quadrant 92 percent (46 out of 50 subjects). As with the

PMs at the group level, peak vertex distribution maps at the single-subject level (Fig-

ure 7) showed less spatial variability for CBA compared to SBAV. Furthermore, for

CBA compared to SBAV we observed an increase in the number of multiple overlap-

ping single-subject ROI peak vertices per vertex for each visual quadrant and a cor-

responding decrease in the number of single overlapping single-subject ROI peak

vertices per vertex (Table 6).

DISCUSSION

The aim of our study was to evaluate the utility of CBA for an attention-enhanced vis-

ual field localizer paradigm used to map a circumscribed region within retinotopically

organized visual areas. Our paradigm mapped homologous regions in each visual

quadrant reliably across early visual areas. As expected, CBA led to a marked reduc-

tion in macroanatomical variability with a number of beneficial effects on the func-

tional level, which clearly exceeded those observed for SBAV.

Compared to VBA and SBAV, CBA resulted in the largest improvement in the group

ROI analysis across visual quadrants (Figure 4). This was reflected in the consistent

size increase of the resulting group ROIs compared to VBA and SBAV (Table 1),

which indicates an increased power to detect subregions of early visual areas show-

ing position selectivity. Unsurprisingly, we did not observe an increase of position se-

lectivity when comparing visual quadrant ROIs across alignment techniques (Table

2). Importantly, only CBA but not SBAV preserved the vertical symmetry of group

ROIs characteristic for early visual areas, which was already evident for VBA (Table

3).

Regarding changes in the probability of activation overlap across the three alignment

techniques as reflected in the PMs, a clear pattern emerged. The probability of acti-

vation overlap increased gradually with each step, peaking for CBA with a value of

about 75 percent. For SBAV, effects were weaker and considerably more wide-

spread, favoring more peripheral brain regions. Similarly, PDMs showed an increase

in the probability of activation overlap of up to 40 percent in the central region of in-

terest for CBA, resulting in considerably more focused activation patterns, while the

opposite effect emerged in more peripheral vertices (Figure 6). This is most likely not

attributable to a decreased spatial overlap in the periphery of early visual areas. Ra-

ther, it demonstrates that CBA consistently reduces spurious spread-out activation

resulting from poor macroanatomical correspondence after VBA and a generalized

increase in the SNR due to SBAV. It also suggests that VBA and SBAV might mis-

represent the location and extent of early visual areas. This is notion is supported by

changes of the center gravity of group ROIs between SBAV and CBA, which was

particularly pronounced in the lower left visual quadrant (Table 1). Together, these

findings indicate that CBA substantially increases statistical power when studying

early visual areas at the group level. Naturally, this effect of CBA should also extend

to studies with a more global focus, such as connectivity analyses[35,55].

The specific advantages of CBA compared to SBAV were also evident in the mark-

edly decreased variability of single-subject ROI peak vertex locations for each visual

quadrant (Figure 7, Table 6). This indicates that the reduction of macroanatomical

and functional inter-subject variability achieved by CBA is the main reason for our im-

proved group-level results. Our findings confirm the notion that transforming func-

tional data from volume-space into surface space already increases statistical power

by reducing signal contamination from non-neuronal tissue, thus improving the sig-

nal-to-noise ratio. Therefore, using SBAV as a proxy for VBA would underestimate

the actual benefits of CBA, which clearly exceed those of SBAV. Overall, the com-

bined beneficial effects of an improved signal-to-noise ratio and reduced macroana-

tomical alignment inherent in the CBA approach and evident in our data support the

notion that CBA is clearly the superior alignment technique when studying the visual

system.

For VBA we observed the largest group ROIs for the right and left lower visual quad-

rant, an effect that changed after SBAV and CBA (Figure 4a, Table 1). For SBAV we

observed the largest group ROIs for the left upper and right lower visual quadrant,

which did not persist after CBA (Figure 4b & 4c, Table 1). Notably, several studies

using volume-based fMRI and magnetoencephalographic analyses reported lateral-

ized effects on neurophysiological parameters in early visual areas[56,57]. Our observa-

tion raises the question, whether these findings could at least partly be explained by

lateralized differences in macro-anatomical variability rather than true functional dif-

ferences.

Conversely, our CBA-aided group analysis allowed us to compare the response prop-

erties of each visual quadrant in a more unbiased way. We observed larger group

ROIs for the lower visual hemifield. In a CBA-based probabilistic atlas of the visual

system, which included all regions that could be defined in more than fifty percent of

subjects, probabilistic ROIs for dorsal V1 and V2 were also noticeably larger probabil-

istic ROIs for ventral V1 and V2, whereas this effect was less clear for V3[44]. These

results are in line with our own findings and could be attributable to higher residual

anatomical variability after CBA in ventral occipital cortex representing the upper vis-

ual hemifield. Alternatively, they could be due to true differences in response proper-

ties such as receptive field size or overall area size. The latter interpretation is sup-

ported by studies showing functional differences between upper and lower visual

hemifields already at the retinal level in the form of differences in receptor densities.

Cone density was higher in the superior parts of the retina, which processes infor-

mation from lower visual fields. Conversely, higher rod density was observed in the

inferior parts[58,59]. Moreover, Eickhoff et al. observed dorso-ventral asymmetries in re-

ceptor densities in V2 and V3[58] and reported a higher density of GABA-A, and mus-

carinic M3-receptors in ventral parts of V2 and V3. Furthermore, there is evidence for

fundamental differences in receptive field shape from a pRF mapping study. Estimat-

ing both the aspect ratios and the size of the mapped areas, a more elliptical recep-

tive field shape was observed for the upper visual hemifield represented by ventral

parts of the visual cortex being compared to the lower visual hemifield represented by

dorsal parts of the visual cortex[60]. Additionally, there is evidence for a behavioral ad-

vantage in the lower visual hemifield for shape discrimination as well as higher

BOLD-signal changes and peak amplitudes of MEG/EEG responses[51,53,54,61,62]. To-

gether, these findings demonstrate clear differences in the functional architecture of

early visual areas representing the upper and lower visual hemifield. This has been

attributed to the fact that the lower visual hemifield represented by dorsal parts of the

occipital lobe is more closely linked to the dorsal visual pathway, while the upper vis-

ual hemifield represented by ventral parts of the occipital lobe is more closely linked

to the ventral visual pathway[63,64]. Furthermore, there is evidence for fundamental dif-

ferences in receptive field shape from a pRF mapping study. For the upper visual

hemifield represented by ventral parts of the visual cortex an increased size and

more elliptical shape of receptive fields was observed compared to the lower visual

hemifield represented by dorsal parts of the visual cortex[60]. This implies, that the

lower visual field is more specialized for the precise localization and representation of

space. Our observation of larger ROIs in the lower visual hemifield is in line with

these findings. Thus, our results imply that CBA is a useful tool to extend the study of

functional and behavioral asymmetries in early visual areas to the group level.

One important limitation of the current study is the lack of a complementary retino-

topic mapping data set due to time constraints. This data would have allowed us to

delineate the boundaries of early visual areas and pinpoint the exact visual area con-

taining each individual single-subject ROI. Retinotopic mapping studies indicate that

peak activation of single subjects assessed by visual localizers are not consistently

located in the same visual area. Most localizer tasks show peak activation not in V1

but rather in V2 or V3[12]. It is therefore highly likely that our single-subject peak activa-

tion did not consistently belong to the same visual cortical area. With the current data

set we cannot determine how precisely individual visual areas were aligned with

CBA, and whether individual levels of the visual cortical hierarchy were differentially

affected. However, the position of our group ROIs, which bordered the calcarine sul-

cus and spanned the occipital pole, indicate that they mainly comprised V2 and V3.

Similarly, after CBA a comparable increase in the probability of overlap was observed

in the same part of occipital cortex. While this is at least suggestive of a relatively

consistent benefit of CBA across visual areas, more fine-grained studies including

retinotopic mapping are required to definitively address this question.

Furthermore, we did not use eye tracking to insure sufficient fixation. We also did not

include an additional central attentional control task on the fixation cross, which

would have further encouraged continuous fixation. This was done deliberately in or-

der to keep the difficulty level adequate for psychiatric patient populations. Our aver-

age success rate for finding reliable activation in early visual areas across all four vis-

ual quadrants was 97 (92 - 100) percent. Insufficient fixation might explain our failure

to find reliable activation in a small fraction of subjects in the upper right and lower

left visual quadrant.

Finally, several properties of the VBA data set differed from the SBAV and CBA data

sets. We could not match volume-based and surface-based preprocessing parame-

ters completely due to inherent differences between the three-dimensional and two-

dimensional spatial smoothing algorithms employed. The number of voxels and verti-

ces containing functional data was not identical, differentially affecting Bonferroni cor-

rection of group results. However, the smaller analysis space of the VBA data set, 69

percent the size of the SBAV and CBA data set, should only underestimate the effect

of the additional processing steps featured in the SBAV and CBA analysis.

Our study also has implications beyond mapping the visual system in healthy popula-

tions. Visual processing deficits are a prominent feature of neurodevelopmental psy-

chiatric disorders such as ADHD, schizophrenia and autism spectrum disorders[7,8,65-

72], which can also perturb crucial higher order cognitive processes including working

memory[73-75]. The current localizer paradigm will be useful to investigate local impair-

ments of visual information processing as well as disturbances in the interplay be-

tween early visual areas and brain networks supporting higher-order cognitive pro-

cesses. Here, CBA will be particularly relevant to reduce the confounding effects of

increased macroanatomical variability in disorders such as schizophrenia in order to

measure true group differences and true functional variability[38,76]. On the other hand,

CBA might also be crucial for investigating the neurodevelopmental underpinnings of

increased macroanatomical variability itself. To this end, the inclusion of probabilistic

atlases containing information about gene expression profiles[77] as well as cyto- and

receptor architectonics[78,79] will be valuable.

Our CBA approach relied solely on cortical curvature information to reduce macroan-

atomical variability. The advantage of this method is its feasibility for the vast majority

of fMRI data sets, since it only requires a structural brain scan of sufficient quality and

resolution. Among comparable methods, it is the most data driven and objective ap-

proach. However, the achievable reduction of macroanatomical variability is limited

by the variable and imperfect correlation between brain structure and brain func-

tion[34,40]. Therefore, more advanced methods additionally utilize orthogonal functional

data to further reduce anatomical variability, including the use of functional activation

or connectivity patterns to improve macroanatomical alignment across the whole

brain[20,80,81]. Additionally, a more complex approach has been proposed, which aligns

cortical data using 'areal features' more closely tied to cortical areas than cortical

folding patterns, including maps of relative myelin content and functional resting state

networks[82]. These methods have shown to provide a relevant additional reduction of

macroanatomical variability for a variety of paradigms including visual functional lo-

calizers. Future studies should also evaluate these methods for retinotopic mapping

and visual field localizers. Moreover, it has been demonstrated for early auditory ar-

eas, that the additional use of a probabilistic atlas of cytoarchitectonically defined ar-

eas further enhance standard CBA results[83]. In principle, such an approach would

easily be feasible for the visual system.

To summarize, we demonstrated the clear advantages of CBA compared to VBA for

the analysis of visual field localizer data at the group level, signified by a considerable

reduction of spatial variability across subjects of up to forty percent across early vis-

ual areas. Our findings extend previous CBA studies examining other major catego-

ries of visual mapping techniques. They underscore the loss of information and statis-

tical power incurred by the use of VBA methods in the majority of fMRI studies.

Therefore, CBA and comparable methods should be seriously considered as a stand-

ard procedure for the detailed study of visual information processing and its disturb-

ance in mental disorders.

METHODS AND MATERIALS

Participants.

All participants gave their written informed consent to participate in the study in ac-

cordance with the study protocol approved by the ethical review board of the Faculty

of Medicine at Goethe University. The experimental procedures were conducted in

conformity with the approved guidelines and the Declaration of Helsinki. Individuals

received compensation for their participation. We recruited 51 healthy volunteers (fe-

male : male = 28 : 23) with age ranging between 18-43 years (mean = 24). All partici-

pants were non-smokers, had no history of neurological or psychiatric illness and re-

ported normal or corrected-to-normal visual acuity. One participant was left-handed

as assessed by the German version of the Edinburgh Handedness Inventory[84].

Stimuli and task.

Subjects performed a visual field localizer paradigm (Figure 1a) implemented using

Presentation (Neurobehavioral Systems, Version 18.0) as part of a larger study in-

vestigating the role of visual areas for higher cognitive functions. The task consisted

of a series of flickering black-and-white-colored round shaped checkerboard stimuli

(flicker frequency = 7.5 Hz). Checkerboard stimuli appeared for 2000 ms randomly at

one of four different locations (non-target trial). Each location mapped a homologous

position in one of the four visual quadrants. The regular inter-trial interval (ITI) was 0

ms. However, once every 10 to 14 trials (11 times overall), the ITI increased to 2000

ms (prolonged ITI). Our paradigm featured a simple target-detection task. During 36

trials, the two centrally located squares of the checkerboard changed their color to

yellow for 133 ms (target trial). Participants were instructed to press a response box

button with their left thumb as quickly as possible if they detected a target. The task

consisted of a total of 144 trials (36 target trials, 108 standard trials). This target prob-

ability of 25 percent resulted in one target trial every fourth trial on average (range 3 to

5 trials) (Figure 1a). Throughout the task a black, x-shaped fixation cross was dis-

played at the center of the screen. Participants were instructed to continuously fixate

on the fixation cross. Before the first trial, only the fixation cross was displayed for 10

seconds. After the last trial, only the fixation cross was displayed for 20 seconds. The

total duration of the paradigm was 340 seconds (Figure 1b). For the purpose of our

analyses we defined a total of four conditions, one for each of the four stimulus loca-

tions. Each participant practiced the task prior to the measurement.

Acquisition and analysis of fMRI data.

We acquired functional MRI data on a Siemens 3T MAGNETOM Trio scanner at the

Goethe University Brain Imaging Centre using a gradient-echo 2D EPI sequence (32

axial slices, TR = 2000 ms, TE = 30 ms, FA = 90°, FoV = 192 x 192 mm2, voxel size

= 3 x 3 x 3 mm^3, gap = 1 mm, effective slice thickness = 4 mm). Slices were posi-

tioned parallel to the anterior- and posterior commissure. Functional images were ac-

quired in a single run comprising the acquisition of 170 volumes. Immediately before

each functional run, 6 volumes of this 2D EPI sequence were acquired with identical

parameters except for a switch of phase encoding direction (posterior to anterior in-

stead of anterior to posterior) for EPI distortion correction. Anatomical MRI data for

cortex reconstruction and co-registration with functional MRI data was acquired with

a high-resolution T1-weighted 3D volume using a Magnetization-Prepared Rapid

Gradient-Echo (MP-RAGE) sequence (192 sagittal slices, TR = 1900 ms, TE = 3.04

ms, TI 900 ms, FA = 9°, FoV = 256 x 256 mm^2, voxel size = 1 x 1 x 1 mm^3). Stimulus

presentation was constantly synchronized with the fMRI sequence. Head motion was

minimized with pillows. The task was projected by a beamer onto a mirror attached

on the head coil. MRI data were pre-processed and analyzed using BrainVoyager

20.6[85], the NeuroElf Matlab toolbox (www.neuroelf.net) and custom software written

in Matlab. One subject had to be excluded due to excessive intra-scan motion.

Structural image pre-processing.

Structural data pre-processing included background cleaning, brain extraction and

bias field correction to minimize image intensity inhomogeneities[85]. Bias field correc-

tion employed a "surface fitting" approach using singular value decomposition based

least squares low-order (Legendre) polynomials to model low-frequency variations

across 3D image space[86]. We used polynomials with an order of three, which were

fitted to a subset of voxels labeled as belonging to white matter. The estimated pa-

rameters of the polynomials were used to construct a bias field which was removed

from the data. Our approach comprised of one iteration using automatic white matter

labeling[87] and four iterations using manual white matter labeling.

Subsequently, structural data were transformed into Talairach coordinate space[29].

This comprised manual labeling of the anterior commissure (AC) and posterior com-

missure (PC) as well as the borders of the cerebrum. These landmarks were then

used to rotate each brain in the AC-PC plane followed by piece-wise, linear transfor-

mations to fit each brain in the common Talairach "proportional grid" system[19]. Trans-

formation into Talairach coordinate space was performed because the subsequent

automatic segmentation procedure exploits anatomical knowledge for initial brain

segmentation including removing subcortical structures and disconnection of cortical

hemispheres. To prepare the data for this procedure, we performed a manual filling

of the lateral ventricles. Based on the automatic segmentation of the structural scans

along the white-gray matter boundary[88], cortical hemispheres were reconstructed into

folded, topologically correct folded mesh representations, which were tessellated to

produce surface reconstructions and calculate curvature maps reflecting the individ-

ual cortical folding patterns. Surface reconstructions were subsequently morphed into

distortion corrected spherical representations. Finally, both folded and spherical

mesh representations were downsampled to a standard number of vertices (40962

vertices per hemisphere, mean vertex distance: 1.5 mm). We used these standard-

ized mesh representations for all surface-based processing steps.

Cortex-based alignment of structural data.

We then applied for each hemisphere separately a high-resolution, multiscale cortex-

based alignment procedure based on the individual curvature maps of all 50 partici-

pants. This CBA approach, which reliably aligns corresponding gyri and sulci across

subjects[85], consists of an initial rigid and a subsequent non-rigid alignment step[19]

(Figure 2a & 2b). During the initial step, cortical folding patterns of each sphere are

aligned rigidly to the cortical folding pattern of a single target sphere by global rota-

tion. Rigid CBA operates only on highly smoothed curvature maps containing only the

most prominent anatomical landmarks. We used the rotation parameters with the

highest degree of overlap between the curvature of each individual sphere and the

target sphere as the starting point for the subsequent non-rigid CBA.

Non-rigid CBA employs a coarse-to-fine matching strategy, which operates sequen-

tially at four levels of curvature smoothing starting with the detail level used during

rigid CBA. Each subsequent level includes increasingly finer anatomical details up to

almost the full curvature information. Importantly, non-rigid CBA aligns each cortical

folding pattern to a dynamically updated group average through iterative morphing.

This moving target approach, which generates the target curvature map from the av-

erage curvature across all hemispheres at a given alignment stage avoids the possi-

ble confounding effects of a suboptimal selection of an individual target brain, whose

folding pattern might deviate considerably from the cohort average.

Notably, rigid CBA typically utilizes a single brain randomly drawn from the cohort to

be aligned as its target brain. However, the folding pattern of this brain might also de-

viate considerably from the cohort average. To also address this potential confound,

we first conducted a preliminary CBA encompassing both rigid and non-rigid macro-

anatomical alignment (Figure 2a). We then conducted a second, final CBA. Here, we

used the aligned average brain derived from the preliminary CBA as an unbiased tar-

get for the rigid alignment step (Figure 2b). After the final non-rigid CBA, we merged

both hemispheres of each individual brain to create a global surface-based analysis

space.

Furthermore, for each hemisphere we created average surface representations from

the original, non-aligned folded mesh representations, which were subsequently

merged, inflated and used for data analysis and visualization. We repeated these

steps after applying the transformation matrix of the final rigid and non-rigid CBA to

the folded mesh representations, yielding an accurate representation of the structural

effects of macroanatomical alignment (Figure 2c).

Functional image pre-processing.

The first four volumes of each functional run were discarded to allow for T1 equilibra-

tion. Initial volume-based pre-processing of functional MRI data comprised slice tim-

ing correction using sinc interpolation and 3D motion correction using sinc interpola-

tion. Next, we performed echo-planar imaging distortion correction using the Correc-

tion based on Opposite Phase Encoding method[89,90]. EPI distortion corrected func-

tional data were co-registered to the untransformed extracted brains. This was ac-

complished utilizing a boundary-based registration algorithm optimized for surface-

based analyses[91]. After co-registration to the fully cleaned but untransformed struc-

tural data, functional data were transformed into Talairach coordinate space by apply-

ing the transformation matrix generated during Talairach transformation of the ana-

tomical data using sinc interpolation. This transformation preserved the original voxel

size of the functional data ($3 \times 3 \times 3$ mm^3) (Figure 3).

Surface-based pre-processing.

The volumetric functional data were then transformed into surface space by sampling

on the individual cortical surface reconstructions incorporating data from −1 to +3 mm

along vertex normals using trilinear interpolation. Subsequent pre-processing of func-

tional MRI data in surface space started with spatial smoothing using a nearest

neighbor interpolation (1 iteration). Based on the standardized vertex distance of 1.5

mm this approximates a 2D Gaussian smoothing kernel with a full width at half maxi-

mum (FWHM) of 3 mm. We opted for minimal spatial smoothing to prevent a loss of

accuracy for our visual field localizer. Spatial smoothing was followed by linear trend

removal and temporal high-pass filtering using fast Fourier transformation (high-pass

0.00903 Hz). Based on the vertex-to-vertex referencing from the folded, topologically

correct surface reconstructions to the spherical representations, we mapped the fully

preprocessed functional data into a common spherical coordinate system (Figure 3).

Finally, we applied surface-based anatomical masks that only included cortical verti-

ces in our analysis to the functional data. These masks excluded those subcortical

structures, i.e., parts of thalamus and the basal ganglia, which mapped onto the mid-

line of our surface reconstructions. For functional data analysis and subsequent Bon-

ferroni correction in surface space, this yielded a total number of 76132 vertices.

Full volume-based pre-processing.

To generate a purely volumetric data set for the comparison of VBA and SBAV, pre-

processing after EPI distortion correction was also conducted in volume space mirror-

ing as closely as possible the steps and parameters outlined above for surface-based

pre-processing. First, we applied spatial smoothing using a 3D Gaussian smoothing

kernel with a FWHM of 3 mm, which approximates the degree of surface-based spa-

tial smoothing. Second, we applied linear trend removal and temporal high-pass filter-

ing using fast Fourier transformation (high-pass 0.00903 Hz) using parameters ex-

actly matching surface-based pre-processing (Figure 3). This data set was not trans-

formed into surface space and did not include an anatomical mask. For functional

data analysis and subsequent Bonferroni correction in volume space, this yielded a

total number of 52504 voxels. Thus, the analysis space for VBA was 69 percent the

size of the analysis space for SBAV and CBA (52504 voxels vs. 76132 vertices).

Final functional data sets.

Overall, we generated three different functional data sets: a volume-based data set,

which was fully pre-processed and aligned in volume-space (VBA); a surface-based

data set, for which spatial smoothing and temporal filtering were only applied after

transformation in surface space, but without macroanatomical alignment (SBAV); a

surface-based data set, which was pre-processed in the same way as the SBAV data

set and also utilized macroanatomical alignment (CBA) (Figure 3). Accordingly, pri-

mary analysis of these datasets was carried out in volume space (VBA) and surface

space (SBAV, CBA) respectively.

Planned direct comparisons between these three data sets allowed us to evaluate

the effects of different steps of our macroanatomical alignment approach. We com-

pared the VBA and SBAV data sets to assess in isolation the impact of surface-

based preprocessing while keeping anatomical alignment constant. We compared

the SBAV and CBA data sets to assess in isolation the impact of macroanatomical

alignment while keeping preprocessing parameters constant. Finally, we compared

the VBA and CBA data sets to assess the combined impact of both surface-based

preprocessing and macroanatomical alignment.

fMRI group analysis of visual quadrants.

We performed multi-subject statistical analyses using multiple linear regression of the

BOLD signal. The presentation of each checkerboard stimulus sequence at a single

location, was modelled by an ideal box-car function, which covered the volume of

each trial, convolved with a synthetic two-gamma function. These predictors were

used to build the design matrix of the experiment. Individual statistical maps were

generated by associating each voxel with the beta-value corresponding to the spe-

cific set of predictors and calculated on the basis of the least mean squares solution

of the general linear model. The resulting individual statistical maps were entered into

a second-level random-effects group analysis using a summary statistic approach.

We performed analyses focusing on the mapping of the four visual quadrants at the

group level. To define the group-level ROI for each visual quadrant, we computed

separate weighted contrasts for each quadrant against the other three quadrants. We

assigned a weight of three to the position of interest, e.g. $(\beta^{Quad1} \times 3) / (\beta^{Quad2} + \beta^{Quad3}$

$+ \beta^{Quad4})$ ($p < 0.05$, Bonferroni corrected). This allowed us to detect brain regions

showing significant position selectivity. For each resulting group-level ROI, we ex-

tracted average time courses (incl. standard errors of the mean = SEM) for all four

conditions. We conducted this analysis for all three data sets (VBA, SBAV, CBA). For

the VBA data set, we computed this analysis completely in volume space using the

original resolution of the functional data (voxel size: $3 \times 3 \times 3$ mm^3). We projected the

resulting maps on the non-aligned average surface representation to achieve an ap-

propriate visualization and to allow a comparable quantification of cluster size for the

resulting ROIs, i.e., vertices instead of voxels. To this end, volumetric functional

maps were transformed into surface space by sampling on the average cortical sur-

face incorporating data from −1 to +3 mm along vertex normals of the group average

surface brain using trilinear interpolation.

We used two approaches to determine, whether position selectivity differed between

our three data sets: First, to assess differences in the extent of position selectivity

across early visual areas, we compared the ROI size, i.e. the number of vertices, for

each position of interest across data sets. Second, to test for differences in the

strength of position selectivity within corresponding ROIs we conducted separate lin-

ear mixed models with random intercept for each visual quadrant using the t-values

of each subject as the dependent variable and the alignment methods (VBA, SBAV

and CBA) as the independent variable. To correct for multiple comparisons, p-values

were adjusted using Bonferroni correction.

Finally, to assess the impact of the three alignment approaches on horizontal and

vertical symmetry of our group-level ROIs, we computed an asymmetry index (AI)

based on ROI size, i.e the number of vertices, following an established procedure[92],

between each pair of ROIs using the following formula: (| size^{ROI1} - size^{ROI2} | /

size^{ROI1} + size^{ROI2}) * 100.

Probability maps.

To quantify and visualize variability of functional activation and possible changes due

to macroanatomical alignment, the use of PMs has been proposed. PMs are specifi-

cally useful to assess inconsistencies, i.e., disparities between individuals regarding

the location of a particular (visual) area[93,94]. To quantify the spatial consistency of po-

sition selective activation patterns, we generated PMs for each visual quadrant for all

three data sets (VBA, SBAV, CBA). These maps represent the relative number of

subjects showing significant task activity in our single-subject analysis. To this end,

we generated single-subject t-maps based on the same weighted contrasts employed

in the group analysis but set at a more lenient statistical threshold ($p < 0.05$ uncor-

rected). PMs were calculated by counting the number of subjects showing above-

threshold activation in their individual t-maps at a given vertex, dividing this value by

the total number of subjects, and multiplying the result by 100. For the VBA data sets,

we computed all of these steps in volume space and transformed the final PM into

surface space using the same parameters outlined above for the volumetric group

maps. Finally, all PMs were thresholded at a minimum of ten percent probability of

activation overlap. We also applied a cluster level threshold of 100 vertices to focus

on the main areas of interest, i.e., the visual quadrants. Additionally, we counted the

number of vertices in the corresponding probability maps exceeding the threshold of

ten percent probability of activation overlap for each visual quadrant and analysis

methods. Our goal was to quantify and compare the extent of early visual cortex,

where each analysis method had a relevant impact on the probability of activation

overlap.

Probability difference maps.

Additionally, we aimed to quantify changes in spatial consistency of position selective

activation patterns as a result of the different alignment methods. To this end, using

the original unthresholded PMs we calculated PDMs for each visual quadrant,

thresholded at a minimum probability difference of five percent. The resulting three

PDMs capture different aspects of our overall approach: the impact of surface-based

functional data readout and pre-processing compared to volume-based alignment

(SBAV minus VBA), the additional impact of applying macroanatomical alignment

(CBA minus SBAV) and the additive impact of both methods (CBA minus VBA). Addi-

tionally, we counted the number of vertices in the corresponding PDMs exceeding the

threshold of 5 percent difference in probability of activation overlap for each visual

quadrant. Our goal was to quantify and compare the extent of early visual cortex,

where we observed a difference in the probability of activation overlap for a compari-

son of analysis method.

699 Single-subject ROI peak vertex distribution mapping.

For single-subject level analyses, we first defined ROIs for each subject inde-

pendently before and after macroanatomical alignment, i.e., for SBAV and CBA, us-

ing the same weighted contrasts employed in the group analysis. We applied a more

lenient statistical threshold ($p < 0.05$ uncorrected). Next, we determined the peak ver-

tex for each subject's four visual quadrant ROIs, i.e., the vertex with the highest t-

value, for SBAV and CBA. To specifically assess the impact of macroanatomical

alignment on the overlap of single-subject ROI peak vertices for each visual quad-

rant, we mapped all peak vertices per visual quadrant for SBAV and CBA. To quan-

tify changes in the number of precisely overlapping single-subject peak vertices, we

counted for each occipital vertex the number of peak vertices for SBAV and CBA. We

performed this analysis in addition to the PM- and PDM-analysis to provide a more

direct assessment and visualization of the effects of macroanatomical alignment on

the spatial correspondence of single-subject ROIs.

REFERENCES

Wandell, B. A., Dumoulin, S. O. & Brewer, A. A. Visual field maps in human cortex. *Neuron* **56**, 366-383, doi:10.1016/j.neuron.2007.10.012 (2007).

Corbetta, M. & Shulman, G. L. Control of goal-directed and stimulus-driven attention in the brain. *Nature Reviews Neuroscience* **3**, 201-215, doi:10.1038/nrn755 (2002).

Das, M., Bennett, D. M. & Dutton, G. N. Visual attention as an important visual function: an outline of manifestations, diagnosis and management of impaired visual attention. *Br J Oph-thalmol* **91**, 1556-1560, doi:10.1136/bjo.2006.104844 (2007).

de Haan, B., Bither, M., Brauer, A. & Karnath, H. O. Neural Correlates of Spatial Attention and Target Detection in a Multi-Target Environment. *Cereb Cortex* **25**, 2321-2331, doi:10.1093/cercor/bhu046 (2015).

Goodale, M. & Milner, D. One brain - Two visual systems. *Psychologist* **19**, 660-663 (2006).

Bergmann, J., Genç, E., Kohler, A., Singer, W. & Pearson, J. Neural Anatomy of Primary Visual Cortex Limits Visual Working Memory. *Cerebral Cortex* **26**, 43-50, doi:10.1093/cercor/bhu168 (2014).

Lee, J. *et al.* fMRI evidence of aberrant neural adaptation for objects in schizophrenia and bipolar disorder. *Hum Brain Mapp* **40**, 1608-1617, doi:10.1002/hbm.24472 (2019).

Silverstein, S. M. *et al.* An fMRI examination of visual integration in schizophrenia. *J Integr Neurosci* **8**, 175-202, doi:10.1142/s0219635209002113 (2009).

Dumoulin, S. O. & Wandell, B. A. Population receptive field estimates in human visual cortex. *Neuroimage* **39**, 647-660, doi:10.1016/j.neuroimage.2007.09.034 (2008).

Sereno, M. I. *et al.* Borders of multiple visual areas in humans revealed by functional magnetic resonance imaging. *Science (New York, N.Y.)* **268**, 889-893, doi:10.1126/science.7754376 (1995).

Harrison, S. A. & Tong, F. Decoding reveals the contents of visual working memory in early visual areas. *Nature* **458**, 632-635, doi:10.1038/nature07832 (2009).

Peters, B., Kaiser, J., Rahm, B. & Bledowski, C. Activity in Human Visual and Parietal Cortex Reveals Object-Based Attention in Working Memory. *The Journal of Neuroscience* **35**, 3360-3369, doi:10.1523/jneurosci.3795-14.2015 (2015).

Downing, P. E., Chan, A. W., Peelen, M. V., Dodds, C. M. & Kanwisher, N. Domain specificity in visual cortex. *Cereb Cortex* **16**, 1453-1461, doi:10.1093/cercor/bhj086 (2006).

Kanwisher, N., McDermott, J. & Chun, M. M. The fusiform face area: a module in human extrastriate cortex specialized for face perception. *J Neurosci* **17**, 4302-4311, doi:10.1523/jneurosci.17-11-04302.1997 (1997).

Brett, M., Johnsrude, I. S. & Owen, A. M. The problem of functional localization in the human brain. *Nat Rev Neurosci* **3**, 243-249, doi:10.1038/nrn756 (2002).

Desai, R., Liebenthal, E., Possing, E. T., Waldron, E. & Binder, J. R. Volumetric vs. surface-based alignment for localization of auditory cortex activation. *Neuroimage* **26**, 1019-1029, doi:10.1016/j.neuroimage.2005.03.024 (2005).

Dougherty, R. F. *et al.* Visual field representations and locations of visual areas V1/2/3 in human visual cortex. *J Vis* **3**, 586-598, doi:10.1167/3.10.1 (2003).

Fischl, B., Sereno, M. I. & Dale, A. M. Cortical surface-based analysis. II: Inflation, flattening, and a surface-based coordinate system. *Neuroimage* **9**, 195-207, doi:10.1006/nimg.1998.0396 (1999).

Frost & Goebel. Measuring structural-functional correspondence: spatial variability of specialised brain regions after macro-anatomical alignment. *Neuroimage* **59**, 1369-1381, doi:10.1016/j.neuroimage.2011.08.035 (2012).

20   Frost & Goebel. Functionally informed cortex based alignment: an integrated approach for
whole-cortex macro-anatomical and ROI-based functional alignment. *NeuroImage* **83**, 1002-
1010, doi:10.1016/j.neuroimage.2013.07.056 (2013).
21   Steinmetz, H., Fürst, G. & Freund, H. J. Variation of perisylvian and calcarine anatomic land-
marks within stereotaxic proportional coordinates. *AJNR Am J Neuroradiol* **11**, 1123-1130
(1990).
22   Van Essen, D. C. *et al.* An integrated software suite for surface-based analyses of cerebral
cortex. *J Am Med Inform Assoc* **8**, 443-459, doi:10.1136/jamia.2001.0080443 (2001).
23   Zilles, K. *et al.* Quantitative analysis of sulci in the human cerebral cortex: Development, re-
gional heterogeneity, gender difference, asymmetry, intersubject variability and cortical ar-
chitecture. *Human Brain Mapping* **5**, 218-221, doi:https://doi.org/10.1002/(SICI)1097-
0193(1997)5:4<218::AID-HBM2>3.0.CO;2-6 (1997).
24   Bridge, H. Mapping the visual brain: how and why. *Eye (Lond)* **25**, 291-296,
doi:10.1038/eye.2010.166 (2011).
25   Fedorenko, E., Hsieh, P. J., Nieto-Castañón, A., Whitfield-Gabrieli, S. & Kanwisher, N. New
method for fMRI investigations of language: defining ROIs functionally in individual subjects.
J Neurophysiol **104**, 1177-1194, doi:10.1152/jn.00032.2010 (2010).
26   Julian, J. B., Fedorenko, E., Webster, J. & Kanwisher, N. An algorithmic method for function-
ally defining regions of interest in the ventral visual pathway. *NeuroImage* **60**, 2357-2364,
doi:10.1016/j.neuroimage.2012.02.055 (2012).
27   Nieto-Castañón, A. & Fedorenko, E. Subject-specific functional localizers increase sensitivity
and functional resolution of multi-subject analyses. *Neuroimage* **63**, 1646-1669,
doi:10.1016/j.neuroimage.2012.06.065 (2012).
28   Smith, S. M. *et al.* Functional connectomics from resting-state fMRI. *Trends in cognitive sci-
ences* **17**, 666-682, doi:10.1016/j.tics.2013.09.016 (2013).
29   Talairach, J., & Tournoux, P. *Co-planar stereotaxic atlas of the human brain: 3-dimensional
proportional system : an approach to cerebral imaging.* (Thieme, 1988).
30   Evans, A. C. *et al.* 3D statistical neuroanatomical models from 305 MRI volumes. *1993 IEEE
Conference Record Nuclear Science Symposium and Medical Imaging Conference*, 1813-1817
vol.1813 (1993).
31   Klein, A. *et al.* Evaluation of volume-based and surface-based brain image registration meth-
ods. *Neuroimage* **51**, 214-220, doi:10.1016/j.neuroimage.2010.01.091 (2010).
32   Ghosh, S. S. *et al.* Evaluating the validity of volume-based and surface-based brain image reg-
istration for developmental cognitive neuroscience studies in children 4 to 11 years of age.
NeuroImage **53**, 85-93, doi:10.1016/j.neuroimage.2010.05.075 (2010).
33   Nieto-Castanon, A., Ghosh, S. S., Tourville, J. A. & Guenther, F. H. Region of interest based
analysis of functional imaging data. *Neuroimage* **19**, 1303-1316, doi:10.1016/s1053-
8119(03)00188-5 (2003).
34   Fischl, B., Sereno, M. I., Tootell, R. B. & Dale, A. M. High-resolution intersubject averaging and
a coordinate system for the cortical surface. *Hum Brain Mapp* **8**, 272-284,
doi:10.1002/(sici)1097-0193(1999)8:4<272::aid-hbm10>3.0.co;2-4 (1999).
35   Brodoehl, S., Gaser, C., Dahnke, R., Witte, O. W. & Klingner, C. M. Surface-based analysis in-
creases the specificity of cortical activation patterns and connectivity results. *Scientific Re-
ports* **10**, 5737, doi:10.1038/s41598-020-62832-z (2020).
36   Blazejewska, A. I., Fischl, B., Wald, L. L. & Polimeni, J. R. Intracortical smoothing of small-voxel
fMRI data can provide increased detection power without spatial resolution losses compared
to conventional large-voxel fMRI data. *Neuroimage* **189**, 601-614,
doi:10.1016/j.neuroimage.2019.01.054 (2019).

37   Fischl, B. *et al.* Cortical folding patterns and predicting cytoarchitecture. *Cereb Cortex* **18**,
1973-1980, doi:10.1093/cercor/bhm225 (2008).
38   Anticevic, A. *et al.* Comparing surface-based and volume-based analyses of functional
neuroimaging data in patients with schizophrenia. *Neuroimage* **41**, 835-848,
doi:10.1016/j.neuroimage.2008.02.052 (2008).
39   Pantazis, D. *et al.* Comparison of landmark-based and automatic methods for cortical surface
registration. *Neuroimage* **49**, 2479-2493, doi:10.1016/j.neuroimage.2009.09.027 (2010).
40   Van Essen, D. C. & Drury, H. A. Structural and functional analyses of human cerebral cortex
using a surface-based atlas. *J Neurosci* **17**, 7079-7102, doi:10.1523/jneurosci.17-18-
07079.1997 (1997).
41   Frost, M. A., Esposito, F. & Goebel, R. Improved correspondence of resting-state networks
after macroanatomical alignment. *Hum Brain Mapp* **35**, 673-682, doi:10.1002/hbm.22191
(2014).
42   Hinds, O. P. *et al.* Accurate prediction of V1 location from cortical folds in a surface
coordinate system. *NeuroImage* **39**, 1585-1599, doi:10.1016/j.neuroimage.2007.10.033
(2008).
43   Weiner, K. S. *et al.* Defining the most probable location of the parahippocampal place area
using cortex-based alignment and cross-validation. *Neuroimage* **170**, 373-384,
doi:10.1016/j.neuroimage.2017.04.040 (2018).
44   Rosenke, M., van Hoof, R., van den Hurk, J., Grill-Spector, K. & Goebel, R. A Probabilistic
Functional Atlas of Human Occipito-Temporal Visual Cortex. *Cerebral Cortex*,
doi:10.1093/cercor/bhaa246 (2020).
45   Huang, T., Chen, X., Jiang, J., Zhen, Z. & Liu, J. A probabilistic atlas of the human motion
complex built from large-scale functional localizer data. *Hum Brain Mapp* **40**, 3475-3487,
doi:10.1002/hbm.24610 (2019).
46   Di Russo, F., Martínez, A. & Hillyard, S. A. Source analysis of event-related cortical activity
during visuo-spatial attention. *Cereb Cortex* **13**, 486-499, doi:10.1093/cercor/13.5.486
(2003).
47   Di Russo, F., Martínez, A., Sereno, M. I., Pitzalis, S. & Hillyard, S. A. Cortical sources of the
early components of the visual evoked potential. *Hum Brain Mapp* **15**, 95-111,
doi:10.1002/hbm.10010 (2002).
48   Shigihara, Y., Hoshi, H. & Zeki, S. Early visual cortical responses produced by checkerboard
pattern stimulation. *NeuroImage* **134**, doi:10.1016/j.neuroimage.2016.03.078 (2016).
49   Kraft, A. *et al.* fMRI localizer technique: efficient acquisition and functional properties of
single retinotopic positions in the human visual cortex. *NeuroImage* **28**, 453-463,
doi:10.1016/j.neuroimage.2005.05.050 (2005).
50   Bressler, D. W. & Silver, M. A. Spatial attention improves reliability of fMRI retinotopic
mapping signals in occipital and parietal cortex. *Neuroimage* **53**, 526-533,
doi:10.1016/j.neuroimage.2010.06.063 (2010).
51   Anderson, J., Cameron, E. & Levine, M. A method for quantifying visual field
inhomogeneities. *Vision Research* **105**, doi:10.1016/j.visres.2014.09.010 (2014).
52   Liu, T., Heeger, D. J. & Carrasco, M. Neural correlates of the visual vertical meridian
asymmetry. *J Vis* **6**, 1294-1306, doi:10.1167/6.11.12 (2006).
53   O'Connell, C. *et al.* Structural and functional correlates of visual field asymmetry in the
human brain by diffusion kurtosis MRI and functional MRI. *Neuroreport* **27**, 1225-1231,
doi:10.1097/wnr.0000000000000682 (2016).
54   Rubin, N., Nakayama, K. & Shapley, R. Enhanced perception of illusory contours in the lower
versus upper visual hemifields. *Science* **271**, 651-653, doi:10.1126/science.271.5249.651
(1996).

55 Bullmore, E. & Sporns, O. Complex brain networks: graph theoretical analysis of structural
and functional systems. *Nature Reviews Neuroscience* **10**, 186-198, doi:10.1038/nrn2575
(2009).
56 Chen, H., Yao, D. & Liu, Z. A study on asymmetry of spatial visual field by analysis of the fMRI
BOLD response. *Brain Topogr* **17**, 39-46, doi:10.1023/b:brat.0000047335.00110.6a (2004).
57 Loughnane, G. M., Shanley, J. P., Lalor, E. C. & O'Connell, R. G. Behavioral and
electrophysiological evidence of opposing lateral visuospatial asymmetries in the upper and
lower visual fields. *Cortex* **63**, 220-231, doi:10.1016/j.cortex.2014.09.003 (2015).
58 Eickhoff, S. B., Rottschy, C., Kujovic, M., Palomero-Gallagher, N. & Zilles, K. Organizational
Principles of Human Visual Cortex Revealed by Receptor Mapping. *Cerebral Cortex* **18**, 2637-
2645, doi:10.1093/cercor/bhn024 (2008).
59 Curcio, C. A. & Allen, K. A. Topography of ganglion cells in human retina. *J Comp Neurol* **300**,
5-25, doi:10.1002/cne.903000103 (1990).
60 Silson, E. H., Reynolds, R. C., Kravitz, D. J. & Baker, C. I. Differential Sampling of Visual Space
in Ventral and Dorsal Early Visual Cortex. *The Journal of Neuroscience* **38**, 2294-2303,
doi:10.1523/jneurosci.2717-17.2018 (2018).
61 Hagler, D. J., Jr. Visual field asymmetries in visual evoked responses. *J Vis* **14**, 13,
doi:10.1167/14.14.13 (2014).
62 Schmidtmann, G., Logan, A. J., Kennedy, G. J., Gordon, G. E. & Loffler, G. Distinct lower visual
field preference for object shape. *J Vis* **15**, 18, doi:10.1167/15.5.18 (2015).
63 Thomas, N. A. & Elias, L. J. Upper and lower visual field differences in perceptual
asymmetries. *Brain Res* **1387**, 108-115, doi:10.1016/j.brainres.2011.02.063 (2011).
64 Zito, G. A., Cazzoli, D., Müri, R. M., Mosimann, U. P. & Nef, T. Behavioral Differences in the
Upper and Lower Visual Hemifields in Shape and Motion Perception. *Front Behav Neurosci*
10, 128-128, doi:10.3389/fnbeh.2016.00128 (2016).
65 Bakroon, A. & Lakshminarayanan, V. Visual function in autism spectrum disorders: a critical
review. *Clin Exp Optom* **99**, 297-308, doi:10.1111/cxo.12383 (2016).
66 Butler, P. D. *et al.* Dysfunction of early-stage visual processing in schizophrenia. *Am J*
Psychiatry **158**, 1126-1133, doi:10.1176/appi.ajp.158.7.1126 (2001).
67 Butler, P. D., Silverstein, S. M. & Dakin, S. C. Visual perception and its impairment in
schizophrenia. *Biol Psychiatry* **64**, 40-47, doi:10.1016/j.biopsych.2008.03.023 (2008).
68 Chen, C. Y. *et al.* Visual spatial attention in children with attention deficit hyperactivity
disorder. *Chang Gung Med J* **25**, 514-521 (2002).
69 Hale, T. S. *et al.* Visual Network Asymmetry and Default Mode Network Function in ADHD: An
fMRI Study. *Front Psychiatry* **5**, 81, doi:10.3389/fpsyt.2014.00081 (2014).
70 Sanz-Cervera, P., Pastor-Cerezuela, G., González-Sala, F., Tárraga-Mínguez, R. & Fernández-
Andrés, M.-I. Sensory Processing in Children with Autism Spectrum Disorder and/or Attention
Deficit Hyperactivity Disorder in the Home and Classroom Contexts. *Front Psychol* **8**, 1772-
1772, doi:10.3389/fpsyg.2017.01772 (2017).
71 Seymour, R. A., Rippon, G., Gooding-Williams, G., Schoffelen, J. M. & Kessler, K. Dysregulated
oscillatory connectivity in the visual system in autism spectrum disorder. *Brain* **142**, 3294-
3305, doi:10.1093/brain/awz214 (2019).
72 Shimizu, V. T., Bueno, O. F. & Miranda, M. C. Sensory processing abilities of children with
ADHD. *Braz J Phys Ther* **18**, 343-352, doi:10.1590/bjpt-rbf.2014.0043 (2014).
73 Bittner, R. A. *et al.* The When and Where of Working Memory Dysfunction in Early-Onset
Schizophrenia-A Functional Magnetic Resonance Imaging Study. *Cereb Cortex* **25**, 2494-2506,
doi:10.1093/cercor/bhu050 (2015).

74   Butler, P. D., Thompson, J. L., Seitz, A. R., Deveau, J. & Silverstein, S. M. Visual perceptual
905   remediation for individuals with schizophrenia: Rationale, method, and three case studies.
906   *Psychiatr Rehabil J* **40**, 43-52, doi:10.1037/prj0000212 (2017).
907   75   Haenschel, C. *et al.* Contribution of impaired early-stage visual processing to working
908   memory dysfunction in adolescents with schizophrenia: a study with event-related potentials
909   and functional magnetic resonance imaging. *Arch Gen Psychiatry* **64**, 1229-1240,
910   doi:10.1001/archpsyc.64.11.1229 (2007).
911   76   Manoach, D. S. Prefrontal cortex dysfunction during working memory performance in
912   schizophrenia: reconciling discrepant findings. *Schizophr Res* **60**, 285-298,
913   doi:10.1016/s0920-9964(02)00294-3 (2003).
914   77   French, L. & Paus, T. A FreeSurfer view of the cortical transcriptome generated from the Allen
915   Human Brain Atlas. *Frontiers in Neuroscience* **9**, doi:10.3389/fnins.2015.00323 (2015).
916   78   Gulban, O. F. *et al.* Improving a probabilistic cytoarchitectonic atlas of auditory cortex using a
917   novel method for inter-individual alignment. *eLife* **9**, e56963, doi:10.7554/eLife.56963
918   (2020).
919   79   Hawrylycz, M. J. *et al.* An anatomically comprehensive atlas of the adult human brain
920   transcriptome. *Nature* **489**, 391-399, doi:10.1038/nature11405 (2012).
921   80   Sabuncu, M. R. *et al.* Function-based Intersubject Alignment of Human Cortical Anatomy.
922   *Cerebral Cortex* **20**, 130-140, doi:10.1093/cercor/bhp085 (2009).
923   81   Conroy, B. R., Singer, B. D., Guntupalli, J. S., Ramadge, P. J. & Haxby, J. V. Inter-subject
924   alignment of human cortical anatomy using functional connectivity. *NeuroImage* **81**, 400-
925   411, doi:https://doi.org/10.1016/j.neuroimage.2013.05.009 (2013).
926   82   Glasser, M. F. *et al.* A multi-modal parcellation of human cerebral cortex. *Nature* **536**, 171-
927   178, doi:10.1038/nature18933 (2016).
928   83   Gulban, O. F. *et al.* Improving a probabilistic cytoarchitectonic atlas of auditory cortex using a
929   novel method for inter-individual alignment. *Elife* **9**, doi:10.7554/eLife.56963 (2020).
930   84   Oldfield, R. C. The assessment and analysis of handedness: the Edinburgh inventory.
931   *Neuropsychologia* **9**, 97-113, doi:10.1016/0028-3932(71)90067-4 (1971).
932   85   Goebel, R., Esposito, F. & Formisano, E. Analysis of functional image analysis contest (FIAC)
933   data with brainvoyager QX: From single-subject to cortically aligned group general linear
934   model analysis and self-organizing group independent component analysis. *Hum Brain Mapp*
935   **27**, 392-401, doi:10.1002/hbm.20249 (2006).
936   86   Sled, J. G., Zijdenbos, A. P. & Evans, A. C. in *Proceedings of the 15th International Conference*
937   *on Information Processing in Medical Imaging*   459–464 (Springer-Verlag, 1997).
938   87   Hou, Z., Huang, S., Hu, Q. & Nowinski, W. L. A fast and automatic method to correct intensity
939   inhomogeneity in MR brain images. *Med Image Comput Comput Assist Interv* **9**, 324-331,
940   doi:10.1007/11866763_40 (2006).
941   88   Kriegeskorte, N. & Goebel, R. An Efficient Algorithm for Topologically Correct Segmentation
942   of the Cortical Sheet in Anatomical MR Volumes. *NeuroImage* **14**, 329-346 (2001).
943   89   Fan, X. *et al.* Metabolic effects of adjunctive aripiprazole in clozapine-treated patients with
944   schizophrenia. *Acta Psychiatr Scand* **127**, 217-226, doi:10.1111/acps.12009 (2013).
945   90   Breman, H. *et al.* in *Biomedical Image Registration.* (eds Žiga Špiclin, Jamie McClelland, Jan
946   Kybic, & Orcun Goksel) 122-130 (Springer International Publishing).
947   91   Greve, D. N. & Fischl, B. Accurate and robust brain image alignment using boundary-based
948   registration. *Neuroimage* **48**, 63-72, doi:10.1016/j.neuroimage.2009.06.060 (2009).
949   92   Heckemann, R. A. *et al.* Automatic morphometry in Alzheimer's disease and mild cognitive
950   impairment. *NeuroImage* **56**, 2024-2037, doi:10.1016/j.neuroimage.2011.03.014 (2011).
951   93   Wang, L., Mruczek, R. E., Arcaro, M. J. & Kastner, S. Probabilistic Maps of Visual Topography
952   in Human Cortex. *Cereb Cortex* **25**, 3911-3931, doi:10.1093/cercor/bhu277 (2015).

953 94   Yamamoto, H. *et al.* Inconsistency and uncertainty of the human visual area loci following
954   surface-based registration: Probability and Entropy Maps. *Hum Brain Mapp* **33**, 121-129,
955   doi:10.1002/hbm.21200 (2012).
956

ACKNOWLEDGEMENTS

The authors are most grateful to Deliah Macht, Christina Raab and Leonie Winkler-

Lauble for help with data acquisition. C.V Barnes-Scheufler was supported by a

"main doctus" scholarship from The Polytechnic Foundation of Frankfurt am Main. E.

Raspor was supported by a Research Grant for Doctoral Programmes in Germany

from the German Academic Exchange Service (DAAD).

AUTHOR CONTRIBUTIONS

All authors made substantial contributions to the conception or design of the work, or

the acquisition, analysis, or interpretation of data. AR and RAB acquired funding.

RAB, BP, MS and LR designed the experiment. MQ, CVB-S, LR and RAB acquired

the data. ER and RG provided analytical tools. MQ, CS and RAB analyzed the data.

MQ and RAB undertook the literature searches and wrote the first draft of the manu-

script. All authors contributed to and revised the manuscript. All authors read and ap-

proved the final manuscript.

DISCLOSURES

The authors have declared that there are no conflicts of interest in relation to the sub-

ject of this study.

DATA AVAILABILITY STATEMENT

The data that support the findings of this study are available from the corresponding

author, R.A.B., upon reasonable request.

FIGURE LEGENDS

Figure 1.

Visual field Localizer Paradigm. (a) The task consisted of flickering, black-and-white

colored checkerboards that appeared randomly at homologous positions of the par-

ticipant's visual quadrant. Only in 25 percent of the trials the two centrally located

squares changed their color into yellow for 133 ms. Participants were required to

press a response box button when noticing that. During the whole task participants

were instructed to fixate a black, x-shaped fixation cross presented in the center of

the screen. Checkerboards appeared for 2000 ms. The regular inter-stimulus interval

was 0 ms. (b) Every 10 to 14 trials, the ITI extended to 2000 ms. The task comprised

144 trials (25 percent target trials). It was preceded and followed by a presentation of

the fixation cross for 10 seconds.

Figure 2.

CBA consisted of a rigid alignment to a single target brain and a non-linear alignment

to an iteratively updated group average brain. **(a)** We carried out an initial CBA solely

to generate an unbiased average target brain for the final CBA. A randomly selected

brain from among all participants was used for the initial rigid CBA. **(b)** For the final

CBA we used the unbiased average target brain created during the initial CBA for

rigid CBA. **(c)** We generated average surface representations before and after

macroanatomical alignment for each hemisphere, which we subsequently merged,

inflated and used for analysis and visualization of the appropriate data sets.

1002

Figure 3.

Sequences of functional data pre-processing, coregistration of structural and func-

tional data and spatial transformation operations used to generate the three func-

tional data sets used in our study: VBA, SBAV and CBA. For VBA we conducted all

data pre-processing operations in volume space, including slice-scan-time correction,

3D motion correction, echo-planar imaging distortion correction, 3D spatial smoothing

and linear trend removal with temporal high-pass filtering. Finally, functional data

were co-registered to the structural data and transformed into Talairach space. For

SBAV and CBA, we conducted all data pre-processing operations up to echo-planar

imaging distortion correction in volume space. Here, co-registration of functional data

to the structural data and transformation into Talairach space was followed by trans-

formation into surface space. We then conducted 2D spatial smoothing and linear

trend removal with temporal high-pass filtering in surface space. For CBA only, we

then applied macroanatomical alignment.

1017

Figure 4.

Group analysis of visual quadrants. **(a)** VBA results. Maps and average timecourses

were computed in volume space; maps were projected on the non-aligned average

surface representation. **(b)** SBAV results. Maps and average timecourses were com-

puted in surface space; maps were projected on the non-aligned surface representa-

tion. **(c)** CBA results. Maps and average timecourses were computed in surface

space; maps were projected on the aligned average surface representation. Overall,

three out of four visual quadrant ROIs exhibited a pattern of continuously increasing

cluster size reflecting an increasing extent of significant position selectivity across

alignment techniques. Conversely, the ROI of the lower right visual quadrant showed

a cluster size decrease after CBA. Average timecourses (incl. SEM) showed clear

position selectivity with a strong BOLD signal increase for the position of interest and

no BOLD signal increase for the other three positions. ROI/graph colors: light-blue =

lower right (LR) visual quadrant, orange = lower left (LL) visual quadrant, red = upper

left (UL) visual quadrant, dark-blue = upper right (UR) visual quadrant.

Figure 5.

Probability maps (PMs). PMs indicating the probability of activation overlap across

subjects for each visual quadrant. The color code gray-to-white indicates the proba-

bility of activation overlap of single-subject maps thresholded at a minimum of ten

percent probability of activation overlap. Single-subject maps were thresholded at

p<0.05 (uncorr.). PMs were thresholded at a minimum of ten percent probability of

activation overlap. We also applied a cluster level threshold of 100 vertices. **(a)** PMs

for VBA showed a maximum probability of activation overlap of up to 50 percent. **(b)**

PMs for SBAV showed a maximum probability of activation overlap of up to 45 per-

cent. **(c)** PMs for CBA showed a maximum probability of activation overlap of up to

75 percent.

1044

Figure 6.

Probability Difference Maps (PDMs). PDMs indicating the differential impact of the in-

dividual steps of our overall macroanatomical alignment approach for each visual

quadrant. PDMs were generated using PMs derived from single-subject maps

thresholded at minimum of zero percent probability. The color code indicates the dif-

ference of activation overlap. The color code brown-to-white indicates a higher de-

gree of functional activation overlap for the more advanced alignment method. The

color code blue-to-green indicates a higher degree of functional activation overlap for

the less advanced alignment method. PDMs were thresholded at a minimum proba-

bility difference of five percent. **(a)** The impact of surface-based functional data

readout and pre-processing compared to standard volume-based alignment (SBAV

minus VBA) was characterized by a widespread activation with an increase in the

probability of activation overlap of up to 15 percent around the central ROIs and a de-

crease in the probability of activation overlap of up to 10 at the location of the central

ROIs. **(b)** The additional impact of macroanatomical alignment (CBA minus SBAV)

was less widespread but characterized by an increase in the probability of activation

overlap of up to 35 percent at the location of the central ROIs and a decrease in the

probability of activation overlap of up to 35 percent around the central ROIs. **(c)** The

additive impact of both methodological elements (CBA minus VBA) was character-

ized by an increase in the probability of activation overlap of up to 30 percent at the

location of the central ROIs and a decrease in the probability of activation overlap of

up to 20 percent around the central ROIs.

1067

Figure 7.

Single-subject peak vertex distribution maps for SBAV and CBA data sets. We

mapped single-subject peak vertices for each visual quadrant in surface space for

SBAV and CBA data. We then computed the vertex-wise number of single-subject

peak vertices. The color code indicates the number of overlapping single-subject

peak vertices per vertex. An increase in the number of overlapping single-subject

ROI peak vertices per vertex was observed after macroanatomical alignment (CBA).

For each visual quadrant, single subject peak vertices were mapped in surface space

and the vertex-wise number of single subject peak vertices was computed before and

after macroanatomical alignment. The color code indicates the number of overlapping

single-subject peak vertices per vertex. A nominal increase in the degree of overlap

single subject peak vertices was observed after macroanatomical alignment (CBA).

TABLE LEGENDS

Table 1.

Group ROIs. Size and Talairach coordinates of the group ROIs of the corresponding

visual quadrants for the VBA, SBAV and CBA data sets. Three out of four visual

quadrant ROIs showed a consistent pattern of increasing cluster, i.e., an increasing

extent of position selectivity across alignment techniques.

1086

Table 2.

Linear mixed model results. To test for differences in the strength of position selectiv-

ity within corresponding ROIs across alignment techniques, we conducted separate

linear mixed models with random intercept for each visual quadrant. We used each

subject's t-values as the dependent variable and the alignment techniques (VBA,

SBAV and CBA) as the independent variable. We adjusted p-values using Bonferroni

correction. We did not observe any significant effect, indicating that this measure off

position selectivity was not affected by alignment technique.

* Random effect variance estimate at the subject-level for the LR ROI was 0, result-

ing in a singular fit when using lmer. Therefore, results for LR were estimated without

a random intercept for ID and are thus equivalent to a regular ANOVA.

1098

Table 3.

Vertical and horizontal asymmetry indices (AIs). To assess the impact of the three

alignment techniques on horizontal and vertical symmetry of our group-level ROIs,

we computed an ROI size asymmetry index between each pair of ROIs. AIs revealed

greater vertical symmetry of the upper and lower hemifield ROIs for VBA and CBA

compared to SBAV.

1105

Table 4.

Extent of probability maps. For each visual quadrant and analysis methods, we

counted the number of vertices in the corresponding probability maps exceeding the

threshold of 10 percent probability of activation overlap. For VBA, maximum probabil-

ity of activation overlap was around 50 percent. For SBAV, maximum probability of

activation overlap was around 45 percent. For CBA, maximum probability of activa-

tion overlap was around 75 percent.

1113

Table 5.

Extent of probability difference maps (PDMs). For each visual quadrant and compari-

son between analysis methods (SBAV minus VBA, CBA minus SBAV, CBA minus

VBA), we counted the number of vertices in the corresponding PDMs exceeding the

threshold of 5 percent difference in probability of activation overlap. Overall, the ex-

tent of PDMs was consistently greatest for the CBA minus SBAV comparison, i.e.,

the isolated effect of effect of macroanatomical alignment.

1121

Table 6.

Single-subject ROI peak vertex distribution mapping. Number of overlapping single-

subject peak vertices for each visual quadrant before and after macroanatomical

alignment (SBAV and CBA). Overall, there was an increase in the number of multiple

overlapping single-subject ROI peak vertices per vertex after macroanatomical align-

ment (CBA) for each visual quadrant and a corresponding decrease in the number of

single overlapping single-subject ROI peak vertices per vertex. PV = ROI peak ver-

tex.

a

50 participants   Target (randomly drawn)   Rigid CBA   Non-rigid CBA   Final unbiased target for rigid CBA

b

50 participants   Rigid CBA   Non-rigid CBA   Final aligned group average

c

LH   RH

a   VBA

b   SBAV

c   CBA

LR Quadrant   LL Quadrant   UL Quadrant   UR Quadrant

a   VBA

UR Quadrant   UL Quadrant   LR Quadrant   LL Quadrant

b   SBAV

UR Quadrant   UL Quadrant   LR Quadrant   LL Quadrant

c   CBA

UR Quadrant   UL Quadrant   LR Quadrant   LL Quadrant

100%   0%

a SBAV minus VBA

UR Quadrant   UL Quadrant   LR Quadrant   LL Quadrant

b CBA minus SBAV

UR Quadrant   UL Quadrant   LR Quadrant   LL Quadrant

c CBA minus VBA

UR Quadrant   UL Quadrant   LR Quadrant   LL Quadrant

40%   0%   - 40%

SBAV   CBA

SBAV   CBA

LR Quadrant

LL Quadrant

UR Quadrant

UL Quadrant

SBAV   CBA

SBAV   CBA

1   5

Table 1

Region of interest	Analysis method	Number of vertices	Number of voxels	TAL coordinates		
				x	y	z
lower right visual quadrant	VBA	47	953	-22	-90	2
	SBAV	295	*NA*	-42	-64	5
	CBA	161	*NA*	-24	-91	-4
lower left visual quadrant	VBA	46	828	19	-94	2
	SBAV	28	*NA*	23	-90	3
	CBA	127	*NA*	9	-94	-5
upper left visual quadrant	VBA	4	*55*	4	-87	-9
	SBAV	47	*NA*	24	-75	-14
	CBA	82	*NA*	18	-77	-15
upper right visual quadrant	VBA	3	1	-5	-88	-13
	SBAV	6	*NA*	-18	-80	-15
	CBA	58	*NA*	-18	-80	-15

Table 2

	Sum Sq	Mean Sq	NumDF	DenDf	F value	Pr (>F)	p-value (corr.)
LR	183.2	92	2	147	0.31	0.737	1.000
LL	2660.2	1330.1	2	98	1.25	0.291	1.000
UL	2546.2	1273.1	2	98	3.36	0.039	0.156
UR	73678	36839	2	98	1.19	0.308	1.000

Table 3

Symmetry	ROI comparison	AI (VBA)	AI (SBAV)	AI (CBA)
vertical	LR and LL	1.1	82.7	11.8
	UL and UR	14.3	77.4	17.1
horizontal	LR and UR	88.0	96.0	47.0
	LL and UL	84.0	25.3	21.5

Table 4

Probability map	Analysis method	Number of vertices
lower right visual quadrant	VBA	1243
	SBAV	2074
	CBA	1391
lower left visual quadrant	VBA	853
	SBAV	1396
	CBA	918
upper left visual quadrant	VBA	958
	SBAV	1401
	CBA	1162
upper right visual quadrant	VBA	965
	SBAV	1897
	CBA	1445

Table 5

Probability difference map	Analysis method	Number of vertices
lower right visual quadrant	SBAV minus VBA	1169
	CBA minus SBAV	1341
	CBA minus VBA	739
lower left visual quadrant	SBAV minus VBA	570
	CBA minus SBAV	694
	CBA minus VBA	402
upper left visual quadrant	SBAV minus VBA	612
	CBA minus SBAV	680
	CBA minus VBA	546
upper right visual quadrant	SBAV minus VBA	689
	CBA minus SBAV	964
	CBA minus VBA	795

Table 6

# of overlapping PV	LR		LL		UL		UR		Overall	
	SBAV	CBA	SBAV	CBA	SBAV	CBA	SBAV	CBA	SBAV	CBA
1	46	42	48	36	40	40	41	12	175	130
2	2	4	0	4	3	5	1	3	6	16
3	0	0	0	0	0	0	1	1	1	1
4	0	0	0	1	1	0	0	0	1	1
5	0	0	0	0	0	0	0	0	0	5