Improved correspondence of fMRI visual field localizer data after macroana-

tomical alignment

Running Title: Visual field localizer & CBA

Qubad M.[1], Barnes-Scheufler C.V.[1], Schaum M.[2], Raspor E.[1], Rösler L.[1,3], Peters B.[4,5], Goe-

bel R.[3,6], Reif A.[1], Bittner R.A.[1,7]*

[1] Department of Psychiatry, Psychosomatic Medicine and Psychotherapy and Brain Imaging

Center, University Hospital, Goethe University, Frankfurt, Germany,

[2] Leibniz Institute for Resilience Research, Mainz, Germany

[3] Netherlands Institute for Neuroscience, Amsterdam, The Netherlands

[4] Institute of Medical Psychology, University Hospital, Goethe University, Frankfurt, Germany,

[5] Zuckerman Mind Brain Behavior Institute, Columbia University, New York, NY, USA

[6] Department of Cognitive Neuroscience, Faculty of Psychology and Neuroscience, Maas-

tricht University, Maastricht, The Netherlands

[7] Ernst Strüngmann Institute for Neuroscience (ESI) in Cooperation with Max Planck Society,

Frankfurt am Main, Germany

*Corresponding author. Department of Psychiatry, Psychosomatic Medicine and Psychother-

apy, University Hospital Frankfurt, Goethe University, Heinrich-Hoffmann-Str. 10, D-60528

Frankfurt am Main, Germany.

Phone: +49 69 6301 84713. Fax: +49 69 6301 81775.

Email: bittner@med.uni-frankfurt.de

Total number of words in abstract: 200/200

Total number of words in main text: 3444/4500

ORCID IDs

M.Q.: orcid.org/0000-0002-9291-5739

C.V.B.-S.: orcid.org/0000-0002-4730-1981

M.S.: orcid.org/0000-0002-6589-4530

E.R.: orcid.org/0000-0003-4418-7084

L.R.: orcid.org/0000-0003-1046-1073

B.P.: orcid.org/0000-0002-0948-8976

R.G.: orcid.org/0000-0003-1780-2467

A.R.: orcid.org/0000-0002-0992-634X

R.A.B.: orcid.org/0000-0003-2021-0358

ABSTRACT

The study of the visual system and its role for human cognition with fMRI often requires the use of localizer paradigms to define regions of interest (ROIs). However, the considerable degree of interindividual variability of the cerebral cortex represents an important confound, when analyzing data on the group level. Cortex-based alignment (CBA) techniques lead to a reliable reduction of interindividual macroanatomical variability. Yet, CBA has not been evaluated for visual field localizer paradigms used to map specific parts of the visual field within retinotopically organized early visual areas. We evaluated CBA for an attention-enhanced visual field localizer mapping a homologous part of each visual quadrant in 50 participants. After CBA, group ROIs showed markedly increased spatial consistency. CBA also led to an increase in the probability of activation overlap of up to forty percent. Furthermore, after CBA group ROI size for the lower visual hemifield was larger than for the upper. This asymmetry, which mirrors previous mapping studies, was not detectable before CBA. Our results confirm and extend the utility of CBA for the study of the visual system in the context of group analyses. CBA should be particularly relevant when studying neuropsychiatric disorders with abnormally increased interindividual macroanatomical variability.

KEYWORDS

INTRODUCTION

The visual system includes a multitude of topographical representations of varying resolution

across increasingly specialized[1] visual areas. Functional magnetic resonance imaging (fMRI)

offers a variety of methods either to map these topographical representations in full, or to lo-

calize specific visual areas or retinotopic positions within their topography. These ap-

proaches are essential not only for the fine-grained study of fundamental properties of the

visual system[1], but also for investigating the role of these areas for higher-order cognitive

processes such as visual attention and working memory[2-6]. This also extends to translational

studies of visual dysfunction and its cognitive consequences in neuropsychiatric disorders[7,8].

Methods for fMRI-based visual mapping, i.e. techniques to define regions of interest in the

visual system based on specific functional properties, fall in in three broad categories: retino-

topic mapping, visual field localizer and functional localizer paradigms. Retinotopic mapping

and the more advanced population receptive field mapping allow the complete delineation of

early visual areas[1,9,10]. Conversely, visual field localizer paradigms can map a circumscribed

region within a retinotopically organized visual area[11,12]. Finally, functional localizers can de-

tect higher-order visual areas such as the fusiform face area, parahippocampal place area

(PPA), extrastriate body area (EBA), lateral occipital complex (LOC), which are clustered and

show specialization for the processing of specific categories of complex visual infor-

mation[1,13,14]. In most fMRI studies, high interindividual anatomical variability of cortical areas

in terms of both size and location constitutes an important challenge[15-23]. For instance, it has

been shown that primary visual cortex (V1) can differ in size by about 2-fold between individ-

uals[17]. Furthermore, anatomical variability in terms of location has been shown to be particu-

larly pronounced in extrastriate visual areas[24]. This crucial confound reduces the power to re-

liably map visual areas at the group level.

One way to mitigate this problem for the visual system is to pool single-subject ROIs, while

simultaneously using the overall group-based probability for that ROI at each point in a Car-

tesian coordinate system as a constraint[25-27]. While such a single-subject-based analysis im-

proves sensitivity and functional resolution compared to a standard group-based approach, it

does not actually reduce macroanatomical variability. Additionally, studying the interplay be-

tween visual areas and other cortical areas more directly involved in higher-order cognitive

processes with whole-brain methods such as functional connectomics network analyses[28]

might preclude a single-subject based strategy.

Such group-based analyses typically require spatial normalization of structural and functional

imaging data to a common Cartesian coordinate system such as Talairach[29] or MNI[30] space.

In its most basic form, volume-based spatial normalization employs a linear transformation

that matches the overall extents of the brains to a standard brain template. While transfor-

mation into Talairach space relies on anatomical landmarks, transformation into MNI space

utilizes fully data driven registration of structural images to an average template brain[30].

While these spatial normalization approaches inherently lead to an alignment of brains, the

underlying algorithms are not specifically geared toward aligning homologous brain struc-

tures. Conversely, more refined methods employ non-linear warping algorithms guided by in-

tensity differences to improve macroanatomical alignment[31]. Thus, all of these methods can

be categorized as volume-based alignment (VBA) techniques.

However, both linear and nonlinear VBA mostly disregard the topological properties of the

cerebral cortex and its geometric features such as sulci and gyri. Consequently, VBA meth-

ods result in a considerable amount of residual inter-subject anatomical variability[32,33].

Surface-based procedures constitute an important alternative approach. Surface-based nor-

malization typically uses a geodesic coordinate system, which allows for a two-dimensional

representation of the cerebral cortex and respects the cortical topography to a much larger

degree than a traditional Cartesian coordinate system[18,34]. Surface-based spatial normaliza-

tion allows to constrain data preprocessing such as spatial smoothing to cortical tissue,

greatly reducing signal contamination by white matter and cerebrospinal fluid. Another ad-

vantage of surface-based spatial normalization is the possibility to use individual cortical fold-

ing patterns for an additional fully data-driven macroanatomical alignment. Compared to VBA

techniques, these cortex-based alignment (CBA) methods considerably improve anatomical

correspondence of cortical structures thus leading to a notable reduction of residual inter-

subject anatomical variability[18,34-37]. CBA has also been shown to correctly align cyto-archi-

tectonic boundaries[38].

Due to these properties CBA methods have been proposed as an alternative approach spe-

cifically for the visual system[26]. Several studies have compared the impact of VBA and CBA

methods on specific visual mapping techniques. For retinotopic mapping an improvement of

functional overlap in both V1 and V2 after CBA has been demonstrated[34,39]. Furthermore, for

functional localizer data CBA substantially increases the overlap of object processing areas

LOC, FFA and PPA across subjects[19,40-42]. Conversely, the effects of CBA on visual field lo-

calizer paradigms mapping specific retinotopic positions have not been studied. Thus, the

utility of CBA has been demonstrated for two of the three main categories of visual mapping

methods, i.e. those methods, which map whole areas, either defined by cytoarchitectonic

(e.g. V1) or functional (e.g. FFA) properties. Conversely, it remains unclear, to which degree

CBA can improve the alignment of ROIs mapped by visual field localizer paradigms. Such

paradigms are required for the detailed study of the local processing of simple visual stimuli

in early visual areas[11,12,43-45]. Flashing chequerboards covering the exact area of interest

within the visual field are primarily used for this purpose. Chequerboards lead to a particu-

larly strong BOLD-signal increase in early visual areas (V1-V3)[46]. To maximize the fidelity of

the resulting localizer maps, visual field localizer paradigms typically utilize the fact that at-

tentional modulation induced by task demands significantly enhances response reliability

across visual areas. This can be accomplished by adding a simple target detection task[47].

We used such an attention-enhanced visual field localizer paradigm to map a circumscribed

location in each visual quadrant across early visual areas in order to define ROIs to be used

for the study of higher cognitive processes. We chose a CBA method using a dynamic group

average as the target brain[19]. Thus, we eliminated the possible confound of a static CBA tar-

get based on an individual brain, whose cortical folding pattern might by chance deviate con-

siderably from the group average.

Our primary goal was to examine the effects of CBA for a visual field localizer paradigm.

More specifically, we aimed to determine, whether CBA can improve the reliability of map-

ping subregions within retinotopically organized visual areas delineated by such a paradigm

at the group level. Based on previous findings for other localizer paradigm classes and the

relatively good structural-functional correspondence in posterior occipital cortex, we expected

to observe a benefit of CBA when aligning subregions within early visual cortex.

Additionally, we compared two methods for the definition of single subject ROIs: the use of

extended peak vertices, i.e. single vertices showing the strongest level of activation in a cer-

tain brain region and their immediately surrounding vertices, and the delineation of a larger

ROI, which includes all vertices above a prespecified threshold, which is kept constant

across brain regions[48]. Our goal was to compare these two approaches and to assess

changes in spatial correspondence after CBA at the single subject level. Mirroring our hy-

potheses for the group analyses, we expected to observe an improved correspondence of

single-subject peak-vertex ROIs after CBA.

Interestingly, several studies have shown differential response properties such as receptive

field size for homologous early visual areas by visual quadrant or hemifield. For instance, im-

proved behavioral performance and higher BOLD-signal amplitudes were observed in the

lower visual hemifield[49-52]. We were therefore also interested, whether we could observe dif-

ferences between upper and lower visual hemifields in our group analysis after CBA. Overall,

the aim of the study was to close an important gap in the evaluation of CBA for the study of

the visual system. Since visual field localizers are crucial for investigating the contribution of

the visual system to higher-order cognitive processes, our results should have implications

for the study of visual cognition in both for basic and translational research.

RESULTS

<u>Impact of CBA on spatial consistency of ROIs (group level)</u>

Group-level mapping of the four visual quadrants revealed considerable differences before

and after CBA. For the right lower visual quadrant, our analysis before CBA yielded a large

cluster in the left upper occipital cortex, also encompassing posterior parts of temporal cor-

tex, which decreased considerably in size after CBA and was much more circumscribed.

Conversely, for the other three visual quadrants, our analysis before CBA yielded clusters in

the corresponding parts of the occipital pole, which increased in size after CBA without a no-

table change of their center of gravity (Figure 2, Table 1). Overall, after CBA group ROIs

showed markedly greater spatial consistency and vertical symmetry. Additionally, after CBA,

ROIs for the lower visual hemifield were considerably larger in size than ROIs for the upper

visual hemifield. Before and after CBA, average time courses showed clear position selectiv-

ity which increased after CBA (Figure 2).

<u>Probability maps (group level)</u>

Before CBA, probability maps showed a relatively wide spread of functional activation around

the core region of interest defined in our previous group analysis (Figure 3). Here, the maxi-

mum probability of overlap of around 50 % was consistently located at the center of each

ROI. After CBA, probability maps showed a noticeable decrease in the spread of functional

activation around the core region of interest with a corresponding increase in the maximum

probability of overlap. Consequently, probability difference maps (Figure 4) showed an in-

crease in the probability of overlap of up to 40 % around the core region of interest and a cor-

responding peripheral decrease in the probability of overlap of up to 40 %.

<u>Comparison of full ROIs and extended peak-vertex ROIs (single-subject level)</u>

The comparison of time courses averaged over all full single subject ROIs with time courses

averaged over all single subject extended peak-vertex ROIs showed consistently larger

BOLD signal amplitudes for extended peak-vertex ROIs (Figure 5). There was no difference

in BOLD signal amplitude before and after CBA for both types of ROI. Before and after CBA,

all ROIs showed comparable clear position selectivity.

The rate of success for finding a ROI for each subject was as follows: upper right visual

quadrant 46 out of 50 subjects, upper left visual quadrant 50 out of 50 subjects, lower left vis-

ual quadrant 48 out of 50 subjects, lower right visual quadrant 50 out of 50 subjects.

<u>Impact of CBA on spatial variability of peak vertex distribution (single-subject level)</u>

As with the probability maps at the group level, peak vertex distribution maps at the single-

subject level (Figure 6) showed less spatial variability after CBA with an increased probability

of overlap. Both before and after CBA, the spread of peak vertices was larger for the lower

visual quadrants.

DISCUSSION

The aim of our study was to evaluate the utility of CBA for an attention-enhanced visual field

localizer paradigm used to map a circumscribed region within retinotopically organized visual

areas. Our paradigm mapped homologous regions in each visual quadrant reliably across

early visual areas. As expected, CBA led to a marked reduction in macroanatomical variabil-

ity. On the functional level, CBA had a number of beneficial effects.

CBA improved the results of the group ROI analysis for all visual quadrants (Figure 2). This

was reflected in both the spatial consistency of the resulting group ROIs and in the signal-to-

noise ratio exemplified by the difference between the BOLD signal amplitude for the stimu-

lated location compared to the other locations. Probability difference maps showed an in-

crease in the probability of overlap of up to forty percent in the central region of interest,

which resulted in considerably more focused activation patterns. However, the opposite ef-

fect was observed in more peripheral vertices (Figure 4). The latter effect is most likely not

attributable to a decrease of spatial overlap in the periphery of early visual areas. Rather it

demonstrates that CBA consistently reduces spurious spread-out activation resulting from

poor macroanatomical correspondence after VBA. This observation suggests that volume-

based group analyses generally overestimate the extent of visual areas. Together these find-

ings indicate that CBA substantially increases statistical power when studying early visual ar-

eas at the group level. Naturally, this effect of CBA also extends to studies with a more global

focus, such as connectivity analyses[53,54].

Regarding the comparison between full single-subject ROIs and extended peak-vertex sin-

gle-subject ROIs, we made the following observations. As expected, BOLD signal amplitudes

for single subject ROIs were not affected by CBA (Figure 5). Conversely, the variability of

these ROIs as signified by the peak vertex location decreased after CBA (Figure 6). This un-

derscores that the reduction of functional inter-subject variability due to CBA is the main rea-

son for our improved results at the group level. Furthermore, BOLD signal amplitudes for the

full single-subject ROIs were consistently lower compared to our extended single-subject

peak vertex ROIs (Figure 5). This is well in line with previous findings in volume space of

larger effect sizes for ROI definition strategies using peak-voxels with or without including di-

rectly neighboring voxels[48].

Before CBA we observed the strongest group effects, i.e. the largest group ROIs, for the left

upper and right lower visual quadrant, an effect that clearly did not persist after CBA (Figure

2). Notably, several studies using volume-based fMRI and magnetoencephalographic anal-

yses reported lateralized effects on neurophysiological parameters in early visual areas [55,56].

Our observation raises the question, whether these findings could at least partly be explained

by lateralized differences in macro-anatomical variability rather than true functional differ-

ences.

Conversely, our CBA-aided group analysis allowed us to compare the response properties of

each visual quadrant in a more unbiased way. We observed larger group ROIs for the lower

visual hemifield. In a CBA-based probabilistic atlas of the visual system, which included all

regions that could be defined in more than 50% of subjects, probabilistic ROIs for dorsal V1

and V2 were also noticeably larger than probability maps for ventral V1 and V2, whereas this

effect was less clear for V3[41]. These results are in line with our own findings and could be at-

tributable to higher residual anatomical variability after CBA in ventral occipital cortex repre-

senting the upper visual hemifield. Alternatively, they could be due to true differences in re-

sponse properties such as receptive field size or overall area size. The latter interpretation is

supported by studies showing functional differences between upper and lower visual hemi-

fields already at the retinal level in the form of differences in receptor densities. Cone density

was higher in the superior parts of the retina which processes information from lower visual

fields. Conversely, higher rod density was observed in the inferior parts[57,58]. Moreover, Eick-

hoff et al. observed dorso-ventral asymmetries in receptor densities in V2 and V3[57]. They

found a higher density of GABA-A, benzodiazepine- as well as muscarinic M3-receptors in

ventral parts of V2 and V3. Furthermore, there is evidence for fundamental differences in re-

ceptive field shape from a pRF mapping study. Estimating both the aspect ratios and the size

of the mapped areas, a more elliptical receptive field shape was observed for the upper vis-

ual hemifield represented by ventral parts of the visual cortex being compared to the lower

visual hemifield represented by dorsal parts of the visual cortex[59]. Additionally, there is evi-

dence for a behavioral advantage in the lower visual hemifield for shape discrimination as

well as higher BOLD-signal changes and peak amplitudes of MEG/EEG responses[49,51,52,60,61].

Together, these findings demonstrate clear differences in the functional architecture of early

visual areas representing the upper and lower visual hemifield. This has been attributed to

the fact that the lower visual hemifield represented by dorsal parts of the occipital lobe is

more closely linked to the dorsal visual pathway, while the upper visual hemifield represented

by ventral parts of the occipital lobe is more closely linked to the ventral visual pathway[62,63].

Furthermore, there is evidence for fundamental differences in receptive field shape from a

pRF mapping study. For the upper visual hemifield represented by ventral parts of the visual

cortex an increased size and more elliptical shape of receptive fields was observed com-

pared to the lower visual hemifield represented by dorsal parts of the visual cortex[59]. This im-

plies, that the lower visual field is more specialized for the precise localization and represen-

tation of space. Our observation of larger ROIs in the lower visual hemifield is in line with

these findings. Thus, our results imply that CBA is a useful tool to extend the study of func-

tional and behavioral asymmetries in early visual areas to the group level.

One important limitation of the current study is the lack of a complementary retinotopic map-

ping data set due to time constraints. This data would have allowed us to delineate the

boundaries of early visual areas and pinpoint the exact visual area containing each individual

single-subject ROI. Retinotopic mapping studies indicate that peak activation of single sub-

jects assessed by visual localizers are not consistently located in the same visual area. Most

localizer tasks show peak activation not in V1 but rather in V2 or V3[12]. It is therefore very

likely that our single-subject peak activation did not consistently belong to the same visual

cortical area. With the current data set we cannot determine how precisely individual visual

areas were aligned with CBA, and whether individual levels of the visual cortical hierarchy

were differentially affected. However, the position of our group regions of interest, which bor-

dered the calcarine sulcus and spanned the occipital pole, indicate that they mainly com-

prised V2 and V3. Similarly, after CBA a comparable increase in the probability of overlap

was observed in the same part of occipital cortex. While this is at least suggestive of a rela-

tively consistent benefit of CBA across visual areas, more fine-grained studies including reti-

notopic mapping are required to definitively address this question.

Furthermore, we did not use eye tracking to insure sufficient fixation. We also did not include

an additional central attentional control task on the fixation cross, which would have further

encouraged continuous fixation. This was done deliberately in order to keep the difficulty

level adequate for psychiatric patient populations, but it might explain our failure to find relia-

ble activation in early visual areas in a fraction of our subjects.

Our study also has implications beyond mapping the visual system in healthy populations.

Visual processing deficits are a prominent feature of neurodevelopmental psychiatric disor-

ders such as ADHD, schizophrenia and autism spectrum disorders[7,8,64-71], which can also

perturb crucial higher order cognitive processes including working memory[72-74]. The current

localizer paradigm will be useful to investigate local impairments of visual information pro-

cessing as well as disturbances in the interplay between early visual areas and brain net-

works supporting higher-order cognitive processes. Here, CBA will be particularly relevant to

reduce the confounding effects of increased macroanatomical variability in disorders such as

schizophrenia in order to detect true group differences and true functional variability[35,75]. On

the other hand, CBA should also be crucial for investigating the neurodevelopmental under-

pinnings of increased macroanatomical variability itself. To this end the inclusion of probabil-

istic atlases containing information about gene expression profiles[76] as well as cyto- and re-

ceptor architectonics[77,78] will be valuable.

Our CBA approach relied solely on cortical curvature information to reduce macroanatomical

variability. The advantage of this method is its feasibility for the vast majority of fMRI data

sets, since it only requires a structural brain scan of sufficient quality. Among comparable

methods it is the most data driven and objective approach. However, the achievable reduc-

tion of macroanatomical variability is limited by the variable and imperfect correlation be-

tween brain structure and brain function[34,37]. Therefore, more advanced methods additionally

utilize orthogonal functional data to further reduce anatomical variability. This includes the

use of activation patterns or functional connectivity patterns to improve macroanatomical

alignment across the whole brain[20,79,80]. Furthermore, a more complex approach has been

proposed, which aligns cortical data using 'areal features' more closely tied to cortical areas

than cortical folding patterns, including maps of relative myelin content and functional resting

state networks[81]. These methods have shown to provide a relevant additional reduction of

macroanatomical variability for a variety of paradigms including visual functional localizers.

Future studies should also evaluate these methods for retinotopic mapping and visual field

localizers. Moreover, it has been demonstrated for early auditory areas, that the additional

use of a probabilistic atlas of cytoarchitectonically defined areas further improves standard

CBA results[82]. Such an approach would easily be feasible for the visual system.

To summarize, we demonstrated the clear superiority of CBA compared to VBA for the anal-

ysis of visual field localizer data on the group level indicated by a forty percent increase of

overlap of ROIs. Our findings extend previous studies examining other major categories of

visual mapping techniques. They underscore the comparable loss of information and statisti-

cal power incurred by the use of VBA methods in the majority of fMRI studies. Therefore,

CBA and other comparable methods should be seriously considered as a standard proce-

dure for the detailed study of visual information processing and its disturbance in mental dis-

orders.

METHODS AND MATERIALS

Participants:

All participants gave their written consent to participate in the study. The ethical review board

of the Faculty of Medicine at Goethe University granted ethic approval. We recruited 51

healthy volunteers (male : female = 23 : 28) with age ranging between 18-43 years (mean =

24). All participants were non-smokers, had no history of neurological or psychiatric illness

and reported normal or corrected-to-normal visual acuity. One participant was left-handed as

assessed by the German version of the Edinburgh Handedness Inventory[83].

Stimuli and task:

Subjects performed a visual field localizer paradigm (Figure 1) implemented using Presenta-

tion® (Neurobehavioral Systems, Version 18.0) as part of a larger study investigating the role

of visual areas for higher cognitive functions. The task consisted of a series of flickering

black-and-white-colored round shaped chequerboard stimuli (flicker frequency = 7.5 Hz).

Chequerboard stimuli appeared for 2000 ms randomly at one of four different locations (non-

target trial). Each location mapped a homologous position in one of the four visual quadrants.

Our paradigm featured a simple target-detection task. During thirty-six trials, the two centrally

located squares of the chequerboard changed their color to yellow for 133 ms (target trial).

Participants were instructed to press a response box button with their left thumb as quickly

as possible if they detected a target. The task consisted of a total of 144 trials (36 target tri-

als, 108 standard trials). Thus, target probability was 25 %. The regular inter-stimulus interval

was 0 ms. However, once every 10 to 14 trials the inter-stimulus interval increased to 2000

ms. Throughout the task a black, x-shaped fixation cross was displayed at the center of the

screen. Participants were instructed to continuously fixate on the fixation cross. Before the

first trial, only the fixation cross was displayed for 10 seconds. After the last trial, only the fix-

ation cross was displayed for 20 seconds. The total duration of the paradigm was 340 sec-

onds. For the purpose of our analyses we defined a total of four conditions, one for each of

the four stimulus locations. Each participant practiced the task prior to the measurement.

<u>Acquisition and analysis of fMRI data:</u>

We acquired functional MRI data on a Siemens 3T MAGNETOM Trio scanner at the Goethe

University Brain Imaging Centre using a gradient-echo 2D EPI sequence (32 axial slices, TR

= 2000 ms, TE = 30 ms, FA = 90°, FoV = 192 x 192 mm2, voxel size = 3 x 3 x 3 mm^3, gap =

1 mm, effective slice thickness = 4 mm). Slices were positioned parallel to the anterior- and

posterior commissure. Functional images were acquired in a single run comprising the acqui-

sition of 170 volumes. Immediately before each functional run, 6 volumes of this 2D EPI se-

quence were acquired with identical parameters except for a switch of phase encoding direc-

tion (posterior to anterior instead of anterior to posterior) for EPI distortion correction. Ana-

tomical MRI data for cortex reconstruction and co-registration with functional MRI data was

acquired with a high-resolution T1-weighted 3D volume using a Magnetization-Prepared

Rapid Gradient-Echo (MP-RAGE) sequence (192 sagittal slices, TR = 1900 ms, TE = 3.04

ms, TI 900 ms, FA = 9°, FoV = 256 x 256 mm^2, voxel size = 1 x 1 x 1 mm^3). Stimulus presen-

tation was constantly synchronized with the fMRI sequence. Head motion was minimized

with pillows. The task was projected by a beamer onto a mirror attached on the head coil.

MRI data were pre-processed and analyzed using Brain Voyager 20.6[84], the NeuroElf Matlab

toolbox (www.neuroelf.net) and custom software written in Matlab. One subject had to be ex-

cluded due to excessive intra-scan motion.

<u>Structural image pre-processing:</u>

Structural data pre-processing included background cleaning, brain extraction and bias field

correction to minimize image intensity inhomogeneities[84]. Bias field correction employed a

"surface fitting" approach using singular value decomposition based least squares low-order

(Legendre) polynomials to model low-frequency variations across 3D image space[85]. We

used polynomials with an order of 3, which were fitted to a subset of voxels labeled as be-

longing to white matter. The estimated parameters of the polynomials were used to construct

a bias field which was removed from the data. Our approach comprised of one iteration using

automatic white matter labeling[86] and four iterations using manual white matter labeling.

Subsequently, structural data were transformed into Talairach coordinate space[29]. This com-

prised manual labeling the anterior commissure (AC) and posterior commissure (PC) as well

as the borders of the cerebrum. These landmarks were then used to rotate each brain in the

AC–PC plane followed by piece-wise, linear transformations to fit each brain in the common

Talairach "proportional grid" system[19]. Transformation into Talairach coordinate space was

performed because the subsequent automatic segmentation procedure exploits anatomical

knowledge for initial brain segmentation including removing subcortical structures and dis-

connection of cortical hemispheres. To prepare the data for this procedure, we performed a

manual filling of the lateral ventricles. Based on the automatic segmentation of the structural

scans along the white–grey matter boundary[87], cortical hemispheres were reconstructed into

folded, topologically correct mesh representations, which were tessellated to produce surface

reconstructions and calculate curvature maps reflecting the individual cortical folding pat-

terns. Surface reconstructions were subsequently morphed into distortion corrected spherical

representations.

<u>Cortex-based alignment of structural data:</u>

We then applied for each hemisphere separately a high-resolution, multiscale cortex-based

alignment procedure based on the individual curvature maps of all 50 participants. This CBA

approach, which reliably aligns corresponding gyri and sulci across subjects[84], consists of an

initial rigid and a subsequent non-rigid alignment step[19]. Non-rigid CBA employs a coarse-to-

fine matching strategy. Initially, curvature maps contain only smoothed versions of major an-

atomical landmarks, adding increasingly more anatomical detail until all curvature information

from each hemisphere is included. It aligns each cortical folding pattern to a dynamically up-

dated group average through iterative morphing. This moving target approach, which gener-

ates the target curvature map from the average curvature across all hemispheres at a given

alignment stage avoids the possible confounding effects of a suboptimal selection of an indi-

vidual target brain, whose folding pattern might deviate considerably from the cohort aver-

age.

We first conducted a preliminary CBA encompassing both steps to generate an aligned aver-

age brain, which we used for a second, final CBA. During the initial step, we aligned the corti-

cal folding pattern of each sphere rigidly to the cortical folding pattern of a randomly selected

target sphere by global rotation. We used the rotation parameters with the highest degree of

overlap between the curvature of each individual sphere and the target sphere as the starting

point for the subsequent preliminary non-rigid CBA. The results of this preliminary non-rigid

CBA procedure were only used to generate an average brain to be used as an unbiased tar-

get for the final rigid alignment step. This procedure allowed us to minimize the influence of a

suboptimal target brain during the rigid alignment step. The resulting optimal rotation param-

eters were used as the starting point for the subsequent final non-rigid CBA. After the final

non-rigid CBA, we merged both hemispheres of each individual brain to create a global sur-

face-based analysis space.

Furthermore, we created average hemispheres from the original folded mesh representa-

tions, which were subsequently merged, inflated and used for data analysis and visualization.

Averaging of hemispheres was repeated after applying the transformation matrix of the final

rigid and non-rigid CBA to the folded mesh representations, resulting in an accurate repre-

sentation of the effects of macroanatomical alignment.

Functional image pre-processing:

Initial volume-based pre-processing of functional MRI data comprised slice timing correction

using sinc interpolation and 3D motion correction using sinc interpolation. Next, we per-

formed echo-planar imaging distortion correction using the Correction based on Opposite

Phase Encoding (COPE) method[88]. The EPI distortion corrected functional data were co-reg-

istered to the untransformed extracted brains utilizing a boundary-based registration algo-

rithm optimized for the use in surface-based analyses[89]. After co-registration to volume

space, functional data were transformed into Talairach coordinate space using the transfor-

mation matrix generated during Talairach transformation of the anatomical data. The volu-

metric functional data were then transformed into surface space by sampling on the individ-

ual cortical surface incorporating data from −1 to +3 mm along vertex normals. Subsequent

pre-processing of functional MRI data in surface space comprised spatial smoothing using a

nearest neighbor interpolation (1 iteration, approximating a 2D Gaussian smoothing kernel

with a FWHM of 1 mm), linear trend removal and temporal high-pass filtering (high-pass

0.00903 Hz). Spatial smoothing in surface space is clearly superior to spatial smoothing in

volume space by avoiding contamination from cortical areas proximal in volume space but

more distant in surface space[19,53]. However, we still opted for minimum spatial smoothing to

prevent a loss of accuracy for our visual field localizer. Based on the vertex-to-vertex refer-

encing from the folded, topologically correct meshes to the spherical representations, we

mapped the fully preprocessed functional data into a common spherical coordinate system.

Finally, we applied surface-based anatomical masks that only included cortical vertices in our

analysis to the functional data. These masks excluded those subcortical structures, i.e. parts

of thalamus and the basal ganglia, which mapped onto the midline of our surface reconstruc-

tions.

<u>fMRI group analysis:</u>

We performed multi-subject statistical analyses using multiple linear regression of the BOLD

signal. The presentation of each chequerboard stimulus sequence at a single location, was

modelled by an ideal box-car function, which covered the volume of each trial, convolved

with a synthetic two-gamma function. These predictors were used to build the design matrix

of the experiment. Individual statistical maps were generated by associating each voxel with

the beta-value corresponding to the specific set of predictors and calculated on the basis of

the least mean squares solution of the general linear model. The resulting individual statisti-

cal maps were entered into a second-level random-effects group analysis using a summary

statistic approach.

To study the effects of surface-based macroanatomical alignment, we computed the same

analysis steps in surface space before and after CBA, following the protocol established by

previous works evaluating this particular method[19,90]. Thus, we essentially compared the im-

pact of pure surface-based registration, referred to as "pre-CBA", with explicit surface-based

macroanatomical alignment, referred to as "post-CBA". However, for the purpose of our

study, we consider the former (pre-CBA) to be a good proxy for VBA, while allowing to di-

rectly contrast its results with those of our post-CBA analysis within the same coordinate sys-

tem.

<u>Group analysis of visual quadrants before and after CBA:</u>

We performed analyses focusing on the mapping of the four visual quadrants at the group

level and compared these with results at single subject level. To define the group level ROI

for each visual quadrant, we computed separate weighted contrasts for each quadrant

against the other three quadrants. We assigned a weight of three to the position of interest,

e.g. $(\beta^{Quad1} \times 3) / (\beta^{Quad2} + \beta^{Quad3} + \beta^{Quad4})$ ($p < 0.5$, Bonferroni corrected). For each resulting

group level ROI, we extracted average time courses (incl. standard errors of the mean =

SEM) for all four conditions. We conducted this analysis both before and after CBA.

<u>Probability maps before and after CBA:</u>

To quantify and visualize variability of functional activation and possible changes due to

macroanatomical alignment, the use of probability maps has been proposed. Probability

maps are specifically useful to quantify inconsistency i.e. the disparity between individuals

regarding the location of a particular (visual) area. To quantify the spatial consistency of acti-

vation patterns before and after CBA, we generated probability maps for each visual quad-

rant [91,92]. These maps represent the relative number of subjects showing significant task ac-

tivity in our single-subject analysis. Probability maps for early visual areas before and after

CBA were created based on the previously generated single subject t-maps, with a threshold

of a minimum of ten percent probability. To quantify the change in spatial consistency of acti-

vation patterns as a result of CBA, we created probability difference maps for each position

(post-CBA minus pre-CBA). This allowed us to determine both increases and decreases of

spatial consistency due to CBA.

Single-subject ROIs of visual quadrants before and after CBA including peak vertex map-

ping:

For analyses at single-subject level, we first defined ROIs for each subject independently us-

ing the same weighted contrasts employed in the group analysis. We applied a more lenient

statistical threshold ($p < 0.05$ uncorrected). For each resulting single-subject ROI we ex-

tracted average time courses for all four conditions. We then averaged these time courses

across all subjects. Thus, we generated group level average time courses reflecting each

subject's region of maximum activation. We conducted this analysis both before and after

CBA. Additionally, we determined the peak vertex for each subject's four visual quadrant

ROIs, i.e. the vertex with the highest t-value. We then defined extended single-subject peak

vertex ROIs, which also included the six vertices adjacent to the peak vertex [48]. For each re-

sulting single-subject peak vertex ROI we extracted average time courses (incl. SEM) for all

four conditions. We then averaged these timecourses across all subjects. We conducted this

analysis both before and after CBA.

Peak vertex distribution maps:

We mapped all peak vertices per visual quadrant before and after CBA and computed the

vertex-wise number of peak vertices to estimate the degree of overlap between subjects.

REFERENCES

541
542   1   Wandell, B. A., Dumoulin, S. O. & Brewer, A. A. Visual field maps in human cortex. *Neuron* **56**,
543   366-383, doi:10.1016/j.neuron.2007.10.012 (2007).
544   2   Corbetta, M. & Shulman, G. L. Control of goal-directed and stimulus-driven attention in the
545   brain. *Nature Reviews Neuroscience* **3**, 201-215, doi:10.1038/nrn755 (2002).
546   3   Das, M., Bennett, D. M. & Dutton, G. N. Visual attention as an important visual function: an
547   outline of manifestations, diagnosis and management of impaired visual attention. *Br J*
548   *Ophthalmol* **91**, 1556-1560, doi:10.1136/bjo.2006.104844 (2007).
549   4   de Haan, B., Bither, M., Brauer, A. & Karnath, H. O. Neural Correlates of Spatial Attention and
550   Target Detection in a Multi-Target Environment. *Cereb Cortex* **25**, 2321-2331,
551   doi:10.1093/cercor/bhu046 (2015).
552   5   Goodale, M. & Milner, D. One brain - Two visual systems. *Psychologist* **19**, 660-663 (2006).
553   6   Bergmann, J., Genç, E., Kohler, A., Singer, W. & Pearson, J. Neural Anatomy of Primary Visual
554   Cortex Limits Visual Working Memory. *Cerebral Cortex* **26**, 43-50, doi:10.1093/cercor/bhu168
555   (2014).
556   7   Lee, J. *et al.* fMRI evidence of aberrant neural adaptation for objects in schizophrenia and
557   bipolar disorder. *Hum Brain Mapp* **40**, 1608-1617, doi:10.1002/hbm.24472 (2019).
558   8   Silverstein, S. M. *et al.* An fMRI examination of visual integration in schizophrenia. *J Integr*
559   *Neurosci* **8**, 175-202, doi:10.1142/s0219635209002113 (2009).
560   9   Dumoulin, S. O. & Wandell, B. A. Population receptive field estimates in human visual cortex.
561   *Neuroimage* **39**, 647-660, doi:10.1016/j.neuroimage.2007.09.034 (2008).
562   10   Sereno, M. I. *et al.* Borders of multiple visual areas in humans revealed by functional
563   magnetic resonance imaging. *Science (New York, N.Y.)* **268**, 889-893,
564   doi:10.1126/science.7754376 (1995).
565   11   Harrison, S. A. & Tong, F. Decoding reveals the contents of visual working memory in early
566   visual areas. *Nature* **458**, 632-635, doi:10.1038/nature07832 (2009).
567   12   Peters, B., Kaiser, J., Rahm, B. & Bledowski, C. Activity in Human Visual and Parietal Cortex
568   Reveals Object-Based Attention in Working Memory. *The Journal of Neuroscience* **35**, 3360-
569   3369, doi:10.1523/jneurosci.3795-14.2015 (2015).
570   13   Downing, P. E., Chan, A. W., Peelen, M. V., Dodds, C. M. & Kanwisher, N. Domain specificity in
571   visual cortex. *Cereb Cortex* **16**, 1453-1461, doi:10.1093/cercor/bhj086 (2006).
572   14   Kanwisher, N., McDermott, J. & Chun, M. M. The fusiform face area: a module in human
573   extrastriate cortex specialized for face perception. *J Neurosci* **17**, 4302-4311,
574   doi:10.1523/jneurosci.17-11-04302.1997 (1997).

575   15   Brett, M., Johnsrude, I. S. & Owen, A. M. The problem of functional localization in the human
576   brain. *Nat Rev Neurosci* **3**, 243-249, doi:10.1038/nrn756 (2002).
577   16   Desai, R., Liebenthal, E., Possing, E. T., Waldron, E. & Binder, J. R. Volumetric vs. surface-
578   based alignment for localization of auditory cortex activation. *Neuroimage* **26**, 1019-1029,
579   doi:10.1016/j.neuroimage.2005.03.024 (2005).
580   17   Dougherty, R. F. *et al.* Visual field representations and locations of visual areas V1/2/3 in
581   human visual cortex. *J Vis* **3**, 586-598, doi:10.1167/3.10.1 (2003).
582   18   Fischl, B., Sereno, M. I. & Dale, A. M. Cortical surface-based analysis. II: Inflation, flattening,
583   and a surface-based coordinate system. *Neuroimage* **9**, 195-207,
584   doi:10.1006/nimg.1998.0396 (1999).
585   19   Frost & Goebel. Measuring structural-functional correspondence: spatial variability of
586   specialised brain regions after macro-anatomical alignment. *Neuroimage* **59**, 1369-1381,
587   doi:10.1016/j.neuroimage.2011.08.035 (2012).
588   20   Frost & Goebel. Functionally informed cortex based alignment: an integrated approach for
589   whole-cortex macro-anatomical and ROI-based functional alignment. *NeuroImage* **83**, 1002-
590   1010, doi:10.1016/j.neuroimage.2013.07.056 (2013).
591   21   Steinmetz, H., Fürst, G. & Freund, H. J. Variation of perisylvian and calcarine anatomic
592   landmarks within stereotaxic proportional coordinates. *AJNR Am J Neuroradiol* **11**, 1123-
593   1130 (1990).
594   22   Van Essen, D. C. *et al.* An integrated software suite for surface-based analyses of cerebral
595   cortex. *J Am Med Inform Assoc* **8**, 443-459, doi:10.1136/jamia.2001.0080443 (2001).
596   23   Zilles, K. *et al.* Quantitative analysis of sulci in the human cerebral cortex: Development,
597   regional heterogeneity, gender difference, asymmetry, intersubject variability and cortical
598   architecture. *Human Brain Mapping* **5**, 218-221, doi:https://doi.org/10.1002/(SICI)1097-
599   0193(1997)5:4<218::AID-HBM2>3.0.CO;2-6 (1997).
600   24   Bridge, H. Mapping the visual brain: how and why. *Eye (Lond)* **25**, 291-296,
601   doi:10.1038/eye.2010.166 (2011).
602   25   Fedorenko, E., Hsieh, P. J., Nieto-Castañón, A., Whitfield-Gabrieli, S. & Kanwisher, N. New
603   method for fMRI investigations of language: defining ROIs functionally in individual subjects.
604   *J Neurophysiol* **104**, 1177-1194, doi:10.1152/jn.00032.2010 (2010).
605   26   Julian, J. B., Fedorenko, E., Webster, J. & Kanwisher, N. An algorithmic method for
606   functionally defining regions of interest in the ventral visual pathway. *NeuroImage* **60**, 2357-
607   2364, doi:10.1016/j.neuroimage.2012.02.055 (2012).
608   27   Nieto-Castañón, A. & Fedorenko, E. Subject-specific functional localizers increase sensitivity
609   and functional resolution of multi-subject analyses. *Neuroimage* **63**, 1646-1669,
610   doi:10.1016/j.neuroimage.2012.06.065 (2012).
611   28   Smith, S. M. *et al.* Functional connectomics from resting-state fMRI. *Trends in cognitive
612   sciences* **17**, 666-682, doi:10.1016/j.tics.2013.09.016 (2013).
613   29   Talairach, J., & Tournoux, P. *Co-planar stereotaxic atlas of the human brain: 3-dimensional
614   proportional system : an approach to cerebral imaging.* (Thieme, 1988).
615   30   Evans, A. C. *et al.* 3D statistical neuroanatomical models from 305 MRI volumes. *1993 IEEE
616   Conference Record Nuclear Science Symposium and Medical Imaging Conference*, 1813-1817
617   vol.1813 (1993).
618   31   Klein, A. *et al.* Evaluation of volume-based and surface-based brain image registration
619   methods. *Neuroimage* **51**, 214-220, doi:10.1016/j.neuroimage.2010.01.091 (2010).
620   32   Ghosh, S. S. *et al.* Evaluating the validity of volume-based and surface-based brain image
621   registration for developmental cognitive neuroscience studies in children 4 to 11 years of
622   age. *NeuroImage* **53**, 85-93, doi:10.1016/j.neuroimage.2010.05.075 (2010).
623   33   Nieto-Castanon, A., Ghosh, S. S., Tourville, J. A. & Guenther, F. H. Region of interest based
624   analysis of functional imaging data. *Neuroimage* **19**, 1303-1316, doi:10.1016/s1053-
625   8119(03)00188-5 (2003).

34   Fischl, B., Sereno, M. I., Tootell, R. B. & Dale, A. M. High-resolution intersubject averaging and
a coordinate system for the cortical surface. *Hum Brain Mapp* **8**, 272-284,
doi:10.1002/(sici)1097-0193(1999)8:4<272::aid-hbm10>3.0.co;2-4 (1999).
35   Anticevic, A. *et al.* Comparing surface-based and volume-based analyses of functional
neuroimaging data in patients with schizophrenia. *Neuroimage* **41**, 835-848,
doi:10.1016/j.neuroimage.2008.02.052 (2008).
36   Pantazis, D. *et al.* Comparison of landmark-based and automatic methods for cortical surface
registration. *Neuroimage* **49**, 2479-2493, doi:10.1016/j.neuroimage.2009.09.027 (2010).
37   Van Essen, D. C. & Drury, H. A. Structural and functional analyses of human cerebral cortex
using a surface-based atlas. *J Neurosci* **17**, 7079-7102, doi:10.1523/jneurosci.17-18-
07079.1997 (1997).
38   Fischl, B. *et al.* Cortical folding patterns and predicting cytoarchitecture. *Cereb Cortex* **18**,
1973-1980, doi:10.1093/cercor/bhm225 (2008).
39   Hinds, O. P. *et al.* Accurate prediction of V1 location from cortical folds in a surface
coordinate system. *NeuroImage* **39**, 1585-1599, doi:10.1016/j.neuroimage.2007.10.033
(2008).
40   Weiner, K. S. *et al.* Defining the most probable location of the parahippocampal place area
using cortex-based alignment and cross-validation. *Neuroimage* **170**, 373-384,
doi:10.1016/j.neuroimage.2017.04.040 (2018).
41   Rosenke, M., van Hoof, R., van den Hurk, J., Grill-Spector, K. & Goebel, R. A Probabilistic
Functional Atlas of Human Occipito-Temporal Visual Cortex. *Cerebral Cortex*,
doi:10.1093/cercor/bhaa246 (2020).
42   Huang, T., Chen, X., Jiang, J., Zhen, Z. & Liu, J. A probabilistic atlas of the human motion
complex built from large-scale functional localizer data. *Hum Brain Mapp* **40**, 3475-3487,
doi:10.1002/hbm.24610 (2019).
43   Di Russo, F., Martínez, A. & Hillyard, S. A. Source analysis of event-related cortical activity
during visuo-spatial attention. *Cereb Cortex* **13**, 486-499, doi:10.1093/cercor/13.5.486
(2003).
44   Di Russo, F., Martínez, A., Sereno, M. I., Pitzalis, S. & Hillyard, S. A. Cortical sources of the
early components of the visual evoked potential. *Hum Brain Mapp* **15**, 95-111,
doi:10.1002/hbm.10010 (2002).
45   Shigihara, Y., Hoshi, H. & Zeki, S. Early visual cortical responses produced by checkerboard
pattern stimulation. *NeuroImage* **134**, doi:10.1016/j.neuroimage.2016.03.078 (2016).
46   Kraft, A. *et al.* fMRI localizer technique: efficient acquisition and functional properties of
single retinotopic positions in the human visual cortex. *NeuroImage* **28**, 453-463,
doi:10.1016/j.neuroimage.2005.05.050 (2005).
47   Bressler, D. W. & Silver, M. A. Spatial attention improves reliability of fMRI retinotopic
mapping signals in occipital and parietal cortex. *Neuroimage* **53**, 526-533,
doi:10.1016/j.neuroimage.2010.06.063 (2010).
48   Tong, Y. *et al.* Seeking Optimal Region-Of-Interest (ROI) Single-Value Summary Measures for
fMRI Studies in Imaging Genetics. *PLoS One* **11**, e0151391,
doi:10.1371/journal.pone.0151391 (2016).
49   Anderson, J., Cameron, E. & Levine, M. A method for quantifying visual field
inhomogeneities. *Vision Research* **105**, doi:10.1016/j.visres.2014.09.010 (2014).
50   Liu, T., Heeger, D. J. & Carrasco, M. Neural correlates of the visual vertical meridian
asymmetry. *J Vis* **6**, 1294-1306, doi:10.1167/6.11.12 (2006).
51   O'Connell, C. *et al.* Structural and functional correlates of visual field asymmetry in the
human brain by diffusion kurtosis MRI and functional MRI. *Neuroreport* **27**, 1225-1231,
doi:10.1097/wnr.0000000000000682 (2016).
52   Rubin, N., Nakayama, K. & Shapley, R. Enhanced perception of illusory contours in the lower
versus upper visual hemifields. *Science* **271**, 651-653, doi:10.1126/science.271.5249.651
(1996).

53 Brodoehl, S., Gaser, C., Dahnke, R., Witte, O. W. & Klingner, C. M. Surface-based analysis
increases the specificity of cortical activation patterns and connectivity results. *Scientific*
Reports **10**, 5737, doi:10.1038/s41598-020-62832-z (2020).
54 Bullmore, E. & Sporns, O. Complex brain networks: graph theoretical analysis of structural
and functional systems. *Nature Reviews Neuroscience* **10**, 186-198, doi:10.1038/nrn2575
(2009).
55 Chen, H., Yao, D. & Liu, Z. A study on asymmetry of spatial visual field by analysis of the fMRI
BOLD response. *Brain Topogr* **17**, 39-46, doi:10.1023/b:brat.0000047335.00110.6a (2004).
56 Loughnane, G. M., Shanley, J. P., Lalor, E. C. & O'Connell, R. G. Behavioral and
electrophysiological evidence of opposing lateral visuospatial asymmetries in the upper and
lower visual fields. *Cortex* **63**, 220-231, doi:10.1016/j.cortex.2014.09.003 (2015).
57 Eickhoff, S. B., Rottschy, C., Kujovic, M., Palomero-Gallagher, N. & Zilles, K. Organizational
Principles of Human Visual Cortex Revealed by Receptor Mapping. *Cerebral Cortex* **18**, 2637-
2645, doi:10.1093/cercor/bhn024 (2008).
58 Curcio, C. A. & Allen, K. A. Topography of ganglion cells in human retina. *J Comp Neurol* **300**,
5-25, doi:10.1002/cne.903000103 (1990).
59 Silson, E. H., Reynolds, R. C., Kravitz, D. J. & Baker, C. I. Differential Sampling of Visual Space
in Ventral and Dorsal Early Visual Cortex. *The Journal of Neuroscience* **38**, 2294-2303,
doi:10.1523/jneurosci.2717-17.2018 (2018).
60 Hagler, D. J., Jr. Visual field asymmetries in visual evoked responses. *J Vis* **14**, 13,
doi:10.1167/14.14.13 (2014).
61 Schmidtmann, G., Logan, A. J., Kennedy, G. J., Gordon, G. E. & Loffler, G. Distinct lower visual
field preference for object shape. *J Vis* **15**, 18, doi:10.1167/15.5.18 (2015).
62 Thomas, N. A. & Elias, L. J. Upper and lower visual field differences in perceptual
asymmetries. *Brain Res* **1387**, 108-115, doi:10.1016/j.brainres.2011.02.063 (2011).
63 Zito, G. A., Cazzoli, D., Müri, R. M., Mosimann, U. P. & Nef, T. Behavioral Differences in the
Upper and Lower Visual Hemifields in Shape and Motion Perception. *Front Behav Neurosci*
10, 128-128, doi:10.3389/fnbeh.2016.00128 (2016).
64 Bakroon, A. & Lakshminarayanan, V. Visual function in autism spectrum disorders: a critical
review. *Clin Exp Optom* **99**, 297-308, doi:10.1111/cxo.12383 (2016).
65 Butler, P. D. *et al.* Dysfunction of early-stage visual processing in schizophrenia. *Am J*
Psychiatry **158**, 1126-1133, doi:10.1176/appi.ajp.158.7.1126 (2001).
66 Butler, P. D., Silverstein, S. M. & Dakin, S. C. Visual perception and its impairment in
schizophrenia. *Biol Psychiatry* **64**, 40-47, doi:10.1016/j.biopsych.2008.03.023 (2008).
67 Chen, C. Y. *et al.* Visual spatial attention in children with attention deficit hyperactivity
disorder. *Chang Gung Med J* **25**, 514-521 (2002).
68 Hale, T. S. *et al.* Visual Network Asymmetry and Default Mode Network Function in ADHD: An
fMRI Study. *Front Psychiatry* **5**, 81, doi:10.3389/fpsyt.2014.00081 (2014).
69 Sanz-Cervera, P., Pastor-Cerezuela, G., González-Sala, F., Tárraga-Mínguez, R. & Fernández-
Andrés, M.-I. Sensory Processing in Children with Autism Spectrum Disorder and/or Attention
Deficit Hyperactivity Disorder in the Home and Classroom Contexts. *Front Psychol* **8**, 1772-
1772, doi:10.3389/fpsyg.2017.01772 (2017).
70 Seymour, R. A., Rippon, G., Gooding-Williams, G., Schoffelen, J. M. & Kessler, K. Dysregulated
oscillatory connectivity in the visual system in autism spectrum disorder. *Brain* **142**, 3294-
3305, doi:10.1093/brain/awz214 (2019).
71 Shimizu, V. T., Bueno, O. F. & Miranda, M. C. Sensory processing abilities of children with
ADHD. *Braz J Phys Ther* **18**, 343-352, doi:10.1590/bjpt-rbf.2014.0043 (2014).
72 Bittner, R. A. *et al.* The When and Where of Working Memory Dysfunction in Early-Onset
Schizophrenia-A Functional Magnetic Resonance Imaging Study. *Cereb Cortex* **25**, 2494-2506,
doi:10.1093/cercor/bhu050 (2015).

73   Butler, P. D., Thompson, J. L., Seitz, A. R., Deveau, J. & Silverstein, S. M. Visual perceptual
remediation for individuals with schizophrenia: Rationale, method, and three case studies.
Psychiatr Rehabil J **40**, 43-52, doi:10.1037/prj0000212 (2017).
74   Haenschel, C. *et al.* Contribution of impaired early-stage visual processing to working
memory dysfunction in adolescents with schizophrenia: a study with event-related potentials
and functional magnetic resonance imaging. *Arch Gen Psychiatry* **64**, 1229-1240,
doi:10.1001/archpsyc.64.11.1229 (2007).
75   Manoach, D. S. Prefrontal cortex dysfunction during working memory performance in
schizophrenia: reconciling discrepant findings. *Schizophr Res* **60**, 285-298,
doi:10.1016/s0920-9964(02)00294-3 (2003).
76   French, L. & Paus, T. A FreeSurfer view of the cortical transcriptome generated from the Allen
Human Brain Atlas. *Frontiers in Neuroscience* **9**, doi:10.3389/fnins.2015.00323 (2015).
77   Gulban, O. F. *et al.* Improving a probabilistic cytoarchitectonic atlas of auditory cortex using a
novel method for inter-individual alignment. *eLife* **9**, e56963, doi:10.7554/eLife.56963
(2020).
78   Hawrylycz, M. J. *et al.* An anatomically comprehensive atlas of the adult human brain
transcriptome. *Nature* **489**, 391-399, doi:10.1038/nature11405 (2012).
79   Sabuncu, M. R. *et al.* Function-based Intersubject Alignment of Human Cortical Anatomy.
Cerebral Cortex **20**, 130-140, doi:10.1093/cercor/bhp085 (2009).
80   Conroy, B. R., Singer, B. D., Guntupalli, J. S., Ramadge, P. J. & Haxby, J. V. Inter-subject
alignment of human cortical anatomy using functional connectivity. *NeuroImage* **81**, 400-
411, doi:https://doi.org/10.1016/j.neuroimage.2013.05.009 (2013).
81   Glasser, M. F. *et al.* A multi-modal parcellation of human cerebral cortex. *Nature* **536**, 171-
178, doi:10.1038/nature18933 (2016).
82   Gulban, O. F. *et al.* Improving a probabilistic cytoarchitectonic atlas of auditory cortex using a
novel method for inter-individual alignment. *Elife* **9**, doi:10.7554/eLife.56963 (2020).
83   Oldfield, R. C. The assessment and analysis of handedness: the Edinburgh inventory.
Neuropsychologia **9**, 97-113, doi:10.1016/0028-3932(71)90067-4 (1971).
84   Goebel, R., Esposito, F. & Formisano, E. Analysis of functional image analysis contest (FIAC)
data with brainvoyager QX: From single-subject to cortically aligned group general linear
model analysis and self-organizing group independent component analysis. *Hum Brain Mapp*
27, 392-401, doi:10.1002/hbm.20249 (2006).
85   Sled, J. G., Zijdenbos, A. P. & Evans, A. C. in *Proceedings of the 15th International Conference
on Information Processing in Medical Imaging*   459–464 (Springer-Verlag, 1997).
86   Hou, Z., Huang, S., Hu, Q. & Nowinski, W. L. A fast and automatic method to correct intensity
inhomogeneity in MR brain images. *Med Image Comput Comput Assist Interv* **9**, 324-331,
doi:10.1007/11866763_40 (2006).
87   Kriegeskorte, N. & Goebel, R. An Efficient Algorithm for Topologically Correct Segmentation
of the Cortical Sheet in Anatomical MR Volumes. *NeuroImage* **14**, 329-346 (2001).
88   Breman, H. *et al.* in *Biomedical Image Registration.* (eds Žiga Špiclin, Jamie McClelland, Jan
Kybic, & Orcun Goksel) 122-130 (Springer International Publishing).
89   Greve, D. N. & Fischl, B. Accurate and robust brain image alignment using boundary-based
registration. *Neuroimage* **48**, 63-72, doi:10.1016/j.neuroimage.2009.06.060 (2009).
90   Frost, M. A., Esposito, F. & Goebel, R. Improved correspondence of resting-state networks
after macroanatomical alignment. *Hum Brain Mapp* **35**, 673-682, doi:10.1002/hbm.22191
(2014).
91   Wang, L., Mruczek, R. E., Arcaro, M. J. & Kastner, S. Probabilistic Maps of Visual Topography
in Human Cortex. *Cereb Cortex* **25**, 3911-3931, doi:10.1093/cercor/bhu277 (2015).
92   Yamamoto, H. *et al.* Inconsistency and uncertainty of the human visual area loci following
surface-based registration: Probability and Entropy Maps. *Hum Brain Mapp* **33**, 121-129,
doi:10.1002/hbm.21200 (2012).
779

ACKNOWLEDGEMENTS

786 The authors are most grateful to Deliah Macht, Christina Raab and Leonie Winkler-Lauble for

787 help with data acquisition. C.V Barnes-Scheufler was supported by a "main doctus" scholar-

788 ship from The Polytechnic Foundation of Frankfurt am Main. E. Raspor was supported by a

789 Research Grant for Doctoral Programmes in Germany from the German Academic Exchange

790 Service (DAAD).

AUTHOR CONTRIBUTION

All authors made substantial contributions to the conception or design of the work, or the ac-

quisition, analysis, or interpretation of data. AR and RAB acquired funding. RAB, BP, MS and

LR designed the experiment. MQ, CVB-S, LR and RAB acquired the data. ER and RG pro-

vided analytical tools. MQ and RAB analyzed the data. MQ and RAB undertook the literature

searches and wrote the first draft of the manuscript. All authors contributed to and revised

the manuscript. All authors read and approved the final manuscript.

DISCLOSURES

801 The authors have declared that there are no conflicts of interest in relation to the subject of

802 this study.

DATA AVAILABILITY STATEMENT

805 The data that support the findings of this study are available from the corresponding author

806 upon reasonable request.

FIGURE LEGENDS

Figure 1 – Visual field Localizer Paradigm

The task consisted of flickering, black-and-white colored checkerboards that appeared ran-
domly at homologous positions of the participant's visual quadrant. Only in 25 % of the trials
the two centrally located squares changed their color into yellow for 133 ms. Participants
were required to press a response box button when noticing that. During the whole task par-
ticipants were instructed to fixate a black, x-shaped fixation cross presented in the center of
the screen. Checkerboards appeared for 2000 ms. The regular inter-stimulus interval was 0
ms. Every 10 to 14 trial the ITI extended to 2000 ms.

Figure 2 – Group analysis of visual quadrants

Surface-based group results (a) before and (b) after CBA (pre-CBA and post-CBA) and aver-
aged time courses. After CBA, group ROIs showed greater spatial consistency and ROIs for
the lower visual hemifield were larger in size than ROIs for the upper visual hemifield. Aver-
age timecourses (incl. SEM) showed clear position selectivity which was higher after CBA.
ROI/graph colors: red = upper left (UL) visual quadrant, orange = lower left (LL) visual quad-
rant, light-blue = lower right (LR) visual quadrant, dark-blue = upper right (UR) visual quad-
rant.

Figure 3 – Probability Maps

Probability Maps (a) before and (b) after CBA (pre-CBA and post-CBA) for each visual quad-
rant. The color code grey-to-white indicates an increase in the probability of overlap. Proba-

bility maps (a) before CBA showed a maximum probability of overlap of up to 50 %. Probabil-

ity maps (b) after CBA showed a maximum probability of overlap of up to 90 %.

Figure 4 – Probability Difference Maps

Probability difference maps (post-CBA minus pre-CBA). The color code dark-red-to-white in-

dicates an increase of activation overlap. The color code blue-to-green indicate a decrease

of activation overlap. Overall, at the center of each ROI the probability of overlap increased

by up to forty percent after CBA.

Figure 5 – Average timecourses of visual quadrants

Average timecourses (incl. SEM) of visual quadrants for full single-subject and single-subject

extended peak vertex ROIs before and after CBA (pre-CBA and post-CBA). Extended peak-

vertex ROIs showed consistently larger BOLD-signal amplitudes. Average timecourses

showed clear position selectivity which was unaffected by CBA.

ROI/graph colors: red = upper left (UL) visual quadrant, orange = lower left (LL) visual quad-

rant, light-blue = lower right (LR) visual quadrant, dark-blue = upper right (UR) visual quad-

rant. Dashed lines = single subject extended peak-vertex ROIs, solid lines = full single-sub-

ject ROIs.

Figure 6 – Peak vertex distribution maps

For each visual quadrant single-subject peak vertices were mapped (a) before and (b) after

CBA (pre-CBA and post-CBA) and a vertex-wise number of peak vertices were computed to

estimate the degree of overlap. The color code turquoise-to-red indicates an increase in the

degree of overlap. After CBA, we observed reduced spatial variability of peak vertex location.

Table 1 – Talairach coordinates

Table 1 depicts the Talairach coordinates of the group ROIs of the corresponding visual

quadrants before and after CBA.

Table 1					
Region of interest		**Number of vertices**	**Talairach coordinates (in mm)**		
			x	**y**	**z**
upper right visual quadrant	pre CBA	6	-18	-80	-15
	post CBA	58	-18	-80	-15
upper left visual quadrant	pre CBA	47	24	-75	-14
	post CBA	82	18	-77	-15
lower left visual quadrant	pre CBA	295	-42	-64	5
	post CBA	161	-24	-91	-4
lower right visual quadrant	pre CBA	28	23	-90	3
	post CBA	127	9	-94	-5

(a) pre-CBA

(b) post-CBA

UR Quadrant   UL Quadrant   LL Quadrant   LR Quadrant

Figure 2

(a)   pre-CBA

UR Quadrant   UL Quadrant   LR Quadrant   LL Quadrant

100%   0%

(b)   post-CBA

UR Quadrant   UL Quadrant   LR Quadrant   LL Quadrant

Figure 3

UR Quadrant   UL Quadrant   LR Quadrant   LL Quadrant

40%   0%   - 40%

Figure 4

(a) pre-CBA

(b) post-CBA

UR Quadrant — UL Quadrant — LL Quadrant — LR Quadrant

- - - Full ROI — Extended Peak Vertex ROI

Figure 5

(a) pre-CBA

UR Quadrant   UL Quadrant   LR Quadrant   LL Quadrant

(b) post-CBA

UR Quadrant   UL Quadrant   LR Quadrant   LL Quadrant

Figure 6