A unicellular walker controlled by a microtubule-based finite state machine

Ben T. Larson[1], Jack Garbus[2], Jordan B. Pollack[2], and Wallace F. Marshall[1]*

Summary

Cells are complex biochemical systems whose behavior emerges from interactions among myriad molecular components. Computation is often invoked as a general framework for navigating this cellular complexity. However, it is unclear how cells might embody computational processes such that theories of computation, including finite state machine models, could be productively applied. Here, we demonstrate finite state machine-like processing embodied in cells using the walking behavior of *Euplotes eurystomus*, a ciliate that walks across surfaces using fourteen motile appendages (cirri). We found that cellular walking entails regulated transitions between a discrete set of gait states. The set of observed transitions decomposes into a small group of high-probability, temporally irreversible transitions and a large group of low-probability time-symmetric transitions, thus revealing stereotypy in sequential patterns of state transitions. Simulations and experiments suggest that the sequential logic of the gait is functionally important. Taken together, these findings implicate a finite state machine-like process. Cirri are connected by microtubule bundles (fibers), and we found that the dynamics of cirri involved in different state transitions are associated with the structure of the fiber system. Perturbative experiments revealed that the fibers mediate gait coordination, suggesting a mechanical basis of gait control.

[1] Department of Biochemistry and Biophysics, University of California, San Francisco, California, USA
[2] Computer Science Department, Brandeis University, Waltham, Massachusetts, USA
* Correspondence: wallace.ucsf@gmail.com

Keywords

cellular behavior, information processing, computation, decision-making, broken detailed balance, gait coordination, cell motility, cilia, ciliate, tubulin

Introduction

Cells are complex physical systems controlled by networks of signaling molecules. Single cells can display remarkably sophisticated, animal-like behaviors [1–3], orchestrating active processes far from thermodynamic equilibrium in order to properly carry out biological functions [4,5]. Indeed, single cells can make decisions by sensing and responding to diverse cues and signals [6], execute coordinated movements [7,8] and directed motility [9–12], and even solve mazes [13,14] and possibly learn [15–18]. Such behaviors in animals arise from neural activity and have been studied extensively, but we know comparatively little about the mechanisms of cellular behavior [19,20]. In individual cells, behaviors emerge directly through the joint action of chemical reactions [21], cellular architecture [3], physical mechanisms and constraints within the cell [22,23], and interactions of the cell with its local environment [24]. The links between information processing, decision making, and the physical manifestation of those processes as cell state transitions suggest that cellular behavior might be understood as an embodied computation [25,26]. The theory of computation has often been invoked as a general framework for understanding cellular dynamics [25,27–32], environmental sensing by bacteria being a deeply studied example [31–33], and has been used to engineer programmable cell states [34]. However, the manner and extent to which cells might embody functional, computational processes as well as the extent to which a computational perspective on cellular behavior might prove productive remains to be seen.

Among the microbial eukaryotes (protists), ciliates display some of the most striking examples of unicellular behavior, including hunting [3], sensorimotor navigation [10], and predator

avoidance [35]. Spirotrichous ciliates of the genus *Euplotes* are notable for their complex

locomotion [36–38], using bundles of specialized cilia (cirri) to walk across surfaces [36,37] (Figure 1A,

Videos S1 and S2). Depending on the species, these cells generally have 14 to 15 ventral cirri

arranged in a highly consistent pattern used for walking locomotion [39]. *Euplotes* live in aquatic

environments, and in addition to walking, use their cirri for swimming and rapid escape

responses [40] (Video S2). Oral membranelles (Figure 1B) generate feeding currents to capture

bacteria and small protistan prey and are also used for swimming. Early 20th century

protistologists were so impressed by the apparent coordination of cirri that they proposed the

existence of a rudimentary nervous system, the neuromotor apparatus, to account for their

observations [38]. This theory was motivated by the presence of intracellular fibers connecting

various cirri (Figure 1C), now known to be to be tubulin-based structures [41,42].

How can a single cell coordinate a gait without a nervous system? Although the walking

movements of *Euplotes* are superficially similar to those of animals such as insects, the low

Reynolds environment of aquatic microorganisms, where viscous forces dominate over inertial

forces, imposes significant physical constraints on all movements that do not impinge on the

movements of larger terrestrial animals [43]. Coordination, to the extent that it exists in the gait of

Euplotes, would require some kind of dynamical coupling among cirri or between cirri and some

shared external influence. Recently, analytical techniques from statistical physics have been used

to characterize, understand, and predict mesoscale dynamics in biological systems, including

cellular behavior [4,5,44,45]. These approaches rely on coarse-graining the complexity of biological

dynamics into states and analyzing the transitions among states. In particular, a state

representation allows us to ask whether forward and reverse transitions among states are equal, a

condition known as detailed balance [4,46]. Systems that violate detailed balance operate in a non-

equilibrium mode, display net probability flows, and can produce directed cycles in state space

[4,47]. Broken detailed balance has been observed in the motility dynamics of cultured mammalian

cells as well as the motility dynamics of a freely behaving flagellate protist [5,44] and implies that

non-equilibrium models are most applicable to such systems [45]. Identification of broken detailed

balance, therefore, highlights temporal irreversibility and can indicate active control of biological

dynamics.

76   When information processing drives patterns of state transitions, such a system can be

analyzed using automata theory, a fundamental level in the theory of computation [48–50].

Automata theory can be used to address problems of decision-making and control in complex

systems by providing predictive understanding that is independent of the underlying details of

how a given process is implemented [49]. Inspired by work considering cellular behavior in the

context of the theory of computation [25], we hypothesized that walking cells might be governed

by finite state automata with directed, processive movement arising from reproducible patterns

of state transitions. Although the behavioral repertoire of *Euplotes* in different environmental

conditions represents a rich and complex phenomenology involving information processing [51],

for experimental tractability, we chose to focus on the reproducible, spontaneous linear walking

behavior of cells. We reasoned that cellular walking might require some form of information

processing to properly coordinate the movements of cirri.

88   The consistent structure of *Euplotes*, its mode of motility, and its ease of observation

make these cells an ideal biological test-bed in which to apply theories of non-equilibrium

statistical mechanics and embodied computation, both of which rely on describing a system in

terms of discrete state transitions. Here, we use time-lapse microscopy and quantitative analyses

to show that *Euplotes eurystomus* walks with a cyclic stochastic gait displaying broken detailed

balance and exhibiting elements of stereotypy and variability, in accord with a finite state

automaton representation. The observed dynamics are reminiscent of behavioral regulation in

some cells and animals [5,52] but contrast with many well-characterized examples of cellular and organismal motility [7,9,59,10,12,53–58]. Our results provide a demonstration of state machine-like processes governing cellular state transitions as well as an illustration of how such a computational perspective can drive mechanistic insight and serve as a framework for investigating the principles of behavioral control and non-equilibrium dynamics in single cells.

Results

A reduced state space is sufficient to describe walking dynamics

In order to ask whether cell behavior is governed by a finite state machine, we analyzed the walking behavior of *Euplotes eurystomus* cells, [40], focusing on the simplest case of uninterrupted, linear walking trajectories (Figure 2A,B, Video S1). Cells were placed onto coverslips on which free, spontaneous walking behavior was observed by microscopy (imaged at 33 frames/s). A focal plane at the cirrus-coverslip interface was chosen in order to clearly observe cirral dynamics (Figure 2A). The consistency of the relative spatial positioning of cirri across cells allowed us to give each of the 14 cirri an alphabetic label from a-n (Figure 2C). During walking, cirri move in a manner analogous to the recovery stroke-power stroke cycle executed by many eukaryotic flagella, first lifting off the substrate and sweeping close to the cell body before extending in roughly the direction of cell orientation before sweeping downward to reestablish contact with the substrate [37,60,61] (Figure 2B, Video S1). In each video frame, the walking state of the cell was encoded as a 14-element binary vector, with each element corresponding to a cirrus and receiving a value of "0" if the cirrus was in contact with the coverslip and stationary and a "1" if the cirrus was in motion or had moved in the preceding interval between frames (instances of stationary cirri held above the coverslip for a sustained period of time were not observed). The trajectories of 13 cells were manually tracked and

annotated for a total of 2343 time points. This quantitative analysis revealed stepping-like cirral

dynamics: cirri tend to undergo rapid movements followed by longer periods of quiescence

(Figure 2D). Cirral dynamics seemed to lack any obvious patterns such as periodicity or

repeating sequences of states (e.g. Figure 2D), implying that the state sequences are generated

either by stochastic processes or complex deterministic mechanisms. This lack of periodicity

(confirmed by autocorrelation analysis, Figure S1) or fixed phase relationships between

appendage movements is different from those reported for various unicellular flagellates and the

gaits of most animals [58,62,63].

Despite the apparent complexity of cirral dynamics, we wondered whether there might be

some underlying structure, which would allow us to effectively coarse-grain the dynamics in a

principled manner. We first sought to obtain a reduced state space that could accurately describe

the dynamics, as has proven successful in behavioral analysis of diverse living systems [44,45,64–67].

Because our ultimate goal was to identify interpretable motifs among the patterns of cirral

activity, which entail strictly nonnegative values, we chose to perform dimensionality reduction

by non-negative matrix factorization (NMF), a technique that has been used to identify patterns

for textual analysis, natural language processing, neural activity analysis, and gene expression

analysis [68,69]. NMF revealed the cirral states to be well-described by a three-dimensional NMF

space (Figure 2E-G, see Method Details and Figure S1 for more details). Factors of the NMF

(H1, H2, H3) constitute the basis of the NMF space and correspond to non-overlapping groups of

cirri. These factors indicate features of cirral activity that in various weighted combinations can

be used to compactly represent all of the cirral activity measurements. These groups of cirri

constitute spatially distinct partitions of cirri with respect to their positions on the cell body

(Figure 2H). The dimensionality reduction of the gait state space arises in part from shared

pairwise mutual information between groups of cirri (Figure 2I). Here, mutual information

quantifies the amount of information about the patterns of activity of one cirrus gained by

measuring the activity of another, and in this way acts like a generalized measurement of

correlation. Therefore, dimensionality reduction by NMF reflects correlations in cirral activity.

Noting the apparent structure of the data visualized in the reduced gait state space in the

form of clusters of points in NMF space (Figure 2E-G), we next sought to identify individual gait

states defined in terms of similar patterns of cirral activity indicated by these clusters. To do this,

we applied the density-based spatial clustering of applications with noise (DBSCAN) algorithm

[70] to group the output of NMF into clusters in an unbiased fashion, with members of a given

cluster sharing similar patterns of cirral activity (see Method Details and Figure S1 for more

details). Although NMF can itself be used to define clusters, the advantage of clustering in two

steps, NMF followed by DBSCAN, is that we could use the visual observation of clustering in

the NMF output to confirm correct performance of the subsequent clustering step. The effect of

this two-step approach to coarse graining, which is a commonly used in clustering of single-cell

transcription data, is to reduce noise while retaining features that can accurately capture the

complexity of cirral activity. Visual inspection in conjunction with silhouette coefficient (a

metric of cluster cohesion and separation) analysis [71] revealed that 32 clusters accurately

captured the visible structure in the reduced state space without overfitting (Figure 2E-G,

Method Details and Figure S1). These reduced gait states correspond to distinct patterns of cirral

activity (Figure 2J).

Taken together, our results reveal stereotypy in the spatiotemporal patterns of cirral

activity, consistent with the behavior of a finite state machine. The discrete set of gait states,

which exist in the reduced gait state space, demonstrate that cells make use of a subset of the

possible patterns of appendage movement during walking locomotion.

***Euplotes* walks with a cyclic stochastic gait**

In order to relate the gait states identified in our cluster analysis to cell motility, we asked

how changes in the number of active cirri may relate to cell movement. At low Reynolds

number, velocity of a suspended particle is proportional to the difference between the net force

acting on it and the opposing drag [43], so cell velocity provides a relative readout of net

locomotive force. Naively, one might expect that the force associated with locomotion is roughly

proportional to the number of moving appendages [72]. Alternatively, the velocity might inversely

correlate with the net change in cirral activity, which would be expected if stationary cirri were

generating a pushing traction force as in crawling or climbing animals [62,73] or if cirri execute a

power stroke just before coming to rest as has been suggested previously [37]. However, we found

that data supported neither expectation: cell velocity was only weakly correlated with number of

active cirri (R^2=0.03), and instead, the largest cell velocities corresponded to small-to-moderate

changes in the number of active cirri (Figure 3A). We hypothesized that transitions between gait

states must be important to driving the forward progression of walking cells, and thus sought to

determine whether such active coordination might manifest in the observed gait dynamics.

To search for evidence of active coordination dictating gait state transitions, we

calculated the forward and reverse transition rates between states from the 1423 pairwise

transitions in our dataset as N_{ij}/T where N_{ij} is the total number of transitions observed between

states i and j, and T is the total observation time (Figure 3B, see Method Details). We note that

the transition rate we define here is proportional to the probability current $J_{ij} = P_i k_{ij}$ where P_i is

the probability of state i and k_{ij} is the conditional transition probability analogous to a chemical

reaction rate. Although walking, and cilia-based motility in general, is fundamentally non-

equilibrium in the sense that it requires energy input, a coarse-grained representation of motility

such as the one we pursue here need not exhibit broken detailed balance [5]. However, broken

detailed balance in gait state transitions (unbalanced forward and reverse transitions) would

indicate a non-equilibrium process dictating temporal directedness (irreversibility) to gait state

transitions, thus implicating active coordination in terms of stereotyped sequential changes in

cirral activity. Importantly, broken detailed balance in gait state transitions define an identifiable

temporal direction in sequences of gait state transitions. The presence of strongly unbalanced

transitions such as from gait state 3 to 17 versus 17 to 3 suggested broken detailed balance, and

indeed, a number of forward and reverse transitions were found to be significantly unbalanced by

the binomial test (see Method Details). Unbalanced transitions can also arise in equilibrium

systems that have not yet reached steady state, in which case transition rates may change in time.

To test whether the gait operates at steady state, we checked whether the total number of

transitions into each state were balanced by total transitions out of that state: $\sum_j N_{ij} = \sum_j N_{ji}$.

Consistent with steady state dynamics, we found that this condition held to within a difference of

at most a single transition. Although there are cases where non-equilibrium state transitions can

nevertheless exhibit balanced transitions, the presence of unbalanced transitions existing as part

of loops in state space unambiguously indicated non-equilibrium dynamics. To illustrate this

non-equilibrium or temporally irreversible character of gait state transitions, we apply the

Kolmogorov criterion, which specifies the necessary and sufficient condition for reversibility

that the product of transition probabilities traversing any closed loop in state space must equal

the product of the transition probabilities in the reverse direction [5,74]. Due to the presence of

unbalanced transitions, the gait of *Euplotes* clearly violates this condition; for example,

$k_{1,2}k_{2,3}k_{3,17}k_{17,1} = 0.003 \neq k_{1,17}k_{17,3}k_{3,2}k_{2,1} = 0.0005$, where each $k_{i,j}$ is the conditional probability of

transitioning from state i to state j estimated as N_{ij}/N_i with N_i the total number of transitions from

state i. To further understand the degree to which detailed balance was broken, or, similarly, the

distance from equilibrium, we calculated the entropy production rate [5]. Following the procedure

detailed by Wan and Goldstein [5], we obtained a lower bound estimate for an entropy production

rate of 0.4 nats, similar to the value reported for strongly non-equilibrium gait transitions

observed in a flagellate [5]. Walking *Euplotes* cells, therefore, have a non-equilibrium gait,

displaying temporal order in sequences of appendage movement, despite a lack of standard gait

periodicity.

We next sought to better understand the organization and sequential logic of this unusual

gait. First, we noted that gait state transitions appear constrained: only 322 of the 1024 possible

types of transitions were observed to occur at least once, and within this restricted set, only 173

occurred more than once (Figure 3C). Crucially, the presence of broken detailed balance

revealed directed cycles of cirral activity during locomotion pushing the system away from

equilibrium, yielding a temporal directedness or irreversibility to sequences of gait state

transitions in spite of the lack of standard gait periodicity. To investigate the nature of these

cycles, we grouped transitions into two categories: balanced transitions, which satisfy detailed

balance, and unbalanced transitions, which do not (see Method Details). This partitioning

allowed us to separately investigate unbalanced, non-equilibrium-like and balanced, equilibrium-

like transitions (Method Details, Figure 3D,E). We found that unbalanced transitions occur at

relatively high frequency but involve a small number of states (Figure 3D). Of the 32 gait states,

only states 2, 3, 4, 7, 17, and 27 (Figure 3F) are associated with unbalanced transitions, and

among these states, three formed a directed cycle following 2→3→17→2. We had expected that

unbalanced transitions might be associated with a "power stroke" (in the sense of occurring

simultaneously with cell movement) but found instead that high cellular velocities tend to be

associated with balanced transitions (Figure 3D,E) and that relatively few transitions

corresponded to substantial cell movement (Figure 3C). Additionally, we found that, with the

exception of transitions between states 1 and 2, the highest frequency transitions are unbalanced

(Figure 3D).

Notably, the most frequent balanced transitions were associated with transitions into and out of gait state 1, a unique "rest state" which involves no cirral movement (Figure 2J). Furthermore, we found by autocorrelation analysis of gait trajectories that gait state 1 uniquely exhibited significant positive autocorrelation (see Method Details, Figure S1). Investigation of transitions to and from gait state 1 revealed that although transitions between gait states 1 and 2 are balanced, the most frequent transitions out of gait state 2 are strongly biased toward transitions into gait state 3, from which other strongly biased transitions also frequently occur, including the cycle of biased transitions mentioned above. Despite the presence of high frequency unbalanced transitions, the gait of *Euplotes* involves highly variable trajectories through gait state space. The picture of walking trajectories that emerges is one of stochastic excursions from gait state 1 into non-determinate paths through state space involving a mix of balanced and unbalanced transitions. The majority of cell movement occurs during infrequent, equilibrium-like (balanced) transitions. These balanced transitions are temporally reversible, but we wondered whether there might be some asymmetry in the amount of cell movement due to transitions leading toward versus away from the cycle states. We found that although balanced transitions leading away from the cycle states account for 7.6% more cell movement, the difference in these transitions is not statistically significant (p=0.09 by two-sample Kolmogorov-Smirnov test). Temporal irreversibility, or directedness, in sequential changes in cirral activity arises from biased, non-equilibrium-like (unbalanced) transitions, occurring at relatively high frequency from a small subset of states. A natural question arising from the observation of these complex patterns is whether gait state transitions conform to a first-order Markov process, often referred to as "memoryless" [75], which entails that transition probabilities are determined completely by the present state, and that previous dynamics contribute no additional predictive information [76,77]. Does the cell retain some "memory" of past cirral activity that might influence

future cirral activity? While not conclusive, several analyses (see Method Details and Figure S1) suggested that *Euplotes* retains some "memory" of the prior sequence of cirral movements during locomotion in that the gait may not conform to a continuous- or discrete-time first-order Markov process.

Taken together, our analysis revealed a mixture of unbalanced transitions driving cycles and balanced transitions arranged as networks, for which we propose to apply the term "cyclic stochastic gait". The cyclic stochastic gait of *Euplotes eurystomus* incorporates elements of both stereotypy and variability in gait dynamics, in terms of biased transitions and non-determinate sequences of gait state transitions respectively. Forward progress of the cell is not produced merely by a physical ratchetting process driven by unpatterned fluctuations in cirral activity, nor is it produced by a highly regular, deterministic process like a clock. It has been argued that significant computation arises in physical systems exhibiting such a mix of stereotypy and variability [48,78–80] in the sense that the time-evolution of the system is most compactly described as the result of a computational process involving state transitions, memory, and decision rules, rather than periodic oscillations or random coin flips.

Our analysis suggested a computational underpinning of gait, so we sought to better understand the functional organization of the dynamical patterns, the sequential logic of the gait. Focusing on the dominant structure of gait transitions in terms of transition probabilities (Figure 3G, Method Details) allowed us to derive a simplified, functional representation of stereotypy in gait dynamics as depicted in Figure 3H. We found that few states were the recipients of the majority of the highest probability transitions and that many received no high probability transitions (Figure 3G). Additionally, we found a "cloud" of states linked by low-probability balanced, equilibrium-like fluctuations. Nearly all of the states receiving high probability transitions were either the three "cycle states" or else fed cycle states with their highest

probability transitions, with the majority feeding gait state 17. Although gait state 1 is not the

recipient of any individual high probability transitions, we identified it as the unique "start" state

from which cells initiate walking. Beginning with this start state, cells transition with high

probability to gait state 2, also one of the highest frequency transitions and the first state in the

2→3→17→2 cycle of unbalanced transitions. From this first cycle state, cells transition to gait

state 3, the second cycle state, with highest probability and frequency and then similarly on to

gait state 17, the third cycle state. This sequence from the start state through the cycle states

corresponds to increasing amounts of cirral activity. Although the highest probability transitions

from the third cycle state to any single gait state tend to return to the first or second cycle state

with equal probability, cells in fact transition to the equilibrium "cloud" of motility-associated

states with overall higher probability. Return to the cycle states tend to occur through various

moderately high probability transitions from the motility state cloud or through a restricted set of

intermediate states. In conjunction with this set of transitions, we also noted unbalanced

transitions stemming from the cycle states to the motility state as well as the presence of

intermediate states from a given cycle state that subsequently feed the next cycle state.

Altogether, the picture of stereotypical gait dynamics that emerges is of biased transitions

involving cycle states preceding relatively low probability, unbiased transitions associated with

substantial cell movement before returning to the start or cycle states and beginning the sequence

again. While this general sequence is repeated during walking, there is considerable variability or

apparent stochasticity in the details of gait state transitions departing from the start state with

increasingly variable transitions as any given sequence progresses. Given that the motion of the

cell alternates between periods of highly processive forward motion and other periods in which

the cirri seem to flail ineffectually without driving the cell forward, one might assume that the

periods of effective motion would correspond to the highly directed cycle of unbalanced

transitions and the periods of flailing would correspond to the cloud of balanced, random

transitions, but in fact the opposite is the case. We hypothesized that sequences involving the

cycle states serve to establish configurations of cirri necessary for cells to later transition

between states from which substantial forward progress of the cell is generated. Many state

transitions along any instance of the stereotyped sequence are unbiased; however, biased, high

probability transitions, presumably resulting from active cellular control, give temporal

irreversibility to the sequence. We note here that although our coarse-graining procedure to

identify gait states does not constitute a unique representation of gait structure, we can be

confident that our analysis captures structure in gait dynamics, the presence of broken detailed

balance in particular. In general, coarse-graining of the state space of a system can obscure

broken detailed balance, but the net flux of transitions in the state space of a system should not

arise in an illusory fashion based on a coarse-graining procedure or partial observation of a

system [81–83]. Additionally, we have chosen a coarse-graining procedure based on the properties

of our particular data and demonstrate its performance on simulated cirral dynamics with varying

noise and for the case of unpatterned, random cirral fluctuations (Figure S1, Method Details).

Finally, we sought to investigate the functional significance of the sequential logic of the

gait in driving processive cell movement. To do so, we developed a simple model based on a

coarse-grained physical picture of cellular walking (Figure 3I). Briefly, we consider a 2D system

where a cell walks using its 14 cirri, which can exist in two states: moving or not moving. As we

are agnostic to the details by which cirri generate forces involved in motility, we model cirri as

producing constant force in the direction of cell orientation while moving and resisting

displacement by acting as linear springs when not moving. Cell position and orientation is

defined in terms of the equilibrium positions of the cirri. For simulations, we use relative

positions of cirri taken from stationary cell measurements (Figure 3J). In each timestep of the

simulation, we calculate a cell displacement based on the sum of forces due to the cirri and

changes in cell orientation from the sum of torques (see Method Details). We found that this

simple model was sufficient to qualitatively and semi-quantitatively reproduce the linear

trajectories of walking cells when we ran simulations using the actual patterns of cirral activity

from walking cells (Figure 3K, Figure S2). When we ran simulations with either the same gait

states as those from actual cells but in a shuffled order or random cirral activity with the same

average cirral activity as actual cells, we found that path straightness significantly decreased in

the case of shuffled transitions (p=0.04 by Wilcoxon rank sum test) and scaled cell velocity

significantly decreased in the case of random activity (p=0.003 by Wilcoxon rank sum, Figure

3K-O). Note that in both the case of shuffled transitions and random cirral activity, gait state

transitions satisfy detailed balance (Figure S2). These results suggest that the sequential logic of

the gait is functionally significant in a manner consistent with our biophysical picture of walking.

The microtubule-based fiber system mediates gait coordination

The complex yet sequentially structured gait patterns in conjunction with our simulation

results are consistent with the existence of some form of nontrivial gait coordination. What

physical machinery could embody the information processing required to generate the stochastic

cyclic state transitions seen during *Euplotes*' walking? We reasoned that there must be some

form of coupling or communication between cirri or feedback between gait states and cirral

dynamics. Since the early 1900s , the role of the system of cytoskeletal fibers associated with

cirri as conduits of information between cirri during cellular locomotion, supported by

microsurgical experiments, has been a dominant yet contentious hypothesized mechanism of gait

coordination [84–86]. We wondered whether the structure of the cytoskeletal fiber system associated

with cirri (Figure 4A) could give some insight into how cirri might be coordinated.

We reconstructed in 3D the microtubule-based fiber system of *Euplotes* associated with

cirri and lying just beneath the cell cortex [38,41,42]. Upon inspection of our confocal

reconstructions of SiR-tubulin labeled cells (Figure 4A, Figure S3), we noted the presence of two

morphologically distinct classes of fibers, one thicker and linear and the other thinner, splayed,

and less linear, consistent with previous observations (Figure 1C, [38,41,42]). Fibers emanate from

the base of all cirri, appear to intersect one another, and also connect to the cortex of the cell at

various points. Some cirri were found to be associated with only thick fibers, while others have

both or only thin fibers. Based on apparent fiber intersections and convergences, we found the

fiber system forms a continuous network between all cirri, with the fibers associated with the

base of each cirrus intersecting the fiber system of at least one other cirrus (Figure 4A,B).

Contrary to the long-standing standing hypothesis from the literature [84], the functional

modules (groups of co-varying cirri) identified in our dynamical analysis were not exclusively

linked by dense fiber intersections (Figure 4A,B) [38,42,84]. In fact, connections between cirri are

not generally associated with any statistically significant difference in mutual information

(defined in terms of the information that the activation state of one cirrus has concerning the

other) compared to unlinked pairs of cirri (p=0.14 by Wilcoxon rank sum test, Figure 4C).

However, information flow became apparent when fiber-fiber links were grouped by type (i.e.

thick to thick fiber, thick to thin fiber, or thin to thin fiber). Under this grouping, we found that

pairs of cirri associated with only thick to thick fiber and only thin to thin fiber links have

increased mutual information compared to those without links (Figure S3). Interestingly, we

found that cirri nearby one another and connected by fibers to similar regions of the cell cortex

shared the most mutual information (Figure 2C,G, 4D,E), suggesting that if the fibers play a role

in cirral coordination, coupling may also be mediated by mechanisms involving the cirrus and

fiber-cortex interface. Cirri d, e, h, i, for example, share very little mutual information with any

of the other cirri, and fibers emanating from the base of these cirri contact the cell cortex and

other fibers at various unique points. On the other hand, cirri g and f, which share more mutual

information than any other pair, are associated with both thick and thin fibers terminating at

similar regions of the cell cortex. Indeed, distances between pairs of cirri and cross nearest-

neighbor distance [87] between paired sets of cirrus-cortex contact points both show significant

Spearman correlations (-0.49, p<0.001 and -0.62, p<0.001 respectively) to mutual information

(Figure 4D,E). These correlations indicate that mutual information between pairs of cirri tends to

increase with proximity and also tends to increase with similarity between fiber-cortex contact

locations. Thus, the cirri with the highest mutual information are those that are close together

with similar fiber-cortex connections (Figure 4D-F).

Together, these observations suggest a mechanism of mechanical coordination in which

microtubule bundles allow groups of cirri to influence successive behavior of other groups of

cirri. We first sought to test this hypothesized mechanism by perturbing the fiber system using

drug treatments. We observed that nocodazole, a drug that blocks the polymerization of

microtubules, affected walking motility, causing cells to walk along more compact trajectories

due to increased turning (Figure 5A,B, Videos S3 and S4) in a manner reminiscent of historical

reports of altered walking due to microsurgery in ciliate *Stylonychia* [88]. In contrast to the effect

of nocodazole, we found that the microtubule stabilizer paclitaxel caused cells to walk along less

convoluted trajectories compared to controls (Figure 5C). Quantifying these effect from video

microscopy of cells under darkfield illumination in terms of a scaled path length defined as the

total integrated path length walked by cells scaled by the maximum radial distance traversed,

where a decrease in linear runs decreases this scaled path length, we found that nocodazole

significantly decreased and that paclitaxel significantly increased the scaled path length of cells

compared to controls (Figure 5D). We further noted that nocodazole acted in a dose dependent

and reversible manner (Figure 5D and Figure S4). To test whether the actin cytoskeleton might

also regulate motility, we treated cells with actin inhibitors including latrunculin, cytochalasin,

and jasplakinolide and observed no effect on motility (Figure S4). The specific and opposing

effects of a microtubule polymerization inhibiting drug (nocodazole) versus a microtubule

stabilizing drug (paclitaxel) were consistent with the microtubule cytoskeleton playing a key role

in coordinating walking motility. Next, we checked whether nocodazole treatment had an

observable effect on the fiber system. By analyzing 3D reconstructions of confocal images of

SiR-tubulin labeled cells, we found that fiber length significantly decreased compared to controls

in cells where microtubule polymerization was disrupted by nocodazole (Figure 5E-G). Further,

we were unable to detect the presence of any thin fibers in four out of seven nocodazole treated

cells used for fiber analysis. Of the three cells with detectable thin fibers, we never observed

connections between fibers associated with cirri a, b, and c and any other fibers. Additionally, we

did not observe any thin fibers making distal cortical contacts. We did note, however, that when

thin fibers were visible, connections appeared to be the same as those in Figure 4A,B. We then

investigated how cirral dynamics were affected. Following the gait annotation procedure detailed

previously, we characterized the walking dynamics of 6 nocodazole treated cells for a total of

1133 timepoints. Of those timepoints, 681 corresponded to cirral configurations never observed

in untreated cells with a total of 391 new unique configurations. Projecting these new

configurations onto the NMF space we obtained previously, however, revealed that most of the

cirral configurations in nocodazole treated cells were near or within the clusters corresponding to

the gait states we obtained from untreated cells (Figure S4). This allowed us to map new cirral

configurations onto the gait states (see Method Details). We did note that while close to the

original gait states, new cirral configurations tended to skew towards more active cirri, and we

also noted the presence of a new cluster involving movement in all or nearly all cirri, to which

we assigned a new gait state (Figure S4). We also found that mutual information between cirri

was higher in general, with many pairs of cirri sharing higher mutual information than the

highest values obtain in untreated cells (Figure 5H, Figure S4). This increase in cirral activity

and correlations is consistent with the fibers playing a role in conveying inhibitory information

during unperturbed walking. In further support of this inhibitory role, we found that paclitaxel

treatment also caused an overall increase in mutual information of pairs of cirri (Figure 5H).

Next, we investigated how the dynamics of gait state transitions were affected. Following

our previous analysis, we evaluated gait state transitions for the presence of broken detailed

balance, separating state transitions into balanced and unbalanced transitions and found that the

structure of state transitions differed greatly from that of unperturbed cells. Although some gait

state such as 1, 2, and 17 were involved in high frequency transitions in both conditions, gait

state transitions of cells with perturbed fiber systems exhibited less broken detailed balance and

closer to equilibrium-like dynamics as indicated by an entropy production rate of 0.1 nats

(compared to 0.4 nats in the unperturbed case), including the loss of the unbalanced, cyclic

transitions (Figure S4). Figure 5I summarizes the change in the structure of gait state transitions

including changes in broken detailed balance, reduction in transitions toward the states

previously involved in cyclic transitions, and the reduction in the occupancy of states associated

with the cloud of states involved in substantial cell movement, with only 10 of the original 21

cloud states sampled. We also noted that many of the new highest probability transitions feed

gait state 18 (Figure 5I, Figure S4), which involves nearly exclusive activity in cirri f and g

located close to one another at the edge of the cell. A persistent bias toward activation of these

cirri, which are farthest from the central axis of the cell, may in part explain the increase in

turning in trajectories. These results, in conjunction with the fact that cell velocities were

indistinguishable control conditions (Figure S4) are consistent with predictions from simulations

in which shuffling the order of gait state transitions reduced processivity without affecting speed

(Figure 3N,O).

Although the result that disrupted gait coordination involves increased mutual

information between cirri, note that this increase is due to more frequent joint activation or

synchronous movements among cirri and so stems from reduced complexity of gait dynamics.

For a complementary perspective linking these changes in gait dynamics to an underlying

computational process, we applied the Causal State Splitting Reconstruction (CSSR) algorithm

to construct ϵ-machines corresponding to walking cells [48,89]. ϵ-machines are automaton models

consisting of a set of causal states with transitions between them and represent the minimal

model consistent with accurate prediction of a stochastic process [48]. The causal states of an ϵ-

machine indicate how the process from which it is constructed stores information, and state

transitions indicate how the process transforms information [48]. We found that ϵ-machines

constructed from the cirral activity of untreated cells tended to be similar to one another and

were more complex in terms of having more causal states and transitions than those constructed

from nocodazole treated cells, which also tended to be similar to one another (Figure S5, Method

Details). This reduction in complexity may reflect reduced computational capacity of the

nocodazole treated cells. Further, in conjunction with the decrease in entropy production under

nocodazole treatment, these results are consistent with recent findings linking increases in

broken detailed balance to increases in information processing in biological systems [90].

As a final additional test of the role of the fibers in mediating gait coordination, we

revisited historical microdissection experiments. Although Taylor first reported in 1921 results of

microdissection experiments indicating a role of the fiber system in mediating gait coordination

[84], a reproduction of these experiments in 1966 with improved physiological conditions and

quantitative analysis failed to observe disrupted coordination among cirri upon bisection of cells

[85]. Importantly, neither of these reports involved analysis of walking behavior. We performed microdissections on cells using pulled quartz microneedles, severing cells transversely just in front of cirrus h, ensuring that we had severed all fibers associated with cirri j-n (Figure 5J,K). Similar to previous reports, we found that cell fragments regained spontaneous cirral activity after a brief recovery period. After 24 hr, we found that anterior portions of cells began to exhibit spontaneous walking activity (Figure 5K,L, Video S6), which persisted for up to 72 hr subsequently and has not been previously reported. Although these fragments were able to walk across surfaces, in all cases, they walked along curling or circular trajectories reminiscent of nocodazole treated cells except in the reverse direction (Figure 5L, Video S6). We analyzed the cirral dynamics of 10 fragments and found significantly higher mutual information shared between cirri indicating more synchronous cirral activity, similarly microtubule inhibitor treated cells, compared to cells that were merely wounded and displaying apparently normal motility (Video S7) or unperturbed (p<0.001 in both cases by Wilcoxon rank sum test), which displayed values consistent with one another (Figure 5M). As with the microtubule inhibitor experiments, these results are consistent the fibers playing an inhibitory role in gait coordination.

Taken together, these results are consistent with our hypothesis that the fiber system mechanically mediates gait coordination. Further, these results provide additional evidence that proper, active coordination of cirral dynamics and emergent gait states and gait state transitions is required for proper cellular walking.

Discussion

Faced with the challenge of accounting for the emergence of apparently sophisticated cellular behavior, the directed walking of *Euplotes* driven by a seemingly disordered gait, we conceptualized the cell as a finite state machine. Traditionally, studies of computational processes performed by cells have focused on combinatorial logic, where the output of a

computation depends only on the current input, performed by networks of molecules in cells

[25,31–33]. We have focused on sequential logic, where outputs depend on the system state as well,

an equally important aspect of the theory of computation with notable yet less developed

representation in studies of cellular dynamics [16,20,91]. Automata theory, which includes finite state

machine models and necessarily involves sequential logic, provides tools for understanding

structure and stereotypy in transitions between dynamical states, increasingly appreciated as

features of the behavior of eukaryotic cells. Our approach revealed modularity in cellular

dynamics associated with structural modularity of the cell (Figure 2, 4) in addition to stereotyped

patterns of sequential activity (Figure 3) and yielded new insight into regulatory mechanisms

(Figure 3, 5).

Related approaches involving the coarse-graining of complex dynamics and subsequent

analysis have proved useful in revealing simplicity and stereotypy in the behavior and movement

patterns of various organisms including flies, worms, and protists[3,5,65,66,92]. Reducing the

complexity of biological dynamics in this way can be an important step toward making

quantitative predictions and for interpreting the effects of perturbations. In general, particular

approaches to coarse-graining will depend on details of the biological system, type of data

collected, and underlying patterns in the data. The specific steps we have taken in the present

study are best suited to biological processes that can be characterized in terms of discrete events.

The analyses following coarse graining stand to provide the most insight into processes

involving information processing where the sequential order of events may be important to the

resultant function. Additionally, our perturbative experiments demonstrated how these

approaches can drive mechanistic insight by helping connect function and computation to

underlying mechanics and physiology.

Although there are examples of locomotor coordination reminiscent of the stochastic, non-equilibrium gait dynamics of *Euplotes*, such as gait switching in Kangaroo rats [52] or, most saliently, gait switching in an octoflagellate [5] and motility in cultured mammalian cells [44], walking locomotion in *Euplotes* represents a departure from many of the best studied appendage-based locomotor systems. For example, limbed locomotion in animals tends to proceed by highly stereotyped, determinate patterns of activity [58,62], and many small, aquatic animals exhibit periodic movements of appendages, often cilia, during locomotion [7,63,93]. Many forms of unicellular locomotion involve such dynamics as well including in sperm cells [94], diverse flagellates with various numbers of flagella [63], and ciliates [63,95,96]. Even in cases where cellular locomotion involves fundamentally stochastic dynamics such as in run-and-tumble motility in *E. coli* [12] or analogous behaviors observed in protists [11,97–99], motility can be described by equilibrium processes [5], in contrast to the non-equilibrium character of the gait of *Euplotes*.

Here, broken detailed balance in gait state transitions revealed cyclic activity with a mixture of stereotypy and variability in the gait of a single cell (Figure 3). To explain how these dynamics give rise to directed walking, we propose a mechanism, captured by our biophysical model, in which biased, actively controlled cyclic transitions serve to establish strain, effectively storing stress, in certain cirri, and the spontaneous release of these cirri from the substrate, during a series of unbiased gait state transitions, allows the cell to move forward. The cloud of unbiased transitions associated with substantial cellular movement is consistent with motility generation not depending on the precise order in which the strained cirri are released from the substrate. Return to the cycle states then are necessary to establish this process anew by winding up the system for continued, proper cell movement. Disruption in this resetting may lead to defects in walking as predicted by simulations (Figure 3K-O, 5) and with the reduction in entropy production and loss of the cycle states in irregularly walking cells with disrupted fiber systems,

which appeared unable to consistently maintain linear trajectories (Figure 5B,D,I, Video S4). We

find additional experimental support for our proposed mechanism in previously reported

observations of cyclic velocity fluctuations in the trajectories of walking *Euplotes* [36].

We argue that subcellular processes are involved in actively coordinating cirri and

propose that broken detailed balance in the gait of *Euplotes* indicates this active coordination.

The results of experiments perturbing the tubulin-based cytoskeletal fiber system are consistent

with its role in mechanically mediating communication both among cirri and between cirri and

the cell cortex (Figure 4,5). We conjecture that movement of cirri relative to one another can

establish tension in the fiber system and that the tension state of fibers associated with each

cirrus may then modulate cirral activity in a manner reminiscent of basal coupling in flagellates

[22]. Further, It is possible that microtubules may mechanically mediate the coordination of cellular

processes in other eukaryotes, and microtubules in cells have recently been shown to respond

directly to mechanical forces [100]. Microtubules have also been shown and proposed to be

involved in more complex signal transduction pathways[101,102]. Changes in fiber length, as

observed under nocodazole treatment, change the relative positions of cirri, which could lead to

different distributions of tension and thereby induce the altered pattern of state transitions and

associated walking defects. Our results show that perturbation of the fiber system shift the gait of

Euplotes from a regime of asynchronous yet coordinated movement to a dysregulated regime

with synchronous yet improperly coordinated movement. It should be noted, however, that the

fact that cells with perturbed fiber systems, including bisected cells, display some walking ability

implicates additional robustness and complexity in mechanisms of gait coordination and cell

motility. Although details of the mechanism remain to be fully explored and tested, our results

are consistent with the fibers system playing an active and inhibitory role in gait coordination.

Thus, by combining information processing to properly dictate patterns of cirral activity and the

mechanical actions of cirral movement, walking *Euplotes* embodies the sequential computation of a finite state machine. Furthermore, our approach to understanding a complex cellular behavior, grounded in theory of computation, allowed us to derive mechanistic insight from observational and perturbative experiments. Because reproducible biological function often emerges from the productive management of stochastic fluctuations, we expect our conceptual and analytical approaches may be useful in studying other living systems.

Our work lays a foundation for studies of sensorimotor behavior in *Euplotes*, which we believe will shed light on principles of cellular behavior. We have focused here on a single behavior (spontaneous, linear walking), but *Euplotes* exhibits a complex behavioral repertoire in response to diverse environmental cues, contexts, and physiological states[39,40,103104]. Many of these behaviors involve different patterns of cell motility. Our biophysical model of cell walking could help predict and interpret how changes in gait state transitions or the emergence of new gait states might give rise to these different patterns of movement. We speculate that the unusual gait of *Euplotes* and associated asymmetric cell structure, contrasting sharply with deterministic appendage movements and symmetric structure in animals and many swimming flagellates and ciliates, may reflect physical and phylogenetic constraints. The framework we have developed integrates across scales of space, time, and biological organization to link mechanics and subcellular dynamics and structure to overall cell movement patterns. Therefore, our work provides a way to begin to address the functional significance of cell morphology, gait patterns, and mechanisms of gait control, and in conjunction with comparative approaches, could provide new insight into how cellular behavior evolves.

Among the domains of life, eukaryotes uniquely display remarkable complexity and diversity in cellular behavior [105]. Our approach, grounded in finite state machine analysis, has revealed modularity and stereotypy underlying complex cellular behavior and has provided

insight into regulatory mechanisms across scales of biological organization. Our results suggest

that integrating approaches from theoretical computer science, non-equilibrium statistical

physics, and cell biology stands to shed light on the regulation of cellular behavior in eukaryotes

more broadly. By revealing principles of cellular behavior, the line of research established here

stands to advance our ability to predict, understand, and even one day engineer cellular behavior

across diverse eukaryotic systems.

Figures

Figure 1. ***Euplotes* exhibits highly polarized, complex cellular architecture and walks across surfaces using microtubule-based organelles called cirri, some of which are physically linked.** (A) A single *Euplotes eurystomus* cell in profile displays its ventral cirri, which are used for walking locomotion across surfaces (arrowhead indicates a single cirrus stretching out from the cell). Scale bar is 10 μm. (B) A drawing of a *Euplotes* cell, viewed from the ventral surface, highlighting the complex, asymmetric structure of the cell. Notable features include the cirri (ci) and the membranellar band (m), wrapping from the top of the cell to the center, which is used to generate a feeding current to draw in prey items. Drawing adapted and obtained from Wikimedia Commons, from original source [106]. (C) A drawing of a *Euplotes* cell, highlighting the fiber system associated with the cirri, historically referred to as the neuromotor apparatus. Drawing adapted from [107].

Figure 2. *Euplotes* walks across surfaces according to a complex gait, which can be described in a discrete, reduced state space with gait states corresponding to identifiable patterns of cirral activity. (A-A''), The movements of cirri during walking locomotion are

clearly visible by brightfield microscopy by focusing on a plane at the surface of the coverslip on

which cells are walking. Three snapshots depict different time points during a single walking

trajectory, and white arrowheads indicate cirri. In the panels from left to right, the cirrus

indicated by the arrowhead on the left is stationary, stationary, and then moving, and the cirrus

indicated by the arrowhead on the right is stationary, moving, and then stationary. Scale bar is 15

μm. (B) The trajectory of a cell during a single recorded trajectory as the cell walked across a

coverslip from left to right. The cell position was manually tracked in each frame. (C) The

scheme for encoding cirral dynamics during walking involved labeling each of the 14

distinguishable ventral cirri (a-n), and recording cirral activity in each frame, corresponding to

timepoints $(t_1,...,t_n)$, of recordings of walking cells as a 14-bit binary vector. Each entry in each

vector is given a value of either 0 if the cirrus is not moving and in contact with the coverslip or

1 if the cirrus is moving. Scale bar is 15 μm. (D) Representative visualization of cirral dynamics

for a single trajectory of a walking cell. These dynamics correspond to the walking trajectory in

panel E. Each row corresponds to a cirrus and each column is a single video frame. White

denotes cirral activity, a value of 1, in the vector encoding of dynamics from panel F. Note the

dynamical complexity and discrete, stepping-like nature of cirral movements. (E-G) Three

roughly orthogonal views of a plot displays the structure of all recorded cirral dynamics, encoded

as in panel E, from 13 cells over 2343 timepoints in a reduced state space obtained by non-

negative matrix factorization (NMF). Axes correspond to the components of the NMF (H1, H2,

H3), and each point is a single timepoint. Randomized colors highlight the 32 clusters identified

using the density-based spatial clustering of applications with noise (DBSCAN) algorithm [70]. We

refer to these clusters as gait states, and they correspond to unique configurations of cirral

activity during walking locomotion. (H) A plot of the magnitudes associated with each cirrus

corresponding to the components of the NMF of cirral dynamics shows distinct contributions

from spatially distinct groups of cirri. Component H1, for example, is associated with activity in

cirri a, b, and c. The tracing of a cell including the position of cirri has the same color map as the

plot above and shows the grouping of the cirri corresponding to each component. (I) A heatmap

of mutual information between all pairs of cirri shows that correlations in cirral activity

correspond to the NMF components displayed in panel K. For example, cirri a, b, and c share

mutual information with one another and are the cirri contributing to component H1. (J) A

heatmap representation of the cirral activity associated with each of the 32 gait states. Values for

each cirrus are the mean over all instances of the gait state. Note that each gait state has a unique

signature of cirral activity. See also Figure S1, Video S1, and Video S2.

Figure 3. *Euplotes* **walks with a cyclic stochastic gait exhibiting broken detailed**

balance, stereotypy, and state machine-like dynamics. All data for panels A-H are pooled

from the walking trajectories of 13 different cells over 2343 timepoints and 1423 pairwise gait state transitions. (A) A plot of the mean net change in cirral activity versus the net scaled cell velocity associated with all transitions between the 32 gait states identified in Figure 2 shows that the change in number of active cirri is not strongly correlated with cell velocity (R^2=0.03). Cell velocities were obtained from manually tracked walking trajectories and then scaled by dividing frame to frame displacements for each trajectory by the length of the cell being tracked and also dividing by the average frequency of cirral inactivity. Scaling provided a non-dimensional velocity scaled by natural units of the system. Note that at low Reynolds number, velocity should be proportional to force [43], so this plot also reflects the net walking force generated by the cell. Net change in cirral activity was computed using the data presented in Figure 2. Note that the largest velocities are associated with small negative and small to moderate positive net changes in cirral activity. (B) The transition matrix of all gait state transitions, with rows representing the starting state and columns indicating the ending state, exhibits broken detailed balance. Rates were estimated by dividing the total number of observed transitions between each state pair and dividing by the total time observed. Under detailed balance or equilibrium conditions, transitions from one state to another should be balanced by reverse transitions. Lack of this kind of reversibility, as seen by the lack of symmetry of the heatmap across the diagonal, indicates broken detailed balance and non-equilibrium dynamics. (C) A directed graph representation of all gait state transitions. Nodes correspond to the 32 gait states, and node sizes are scaled by the proportion of total time cells spent in each state. Directed edges are represented by arrows between nodes and signify state transitions. The size of the arrows is scaled by transition rates as in panel b. Edge color represents scaled cell velocity as in panel A according to the indicated color scale. (D) A subset of transitions visualized as in panel C shows the restricted and relatively high frequency nature of unbalanced, non-equilibrium-like

transitions. Only transitions that were observed to happen more than one time and exhibiting a

significant difference between forward and reverse transitions (p<0.05 by binomial test, see

Method Details) are displayed. (E) A subset of transitions, similarly to panel d, except that only

the balanced transitions, lacking a significant difference between forward and reverse transitions

(p<0.05 by binomial test) are displayed, also show a complex and widespread structure, this time

of balanced, equilibrium transitions. Note that the majority of transitions associated with high

cell velocity involve equilibrium-like dynamics. (F) Examples illustrating the spatial

organization of cirral activity corresponding to gait states. Some states, such as 7, correspond to

activity in spatially discrete groups of cirri, while others, such as 17, correspond to cirral activity

across the cell. The gait states displayed here are those involved in unbalanced transitions. (G) A

heatmap of transition probabilities between states, showing only the most probable transitions

from a given state with all others set to zero, shows distinct structure. In cases were multiple

state transitions from a state were tied for the highest probability, all of these transitions are

displayed. Fewer than half of the total states are recipients of multiple high probability

transitions, and many states are the recipients of no high probability transitions. (H) A

representation of functional states and transitions between them highlights the machine-like

nature of the gait of *Euplotes*. Gait states are represented as circles with numerical labels. Blue

circles represent states that are both recipients and sources of unbalanced transitions as identified

in panel D and constitute the three cycle states. Red circles represent states that are recipients but

not sources of unbalanced transitions as identified in panel D. Black circles correspond to gait

states that are associated only with balanced transitions as in panel E. States receiving no more

than one unique high probability transition from states with only a single highest one as

identified in panel g were grouped together into a compound state represented by the dark gray

blob. The blue background behind states 4, 16, 18, 26, and 28 indicates that these states all share

the same highest probability transitions between states identified in this panel, and thus, the

group constitutes a single compound functional state. Arrows represent the highest probability

transitions between the states, including compound states composed of multiple gait states as

identified in Figure 2 (dark gray blob and blue background). Gait state 1 is also depicted, as it is

the state in which cells spent the most time over all walking trajectories and also is uniquely the

state from which cells begin walking. Cells also frequently return to the state during walking.

Further, transitions from gait state 2 from gait state 1 constitute the single highest frequency

transition. Together, all identified states in this panel constitute functional states. Arrows

represent the most probable transitions between functional states, and all unbalanced transitions

are also represented with size scaled by their proportional probability compared to all other

transitions emanating from the source functional state. Cartoons are a walking cell in profile with

cirri in a configuration representative of the corresponding functional state. Labels refer to the

apparent functional role of states and their associated transitions. Beginning from gait state 1, the

resting/start state, cells are most likely to follow transitions from gait state 2 to 3 to 17 at which

point cells are likely to enter the functional state associated with substantial cell movement

involving variable balanced transitions between a number of gait states. Transitions are then

likely to lead back toward the cycle states. Note that while this representation of gait dynamics

highlights the most probable transitions, substantial variability, primarily involving reversible

transitions, occurs during walking trajectories. (I) A cartoon depicts the basic features of our

model of a walking cell. Cirri generate a constant force when moving and resist displacement in

a spring-like fashion when in contact with the substrate. Force and torque balance yield cell

displacement and orientation in each simulation timestep. (J) A plot of the relative average

resting cirral-surface contact positions measured from the 13 cells used for gait analysis, which

were used for all simulations and define simulated cell position and orientation. (K) A

representative trajectory of a simulation of a walking cell using experimentally recorded cirral

activity illustrates the linear, processive motion, which matches experimentally measured cell

walking dynamics qualitatively and semi-quantitatively. (L) A representative trajectory of a

simulation of a walking cell using the same experimentally recorded cirral activity as in panel K

but in a shuffled order illustrates a decrease in processivity of cell movement. (M) A

representative trajectory of a simulation of a walking cell with random patterns of cirral activity

with the same average level of cirral activity and the same number of timesteps as in panels k

and l illustrates the decrease in cell speed. (N) A plot of path straightness measured from

simulations using actual cirral pattern measured experimentally (Actual), actual cirral patterns in

a randomly shuffled order (Shuffle), and randomly generated cirral activity with the same

average cirral activity as actual patterns (Random) suggest that the order of gait state transitions

matters for processivity of cell movement due to a predicted decrease in path straightness for

Shuffle trajectories. Path straightness is the distance between the start and end of the cell

trajectory divided by the sum total path length. The trajectories of the 13 cells analyzed for

Figure 2 were used for the Actual simulations. For the Shuffle simulations, cirral activity patterns

from each of the 13 cells were shuffled to generate a random order five times. For the Random

simulations, five sets of patterns of lengths equal to each of the 13 cell trajectories were

generated. The asterisk indicates p=0.04 by the Wilcoxon rank sum test. (O) A plot of scaled cell

velocity for the simulations described for panel k predict that the gait states are responsible for

generating cell velocities as indicated by decreased scaled cell velocity for random patterns of

cirral activity. Scaled velocity was computed as described for panel a and averaged over the

entire cell trajectory. Three asterisks indicates p<0.001 by the Wilcoxon rank sum test. See also

Figure S1 and Figure S2.

Figure 4. The structure of the complex, interconnected fiber system of *Euplotes* correlates with dynamical associations between cirri. (A) The SiR-tubulin labeled cell (faint, dark blue) was imaged by confocal microscopy, and a 3D reconstruction as obtained from serial confocal slices. Fibers were manually traced in each slice using TrakEM2 in FIJI. Two morphologically distinct classes of fibers were observed and are indicated as follows: thick, linear fibers are cyan and thinner, filamentous fibers are magenta (see Figure S3 for more image data). Fibers emanate from the base of each cirrus and form a connected network between all cirri. The base of each cirrus is indicated by corresponding letters (as in Figure 2C). Gray shading indicates the dynamical groups identified by dimensionality reduction and follows the same color scheme as in Figure 2H. Scale bar is 10 μm. (B) A graph representation of fiber-fiber connections illustrates the complex and interconnected nature of cirrus associated fiber topology. Nodes correspond to the cirri to which each fiber system is associated, and edges indicate

connections between fiber systems. Colors of nodes indicate the same groups as in panel a, and colors of edges indicate the types of fibers connecting to one another, cyan for thick fiber connections, magenta for thin fiber connections, and purple for thick to thin fiber connections. (C) Pairs of cirri that are linked by fiber-fiber contacts show no statistically significant difference in mutual information compared to those lacking fiber-fiber contacts. The plot displays mutual information between all pairs of cirri grouped by the absence (Not linked) or presence (Linked) of associated fiber-fiber connections. Statistical significance was evaluated by the Wilcoxon rank sum test. Note that when pairs of cirri were grouped by fiber-fiber connection type, we did observe an increase in mutual information for cirri associated with only thin fiber-fiber connection and only thick fiber-fiber connections compared to those lacking fiber-fiber connections (see Figure S3). (D) A plot of mutual information as a function of inter-cirrus distance displays negative correlation, with a Spearman correlation coefficient of -0.49 (p<0.001). Plotted values are defined with respect to pairs of cirri. (E) A plot of mutual information as a function of fiber-cortex contact distance grouped by fiber type similarity and lack thereof displays negative correlation, with a Spearman correlation coefficient of -0.62 (p<0.001) for pairs of cirri with similar fiber types and no significant correlation for those with dissimilar fiber types. Similarity of fiber types is defined in terms of sharing at least some fiber types as defined in panel A. Fiber-cortex contact difference is measured by the mean cross nearest neighbor distance (see Method Details) for all fiber-cortex contact points associated with each cirrus. The negative correlation values from the data plotted in panels D and E indicate that cirri that are closer to one another and also cirri with fiber-cortex contacts in nearby regions of the cell tend to have higher mutual information, and indeed cirri that are both close to one another and with similar patterns of fiber-cortex contacts display the highest mutual information. (F) A plot of fiber-cortex contact difference versus inter-cirrus difference (as in panels D and E)

792   illustrates that nearby cirri tend to have similar associated fiber-cortex contacts, highlighting that

793   nearby cirri with similar fiber-cortex contacts share the most mutual information. See also Figure

794   S3.

Figure 5. The fiber system of *Euplotes* mediates gait coordination. (A-C) Representative cell

motility trajectories of cells imaged under darkfield illumination and tracked using the

TrackMate plugin in FIJI [108] in control (A), nocodazole treated (B), and paclitaxel treated (C)

cells highlight the increased turning causing curved and curled trajectories in cells treated with

the microtubule polymerization inhibitor nocodazole and decrease in turns leading to smoother,

less convoluted trajectories in cells treated with the microtubule stabilizer paclitaxel. Different

colors represent the trajectories of different cells. Scale bar is 500 μm. (D) Nocodazole affects

cell motility in a dose dependent and specific manner. Motility was quantified by the scaled path

length, which is the total integrated distance walked by a cell scaled by the maximum distance

the cell travelled from its starting point. This scaled path length decreases with a decrease in

long, straight segments of trajectories corresponding to normal cell walking. Scaled path length

decreased with increased nocodazole concentration, becoming significantly less than controls by

0.2 μM nocodazole. Note that at 2 μM nocodazole, cells were often observed to swim instead of

walk, which may account for the lack of decrease in scaled path length compared to 0.2 μM

treatment. In contrast to nocodazole, treatment of cells with 0.02 μM paclitaxel, which stabilizes

microtubules, increases scaled path length compared to the control. The trajectories of at least 20

cells were analyzed for each condition. The black bars are median values. A single asterisk

indicates $p<0.05$, and a double asterisk indicates $p<0.005$, as computed by a Wilcoxon rank-sum

test. (E,F) Representative images illustrating the effect of the inhibition of microtubule

polymerization by nocodazole on the fiber system and approach to quantifying this effect.

Images are maximum intensity projections of confocal z-stack images of cells labeled by SiR-

tubulin. Scale bar is 20 μm. (G) A plot showing that nocodazole treatment shortens fibers

compared to controls. Scaled fiber length was measured by dividing the length of the longest

fiber by the length of the entire cell. Nine cells were analyzed for each condition. The black bars

are median values. A single asterisk indicates $p<0.05$ as computed by a Wilcoxon rank-sum test.

(H) A plot of mutual information of all pairs of cirri shows that treatment of cells with the

microtubule inhibitors nocodazole and paclitaxel change the distribution of mutual information

due to an overall increase in cirral synchrony compared to untreated cells. The data for untreated

cells is that from Figures 2 and 3. Nocodazole treatment was 0.2 μM, and 6 cells over 1133

timepoints were analyzed. Paclitaxel treatment was 0.02 μM, and 6 cells over 441 timepoints were analyzed. Three asterisks indicates $p<0.001$ as computed by a two sample Kolmogorov-Smirnov test. (I) A representation of the changes in the nature and organization of functional states as well as transitions between them highlights the effects of nocodazole treatment. This panel is partially adapted from Figure 3H and was produced by following the same analysis procedure. Any functional states and transitions depicted in Figure 3H no longer observed under nocodazole treatment appear faded. Arrows that are not faded represent the new highest probability transitions associated with the states. Outer colors of the nodes corresponding to the numbered gait states denote the type or absence of broken detailed balance associated with the gait state for untreated cells while the core color represents that for the treated cells. As in Figure 3H, black denotes the lack of unbalanced transitions, red denotes the state is only a source or only a recipient of unbalanced transitions, while blue denotes states which are both sources and recipients of unbalanced transitions. Note the loss of the cycle states as well as the functional states directing return to the cycle states, the emergence of the high cirral activity state involving activity in nearly all cirri, the reduction in gait states associated with the cloud of states of substantial cell movement (indicated by the reduction in the size of the black blob), and the emergence of many high probability transitions to gait state 18. (J) A cartoon illustrates the location of the cuts made for microsurgery experiments (dashed gray line). (K) A representative image shows a recovered bisected cell and a fully intact, unperturbed cell for reference. Scale bar is 50 μm. (L) A plot of the manually tracked trajectory of a walking, bisected cell illustrates the unsteady, curving manner of cell movement. (M) A plot of mutual information of all pairs of cirri h-n for unperturbed, wounded, and bisected cells demonstrating a change in the distribution of mutual information only in the case of bisected cells where fibers associated with h-n have been severed. Three asterisks indicates $p<0.001$ as computed by a two sample Kolmogorov-

849   Smirnov test. The data for unperturbed cells is that from Figure 2, and data for wounded and

850   bisected conditions involved 10 cells over 1308 timepoints in the case of the wounded cells and

851   1815 timepoints for bisected cells. See also Figure S4, Figure S5, and Videos S3-S7.

STAR Methods

Resource Availability

Lead contact

Further information and requests for resources, data, and code should be directed to and will be

fulfilled by the lead contact, Wallace F. Marshall (wallace.ucsf@gmail.com).

Materials availability

This study did not generate new unique reagents.

Data and code availability

All data reported in this paper will be shared by the lead contact upon request.

All original code has been deposited at GitHub and is publicly available as of the date of

publication. DOIs are listed in the key resources table.

Any additional information required to reanalyze the data reported in this paper is available from

the lead contact upon request.

Experimental Model and Subject Details

<u>Cell lines</u>

Cultures of *Euplotes eurystomus* were obtained from Carolina Biological Supply Company (Item

#131480) and were kept at room temperature under ambient light conditions.

Method Details

<u>Cell husbandry</u>

Individual cells were isolated from cultures, which contained other protists and

meiofauna, by pipetting and placed in non-treated 6-well plates (Thermo Fischer Scientific 08-

772-49) containing spring water taken from cultures. Cells were kept in wells for no longer than

five days before imaging, and if cells were to be kept for longer than 48 hours, wells containing

cells were supplemented with 1% Cereal Grass Medium [109] (Thermo Fischer Scientific S25242)

to prevent depletion of prey bacteria and otherwise maintain *Euplotes* under constant growth

conditions.

<u>Live cell brightfield microscopy</u>

Cells were concentrated by centrifugation (500×g for 5 min) and resuspended either in

0.5 mL of spring water in coverglass bottomed FluoroDishes (World Precision Instruments

FD35-100) or in 0.2 mL spring water on a coverslip (FisherScientific, 12-545-D) for imaging.

No more than three cells were kept in 0.5 mL imaging samples and only one cell was ever kept

in 0.2 mL imaging samples in order to minimize cell-cell interactions. Cells were observed to

exhibit spontaneous walking activity on coverglass. Walking cells in FluoroDishes were imaged

under brightfield illumination using a Zeiss Z.1 Observer and Hamamatsu Orca Flash 4.0 V2

CMOS camera (C11440-22CU) with a 20x, 0.8 NA Plan-Apochromat (Zeiss) objective. Cells on

coverslips were imaged under brightfield illumination with coverslips inverted over a well

containing a small amount of distilled water to reduce evaporation using a Zeiss Axio Zoom.V16

and a PCO pco.dimax S1 camera. Importantly, in both imaging systems, the focal plane was set

at the interface between cirri of walking cells and the glass surface upon which they were

walking. Images were acquired at 0.033 seconds per frame with a 0.005 second exposure in order

to capture all cirral dynamics during walking with minimal blur.

<u>Quantification of walking dynamics</u>

Movies of walking cells were viewed using FIJI [110]. Movement of cirri, or lack thereof

was clearly visible in each movie frame (see Figure 2A and Video S1). The dynamical state of

each cirrus in each movie frame was manually annotated. For each frame, each cirrus received a

label of "1" if the cirrus was in motion and "0" if the cirrus was not moving and in contact with

the coverslip. Motion of cirri was evident in terms of a change in cirrus shape or tip position

often in addition to blur due to motion during image acquisition or position out of the focal plane

(see Figure 2A and Video S1). While only slowly walking cells were recorded, sometimes cells

nevertheless exhibit brief, spontaneous departures from slow walking during the course of movie

acquisition. Any frame in which the movement of the cell and/or cirri were too fast to be

resolved, such as during spontaneous escape responses [40] (Video S2), was excluded from

analysis such that some videos were split into a number of separate continuous sequences. Thus,

each movie frame associated with a particular time point in the walking trajectory, with the

exception of those excluded from analysis as described, yielded a corresponding 14-element

binary vector encoding the motility state of the cell in terms of the movement of cirri. Cell

movement was tracked using the manual tracking feature of the TrackMate plugin in FIJI [108].

The center of each cell was used as the reference feature for tracking. We analyzed the walking

dynamics of 13 different cells.

Dimensionality reduction

Dimensionality reduction was performed by non-negative matrix factorization (NMF)

implemented in MATLAB release 2019b (Mathworks, Natick). NMF was chosen as a

dimensionality reduction technique to allow us to obtain a reduced, sparse, and interpretable

representation of walking dynamics. Because NMF derives non-negative factors, the basis

vectors in NMF space correspond straightforwardly to patterns of cirral activity. NMF involves

factoring data, A, an n by m matrix, into non-negative factors W, an n by k matrix, and H, a k by

m matrix where the product $W*H$ approximates A. To determine the appropriate number of

dimensions or rank, k, that are necessary to accurately represent the data without overfitting, we

performed cross-validation by imputation with random holdouts [111,112], also implemented in

MATLAB. We randomly held out 15% of our walking dynamics data, performed NMF for a

given k, and then used the NMF reconstruction $W*H$, to update the missing data entries. This

process of updating is known as imputation, and we repeated the imputation process 50 times, by

which point the imputed values had stabilized, to obtain a final NMF reconstruction. We then

computed the root mean squared residual (RMSR) between the final NMF reconstruction, $W*H$,

and our dataset, A. We performed this entire process 100 times for each value of k. As is

generally the case for NMF, we observed a monotonic decrease in reconstruction error with

increasing k without performing the imputation procedure [113] (Figure S1). In contrast to this

trend, we observed an increase in RMSR of imputed values with increasing k indicating

overfitting [111] (Figure S1). We chose $k=3$ because this value was the highest value before a

notable increase in imputation error (Figure S1), which would indicate overfitting [111,112]. Thus,

our choice of rank 3 selects the lowest rank approximation that captures structure of the dataset

without overfitting that structure. Further, our choice facilitated the visual inspection of the

structure of data in the reduced dimensional reconstruction.

Finally, we noted that for our chosen value of k, due to the stochastic nature of the NMF

algorithm, which involves a random initialization step, we obtained slightly different solutions

for different iterations [111]. In order to choose the best reduced dimensional approximation,

therefore, we performed NMF 500 times and chose the particular solution corresponding to the

lowest RMSR compared to our dataset.

Clustering

949    Clustering on the dataset obtained using NMF was performed by density-based spatial

950 clustering of applications with noise (DBSCAN) algorithm [70] implemented in MATLAB release

951 2019b (Mathworks, Natick). Structure in NMF space was clearly visible (Figure 2E-G), and

952 DBSCAN using a Euclidean distance metric, was initially chosen as a clustering method because

953 it yielded qualitatively good partitioning of the data. The DBSCAN algorithm involves stochastic

954 search within neighborhoods of a given radius ε around datapoints, and points with a minimum

955 number of neighbors, n, within their neighborhood are grouped as belonging to the same cluster,

956 leaving two free parameters to determine. We set ε by first using the

957 clusterDBSCAN.estimateEpsilon function in MATLAB (release 2020b, Phased Array System

958 Toolbox), which yielded a value of 0.15. We next set about determining the minimum neighbor

959 number, n. To do so, we computed the average Silhouette coefficient, a commonly used measure

960 of clustering quality that indicates how well-separated clusters are [114], for various values of n.

961 The results of this analysis are plotted in Figure S1. Higher Silhouette coefficients indicate better

962 clustering, and we found that a value of $n=8$ maximized the mean Silhouette coefficient (Figure

963 S1). We also noted, however, that for this value, many datapoints were found to be outliers, not

964 belonging to any cluster due to having too few points within a distance of ε. Figure S1 displays

965 percentage of datapoints found to be outliers as a function of n. In order to avoid categorizing

966 more than 5% of datapoints as outliers, we chose to settle on $n=4$, which does not have a

967 significantly different mean Silhouette coefficient compared to any of the others in the range

968 $n=2\text{-}7$. This choice was further supported by the fact that major clusters involving more than 5

969 datapoints identified with $n=8$ were also identified with $n=4$.

970    Although this set of parameters gave qualitatively and quantitatively reasonable

971 clustering results, we sought to further refine our clusters and to further reduce the outlier

972 datapoints. We noted the obvious partitioning of the NMF dataset into three groups along the H2

axis (Figure 2E). We found the previously determined parameter values to yield good clustering

for the top and middle partitions (H2≤1.1 and 0.2<H2<1.1), with no outliers. For the lower

partition (H2≤0.2), however, we found that we were able to improve clustering by using

$\varepsilon=0.1182$. With this updated value, we found no statistically significant change in Silhouette

coefficient and reduced outliers to 0%. The clusters obtained by this process constituted the

identification of the 32 gait states. We note here that the problem of determining the true or

optimal number of clusters is an unresolved problem [115], and we note that we have followed

standard methods to determine cluster number, and we found that our key results do not depend

sensitively on the precise number of clusters identified (see following section and Figure S1 for

more details).

<u>State transition analysis</u>

Following dimensionality reduction and clustering to identify gait states, we proceeded to

characterize state transition dynamics. For each cell trajectory, we identified all unique gait state

transitions for a total of 1423 unique pairwise transitions over the cumulative 2343 video frames

for 77.14 s of recording. We computed empirical transition rates between states as the total

number of observed transitions divided by the total time of observation. In order to determine

which transitions were balanced and which were unbalanced, we followed Chang and Marshall

[45], and performed binomial tests of statistical significance. Assuming a system at equilibrium,

with all transitions obeying detailed balance, we expect to observe some deviation from exactly

reciprocal transitions and can calculate the probability of observing a given set of ratios given

underlying probabilities of forward and reverse transitions. The binomial probability of

observing a set of transitions with known forward and reverse probabilities is given by

$$P(X = f) = \binom{n}{f} p_{forward}^{f} p_{reverse}^{n-f}$$

where $\binom{n}{f} = \frac{n!}{f!(n-f)!}$ is the number of combinations, f is the number of forward transitions, n is

the total number of transitions (such that n-f is the number of reverse transitions), and the

probabilities $p_{forward}$ and $p_{reverse}$ are the forward and reverse probabilities. Considering only

the set of transitions involving a specific pair of states, and calculating the probability that a

transition between those states is either in the forward or reverse direction, the values of forward

and reverse probabilities in the balanced case must be equal such that $p_{forward} = p_{reverse} = 0.5$.

With an α level of 0.05, we then considered reciprocal transition pairs with binomial

probabilities less than 0.05 to be significantly unbalanced. Figure S1 displays the binomial

probabilities associated with all transitions.

In order to calculate the estimated entropy production rate, we followed Wan and

Goldstein [5], where the entropy production rate is defined as

$$\dot{S} = \frac{1}{2} \sum_{i \neq j} J_{ij} A_{ij}$$

1008

with conjugate fluxes $J_{ij} = p_i k_{ij} - p_j k_{ji}$ and forces $A_{ij} = \ln \left(\frac{p_i k_{ij}}{p_j k_{ji}} \right)$ where the p_l are the

probabilities of being in state l at steady state and the k_{ij} are the transition probabilities from

states i to j. We estimate the state occupancy probabilities p_l as $\frac{T_l}{T_{Total}}$, where T_l is the amount of

time spent in state l over all trajectories and T_{Total} is the total recorded time, and the transition

probabilities k_{ij} as $\frac{N_{ij}}{N_i}$, where N_{ij} is the total number of observed transitions from state i to state j

and N_i is the total number of transitions emanating from state i. To avoid $k_{ji} = 0$ for pairs of

states for which we did not observe any transitions during our experiments, we let $k_{ji} = \frac{1}{p_j T_{max}}$

where $T_{max} = 238$ is the maximum number of observed transitions for any single recorded

walking trajectory.

In the course of our state transition analysis, we also checked whether the waiting times between instances of each state might be non-exponentially distributed, with exponential distributions indicative of an embedded Markov process or possibly self-organized criticality [116,117]. Using the Lilliefors test implemented in MATLAB, we found that in general, waiting times were not exponentially distributed, although states 2, 3, 6, 16, 17, 18, 25, 27, 28, 32 were found to have waiting times consistent with exponential distributions with Benjamini-Hochburg corrected p-values of 0.046, 0.046, 0.022, 0.008, 0.046, 0.017, 0.046, 0.0081, 0.0046, 0.0046 respectively. Interestingly, none of the waiting times between the movements of individual cirri were found to be consistent with exponential distributions. These results are consistent with mechanisms constraining the temporal dynamics of cirri and state transitions.

In order to begin evaluating whether state transitions obeyed the Markov property for a discrete-time, first-order Markov process, where the transition probabilities from one state to the next are completely determined by current state [75,76], we estimated the transition matrix for walking dynamics, consisting of the transition probabilities between all states. We estimate the transition probability from state i to state j as $k_{ij} = \frac{N_{ij}}{\sum_k N_{ik}}$ such that $\sum_i k_{ij} = 1$. The entries of the transition matrix, P, are these transition probabilities with indices i for rows and j for columns. If gait state transitions obeyed the Markov property, we expect that the product of the transition matrix with itself, P^2, would be equivalent to the two-step transition matrix where transition probabilities are computed as before except that state j is the state to which i has transitioned after an intervening transition. Figure S1 displays the results of this analysis showing that the two matrices show some quantitative and qualitative differences. Although these results strongly suggest violation of the Markov property, we applied the Billingsley test for a more statistically rigorous evaluation [118,119]. This test was implemented and performed in MATLAB. The Billingsley test gives a χ^2 metric with M^2-$2M$ degrees of freedom given by

$\sum_{i=1}^{M} \sum_{j=1}^{M} \frac{\left(N_{ij} - R_{ij} \sum_{j=1}^{M} N_{ij}\right)^2}{R_{ij} \sum_{j=1}^{M} N_{ij}},$

where R_{ij}, the independent trials probability matrix, is given by

$R_{ij} = \sum_{k=1}^{M} N_{kj} / (\sum_{h=1}^{M} \sum_{l=1}^{M} N_{hl} - \sum_{k=1}^{M} N_{ik})$. Applying this test to our gait state transition data,

we found that the null hypothesis that the gait conforms to a first-order discrete time Markov

process was rejected (p=0.005).

Importantly, we also noted that the key qualitative results of our state transition analysis

are robust to the details of clustering results. In particular, we find that strongly unbalanced

transitions and violation of the Markov property exist for a range of clustering parameters.

Figure S1 displays the transition matrices for different clustering results.

To arrive at the simplified, state machine representation of the gait, we focused on the

highest probability transitions emanating from each state. Transition probabilities were estimated

as k_{ij} (as defined above). This allowed us to prune away rare transitions in order to reveal the

dominant structure of gait state transitions. Figure 3G displays the pruned transition matrix as a

heatmap. We found that relatively few states were the recipients of the majority of high

probability transitions, and many states received none. To more clearly visualize the structure of

transitions, we grouped together all gait states receiving no more than one unique high

probability transition based on the idea that state transitions into this group show little bias in

terms of source state, and within the group, transitions between states exhibit low probability,

time unbiased, equilibrium-like fluctuations.

1061

1062   <u>Biophysical model and simulations</u>

1063   For our simple biophysical model, we consider a 2D system in which a *Euplotes* cell is

1064   walking across a surface in a low Reynolds number environment [43]. The cell has 14 cirri, which

exist in one of two states: actively moving or not actively moving, following our quantitative gait

characterization. Cell position and orientation is defined in terms of equilibrium position of the

cirri. Cirri can generate a motive force to drive cell motility when moving and resist

displacement when not moving and in contact with the surface. For our model, we remain

agnostic to the details by which cirri produce generate force noting only that in our experiments,

no cell displacement was observed when cirri were not moving and in contact with the coverslip.

We therefore let cirri generate a constant force in the direction dictated by cell orientation when

moving. We conceptualize the resistance to displacement of unmoving cirri as stemming from

the adhesive interaction between the cirrus and the substrate on which the cell is walking and the

energy required to bend or deflect a cirrus. Consistent with experimental observations, we do not

allow for translation of a cirrus-substrate contact point while a cirrus is not actively moving.

For a particle moving through a fluid at low Reynolds number, such as our cell, velocity

$\vec{v}$ will be given by

$$\vec{v} = \frac{\sum \vec{F}}{\gamma},$$

where $\sum \vec{F}$ is the sum of the external forces acting on the particle, and γ is constant related to the

geometry of the particle and the viscosity of the fluid [120,121] accounting for drag. In our model,

cirri are responsible for the forces involved in motility, so $\sum \vec{F} = \vec{F_a} + \vec{F_i}$ with the motive force

$\vec{F_a} = nF_a\vec{o}$ where n is the number of active cirri, F_a is the magnitude of the constant force

generated by active cirri, and $\vec{o}$ is the unit vector in the direction of cell orientation, and the

resistive force $\vec{F_i} = \sum_i k\vec{d_i}$ is a sum over the inactive cirri where k is a constant controlling the

resistance of a cirrus to deformation and $\vec{d_i}$ is the displacement vector of inactive cirrus i. We

note that the forces driving cell motility in *Euplotes* stem from complex mechanical interactions,

but for our model, we have chosen simple, first order expression to capture very basic features.

Similar to the expression for velocity above, angular velocity of a walking cell in our

model is given by

$$\vec{\omega} = \frac{\sum \vec{F} \times \vec{r}}{\beta},$$

where β is a constant related to the geometry of the cell and viscosity of the fluid, $\vec{F_i}$ is the force

due to cirrus i ($F_a\vec{o}$ for active cirri and $k\vec{d}$ for inactive cirri), and $\vec{r_i}$ is the vector pointing from

the center of the cell to cirrus i.

In addition to the relative positions of the cirri and patterns of cirral activity, the four

parameters γ, β, F_a and k govern cell motility in our model. From these four parameters, we

obtain three related dimensionless parameters: $\alpha = \beta/\gamma l$ where l is the maximum distance

between cirri, which can be thought of as characterizing the unsteadiness of the cell the intrinsic

susceptibility of the cell to turning due to cell geometry; $M = F_a/kl$, which can be thought of as

characterizing degree to which cirral activity will tend to induce cell movement in opposition to

inactive cirri; and $\mathcal{F} = F_a t/\gamma l$ where t is the duration of a timestep in simulations, which can be

thought of as the strength of the cirral motive force relative to the viscous drag experienced by

the cell due to its fluid environment.

For all simulations, relative equilibrium cirral positions calculated from the average cirral

positions in a video frame with no cirral activity over the 13 cells used for gait analysis (Figure

2) were used. In each simulation timestep, defined by the timestep used for recording videos for

gait analysis, velocity and angular velocity are calculated based on the positions and activity of

cirri, and the positions of all cirri are updated accordingly before proceeding. To calibrate the

parameters of the model, we used the cirral patterns recorded from the cells used for gait

analysis. We swept parameter space and found that simulations qualitatively and semi-

quantitatively recapitulated experimentally measured cell motility with $\alpha = 0.001$, M $= 0.26$,

and $\mathcal{F} = 0.008$ (Figure 3K, Figure S1). These parameter values were subsequently used for all

simulations.

For simulations with shuffled gait state transitions, we used MATLAB's shuffle

command on the cirral dynamics of actual cells to obtain sequences of gait state transitions of the

same length as those that were experimentally obtained except in a random order. To obtain

random patterns of cirral activity similar to those measured experimentally, we generated cirral

activity according to a process defined by two probabilities: p_a, the probability of transitioning

from inactive to active at each timestep and p_i, the probability of transitioning from inactive to

active at each timestep. We initialized sequences with no cirral activity and then updated cirral

activity according to these probabilities for each timestep in the sequence. We found that setting

p_a=0.3 and p_i=0.1 yielded the same average cirral activity as that recorded experimentally, $0.23 \pm$

0.42 per frame and 0.24 ± 0.43 per frame respectively where the values are mean $\pm$ standard

deviation. All simulations were performed in MATLAB.

1124

Confocal microscopy

Cells were prepared for imaging and placed into a FluoroDish as described in the Live

Cell Brightfield Microscopy section. Cells were then labeled with SiR-tubulin (Spirochrome

provided by Cytoskeleton, Inc, CY-SC002) at 1 µM concentration. Cells were imaged using a

Zeiss LSM 880 AxioExaminer and a 40x, 1.2 NA C-Apochromat water immersion objective

(Zeiss) and excitation provided by a 633 nm laser (Zeiss). Only one full confocal z-stack of a

complete cell was obtained during imaging to avoid effects of photodamage.

1132

1133 Fiber reconstruction and analysis

The image stack resulting from confocal imaging was first aligned in FIJI using the

StackReg plugin [110]. Next, fibers were manually segmented in each of the aligned z-stack images

using the TrakEM2 plugin in FIJI [122,123]. Thick and thin fibers (Figure 4A) were morphologically

distinguished, with thick fibers having a diameter of no less than 5 μm at the thinnest point.

Fibers were traced from their distal termini to their convergences at the base of the cirri with

which they were associated. Following segmentation, 3D surfaces were reconstructed in

TrakEM2. Inter-fiber contacts were then found by inspection of 3D reconstructions and verified

by examining individual z-stack frames to confirm intersections between fibers.

Drug treatment experiments and analysis

For all cytoskeleton inhibitor treatment experiments, 1 mL of cells in culture were placed

in wells of 12-well plates (Thermo-Scientific, 12-565-321). Nocodazole (Sigma-Aldrich,

M1404) as a stock solution of 6.64 mM in DMSO diluted further in distilled water, was added to

achieve appropriate concentrations, with no more than 1 μL of additional volume added, and 1

μL of distilled water with equivalent DMSO concentration to nocodazole treatments added to

controls. Cells were incubated for 1 hr before initiating experiments. No cell death was observed

at any concentration of nocodazole in the 6 hours following nocodazole treatment. Paclitaxel

(Sigma-Aldrich, T7191) as a stock solution of 2.23 mM in DMSO diluted further to 20 μM in

distilled water, was added to achieve a final concentration of 20 nM to cells in solution.

Latrunculin B (Thomas Scientific, C834E37) as a stock solution of 1.1 mM in ethanol was

further diluted in distilled water and added to achieve a final concentration of 10 μM with cells

in solution. Cytochalasin B (Fisher Scientific 1493-96-2) as a stock solution of 2.1 mM in

ethanol was further diluted in distilled water was added to achieve a final concentration of 50

μM with cells in solution. Jasplakinolide (Fisher Scientific 42-012-750UG) as a stock solution of

1 mM in DMSO was further diluted in distilled water to achieve a final concentration of 10 µM

with cells in solution. No cell death was observed in the 6 hours following treatment with any of

the actin inhibitors. For the control condition for actin inhibitor experiments, both DMSO and

ethanol was added to match the concentrations added in the cytochalasin B and jasplakinolide

conditions.

Washout experiments

For nocodazole washout experiments, motility assays (described below) were also

performed after placement of cells into well plates and before nocodazole treatment. Following

0.2 µM nocodazole treatment and another motility analysis, cells were picked in 5 µL of media

and placed into wells of 6 well plates (Corning, CLS3736) each containing 2mL of fresh media.

No more than 20 µL of nocodazole treatment media was added to any well so that the resultant

nocodazole concentration in the washed condition was no more than 2 nM. Cells were allowed to

recover in this condition for 4 hr, and then a final motility assay was performed.

Motility analysis

For motility analysis, cells were picked from well-plates and placed onto well-slides

created by using a paper hole punch tool to punch a hole in 0.25 mm thick silicone spacer

material (CultureWell, 664475), and adhering the spacer material to a glass slide (Corning, 2947-

75X25). A total volume of 20 µL including up to 4 cells was added to well slides. A glass

coverslip (FisherScientific, 12-545-D) was placed atop the well slide, sealing the well and

creating an imaging chamber. After creating the imaging chamber, cells were allowed to

acclimate to their new environment for 10 min. Cells were then imaged on a Zeiss Axio

Zoom.V16 microscope under darkfield illumination with a Canon EOS T5i DSLR camera

recording at 30 fps for 2 min. Videos were then processed using FIJI [110]. First, images were

cropped to remove extraneous parts of the field of view that did not contain the imaging

chamber, and then background subtraction was performed by creating an image composed of

mean pixel intensity values over all frames of the video and subtracting this mean image from all

frames of the video. A mean filter with a four-pixel radius was then applied to each frame of the

video for the purpose of smoothing. After processing, tracking was performed using the

TrackMate FIJI plugin [108]. For detection of objects (cells), a Laplacian of Gaussian filter was

applied with an estimated blob diameter of 25 pixels and threshold of 0.1. A quality threshold

was set manually when necessary to filter out any detected objects that were not cells. The

Linear Assignment Tracker with a linking max distance of 15 pixels, gap-closing max distance

of 150 pixels, and a gap-closing max frame gap of 200 frames was then used to generate linked

tracks (trajectories) of detected cells. Trajectories were analyzed in MATLAB. Scaled path

length for each tracked cell was calculated by summing the length of all segments of the track

and dividing by the maximum distance the cell traveled from its starting point.

Fiber length analysis

Just prior to confocal imaging, cells were washed by picking up to five cells in 10 μL

volume each and placing into 1 mL fresh media in a 12-well plate (Thermo-Scientific, 12-565-

321). Cells were then prepared for imaging as described in the Confocal Imaging section of

Method Details. Confocal z-stacks were then loaded into FIJI and aligned using the StackReg

plugin [110]. Cell lengths were determined by finding the maximal distance between two points on

the front and rear ends of the cell. Because of variability in the detectable fibers in nocodazole

treated cells, only fibers associated with the rear cirri (j-n), which were visible in all cells, were

used for analysis. All of these rear fibers were measured, and the reported scaled fiber length was

obtained by dividing the length of the longest fiber by the corresponding cell length. In all cases,

the fiber associated with cirrus m was the longest fiber.

Analysis of cirral dynamics in nocodazole and paclitaxel treated cells

Cells were prepared for imaging and imaged as described in the Live Cell Brightfield

Microscopy section of the Method Details with the exception that a Canon EOS T5i DSLR

mounted on a Zeiss Axio Zoom.V16 microscope was used to record movements. Additionally,

video was recorded at 0.066 seconds per frame to avoid blurring and then videos were

downsampled to 0.033 seconds per frame for analysis. Cirral dynamics were quantified as

described in the Quantification of Walking Dynamics section of the Method Details.

To assign cirral configurations of nocodazole treated cells to previously identified gait

states, we first matched any cirral configurations with known gait state identity. Next, due to

proximity of new cirral configurations not observed in untreated cells (Figure S1), we were able

to map the new cirral configurations onto the clusters defining the gait states by determining the

nearest cluster to the new cirral configuration. Distance between new cirral configurations and

clusters were determined by finding the shortest distances to points defining convex hulls of each

cluster. The shortest of all these distances then indicated the nearest cluster and corresponding

gait state to which the new cirral configuration was assigned. We noted an obvious dense cluster

of points corresponding to activation of nearly all cirri, and we identified a new cluster and

corresponding gait state by applying the DBSCAN algorithm as described in the Clustering

section of the Method Details. Evaluation of transition dynamics was performed as described in

the State Transition Analysis section of the Method Details. This analysis was all performed in

MATLAB.

1226

ϵ-machine construction

Our representation of the Euplotes at timestep t takes the form of a length 14

binary string $x^t = [x_1^t, x_2^t, \ldots, x_{14}^t]$ where $x_i^t \in \{0, 1\}$. For reducing dimensions,

we found that bigrams of the 14-dimensional strings, yielded more consistent, interpretable

results than unigrams, so $x'^t = [x_1^t, x_2^t, \ldots, x_{14}^t, x_1^{t+1}, x_2^{t+1}, \ldots, x_{14}^{t+1}] \in \{0,1\}^{28}$.

In order to learn latent states, we used Variational AutoEncoders (VAEs) [124] to reduce each

fourteen dimensional timestep to three dimensions. VAEs in particular are used for their ability

to learn an interpolatable latent space where high-dimensional training data points are mapped to

points in low-dimensional space that mimic a normal distribution centered around the origin.

This processes yields a smooth latent space with dimensions that represent core aspects of the

data. We used a very minimal VAE with an Adam optimizer [125], consisting of one dense layer

for the generating encoder means and one dense layer for generating encoder variance. This

creates a 3-dimensional normal-like distribution, which we sample from and decode with a one

layer decoder.

$\text{VariationalEncoder}(x'^t) \rightarrow z^t \in \mathbb{R}^3$

After reducing the dimensionality of our data, we perform spectral clustering [126] using

scikit-learn [127] with 100 k-means runs and 4 clusters, where the cluster count was decided by

inspection of the resulting classes.

$\text{SpectralClustering}(z^t) \rightarrow c^t \in \{1, 2, 3, 4\}$

Applying this procedure to the data from the cells used for gait analysis (the same as used for

Figure 2) with bigram training parameters lr=0.0011, weight_decay=0.000727, k=3,

hidden_dim=5, beta1=0.5, and beta2=0.900, we found the following four clusters corresponding

to particular patterns of cirral ativity: 1) Front cirral activity 2) High overall cirral activity 3)

Rear cirral activity and 4) Little cirral activity (Figure S5). When we applied this procedure to

the data from cells treated with 0.2 µM nocodazole (the same data used for Figure 5) with

bigram training parameters lr=0.0026, weight_decay=0.000106, k=3, hidden_dim=3, beta1=0.5,

and beta2=0.999, we obtained only three distinct clusters with similar cirral activity patterns to

those of untreated cells except that the rear cirral activity cluster (3 above) disappeared (Figure

S5). These clusters represent an independent, alternative coarse-graining of patterns of cirral

activity into gait states to that presented in Figure 2. Although independent, this procedure

identified the same underlying structure in the data. In particular, the alternative gait states here

involve similar patterns of cirral activity to those defining the NMF components depicted in

Figure 2H. Whereas the spatially distinct groups of cirri sharing the most mutual information

define the space of gait states under the NMF-based approach, under this alternative, activity or

complete lack thereof in those same groups of cirri defines the gait states. Furthermore, the loss

of cluster 4 under nocodazole treatment is consistent with the loss of gait states as depicted in

Figure 5I.

To construct ϵ-machines [79], characterized by a set of causal-states and transitions between

them, we feed a stream of behavioral symbols $c_0, c_1, . . ., c_T$ into a Causal-State Splitting

Reconstruction (CSSR) [89,128] algorithm. Due to the nature of the CSSR algorithm, we can only

construct ϵ-machines from unbroken data streams, so we constructed ϵ-machines for each cell

from a given dataset. Performing this process over gait state data streams derived from cirral

activity recordings as described above for each cell from the untreated and nocodazole datasets

yielded a set of ϵ machines. Representative examples of ϵ-machines obtained from each dataset

are depicted in Figure S5.

Microsurgery experiments

All microsurgeries were performed by hand under observation with a Zeiss Axio

Zoom.V16 using quartz microneedles pulled from quarts rods (Sutter Instrument, QR100-10)

using a Sutter Instrument P-2000 Laser-based micropipette puller. Individual cells were picked

and placed in a 5 μL droplet at the center of an imaging chamber as described in the Motility

Analysis section of the Method Details. For bisections, microneedles were quickly and firmly pressed onto cells just anterior to cirrus h, cleanly severing the cell including the fibers associated with cirri j-n and sealing the cell membrane. Cells were allowed to recover for at least 5 minutes until some motility resumed, and then 15 µL of fresh culture media was added to the imaging chamber, which was subsequently sealed with a glass coverslip (FisherScientific, 12-545-D) to minimize evaporation. After 24 hr, once cell fragments resumed walking motility, gait analysis was performed as described in the quantification of walking dynamics section of the Method Details, except that cell fragments only had 7 cirri (j-n) instead of the full 14. Experiments with wounded cells were conducted in the same fashion as described for bisections except that instead of cutting cells in half, wounding was performed by stabbing with a microneedle a portion of the cell that does not have any fibers and was confirmed visually by the loss of some amount of cytoplasm.

Quantifications and statistical analysis

Statistical details of the experiments can be found in the figure legends, the main text, or the Method Details section. Statistical details include exact value of n, what n represents (generally the number of cells measured), definitions of center, and dispersion and precision measures. Statistical tests and fits were performed as described in the Method Details section using MATLAB release 2019b or 2020b (Mathworks, Natick).

Supplemental Video Captions

Video S1. **A walking cell.** A representative example of a single *Euplotes eurystomus* cell walking across a coverslip using motile cirri illustrates the walking gait. Note that the movement of cirri, or lack thereof is clearly visible in each frame of the movie. This cell was imaged using a

Zeiss Axio Zoom.V16 microscope under brightfield illumination. The video is slowed down by a factor of 4 from real time to show that the movements of cirri are clearly visible.

Video S2. **Walking motility including escape responses.** A representative example of a single *Euplotes eurystomus* cell walking across a coverslip using motile cirri including escape responses, which were excluded from gait analysis. Escape responses occur at t= 18s, 20s, and 25s. This cell was imaged under brightfield illumination using a Zeiss Z.1 Observer with a 20x, 0.8 NA Plan-Apochromat (Zeiss) objective. The video is slowed down by a factor of 4 from real time so that the movements are clearly perceivable.

Video S3. Cell tracking for motility analysis. A representative example of the trajectories of cells imaged at low magnification under darkfield illumination using a Zeiss Axio Zoom.V16 microscope. Cells appear as bright objects with magenta circles around them. Different colored lines correspond to the tracks of different cells tracked using the FIJI plugin TrackMate.

Video S4. Motility of nocodazole treated cells. A representative example of the trajectories of cells treated with 200 nM nocodazole imaged at low magnification under darkfield illumination. Cells appear as bright objects with magenta circles around them. Different colored lines correspond to the tracks of different cells tracked using the FIJI plugin TrackMate. Note the confined movements of cells due to a decrease in long, linear runs. Compare to Movie 3 for the control condition.

Video S5. Motility of paclitaxel cells. A representative example of the trajectories of cells treated with 20 nM paclitaxel imaged at low magnification under darkfield illumination. Cells

appear as bright objects with magenta circles around them. Different colored lines correspond to the tracks of different cells tracked using the FIJI plugin TrackMate. Note the increase in long, linear runs and decrease in abrupt changes in direction and tight turns. Compare to Movie 3 for the control condition.

Video S6. A walking, bisected cell. A representative example of a bisected cell analyzed in microsurgery experiments displaying the characteristic backwards, spiral walking motility defect. This cell fragment was imaged using a Zeiss Axio Zoom.V16 microscope under brightfield illumination. The video is slowed down by a factor of 4 from real time to clearly show the movements of cirri.

Video S7. A walking, wounded cell. A representative example of a wounded cell analyzed in microsurgery experiments. This cell was imaged using a Zeiss Axio Zoom.V16 microscope under brightfield illumination. The video is slowed down by a factor of 4 from real time to clearly show the movements of cirri.

Acknowledgements

We would like to thank Nicole King for the use of the confocal microscope and Adair Oesterle for assistance with microsurgery. We would like to thank the students and faculty of the 2016 Marine Biological Laboratory Physiology Course and current and former members of the Marshall Lab and King Lab for comments, critiques, and encouragement during the development of this project. We would also like to thank David Booth, Greyson Lewis, Dennis Bray, and members of the Fourmentin-Guilbert Scientific Foundation for comments on the manuscript. This work was funded by the I2CELL Seed Award of the Fourmentin-Guilbert Scientific

1350 Foundation (WFM, JBP). Additional funding was provided by NIH grant R35 GM130327

1351 (WFM), NSF grant MCB- 2012647 (WFM), and a Merck Fellowship of the Jane Coffin Childs

1352 Memorial Fund for Medical Research (BTL).

1353

1354 **Author Contributions**

1355 BTL, JDP, and WFM conceived of the study. BTL, JG, and WFM developed methodology. BTL

1356 performed the experiments, and all authors analyzed and interpreted data. BTL and WFM

1357 originally drafted the manuscript, and all authors reviewed and edited the work.

1358

1359 **Declaration of Interests**

1360 The authors declare no competing interests.

1361

References

1. Jennings, H.S. (1906). Behavior of the lower organisms. (Columbia University Press).

2. Reid, C.R., MacDonald, H., Mann, R.P., Marshall, J.A.R., Latty, T., and Garnier, S.

(2016). Decision-making without a brain: how an amoeboid organism solves the two-

armed bandit. J. R. Soc. Interface *13*, 20160030.

3. Coyle, S.M., Flaum, E.M., Li, H., Krishnamurthy, D., and Prakash, M. (2019). Coupled

Active Systems Encode an Emergent Hunting Behavior in the Unicellular Predator

Lacrymaria olor. Curr. Biol. *29*, 3838-3850.e3.

4. Battle, C., Broedersz, C.P., Fakhri, N., Geyer, V.F., Howard, J., Schmidt, C.F., and

MacKintosh, F.C. (2016). Broken detailed balance at mesoscopic scales in active

biological systems. Science *352*, 604–7.

5.   Wan, K.Y., and Goldstein, R.E. (2018). Time Irreversibility and Criticality in the Motility

of a Flagellate Microorganism. Phys. Rev. Lett. *121*, 58103.

6.   Woznica, A., Cantley, A.M., Beemelmanns, C., Freinkman, E., Clardy, J., and King, N.

(2016). Bacterial lipids activate, synergize, and inhibit a developmental switch in

choanoflagellates. Proc. Natl. Acad. Sci. U. S. A. *113*, 7894–7899.

7.   Wan, K.Y. (2018). Coordination of eukaryotic cilia and flagella. Essays Biochem. *62*,

829–838.

8.   Nomura, M., and Ishida, K.I. (2016). Fine-structural Observations on Siliceous Scale

Production and Shell Assembly in the Testate Amoeba Paulinella chromatophora. Protist

167, 303–318.

9.   Bennett, R.R., and Golestanian, R. (2015). A steering mechanism for phototaxis in

Chlamydomonas. J. R. Soc. Interface *12*, 20141164–20141164.

10.   Machemer, H. (2001). The swimming cell and its world: Structures and mechanisms of

orientation in protists. Eur. J. Protistol. *37*, 3–14.

11.   Almagor, M., Ron, A., and Bar-Tana, J. (1981). Chemotaxis in Tetrahymena thermophila.

Cell Motil. *1*, 261–268.

12.   Berg, H.C., and Brown, D.A. (1972). Chemotaxis in Escherichia coli analysed by three-

dimensional tracking. Nature *239*, 500–504.

13.   Nakagaki, T., Yamada, H., and Tóth, Á. (2000). Maze-solving by an amoeboid organism.

Nature *407*, 470–470.

14.   Tweedy, L., Thomason, P.A., Paschke, P.I., Martin, K., Machesky, L.M., Zagnoni, M.,

and Insall, R.H. (2020). Seeing around corners: Cells solve mazes and respond at a

distance using attractant breakdown. Science (80-.). *369*.

15.   Applewhite, P.B. (1979). Learning in Protozoa. In Biochemistry and Physiology of

Protozoa (Elsevier), pp. 341–355.

16.   Dexter, J.P., Prabakaran, S., and Gunawardena, J. (2019). A Complex Hierarchy of

Avoidance Behaviors in a Single-Cell Eukaryote. Curr. Biol. *29*, 4323-4329.e2.

17.   Jennings, H.S. (1902). Studies on reactions to stimuli in unicellular organisms. IX—On

the behavior of fixed infusoria (Stentor and Vorticella), with special reference to the

modifiability of protozoan reactions. Am. J. Physiol. Content *8*, 23–60.

18.   Tang, S.K.Y., and Marshall, W.F. (2018). Cell learning. Curr. Biol. *28*, R1180–R1184.

19.   Lyon, P. (2015). The cognitive cell: Bacterial behavior reconsidered. Front. Microbiol. *6*.

20.   Marshall, W.F. (2019). Cellular Cognition: Sequential Logic in a Giant Protist. Curr. Biol.

29, R1303–R1305.

21.   Greenfield, D., McEvoy, A.L., Shroff, H., Crooks, G.E., Wingreen, N.S., Betzig, E., and

Liphardt, J. (2009). Self-Organization of the Escherichia coli Chemotaxis Network

Imaged with Super-Resolution Light Microscopy. PLoS Biol. *7*, e1000137.

22.   Wan, K.Y., and Goldstein, R.E. (2016). Coordinated beating of algal flagella is mediated

by basal coupling. Proc. Natl. Acad. Sci. U. S. A. *113*, E2784-93.

23.   Höfer, T., Sherratt, J.A., and Maini, P.K. (1995). Dictyostelium discoideum : cellular self-

organization in an excitable biological medium. Proc. R. Soc. London. Ser. B Biol. Sci.

259, 249–257.

24.   Kantsler, V., Dunkel, J., Polin, M., and Goldstein, R.E. (2013). Ciliary contact interactions

dominate surface scattering of swimming eukaryotes. Proc. Natl. Acad. Sci. U. S. A. *110*,

1187–1192.

25.   Bray, D. (2009). Wetware (Yale University Press).

26.   Pfeifer, R., Lungarella, M., and Iida, F. (2007). Self-organization, embodiment, and

biologically inspired robotics. Science (80-.). *318*, 1088–1093.

27.   Regev, A., and Shapiro, E. (2002). Cellular abstractions: Cells as computation. Nature

419, 343–343.

28.   Mitchell, M. (2012). Biological Computation. Comput. J. *55*, 852–855.

29.   Navlakha, S., and Bar-Joseph, Z. (2011). Algorithms in nature: the convergence of

systems biology and computational thinking. Mol. Syst. Biol. *7*, 546.

30.   Savin, T., Kurpios, N.A., Shyer, A.E., Florescu, P., Liang, H., Mahadevan, L., and Tabin,

C.J. (2011). On the growth and form of the gut. Nature *476*, 57–62.

31.   Mehta, P., and Schwab, D.J. (2012). Energetic costs of cellular computation. Proc. Natl.

Acad. Sci. U. S. A. *109*, 17978–17982.

32.   Berg, H.C., and Purcell, E.M. (1977). Physics of chemoreception. Biophys. J. *20*, 193–

219.

33.   Bialek, W., and Setayeshgar, S. (2005). Physical limits to biochemical signaling. Proc.

Natl. Acad. Sci. U. S. A. *102*, 10040–5.

34.   Andrews, L.B., Nielsen, A.A.K., and Voigt, C.A. (2018). Cellular checkpoint control

using programmable sequential logic. Science (80-.). *361*.

35.   Gilbert, J.J. (1994). Jumping behavior in the oligotrich ciliates Strobilidium velox and

Halteria grandinella, and its significance as a defense against rotifer predators. Microb.

Ecol. *27*, 189–200.

36.   Lueken, W., Ricci, N., and Krüppel, T. (1996). Rhythmic spontaneous depolarizations

determine a slow-and-fast rhythm in walking of the marine hypotrich Euplotes vannus.

Eur. J. Protistol. *32*, 47–54.

37.   Erra, F., Iervasi, A., Ricci, N., and Banchetti, R. (2001). Movement of the cirri during the

creeping of Euplotes crassus (Ciliata, Hypotrichida). Can. J. Zool. *79*, 1353–1362.

38.   Taylor, C. V. (1919). The neuromotor system of Euplotes. Exp. Biol. Med. *16*, 101–103.

39. Syberg-Olsen, M.J., Irwin, N.A.T., Vannini, C., Erra, F., Di Giuseppe, G., Boscaro, V., and Keeling, P.J. (2016). Biogeography and character evolution of the ciliate genus Euplotes (Spirotrichea, Euplotia), with description of Euplotes curdsi sp. nov. PLoS One *11*.

40. Ricci, N., Giannetti, R., and Miceli, C. (1988). The ethogram of Euplotes crassus (ciliata, hypotrichida): I. The wild type. Eur. J. Protistol. *23*, 129–140.

41. Li, Y., Shi, L., and Gu, F. (2010). Comparison of ciliature microtubule organelles in three hypotrichous ciliate species*. Chinese J. Oceanol. Limnol. *28*, 443–450.

42. Fleury, A. (1991). Dynamics of the cytoskeleton during morphogenesis in the ciliate Euplotes I. Basal bodies related microtubular system. Eur. J. Protistol. *27*, 99–114.

43. Purcell, E.M. (1977). Life at low Reynolds number. Am. J. Phys. *45*, 3–11.

44. Kimmel, J.C., Chang, A.Y., Brack, A.S., and Marshall, W.F. (2018). Inferring cell state by quantitative motility analysis reveals a dynamic state system and broken detailed balance. PLoS Comput. Biol. *14*, e1005927.

45. Chang, A.Y., and Marshall, W.F. (2019). Dynamics of living cells in a cytomorphological state space. Proc. Natl. Acad. Sci. U. S. A. *116*, 21556–21562.

46. Boltzmann, L. (1872). Weirere Studien uber das warmegleich-gewich unter gasmolekulen. Sitzungsberichte der Akad. der Wissenschaften, Vienna, part II *66*, 275–370.

47. Alberts, B., Johnson, A., Lewis, J., Raff, M., Roberts, K., and Walter, P. (2002). Molecular biology of the cell 4th ed. (Garland Science).

48. Shalizi, C.R., and Crutchfield, J.P. (2001). Computational Mechanics: Pattern and Prediction, Structure and Simplicity. J. Stat. Phys. *104*, 817–879.

49. Crutchfield, J.P. (1994). The calculi of emergence: computation, dynamics and induction. Phys. D Nonlinear Phenom. *75*, 11–54.

1469   50.   Hopcroft, J.E., Motwani, R., and Ullman, J.D. (2008). Introduction to Automata Theory,

1470   Languages, and Computation 3rd ed. (Pearson Education).

1471   51.   Ricci, N., Giannetti, R., and Miceli, C. (1988). The ethogram of Euplotes crassus (ciliata,

1472   hypotrichida): I. The wild type. Eur. J. Protistol. *23*, 129–140.

1473   52.   Moore, T.Y., Cooper, K.L., Biewener, A.A., and Vasudevan, R. Unpredictability of

1474   escape trajectory explains predator evasion ability and microhabitat preference of desert

1475   rodents.

1476   53.   Goldstein, R.E. (2015). Green Algae as Model Organisms for Biological Fluid Dynamics.

1477   Annu. Rev. Fluid Mech. *47*, 343–375.

1478   54.   Simon, M., and Plattner, H. (2014). Unicellular Eukaryotes as Models in Cell and

1479   Molecular Biology: Critical Appraisal of Their Past and Future Value. In International

1480   Review of Cell and Molecular Biology (Elsevier Inc.), pp. 141–198.

1481   55.   Durham, W.M., Climent, E., Barry, M., De Lillo, F., Boffetta, G., Cencini, M., and

1482   Stocker, R. (2013). Turbulence drives microscale patches of motile phytoplankton. Nat.

1483   Commun. *4*.

1484   56.   Mogilner, A., and Oster, G. (1996). Cell motility driven by actin polymerization. Biophys.

1485   J. *71*, 3030–3045.

1486   57.   Collins, J.J., and Richmond, S.A. (1994). Biological cybemet Hard-wired central pattern

1487   generators for quadrupedal locomotion.

1488   58.   Collins, J.J., and Stewart, I.N. (1993). Coupled nonlinear oscillators and the symmetries of

1489   animal gaits. J. Nonlinear Sci. *3*, 349–392.

1490   59.   Barnhart, E.L., Allen, G.M., Jülicher, F., and Theriot, J.A. (2010). Bipedal locomotion in

1491   crawling cells. Biophys. J. *98*, 933–942.

1492   60.   Machemer, H., and Sugino, K. ELECTROPHYSIOLOGICAL CONTROL OF CILIARY

BEATING: A BASIS OF MOTILE BEHAVIOUR IN CILIATED PROTOZOA. Comp. Biochem. Physiol *94*, 365.

61. Mogami, Y., Pernberg, J., and Machemer, H. (1992). Ciliary beating in three dimensions: Steps of a quantitative description. J. Math. Biol. 1992 303 *30*, 215–249.

62. McNeill Alexander, R. (2003). Principles of Animal Locomotion (Princeton University Press).

63. Wan, K.Y., and Jékely, G. (2020). On the unity and diversity of cilia. Philos. Trans. R. Soc. Lond. B. Biol. Sci. *375*, 20190148.

64. Ahamed, T., Costa, A.C., and Stephens, G.J. Capturing the Continuous Complexity of Behavior in C. elegans.

65. Stephens, G.J., Johnson-Kerner, B., Bialek, W., and Ryu, W.S. (2008). Dimensionality and Dynamics in the Behavior of C. elegans. PLoS Comput. Biol. *4*, e1000028.

66. Berman, G.J., Choi, D.M., Bialek, W., and Shaevitz, J.W. (2014). Mapping the stereotyped behaviour of freely moving fruit flies. J. R. Soc. Interface *11*, 20140672.

67. Stephens, G.J., Osborne, L.C., and Bialek, W. (2010). Searching for simplicity: Approaches to the analysis of neurons and behavior. Proc. Natl. Acad. Sci. U. S. A. *108*, 15565–15571.

68. Devarajan, K. (2008). Nonnegative Matrix Factorization: An Analytical and Interpretive Tool in Computational Biology. PLoS Comput. Biol. *4*.

69. Lee, D.D., and Seung, H.S. (1999). Learning the parts of objects by non-negative matrix factorization. Nat. 1999 4016755 *401*, 788–791.

70. Ester, M., Kriegel, H.-P., Sander, J., and Xu, X. (1996). A Density-Based Algorithm for Discovering Clusters in Large Spatial Databases with Noise.

71. Kauffman, L., and Rousseeuw, P.J. (1990). Finding Groups in Data: and introduction to

cluster analysis (John Wiley & Sons, Inc.).

72.   Kirkegaard, J.B., Marron, A.O., and Goldstein, R.E. (2016). Motility of Colonial

Choanoflagellates and the Statistics of Aggregate Random Walkers. Phys. Rev. Lett. *116*,

038102.

73.   M Cartmill (1985). Climbing. In Functional Vertebrate Morphology (Harvard University

Press), pp. 73–88.

74.   Kolmogoroff, A. (1936). Zur Theorie der Markoffschen Ketten. Math. Ann. *112*, 155–

160.

75.   Gagniuc, P. (2017). Markov Chains: From Theory to Implementation and

Experimentation (Wiley).

76.   Markov, A.A. (1907). Extension of the Limit Theorems of Probability Theory to a Sum of

Variables Connected in a Chain. Notes Imp. Acad. Sci. St. Petersbg. VIII Ser. Physio-

Mathematical Coll. *22*.

77.   Howard, R.A. (1971). Dynamic and Probabilistic Systems, Volume 1: Markov models

(Dover Publications).

78.   Crutchfield, J.P., and Young, K. (1989). Computation at the Onset of Chaos. In

Complexity, Entropy, and Physics of Information, W. Zurek, ed. (Addison-Wesley).

79.   Crutchfield, J.P., and Young, K. (1989). Inferring statistical complexity. Phys. Rev. Lett.

63, 105–108.

80.   Moore, C. (1998). Dynamical recognizers: Real-time language recognition by analog

computers. Theor. Comput. Sci.

81.   Skinner, D.J., and Dunkel, J. (2021). Improved bounds on entropy production in living

systems. Proc. Natl. Acad. Sci. U. S. A. *118*.

82.   Martínez, I.A., Bisker, G., Horowitz, J.M., and Parrondo, J.M.R. Inferring broken detailed

balance in the absence of observable currents.

83.   Esposito, M. (2012). Stochastic thermodynamics under coarse graining. Phys. Rev. E - Stat. Nonlinear, Soft Matter Phys. *85*, 041125.

84.   Taylor, C.V. (1921). Demonstration of the function of the neuromotor apparatus in Euplotes by the method of microdissection. Naturwissenschaften *9*, 910.

85.   Okajima, A., and Kinosita, H. (1966). Ciliary activity and coordination in Euplotes eurystomus-I. Effect of microdissection of neuromotor fibres. Comp. Biochem. Physiol. *19*, 115–131.

86.   Naitoh, Y., and Eckert, R. (1969). Ciliary Orientation: Controlled by Cell Membrane or by Intracellular Fibrils? Science (80-.). *166*, 1633–1635.

87.   Okabe, A., and Miki, F. (1984). A Conditional Nearest-Neighbor Spatial-Association Measure for the Analysis of Conditional Locational Interdependence. Environ. Plan. A Econ. Sp. *16*, 163–171.

88.   Jennings, H.S., and Jamieson, C. (1902). Studies on reactions to stimuli in unicellular organisms. X. The movements and reactions of pieces of ciliate infusoria. Biol. Bull. *3*, 225–234.

89.   Shalizi, C.R., Shalizi, K.L., and Crutchfield, J.P. (2002). An Algorithm for Pattern Discovery in Time Series. ArXiv.

90.   Lynn, C.W., Cornblath, E.J., Papadopoulos, L., Bertolero, M.A., and Bassett, D.S. (2021). Broken detailed balance and entropy production in the human brain. Proc. Natl. Acad. Sci. U. S. A. *118*.

91.   Bar-Ziv, R., Tlusty, T., and Libchaber, A. (2002). Protein-DNA computation by stochastic assembly cascade. Proc. Natl. Acad. Sci. U. S. A. *99*, 11589–11592.

92.   Berman, G.J., Bialek, W., and Shaevitz, J.W. (2016). Predictability and hierarchy in

Drosophila behavior. Proc. Natl. Acad. Sci. U. S. A. *113*, 11943–11948.

93. Guasto, J.S., Rusconi, R., and Stocker, R. (2012). Fluid Mechanics of Planktonic
     Microorganisms. Annu. Rev. Fluid Mech. *44*, 373–400.

94. Goldstein, R.E., Polin, M., and Tuval, I. (2009). Noise and Synchronization in Pairs of
     Beating Eukaryotic Flagella. Phys. Rev. Lett. *103*, 168103.

95. Párducz, B. (1967). Ciliary Movement and Coordination in Ciliates. Int. Rev. Cytol. *21*,
     91–128.

96. Funfak, A., Fisch, C., Abdel Motaal, H.T., Diener, J., Combettes, L., Baroud, C.N., and
     Dupuis-Williams, P. (2015). Paramecium swimming and ciliary beating patterns: A study
     on four RNA interference mutations. Integr. Biol. (United Kingdom) *7*, 90–100.

97. Kirkegaard, J.B., Bouillant, A., Marron, A.O., Leptos, K.C., and Goldstein, R.E. (2016).
     Aerotaxis in the closest relatives of animals. Elife *5*, e18109.

98. Miño, G.L., Koehl, M.A.R., King, N., and Stocker, R. (2017). Finding patches in a
     heterogeneous aquatic environment: pH-taxis by the dispersal stage of choanoflagellates.
     Limnol. Oceanogr. Lett. *2*, 37–46.

99. Polin, M., Tuval, I., Drescher, K., Gollub, J.P., and Goldstein, R.E. (2009).
     Chlamydomonas Swims with Two "Gears" in a Eukaryotic Version of Run-and-Tumble
     Locomotion. Science (80-.). *325*, 487–490.

100. Li, Y., Kučera, O., Cuvelier, D., Rutkowski, D.M., Deygas, M., Rai, D., Pavlovič, T.,
     Vicente, F.N., Piel, M., Giannone, G., et al. (2022). Compressive forces stabilise
     microtubules in living cells. bioRxiv, 2022.02.07.479347.

101. Maniotis, A.J., Chen, C.S., and Ingber, D.E. (1997). Demonstration of mechanical
     connections between integrins, cytoskeletal filaments, and nucleoplasm that stabilize
     nuclear structure. Proc. Natl. Acad. Sci. U. S. A. *94*, 849–854.

1589   102.   Craddock, T.J.A., and Tuszynski, J.A. (2010). A critical assessment of the information

1590   processing capabilities of neuronal microtubules using coherent excitations. J. Biol. Phys.

1591   *36*, 53–70.

1592   103.   Stock, C., Krüppel, T., Key, G., and Lueken, W. (1999). Sexual behaviour in Euplotes

1593   raikovi is accompanied by pheromone-induced modifications of ionic currents. J. Exp.

1594   Biol. *202*.

1595   104.   Tuffrau, M., Fryd-Versavel, G., Tuffrau, H., and Génermont, J. (2000). Description of

1596   Euplotes versatilis n. sp., a marine tropical ciliate exhibiting an unusually extensive

1597   phenotypic plasticity. Eur. J. Protistol. *36*, 355–366.

1598   105.   Keeling, P.J. (2019). Combining morphology, behaviour and genomics to understand the

1599   evolution and ecology of microbial eukaryotes. Philos. Trans. R. Soc. B Biol. Sci. *374*.

1600   106.   Kahl, A. (1935). Urtiere oder Protozoa. (Fischer).

1601   107.   Yocom, H.B. (1917). The neuromotor apparatus of Euplotes patella.

1602   108.   Tinevez, J.Y., Perry, N., Schindelin, J., Hoopes, G.M., Reynolds, G.D., Laplantine, E.,

1603   Bednarek, S.Y., Shorte, S.L., and Eliceiri, K.W. (2017). TrackMate: An open and

1604   extensible platform for single-particle tracking. Methods *115*, 80–90.

1605   109.   King, N., Young, S.L., Abedin, M., Carr, M., and Leadbeater, B.S.C.C. (2009). Starting

1606   and maintaining Monosiga brevicollis cultures. Cold Spring Harb. Protoc. *4*,

1607   pdb.prot5148.

1608   110.   Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T.,

1609   Preibisch, S., Rueden, C., Saalfeld, S., Schmid, B., et al. (2012). Fiji: an open-source

1610   platform for biological-image analysis. Nat. Methods *9*, 676–682.

1611   111.   Kanagal, B., and Sindhwani, V. (2010). Rank Selection in Low-rank Matrix

1612   Approximations: A Study of Cross-Validation for NMFs. In Advances in Neural

Information Processing Systems.

112.   Owen, A.B., and Perry, P.O. (2009). Bi-Cross-Validation of the SVD and the Nonnegative

Matrix Factorization. Source Ann. Appl. Stat. *3*, 564–594.

113.   Wold, S. (1978). Cross-Validatory Estimation of the Number of Components in Factor

and Principal Components Models. Technometrics *20*, 397–405.

114.   Ronan, T., Qi, Z., and Naegle, K.M. (2016). Avoiding common pitfalls when clustering

biological data. Sci. Signal. *9*, re6–re6.

115.   Kawamoto, T., and Kabashima, Y. (2017). Cross-validation estimate of the number of

clusters in a network. Sci. Rep. *7*, 1–17.

116.   Bak, P., Tang, C., and Wiesenfeld, K. (1987). Self-organized criticality: An explanation of

the 1/f noise. Phys. Rev. Lett. *59*, 381–384.

117.   Sánchez, R., Newman, D.E., and Carreras, B.A. (2002). Waiting-Time Statistics of Self-

Organized-Criticality Systems. Phys. Rev. Lett. *88*, 4.

118.   Billingsley, P. (1961). Statistical Methods in Markov Chains (Institute of Mathematical

Statistics).

119.   Hiscott, R.N. (1981). Chi-Square Tests for Markov Chain Analysis.

120.   Happel, J., and Brenner, H. (1981). Low Reynolds number hydrodynamics. *1*.

121.   Lighthill, J. (1976). Flagellar Hydrodynamics. SIAM Rev. *18*, 161–230.

122.   Cardona, A., Saalfeld, S., Schindelin, J., Arganda-Carreras, I., Preibisch, S., Longair, M.,

Tomancak, P., Hartenstein, V., and Douglas, R.J. (2012). TrakEM2 Software for Neural

Circuit Reconstruction. PLoS One *7*, e38011.

123.   Thévenaz, P., Ruttimann, U.E., and Unser, M. (1998). A pyramid approach to subpixel

registration based on intensity. IEEE Trans. Image Process. *7*, 27–41.

124.   Kingma, D.P., and Welling, M. (2013). Auto-Encoding Variational Bayes.

125.   Kingma, D.P., and Ba, L.J. (2015). Adam: A Method for Stochastic Optimization.

126.   Bach, F.R., and Jordan, M.I. (2003). Learning Spectral Clustering. Adv. Neural Inf.

1639   Process. Syst. *16*.

1640   127.   Pedregosa, F., Michel, V., Grisel, O., Blondel, M., Prettenhofer, P., Weiss, R.,

1641   Vanderplas, J., Cournapeau, D., Pedregosa, F., Varoquaux, G., et al. (2011). Scikit-learn:

1642   Machine Learning in Python. J. Mach. Learn. Res. *12*, 2825–2830.

1643   128.   Darmon, D. (2020). transCSSR. GitHub Repos.

1644