**A unicellular walker embodies a finite state machine**

Authors: Ben T. Larson[1], Jack Garbus[2], Jordan B. Pollack[2], and Wallace F. Marshall[1]

[1] Department of Biochemistry and Biophysics, University of California, San Francisco,

California, USA

[2] Computer Science Department, Brandeis University, Waltham, Massachusetts, USA

**Cells are complex biochemical systems whose behavior emerges from interactions**

**among myriad molecular components. The idea that cells execute computational processes**

**is often invoked as a general framework for understanding cellular complexity. However,**

**the manner in which cells might embody computational processes in a way that the**

**powerful theories of computation, such as finite state machine models, could be**

**productively applied, remains to be seen. Here we demonstrate finite state machine-like**

**processing embodied in cells, using the walking behavior of *Euplotes eurystomus*, a ciliate**

**that walks across surfaces using fourteen motile appendages called cirri. We found that**

**cellular walking entails a discrete set of gait states. Transitions between these states are**

**highly regulated, with distinct breaking of detailed balance and only a small subset of**

**possible transitions actually observed. The set of observed transitions decomposes into a**

**small group of high-probability unbalanced transitions forming a cycle and a large group**

**of low-probability balanced transitions, thus revealing stereotypy in sequential patterns of**

**state transitions. Taken together these findings implicate a machine-like process. Cirri are**

**connected by microtubule bundles, and we find an association between the involvement of**

**cirri in different state transitions and the pattern of attachment to the microtubule bundle**

**system, suggesting a mechanical basis for the regularity of state transitions. We propose a**

**model where the actively controlled, unbalanced transitions establish strain in certain cirri,**

**the release of which from the substrate causes the cell to advance forward along a linear**

**trajectory. This demonstration of a finite state machine embodied in a living cell opens up new links between theoretical computer science and cell biology and may provide a general framework for understanding and predicting cell behavior at a super-molecular level.**

**Introduction**

Cells are complex physical systems controlled by networks of signaling molecules. Single cells can display remarkably sophisticated, seemingly animal-like behaviors [1–3], orchestrating active processes far from thermodynamic equilibrium in order to carry out proper biological functions [4,5]. Indeed, single cells can make decisions by sensing and responding to diverse cues and signals [6], execute coordinated movements [7,8] and directed motility [9–12], and even solve mazes [13,14] and possibly learn [15–18]. Such behaviors in animals arise from neural activity and have been studied extensively, but we know comparatively little about the mechanisms of cellular behavior [19,20]. In individual cells, behaviors emerge directly through the joint action of chemical reactions [21], cellular architecture [3], physical mechanisms and constraints within the cell [22,23] and interactions of the cell with its local environment [24]. The links between information processing, decision-making, and their physical manifestation as cell state transitions suggest that cellular behavior might be understood as an embodied computation [25,26]. The theory of computation has often been invoked as a general framework for understanding cellular dynamics [25,27–32], with environmental sensing by bacteria being a particularly deeply studied example [32–34], and has been used to engineer programmable cell states [35], but the manner and extent to which cells might embody functional, computational processes as well as the extent to which a computational perspective on cellular behavior might prove productive remains to be seen.

Among the microbial eukaryotes (protists), ciliates display striking examples of

unicellular behavior including hunting [3], sensorimotor navigation [10], and predator avoidance [36].

Spirotrichous ciliates of the genus *Euplotes* are notable for their complex locomotion [37–39], using

bundles of specialized cilia called cirri to walk across surfaces [37,38] (Fig. 1a, Movies S1 and S2).

Depending on the species, cells generally have 14 to 15 ventral cirri arranged in a highly

stereotyped pattern used for walking locomotion [40]. *Euplotes* live in aquatic environments, and in

addition to walking, their cirri can be used for swimming and rapid escape responses [41] (Movie

S2). Oral membranelles (Fig. 1b) generate feeding currents to capture bacteria and small

protistan prey and are also used for swimming. Early 20th century protistologists were so

impressed by the apparent coordination of cirri that they proposed the existence of a rudimentary

nervous system, the neuromotor apparatus, to account for their observations [39]. This theory was

motivated in part by the presence of intracellular fibers connecting various cirri (Fig. 1C), now

known to be to be tubulin-based structures [42,43]. Although the walking movements of *Euplotes*

are superficially similar to those of animals such as insects, the low Reynolds environment of

aquatic microorganisms, where viscous forces dominate over inertial forces, imposes significant

physical constraints on all movements that do not impinge on the movements of larger terrestrial

animals [44].

How can a single cell coordinate a gait without a nervous system? Coordination, to the

extent that it exists in the gait of *Euplotes*, would require some kind of dynamical coupling

among cirri or between cirri and some shared external influence. Recently, analytical techniques

from statistical physics have been used to characterize, understand, and predict mesoscale

dynamics in biological systems, including cellular behavior [4,5,45,46]. These approaches rely on

coarse-graining the complexity of biological dynamics into states and analyzing the nature of

transitions between states. For example, a state representation allows us to ask whether forward

and reverse transitions between pairs of states are equal, a condition known as detailed balance

[4,47]. Systems that violate detailed balance operate in a non-equilibrium mode and can produce

directed cycles in state space [4,48]. Broken detailed balance has been observed in the motility

dynamics of cultured mammalian cells as well as the motility dynamics of a freely behaving

flagellate protist [5,45] and implies that non-equilibrium models are most applicable to such

systems [46].

When information processing drives patterns of state transitions, such a system can be

viewed in terms of automata theory, a fundamental level in the theory of computation [49–

51].Automata theory can be used to address problems of decision-making and control in complex

systems by providing predictive understanding independently of the underlying details of how a

given process is implemented [50]. Inspired by work considering cellular behavior in the context of

the theory of computation [25], we hypothesized that walking cells might be governed by finite

state automata with directed, processive movement arising from reproducible patterns of state

transitions.

The consistent structure of *Euplotes*, its mode of motility, and its ease of observation

makes these cells an ideal biological test-bed in which to apply theories of non-equilibrium

statistical mechanics and embodied computation, both of which rely on describing a system in

terms of discrete state transitions. Here, we use time-lapse microscopy and quantitative analyses

to show that *Euplotes eurystomus* walks with a cyclic stochastic gait displaying broken detailed

balance and exhibiting elements of stereotypy and variability, consistent with a finite state

automaton representation. The observed dynamics are reminiscent of behavioral regulation in

some cells and animals [5,52] but  contrast with many well-characterized examples of cellular

motility [7,9,10,12,53–56]. Our results provide a clear demonstration of machine-like processes

governing cellular state transitions and serve as a framework for investigating the principles of

behavioral control and non-equilibrium dynamics in single cells.

**Results**

***Walking dynamics can be described in a reduced state space***

In order to ask whether cell behavior is governed by a finite state machine, we analyzed

the walking behavior of *Euplotes eurystomus* cells, [41], focusing on the simplest case of

uninterrupted, linear walking trajectories (Fig. 2a, b, Movie S1). Cells were placed onto

coverslips on which free, spontaneous walking behavior was observed by microscopy. A focal

plane at the cirrus-coverslip interface was chosen in order to clearly observe cirral dynamics

(Fig. 2a). The relative spatial positioning of cirri is highly stereotyped from cell to cell, allowing

us to give each of the 14 cirri an alphabetic label from a-n (Figure 2c). In each video frame (33

frames/s), the walking state of the cell was encoded as a 14-bit binary vector, with each bit

corresponding to a cirrus and receiving a value of "0" if the cirrus was in contact with the

coverslip and stationary and a "1" if the cirrus was in motion (instances of stationary cirri held

above the coverslip for a sustained period of time were not observed). The trajectories of 13 cells

were manually tracked and annotated for a total of 2343 time points. This quantitative analysis

revealed stepping-like cirral dynamics in that cirri tend to undergo rapid movements followed by

longer periods of quiescence (Fig. 2d). Cirral dynamics appeared to lack any obvious patterns

such as periodicity or repeating sequences of states (e.g. Fig. 2d), implying that the state

sequences are either stochastic, or generated by complex deterministic processes. This lack of

periodicity or fixed phase relationships between appendage movements is different from the gaits

of most animals or those reported for various flagellates [57–59]. Autocorrelation analysis confirmed

the observed lack of clear periodicity (Fig. S1).

Despite the apparent complexity of cirral dynamics, we suspected that discernable

structure might exist, which would allow us to obtain a reduced state space that accurately

described the dynamics as has proved successful in behavioral analysis of diverse living systems

[45,46,60–63]. We performed dimensionality reduction using non-negative matrix factorization

(NMF), and cross-validation by imputation [64,65] (see Methods and Fig. S2 for more details)

revealed the cirral states to be well-described in three dimensions (Fig. 2e-g). The components of

the NMF analysis correspond to distinct groups of cirri, and these groups constitute spatially

distinct partitions of cirri with respect to their positions on the cell body (Fig. 3h). The

dimensionality reduction of the gait state space arises at least in part from shared pairwise mutual

information between groups of cirri (Fig. 3i).

We next used the density-based spatial clustering of applications with noise (DBSCAN)

algorithm [66] to group our data into clusters in an unbiased fashion, with members of a given

cluster sharing similar patterns of cirral activity. Visual inspection in conjunction with silhouette

coefficient [67] (a metric of cluster cohesion and separation) analysis revealed that 32 clusters

accurately captured the visible structure in the reduced state space without overfitting (Fig. 2e-g,

see Methods and Fig. S3 for more details)[*]. These reduced gait states correspond to distinct

patterns of cirral activity (Fig. 2j). Taken together, our results reveal stereotypy in the

spatiotemporal patterns of cirral activity. The discrete, reduced state space, or set of gait states,

---

[*] The problem of determining the true number of clusters is an unresolved problem [89]. We have followed standard methods to determine cluster number but have found that our key results do not depend sensitively on the precise number of clusters identified (see SI and Fig. S7).

demonstrates that cells make use of a subset of the possible patterns of appendage movement

during walking locomotion.

***Euplotes walks with a cyclic stochastic gait***

In order to relate the gait states identified in our cluster analysis, we asked how changes

in the number of active cirri may relate to cell movement. Naively, one might expect that the

force associated with locomotion is roughly proportional to the number of moving appendages [68].

Alternatively, we velocity might inversely correlate with the net change in cirral activity, which

would be expected if stationary cirri were generating a pushing traction force as in crawling or

climbing animals [58,69] or if cirri execute a power stroke just before coming to rest as has been

suggested by Erra et al. [38]. At low Reynolds number, velocity should be proportional to the

difference between the net force generated by the cell and the opposing drag [44]. Examining cell

velocity versus the net change in cirral activity, however, showed that neither of these

expectations were in fact the case (Fig. 3a). Cell velocity was only weakly correlated with

number of active cirri ($R^2=0.03$), and instead, small to moderate positive and negative changes in

the number of active cirri corresponded to the largest cell velocities (Fig. 3a). We reasoned that

transitions between gait states must be important to driving the forward progression of walking

cells, and so sought to determine whether this active coordination might manifest in the observed

gait dynamics.

Analysis of the 1423 pairwise transitions in our dataset yielded the transition matrix

displayed in Fig. 3b (see Methods for more details). The presence of strongly unbalanced

transitions such as from gait state 3 to 17 versus 17 to 3 suggested broken detailed balance, and

indeed, a number of forward and reverse transitions were found to be significantly unbalanced by

the binomial test (see Methods). Entropy production rate has been used to quantify the degree of

broken detailed balance, or, similarly, the distance from equilibrium where the entropy

production rate will be zero [5]. Following the procedure detailed in [5], we obtain a lower bound

estimate for an entropy production rate of 0.4, similar to the value reported for strongly non-

equilibrium gait transitions observed in a flagellate [5]. Walking *Euplotes* cells, therefore, have a

strongly non-equilibrium gait despite lack of periodicity.

Only 322 of the 1024 possible types of transitions were observed to occur at least once,

and within this restricted set, only 173 occurred more than once (Fig. 3c). Also, we found that

relatively few transitions corresponded to substantial cellular movement (Fig. 3c). Crucially, the

presence of broken detailed balance revealed the existence of directed cycles of cirral activity

during locomotion. To get a better understanding of the nature of these cycles, we grouped

transitions into two categories: balanced transitions, which satisfy detailed balance, and

unbalanced transitions, which do not (see Methods for details). This partitioning allowed us to

separately investigate unbalanced, non-equilibrium-like and balanced, equilibrium-like

transitions (Fig. 3d, e). Significantly unbalanced transitions ($p<0.05$ by binomial test) are among

the most frequent transitions, but only involve a small number of states (Fig. 3d). Of the 32 gait

states, we found that only states 2, 3, 4, 7, 17, and 27 were associated with unbalanced

transitions. We noted the presence of one complete cycle with unbalanced transitions following

2→3→17→2. We had expected that unbalanced transitions might be associated with a "power

stroke" in the sense of occurring simultaneously with cell movement, but in fact high cellular

velocities tended to be associated with balanced transitions (Fig. 3d, e). Additionally, we found

that, with the exception of transitions between states 1 and 2, transitions occurring at the highest

frequencies were unbalanced (Fig. 3d).

Notably, the most frequent balanced transitions were associated with transitions into and

out of gait state 1, a unique "rest state" which involves no cirral movement (Fig. 3c, d).

Furthermore, we found by computing the autocorrelation function for a binarized sequence of

each state that gait state 1 has the most predictable dynamics in terms of significant positive

autocorrelation in contrast to the lack of significant autocorrelation seen for other states (see Fig.

S4). Although transitions between states 1 and 2 are balanced, the most frequent transitions out

of state 2 are strongly biased toward transitions into state 3, from which other strongly biased

transitions also frequently occur, including the cycle of biased transitions mentioned in the

preceding paragraph. The presence of high frequency unbalanced transitions does not preclude

the existence of highly variable trajectories through state space. The picture of walking

trajectories that emerges is of stochastic excursions from gait state 1 into a stochastic cycle

involving a mix of balanced and unbalanced transitions, with the majority of cell movement

occurring during infrequent, equilibrium-like transitions. Biased transitions, occurring at

relatively high frequency from a subset of states, introduce temporal irreversibility to the gait of

*Euplotes* due to their strongly non-equilibrium character.

Finally, we checked whether gait state transition dynamics had the Markov property,

which entails that transition probabilities are determined completely by the present state, and that

previous dynamics contribute no additional predictive information [70,71]. Lack of past dependence

has led to Markov processes often being referred to as "memoryless" [72]. When we compared the

gait transition matrix (Fig. 3b) with a computed transition matrix over two timesteps, we

observed these matrices to be different from one another, which is inconsistent with the Markov

property (see Fig. S5). Finally, we applied a Billingsley test, a chi-squared measure for

Markovness [73], which revealed that the null hypothesis that the process was Markov could be

rejected (p=0.005). These analyses showed that *Euplotes* retains some "memory" of the prior

sequence of cirral movements during locomotion.

Taken together, our analysis revealed a mixture of unbalanced transitions arranged in

cycles and balanced transitions arranged as networks, for which we propose to apply the term

"cyclic stochastic gait". The cyclic stochastic gait of *Euplotes eurystomus* incorporates elements

of both stereotypy and variability in gait dynamics. Forward progress of the cell is not produced

merely by a physical ratchetting process driven by unpatterned fluctuations in cirral activity, nor

is it produced by a highly regular, deterministic process like a clock. It has been argued that

significant computation arises in physical systems exhibiting such a mix of stereotypy and

variability [49,74,75] in the sense that the time-evolution of the system is most compactly described

by the result of a computation involving state transitions, memory, and decision rules, rather than

by a periodic oscillation or a random coin flip.

While our analysis revealed a computational underpinning of gait, we sought to better

understand the functional organization of the dynamical patterns driving processive motion of

the cell. To do so, we first focused on the highest transition probabilities emanating from each

state. Transition probabilities were estimated as $N_{ij}/N_i$ where $N_{ij}$ is the number of transitions from

state $i$ to state $j$, and $N_i$ is the total number of transitions from state $i$. This allowed us to prune

away rare transitions in order to reveal the dominant structure of gait state transitions. Figure 3g

displays the pruned transition matrix as a heatmap. We found that relatively few states were the

recipients of the majority of high probability transitions, and many states received none. To more

clearly visualize the structure of transitions, we grouped together all gait states receiving no more

than one unique high probability transition with the idea being that state transitions into this

group show little bias in terms of source state, and within the group, transitions between states

exhibit low probability, time unbiased, equilibrium-like fluctuations. the majority of cell

movement was associated with transitions between states within this group. In contrast to the

"cloud" of states linked by low-probability, balanced transitions, nearly all of the states receiving

high probability transitions were either the three "cycle" states, or else fed cycle states with their

highest probability transitions, with the majority feeding gait state 17.

Focusing on the structure of transitions in this way allowed us to derive a simplified,

functional representation of stereotypy in gait dynamics as depicted in Fig. 3h. Although gait

state 1 is not the recipient of any individual high probability transitions, we identified it as the

unique "start" state from which cells initiate walking. Beginning with this start state, cells

transition with high probability to gait state 2, also one of the highest frequency transitions and

the first state in the 2→3→17→2 cycle of unbalanced transitions. From this first cycle state,

cells transition to gait state 3, the second cycle state, with highest probability and frequency and

then similarly on to gait state 17, the third cycle state. This sequence from the start state through

the cycle states corresponds to increasing amounts of cirral activity. Although the highest

probability transitions from the third cycle state to any single gait state return to the first and

second cycle state with equal probability and return to the first cycle state also being unbalanced,

cells in fact transition to the equilibrium "cloud" of motility-associated states with overall higher

probability. Return to the cycle states tend to occur through various moderately high probability

transitions from the motility state cloud or through intermediate states. In conjunction with this

set of transitions, we also noted unbalanced transitions stemming from the cycle states to the

motility state as well as intermediate state subsequently feeding the next cycle state.

Altogether, the picture that emerges of stereotypical gait dynamics is of biased transitions

through cycle states before relatively low probability, unbiased transitions associated with

substantial cell movement before returning to the start or cycle states and beginning the sequence

again. While this general sequence is repeated during walking, there is variability or apparent

stochasticity in the details of gait state transitions with increasingly variable transitions as any

given sequence progresses. We propose that the cycle states serve to establish configurations of

cirri necessary for cells to later transition to between states from which forward progress of the

cell is generated. Many state transitions along any instance of the stereotyped sequence are

unbiased, but biased, high probability transitions, presumably resulting from active cellular

control, give temporal irreversibility to the sequence.

***The fiber system of Euplotes constrains models of gait coordination***

What physical machinery could embody this information processing required to generate

the stochastic cyclic state transitions seen during *Euplotes*' walking? We reasoned that there

must be some form of coupling or communication between cirri or feedback between gait state

and cirral dynamics. The role of the system of cytoskeletal fibers associated with cirri as conduits

of information between cirri during cellular locomotion, supported by microsurgical

experiments, has been a key hypothesized mechanism of gait coordination since the early 1900s

[76,77]. We wondered whether the structure of the cytoskeletal fiber system associated with cirri

(Fig. 4a) could give some insight into how cirri might be coordinated.

We sought to characterize and reconstruct in 3D the tubulin-based fiber system of

*Euplotes* associated with cirri and lying just beneath the cell cortex [39,42,43]. Upon inspection of

our confocal reconstructions of SiR-tubulin labeled cells (Fig. 4a, Fig. S6), we noted the

presence of two morphologically distinct classes of fibers, one thicker, linear class and the other

more filamentous and less linear, consistent with previous observations (Fig. 1c, [39,42,43]).

Additionally, we observed a group of thick linear fibers associated with some of the frontal cirri,

which to our knowledge has not been previously reported. Fibers emanate from the base of all

cirri, appear to intersect one another, and also connect to the cortex of the cell at various points

(Fig. 3g). Some cirri were found to be associated with only thick fibers while others have both or

only thin fibers. Based on apparent fiber intersections and convergences, we found the fiber

system to constitute a continuous network between all cirri, with the fibers associated with the

base of each cirrus intersecting the fiber system associated with at least one other cirrus (Fig. 4a,

b). Contrary to the long-standing standing hypothesis from the literature [76], the functional

modules (groups of co-varying cirri) identified in our dynamical analysis were not exclusively

linked by dense fiber intersections [39,43,76] (Fig. 4a, b). In fact, connections between cirri are not

generally associated with any statistically significant difference in mutual information (defined

in terms of the information that the activation state of one cirrus has concerning the other)

compared to unlinked pairs of cirri (p=0.14 by Wilcoxon rank sum test, Fig. 4C). However,

information flow became apparent when fiber-fiber links are grouped by type (i.e. thick to thick

fiber, thick to thin fiber, or thin to thin fiber). We found that pairs of cirri associated with only

thick fiber to thick fiber and only thin fiber to thin fiber links have increased mutual information

compared to those without links (Fig. S6). Interestingly, we found that cirri nearby one another

and connected by fibers to similar regions of the cell cortex shared the most mutual information

(Fig. 2i, 3c, 4d, e), suggesting that if the fibers play a role in cirral coordination, coupling may

also be mediated by mechanisms at the cirrus and fiber-cortex interface. Cirri d, e, h, i, for

example, share very little mutual information with any of the other cirri, and fibers emanating

from the base of these cirri contact the cell cortex and other fibers at various unique points. Cirri

g and f, on the other hand, which share more mutual information than any other pair, are

associated with both thick and thin fibers terminating at similar regions of the cell cortex. Indeed,

both distances between pairs of cirri and cross nearest neighbor distance (a measure of similarity

between discrete spatial distributions) between paired sets of cirrus-cortex contact points show

significant Spearman correlations to mutual information (-0.49, p<0.001 and -0.62, p<0.001

respectively) (Fig. 4d, e). These correlations indicate that mutual information between pairs of

cirri tends to increase with proximity and also tends to increase with similarity between fiber-

cortex contact locations, so the cirri with the highest mutual information are those that are close

together with similar fiber-cortex connections (Fig. 4d-f).  Together, these observations strongly

support a mechanical coordination mechanism in which microtubule bundles allow groups of

cirri to influence successive behavior of other groups of cilia.

**Discussion**

In order to meet the challenge of accounting for the emergence of apparently

sophisticated cellular behavior, we conceptualized the cell as a finite state machine.

Traditionally, studies of computational processes performed by cells have tended to focused on

combinatorial logic, where the output of a computational process depends only on the current

input, performed by networks of molecules in bacterial cells [25,32–34]. We have focused on

sequential logic, where outputs depend on the system state as well, an equally important aspect

of the theory of computation with notable yet less developed representation in studies of cellular

and sub-cellular dynamics [16,20,78]. Behavior of eukaryotes has frequently been observed to

involve stereotyped transitions between dynamical states, and our results suggest that automata

theory, which includes finite state machine models and necessarily involves sequential logic,

may be particularly well-suited to studying the behavior of eukaryotic cells. Our approach,

relying on dimensionality reduction to identify dynamical states, revealed modularity in cellular

dynamics associated with structural modularity of the cell (Fig. 2, 4) in addition to cyclic

patterns of sequential dynamical activity (Fig. 3).

Walking locomotion in *Euplotes* represents a departure from many of the best studied

appendage-based locomotor systems. For example, limbed locomotion in animals tends to

proceed by highly stereotyped, determinate patterns of activity [57,58], and many small, aquatic

animals exhibit periodic movements of appendages, often cilia, during locomotion [7,59,79]. Many

forms of unicellular locomotion involve such dynamics as well including in sperm cells [80],

diverse flagellates with various numbers of flagella [59], and ciliates [59,81,82]. Even in cases where

cellular locomotion involves fundamentally stochastic dynamics such as in run-and-tumble

motility in *E. coli* [12] or analogous behaviors observed in protists [11,83–85], motility can be described

by equilibrium processes [5], in contrast to the non-equilibrium character of the gait of *Euplotes*.

There are examples, however, of locomotor dynamics in both animals and unicellular organisms

that are reminiscent of those we have observed in *Euplotes*. Gait switching in kangaroo rats has

been shown to have a stochastic, non-equilibrium character, perhaps to facilitate predator

avoidance by being difficult to predict [52]. Most saliently, gait switching in an octoflagellate [5] and

motility dynamics in cultured mammalian cells [45] have been shown to exhibit broken detailed

balance. We propose that broken detailed balance in the gait of *Euplotes* indicates active

coordination of motility processes. Here, broken detailed balance in gait state transitions revealed

cyclic activity, characterized by transitions into and out of a resting state with a mixture of

stereotypy and variability in the intervening steps, in the gait of a single cell (Fig. 3, 4). To

explain how these dynamics give rise to directed walking, we propose a mechanism in which

biased, actively controlled cyclic transitions serve to establish strain, effectively storing stress, in

certain cirri, and the spontaneous release of these cirri from the substrate, during a series of

unbiased gait state transitions, allows the cell to move forward. The cloud of unbiased transitions

associated with substantial cellular movement is consistent with the fact that the details of the

precise order in which the strained cirri are released does not matter for generating motility.

Consistent with this mechanism, inspection of videos revealed that substantial cell movement

appears to be correlated with the movement of notably bent inactive cirri. Return to the cycle

states then are necessary to establish this process anew by winding up the system for continued

cell movement. This mechanism and these patterns of cirral activity are consistent with previous

observations of cyclic velocity fluctuations in the trajectories of walking *Euplotes* [37]. We argue

that subcellular processes must be involved in actively coordinating cirri in order to accomplish

the observed stereotypy in biased sequential activity. Our analysis of the tubulin-based

cytoskeletal fiber system is consistent with its role in mechanically mediating communication

among cirri and between cirri and cell cortex (Fig. 4). Thus, by combining information

processing to properly dictate patterns of cirral activity and the mechanical actions of cirral

movement, walking *Euplotes* embodies the sequential computation of a finite state machine.

Among the domains of life, eukaryotes uniquely display remarkable complexity and

diversity in cellular behavior [86]. Our approach, grounded in finite state-machine analysis, has

revealed modularity and stereotypy underlying a complex cellular behavior, implicating a

machine-like process. Our results suggest that integrating approaches from theoretical computer

science, non-equilibrium statistical physics, and cell biology stands to shed light on the

regulation of cellular behavior in eukaryotes more broadly. By revealing principles of cellular

behavior, the line of research established here stands to advance our ability to predict and even

one day engineer cellular behavior across diverse eukaryotic systems.

**References**

1.    Jennings, H. S. *Behavior of the lower organisms. Behavior of the lower organisms.*

(Columbia University Press, 1906). doi:10.1037/10817-000

2.    Reid, C. R. *et al.* Decision-making without a brain: how an amoeboid organism solves the

two-armed bandit. *J. R. Soc. Interface* **13**, 20160030 (2016).

3.    Coyle, S. M., Flaum, E. M., Li, H., Krishnamurthy, D. & Prakash, M. Coupled Active

Systems Encode an Emergent Hunting Behavior in the Unicellular Predator Lacrymaria

olor. *Curr. Biol.* **29**, 3838-3850.e3 (2019).

4.    Battle, C. *et al.* Broken detailed balance at mesoscopic scales in active biological systems.

*Science* **352**, 604–7 (2016).

5.    Wan, K. Y. & Goldstein, R. E. Time Irreversibility and Criticality in the Motility of a

Flagellate Microorganism. *Phys. Rev. Lett.* **121**, 58103 (2018).

6.    Woznica, A. *et al.* Bacterial lipids activate, synergize, and inhibit a developmental switch

in choanoflagellates. *Proc. Natl. Acad. Sci. U. S. A.* **113**, 7894–7899 (2016).

7.    Wan, K. Y. Coordination of eukaryotic cilia and flagella. *Essays Biochem.* **62**, 829–838

(2018).

8.    Nomura, M. & Ishida, K. I. Fine-structural Observations on Siliceous Scale Production

and Shell Assembly in the Testate Amoeba Paulinella chromatophora. *Protist* **167**, 303–

318 (2016).

9.    Bennett, R. R. & Golestanian, R. A steering mechanism for phototaxis in

Chlamydomonas. *J. R. Soc. Interface* **12**, 20141164–20141164 (2015).

10.   Machemer, H. The swimming cell and its world: Structures and mechanisms of orientation

in protists. *Eur. J. Protistol.* **37**, 3–14 (2001).

11.    Almagor, M., Ron, A. & Bar-Tana, J. Chemotaxis in Tetrahymena thermophila. *Cell*

*Motil.* **1**, 261–268 (1981).

12.    Berg, H. C. & Brown, D. A. Chemotaxis in Escherichia coli analysed by three-

dimensional tracking. *Nature* **239**, 500–504 (1972).

13.    Nakagaki, T., Yamada, H. & Tóth, Á. Maze-solving by an amoeboid organism. *Nature*

**407**, 470–470 (2000).

14.    Tweedy, L. *et al.* Seeing around corners: Cells solve mazes and respond at a distance

using attractant breakdown. *Science (80-. ).* **369**, (2020).

15.    Applewhite, P. B. Learning in Protozoa. in *Biochemistry and Physiology of Protozoa* 341–

355 (Elsevier, 1979). doi:10.1016/b978-0-12-444601-4.50018-7

16.    Dexter, J. P., Prabakaran, S. & Gunawardena, J. A Complex Hierarchy of Avoidance

Behaviors in a Single-Cell Eukaryote. *Curr. Biol.* **29**, 4323-4329.e2 (2019).

17.    Jennings, H. S. Studies on reactions to stimuli in unicellular organisms. IX—On the

behavior of fixed infusoria (Stentor and Vorticella), with special reference to the

modifiability of protozoan reactions. *Am. J. Physiol. Content* **8**, 23–60 (1902).

18.    Tang, S. K. Y. & Marshall, W. F. Cell learning. *Current Biology* **28**, R1180–R1184

(2018).

19.    Lyon, P. The cognitive cell: Bacterial behavior reconsidered. *Frontiers in Microbiology* **6**,

(2015).

20.    Marshall, W. F. Cellular Cognition: Sequential Logic in a Giant Protist. *Current Biology*

**29**, R1303–R1305 (2019).

21.    Greenfield, D. *et al.* Self-Organization of the Escherichia coli Chemotaxis Network

412  Imaged with Super-Resolution Light Microscopy. *PLoS Biol.* **7**, e1000137 (2009).

413 22. Wan, K. Y. & Goldstein, R. E. Coordinated beating of algal flagella is mediated by basal

414  coupling. *Proc. Natl. Acad. Sci. U. S. A.* **113**, E2784-93 (2016).

415 23. Höfer, T., Sherratt, J. A. & Maini, P. K. Dictyostelium discoideum : cellular self-

416  organization in an excitable biological medium. *Proc. R. Soc. London. Ser. B Biol. Sci.*

417  **259**, 249–257 (1995).

418 24. Kantsler, V., Dunkel, J., Polin, M. & Goldstein, R. E. Ciliary contact interactions

419  dominate surface scattering of swimming eukaryotes. *Proc. Natl. Acad. Sci. U. S. A.* **110**,

420  1187–1192 (2013).

421 25. Bray, D. *Wetware*. (Yale University Press, 2009).

422 26. Pfeifer, R., Lungarella, M. & Iida, F. Self-organization, embodiment, and biologically

423  inspired robotics. *Science* **318**, 1088–1093 (2007).

424 27. Regev, A. & Shapiro, E. Cellular abstractions: Cells as computation. *Nature* **419**, 343–343

425  (2002).

426 28. Mitchell, M. Biological Computation. *Comput. J.* **55**, 852–855 (2012).

427 29. Navlakha, S. & Bar-Joseph, Z. Algorithms in nature: the convergence of systems biology

428  and computational thinking. *Mol. Syst. Biol.* **7**, 546 (2011).

429 30. Savin, T. *et al.* On the growth and form of the gut. *Nature* **476**, 57–62 (2011).

430 31. Mehta, P. & Schwab, D. J. Energetic costs of cellular computation. *Proc. Natl. Acad. Sci.*

431  *U. S. A.* **109**, (2012).

432 32. Berg, H. C. & Purcell, E. M. Physics of chemoreception. *Biophys. J.* **20**, 193–219 (1977).

433 33. Bialek, W. & Setayeshgar, S. Physical limits to biochemical signaling. *Proc. Natl. Acad.*

434  *Sci. U. S. A.* **102**, 10040–5 (2005).

435    34.    Mehta, P. & Schwab, D. J. Energetic costs of cellular computation. *Proc. Natl. Acad. Sci.*

436           *U. S. A.* **109**, 17978–17982 (2012).

437    35.    Andrews, L. B., Nielsen, A. A. K. & Voigt, C. A. Cellular checkpoint control using

438           programmable sequential logic. *Science (80-. ).* **361**, (2018).

439    36.    Gilbert, J. J. Jumping behavior in the oligotrich ciliates Strobilidium velox and Halteria

440           grandinella, and its significance as a defense against rotifer predators. *Microb. Ecol.* **27**,

441           189–200 (1994).

442    37.    Lueken, W., Ricci, N. & Krüppel, T. Rhythmic spontaneous depolarizations determine a

443           slow-and-fast rhythm in walking of the marine hypotrich Euplotes vannus. *Eur. J.*

444           *Protistol.* **32**, 47–54 (1996).

445    38.    Erra, F., Iervasi, A., Ricci, N. & Banchetti, R. Movement of the cirri during the creeping

446           of Euplotes crassus (Ciliata, Hypotrichida). *Can. J. Zool.* **79**, 1353–1362 (2001).

447    39.    Taylor, C. V. The neuromotor system of Euplotes. *Exp. Biol. Med.* **16**, 101–103 (1919).

448    40.    Syberg-Olsen, M. J. *et al.* Biogeography and character evolution of the ciliate genus

449           Euplotes (Spirotrichea, Euplotia), with description of Euplotes curdsi sp. nov. *PLoS One*

450           **11**, (2016).

451    41.    Ricci, N., Giannetti, R. & Miceli, C. The ethogram of Euplotes crassus (ciliata,

452           hypotrichida): I. The wild type. *Eur. J. Protistol.* **23**, 129–140 (1988).

453    42.    Li, Y., Shi, L. & Gu, F. Comparison of ciliature microtubule organelles in three

454           hypotrichous ciliate species*. *Chinese J. Oceanol. Limnol.* **28**, 443–450 (2010).

455    43.    Fleury, A. Dynamics of the cytoskeleton during morphogenesis in the ciliate Euplotes I.

456           Basal bodies related microtubular system. *Eur. J. Protistol.* **27**, 99–114 (1991).

457    44.    Purcell, E. M. Life at low Reynolds number. *Am. J. Phys.* **45**, 3–11 (1977).

458    45.    Kimmel, J. C., Chang, A. Y., Brack, A. S. & Marshall, W. F. Inferring cell state by

459            quantitative motility analysis reveals a dynamic state system and broken detailed balance.

460            *PLoS Comput. Biol.* **14**, e1005927 (2018).

461    46.    Chang, A. Y. & Marshall, W. F. Dynamics of living cells in a cytomorphological state

462            space. *Proc. Natl. Acad. Sci. U. S. A.* **116**, 21556–21562 (2019).

463    47.    Boltzmann, L. Weirere Studien uber das warmegleich-gewich unter gasmolekulen.

464            *Sitzungsberichte der Akad. der Wissenschaften, Vienna, part II* **66**, 275–370 (1872).

465    48.    Alberts, B. *et al. Molecular biology of the cell*. (Garland Science, 2002).

466    49.    Shalizi, C. R. & Crutchfield, J. P. Computational Mechanics: Pattern and Prediction,

467            Structure and Simplicity. *J. Stat. Phys.* **104**, 817–879 (2001).

468    50.    Crutchfield, J. P. The calculi of emergence: computation, dynamics and induction. *Phys.*

469            *D Nonlinear Phenom.* **75**, 11–54 (1994).

470    51.    Hopcroft, J. E., Motwani, R. & Ullman, J. D. *Introduction to Automata Theory,*

471            *Languages, and Computation*. (Pearson Education, 2008).

472    52.    Moore, T. Y., Cooper, K. L., Biewener, A. A. & Vasudevan, R. Unpredictability of escape

473            trajectory explains predator evasion ability and microhabitat preference of desert rodents.

474            doi:10.1038/s41467-017-00373-2

475    53.    Goldstein, R. E. Green Algae as Model Organisms for Biological Fluid Dynamics. *Annu.*

476            *Rev. Fluid Mech.* **47**, 343–375 (2015).

477    54.    Simon, M. & Plattner, H. Unicellular Eukaryotes as Models in Cell and Molecular

478            Biology: Critical Appraisal of Their Past and Future Value. in *International Review of*

479            *Cell and Molecular Biology* **309**, 141–198 (Elsevier Inc., 2014).

55.    Durham, W. M. *et al.* Turbulence drives microscale patches of motile phytoplankton. *Nat.*

*Commun.* **4**, (2013).

56.  Mogilner, A. & Oster, G. Cell motility driven by actin polymerization. *Biophys. J.* **71**,

3030–3045 (1996).

57.  Collins, J. J. & Stewart, I. N. Coupled nonlinear oscillators and the symmetries of animal

gaits. *J. Nonlinear Sci.* **3**, 349–392 (1993).

58.  McNeill Alexander, R. *Principles of Animal Locomotion*. (Princeton University Press,

2003).

59.  Wan, K. Y. & Jékely, G. On the unity and diversity of cilia. *Philos. Trans. R. Soc. Lond.*

*B. Biol. Sci.* **375**, 20190148 (2020).

60.  Ahamed, T., Costa, A. C. & Stephens, G. J. Capturing the Continuous Complexity of

Behavior in C. elegans. doi:10.1101/827535

61.  Stephens, G. J., Johnson-Kerner, B., Bialek, W. & Ryu, W. S. Dimensionality and

Dynamics in the Behavior of C. elegans. *PLoS Comput. Biol.* **4**, e1000028 (2008).

62.  Berman, G. J., Choi, D. M., Bialek, W. & Shaevitz, J. W. Mapping the stereotyped

behaviour of freely moving fruit flies. *J. R. Soc. Interface* **11**, 20140672 (2014).

63.  Stephens, G. J., Osborne, L. C. & Bialek, W. Searching for simplicity: Approaches to the

analysis of neurons and behavior. *Proc. Natl. Acad. Sci. U. S. A.* **108**, 15565–15571

(2010).

64.  Kanagal, B. & Sindhwani, V. Rank Selection in Low-rank Matrix Approximations: A

Study of Cross-Validation for NMFs. in *Advances in Neural Information Processing*

*Systems* **23**, (2010).

65.  Owen, A. B. & Perry, P. O. Bi-Cross-Validation of the SVD and the Nonnegative Matrix

Factorization. *Source Ann. Appl. Stat.* **3**, 564–594 (2009).

66.    Ester, M., Kriegel, H.-P., Sander, J. & Xu, X. *A Density-Based Algorithm for Discovering*

505           *Clusters in Large Spatial Databases with Noise*. (1996).

506    67.    Kauffman, L. & Rousseeuw, P. J. *Finding Groups in Data: and introduction to cluster*

507           *analysis*. (John Wiley & Sons, Inc., 1990). doi:10.1002/9780470316801

508    68.    Kirkegaard, J. B., Marron, A. O. & Goldstein, R. E. Motility of Colonial

509           Choanoflagellates and the Statistics of Aggregate Random Walkers. *Phys. Rev. Lett.* **116**,

510           038102 (2016).

511    69.    M Cartmill. Climbing. in *Functional Vertebrate Morphology* 73–88 (Harvard University

512           Press, 1985).

513    70.    Markov, A. A. Extension of the Limit Theorems of Probability Theory to a Sum of

514           Variables Connected in a Chain. *Notes Imp. Acad. Sci. St. Petersbg. VIII Ser. Physio-*

515           *Mathematical Coll.* **22**, (1907).

516    71.    Howard, R. A. *Dynamic and Probabilistic Systems, Volume 1: Markov models*. (Dover

517           Publications, 1971).

518    72.    Gagniuc, P. *Markov Chains: From Theory to Implementation and Experimentation*.

519           (Wiley, 2017).

520    73.    Billingsley, P. *Statistical Methods in Markov Chains*. *The Annals of Mathematical*

521           *Statistics* **32**, 12–40 (Institute of Mathematical Statistics, 1961).

522    74.    Crutchfield, J. P. & Young, K. Computation at the Onset of Chaos. in *Complexity,*

523           *Entropy, and Physics of Information* (ed. Zurek, W.) (Addison-Wesley, 1989).

524    75.    Crutchfield, J. P. & Young, K. Inferring statistical complexity. *Phys. Rev. Lett.* **63**, 105–

525           108 (1989).

76.    Taylor, C. V. Demonstration of the function of the neuromotor apparatus in Euplotes by

the method of microdissection. *Naturwissenschaften* **9**, 910 (1921).

77.    Okajima, A. & Kinosita, H. Ciliary activity and coordination in Euplotes eurystomus-I.

Effect of microdissection of neuromotor fibres. *Comp. Biochem. Physiol.* **19**, 115–131

(1966).

78.    Bar-Ziv, R., Tlusty, T. & Libchaber, A. Protein-DNA computation by stochastic assembly

cascade. *Proc. Natl. Acad. Sci. U. S. A.* **99**, 11589–11592 (2002).

79.    Guasto, J. S., Rusconi, R. & Stocker, R. Fluid Mechanics of Planktonic Microorganisms.

*Annu. Rev. Fluid Mech.* **44**, 373–400 (2012).

80.    Goldstein, R. E., Polin, M. & Tuval, I. Noise and Synchronization in Pairs of Beating

Eukaryotic Flagella. *Phys. Rev. Lett.* **103**, 168103 (2009).

81.    Párducz, B. Ciliary Movement and Coordination in Ciliates. *Int. Rev. Cytol.* **21**, 91–128

(1967).

82.    Funfak, A. *et al.* Paramecium swimming and ciliary beating patterns: A study on four

RNA interference mutations. *Integr. Biol. (United Kingdom)* **7**, 90–100 (2015).

83.    Kirkegaard, J. B., Bouillant, A., Marron, A. O., Leptos, K. C. & Goldstein, R. E.

Aerotaxis in the closest relatives of animals. *Elife* **5**, e18109 (2016).

84.    Miño, G. L., Koehl, M. A. R., King, N. & Stocker, R. Finding patches in a heterogeneous

aquatic environment: pH-taxis by the dispersal stage of choanoflagellates. *Limnol.*

*Oceanogr. Lett.* **2**, 37–46 (2017).

85.    Polin, M., Tuval, I., Drescher, K., Gollub, J. P. & Goldstein, R. E. Chlamydomonas

Swims with Two 'Gears' in a Eukaryotic Version of Run-and-Tumble Locomotion.

*Science (80-. ).* **325**, 487–490 (2009).

86.    Keeling, P. J. Combining morphology, behaviour and genomics to understand the

evolution and ecology of microbial eukaryotes. *Philos. Trans. R. Soc. B Biol. Sci.* **374**,

(2019).

87.   Kahl, A. *Urtiere oder Protozoa.* (Fischer, 1935).

88.   Yocom, H. B. The neuromotor apparatus of Euplotes patella. (University of California,

Berkeley, 1917).

89.   Kawamoto, T. & Kabashima, Y. Cross-validation estimate of the number of clusters in a

network. *Sci. Rep.* **7**, 1–17 (2017).

**Figures**

**Figure 1.** *Euplotes* exhibits highly polarized, complex cellular architecture and walks

across surfaces using microtubule-based organelles called cirri, some of which are physically

linked. Scale bar is 10 μm. **a**, A single *Euplotes eurystomus* cell in profile displays its ventral

cirri, which are used for walking locomotion across surfaces (arrowhead indicates a single cirrus

stretching out from the cell). **b**, A drawing of a *Euplotes* cell, viewed from the ventral surface,

highlighting the complex, asymmetric structure of the cell. Notable features include the cirri (ci)

and the membranellar band (m), wrapping from the top of the cell to the center, which is used to

generate a feeding current to draw in prey items. Drawing adapted and obtained from Wikimedia

Commons, from original source [87]. **c**, A drawing of a *Euplotes* cell, highlighting the fiber system

associated with the cirri, historically referred to as the neuromotor apparatus. Drawing adapted

from [88].

**Figure 2.** The gait of *Euplotes* can be described in a discrete, reduced state space with gait

states corresponding to identifiable patterns of cirral activity. **a-a''**, The movements of cirri

during walking locomotion are clearly visible by brightfield microscopy by focusing on a plane

at the surface of the coverslip on which cells are walking. Three snapshots depict different time

points during a single walking trajectory, and white arrowheads indicate cirri. In the panels from

left to right, the cirrus indicated by the arrowhead on the left is stationary, stationary, and then

moving, and the cirrus indicated by the arrowhead on the right is stationary, moving, and then

stationary. Scale bar is 15 μm. **b**, The trajectory of a cell during a single recorded trajectory as

the cell walked across a coverslip from left to right. The cell position was manually tracked in

each frame. **c**, The scheme for encoding cirral dynamics during walking involved labeling each

of the 14 distinguishable ventral cirri (a-n), and recording cirral activity in each frame,

corresponding to timepoints $(t_1,...,t_n)$, of recordings of walking cells as a 14-bit binary vector.

Each entry in each vector is given a value of either 0 if the cirrus is not moving and in contact

with the coverslip or 1 if the cirrus is moving. Scale bar is 15 μm. **d**, Representative visualization

of cirral dynamics for a single trajectory of a walking cell. These dynamics correspond to the

walking trajectory in **b**. Each row corresponds to a cirrus and each column is a single video

frame. White denotes cirral activity, a value of 1, in the vector encoding of dynamics from **c**.

Note the dynamical complexity and discrete, stepping-like nature of cirral movements. **e-g**,

Three roughly orthogonal views of a plot displays the structure of all recorded cirral dynamics

encoded as in Figure 2C from 13 cells over 2343 timepoints in a reduced state space obtained by

non-negative matrix factorization (NMF). Axes correspond to the components of the NMF (H1,

H2, H3), and each point is a single timepoint. Randomized colors highlight the 32 clusters

identified using the density-based spatial clustering of applications with noise (DBSCAN)

algorithm [66]. We refer to these clusters as gait states, and they correspond to unique

configurations of cirral activity during walking locomotion (see panel F). **h**, Plot of the

magnitudes associated with each cirrus corresponding to the components of the NMF of cirral

dynamics shows distinct contributions from spatially distinct groups of cirri. Component H1, for

example, is associated with activity in cirri a, b, and c. The tracing of a cell including the position

of cirri has the same color map as the plot above and shows the grouping of the cirri

corresponding to each component. **i**, A heatmap of mutual information between all pairs of cirri

shows that correlations in cirral activity correspond to the NMF components displayed in **d**. For

example, cirri a, b, and c share mutual information with one another and are the cirri contributing

to component H1. **j**, A heatmap representation of the cirral activity associated with each of the 32

gait states. Values for each cirrus are the mean over all instances of the gait state. Note that each

gait state has a unique signature of cirral activity.

**a**

Scaled cell velocity vs Net change in cirral activity

**b**

Start gait state vs End gait state — Transition rate (1/s)

**c**

**d** Unbalanced Transitions

**e** Balanced Transitions

**f** Gait state / Gait state — Mean cirral activity

**g** Start gait state vs End gait state — Transition probability

**h**

Resting/start state
2nd cycle state
Return to 1st cycle state
Substantial cell movement
1st cycle state
3rd cycle state
Return to 3rd cycle state

**Figure 3.** *Euplotes* walks with a cyclic stochastic gait exhibiting broken detailed balance,

stereotypy, and state machine-like dynamics. All data is pooled from the walking trajectories of

13 different cells over 2343 timepoints and 1423 pairwise gait state transitions. **a**, A plot of the

mean net change in cirral activity versus the net scaled cell velocity associated with all

transitions between the 32 gait states identified in Fig. 3 shows that the change in number of

active cirri is not strongly correlated with cell velocity ($R^2$=0.03). Cell velocities were obtained

from manually tracked walking trajectories and then scaled by dividing frame to frame

displacements for each trajectory by the length of the cell being tracked and also dividing by the

average frequency of cirral inactivity. Scaling provided a non-dimensional velocity scaled by

natural units of the system. Note that at low Reynolds number, velocity should be proportional to

force [44], so this plot also reflects the net walking force generated by the cell. Net change in cirral

activity was computed using the data presented in Fig. 3F. Note that the largest velocities are

associated with small negative and small to moderate positive net changes in cirral activity. **b**,

The transition matrix of all gait state transitions, with rows representing the starting state and

columns indicating the ending state, exhibits broken detailed balance. Rates were estimated by

dividing the total number of observed transitions between each state pair and dividing by the

total time observed. Under detailed balance or equilibrium conditions, transitions from one state

to another should be balanced by reverse transitions. Lack of this kind of reversibility, as seen by

the lack of symmetry of the heatmap across the diagonal, indicates broken detailed balance and

non-equilibrium dynamics. **c**, A directed graph representation of all gait state transitions. Nodes

correspond to the 32 gait states, and node sizes are scaled by the proportion of total time cells

spent in each state. Directed edges are represented by arrows between nodes and signify state

transitions. The size of the arrows is scaled by transition rates as in **b**. Edge color represents

scaled cell velocity as in **a**, with cool colors (more cyan) representing lower velocity, and

warmer colors (more magenta) representing higher velocity. **d**, A subset of transitions visualized

as in **c** shows the restricted and relatively high frequency nature of unbalanced, non-equilibrium-

like transitions. Only transitions that were observed to happen more than one time and exhibiting

a significant difference between forward and reverse transitions ($p < 0.05$ by binomial test, see

Methods for more details) are displayed. **e**, A subset of transitions, similarly to panel E, except

that only the balanced transitions, lacking a significant difference between forward and reverse

transitions ($p < 0.05$ by binomial test) are displayed, also show a complex and widespread

structure, this time of balanced, equilibrium transitions. Note that the majority of transitions

associated with high cell velocity involve equilibrium-like dynamics. **f,** Examples illustrating the

spatial organization of cirral activity corresponding to gait states. Some states, such as 7,

correspond to activity in spatially discrete groups of cirri, while others, such as 17, correspond to

cirral activity across the cell. The gait states displayed here are those involved in unbalanced

transitions. **g**, A heatmap of transition probabilities between states, showing only the most

probable transitions from a given state with all others set to zero, shows distinct structure. In

cases were multiple state transitions from a state were tied for the highest probability, all of these

transitions are displayed. Fewer than half of the total states are recipients of multiple high

probability transitions, and many states are the recipients of no high probability transitions. **h**, A

representation of functional states and transitions between them highlights the machine-like

nature of the gait of *Euplotes*. Gait states are represented as circles with numerical labels. Blue

circles represent states that are both recipients and sources of unbalanced transitions as identified

in **d** and constitute the three cycle states. Red circles represent states that are recipients but not

sources of unbalanced transitions as identified in **d**. Black circles correspond to gait states that

are associated only with balanced transitions as in **e**. States receiving no more than one unique

high probability transitions from states with only a single highest as identified in **g** were grouped

together into a compound state represented by the dark gray blob. The blue background behind

states 4, 16, 18, 26, and 28 indicates that these states all share the same highest probability

transitions between states identified in this panel, and thus, the group constitutes a single

compound functional state. Arrows represent the highest probability transitions between the

states, including compound states composed of multiple gait states as identified in Fig. 2 and 3

(dark gray blob and blue background). Gait state 1 is also depicted, as it is the state in which

cells spent the most time over all walking trajectories and also is uniquely the state from which

cells begin walking. Cells also frequently return to the state during walking. Further, transitions

from gait state 2 from gait state 1 constitute the single highest frequency transition. Together, all

identified states in this panel constitute functional states. Arrows represent the most probable

transitions between functional states, and all unbalanced transitions are also represented with size

scaled by their proportional probability compared to all other transitions emanating from the

source functional state. Cartoons are a walking cell in profile with cirri in a configuration

representative of the corresponding functional state. Labels refer to the apparent functional role

of states and their associated transitions. Beginning from gait state 1, the resting/start state, cells

are most likely to follow transitions from gait state 2 to 3 to 17 at which point cells are likely to

enter the functional state associated with substantial cell movement involving variable balanced

transitions between a number of gait states. Transitions are then likely to lead back toward the

cycle states. Note that while this representation of gait dynamics highlights the most probable

transitions, substantial variability, primarily involving reversible transitions, occurs during

walking trajectories.

**Figure 4.** The fiber system of *Euplotes* is complex, interconnected, and constrains

mechanistic hypotheses of gait coordination. **a**, The SiR-tubulin labeled cell (faint, dark blue)

was imaged by confocal microscopy, and a 3D reconstruction as obtained from serial confocal

slices. Fibers were manually traced in each slice using TrakEM2 in FIJI. Two morphologically

distinct classes of fibers were observed and are indicated as follows: thick, linear fibers are cyan

and thinner, filamentous fibers are magenta (see Fig. S6 for raw image data). Fibers emanate

from the base of each cirrus and form a connected network between all cirri. The base of each

cirrus is indicated by corresponding letters (as in Fig. 2c). Gray shading indicates the dynamical

groups identified by dimensionality reduction and follows the same color scheme as in Fig. 3d.

Scale bar is 10 μm. **b**, A graph representation of fiber-fiber connections illustrates the complex

and interconnected nature of cirrus associated fiber topology. Nodes correspond to the cirri to

which each fiber system is associated, and edges indicate connections between fiber systems.

Colors of nodes indicate the same groups as in **a**, and colors of edges indicate the types of fibers

connecting to one another, cyan for thick fiber connections, magenta for thin fiber connections,

and purple for thick to thin fiber connections. **c**, Pairs of cirri that are linked by fiber-fiber

contacts show no statistically significant difference in mutual information compared to those

lacking fiber-fiber contacts. The plot displays mutual information between all pairs of cirri

grouped by the absence (Not linked) or presence (Linked) of associated fiber-fiber connections.

Statistical significance was evaluated by the Wilcoxon rank sum test. Note that when pairs of

cirri are grouped by fiber-fiber connection type, we do observe an increase in mutual information

for cirri associated with only thin fiber-fiber connection and only thick fiber-fiber connections

compared to those lacking fiber-fiber connections (see Fig. S6). **d**, A plot of mutual information

as a function of inter-cirrus distance displays negative correlation, with a Spearman correlation

coefficient of -0.49 (p<0.001). Plotted values are defined with respect to pairs of cirri. **e**, A plot

of mutual information as a function of fiber-cortex contact distance grouped by fiber type

similarity and lack thereof displays negative correlation, with a Spearman correlation coefficient

of -0.62 (p<0.001) for pairs of cirri with similar fiber types and no significant correlation for

those with dissimilar fiber types. Similarity of fiber types is defined in terms of sharing at least

some fiber types as defined in **a**. Fiber-cortex contact difference is measured by the mean cross

nearest neighbor distance (see Methods) for all fiber-cortex contact points associated with each

cirrus. The negative correlation values from the data plotted in **d** and **e** indicate that cirri that are

closer to one another and also cirri with fiber-cortex contacts in nearby regions of the cell tend to

have higher mutual information, and indeed cirri that are both close to one another and with

similar patterns of fiber-cortex contacts display the highest mutual information. **f**, A plot of fiber-

cortex contact difference versus inter-cirrus difference (as in panels **d** and **e**) illustrates that

nearby cirri tend to have similar associated fiber-cortex contacts, highlighting that nearby cirri

with similar fiber-cortex contacts share the most mutual information.

**Methods**

*Cell husbandry*

Cultures of *Euplotes eurystomus* were obtained from Carolina Biological Supply Company (Item

#131480). Individual cells were isolated from cultures, which contained other protists and

meiofauna, by pipetting and placed in non-treated 6-well plates (Thermo Fischer Scientific 08-

772-49) containing spring water taken from cultures. Cells were kept in wells for no longer than

five days before imaging, and if cells were to be kept for longer than 48 hours, wells containing

cells were supplemented with 1% Cereal Grass Medium [1] (from Thermo Fischer Scientific

S25242) to prevent depletion of a population of prey bacteria and otherwise maintain *Euplotes*

under constant growth conditions.

*Live cell brightfield microscopy*

Cells were concentrated by centrifugation (500×g for 5 min) and resuspended either in 0.5 mL of

spring water in coverglass bottomed FluoroDishes (World Precision Instruments FD35-100) or

in 0.2 mL spring water on a coverslip (FisherScientific, 12-545-D) for imaging. No more than

three cells were kept in 0.5 mL imaging samples and only one cell was ever kept in 0.2 mL

imaging samples in order to minimize cell-cell interactions. Cells were observed to exhibit

spontaneous walking activity on coverglass. Walking cells in FluoroDishes were imaged under

brightfield illumination using a Zeiss Z.1 Observer and Hamamatsu Orca Flash 4.0 V2 CMOS

camera (C11440-22CU) with a 20x, 0.8 NA Plan-Apochromat (Zeiss) objective. Cells on

coverslips were imaged under brightfield illumination with coverslips inverted over a well

containing a small amount of distilled water to reduce evaporation using a Zeiss Axio Zoom.V16

and a PCO pco.dimax S1 camera. Importantly, in both imaging systems, the focal plane was set

at the interface between cirri of walking cells and the glass surface upon which they were

walking. Images were acquired at 0.033 seconds per frame with a 0.005 second exposure in order

to capture all cirral dynamics during walking with minimal blur.

*Quantification of walking dynamics*

746           Movies of walking cells were viewed using FIJI [2]. Movement of cirri, or lack thereof was

clearly visible in each movie frame (see Fig. 2a and Movie S1). The dynamical state of each

cirrus in each movie frame was manually annotated. For each frame, each cirrus received a label

of "1" if the cirrus was in motion and "0" if the cirrus was not moving and in contact with the

coverslip. Motion of cirri was evident in terms of a change in cirrus shape or tip position often in

addition to blur due to motion during image acquisition or position out of the focal plane (see

Fig. 2a and Movie S1). While only slowly walking cells were recorded, sometimes cells

nevertheless exhibit brief, spontaneous departures from slow walking during the course of movie

acquisition. Any frame in which the movement of the cell and/or cirri were too fast to be

resolved, such as during spontaneous escape responses[3] (Movie S2), was excluded from analysis

such that some videos were split into a number of separate continuous sequences. Thus, each

movie frame associated with a particular time point in the walking trajectory, with the exception

of those excluded from analysis as described, yielded a corresponding 14-element binary vector

encoding the motility state of the cell in terms of the movement of cirri. Cell movement was

tracked using the manual tracking feature of the TrackMate plugin in FIJI [4]. The center of each

cell was used as the reference feature for tracking. We analyzed the walking dynamics of 13

different cells.

***Dimensionality reduction***

Dimensionality reduction was performed by non-negative matrix factorization (NMF)

implemented in MATLAB release 2019b (Mathworks, Natick). NMF was chosen as a

dimensionality reduction technique to allow us to obtain a reduced, sparse, and interpretable

representation of walking dynamics. Because NMF derives non-negative factors, the basis

vectors in NMF space correspond to patterns of cirral activity. NMF involves factoring data, $A$,

an $n$ by $m$ matrix, into non-negative factors $W$, an $n$ by $k$ matrix, and $H$, a $k$ by $m$ matrix where

the product $W*H$ approximates $A$. To determine the appropriate number of dimensions or rank, $k$,

that are necessary to accurately represent the data without overfitting, we performed cross-

validation by imputation with random holdouts [5,6], also implemented in MATLAB. We randomly

held out 15% of our walking dynamics data, performed NMF for a given $k$, and then used the

NMF reconstruction $W*H$, to update the missing data entries. This process of updating is known

as imputation, and we repeated the imputation process 50 times, by which point the imputed

values had stabilize, to obtain a final NMF reconstruction. We then computed the root mean

squared residual (RMSR) between the final NMF reconstruction, $W*H$, and our dataset, $A$. We

performed this entire process 100 times for each value of $k$. As is generally the case for NMF, we

observed a monotonic decrease in reconstruction error with increasing $k$ without performing the

imputation procedure [7] (Fig. S2a). In contrast to this trend, we observed an increase in RMSR of

imputed values with increasing $k$ indicating overfitting [5] (Fig. S2b). We chose $k=3$ because this

value was the highest value before a notable increase in imputation error (Fig. S2b), which

would indicate overfitting [5,6]. Thus, our choice of rank 3 selects the lowest rank approximation

that captures structure of the dataset without overfitting that structure. Further, our choice

facilitated the visual inspection of the structure of data in the reduced dimensional

reconstruction.

Finally, we noted that for our chosen value of $k$, due to the stochastic nature of the NMF

algorithm, which involves a random initialization step, we obtained slightly different solutions

for different iterations [5]. In order to choose the best reduced dimensional approximation,

therefore, we performed NMF 500 times and chose the particular solution corresponding to the

lowest RMSR compared to our dataset.

***Clustering***

Clustering on the dataset obtained using NMF was performed by density-based spatial

clustering of applications with noise (DBSCAN) algorithm [8] implemented in MATLAB release

2019b (Mathworks, Natick). Structure in NMF space was clearly visible (Fig. 2e-g), and

DBSCAN using a Euclidean distance metric, was initially chosen as a clustering method because

it yielded qualitatively good partitioning of the data. The DBSCAN algorithm involves stochastic

search within neighborhoods of a given radius $\varepsilon$ around datapoints, and points with a minimum

number of neighbors, $n$, within their neighborhood are grouped as belonging to the same cluster,

leaving two free parameters to determine. We set $\varepsilon$ by first using the

clusterDBSCAN.estimateEpsilon function in MATLAB (release 2020b, Phased Array System

Toolbox), which yielded a value of 0.15. We next set about determining the minimum neighbor

number, $n$. To do so, we computed the average Silhouette coefficient, a commonly used measure

of clustering quality that indicates how well-separated clusters are [9], for various values of $n$. The

results of this analysis are plotted in Fig. S3. Higher Silhouette coefficients indicate better

clustering, and we found that a value of $n=8$ maximized the mean Silhouette coefficient (Fig.

S3a). We also noted, however, that for this value, many datapoints were found to be outliers, not

belonging to any cluster due to having too few points within a distance of ε. Fig. S3b displays

percentage of datapoints found to be outliers as a function of $n$. In order to avoid categorizing

more than 5% of datapoints as outliers, we chose to settle on $n=4$, which does not have a

significantly different mean Silhouette coefficient compared to any of the others in the range

$n=2$-7. This choice was further supported by the fact that major clusters involving more than 5

datapoints identified with $n=8$ were also identified with $n=4$.

Although this set of parameters gave qualitatively and quantitatively reasonable

clustering results, we sought to further refine our clusters and to further reduce the outlier

datapoints. We noted the obvious partitioning of the NMF dataset into three groups along the H2

axis (Fig. 2e). We found the previously determined parameter values to yield good clustering for

the top and middle partitions (H2≤1.1 and 0.2<H2<1.1), with no outliers. For the lower partition

(H2≤0.2), however, we found that we were able to improve clustering by using ε=0.1182. With

this updated value, we found no statistically significant change in Silhouette coefficient and

reduced outliers to 0%. The clusters obtained by this process constituted the identification of the

32 gait states. We note here that the problem of determining the true or optimal number of

clusters is an unresolved problem [10], and we note that we have followed standard methods to

determine cluster number, and we found that our key results do not depend sensitively on the

precise number of clusters identified (see following section and Fig. S8 for more details).

***State transition analysis***

Following dimensionality reduction and clustering to identify gait states, we proceeded to

characterize state transition dynamics. For each cell trajectory, we identified all unique gait state

transitions for a total of 1423 unique pairwise transitions over the cumulative 2343 video frames

for 77.14 s of recording. We computed empirical transition rates between states as the total

number of observed transitions divided by the total time of observation. In order to determine

which transitions were balanced and which were unbalanced, we followed Chang and Marshall

[11], and performed binomial tests of statistical significance. Assuming a system at equilibrium,

with all transitions obeying detailed balance, we expect to observe some deviation from exactly

reciprocal transitions and can calculate the probability of observing a given set of ratios given

underlying probabilities of forward and reverse transitions. The binomial probability of

observing a set of transitions with known forward and reverse probabilities is given by

841
$$P(X = f) = \binom{n}{f} p_{forward}^{f} p_{reverse}^{n-f}$$

where $\binom{n}{f} = \frac{n!}{f!(n-f)!}$ is the choose function, $f$ is the number of forward transitions, $n$ is the total

number of transitions (such that $n$-$f$ is the number of reverse transitions), and the probabilities

$p_{forward}$ and $p_{reverse}$ are the forward and reverse probabilities. Considering only the set of

transitions involving a specific pair of states, and calculating the probability that a transition

between those states is either in the forward or reverse direction, the values of forward and

reverse probabilities in the balanced case must be equal such that $p_{forward} = p_{reverse} = 0.5$.

With an α level of 0.05, we then considered reciprocal transition pairs with binomial

probabilities less than 0.05 to be significantly unbalanced. Figure S9 displays the binomial

probabilities associated with all transitions.

851            In order to calculate the estimated entropy production rate, we followed Wan and

852    Goldstein [12], where the entropy production rate is defined as

853
$$\dot{S} = \frac{1}{2} \sum_{i \neq j} J_{ij} A_{ij}$$

with conjugate fluxes $J_{ij} = p_i k_{ij} - p_j k_{ji}$ and forces $A_{ij} = \ln\left(\frac{p_i k_{ij}}{p_j k_{ji}}\right)$ where the $p_l$ are the

probabilities of being in state $l$ at steady state and the $k_{ij}$ are the transition rates between states $i$

and $j$. We estimate the state occupancy probabilities $p_l$ as $\frac{T_l}{T_{Total}}$, where $T_l$ is the amount of time

spent in state $l$ over all trajectories and $T_{Total}$ is the total recorded time, and the rates $k_{ij}$ as $\frac{N_{ij}}{T_i}$,

where $N_{ij}$ is the total number of observed transitions from state $i$ to state $j$ and $T_i$ is the total time

spent in state $i$. To avoid $k_{ji} = 0$, we let $k_{ji} = \frac{1}{p_j T_{max}}$ where $T_{max} = 11.55$ seconds is the

maximum duration of any single recorded walking trajectory.

In the course of our state transition analysis, we also checked whether the waiting times

between instances of each state might be non-exponentially distributed, with exponential

distributions indicative of an embedded Markov process or possibly self-organized criticality

[13,14]. Using the Lilliefors test implemented in Matlab, we found that in general, waiting times

were not exponentially distributed, although states 2, 3, 6, 16, 17, 18, 25, 27, 28, 32 were found

to have waiting times consistent with exponential distributions with Benjamini-Hochburg

corrected p-values of 0.046, 0.046, 0.022, 0.008, 0.046, 0.017, 0.046, 0.0081, 0.0046, 0.0046

respectively. Interestingly, none of the waiting times between the movements of individual cirri

were found to be consistent with exponential distributions. These results are consistent with

mechanisms constraining the temporal dynamics of cirri and state transitions.

In order to begin evaluating whether state transitions obeyed the Markov property, where

the transition probabilities from one state to the next are completely determined by current state

[15,16], we estimated the transition matrix for walking dynamics, consisting of the transition

probabilities between all states. We estimate the transition probability from state $i$ to state $j$ as

$p_{ij} = \frac{N_{ij}}{\sum_k N_{ik}}$ such that $\sum_i p_{ij} = 1$. The entries of the transition matrix, $P$, are these transition

probabilities with indices $i$ for rows and $j$ for columns. If gait state transitions obeyed the Markov

property, we expect that the product of the transition matrix with itself, $P^2$, would be equivalent

to the two-step transition matrix where transition probabilities are computed as before except that

state $j$ is the state to which $i$ has transitioned after an intervening transition. Figure S5 displays

the results of this analysis showing that the two matrices show substantial quantitative and

qualitative differences. Although these results strongly suggest violation of the Markov property,

we applied the Billingsley test for a more statistically rigorous evaluation [17,18]. This test was

implemented and performed in Matlab (release 2019b). The Billingsley test gives a $\chi^2$ metric

with $M^2$-$2M$ degrees of freedom given by

$\sum_{i=1}^{M} \sum_{j=1}^{M} \frac{\left(N_{ij} - R_{ij} \sum_{j=1}^{M} N_{ij}\right)^2}{R_{ij} \sum_{j=1}^{M} N_{ij}},$

where $R_{ij}$, the independent trials probability matrix, is given by

$R_{ij} = \sum_{k=1}^{M} N_{kj} / \left(\sum_{h=1}^{M} \sum_{l=1}^{M} N_{hl} - \sum_{k=1}^{M} N_{ik}\right).$

Importantly, we also noted that the key qualitative results of our state transition analysis

are robust to the details of clustering results. In particular, we find that strongly unbalanced

transitions and violation of the Markov property exist for a range of clustering parameters.

Figure S8 displays the transition matrices for different clustering results.

***Confocal microscopy***

Cells were prepared for imaging and placed into a FluoroDish as described in the Live Cell

Brightfield Microscopy section. Cells were then labeled with SiR-tubulin (Spirochrome provided

by Cytoskeleton, Inc, CY-SC002) at 1 µM concentration. Cells were imaged using a Zeiss LSM

880 AxioExaminer and a 40x, 1.2 NA C-Apochromat water immersion objective (Zeiss) and

excitation provided by a 633 nm laser (Zeiss). Only one full confocal z-stack of a complete cell

was obtained during imaging to avoid effects of photodamage.

*Fiber reconstruction and analysis*

The image stack resulting from confocal imaging was first aligned in FIJI using the

StackReg plugin [2]. Next, fibers were manually segmented in each of the aligned z-stack images

using the TrakEM2 plugin in FIJI [19,20]. Thick and thin fibers (Fig. 5a) were morphologically

distinguished, with thick fibers having a diameter of no less than 5 μm at the thinnest point.

Fibers were traced from their distal termini to their convergences at the base of the cirri with

which they were associated. Following segmentation, 3D surfaces were reconstructed in

TrakEM2. Inter-fiber contacts were then found by inspection of 3D reconstructions and verified

by examining individual z-stack frames to confirm intersections between fibers.

**References**

1.     King, N., Young, S. L., Abedin, M., Carr, M. & Leadbeater, B. S. C. Starting and

maintaining Monosiga brevicollis cultures. *Cold Spring Harb. Protoc.* **4**, (2009).

2.     Schindelin, J. *et al.* Fiji: an open-source platform for biological-image analysis. *Nat.*

*Methods* **9**, 676–682 (2012).

3.     Ricci, N., Giannetti, R. & Miceli, C. The ethogram of Euplotes crassus (ciliata,

hypotrichida): I. The wild type. *Eur. J. Protistol.* **23**, 129–140 (1988).

4.     Tinevez, J. Y. *et al.* TrackMate: An open and extensible platform for single-particle

tracking. *Methods* **115**, 80–90 (2017).

5.    Kanagal, B. & Sindhwani, V. Rank Selection in Low-rank Matrix Approximations: A

Study of Cross-Validation for NMFs. in *Advances in Neural Information Processing*

*Systems* **23**, (2010).

6.    Owen, A. B. & Perry, P. O. Bi-Cross-Validation of the SVD and the Nonnegative Matrix

Factorization. *Source Ann. Appl. Stat.* **3**, 564–594 (2009).

7.    Wold, S. Cross-Validatory Estimation of the Number of Components in Factor and

Principal Components Models. *Technometrics* **20**, 397–405 (1978).

8.    Ester, M., Kriegel, H.-P., Sander, J. & Xu, X. *A Density-Based Algorithm for Discovering*

*Clusters in Large Spatial Databases with Noise*. (1996).

9.    Ronan, T., Qi, Z. & Naegle, K. M. Avoiding common pitfalls when clustering biological

data. *Science Signaling* **9**, re6–re6 (2016).

10.    Kawamoto, T. & Kabashima, Y. Cross-validation estimate of the number of clusters in a

network. *Sci. Rep.* **7**, 1–17 (2017).

11.    Chang, A. Y. & Marshall, W. F. Dynamics of living cells in a cytomorphological state

space. *Proc. Natl. Acad. Sci. U. S. A.* **116**, 21556–21562 (2019).

12.    Wan, K. Y. & Goldstein, R. E. Time Irreversibility and Criticality in the Motility of a

Flagellate Microorganism. *Phys. Rev. Lett.* **121**, 58103 (2018).

13.    Bak, P., Tang, C. & Wiesenfeld, K. Self-organized criticality: An explanation of the 1/f

noise. *Phys. Rev. Lett.* **59**, 381–384 (1987).

14.    Sánchez, R., Newman, D. E. & Carreras, B. A. Waiting-Time Statistics of Self-Organized-

Criticality Systems. *Phys. Rev. Lett.* **88**, 4 (2002).

15.    Markov, A. A. Extension of the Limit Theorems of Probability Theory to a Sum of

Variables Connected in a Chain. *Notes Imp. Acad. Sci. St. Petersbg. VIII Ser. Physio-*

*Mathematical Coll.* **22**, (1907).

16.    Gagniuc, P. *Markov Chains: From Theory to Implementation and Experimentation*.

(Wiley, 2017).

17.    Billingsley, P. *Statistical Methods in Markov Chains*. *The Annals of Mathematical*

*Statistics* **32**, 12–40 (Institute of Mathematical Statistics, 1961).

18.    Hiscott, R. N. *Chi-Square Tests for Markov Chain Analysis*. *Mathematical Geology* **13**,

(1981).

19.    Cardona, A. *et al.* TrakEM2 Software for Neural Circuit Reconstruction. *PLoS One* **7**,

e38011 (2012).

20.    Thévenaz, P., Ruttimann, U. E. & Unser, M. A pyramid approach to subpixel registration

based on intensity. *IEEE Trans. Image Process.* **7**, 27–41 (1998).

**Acknowledgements**

We would like to thank the students and faculty of the 2016 Marine Biological Laboratory

Physiology Course and current and former members of the Marshall Lab and King Lab for

comments, critiques, and encouragement during the development of this project. We would also

like to thank David Booth and Dennis Bray for comments on the manuscript. This work was

supported in part by the I2CELL Seed Award of the Fourmentin-Guilbert Scientific Foundation

(to WFM), NIH grant R35 GM130327 (to WFM), NSF grant MCB- 2012647 (to WFM),

and the Merck Fellowship of the Jane Coffin Childs Memorial Fund for Medical Research (to

BTL).

**Author contributions**

BTL, JDP, and WFM conceived of the study and developed analyses. BTL collected data. BTL,

JG, JDP, and WFM interpreted the data. BTL and JG performed analyses. BTL and WFM wrote

the paper.

**Competing interests**

The authors declare no competing interests.

**Supplementary Information** is available for this paper including Figures S1-S9 and Movies S1

and S2.