1   **Allelic and Genotypic Frequencies of NAT2, CYP2E1 and AADAC genes in a cohort of Peruvian**

2   **Tuberculosis Patients**

Kelly S. Levano[1], Luis Jaramillo-Valverde[1], David D. Tarazona[1], Cesar Sanchez[1], Silvia Capristano[1], Lely

Solari[1], Alberto Mendoza-Ticona[2], Alonso Soto[3,4], Christian Rojas[1], Roberto Zegarra-Chapoñan[1] and

Heinner Guio[1,5,6*]

[1]Instituto Nacional de Salud, Lima- Perú

[2]Estrategia Sanitaria Nacional de Prevención y Control de Tuberculosis en el Perú, MINSA – Perú

[3]Instituto de investigación en Ciencias Biomédicas (INICIB). Facultad de Medicina. Universidad Ricardo

Palma. Lima - Perú

[4]Departamento de Medicina. Hospital Nacional Hipólito Unanue. Lima, Perú

[5]Universidad de Huánuco, Huánuco, Perú

[6]Universidad Científica del Sur, Lima, Perú

Corresponding author*:

heinnerguio@gmail.com

Abstract

Background: We determined the frequency of genetic polymorphisms in three anti-TB drug metabolic

proteins previously reported: N-acetyltransferase 2 (NAT2), cytochrome P450 2E1 (CYP2E1) and

arylacetamide deacetylase (AADAC) within a Peruvian population in a cohort of TB patients. We included

395 participants completed their anti-tuberculosis treatment.

Results: ~74% of the participants are carriers of slow metabolizer genotypes: NAT2*5, NAT2*6 and

NAT2*7, which increase the sensitivity of INH at low doses and increase the risk of drug-induced liver

injuries. ~ 64% are homozygous for the wild-type CYP2E1*1A allele, which could increase the risk of

hepatotoxicity. However, 16% had a NAT2 fast metabolizer phenotype which could increase the risk of

acquiring resistance to INH, thereby increasing the risk of multidrug-resistant (MDR) or treatment

failure. The frequency of rs1803155 (AADAC*2 allele) was higher (99.9%) in Peruvians than in in

European American, African American, Japanese, and Korean populations.

Conclusions: This high prevalence of slow metabolizers for Isoniazid in the Peruvian population should

be further studied and considered to help individualize drug regimens, especially in countries with a

great genetic diversity like Peru. These data will help the Peruvian National Tuberculosis Control

Program develop new strategies for therapies.

Keywords: NAT2, CYP2E1, AADAC, Tuberculosis

Background:

Tuberculosis (TB) continues to be a leading cause of global morbidity and mortality, with about 10

million cases and a total of 1.2 million deaths reported in 2019 [1]. Even though the current TB regimen

is highly effective under optimal conditions, there are still many undefined issues including drug

underexposure, high prevalence if drug-related toxicity, selection of resistant strains and variability of

response [2], which could be explained by the variability in the pharmacokinetics of anti-TB drugs.

Mutations or polymorphisms in genes encoding metabolic enzymes, transporters or carries can lead to

this variability in drug pharmacokinetics and pharmacodynamics. The identification of these genetic

variations could help select the right anti-TB drug, with the right dosage increasing efficacy and reducing

drug-related toxicity and preventing drug resistance [2, 3]. To determine if genetic variabilities affecting

drug response were present in the Peruvian population, which has a high TB burden with an estimated

32,970cases in 2019 [1] , including a high prevalence of drug-resistant TB cases, we determined the

frequency of genetic polymorphisms in three anti-TB drug metabolic proteins previously reported: N-

acetyltransferase 2 (NAT2), cytochrome P450 2E1 (CYP2E1) and arylacetamide deacetylase (AADAC) [4].

These three proteins participate in the metabolism of the initial phase anti-TB drugs: isoniazid and

rifampicin. In the liver, isoniazid is acetylated to its major metabolite, N-acetyl-isoniazid by the action of

NAT2. It is then further deactivated by other enzymes including CYP2E1 [5, 6]. Thus, genetic variations in

these two enzymes, leading to alterations in their enzymatic functions can cause variations in isoniazid

pharmacokinetics. AADAC is one of the few known enzymes responsible for the deacetylation of

rifampicin and AADAC allele decreased enzyme activity [7, 8]. Thus, genetic variations in these three

enzymes, leading to alterations in their enzymatic functions could cause variations in isoniazid and

rifampicin pharmacokinetics. For the present study, we selected previously reported single nucleotide

polymorphisms (SNPs) that could alter NAT2, CYP2E1 and AADAC enzyme activity and determine their

frequency within a Peruvian population in a cohort of TB patients.

Methods:

Studied populations:

Our study includes 395 unrelated individuals diagnosed with pulmonary tuberculosis between 2014-

2015 recruited from health establishments of the Minister of Health (MINSA) located in Lima and Callao,

Peru. The 395 participants (217 males and 178 females) completed their anti-tuberculosis treatment.

Our study was approved by the Ethics in Research Committee of the Peruvian National Institute of

Health (INS), and written informed consent was obtained from all the participants.

Genotyping of NAT2, CYP2E1 and AADAC:

Genomic DNA was extracted from peripheral blood of all 395 participants using the genomic DNA

extraction kit QIAamp DNA Blood Mini Kit (Qiagen, Germany). The selected genomic DNA regions for the

analysis of each gene included the most common reported SNPs (For NAT2: rs1041983, rs1801280,

rs1799929, rs1799930, rs1208 and rs1799931; for CYP2E1: rs3813867 and rs2031920; for AADAC:

rs1803155). These regions were amplified by PCR using Platinum Taq DNA polymerase kit (Invitrogen,

USA) using the following primers: For NAT2: 5'-GTCACACGAGGAAATCAAATGCT-3 and 5'-

CGTGAGGGTAGAGAGGATATCTG-3'; for CYP2E1: 5'- CCGTGAGCCAGTCGAGTCTA-3' and 5'-

TTCATTCTGTCTTCTAACTGGCAA-3'; and for AADAC: 5'- TCATTCCTAGCAGAAAGGAGATT-3' and 5'-

GCTCACATTTATTCTCTTGCATCG-3'. PCR-amplified fragments were purified using QIAmp Gel Purification

Kit (Qiagen, USA). SNP genotyping on the purified fragments was performed using sanger sequencing

(Macrogen, South Korea). Nucleotide substitutions were identified and analyzed using the Geneious

version 9.1.5 (Biomatters Ltd., New Zealand).

computational phenotyping for NAT2:

Predicted phenotypes were determined from genotypes as three types of metabolizers: slow

metabolizer (two slow alleles), rapid metabolizer (two rapid alleles) and intermediate metabolizer (one

slow and another rapid acetylator allele). The alleles considered rapid were: wild-type NAT2∗4, 282C>T

(NAT2∗13), 481C>T (NAT2∗11) and 803A>G (NAT2∗12), while the alleles considered slow were: 341T>C

(NAT2∗5), 590G>A (NAT2∗6), 857G>A (NAT2∗7), and 191G>A (NAT2∗14) [9]. The computational inferred

phenotypes using a combination of NAT2 SNPs for the 395 participants were determined using an online

software program, NAT2PRED (nat2pred.rit.albany.edu) [10, 11].

Statistical analysis:

Genotype frequencies in our population were calculated in accordance with the Hardy-Weinberg

equilibrium. Intervals confidence 95% were calculated for phenotypic genotypic and allelic frequencies.

Data analysis was carried out using Stata 15 program (StataCorp. 2016. Stata Statistical Software:

Release 15. College Station, TX, USA).

Results:

In this study, we determined the presence of the six most common SNPs, rs1041983 (282C>T),

rs1801280 (341T>C), rs1799929 (481C>T), rs1799930 (590G>A), rs1208 (803A>G) and rs1799931

(857G>A), of the NAT2 gene in the 395 individuals from Lima and Callao, Peru. No new SNPs were

identified, indicating that the NAT2 gene has no other SNPs in the Peruvian population studied. The

allele frequencies of these major NAT2 SNPs are represented in Table 1. This study found that NAT2∗13,

C283T, is the most frequent genetic variant (~40% of alleles) among our samples. The allele not

harboring any mutation (wild-type NAT2∗4) was present in 44 of the 395 samples (~11% of alleles).

Table 2 shows the frequencies of NAT2 genotype obtained from the studied population. The most

frequent observed heterozygote was also NAT2∗13 C283T (48.6%) followed by NAT2∗11 C481T (34.9%)

and NAT2∗5 T341C (34.7%) among the Peruvian population studied. The lowest frequency of observed

heterozygote genotypes was NAT2∗12 A803G with a frequency of 18.7%. In homozygote, the NAT2∗12

803A>G (31.2%) genotype was the most common one but the lowest homozygote among them was

NAT2∗6 590G>A (12.5%). The linkage disequilibrium (LD) analysis is shown in Figure 1. The six NAT2

variants, 282C>T, 341T>C, 481C>T, 590G>A, 803A>G, and 857G>A were applied to Haploview software.

The LD for each pair of genetic variants was measured using |D☐ | and correlation coefficient (r2 > 0:8). A

haplotype block was found in the following SNP positions 341T>C and 481C>T (D′ :0,808 and r2:0,437) in

Peruvian population samples which is identified as like strong LD. There were no significant differences

observed between NAT2 genotypes with respect to age and gender. The NAT2 inferred metabolizing

status was predicted using the 6 SNPs analyzed as stated above. As a result, the predicted metabolizing

phenotype of fast, intermediate, and slow metabolizers was 14.9%, 38.2% and 46.8%, respectively

(Table 3).

The allelic and genotypic distribution of the rs2031920 (-1053C>T) variant of CYP2E1 among the studied

population is shown in Tables 1 and 2, respectively. The results show an allele frequency of ~21% for the

CYP2E1 variant, and ~79% for the wild-type allele. The observed genotype frequency of the homozygous

and heterozygotes were ~7% and ~36%, respectively.

We also analyzed the allele and genotype frequency of the rs1803155 SNP of the AADAC gene in the 395

individuals of this study (Table 1 and 2). According to our results, the AADAC genetic variant has an allele

frequency of ~99.9%, while the wild-type allele was ~20%. The homozygous and heterozygous genotype

distribution were ~85% and ~10%, respectively.

Discussion:

Studies in different populations have shown ethnic variabilities in both NAT2 and CYP2E1 genotypes and

phenotypes. There is still limited information about the genetic variations in the Peruvian population. In

the current study, we analyzed NAT2, CYP2E1 and AADAC genotypes and allele frequencies in 395

individuals from Peru. As stated above, NAT2 and CYP2E1 are two essential enzymes in the metabolism

of INH. Altered NAT2 and/or CYP2E1 activities due to polymorphic genotypes can result in (a) the

accumulation of toxic substances in the liver, and (b) variations in INH plasma concentrations that can

affect the efficacy of the drug.

In the analysis of the *NAT2* gene, the results showed that *NAT2*13* (39.7% of alleles) and *NAT2*7* (35.6%

of alleles) were the most frequent genetic variants amount the population studied. *NAT2*13* is a silent

mutation, Y94Y, that does not alter the metabolizer phenotype, whereas *NAT2*7* results in an amino

acid substitution, G286E, that leads to a significant decrease in the enzyme's activity [12, 13]. The

distribution of the NAT2 polymorphisms in the population studied were similar to other American

populations in that one of the most frequent alleles was *NAT2*5*. It is established that the frequency of

*NAT2*5* in European populations is ~50%, in African populations is ~33-42% and in Asian populations is

~5% [14–17]. According to our results, the allele frequency of *NAT2*5* is ~25%. The other two slow

metabolizer alleles are *NAT2*6* and *NAT2*7*. The *NAT2*6* is common in all populations mentioned above

with a frequency of ~30%. Conversely, the frequency of NAT2*7 is low in European populations (~2%)

and African populations (~3-6%). In Asian populations, the frequency of NAT2*7 is ~10-12% [17].

Diverging from these reports, in our studied population the allele frequency of NAT2*6 is ~14% and of

NAT2*7 is ~36%. As stated above, reduced NAT2 activity, which is observed in NAT2*7 variants, can lead

to adverse drug reactions due to increased accumulation of toxic metabolites. Additionally, our study

revealed that the genotype frequency (predicted phenotype) of slow metabolizers is ~47%. The

relationship of NAT2 polymorphisms with INH-induced hepatotoxicity in TB patients among different

populations were studied [14–21], but the previously published studies have demonstrated inconsistent

results. Therefore, analysis of the slow genotypes should become part of the dosage regimen of INH in

TB patients undergoing anti-TB treatment to prevent drug-induced liver injuries [18–21].

After NAT2 acetylates INH converting it to acetyl-INH, it can enter the CYP2E1 pathway, which couples

with the glutathione-S-transferase (GST) metabolic pathway to facilitate the elimination of toxic

metabolites [22–24]. Studies have shown that individuals with the CYP2E1 wild-type allele (c1/c1

genotype) have a higher CYP2E1 activity that those with CYP2E1*5B allele (c1/c2 or c2/c2 genotype).

Thus, these individuals can generate more hepatotoxins and therefore increase the risk of drug-induced

liver injuries [23, 25, 26]. In our studied population, the allele frequency of the CYP2E1*5B is ~ 79%,

which increases the risk of hepatotoxicity specially in patients with a slow metabolizer phenotype for

NAT2 [23, 27].

An important enzyme in the metabolism of RIF is AADAC catalyzing its deacetylation to 25-deacetyl-RIF

[7, 28, 29]. Polymorphic variations affecting this enzyme's activity can also result in the accumulation of

toxic substances and variations in RIF plasma concentrations that can affect the efficacy of this drug. In

this study, we analyzed the nonsynonymous SNP rs1803155 (AADAC*2 allele), which leads to a change

in amino acid (V281I) in the coding region [30]. An allele frequency of ~60% for AADAC*2 has been

reported in European American, African American, Japanese, and Korean populations. In our studied

population, the allele frequency of *AADAC*2* is ~99.9%. A limitation in this study is the number of SNPs

analyzed in each gene, especially in AADAC. The analysis of additional genetic variations in AADAC can

provide additional information in the metabolism of RIF. For example, the allele AADAC*3

(g.13651G>A/g.14008T>C), not analyzed in the current study, has shown a reduced metabolizing activity

for RIF [30]. Additionally, studies have reported genetic polymorphisms in other RIF metabolizing

enzymes, including carboxylesterase 1(CES1) and carboxylesterase 2 (CES2) [31], as well as in drug

transporters and /or their transcriptional regulators, including SLCO1B1[32] and ABCB1 [33].

Countries have begun clinical trials focused on personalization of tuberculosis treatment to reduce the

consequences for patients in treatment [21, 34]. In countries like Peru, where high rates of tuberculosis

are recorded and therefore more people in treatment, the pharmacogenomic of individuals becomes a

crucial tool for an optimum tuberculosis treatment. This review highlights the importance of having

pharmacogenomic studies and having the identification of polymorphisms associated to the metabolism

of the anti-tuberculosis drugs in our Peruvian population.

Conclusion:

In conclusion, our study showed the distribution of NAT2, CYP2E1 and AADAC genetic polymorphisms in

a Peruvian population diagnosed with tuberculosis. This is a preliminary study to help understand the

genetic basis of metabolizing polymorphisms in our population, and thus contribute to the use of this

and future data in determining the safe INH and RIF dose in slow and fast metabolizers and thus

minimizing adverse drug reactions. According to our results, ~74% of the participants are carriers of slow

metabolizer genotypes: *NAT2*5*, *NAT2*6* and *NAT2*7*, which increase the sensitivity of INH at low doses

and increase the risk of drug-induced liver injuries. Additionally, ~ 64% are homozygous for the wild-

type CYP2E1*1A allele, which could increase the risk of hepatotoxicity. This high prevalence of slow

metabolizers for Isoniazid in the Peruvian population should be further studied and considered to help

individualize drug regimens, especially in countries with a great genetic diversity like Peru. However,

16% had a NAT2 fast metabolizer phenotype which could increase the risk of acquiring resistance to INH,

thereby increasing the risk of multidrug-resistant (MDR) or treatment failure. The frequency of

rs1803155 (AADAC*2 allele) was higher (99.9%) in Peruvians than in in European American, African

American, Japanese, and Korean populations. These data will help the Peruvian National Tuberculosis

Control Program develop new strategies for therapies.

References:

1. WHO | Global tuberculosis report 2019. WHO. 2020.

2. Motta I, Calcagno A, Bonora S. Pharmacokinetics and pharmacogenetics of anti-tubercular drugs: a

tool for treatment optimization? Expert Opinion on Drug Metabolism and Toxicology. 2018;14:59–82.

doi:10.1080/17425255.2018.1416093.

3. Figueiredo Teixeira RL de, Pires Lopes MQ, Noel P, Rezende A. Tuberculosis Pharmacogenetics: State

of The Art. In: Tuberculosis - Current Issues in Diagnosis and Management. InTech; 2013.

4. Guio H, Levano KS, Sánchez C, Tarazona D. The role of pharmacogenomics in the tuberculosis

treatment regime. Revista Peruana de Medicina Experimental y Salud Publica. 2015.

5. Sotsuka T, Sasaki Y, Hirai S, Yamagishi F, Ueno K. Association of isoniazid-metabolizing enzyme

genotypes and isoniazid-induced hepatotoxicity in tuberculosis patients. In Vivo. 2011;25:803–12.

6. Bao Y, Ma X, Rasmussen TP, Zhong X bo. Genetic Variations Associated with Anti-Tuberculosis Drug-

Induced Liver Injury. Current Pharmacology Reports. 2018;4:171–81.

7. Nakajima A, Fukami T, Kobayashi Y, Watanabe A, Nakajima M, Yokoi T. Human arylacetamide

deacetylase is responsible for deacetylation of rifamycins: Rifampicin, rifabutin, and rifapentine.

Biochemical Pharmacology. 2011;82:1747–56. doi:10.1016/j.bcp.2011.08.003.

8. Lee M-R, Huang H-L, Lin S-W, Cheng M-H, Lin Y-T, Chang S-Y, et al. Isoniazid Concentration and NAT2

Genotype Predict Risk of Systemic Drug Reactions during 3HP for LTBI. Journal of Clinical Medicine.

2019;8:812. doi:10.3390/jcm8060812.

9. Hein DW, Doll MA, Fretland AJ, Leff MA, Webb SJ, Xiao GH, et al. Molecular genetics and epidemiology

of the NAT1 and NAT2 acetylation polymorphisms. Cancer epidemiology, biomarkers & prevention : a

publication of the American Association for Cancer Research, cosponsored by the American Society of

Preventive Oncology. 2000;9:29–42. http://www.ncbi.nlm.nih.gov/pubmed/10667461. Accessed 29 May

2019.

10. Sabbagh A, Darlu P, Vidaud M. Evaluating NAT2PRED for inferring the individual acetylation status

from unphased genotype data. BMC Medical Genetics. 2009;10.

11. Kuznetsov IB, Mcduffie M, Moslehi R. A web server for inferring the human N-acetyltransferase-2

(NAT2) enzymatic phenotype from NAT2 genotype. BIOINFORMATICS APPLICATIONS NOTE.

2009;25:1185–6. doi:10.1093/bioinformatics/btp121.

12. Vatsis KP, Martell KJ, Weber WW. Diverse point mutations in the human gene for polymorphic N-

acetyltransferase. Proceedings of the National Academy of Sciences of the United States of America.

1991;88:6333–7. doi:10.1073/pnas.88.14.6333.

13. Lakkakula S, Pathapati RM, Chaubey G, Munirajan AK, Lakkakula BVKS, Maram R. NAT2 genetic

variations among South Indian populations. Human Genome Variation. 2014;1:1.

doi:10.1038/hgv.2014.14.

14. Cascorbi I, Drakoulis N, Brockmoller J, Maurer A, Sperling K, Roots I. Arylamine N-acetyltransferase

(NAT2) mutations and their allelic linkage in unrelated Caucasian individuals: Correlation with

phenotypic activity. American Journal of Human Genetics. 1995;57:581–92.

/pmc/articles/PMC1801274/?report=abstract. Accessed 22 Dec 2020.

15. Sekine A, Saito S, Iida A, Mitsunobu Y, Higuchi S, Harigae S, et al. Identification of single-nucleotide

polymorphisms (SNPs) of human N-acetyltransferase genes NAT1, NAT2, AANAT, ARD1, and L1CAM in

the Japanese population. Journal of Human Genetics. 2001;46:314–9. doi:10.1007/s100380170065.

16. Borlak J, Reamon-Buettner SM. N-acetyltransferase 2 (NAT2) gene polymorphisms in colon and lung

cancer patients. BMC Medical Genetics. 2006;7:58. doi:10.1186/1471-2350-7-58.

17. Tiis RP, Osipova LP, Lichman DV, Voronina EN, Filipenko ML. Studying polymorphic variants of the

NAT2 gene (NAT2*5 and NAT2*7) in Nenets populations of Northern Siberia. BMC Genetics. 2020;21

Suppl 1. doi:10.1186/s12863-020-00909-4.

18. Zahra MA, Kandeel M, Aldossary SA, Al-Taher A. Study on Genotyping Polymorphism and Sequencing

of N-Acetyltransferase 2 (NAT2) among Al-Ahsa Population. BioMed Research International. 2020;2020.

19. Azuma J, Ohno M, Kubota R, Yokota S, Nagai T, Tsuyuguchi K, et al. NAT2 genotype guided regimen

reduces isoniazid-induced liver injury and early treatment failure in the 6-month four-drug standard

treatment of tuberculosis: A randomized controlled trial for pharmacogenetics-based therapy. European

Journal of Clinical Pharmacology. 2013;69:1091–101. doi:10.1007/s00228-012-1429-9.

20. Ganachari M, Ruiz-Morales JA, Gomez de la Torre Pretell JC, Dinh J, Granados J, Flores-Villanueva PO.

Joint Effect of MCP-1 Genotype GG and MMP-1 Genotype 2G/2G Increases the Likelihood of Developing

Pulmonary Tuberculosis in BCG-Vaccinated Individuals. PLoS ONE. 2010;5:e8881.

doi:10.1371/journal.pone.0008881.

21. Huerta-García AP, Medellín-Garibay SE, Ortiz-Álvarez A, Magaña-Aquino M, Rodríguez-Pinal CJ,

Portales-Pérez DP, et al. Population pharmacokinetics of isoniazid and dose recommendations in

Mexican patients with tuberculosis. International Journal of Clinical Pharmacy. 2020;42:1217–26.

doi:10.1007/s11096-020-01086-1.

22. Teixeira RLDF, Morato RG, Cabello PH, Muniz LMK, Moreira A da SR, Kritski AL, et al. Genetic

polymorphisms of NAT2, CYP2E1 and GST enzymes and the occurrence of antituberculosis drug-induced

hepatitis in Brazilian TB patients. Memorias do Instituto Oswaldo Cruz. 2011;106:716–24.

doi:10.1590/S0074-02762011000600011.

23. Singla N, Gupta D, Birbian N, Singh J. Association of NAT2, GST and CYP2E1 polymorphisms and anti-

tuberculosis drug-induced hepatotoxicity. Tuberculosis. 2014;94:293–8.

24. Guio H, Levano KS, Sanchez C, Tarazona D. [The role of pharmacogenomics in the tuberculosis

treatment regime]. Rev Peru Med Exp Salud Publica. 2015;32:794–800.

25. Vuilleumier N, Rossier MF, Chiappe A, Degoumois F, Dayer P, Mermillod B, et al. CYP2E1 genotype

and isoniazid-induced hepatotoxicity in patients treated for latent tuberculosis. European Journal of

Clinical Pharmacology. 2006;62:423–9. doi:10.1007/s00228-006-0111-5.

26. Huang YS, Chern H Der, Su WJ, Wu JC, Chang SC, Chiang CH, et al. Cytochrome P450 2E1 genotype

and the susceptibility to antituberculosis drug-induced hepatitis. Hepatology. 2003;37:924–30.

27. Guaoua S, Ratbi I, Laarabi FZ, Elalaoui SC, Jaouad IC, Barkat A, et al. Distribution of allelic and

genotypic frequencies of NAT2 and CYP2E1 variants in Moroccan population. BMC Genetics. 2014;15.

28. Thomas L, Miraj SS, Surulivelrajan M, Varma M, Sanju CSV, Rao M. Influence of single nucleotide

polymorphisms on rifampin pharmacokinetics in tuberculosis patients. Antibiotics. 2020;9:1–15.

doi:10.3390/antibiotics9060307.

29. Png E, Alisjahbana B, Sahiratmadja E, Marzuki S, Nelwan R, Balabanova Y, et al. A genome wide

association study of pulmonary tuberculosis susceptibility in Indonesians. BMC medical genetics.

2012;13:5. doi:10.1186/1471-2350-13-5.

30. Shimizu M, Fukami T, Kobayashi Y, Takamiya M, Aoki Y, Nakajima M, et al. A novel polymorphic allele

of human arylacetamide deacetylase leads to decreased enzyme activity. Drug Metabolism and

Disposition. 2012;40:1183–90. doi:10.1124/dmd.112.044883.

31. Sloan DJ, McCallum AD, Schipani A, Egan D, Mwandumba HC, Ward SA, et al. Genetic determinants

of the pharmacokinetic variability of rifampin in Malawian adults with pulmonary tuberculosis.

Antimicrobial Agents and Chemotherapy. 2017;61. doi:10.1128/AAC.00210-17.

32. Yang S, Hwang SJ, Park JY, Chung EK, Lee JI. Association of genetic polymorphisms of CYP2E1, NAT2,

GST and SLCO1B1 with the risk of anti-tuberculosis drug-induced liver injury: A systematic review and

meta-analysis. BMJ Open. 2019;9. doi:10.1136/bmjopen-2018-027940.

33. Pontual Y, Pacheco VSS, Monteiro SP, Quintana MSB, Costa MJM, Rolla VC, et al. ABCB1 gene

polymorphism associated with clinical factors can predict drug-resistant tuberculosis. Clinical Science.

2017;131:1831–40. doi:10.1042/CS20170277.

34. Yoo H, Chun Ji S, Cho J-Y, Kim S-H, Yoon JG, Goo Lee M, et al. A pilot study to investigate the utility of

NAT2 genotype-guided isoniazid monotherapy regimens in NAT2 slow acetylators. Pharmacogenetics

and Genomics. 2020.

Table 1: Allele frequencies of NAT2, CYP2E1 and AADAC polymorphisms in a Peruvian population (n =

395)

Gene	Allele	SNP	position	Substituted amino acid	Allele frequency [288] (95% CI)
NAT2	NAT2*4			Wild-type	0.111 (0.089-0.134)
	NAT2*13	rs1041983	c.283C>T	Y94Y	0.397 (0.363-0.432)
	NAT2*5	rs1801280	c.341T>C	I114T	0.247 (0.216-0.278)
	NAT2*11	rs1799929	c.481C>T	L161L	0.329 (0.296-0.363)
	NAT2*6	rs1799930	c.590G>A	R197Q	0.138 (0.113-0.163)
	NAT2*12	rs1208	c.803A>G	R268K	0.228 (0.198-0.258)
	NAT2*7	rs1799931	c.857G>A	G286E	0.356 (0.322-0.390)
CYP2E1	CYP2E1*1A			Wild-type	0.79359 (0.765-0.823)
	CYP2E1*5B	rs2031920	c.-1053C>T		0.206 (0.177-0.235)
AADAC	AADAC*1			Wild-type	0.199 (0.167-0.230)
	AADAC*2	rs1803155	c.841G>A	V281I	0.999 (0.967-1.03)

290   **Table 2:** Genotype frequency of NAT2, CYP2E1 and AADAC genes in a Peruvian population (n = 395)

Gene	Allele	Wild Type Frequency (proportion, 95% CI)	Heterozygote Frequency (proportion, 95% CI)	Homozygote Frequency (proportion, 95% CI)
NAT2	NAT2*13 (C282T)	C/C: 142 (0.359, 0.312-0.407)	C/T: 192 (0.486, 0.437-0.535)	T/T: 61 (0.154, 0.119-0.190)
	NAT2*5 (T341C)	T/T: 229 (0.580, 0.531-0.628)	T/C: 137 (0.347, 0.300-0.394)	C/C: 29 (0.073, 0.048-0.099)
	NAT2*11 (C481T)	C/C: 196 (0.496, 0.447-0.546)	C/T: 138 (0.349, 0.302-0.396)	T/T: 61 (0.154, 0.119-0.190)
	NAT2*6 (G590A)	G/G: 303 (0.767, 0.725-0.809)	G/A: 75 (0.190, 0.151-0.229)	A/A: 17 (0.043, 0.023-0.063)
	NAT2*12 (A803G)	A/A: 268 (0.678, 0.632-0.725)	A/G: 74 (0.187, 0.149-0.226)	G/G: 53 (0.134, 0.101-0.168)
	NAT2*7 (G857A)	G/G: 210 (0.532, 0.482-0.581)	G/A: 89 (0.225, 0.184-0.267)	A/A: 96 (0.243, 0.201-0.285)
CYP2E1	CYP2E1*5B(C-1053T)	C/C: 250 (0.641, 0.594-0.688)	C/T: 119 (0.359, 0.312-0.406)	T/T: 21 (0.076, 0.032-0.076)
AADAC	AADAC*2 (G841A)	G/G: 19 (0.048, 0.027-0.069)	G/A: 41 (0.104, 0.074-0.134)	A/A: 335 (0.848, 0.813-0.883)

292   **Table 3:** Predicted metabolizing phenotype for NAT2 in a Peruvian population (n = 395)

Gene	Metabolizing Phenotype * Proportion (95% CI)		
NAT2	Fast	Intermediate	Slow
	0.149 (0.114-0.185)	0.382 (0.334-0.430)	0.468 (0.419-0.518)

293   *The   metabolizing   phenotype   was   determined   using   the   online   software

294   http://nat2pred.rit.albany.edu/.

296   **Table 4:** Distribution of NAT2 alleles among the Peruvian population studied compared with various

297   human population.

Population	Peru (current study)	Brazil (2016)	Mexico (2012)	Spain (2011)
NAT2*4 (Wild-type)	0.111	0.258	0.306	0.186
NAT2*13 (C282T)	0.099	0.008	0.008	-
NAT2*5 (T341C)	0.420	0.446	0.312	0.417
NAT2*11 (C481T)	0.035	-	-	-
NAT2*6 (G590A)	0.099	0.150	0.174	0.292
NAT2*12 (A803G)	0.013	0.023	0.048	-
NAT2*7 (G857A)	0.223	0.096	0.140	0.106

301 **Figure 1:** Linkage disequilibrium for NAT2 metabolizer-associated SNPs in Peruvian population studied.