Small Directional Treadmill Perturbations Induce Differential Gait Stability Adaptation

Running title: Gait stability adaptation to small directional perturbations

Jinfeng Li [1],* and Helen J. Huang [1,2]

[1] Department of Mechanical and Aerospace Engineering, University of Central Florida, Orlando, FL 32816, USA;

[2] Disability, Aging, and Technology (DAT) Cluster, University of Central Florida,

Orlando, FL 32816, USA; hjhuang@ucf.edu

* Correspondence: jinfeng@knights.ucf.edu; Tel.: +1-321-666-0567;

12760 Pegasus Dr., Bldg 40, Room 307, Orlando, FL 32816, USA

Abstract: Introducing unexpected perturbations to challenge gait stability is an effective approach to investigate balance control strategies. Little is known about the extent to which people can respond to small perturbations during walking. This study aimed to determine how subjects adapted gait stability to multidirectional perturbations with small magnitudes applied on a stride-by-stride basis. Ten healthy young subjects walked on a treadmill that either briefly decelerated belt speed ("stick"), accelerated belt speed ("slip"), or shifted the platform medial-laterally at right leg mid-stance. We quantified gait stability adaptation in both anterior-posterior and medial-lateral directions using margin of stability and its components, base of support and extrapolated center of mass. Gait stability was disrupted upon initially experiencing the small perturbations as margin of stability decreased in the stick, slip, and medial shift perturbations and increased in the lateral shift perturbation. Gait stability metrics were generally disrupted more for perturbations in the coincident direction. Subjects employed both feedback and feedforward strategies in response to the small perturbations, but mostly use feedback strategies during adaptation. Subjects primarily used base of support (foot placement) control in the lateral shift perturbation and extrapolated center of mass control in the slip and medial shift perturbations. These findings provide new knowledge about the extent of gait stability adaptation to small magnitude perturbations applied on a stride-by-stride basis and reveal potential new approaches for balance training interventions to target foot placement and center of mass control.

NEW & NOTEWORTHY Little is known about if and how humans can adapt to small magnitude perturbations experienced on a stride-by-stride basis during walking. Here, we show that even small perturbations disrupted gait stability and that subjects could still adapt their reactive balance control. Depending on the perturbation direction, subjects might prefer adjusting their foot placement over their center of mass and vice versa. These findings could help potentially tune balance training to target specific aspects of balance.

Keywords: gait; adaptation; perturbation; margin of stability; balance

1. Introduction

Perturbations during walking can induce potential losses of balance and elicit corrective locomotor adaptations, which are useful for understanding human balance control (1, 2). There are multiple approaches to perturb gait stability, including split-belt walking (1, 3), visual flow distortions (4, 5), waist pulling (6, 7), platform displacements (8, 9), and rapid changes in treadmill belt speed (10, 11). Studies using discrete perturbations such as waist pulling, platform displacements, and changes in treadmill belt speeds often apply the perturbation once out of every 5-20 strides (6–11). These large magnitude perturbations are less frequent and often require recovery steps to regain balance. Currently, there is no clear optimal frequency and magnitude for perturbation-based balance training during walking (12, 13), but more frequent exposure to perturbations seems to be beneficial for developing long-term fall-resisting skills (14, 15). Small magnitude perturbations are also more easily navigated by older adults (16, 17). As such, applying frequent repeated perturbations with small magnitudes during walking to create small but frequent losses of balance could be an effective approach for balance training during walking. Little is known, however, about the extent to which humans can adapt to small magnitude discrete balance perturbations applied on a stride-by-stride basis during walking. This approach could reveal stride-to-stride adaptation to almost imperceptible perturbations to gait stability, which would provide additional insight about how humans control, maintain, and adapt their stability during walking.

The margin of stability is a well-accepted metric to quantify stability of human walking (18). The margin of stability is the distance between the edge of the base of support and the extrapolated center of mass (19). People tend to maintain a constant margin of stability when walking on different surfaces (20, 21). When external conditions in the environment disrupt the steady state of walking and its margin of stability, increasing the base of support is effective for maintaining stability based on the inverted pendulum model (22) and has been demonstrated in multiple gait studies (6, 23, 24). For steps before and after unexpected treadmill belt acceleration perturbations during walking, one study showed that margin of stability decreased, and subjects took fewer recovery steps, suggesting improved or preserved stability (25). Treadmill belt speed induced backward loss-of-balance perturbations can lead to larger margin of stability reductions in older adults compared to young adults (26). Interestingly, these margin of stability responses scaled with the perturbation intensity in both the young and older

adults (26). The extent to which humans can control their margin of stability in response to small magnitude

perturbations applied on a stride-by-stride basis remains unknown.

Analyzing the base of support and the extrapolated center of mass components of margin of stability could

provide insights on how the body reacts to perturbations and maintains stability. Controlling foot placement

through the swing leg or controlling center of mass through the stance leg are two main balance strategies for

perturbed walking (6, 27, 28). In young adults, adaptation to multi-directional waist pulling perturbations have

resulted in increased medial-lateral foot placement at perturbation onset, potentially to proactively regulate

balance and also increased anterior-posterior foot placement and margin of stability at contralateral heel strike,

perhaps to reactively regain stability (29). Older adults have demonstrated that they tend to adjust their

extrapolated center of mass, potentially to transfer the improved stability of the trained leg to the untrained leg in

response to treadmill belt acceleration perturbations (30). Treadmill belt slip perturbations have also resulted in

the lengthening of the base of support and shortening the forward excursion of the extrapolated center of mass in

post-stroke individuals, effectively increasing their anterior-posterior margin of stability and overall stability (31).

Thus, separating margin of stability into its components, the base of support and extrapolated center of mass, will

provide greater insight on how humans adapt, if at all, to frequent small magnitude perturbations.

The primary purpose of this study was to determine how small treadmill perturbations applied on a

stride-by-stride basis in healthy young adults affected margin of stability and its components (base of support and

extrapolated center of mass). We rapidly accelerated or decelerated treadmill belt speed to create "slip" or "stick"

perturbations, respectively. We also rapidly shifted (i.e., translated) the treadmill medially or laterally to create

treadmill shift perturbations in the medial or lateral direction, respectively. We applied the treadmill perturbations at

right mid-stance of each stride, except for no-perturbation "catch" strides that occurred randomly in each batch of 5

strides. The purpose of the "catch" strides was to probe whether subjects were potentially adopting a feedforward

strategy in response to the small magnitude perturbations encountered during perturbed walking. Experimental

protocols that apply perturbations trial-by-trial with unexpected catch trials have typically been used in

goal-directed arm reaching studies (32, 33). We are extending that protocol to walking to gain insights on the

adaptability of balance control during walking. Because we applied perturbations on a stride-by-stride basis, the

strengths of the perturbations were limited to avoid the need for recovery steps, which are often necessary during larger treadmill induced trip and slip perturbations that seek to induce a fall.

The first hypothesis was that when subjects first experienced the small perturbations, their gait stability would be disrupted (i.e., become less stable) and then adapt to regain stability as they gained more experience with the perturbations. As such, we expected margin of stability to decrease upon first experiencing the perturbations and then increase over time with more exposure to the perturbations. The second hypothesis was that subjects would adopt more feedforward strategies as they gained more experience with the perturbations. We expected that margin of stability, base of support, and extrapolated center of mass during catch strides would differ from the averaged levels of the unperturbed strides prior to the perturbed period and deviate more as subjects gained more experience with the perturbations.

To assess the extent of the expected adaptation, we also sought to determine whether perturbation direction and size affected the responses and adaptation to the perturbations. We hypothesized that the anterior-posterior margin of stability and its components would be more sensitive to the treadmill belt perturbations, which were in the coincident (anterior-posterior) direction and likewise, medial-lateral margin of stability and its components would be more sensitive to the treadmill shift perturbations, which were in the coincident (medial-lateral) direction. We also hypothesized that adaptation of margin of stability and its components would scale with perturbation size.

2. Methods

2.1. Participants, setup and protocol

Ten healthy young subjects (4 males, 6 females; age: 21.7 ± 2.4 years) participated in this study. The Institutional Review Board of the University of Central Florida approved the experimental protocol (SBE-16-12831). The study met all requirements in accordance with the Declaration of Helsinki, and all subjects provided their written informed consent.

Subjects walked on a dual split-belt instrumented treadmill (M-gait, Motekforce Link, Amsterdam, The Netherlands) operating at 300 Hz as we recorded their lower limb kinematics using the 16-marker Conventional Lower Body Markerset and a passive motion capture system (OptiTrack, NaturalPoint Inc., Corvallis, OR, USA, 13 Prime13W and 9 Prime13 cameras) operating at 240 Hz. The treadmill has independent speed control for each

treadmill belt, and the treadmill is mounted on a moveable platform such that the treadmill can rapidly translate

side-to-side in real-time. These features were used to produce treadmill perturbations.

We programmed the treadmill (D-Flow, version 3.28.0, Motekforce Link, Amsterdam, The Netherlands) to

apply perturbations during mid-stance of the right leg on a stride-by-stride basis. The perturbation onset occurred

when the vertical projection of the estimated center of mass position (the average position of the four pelvis

markers: left anterior superior iliac spine, right anterior superior iliac spine, left posterior superior iliac spine, and

right posterior superior iliac spine) resided between the right toe (2^{nd} metatarsal) and right heel markers in the

anterior-posterior direction and when the left heel marker was at least 5 cm higher than the right toe marker (Fig.

1A). The duration of each perturbation from onset to termination was ~400 ms such that the treadmill could return to

the neutral unperturbed position and baseline belt speed (1.0 m/s) prior to or shortly after left heel strike.

Perturbations were applied in four directions: anterior, posterior, medial, and lateral to examine the effects of

perturbation directions (Fig. 1A). We also used two perturbation sizes for the anterior and posterior directions,

which could be completed in ~400 ms, to examine potential scaling effects with perturbation size (Fig. 1B). The

baseline walking speed was 1.0 m/s, and the belt acceleration was set to 12.5 m/s^2. For anterior "stick"

perturbations, the right belt speed decreased from 1.0 m/s to the target speed (0.6 m/s and 0.8 m/s; Stick0.4 and

Stick0.2, respectively) and then returned to the baseline speed. For posterior "slip" perturbations, the right belt

speed increased to the target speed (1.2 m/s and 1.4 m/s; Slip0.2 and Slip0.4, respectively) and then returned to the

baseline speed. For medial perturbations, the treadmill shifted 1 cm toward the medial side of the right leg (i.e., to

the left) and then returned to the neutral treadmill location (M1). For lateral perturbations, the treadmill shifted 1 cm

toward the lateral side of the right leg (i.e., to the right) and then returned to the neutral treadmill location (L1). The

maximum acceleration for the medial-lateral perturbations was 3.6 m/s^2.

Each subject completed six trials, one trial for each perturbation condition. We randomized the order of the six

trials and did not inform subjects of the perturbation type, timing, direction, and magnitude. Each trial was 8 minutes

long and consisted of 3 blocks. The trial began with 2 minutes of unperturbed walking (pre block), followed by 4

minutes of walking with perturbations (perturbation block) and concluded the trial with another 2 minutes of

unperturbed walking (post block). During the 4-minute perturbation block, no-perturbation "catch" strides occurred

randomly 1 out of every 5 strides to assess whether subjects were anticipating or reacting (Fig. 1C). Subjects wore a safety harness to prevent potential falls during all walking trials.

2.2. Data analysis

We wrote custom MATLAB (version 9.3, R2017b, Mathworks, Natick, MA, USA) scripts to process the kinematic data and calculate the gait stability metrics. We low-pass filtered the three-dimensional coordinates of the markers using a 4th order Butterworth filter with a 12 Hz cut-off frequency, and then extracted the gait events. Heel strike was the instant the vertical position of the heel marker of the foot transitioning from swing to stance was near zero (close to the treadmill) and reached its maximum vertical acceleration (34). Toe off was the instant the toe marker of the foot transitioning from stance to swing reached its maximum vertical acceleration between two consecutive heel strikes (35).

We calculated the margin of stability at every left heel strike (lhs) to assess reactive control and left toe off (lto) to assess anticipatory control as:

$$MOS = BOS - XCOM \tag{1}$$

where MOS is the margin of stability; BOS is the boundary of the base of support; and XCOM is the extrapolated center of mass (19). We calculated the extrapolated center of mass using Equation 2, which also accounts for the movement velocity of the treadmill surface (36, 37).

$$XCOM = COM + (V_{COM} - V_{treadmill})/\sqrt{g/l} \tag{2}$$

COM is the center of mass position, and V_{COM} is the center of mass velocity, which was derived as the first derivative of the center of mass position. $V_{treadmill}$ is the belt velocity for the anterior-posterior perturbations and the platform velocity for the medial-lateral perturbations, respectively. The remaining variables are g, the gravitational constant (9.81 m/s^2) and l, the equivalent pendulum length, which was the distance from the ankle marker (lateral malleolus) of the leading leg to the center of mass at left heel strike and left toe off.

We calculated the anterior-posterior margin of stability (MOS_{ap}) as the anterior-posterior distance between the extrapolated center of mass and the toe marker of the leading leg in the sagittal plane, which defined the anterior-posterior base of support (BOS_{ap}) (Fig. 2A). The anterior-posterior extrapolated center of mass ($XCOM_{ap}$)

was the anterior-posterior distance between the extrapolated center of mass and the toe marker of the trailing leg.

We calculated the medial-lateral margin of stability (MOS_{ml}) as the medial-lateral distance between the extrapolated

center of mass and the ankle marker of the leading leg in the frontal plane, which defined the medial-lateral base of

support (BOS_{ml}) (Fig. 2B). The medial-lateral extrapolated center of mass ($XCOM_{ml}$) was the medial-lateral

distance between the extrapolated center of mass and the ankle marker of the trailing leg.

We quantified margin of stability and its components at different phases throughout the protocol: pre, early

perturbed, late perturbed, early catch, late catch, early post, and late post. Pre referred to all strides during the first

2-minute pre walking block except for the first 11 strides during which the treadmill belts were getting up to speed

(38). Early perturbed was the first 3 perturbed strides during the 4-minute perturbation block. We used the first 3

perturbed strides as people tend to adapt quickly in arm reaching (39) and split-belt walking studies (40). For left toe

off metrics, early perturbed used perturbed strides 2-4, which excluded the first perturbed stride since left toe off

occurred before the perturbation onset. Late perturbed was the last 3 perturbed strides during the perturbation block,

regardless of when the metrics were calculated. Early catch and late catch referred to the first 2 and last 2

unperturbed strides during the 4-minute perturbation block, respectively. Early post was the first 3 strides while late

post was the last 3 strides during the last 2-minute post walking block.

2.3. Statistics

We performed statistical procedures using JMP (JMP Pro 12, SAS Institute Inc., Cary, NC, USA) and set the

significance level to 0.05. First, we checked and excluded outliers as being beyond 3 standard deviations from the

mean for each data set. We also used a Shapiro-Wilk W test, Levene's test, and Mauchly's test to check for the

distribution normality, homogeneity of variance, and sphericity, respectively. For the first three hypotheses related

to adaptation, using a feedforward strategy, and effects of perturbation directions, we analyzed the anterior

(Stick0.4), posterior (Slip0.4), medial (M1), and lateral (L1) perturbation conditions. For the fourth hypothesis

related to perturbation size effect, we analyzed the data of the 2 sizes of the anterior (Stick0.4, Stick0.2) and

posterior (Slip0.4, Slip0.2) perturbations.

For the first hypothesis regarding adaptation, we performed one-way repeated measures analysis of variance

(rANOVA) for the perturbed strides on each metric ($MOS_{ap, lhs}$, $MOS_{ml, lhs}$, $MOS_{ap, lto}$, $MOS_{ml, lto}$, $BOS_{ap, lhs}$, $BOS_{ml, lhs}$,

$BOS_{ap, lto}$, $BOS_{ml, lto}$, $XCOM_{ap, lhs}$, $XCOM_{ml, lhs}$, $XCOM_{ap, lto}$, and $XCOM_{ml, lto}$) with phase (pre, early perturbed, late

perturbed, early post, and late post) as a factor by perturbation condition. When phase was a main effect, we performed post hoc pairwise comparisons using Tukey's Honest Significant Difference (HSD) tests to account for multiple comparisons for the following apriori comparisons: 1) pre to early perturbed to assess the disruption of gait stability, 2) early perturbed to late perturbed to assess adaptation, 3) early post to late post to assess de-adaptation. To determine which component of margin of stability contributed more to changes in margin of stability ($|\Delta MOS|$), we compared the magnitudes of the changes of the base of support ($|\Delta BOS|$) and extrapolated center of mass position ($|\Delta XCOM|$) within a phase comparison (pre and early perturbed; early and late perturbed) using paired t-tests (i.e., for pre and early perturbed, $|\Delta BOS| = |BOS_{pre} - BOS_{early\_perturbed}|$ versus $|\Delta XCOM| = |XCOM_{pre} - XCOM_{early\_perturbed}|$; for early to late perturbed, $|\Delta BOS| = |BOS_{early\_perturbed} - BOS_{late\_perturbed}|$ versus $|\Delta XCOM| = |XCOM_{early\_perturbed} - XCOM_{late\_perturbed}|$).

For the second hypothesis regarding whether subjects used a feedforward strategy for adaptation, we performed one-way rANOVA for the catch strides on each metric with phase (pre, early catch, late catch) as a factor by perturbation condition. If phase was a main effect, we performed Tukey's HSD post hoc tests for the following comparisons: 1) pre to early catch to assess anticipatory responses at early catch, 2) pre to late catch to assess anticipatory responses at late catch, and 3) early catch to late catch to assess if subjects used more feedforward strategies during adaptation.

For the third hypothesis regarding perturbation direction, we also used a one-way rANOVA for each metric (ΔMOS, ΔBOS, and $\Delta XCOM$ for early perturbed minus pre; ΔMOS, ΔBOS, and $\Delta XCOM$ for late perturbed minus early perturbed) with perturbation condition as a factor. Here, the direction of the change, not just the magnitude, from pre to early and from early to late was important. We performed post hoc pairwise comparisons using Tukey's HSD if perturbation condition was a main effect.

For the fourth hypothesis regarding a scaling effect, we used a two-way rANOVA on each metric ($|\Delta MOS|$, $|\Delta BOS|$, and $|\Delta XCOM|$ for pre to early perturbed; $|\Delta MOS|$, $|\Delta BOS|$, and $|\Delta XCOM|$ for early to late perturbed) with perturbation condition (stick, slip) and size (regular, 0.4; half-size, 0.2) as factors. If a significant interaction effect was detected, we performed paired t-tests with Bonferroni correction ($\alpha = 0.025$) to identify differences between sizes for each perturbation condition.

We calculated effect size using partial eta squared (η_p^2) for the rANOVA results and Cohen's d for post hoc

pairwise comparisons and paired t-tests (41). Complete statistical results are provided in the supplemental data

(Supplemental Tables S1-S6; https://doi.org/10.6084/m9.figshare.16910683).

3. Results

3.1. Representative data

Changes of MOS and its components during perturbed walking were evident throughout the gait cycle (Fig. 3).

When perturbations first occurred at early perturbed, MOS_{lhs}, BOS_{lhs}, and $XCOM_{lhs}$ all deviated from the pre

unperturbed strides, indicating that the perturbations disrupted the subjects. As subjects gained more experience

with the perturbations, MOS_{lhs} and BOS_{lhs} of late perturbed generally trended back to pre levels, suggesting

adaptation occurred. When the perturbations were removed at the post block, MOS_{lhs} and its components were

almost the same as pre, suggesting a return to pre levels. The $MOS_{ap,\ lto}$ of the Stick0.4 perturbation, $BOS_{ap,\ lto}$ and

$XCOM_{ap,\ lto}$ of the Slip0.4 perturbation, and $BOS_{ml,\ lto}$ and $XCOM_{ml,\ lto}$ of the L1 perturbation deviated from pre at

early perturbed, suggesting an anticipatory response. The BOS_{lto} and $XCOM_{lto}$ of the Slip0.4 and L1 perturbations

then trended to their pre levels by late perturbed, an indication of adaptation, and returned to the pre levels by the

end of post.

3.2. Adaptation to small anterior-posterior perturbations applied on a stride-by-stride basis (hypothesis 1):

Subjects adapted to the Stick0.4 perturbation after an initial disruption to gait stability (Fig. 4A and

Supplemental Fig. S1A, all supplemental figures are available at https://doi.org/10.6084/m9.figshare.16910713).

Phase had a significant effect on all metrics except for $BOS_{ap,\ lto}$ for the Stick0.4 perturbation (see Supplemental

Table S1 for metric specific F, p, and η_p^2). Upon initial exposure to the perturbations, $MOS_{ap,\ lhs}$ was disrupted and

decreased significantly by 41.3% ($p < 0.001$, Cohen's d = 2.49). After the initial perturbation, subjects immediately

took shorter and wider steps and shifted their XCOM backward at left heel strike (pre to early perturbed: $BOS_{ap,\ lhs}$

decreased, $p < 0.001$, Cohen's d = 4.95; $BOS_{ml,\ lhs}$ increased, $p = 0.03$, Cohen's d = 0.99; $XCOM_{ap,\ lhs}$ decreased, $p <$

0.001, Cohen's d = 2.32). Starting with the second Stick0.4 perturbation, subjects also took wider steps (determined

by the previous right heel strike) and shifted their XCOM forward at left toe off in anticipation of an impending

perturbation at right leg mid-stance (pre to early perturbed: $BOS_{ml,\ lto}$ increased, $p < 0.001$, Cohen's d = 2.21;

$XCOM_{ap, lto}$ increased, $p < 0.001$, Cohen's $d = 1.83$). $MOS_{ap, lhs}$ increased significantly by 44.4% ($p < 0.001$, Cohen's

$d = 1.57$) after the initial disruption and trended toward pre level by late perturbed, indicating adaptation occurred.

Subjects also took longer steps at left heel strike and took narrower steps and shifted their XCOM backward at left

toe off (early to late perturbed: $BOS_{ap, lhs}$ increased, $p = 0.01$, Cohen's $d = 1.09$; $BOS_{ml, lto}$ decreased, $p = 0.001$,

Cohen's $d = 1.37$; $XCOM_{ap, lto}$ decreased, $p < 0.001$, Cohen's $d = 1.77$) as they adapted. $MOS_{ap, lhs}$ continued trending

to pre level during the post block (early post to late post, $p = 0.02$, Cohen's $d = 1.02$).

Like the Stick0.4 perturbation, subjects adapted to the Slip0.4 perturbation at both left heel strike and left toe

off after initially being disrupted (Fig. 4B and Supplemental Fig. S1B). Phase had a significant effect on all metrics

except for $MOS_{ap, lto}$ and $XCOM_{ml, lto}$ for the Slip0.4 perturbation (see Supplemental Table S1 for metric specific F, p,

and η_p^2). Upon initially experiencing the perturbations, anterior-posterior gait stability was disrupted as $MOS_{ap, lhs}$

decreased significantly by 26.6% ($p < 0.001$, Cohen's $d = 2.34$). Subjects also immediately took longer and wider

steps and shifted their XCOM forward and leftward at left heel strike (pre to early perturbed: $BOS_{ap, lhs}$ increased, p

< 0.001, Cohen's $d = 2.92$; $BOS_{ml, lhs}$ increased, $p = 0.02$, Cohen's $d = 1.05$; $XCOM_{ap, lhs}$ increased, $p < 0.001$,

Cohen's $d = 3.39$; $XCOM_{ml, lhs}$ increased, $p < 0.001$, Cohen's $d = 1.53$).   At left toe off, subjects also immediately

increased their step width and shifted their XCOM backward in anticipation of the impending perturbation at

mid-stance (pre to early perturbed: $BOS_{ml, lto}$ increased, $p = 0.02$, Cohen's $d = 1.02$; $XCOM_{ap, lto}$ decreased, $p < 0.001$,

Cohen's $d = 1.98$). The $MOS_{ap, lhs}$ increased significantly by 19.6% ($p = 0.003$, Cohen's $d = 1.27$) and trended toward

pre level from early to late perturbed during adaptation. Subjects also took longer steps and shifted their XCOM

forward at left toe off (early to late perturbed: $BOS_{ap, lto}$ increased, $p < 0.001$, Cohen's $d = 1.49$; $XCOM_{ap, lto}$

increased, $p < 0.001$, Cohen's $d = 1.92$). From early post to late post, $MOS_{ap, lhs}$ continued trending to pre level ($p =$

0.02, Cohen's $d = 1.01$).

3.3. Adaptation to small medial-lateral perturbations applied on a stride-by-stride basis (hypothesis 1):

The L1 perturbation was initially disruptive, but subjects adapted with more experience with the perturbation

(Fig. 5A and Supplemental Fig. S1C). Phase had a significant effect on all metrics except for $XCOM_{ap, lto}$ and

$MOS_{ml, lto}$ for the L1 perturbation (see Supplemental Table S1 for metric specific F, p, and η_p^2). After initial exposure

to the perturbations, $MOS_{ml, lhs}$ was disrupted and increased significantly by 47.3% ($p < 0.001$, Cohen's $d = 4.27$).

Subjects also immediately took shorter and wider steps and shifted their XCOM backward and rightward at left heel

strike (pre to early perturbed: $BOS_{ap, lhs}$ decreased, $p < 0.001$, Cohen's $d = 1.46$; $BOS_{ml, lhs}$ increased, $p = 0.008$,

Cohen's $d = 1.14$; $XCOM_{ap, lhs}$ decreased, $p = 0.004$, Cohen's $d = 1.22$; $XCOM_{ml, lhs}$ decreased, $p < 0.001$, Cohen's d

$= 1.84$). Starting with the second L1 perturbation, subjects showed anticipatory responses based on their wider steps

at left toe off (pre to early perturbed: $BOS_{ml, lto}$ increased, $p = 0.005$, Cohen's $d = 1.20$). From early to late perturbed,

$MOS_{ml, lhs}$ decreased significantly by 8.0% ($p = 0.02$, Cohen's $d = 1.06$) and trended toward pre level, indicating

adaptation to the L1 perturbation. Subjects also took longer and narrower steps at both left heel strike and left toe off

(early to late perturbed: $BOS_{ap, lhs}$ increased, $p = 0.01$, Cohen's $d = 1.09$; $BOS_{ml, lhs}$ decreased, $p = 0.05$, Cohen's $d =$

0.91; $BOS_{ap, lto}$ increased, $p = 0.02$, Cohen's $d = 1.02$; $BOS_{ml, lto}$ decreased, $p = 0.004$, Cohen's $d = 1.22$) as they

adapted. From early post to late post, $MOS_{ml, lhs}$ did not change significantly ($p = 0.87$, Cohen's $d = 0.30$), suggesting

that subjects rapidly restored pre level of MOS_{lhs} preference in the medial-lateral direction during the first few

strides of the post block.

Subjects also adapted to the M1 perturbation that initially disrupted gait stability (Fig. 5B and Supplemental

Fig. S1D). Phase had a significant effect on $MOS_{ap, lhs}$, $BOS_{ap, lhs}$, $XCOM_{ap, lhs}$, $MOS_{ml, lhs}$, $XCOM_{ml, lhs}$, and $BOS_{ap, lto}$

for the M1 perturbation (see Supplemental Table S1 for metric specific F, p, and η_p^2). Upon initially responding to

the perturbations, gait stability was disrupted in both directions as $MOS_{ap, lhs}$ decreased significantly by 20.3% ($p =$

0.002, Cohen's $d = 1.28$) and $MOS_{ml, lhs}$ decreased significantly by 36.3% ($p < 0.001$, Cohen's $d = 2.80$). After the

initial perturbation, subjects immediately shifted their XCOM forward and leftward at left heel strike (pre to early

perturbed: $XCOM_{ap, lhs}$ increased, $p = 0.01$, Cohen's $d = 1.09$; $XCOM_{ml, lhs}$ increased, $p < 0.001$, Cohen's $d = 1.88$).

At left toe off, subjects also immediately took shorter steps (pre to early perturbed: $BOS_{ap, lto}$ decreased, $p = 0.04$,

Cohen's $d = 0.94$). Subjects adapted after the initial disruption as $MOS_{ap, lhs}$ increased significantly by 23.9% ($p =$

0.004, Cohen's $d = 1.21$) and trended toward pre level by late perturbed. Subjects also shifted their XCOM

backward at left heel strike (early to late perturbed: $XCOM_{ap, lhs}$ decreased, $p = 0.01$, Cohen's $d = 1.11$).

3.4. Which component of margin of stability was affected more by the small perturbations?

During disruption, perturbations had different effects on the magnitudes of the changes in the BOS and XCOM

at left heel strike (Fig. 6A) and left toe off (Supplemental Fig. S2A). For the Stick0.4 perturbation, the BOS_{lhs} was

disrupted more than $XCOM_{lhs}$ based on the larger changes in the BOS_{lhs} compared to the changes in $XCOM_{lhs}$ that

were significantly different in the anterior-posterior direction and trending towards significance in the medial-lateral

direction ($|\Delta BOS_{ap, lhs}| > |\Delta XCOM_{ap, lhs}|$ p = 0.001, Cohen's d = 1.60); $|\Delta BOS_{ml, lhs}| > |\Delta XCOM_{ml, lhs}|$ p = 0.08,

Cohen's d = 0.62). For the Slip0.4 and M1 perturbations, the $XCOM_{lhs}$ was disrupted more than BOS_{lhs} as the

change in $XCOM_{lhs}$ was greater than the change in BOS_{lhs} in both directions (Slip0.4: $|\Delta XCOM_{ap, lhs}| > |\Delta BOS_{ap, lhs}|$ p

< 0.001, Cohen's d = 2.00; $|\Delta XCOM_{ml, lhs}| > |\Delta BOS_{ml, lhs}|$ p = 0.04, Cohen's d = 0.73; M1: $|\Delta XCOM_{ap, lhs}| > |\Delta BOS_{ap,}$

$_{lhs}|$ p = 0.06, Cohen's d = 0.68; $|\Delta XCOM_{ml, lhs}| > |\Delta BOS_{ml, lhs}|$ p < 0.001, Cohen's d = 1.93). For the L1 perturbation,

there was no difference between the changes in BOS_{lhs} and $XCOM_{lhs}$ in either direction (p's > 0.21, Cohen's d's <

0.43).

3.5. Which component contributed more to the adaptation of margin of stability?

During adaptation, the component that was the primary contributor to the adaptation of MOS at left heel strike

(Fig. 6B) and left toe off (Supplemental Fig. S2B) depended on the perturbation. The $|\Delta BOS_{ap, lhs}|$ was larger than the

$|\Delta XCOM_{ap, lhs}|$ for the L1 perturbation (p = 0.03, Cohen's d = 0.85). The $|\Delta XCOM_{ap, lhs}|$ was greater than the $|\Delta BOS_{ap,}$

$_{lhs}|$ for the Slip0.4 and M1 perturbations (p's < 0.05, Cohen's d's > 0.72), suggesting that $XCOM_{ap, lhs}$ adapted more

than $BOS_{ap, lhs}$ during those perturbations. There was no significant difference between $|\Delta BOS_{ap, lhs}|$ and $|\Delta XCOM_{ap,}$

$_{lhs}|$ in the Stick0.4 perturbation (p = 0.27, Cohen's d = 0.37). For the medial-lateral metrics, the only significant

difference was the larger $|\Delta BOS_{ml, lhs}|$ compared to the $|\Delta XCOM_{ml, lhs}|$ for the L1 perturbation (p = 0.02, Cohen's d =

0.90), suggesting that $BOS_{ml, lhs}$ adapted more in the lateral shift perturbation. There were no significant differences

between the $|\Delta BOS_{ml, lhs}|$ and $|\Delta XCOM_{ml, lhs}|$ in the other perturbations (p's > 0.12, Cohen's d's < 0.53).

3.6. Did subjects use more feedforward strategies when adapting to small perturbations applied on a

stride-by-stride basis? (hypothesis 2)

Catch strides revealed subjects used feedforward strategies upon initially experiencing the anterior-posterior

perturbations but did not use more anticipatory control as adaptation progressed (Fig. 7A, B and Supplemental Fig.

S1A, B). For the Stick0.4 perturbation, phase had a significant effect on all metrics except for $BOS_{ap, lto}$, $BOS_{ap, lhs}$,

$MOS_{ml, lhs}$ while for the Slip0.4 perturbation phase had a significant effect on $MOS_{ml, lto}$, $BOS_{ml, lto}$, $MOS_{ap, lhs}$,

$XCOM_{ap, lhs}$, $MOS_{ml, lhs}$, $BOS_{ml, lhs}$, and $XCOM_{ml, lhs}$ (see Supplemental Table S4 for metric specific F, p, and η_p^2). At

early catch, $MOS_{ap, lto}$ and $MOS_{ml, lto}$ deviated significantly from their pre levels for the Stick0.4 ($MOS_{ap, lto}$

decreased, p = 0.001, Cohen's d = 1.44; $MOS_{ml, lto}$ increased, p = 0.02, Cohen's d = 0.96) and the Slip0.4 ($MOS_{ml, lto}$

increased, $p = 0.004$, Cohen's $d = 1.20$) perturbations. Subjects also took wider steps for both perturbations (pre to early catch: $BOS_{ml, lto}$ increased, p's < 0.003, Cohen's d's > 1.25) and shifted their XCOM forward and rightward at left toe off for the Stick0.4 perturbation (pre to early catch: $XCOM_{ap, lto}$ increased, $p = 0.02$, Cohen's $d = 0.99$; $XCOM_{ml, lto}$ increased, $p = 0.009$, Cohen's $d = 1.07$). At left heel strike for the Stick0.4 perturbation, subjects marginally shifted their XCOM forward to 0.44 ± 0.05 m (pre to early catch: $XCOM_{ap, lhs}$, $p = 0.130$, Cohen's $d = 0.65$), which was in the opposite direction of the perturbed response of 0.35 ± 0.02 m, revealing evidence of anticipating an impending perturbation. $MOS_{ml, lto}$ and $MOS_{ap, lhs}$ trended to their pre levels by late catch for the Stick0.4 perturbation (early to late catch: $MOS_{ml, lto}$ decreased, $p = 0.02$, Cohen's $d = 0.96$; $MOS_{ap, lhs}$ increased, $p = 0.02$, Cohen's $d = 0.99$). Subjects also took narrower steps for the Stick0.4 perturbation (early to late catch: $BOS_{ml, lto}$ decreased, $p = 0.003$, Cohen's $d = 1.22$; $BOS_{ml, lhs}$ decreased, $p = 0.007$, Cohen's $d = 1.10$) and shifted their XCOM rightward at left heel strike for both perturbations (early to late catch: $XCOM_{ml, lhs}$ decreased, p's < 0.007, Cohen's d's > 1.11). Although $MOS_{ml, lto}$ and $MOS_{ml, lhs}$ increased significantly from pre to late catch for the Slip0.4 perturbation ($MOS_{ml, lto}$, $p = 0.03$, Cohen's $d = 0.91$; $MOS_{ml, lhs}$, $p = 0.03$, Cohen's $d = 0.88$), there were no significant differences between early catch and late catch ($MOS_{ml, lto}$, $p = 0.64$, Cohen's $d = 0.29$; $MOS_{ml, lhs}$, $p = 0.11$, Cohen's $d = 0.67$), suggesting that subjects did not use more feedforward strategies as they experienced more anterior-posterior perturbations.

Based on the responses to the catches, subjects employed feedforward strategies after the initial medial-lateral perturbations, however, more feedforward strategies were not used during adaptation (Fig. 7C, D and Supplemental Fig. S1C, D). Phase had a significant effect on $MOS_{ml, lto}$, $BOS_{ml, lto}$, $MOS_{ap, lhs}$, $XCOM_{ap, lhs}$, $BOS_{ml, lhs}$, and $XCOM_{ml, lhs}$ for the L1 perturbation and $MOS_{ap, lto}$, $BOS_{ap, lto}$, $MOS_{ml, lto}$, $BOS_{ap, lhs}$, $XCOM_{ap, lhs}$, and $MOS_{ml, lhs}$ for the M1 perturbation (see Supplemental Table S4 for metric specific F, p, and η_p^2). From pre to early catch, $MOS_{ml, lto}$ increased significantly for both perturbations (p's < 0.02, Cohen's $d > 1.04$). Subjects also took wider steps at left toe off for the L1 perturbation (pre to early catch: $BOS_{ml, lto}$ increased, $p = 0.004$, Cohen's $d = 1.20$). At left heel strike, subjects shifted their XCOM forward and leftward for the L1 perturbation (pre to early catch: $XCOM_{ap, lhs}$ increased, $p = 0.04$, Cohen's $d = 0.85$; $XCOM_{ml, lhs}$ increased, $p = 0.002$, Cohen's $d = 1.28$) and increased their $MOS_{ml, lhs}$ for the M1 perturbation (pre to early catch, $p = 0.002$, Cohen's $d = 1.27$), these trends were in the opposite direction of the responses for pre to early perturbed. These results suggest that subjects had anticipatory behavior

after the initial medial-lateral perturbations. From early to late catch, $MOS_{ml, lto}$ decreased significantly for both

perturbations (p's < 0.009, Cohen's d's > 1.08). Subjects also took narrower steps at left toe off and shifted their

XCOM rightward at left heel strike for the L1 perturbation (early to late catch: $BOS_{ml, lto}$ decreased, p $= 0.006$,

Cohen's d $= 1.13$; $XCOM_{ml, lhs}$ decreased, p $= 0.01$, Cohen's d $= 1.02$). For the M1 perturbation, $MOS_{ml, lhs}$ increased

and subjects took shorter steps and shifted their XCOM rightward at left heel strike (pre to late catch: $MOS_{ml, lhs}$

increased, p < 0.0001, Cohen's d $= 1.88$; $BOS_{ap, lhs}$ decreased, p $= 0.01$, Cohen's d $= 1.06$; $XCOM_{ap, lhs}$ decreased, p $=$

0.004, Cohen's d $= 1.20$), whereas no significant differences were found between their early catch and late catch

except for $XCOM_{ap, lhs}$ (early to late catch: $MOS_{ml, lhs}$, p $= 0.16$, Cohen's d $= 0.61$; $BOS_{ap, lhs}$, p $= 0.29$, Cohen's d $=$

0.49; $XCOM_{ap, lhs}$ decreased, p $= 0.02$, Cohen's d $= 0.96$). These results suggest that subjects generally did not use

more feedforward strategies as they adapted to the medial-lateral perturbations.

3.7. Are margin of stability and its components sensitive to directions of small perturbations? (hypothesis 3)

Anterior-posterior metrics at left heel strike were more sensitive to the anterior-posterior perturbations when

subjects were disrupted at early perturbed but not during adaptation from early to late perturbed (Fig. 8A). During

disruption, perturbation direction had a significant effect on each anterior-posterior metric at left heel strike (F's

$(3,27) > 7.92$, p's < 0.002, η_p^2's > 0.46). All perturbations produced a decrease in $MOS_{ap, lhs}$, during disruption of

which, the decreases for the Stick0.4 and Slip0.4 perturbations were observed in all subjects and significantly larger

than the decrease for the L1 perturbation (p's < 0.04, Cohen's d's > 0.94). For early perturbed, the $BOS_{ap, lhs}$ and

$XCOM_{ap, lhs}$ had the largest decrease with the Stick0.4 perturbation compared to the decrease with the L1

perturbation (p's < 0.02, Cohen's d's > 1.03). The $BOS_{ap, lhs}$ and $XCOM_{ap, lhs}$ had the largest increase with the Slip0.4

perturbation compared to the increase with the M1 perturbation for early perturbed (p's < 0.001, Cohen's d's $>$

1.72). During adaptation, all perturbations resulted in an increased $MOS_{ap, lhs}$ of similar magnitude (F $(3,27) = 1.36$,

p $= 0.28$, $\eta_p^2 = 0.13$). The Stick0.4 and L1 perturbations resulted in increases of $BOS_{ap, lhs}$ during adaptation, whereas

the only significant difference was between Stick0.4 and Slip0.4 perturbations mainly due to differences in sign (p $=$

0.04, Cohen's d $= 0.92$). Conversely, although perturbation direction did not have a significant main effect on

$XCOM_{ap, lhs}$ (F $(3,27) = 2.83$, p $= 0.06$, $\eta_p^2 = 0.24$), the Slip0.4 and M1 perturbations resulted in decreases of

$XCOM_{ap, lhs}$ during adaptation.

The medial-lateral metrics at left heel strike, $MOS_{ml, lhs}$ and $XCOM_{ml, lhs}$, were more sensitive to the

medial-lateral perturbations during disruption but not during adaptation (Fig. 8B). During disruption, perturbation

direction had a significant effect on each medial-lateral metric at left heel strike (F's $(3,27) > 5.32$, p's < 0.006, η_p^2's

> 0.36). For early perturbed, all subjects increased and decreased $MOS_{ml, lhs}$ in the L1 and M1 perturbations,

respectively. $MOS_{ml, lhs}$ had the largest increase with the L1 perturbation compared to the increase with the Stick0.4

perturbation (p < 0.001, Cohen's d = 2.07), while the $MOS_{ml, lhs}$ had the largest decrease with the M1 perturbation

compared to the decrease with the Slip0.4 perturbation (p < 0.001, Cohen's d = 1.67). All perturbations produced an

increase in $BOS_{ml, lhs}$, during disruption of which, the increase for the M1 perturbation was significantly smaller than

the increases for the Slip0.4 and L1 perturbations (p's < 0.03, Cohen's d's > 0.98). For the $XCOM_{ml, lhs}$ for early

perturbed, the increase of $XCOM_{ml, lhs}$ for all subjects in the M1 perturbation was significantly larger than the

increase in the Stick0.4 perturbation (p < 0.001, Cohen's d = 1.79), while the decrease of $XCOM_{ml, lhs}$ for all subjects

in the L1 perturbation was significantly different than the increases in the Stick0.4 and Slip0.4 perturbations (p's $<$

0.001, Cohen's d's > 1.83). During adaptation, perturbation direction had a significant effect on $\Delta MOS_{ml, lhs}$ (F

$(3,27) = 5.89$, p = 0.003, $\eta_p^2 = 0.40$). The increase of $MOS_{ml, lhs}$ in the M1 perturbation was significantly different

than the decrease in the L1 perturbation (p = 0.002, Cohen's d = 1.32), largely due to differences in sign. For $BOS_{ml,}$

$_{lhs}$ and $XCOM_{ml, lhs}$, nearly all perturbations resulted in a decrease of similar magnitudes during adaptation (F's

$(3,27) < 0.93$, p's > 0.44, η_p^2's < 0.10).

3.8. Do responses in margin of stability and its components scale with perturbation size? (hypothesis 4)

Anterior-posterior metrics at left heel strike scaled with perturbation size when subjects were disrupted at early

perturbed but not during adaptation from early to late perturbed (Fig. 9A). Perturbation size had a main effect on the

changes during disruption for each anterior-posterior metric at left heel strike (F's $(1,27) > 9.51$, p's < 0.006, η_p^2's $>$

0.25). There was a significant interaction effect between perturbation size and perturbation condition for the

changes in $BOS_{ap, lhs}$ during disruption (F $(1,27) = 5.53$, p = 0.03, $\eta_p^2 = 0.17$). Paired t-tests revealed that the half-size

perturbations produced significantly smaller magnitude changes in $BOS_{ap, lhs}$ than the regular perturbations within

each perturbation condition (Stick0.2 vs. Stick0.4 p = 0.001, Cohen's d = 1.61; Slip0.2 vs. Slip0.4 p = 0.01, Cohen's

d = 1.01). These results suggest that subjects scaled $BOS_{ap, lhs}$ more in the stick perturbations than slip perturbations

at early perturbed. During adaptation, perturbation size had a significant effect on $|\Delta BOS_{ap, lhs}|$ (F $(1,27) = 4.75$, p =

0.04, $\eta_p^2 = 0.15$) but not on $|\Delta MOS_{ap, lhs}|$ and $|\Delta XCOM_{ap, lhs}|$ (F's (1,27) < 1.70, p's > 0.20, η_p^2's < 0.07), indicating

most anterior-posterior metrics at left heel strike did not scale with perturbation size during adaptation.

Medial-lateral metrics at left heel strike did not scale with perturbation size when subjects were disrupted or

during adaptation (Fig. 9B). Perturbation size did not have a main effect on the changes during disruption for any

medial-lateral metric at left heel strike (F's (1,27) < 1.19, p's > 0.28, η_p^2's < 0.05). Despite the significant interaction

effect on $|\Delta XCOM_{ml, lhs}|$ during disruption (F (1,27) = 7.96, p = 0.009, $\eta_p^2 = 0.23$), paired t-tests only detected

significant differences between Slip perturbations (p = 0.02, Cohen's d = 0.93) but not between Stick perturbations

(p = 0.21, Cohen's d = 0.43). For the changes of each medial-lateral metric at left heel strike during adaptation, there

was neither a main effect of perturbation size nor a significant interaction effect (F's (1,27) < 1.23, p's > 0.27, η_p^2's

< 0.05). These results suggest that medial-lateral metrics at left heel strike did not scale with perturbation size during

disruption or adaptation.

4. Discussion

We sought to determine how small magnitude multidirectional perturbations affect modulation of margin of

stability, base of support, and extrapolated center of mass on a stride-by-stride basis. In the anterior-posterior

direction, subjects mainly modulated $XCOM_{ap, lhs}$ when adapting to the belt acceleration (Slip0.4) and medial shift

(M1) perturbations. In the medial-lateral direction, subjects primarily modulated $BOS_{ml, lhs}$ when adapting to the

lateral shift (L1) perturbation. Metrics at left toe off and catch strides revealed that subjects immediately took wider

steps for the Stick0.4, Slip0.4, and L1 perturbations and shifted their $XCOM_{ap, lto}$ forward for the Stick0.4

perturbation before perturbation onset, providing evidence of feedforward strategies. Despite the combination of

feedback and feedforward strategies upon initial exposure to the perturbations, subjects did not use more

feedforward strategies during adaptation. When first experiencing the perturbations, anterior-posterior metrics at

left heel strike for perturbed strides were more sensitive to perturbations in the coincident anterior-posterior

direction, and likewise, $MOS_{ml, lhs}$ and $XCOM_{ml, lhs}$ were more sensitive to perturbations in the coincident

medial-lateral direction. Additionally, anterior-posterior metrics at left heel strike scaled with perturbation size upon

initially responding to the perturbations (at disruption). These findings suggest that applying small perturbations on

a stride-by-stride basis led to the adaptation of the margin of stability, providing new insights regarding the extent of

locomotor adaptation to small perturbations. Further, perturbation directions could be used to target specific aspects

of gait stability control (e.g., primarily modulating base of support or center of mass).

One of the main findings was that small magnitude perturbations altered gait stability, i.e., the margin of

stability at left heel strike, in specific directions. For most perturbations, the $MOS_{ap, lhs}$ and $MOS_{ml, lhs}$ initially

decreased before adapting back to pre levels (Fig. 4, 5), suggesting that subjects were destabilized initially before

gradually regaining stability during the Stick0.4, Slip0.4, and M1 perturbations. These results are consistent with

previous studies that showed that treadmill belt and shift perturbations decreased MOS_{ap} (11, 25, 42). Adjusting

foot placement and BOS is a common approach to modulate MOS (6, 23), which was evident in our Stick0.4

perturbation. Other studies demonstrated the importance of center of mass control to modulate MOS (11, 30),

which was evident in the Slip0.4 and M1 perturbations. MOS_{lhs} did not always decrease in response to

perturbations. For the Stick0.4 and L1 perturbation, the $MOS_{ml, lhs}$ initially increased, however, instead of

decreasing as seen in the other perturbations. This increased $MOS_{ml, lhs}$ likely resulted from a combination of a

more lateral (leftward) foot placement (i.e., increased $BOS_{ml, lhs}$) and redirected $XCOM_{lhs}$ rightward during the

lateral shift perturbation. In another study that applied lateral shift perturbations at early single leg support, people

used the gluteus medius of their swing leg to place their subsequent foot more outward (9) in the direction of the

potential fall (27, 43), which would help increase the MOS_{ml} to maintain a more stable body position (19). Larger

MOS_{ml} often occurs in response to medial-lateral treadmill shift perturbations (42, 44, 45), similar to the increased

$MOS_{ml, lhs}$ in our L1 perturbation. Interestingly, subjects maintained this increased $MOS_{ml, lhs}$ for the Stick0.4

perturbation but gradually reduced the $MOS_{ml, lhs}$ for the L1 perturbation, suggesting that subjects either could not

or did not have a reason to decrease $MOS_{ml, lhs}$ while gaining more experience with the Stick0.4 perturbation.

At left heel strike, subjects primarily modulated foot placement compared to the extrapolated center of mass

in the belt deceleration (during disruption) and lateral shift (during adaptation) perturbations. Researchers are

increasingly finding that BOS control is an active neuromechanical balance strategy that is adjusted based on the

state of the center of mass during walking (4, 43, 46, 47). Additionally, the central nervous system is actively

involved in this foot placement strategy (48, 49). In our study, both the $BOS_{ap, lhs}$ and $XCOM_{ap, lhs}$ decreased in

response to the Stick0.4 perturbation, but subjects primarily adjusted $BOS_{ap, lhs}$ by taking shorter steps during

disruption. The reduction and smaller adjustment of $XCOM_{ap, lhs}$ compared to the $BOS_{ap, lhs}$ was likely a

consequence of the rotation and acceleration of the upper body backward relative to the lower body created by the

stick perturbations. The significant $BOS_{ap, lhs}$ adjustment was likely needed to rapidly interrupt the forward

progression of the swing leg to compensate for the interrupted progression of the stance leg caused by the Stick0.4

perturbation (26). These reactions conform to the theory of stepping in the direction of the potential fall and to the

maintenance of forward progression on the treadmill (27, 43, 50). During adaptation to the Stick0.4 perturbation,

there were similar magnitude changes in the $BOS_{ap, lhs}$ and $XCOM_{ap, lhs}$, suggesting a less dominant stepping

strategy during adaptation compared to the dominant foot placement strategy during disruption. For the lateral

shift perturbation, subjects appeared to employ combination of a foot placement strategy through swing leg

control that increased the $BOS_{ml, lhs}$ (9), while also using an ankle strategy with the stance leg that decreased

$XCOM_{ml, lhs}$ by a similar magnitude (51). Changes in the $BOS_{ml, lhs}$ then dominated during adaptation to the L1

perturbation. This aligns with the predominant stepping strategy people often employ, taking shorter and wider

steps during medial waist-pull perturbations that produced relative movements of the BOS and XCOM similar to

our L1 perturbation (52).

Our results also showed that subjects primarily modulated the extrapolated center of mass at left heel strike

in the belt acceleration and medial shift perturbations. Both the $BOS_{ap, lhs}$ and $XCOM_{ap, lhs}$ increased in response to

our Slip0.4 perturbation, but subjects mainly modulated their $XCOM_{ap, lhs}$ while adapting to the Slip0.4

perturbation. The $XCOM_{ap, lhs}$ responses were likely related to the decreased hip flexor moment of the right stance

leg to reduce the forward rotation and acceleration of the upper body (11). Even though the $BOS_{ap, lhs}$ also changed

during adaptation, these changes were smaller than the $XCOM_{ap, lhs}$ and were probably a consequence of the

facilitated forward swing phase induced by the Slip0.4 perturbation. For the medial shift perturbation, subjects

primarily adjusted $XCOM_{lhs}$, interestingly, in the orthogonal anterior-posterior direction, likely by employing an

ankle strategy (27) that incorporated increased tibialis anterior activity of the stance leg (51). Subjects barely

adjusted the $BOS_{ap, lhs}$, likely due to the small magnitude of our M1 perturbation. In other medial-lateral

perturbations, subjects often take crossover steps to adjust BOS in the anterior-posterior direction. (53, 54).

Our results suggest that subjects can flexibly employ balance strategies in response to small magnitude

perturbations applied on a stride-by-stride basis, depending on the perturbation direction. When the central

nervous system detects a perceived fall, the foot placement strategy shifts the swing foot to the direction of the

potential fall, such that at heel strike the undesired movement induced by perturbations could be reversed or at

least mitigated (27, 28). Another strategy is adjusting the extrapolated center of mass to stop the potential fall

through the stance leg, which is similar to the ankle strategy in other studies (27, 52, 55). The foot placement

strategy has the advantage of a large adjustment range, however, it acts slowly due to the time needed for active

integration of sensory information from the proprioceptive, visual, and vestibular systems (27, 28). Conversely,

the extrapolated center of mass strategy allows for quick responses during single stance, whereas its modulation

range is constrained to the relatively small contact area under the stance foot (27, 55, 56). A recent visual

perturbation study shows that foot placement and ankle strategies are interdependent and negatively correlated,

suggesting that these two strategies complement each other (57). Both strategies may increase the energetic cost

as a consequence of the cost of redirecting foot placement and extrapolated center of mass (58–61), but adaptation

over time usually leads to a more economical gait (62, 63). Thus, depending on the perturbation direction, people

appear to recruit more readily available and efficient balance strategies for gait stability when experiencing small

magnitude perturbations applied on a stride-by-stride basis.

In our study, subjects employed a feedforward strategy that involved direction independent foot placement

but direction dependent extrapolated center of mass control after being initially perturbed. Subjects immediately

took wider steps at left toe off for the early catch strides after experiencing the Stick0.4, Slip0.4, and L1

perturbations, indicating a direction independent anticipatory strategy (Fig. 7A, B, C). These changes reflect

anticipatory responses because left toe off occurs immediately before the expected perturbation at right mid-stance.

For pre to early perturbed, like the catches, subjects also took wider steps at left toe off (Fig. 4 and 5A), suggesting

a general feedforward strategy of increasing step width. Taking wider steps can increase the dynamic stability in

the medial-lateral direction (27, 64) and can help prepare for an upcoming perturbation in any direction. For the

Stick0.4 perturbation, $XCOM_{ap, lto}$ shifted forward for early catch (Fig. 7A) and early perturbed (Fig. 4A), which

was opposite to the potential fall direction of the Stick0.4 perturbation, revealing direction dependent anticipatory

XCOM adjustments. Forward shifts of the COM or XCOM position to prepare for impending backward fall

inducing perturbations (65–67) and a reduction of forward COM velocity to prepare for impending forward fall

inducing perturbations (68, 69) are common anticipatory balance strategies in response to anterior-posterior

perturbations. The lack of anterior-posterior XCOM adjustments for the Slip0.4 perturbation was likely due to the

less risk of forward fall compared to backward fall (10, 70).

Although subjects initially anticipated, they did not rely more on feedforward strategies when adapting to

small magnitude perturbations applied on a stride-by-stride basis. In response to repeated perturbations, people

often adapt using feedback (reactive) and feedforward (proactive) control (71, 72). Feedback control relies on

sensory detection of unexpected disturbances to dynamic stability, whereas feedforward control relies on

knowledge about the expected perturbation generated by prior experience (54, 73). In our study, subjects

apparently used feedback control when adapting to the treadmill belt and shift perturbations as MOS and its

components did not deviate more from the pre levels with more experience with catch strides. Despite the

stride-by-stride exposure to the perturbations, subjects did not appear to develop anticipatory responses. The small

perturbations in our study likely contributed to the use of feedback control, as mild medial-lateral perturbations

reduce feedforward control and facilitate feedback control in the medial-lateral direction (74). Additionally, the

repetitive fixed magnitude treadmill belt perturbations also likely contributed to the use of feedback control.

Unlike overground walking where subjects can control the displacement of their stance foot and the magnitude of

a slip perturbation which allows for more proactive responses (65), subjects are unable to control the displacement

of their stance foot during treadmill belt perturbations, resulting in reactive compensatory step lengths and trunk

angles responses (75).

Perturbations in the coincident direction more strongly affected gait stability after being initially perturbed.

As one might expect, the anterior-posterior and medial-lateral perturbations resulted in a stronger effect in their

respective directions (7, 76). In our study, this directional effect was not merely seen in the MOS_{lhs} but was also

reflected in its components. Despite the small magnitude, our perturbations also affected metrics in perpendicular

directions, though not as strongly as perturbations in coincident directions. Perturbations in one direction often

elicit responses perpendicular to the perturbation direction (4, 7, 76, 77) as responses between the sagittal and

frontal planes are dynamically coupled (31, 78). Medial-lateral perturbations are often more challenging than the

anterior-posterior perturbations (8, 76) because medial-lateral balance requires active control during walking (47,

79). In our study, medial-lateral perturbations were not more challenging than the anterior-posterior perturbations

in the anterior-posterior direction, which could be related to the magnitude of the shifts. We used 1 cm

medial-lateral shift perturbations as opposed to the 12-15 cm shifts that were used in other studies (8, 76). This

suggests that there is a threshold, a medial-lateral shift > 1 cm that is needed to elicit a significant effect on gait

stability compared to small anterior-posterior perturbations. We chose relatively small perturbation magnitudes to

avoid falls and extra recovery steps as we applied the perturbations on a stride-by-stride basis.

As expected, there was a scaling effect of perturbation magnitude on the anterior-posterior metrics as people

tend to magnify their responses proportionally with perturbation size (7, 26, 77). Changes in $MOS_{ml, lhs}$ and its

components were similar between the two treadmill belt perturbations sizes. In our study, the $MOS_{ml, lhs}$ did not

respond strongly to the anterior-posterior perturbations (Stick0.4 and Slip0.4) so an even smaller half-sized

perturbation (Stick0.2 and Slip0.2) would not be likely to elicit a strong response. Based on the interaction effect

for $BOS_{ap, lhs}$ observed in our anterior-posterior perturbations (Fig. 9A), we found that the stick perturbations had a

stronger scaling effect than slip perturbations for the same increment (from 0.2 to 0.4). Backward instability

created by belt decelerations (stick perturbations) tend to be more challenging than forward instability created by

the belt accelerations (slip perturbations) (10, 70).

The calculation of MOS is based on an inverted pendulum model and has several assumptions (19). First, the

mass of the whole body is modeled as a single mass point of the inverted pendulum, so the possible effects of

upper body (arm or trunk) movements in response to perturbations are ignored (18). Second, the leg is assumed to

be rigid and with constant length, which does not consider the large effects of the knee joint motions and moments

in response to perturbations (24, 80). Third, the excursions of the COM are small with respect to pendulum length

(19), which was likely met in our study due to the small magnitude perturbations. While all of the assumptions

were not likely met in our study, the MOS is a common metric of balance for studies of walking with large

perturbations (6, 11, 23–26, 29, 30) and the proposed approaches to fulfill the MOS assumptions vary among

studies (81).

The present study had several limitations. One limitation of our study is the small sample size. Post-hoc power

analyses indicated that hypotheses 1 and 3 were adequately powered while hypotheses 2 and 4 were underpowered,

which means we could not reliably detect small statistical differences for all hypotheses. The onsets of the

medial-lateral perturbations had a larger delay compared to the anterior-posterior perturbations. The perturbation

did not always return to the neutral position or baseline speed before the left heel strike, which likely introduced tiny

but extra disturbances. The treadmill system also has a fixed acceleration value for shifting the treadmill platform

medial-laterally, which limited the size of the shift perturbation we could use to ≤ 1 cm for our desired time window

such that the treadmill could return to its neutral position at or near left heel strike. To control for the time of the
perturbation, we set the duration of the anterior-posterior perturbations to be similar to the medial-lateral
perturbations, which again limited the magnitude of the perturbation we could apply. We only applied perturbations
to the right leg to limit the experiment to ~1 hour. We would not have expected a difference between perturbing the
left or right stepping limbs during walking or in forward falls (82, 83). Last, we only applied perturbations at one
instance of the gait cycle, right leg mid-stance. A recent study applied brief treadmill belt accelerations at push off of
each foot on a stride-by-stride basis and that led to sustained increased push-off once the perturbations were
removed (84). Applying perturbations at different points of the gait cycle may reveal phase-dependent modulation
of margin of stability and its components.

Future directions include studies to determine the muscular and cortical mechanisms involved during
adaptation of gait stability in response to small magnitude perturbations during walking. We recently found
significant differences in electrocortical activity of the anterior cingulate and motor cortices in response to small
magnitude mechanical perturbations applied on a stride-by-stride basis during recumbent stepping using
electroencephalography (EEG) and source estimation (85). Recording EEG and performing source estimation for
subjects experiencing small magnitude perturbations on a stride-by-stride basis during walking may reveal specific
cortical networks for modulating foot placement versus center of mass when adapting to perturbations during
walking. Currently, EEG gait adaptation studies often use large infrequent perturbations during walking (86, 87).

5. Conclusions

We investigated how healthy young individuals adjusted their margin of stability, base of support, and
extrapolated center of mass in response to small magnitude multidirectional perturbations applied on a
stride-by-stride basis. Small magnitude perturbations disrupted gait stability by decreasing the $MOS_{ap, lhs}$ in the belt
deceleration (Stick0.4), belt acceleration (Slip0.4), and medial treadmill shift (M1) perturbations, while increasing
the $MOS_{ml, lhs}$ in the lateral treadmill shift (L1) perturbation. During adaptation, BOS (foot placement) control was
primarily used in the L1 perturbations, whereas XCOM control was preferred in the Slip0.4 and M1 perturbations.
Subjects used both feedback and feedforward strategies after being initially perturbed, but primarily used feedback
strategies when adjusting their BOS and XCOM during adaptation. Moreover, MOS_{lhs} and its components were
generally more sensitive to perturbations in the coincident direction and scaled with perturbation size upon initial

exposure to the perturbations. Overall, our findings provide insights into inter-stride gait stability and adaptability to

small magnitude perturbations experienced on a stride-by-stride basis and have implications for the development of

training interventions that could target specific components of gait stability (e.g., base of support or extrapolated

center of mass control) for balance-impaired populations.

SUPPLEMENTAL DATA

Supplemental Tables S1–S6: https://doi.org/10.6084/m9.figshare.16910683

Supplemental Fig. S1–S2: https://doi.org/10.6084/m9.figshare.16910713

ACKNOWLEDGMENTS

The authors thank Alejandro Hurtado, Kevin Graydon, Amy Lebanoff, and Allen Rahrooh for assistance with data

collection.

GRANTS

This study was funded by the National Institute on Aging of the National Institutes of Health (NIH), under award number

R01AG054621 to H.J.H and a University of Central Florida Office of Research In-House Grant to H.J.H.

DISCLOSURES

No conflicts of interest, financial or otherwise, are declared by the authors.

AUTHOR CONTRIBUTIONS

J.L. and H.J.H. conceived and designed research; J.L. performed experiments; J.L. analyzed data; J.L. and H.J.H.

interpreted results of experiments; J.L. prepared figures; J.L. drafted manuscript; J.L. and H.J.H. edited and revised manuscript;

J.L. and H.J.H. approved final version of manuscript.

ENDNOTE

At the request of the authors, readers are herein alerted to the fact that additional materials related to this manuscript may be

found at https://doi.org/10.6084/m9.figshare.c.5285258.v3. These materials are not a part of this manuscript and have not

undergone peer review by the American Physiological Society (APS). APS and the journal editors take no responsibility for

these materials, for the website address, or for any links to or from it.

REFERENCES

1. **Torres-Oviedo G**, **Vasudevan E**, **Malone L**, **Bastian AJ**. Locomotor adaptation. *Prog Brain Res* 191: 65–74, 2011. doi:
10.1016/B978-0-444-53752-2.00013-8.

2. **Mansfield A**, **Wong JS**, **Bryce J**, **Knorr S**, **Patterson KK**. Does perturbation-based balance training prevent falls?
Systematic review and meta-analysis of preliminary randomized controlled trials. *Phys Ther* 95: 700–709, 2015. doi:
10.2522/ptj.20140090.

3. **Buurke TJW**, **Lamoth CJC**, **Vervoort D**, **van der Woude LHV**, **den Otter R**. Adaptive control of dynamic balance in
human gait on a split-belt treadmill. *J Exp Biol* 221, 2018. doi: 10.1242/jeb.174896.

4. **O'Connor SM**, **Kuo AD**. Direction-dependent control of balance during walking and standing. *J Neurophysiol* 102:
1411–1419, 2009. doi: 10.1152/jn.00131.2009.

5. **Franz JR**, **Francis C**, **Allen M**, **Thelen DG**. Visuomotor Entrainment and the Frequency-Dependent Response of
Walking Balance to Perturbations. .

6. **Hof AL**, **Vermerris SM**, **Gjaltema WA**. Balance responses to lateral perturbations in human treadmill walking. *J Exp
Biol* 213: 2655–2664, 2010. doi: 10.1242/jeb.042572.

7. **Vlutters M**, **van Asseldonk EHF**, **van der Kooij H**. Lower extremity joint-level responses to pelvis perturbation during
human walking. *Sci Rep* 8: 14621, 2018. doi: 10.1038/s41598-018-32839-8.

8. **McIntosh EI**, **Zettel JL**, **Vallis LA**. Stepping Responses in Young and Older Adults Following a Perturbation to the
Support Surface During Gait. *J Mot Behav* 49: 288–298, 2017. doi: 10.1080/00222895.2016.1204262.

9. **Afschrift M**, **Pitto L**, **Aerts W**, **van Deursen R**, **Jonkers I**, **De Groote F**. Modulation of gluteus medius activity reflects
the potential of the muscle to meet the mechanical demands during perturbed walking. *Sci Rep* 8: 11675, 2018. doi:
10.1038/s41598-018-30139-9.

10. **Patel PJ**, **Bhatt T**. Fall risk during opposing stance perturbations among healthy adults and chronic stroke survivors. *Exp
Brain Res* 236: 619–628, 2018. doi: 10.1007/s00221-017-5138-6.

11. **Debelle H**, **Harkness-Armstrong C**, **Hadwin K**, **Maganaris CN**, **O'Brien TD**. Recovery From a Forward Falling Slip:
Measurement of Dynamic Stability and Strength Requirements Using a Split-Belt Instrumented Treadmill. *Frontiers in
Sports and Active Living* 2: 82, 2020. doi: 10.3389/fspor.2020.00082.

12. **Gerards MHG**, **Marcellis RGJ**, **Poeze M**, **Lenssen AF**, **Meijer K**, **de Bie RA**. Perturbation-based balance training to
improve balance control and reduce falls in older adults - study protocol for a randomized controlled trial. *BMC Geriatr*
21: 9, 2021. doi: 10.1186/s12877-020-01944-7.

13. **Harper SA**, **Beethe AZ**, **Dakin CJ**, **Bolton DAE**. Promoting Generalized Learning in Balance Recovery Interventions.
Brain Sci 11, 2021. doi: 10.3390/brainsci11030402.

14. **Lee A**, **Bhatt T**, **Liu X**, **Wang Y**, **Wang S**, **Pai Y-CC**. Can Treadmill Slip-Perturbation Training Reduce Longer-Term Fall Risk Upon Overground Slip Exposure? .

15. **Karamanidis K**, **Epro G**, **McCrum C**, **König M**. Improving Trip- and Slip-Resisting Skills in Older People: Perturbation Dose Matters. *Exerc Sport Sci Rev* 48: 40–47, 2020. doi: 10.1249/JES.0000000000000210.

16. **Gerards MHG**, **McCrum C**, **Mansfield A**, **Meijer K**. Perturbation-based balance training for falls reduction among older adults: Current evidence and implications for clinical practice [Online]. *Geriatr Gerontol Int* 17: 2294–2303, 2017. https://onlinelibrary.wiley.com/doi/abs/10.1111/ggi.13082.

17. **Dusane S**, **Bhatt T**. Effect of Multisession Progressive Gait-Slip Training on Fall-Resisting Skills of People with Chronic Stroke: Examining Motor Adaptation in Reactive Stability. *Brain Sciences* 11: 894, 2021. doi: 10.3390/brainsci11070894.

18. **Bruijn SM**, **Meijer OG**, **Beek PJ**, **van Dieën JH**. Assessing the stability of human locomotion: a review of current measures. *J R Soc Interface* 10: 20120999, 2013. doi: 10.1098/rsif.2012.0999.

19. **Hof AL**, **Gazendam MGJ**, **Sinke WE**. The condition for dynamic stability. *J Biomech* 38: 1–8, 2005. doi: 10.1016/j.jbiomech.2004.03.025.

20. **Rosenblatt NJ**, **Grabiner MD**. Measures of frontal plane stability during treadmill and overground walking. *Gait Posture* 31: 380–384, 2010. doi: 10.1016/j.gaitpost.2010.01.002.

21. **Höhne A**, **Stark C**, **Brüggemann G-P**, **Arampatzis A**. Effects of reduced plantar cutaneous afferent feedback on locomotor adjustments in dynamic stability during perturbed walking. *J Biomech* 44: 2194–2200, 2011. doi: 10.1016/j.jbiomech.2011.06.012.

22. **Hof AL**. The "extrapolated center of mass" concept suggests a simple control of balance in walking. *Hum Mov Sci* 27: 112–125, 2008. doi: 10.1016/j.humov.2007.08.003.

23. **Süptitz F**, **Moreno Catalá M**, **Brüggemann G-P**, **Karamanidis K**. Dynamic stability control during perturbed walking can be assessed by a reduced kinematic model across the adult female lifespan. *Hum Mov Sci* 32: 1404–1414, 2013. doi: 10.1016/j.humov.2013.07.008.

24. **Hak L**, **Houdijk H**, **van der Wurff P**, **Prins MR**, **Mert A**, **Beek PJ**, **van Dieën JH**. Stepping strategies used by post-stroke individuals to maintain margins of stability during walking. *Clin Biomech* 28: 1041–1048, 2013. doi: 10.1016/j.clinbiomech.2013.10.010.

25. **McCrum C**, **Karamanidis K**, **Willems P**, **Zijlstra W**, **Meijer K**. Retention, savings and interlimb transfer of reactive gait adaptations in humans following unexpected perturbations. *Commun Biol* 1: 230, 2018. doi: 10.1038/s42003-018-0238-9.

26. **Martelli D**, **Aprigliano F**, **Tropea P**, **Pasquini G**, **Micera S**, **Monaco V**. Stability against backward balance loss: Age-related modifications following slip-like perturbations of multiple amplitudes. *Gait Posture* 53: 207–214, 2017. doi: 10.1016/j.gaitpost.2017.02.002.

27. **Reimann H**, **Fettrow T**, **Jeka JJ**. Strategies for the Control of Balance During Locomotion. *Kinesiology Review* 7: 18–25, 2018. doi: 10.1123/kr.2017-0053.

28. **Bruijn SM**, **van Dieën JH**. Control of human gait stability through foot placement. *J R Soc Interface* 15, 2018. doi: 10.1098/rsif.2017.0816.

29. **Martelli D**, **Kang J**, **Agrawal SK**. A single session of perturbation-based gait training with the A-TPAD improves dynamic stability in healthy young subjects. In: *2017 International Conference on Rehabilitation Robotics (ICORR)*, p. 479–484.

30. **McCrum C**, **Karamanidis K**, **Grevendonk L**, **Zijlstra W**, **Meijer K**. Older adults demonstrate interlimb transfer of reactive gait adaptations to repeated unpredictable gait perturbations. .

31. **Buurke TJW**, **Liu C**, **Park S**, **den Otter R**, **Finley JM**. Maintaining sagittal plane balance compromises frontal plane balance during reactive stepping in people post-stroke. *Clin Biomech*   80: 105135, 2020. doi: 10.1016/j.clinbiomech.2020.105135.

32.   **Huang HJ**, **Kram R**, **Ahmed AA**. Reduction of metabolic cost during motor learning of arm reaching dynamics. *J*
Neurosci 32: 2182–2190, 2012. doi: 10.1523/JNEUROSCI.4003-11.2012.

33.   **Shadmehr R**, **Mussa-Ivaldi FA**. Adaptive representation of dynamics during learning of a motor task. *J Neurosci* 14:
3208–3224, 1994. doi: 10.1523/JNEUROSCI.14-05-03208.1994.

34.   **Hreljac A**, **Marshall RN**. Algorithms to determine event timing during normal walking using kinematic data [Online]. *J*
Biomech 33: 783–786, 2000. https://www.ncbi.nlm.nih.gov/pubmed/10808002.

35.   **De Witt JK**. Determination of toe-off event time during treadmill locomotion using kinematic data. *J Biomech* 43: 3067–
3069, 2010. doi: 10.1016/j.jbiomech.2010.07.009.

36.   **Okubo Y**, **Brodie MA**, **Sturnieks DL**, **Hicks C**, **Lord SR**. A pilot study of reactive balance training using trips and slips
with increasing unpredictability in young and older adults: Biomechanical mechanisms, falls and clinical feasibility. *Clin*
Biomech   67: 171–179, 2019. doi: 10.1016/j.clinbiomech.2019.05.016.

37.   **Beltran EJ**, **Dingwell JB**, **Wilken JM**. Margins of stability in young adults with traumatic transtibial amputation
walking in destabilizing environments. *J Biomech* 47: 1138–1143, 2014. doi: 10.1016/j.jbiomech.2013.12.011.

38.   **Finley JM**, **Bastian AJ**, **Gottschall JS**. Learning to be economical: the energy cost of walking tracks motor adaptation. *J*
Physiol 591: 1081–1095, 2013. doi: 10.1113/jphysiol.2012.245506.

39.   **Pienciak-Siewert A**, **Barletta AJ**, **Ahmed AA**. Transfer of postural adaptation depends on context of prior exposure. *J*
Neurophysiol 111: 1466–1478, 2014. doi: 10.1152/jn.00235.2013.

40.   **Malone LA**, **Bastian AJ**. Age-related forgetting in locomotor adaptation. *Neurobiol Learn Mem* 128: 1–6, 2016. doi:
10.1016/j.nlm.2015.11.003.

41.   **Lakens D**. Calculating and reporting effect sizes to facilitate cumulative science: a practical primer for t-tests and
ANOVAs. *Front Psychol* 4: 863, 2013. doi: 10.3389/fpsyg.2013.00863.

42.   **McAndrew Young PM**, **Wilken JM**, **Dingwell JB**. Dynamic margins of stability during human walking in destabilizing
environments. *J Biomech* 45: 1053–1059, 2012. doi: 10.1016/j.jbiomech.2011.12.027.

43.   **Wang Y**, **Srinivasan M**. Stepping in the direction of the fall: the next foot placement can be predicted from current
upper body state in steady-state walking. *Biol Lett* 10, 2014. doi: 10.1098/rsbl.2014.0405.

44.   **Hak L**, **Houdijk H**, **Steenbrink F**, **Mert A**, **van der Wurff P**, **Beek PJ**, **van Dieën JH**. Speeding up or slowing down?:
Gait adaptations to preserve gait stability in response to balance perturbations. *Gait Posture* 36: 260–264, 2012. doi:
10.1016/j.gaitpost.2012.03.005.

45.   **Onushko T**, **Boerger T**, **Van Dehy J**, **Schmit BD**. Dynamic stability and stepping strategies of young healthy adults
walking on an oscillating treadmill. *PLoS One* 14: e0212207, 2019. doi: 10.1371/journal.pone.0212207.

46.   **van Leeuwen AM**, **van Dieën JH**, **Daffertshofer A**, **Bruijn SM**. Active foot placement control ensures stable gait:
Effect of constraints on foot placement and ankle moments. *PLoS One* 15: e0242215, 2020. doi:
10.1371/journal.pone.0242215.

47.   **Reimold NK**, **Knapp HA**, **Henderson RE**, **Wilson L**, **Chesnutt AN**, **Dean JC**. Altered active control of step width in
response to mediolateral leg perturbations while walking. *Sci Rep* 10: 12197, 2020. doi: 10.1038/s41598-020-69052-5.

48.   **Bruijn SM**, **Van Dieën JH**, **Daffertshofer A**. Beta activity in the premotor cortex is increased during stabilized as
compared to normal walking. *Front Hum Neurosci* 9: 593, 2015. doi: 10.3389/fnhum.2015.00593.

49.   **Sipp AR**, **Gwin JT**, **Makeig S**, **Ferris DP**. Loss of balance during balance beam walking elicits a multifocal theta band
electrocortical response. *J Neurophysiol* 110: 2050–2060, 2013. doi: 10.1152/jn.00744.2012.

50.   **Winter DA**. Human balance and posture control during standing and walking. *Gait Posture* 3: 193–214, 1995. doi:
10.1016/0966-6362(96)82849-9.

51.   **Afschrift M**, **van Deursen R**, **De Groote F**, **Jonkers I**. Increased use of stepping strategy in response to medio-lateral
perturbations in the elderly relates to altered reactive tibialis anterior activity. *Gait Posture* 68: 575–582, 2019. doi:
10.1016/j.gaitpost.2019.01.010.

52.   **Matjačić Z**, **Zadravec M**, **Olenšek A**. Influence of Treadmill Speed and Perturbation Intensity on Selection of
Balancing Strategies during Slow Walking Perturbed in the Frontal Plane. *Appl Bionics Biomech* 2019: 1046459, 2019.
doi: 10.1155/2019/1046459.

53.   **Borrelli J**, **Creath RA**, **Pizac D**, **Hsiao H**, **Sanders OP**, **Rogers MW**. Perturbation-evoked lateral steps in older adults:
Why take two steps when one will do? *Clin Biomech* 63: 41–47, 2019. doi: 10.1016/j.clinbiomech.2019.02.014.

54.   **Major MJ**, **Serba CK**, **Chen X**, **Reimold N**, **Ndubuisi-Obi F**, **Gordon KE**. Proactive Locomotor Adjustments Are
Specific to Perturbation Uncertainty in Below-Knee Prosthesis Users. *Sci Rep* 8: 1863, 2018. doi:
10.1038/s41598-018-20207-5.

55.   **Hof AL**, **Duysens J**. Responses of human ankle muscles to mediolateral balance perturbations during walking. *Hum Mov
Sci* 57: 69–82, 2018. doi: 10.1016/j.humov.2017.11.009.

56.   **Reimann H**, **Fettrow T**, **Grenet D**, **Thompson ED**, **Jeka JJ**. Phase-Dependency of Medial-Lateral Balance Responses
to Sensory Perturbations During Walking. *Frontiers in Sports and Active Living* 1: 25, 2019. doi:
10.3389/fspor.2019.00025.

57.   **Fettrow T**, **Reimann H**, **Grenet D**, **Thompson E**, **Crenshaw J**, **Higginson J**, **Jeka J**. Interdependence of balance
mechanisms during bipedal locomotion. *PLoS One* 14: e0225902, 2019. doi: 10.1371/journal.pone.0225902.

58.   **Donelan JMJM**, **Shipman DWDW**, **Kram R**, **Kuo ADAD**. Mechanical and metabolic requirements for active lateral
stabilization in human walking. *J Biomech* 37: 827–835, 2004. doi: 10.1016/j.jbiomech.2003.06.002.

59.   **Ijmker T**, **Houdijk H**, **Lamoth CJC**, **Beek PJ**, **van der Woude LHV**. Energy cost of balance control during walking
decreases with external stabilizer stiffness independent of walking speed. *J Biomech* 46: 2109–2114, 2013. doi:
10.1016/j.jbiomech.2013.07.005.

60.   **IJmker T**, **Lamoth CJ**, **Houdijk H**, **van der Woude LHV**, **Beek PJ**. Postural threat during walking: effects on energy
cost and accompanying gait changes. *J Neuroeng Rehabil* 11: 71, 2014. doi: 10.1186/1743-0003-11-71.

61.   **Wu M**, **Brown GL**, **Woodward JL**, **Bruijn SM**, **Gordon KE**. A novel Movement Amplification environment reveals
effects of controlling lateral centre of mass motion on gait stability and metabolic cost. *R Soc Open Sci* 7: 190889, 2020.
doi: 10.1098/rsos.190889.

62.   **Selinger JC**, **O'Connor SM**, **Wong JD**, **Donelan JM**. Humans Can Continuously Optimize Energetic Cost during
Walking. *Curr Biol* 25: 2452–2456, 2015. doi: 10.1016/j.cub.2015.08.016.

63.   **Sánchez N**, **Park S**, **Finley JM**. Evidence of Energetic Optimization during Adaptation Differs for Metabolic,
Mechanical, and Perceptual Estimates of Energetic Cost. *Sci Rep* 7: 7682, 2017. doi: 10.1038/s41598-017-08147-y.

64.   **Wu M**, **Matsubara JH**, **Gordon KE**. General and Specific Strategies Used to Facilitate Locomotor Maneuvers. *PLoS
One* 10: e0132707, 2015. doi: 10.1371/journal.pone.0132707.

65.   **Bhatt T**, **Wening JD**, **Pai Y-C**. Adaptive control of gait stability in reducing slip-related backward loss of balance. *Exp
Brain Res* 170: 61–73, 2006. doi: 10.1007/s00221-005-0189-5.

66.   **Yang F**, **Pai CY-C**. Alteration in community-dwelling older adults' level walking following perturbation training. *J
Biomech* 46: 2463–2468, 2013. doi: 10.1016/j.jbiomech.2013.07.025.

67.   **Okubo Y**, **Brodie MA**, **Sturnieks DL**, **Hicks C**, **Carter H**, **Toson B**, **Lord SR**. Exposure to trips and slips with
increasing unpredictability while walking can improve balance recovery responses with minimum predictive gait
alterations. *PLoS One* 13: e0202913, 2018. doi: 10.1371/journal.pone.0202913.

68.   **Wang Y**, **Wang S**, **Bolton R**, **Kaur T**, **Bhatt T**. Effects of task-specific obstacle-induced trip-perturbation training:
proactive and reactive adaptation to reduce fall-risk in community-dwelling older adults. *Aging Clin Exp Res* 32: 893–
905, 2020. doi: 10.1007/s40520-019-01268-6.

69.   **Wang T-Y**, **Bhatt T**, **Yang F**, **Pai Y-C**. Adaptive control reduces trip-induced forward gait instability among young
799   adults. *J Biomech* 45: 1169–1175, 2012. doi: 10.1016/j.jbiomech.2012.02.001.

800   70.   **Roeles S**, **Rowe PJ**, **Bruijn SM**, **Childs CR**, **Tarfali GD**, **Steenbrink F**, **Pijnappels M**. Gait stability in response to
801   platform, belt, and sensory perturbations in young and older adults. *Med Biol Eng Comput* 56: 2325–2335, 2018. doi:
802   10.1007/s11517-018-1855-7.

803   71.   **Kuo AD**. The relative roles of feedforward and feedback in the control of rhythmic movements. *Motor Control* 6: 129–
804   145, 2002. doi: 10.1123/mcj.6.2.129.

805   72.   **Bohm S**, **Mademli L**, **Mersmann F**, **Arampatzis A**. Predictive and Reactive Locomotor Adaptability in Healthy Elderly:
806   A Systematic Review and Meta-Analysis. *Sports Med* 45: 1759–1777, 2015. doi: 10.1007/s40279-015-0413-9.

807   73.   **Patla AE**. Strategies for dynamic stability during adaptive human locomotion. *IEEE Eng Med Biol Mag* 22: 48–52, 2003.
808   doi: 10.1109/memb.2003.1195695.

809   74.   **Wu M**, **Brown G**, **Gordon KE**. Control of locomotor stability in stabilizing and destabilizing environments. *Gait
810   Posture* 55: 191–198, 2017. doi: 10.1016/j.gaitpost.2017.04.021.

811   75.   **Lee A**, **Bhatt T**, **Pai Y-C**. Generalization of treadmill perturbation to overground slip during gait: Effect of different
812   perturbation distances on slip recovery. *J Biomech* 49: 149–154, 2016. doi: 10.1016/j.jbiomech.2015.11.021.

813   76.   **Madehkhaksar F**, **Klenk J**, **Sczuka K**, **Gordt K**, **Melzer I**, **Schwenk M**. The effects of unexpected mechanical
814   perturbations during treadmill walking on spatiotemporal gait parameters, and the dynamic stability measures by which
815   to quantify postural response. *PLoS One* 13: e0195902, 2018. doi: 10.1371/journal.pone.0195902.

816   77.   **Martelli D**, **Vashista V**, **Micera S**, **Agrawal SK**. Direction-Dependent Adaptation of Dynamic Gait Stability Following
817   Waist-Pull Perturbations. *IEEE Trans Neural Syst Rehabil Eng* 24: 1304–1313, 2016. doi:
818   10.1109/TNSRE.2015.2500100.

819   78.   **Bauby CE**, **Kuo AD**. Active control of lateral balance in human walking [Online]. *J Biomech* 33: 1433–1440, 2000.
820   https://www.ncbi.nlm.nih.gov/pubmed/10940402.

821   79.   **Rankin BL**, **Buffo SK**, **Dean JC**. A neuromechanical strategy for mediolateral foot placement in walking humans. *J
822   Neurophysiol* 112: 374–383, 2014. doi: 10.1152/jn.00138.2014.

823   80.   **Yoo D**, **Seo K-H**, **Lee B-C**. The effect of the most common gait perturbations on the compensatory limb's ankle, knee,
824   and hip moments during the first stepping response. *Gait Posture* 71: 98–104, 2019. doi: 10.1016/j.gaitpost.2019.04.013.

825   81.   **Watson F**, **Fino PC**, **Thornton M**, **Heracleous C**, **Loureiro R**, **Leong JJH**. Use of the margin of stability to quantify
826   stability in pathologic gait - a qualitative systematic review. *BMC Musculoskelet Disord* 22: 597, 2021. doi:
827   10.1186/s12891-021-04466-4.

828   82.   **Süptitz F**, **Karamanidis K**, **Moreno Catalá M**, **Brüggemann G-P**. Symmetry and reproducibility of the components of
829   dynamic stability in young adults at different walking velocities on the treadmill. *J Electromyogr Kinesiol* 22: 301–307,
830   2012. doi: 10.1016/j.jelekin.2011.12.007.

831   83.   **Madigan ML**, **Lloyd EM**. Age and stepping limb performance differences during a single-step recovery from a forward
832   fall. *J Gerontol A Biol Sci Med Sci* 60: 481–485, 2005. doi: 10.1093/gerona/60.4.481.

833   84.   **Farrens A**, **Lilley M**, **Sergi F**. Training Propulsion via Acceleration of the Trailing Limb. *IEEE Trans Neural Syst
834   Rehabil Eng* PP, 2020. doi: 10.1109/TNSRE.2020.3032094.

835   85.   **Shirazi SY**, **Huang HJ**. Differential Theta-Band Signatures of the Anterior Cingulate and Motor Cortices During Seated
836   Locomotor Perturbations. *IEEE Trans Neural Syst Rehabil Eng* 29: 468–477, 2021. doi: 10.1109/TNSRE.2021.3057054.

837   86.   **Wagner J**, **Makeig S**, **Gola M**, **Neuper C**, **Müller-Putz G**. Distinct β Band Oscillatory Networks Subserving Motor and
838   Cognitive Control during Gait Adaptation. *J Neurosci* 36: 2212–2226, 2016. doi: 10.1523/JNEUROSCI.3543-15.2016.

839   87.   **Peterson SM**, **Ferris DP**. Group-level cortical and muscular connectivity during perturbations to walking and standing
840   balance. *Neuroimage* 198: 93–103, 2019. doi: 10.1016/j.neuroimage.2019.05.038.

Fig. 1. Schematic of the perturbations and experimental protocol. **(A)** Perturbation onset occurred at right leg mid-stance (black

leg in red outline). There were 4 perturbation directions, and 2 sizes for the anterior-posterior perturbations, for a total of six

perturbation conditions. The red arrows are the initial relative surface displacements, and the faded red arrows are the

displacements to return to the unperturbed state. **(B)** The left belt speed was fixed at 1.0 m/s (blue). For the Stick0.2 and Stick0.4

perturbations, the right belt speed (red) decelerated to 0.8 m/s and 0.6 m/s, respectively, and then returned to the tied belt speed.

For the Slip0.2 and Slip0.4 perturbations, the right belt speed accelerated to 1.2 m/s and 1.4 m/s, respectively, and then returned

to the tied belt speed. For the L1 and M1 perturbations, the treadmill shifted 1 cm laterally and medially, respectively, and then

returned to neutral. The perturbations were ~400 ms in duration. Gait events: right heel strike (rhs), left toe off (lto), left heel

strike (lhs), and right toe off (rto). **(C)** Blocks of the experimental protocol. Each trial started with a 2-minute unperturbed

walking block (pre), followed by a 4-minute perturbed walking block (perturbation), and completed with another 2-minute

unperturbed walking block (post). An unperturbed catch stride occurred randomly 1 out of every 5 strides during the perturbation

block.

Fig. 2. Anterior-posterior and medial-lateral metrics of gait stability at left heel strike and left toe off. **(A)** The anterior-posterior

base of support (BOS_{ap}, orange arrow) was the anterior-posterior distance of the toe marker of the leading leg relative to the toe

marker of the trailing leg. The anterior-posterior extrapolated center of mass ($XCOM_{ap}$, green arrow) was the anterior-posterior

distance of XCOM relative to the toe marker of the trailing leg. The anterior-posterior margin of stability (MOS_{ap}, pink arrow)

was the BOS_{ap} - $XCOM_{ap}$. **(B)** The medial-lateral base of support (BOS_{ml}) was the medial-lateral distance of the ankle marker of

the leading leg relative to the ankle marker of the trailing leg. The medial-lateral extrapolated center of mass ($XCOM_{ml}$) was the

medial-lateral distance of XCOM relative to the ankle marker of the trailing leg. The medial-lateral margin of stability (MOS_{ml})

was the BOS_{ml} - $XCOM_{ml}$.

Fig. 3. Trajectories of the gait stability metrics (MOS, pink; BOS, orange, and XCOM, green) for specific strides of a

representative subject for the Stick0.4, Slip0.4, L1, and M1 perturbations (black dotted line: the 2^{nd} to last stride of pre; colored

thick line: the 2^{nd} stride of early perturbed; colored thin line: the 2^{nd} to last stride of late perturbed; and black dashed line: the 2^{nd}

to last stride of post). The $MOS_{ap, lhs}$, $BOS_{ap, lhs}$, and $XCOM_{ap, lhs}$ of the Stick0.4 and Slip0.4 perturbations, and $MOS_{ml, lhs}$, $BOS_{ml,}$

$_{lhs}$, and $XCOM_{ml, lhs}$ of the L1 and M1 perturbations deviated from pre at early perturbed, then MOS_{lhs} and BOS_{lhs} generally

trended to the pre trajectory by late perturbed and returned to the pre trajectory by post, shown in the exploded view in the inset

circles. The $MOS_{ap, lto}$ of the Stick0.4 perturbation, $BOS_{ap, lto}$ and $XCOM_{ap, lto}$ of the Slip0.4 perturbation, and $BOS_{ml, lto}$ and

$XCOM_{ml, lto}$ of the L1 perturbation deviated from pre at early perturbed, then BOS_{lto} and $XCOM_{lto}$ trended to the pre trajectory by

late perturbed and returned to the pre trajectory by post, shown in the exploded view in the inset circles. Gait events: right heel

strike (rhs), left toe off (lto), left heel strike (lhs), and right toe off (rto). Dark grey area: double support phase. Light grey area:

single support phase. The plots show the MOS and BOS during double support and XCOM for the entire gait cycle.

Fig. 4. Group-averaged (mean and standard deviation indicated by thick horizontal lines and one-sided error bar, n = 10)

anterior-posterior and medial-lateral MOS (pink), BOS (orange), and XCOM (green) at left heel strike and left toe off for pre,

early perturbed, late perturbed, early post, and late post during the anterior-posterior perturbations and the corresponding

footprints with metrics at pre, early perturbed, and late perturbed. Thick arrows indicate significant changes from the previous

phase. In the group-averaged plots, circles are individual subjects. **(A)** Subjects were destabilized at early perturbed then adapted

by late perturbed during the Stick0.4 perturbation. **(B)** Subjects were destabilized and adapted during the Slip0.4 perturbation. *:

$p < 0.05$ and color-coded for the specific metrics with the significant differences between phases indicated by the brackets.

Fig. 5. Group-averaged (mean and standard deviation indicated by thick horizontal lines and one-sided error bar, n = 10)

anterior-posterior and medial-lateral MOS (pink), BOS (orange), and XCOM (green) at left heel strike and left toe off for pre,

early perturbed, late perturbed, early post, and late post during the medial-lateral perturbations and the corresponding footprints

with metrics at pre, early perturbed, and late perturbed. Thick arrows indicate significant changes from the previous phase. In the

group-averaged plots, circles are individual subjects. **(A)** Subjects adapted to the L1 perturbation after initially being disrupted.

(B) Subjects adapted to the M1 perturbation after initially being disrupted. *: $p < 0.05$ and color-coded for the specific metrics

with the significant differences between phases indicated by the brackets.

Fig. 6. Group-averaged (mean and standard deviation, n = 10) $|\Delta BOS_{lhs}|$ (orange) compared with $|\Delta XCOM_{lhs}|$ (green) during

disruption (pre to early perturbed) and adaptation (early to late perturbed). Gray lines between bars are individual subjects. **(A)**

During disruption, BOS_{lhs} was disrupted more by the Stick0.4 perturbation, $XCOM_{lhs}$ was disrupted more by the Slip0.4 and M1

perturbations. **(B)** During adaptation, $BOS_{ap, lhs}$ adapted more in the L1 perturbation, $XCOM_{ap, lhs}$ adapted more in the Slip0.4 and

M1 perturbations, $BOS_{ml, lhs}$ adapted more in the L1 perturbation. *: $p < 0.05$ and significant differences between $|\Delta BOS_{lhs}|$ and

$|\Delta XCOM_{lhs}|$.

Fig. 7. Group-averaged (mean and standard deviation indicated by thick horizontal lines and one-sided error bar, n = 10)

anterior-posterior and medial-lateral MOS (pink), BOS (orange), and XCOM (green) at left heel strike and left toe off for pre,

early catch, late catch for the Stick0.4, Slip0.4, L1, and M1 perturbations. Circles are individual subjects. **(A)** Stick0.4

perturbation, **(B)** Stick0.4 perturbation, **(C)** L1 perturbation **(D)** M1 perturbation induced anticipatory responses at early catch,

but subjects did not use more feedforward strategies by late catch. *: $p < 0.05$ and color-coded for the specific metrics with the

significant differences between phases indicated by the brackets.

Fig. 8. Group-averaged (mean and standard deviation, n = 10) ΔMOS_{lhs} (pink), ΔBOS_{lhs} (orange), $\Delta XCOM_{lhs}$ (green) during

disruption (early perturbed minus pre) and adaptation (late minus early perturbed) for the Stick0.4 (black outline), Slip0.4 (gray

outline), L1 (white fill), and M1 (gray fill) perturbations. Circles are individual subjects. **(A)** Anterior-posterior metrics at left heel strike were more sensitive to the anterior-posterior perturbations during disruption but not adaptation. **(B)** $MOS_{ml, lhs}$ and $XCOM_{ml, lhs}$ were more sensitive to the medial-lateral perturbations during disruption but not adaptation. * with thick brackets: $p < 0.05$ and significant same signed differences between two perturbation conditions, * with thin brackets: $p < 0.05$ and significant opposite signed differences between two perturbation conditions.

Fig. 9. Group-averaged (mean and standard deviation, n = 10) $|\Delta MOS_{lhs}|$ (pink), $|\Delta BOS_{lhs}|$ (orange), $|\Delta XCOM_{lhs}|$ (green) during disruption (pre to early perturbed) and adaptation (early to late perturbed) for the anterior-posterior perturbations. Dark bars indicate the regular size perturbations (Stick0.4 and Slip 0.4) and light bars indicate the half-size perturbations (Stick0.2 and Slip 0.2). Colored lines between bars are individual subjects. **(A)** Anterior-posterior metrics at left heel strike scaled with perturbation size during disruption but not adaptation. **(B)** Medial-lateral metrics at left heel strike did not scale with perturbation size during disruption and adaptation. +: $p < 0.05$, perturbation size effect (perturbation condition effect was not reported). #: $p < 0.05$, interaction effect between perturbation size (regular, 0.4; half-size, 0.2) and perturbation condition (stick, slip). *: $p < 0.025$, significant differences between the regular and half-size perturbations of same condition.

A

Perturbation Onset

Anterior
(Stick0.4, Stick0.2)

Posterior
(Slip0.4, Slip0.2)

Lateral shift
(L1)

Medial shift
(M1)

Belt speed

B

Anterior-posterior Perturbations

Belt Speed (m/s)

slip0.4
slip0.2
stick0.2
stick0.4

Left belt
Right belt
Perturbation onset/termination

~400 ms

Gait Event

rhs   lto   lhs   rto   rhs

Medial-lateral Perturbations

Treadmill Shift (m)

L1

M1

Lateral direction

Medial direction

neutral

~400 ms

Gait Event

rhs   lto   lhs   rto   rhs

C

Trial

| Pre block 2 minutes | Perturbation block 4 minutes | Post block 2 minutes |

Stride type

UUU...UUU   PPCPP PPPPC PPPCP ... PCPPP   UUU...UUU

U = Unperturbed   P = Perturbed   C = Catch, no-perturbation, (random, 1 in 5)

Perturbed Strides for Anterior-posterior Perturbations

Perturbed Strides for Medial-lateral Perturbations

Catch Strides

A Anterior-posterior Metrics

$\triangle MOS_{ap,\ lhs}$ $\triangle BOS_{ap,\ lhs}$ $\triangle XCOM_{ap,\ lhs}$

disruption: early perturbed - pre

early perturbed > pre

closer to 0: less sensitive to perturbation directions

early perturbed < pre

adaptation: late - early perturbed

late > early perturbed

closer to 0: less sensitive to perturbation directions

late < early perturbed

Stick 0.4   Slip 0.4   L1   M1

B Medial-lateral Metrics

$\triangle MOS_{ml,\ lhs}$ $\triangle BOS_{ml,\ lhs}$ $\triangle XCOM_{ml,\ lhs}$

disruption: early perturbed - pre

adaptation: late - early perturbed

Stick 0.4   Slip 0.4   L1   M1

A Anterior-posterior Metrics

$|\triangle MOS_{ap,\,lhs}|$ $|\triangle BOS_{ap,\,lhs}|$ $|\triangle XCOM_{ap,\,lhs}|$

disruption: pre to early perturbed

adaptation: early to late perturbed

B Medial-lateral Metrics

$|\triangle MOS_{ml,\,lhs}|$ $|\triangle BOS_{ml,\,lhs}|$ $|\triangle XCOM_{ml,\,lhs}|$

disruption: pre to early perturbed

adaptation: early to late perturbed

Small Directional Treadmill Perturbations Induce Differential Gait Stability Adaptation

perturbation at every right mid-stance

M1 (medial shift)

Stick (belt deceleration)

Slip (belt acceleration)

L1 (lateral shift)

MOS: margin of stability — baseline

BOS: base of support — initial perturbations

XCOM: extrapolated center of mass — familiar with perturbations

Questions

			Stick 0.4			Slip 0.4			L1			M1		
Are these small perturbations **disruptive**? (baseline to initial)	ap		−	−	−		−	+	+	/	−	−		−	/	+
	ml		+	+	/		/	+	+		+	+	/		−	−	+
Do subjects **adapt** to small perturbations? (initial to familiar)	ap		+	+	/		/	/	/		+	+	/		+	/	−
	ml		/	/	/		/	/	/		−	−	/		/	/	/
Which component of MOS contributed more during **adaptation**? (initial to familiar)				/			$\|\Delta BOS_{ap}\|$ < $\|\Delta XCOM_{ap}\|$			$\|\Delta BOS_{ml}\|$ > $\|\Delta XCOM_{ml}\|$			$\|\Delta BOS_{ap}\|$ < $\|\Delta XCOM_{ap}\|$		

ap: anterior-posterior   ml: medial-lateral   **+** increase   **−** decrease   / no significant difference

CONCLUSION

Small magnitude perturbations applied on a stride-by-stride basis disrupt gait stability and induce direction dependent reactive balance strategies (foot placement or extrapolated center of mass control) during adaptation.