Title

The gene expression profile of the song control nucleus HVC shows sex specificity, hormone responsiveness, and species specificity among songbirds

Author affiliation

Meng-Ching Ko[*] (ORCID 0000-0002-2234-9380)

Carolina Frankl-Vilches[*]

Antje Bakker[*]

Manfred Gahr[*]

[*]Department of Behavioural Neurobiology, Max Planck Institute for Ornithology, Eberhard-Gwinner Straße 6a, 82319 Seewiesen, Germany

Corresponding author

Meng-Ching Ko

E-mail: mcko@orn.mpg.de

Keywords

Sex differences; Songbirds; Gene expression; Brain; HVC; Testosterone

Abstract

Singing occurs in songbirds of both sexes, but some species show typical degrees of sex-specific performance. We studied the transcriptional sex differences in the HVC, a brain nucleus critical for song pattern generation, of the forest weaver (*Ploceus bicolor*), the blue-capped cordon-bleu (*Uraeginthus cyanocephalus*), and the canary (*Serinus canaria*), which are species that show low, medium, and high levels of sex-specific singing, respectively. We observed persistent sex differences in gene expression levels regardless of the species-specific sexual singing phenotypes. We further studied the HVC transcriptomes of defined phenotypes of canary, known for its testosterone-sensitive seasonal singing. By studying both sexes of canaries during both breeding and nonbreeding seasons, nonbreeding canaries treated with testosterone, and spontaneously singing females, we found that the circulating androgen levels and sex were the predominant variables associated with the variations in the HVC transcriptomes. The comparison of natural singing with testosterone-induced singing in canaries of the same sex revealed considerable differences in the HVC transcriptomes. Strong transcriptional changes in the HVC were detected during the transition from nonsinging to singing in canaries of both sexes. Although the sex-specific genes of singing females shared little resemblance with those of males, our analysis showed potential functional convergences. Thus, male and female songbirds achieve comparable singing behaviours with sex-specific transcriptomes.

Introduction

Most of the genomes of male and female individuals of the same species are the same but stark sex differences in physiological, phenotypical, or behavioural traits between the sexes are common and widespread in the animal kingdom. The songbird clade (a suborder of the perching birds) consists of more than 4000 extant avian species, and these exhibit a great diversity of sex differences in singing behaviour. Among domesticated canaries (*Serinus canaria*), males are known for their singing behaviour and have been selected for their sophisticated songs for centuries, whereas female canaries seldom sing (Hartley et al., 1997; Herrick and Harris, 1957; Ko et al., 2020; Pesch and Güttinger, 1985; Shoemaker, 1939; Vallet et al., 1996). Such substantial sex differences in singing behaviour are commonly found in the majority of northern temperate songbird species, even though females of many tropical and southern temperate species sing regularly, and their songs play an important role in inter-sexual communication (Hall et al., 2015; Price, 2019; Price et al., 2009; Slater and Mann, 2004). For example, female blue-capped cordon-bleus (*Uraeginthus cyanocephalus*) use their song as advertising signals and address their songs to their mates (Immelmann, 1968; Ota et al., 2018), although the female songs appear shorter and less complex than those of males (Geberzahn and Gahr, 2011). In another tropical songbird species, forest weavers (*Ploceus bicolor*), males and females develop their songs during pair binding and eventually learn to sing identical duets, which they use to defend their territories (Wickler and Seibt, 1980).

Although the extent of sex differences in singing behaviour varies greatly in the songbird clade (Ball, 2016), the song quality and occurrence in males and females of many songbird species are dependent on testosterone (Dittrich et al., 2014; Fusani et al., 2003; Ko et al., 2020; Leitner et al., 2001b; Nottebohm et al., 1987; Voigt and Leitner, 2008). In canaries that breed seasonally, the optimal breeding conditions are tightly associated with an increase in the day length, which initiates gonadal growth and testosterone production as well as various types of breeding activity, including singing (Leitner et al., 2001b; Nottebohm et al., 1987; Voigt and Leitner, 2008). The length and syllable repetition rate of songs during the breeding season are greater than those of songs in nonbreeding seasons (Leitner et al., 2001a; Leitner et al., 2001b; Voigt and Leitner, 2008). Castrated male canaries sing shorter songs than sham-operated males, and the subcutaneous implantation of testosterone results in the recovery of singing performance (Heid et al., 1985). The local administration of testosterone into the preoptic brain region increases the male canary singing rate by increasing motivation (Alward et al., 2013). Similarly, the systemic administration of testosterone reliably and repeatedly induces

singing behaviour in otherwise nonsinging female canaries (Fusani et al., 2003; Herrick and Harris, 1957; Leonard, 1939; Madison et al., 2015; Nottebohm, 1980; Shoemaker, 1939; Vellema et al., 2019b). Although female canaries rarely exhibit spontaneous singing (Hartley et al., 1997; Herrick and Harris, 1957; Ko et al., 2020; Pesch and Güttinger, 1985; Shoemaker, 1939; Vallet et al., 1996), its occurrence appears to be associated with the plasma androgen levels (Ko et al., 2020).

A set of interconnected neural circuits called the song control system controls the production and learning of singing behaviour (Nottebohm et al., 1976; Wild, 2004). The premotor nucleus HVC (used as the proper name), which is a sensorimotor integration centre in the song control system, is involved in the frequency and temporal modulation of songs in male and female songbirds (Hahnloser et al., 2002; Hoffmann et al., 2019) and in the sexual preferences regarding conspecific song displays (Brenowitz, 1991; Del Negro et al., 1998). The HVC is the only brain nucleus within the song control system that expresses receptors for both androgens and estrogens (Frankl-Vilches and Gahr, 2018; Gahr, 2001). Intriguingly, the anatomical properties of the HVC, such as volume (Figure 1), neuron number, and dendrite complexity, are male-biased (greater, higher and more complex, respectively, in males than in females) in all songbird species that have been examined, irrespective of the existence of sex differences in singing behaviour (Brenowitz et al., 1985; Gahr et al., 2008; Gahr et al., 1998; Gurney and Konishi, 1980; Hall et al., 2010; Lobato et al., 2015; MacDougall-Shackleton and Ball, 1999; Nixdorf et al., 1989; Nottebohm and Arnold, 1976; Schwabl et al., 2015). Testosterone treatment increases the delineable volume of the HVC in both male and female canaries (Fusani et al., 2003; Madison et al., 2015; Nottebohm, 1980) and many other species (Bernard and Ball, 1997; Dittrich et al., 2014; Gulledge and Deviche, 1999; Smith et al., 1997; Van Meir et al., 2004). Nevertheless, the HVC volume of female canaries implanted with testosterone remains markedly smaller than that of males (Nottebohm, 1980) (Figure 1). Testosterone clearly regulates singing behaviour and the anatomy of the HVC in both male and female songbirds. However, an intrinsic limit to the alterations induced by testosterone appears to exist, and this limit prevents female songbirds from reaching the same levels of "maleness" in terms of song characteristics and HVC anatomy.

How are such sex-specific differences in singing and HVC anatomy achieved in different species? We hypothesize the existence of a fundamental difference in gene regulation and expression between male and female conspecific songbirds. Based on this hypothesis, sex differences in the HVC transcriptomes of male and female conspecifics should always be observable, even though both sexes share an almost identical genome.

The female-specific W chromosome harbours 30-50 genes, whereas one and two copies

of the Z chromosome (harbouring approximately 1,080 genes) are present in female and

male birds, respectively (Frankl-Vilches et al., 2015; Sayers et al., 2020; Smeds et al.,

2015). We further hypothesize that such intrinsic differences cannot be overwritten by

manipulating the testosterone level in females. To test our hypotheses, we compared the

transcriptomes of HVCs microdissected from three songbird species—the forest weaver,

the blue-capped cordon-bleu, and the canary. These species represent three categories of

sex differences in singing behaviour (low, medium, and high). Furthermore, we compared

male and female canaries with and without testosterone implantation to evaluate the

effects of testosterone (Supplementary Table 1). The results revealed sex differences in

the HVC transcriptome between- and within-species comparisons. Our results indicate

that although the extent of sex-biased gene expression is context-dependent, it is

persistent regardless of hormonal manipulation, behavioural phenotypes, and distinct

genetic backgrounds.

Results and discussion

The species identity distinguishes its HVC transcriptomes

We quantified the total mRNA from the HVC of male and female birds belonging to three

songbird species (forest weavers, blue-capped cordon-bleus, and canaries) during the

breeding season (Supplementary Table 1). For intraspecies sex comparisons, we included

male and female canaries during the nonbreeding seasons (both nonsinging), and

subgroups were treated with testosterone to induce singing behaviour in both sexes

(Supplementary Figure 1). In addition, we obtained a rare group of nonbreeding female

canaries that sang spontaneously without exogenous testosterone manipulation.

However, their plasma androgen levels were intrinsically higher than those of

nonbreeding and nonsinging females (Ko et al., 2020). To visualize the similarities in the

gene expression patterns among and within the three species, we calculated Spearman's

rank correlation coefficient σ for 12,360 gene expression levels and calculated the

distance matrix based on the coefficient σ. The resulting cladogram indicated that the HVC

transcriptomes of the eleven groups were clustered primarily by phylogenetic

relatedness (Figure 1B and Supplementary Figure 2). We obtained a similar result with

the transcriptomes of another tissue, the entopallium (Supplementary Figure 3), which is

an area of the avian visual system that is functionally equivalent to the primary visual

cortex of mammals. The within-species comparison of the canary HVC showed that the

individual transcriptomes clustered well by sex, with the exception of the nonbreeding

canaries implanted with testosterone (Figure 1B and Supplementary Figure 2). The
subclusters of forest weaver and cordon-bleu HVC transcriptomes were less sexually
differential than those of canaries, i.e., male and female birds were intermittent within the
subcluster of the two species (Supplementary Figure 2 and Supplementary Figure 3).

Figure 1. Species, sex, and plasma androgen level are the major determinants of the HVC gene expression patterns in songbirds.

A, HVC volume of birds used in this study. Forest weavers: males 0.528 mm^3 (mean), females 0.357 mm^3. Mann-Whitney Test, U = 7, P value = 0.0931. Cordon-bleus: males 0.245 mm^3, females 0.136 mm^3. Mann-Whitney Test, U = 0, P value = 0.00217. Breeding canaries: singing males 0.439 mm^3, nonsinging females 0.116 mm^3. Mann-Whitney Test, U = 0, P value = 0.00217. Nonbreeding canaries: nonsinging males 0.228 mm^3, nonsinging females 0.103 mm^3, singing females 0.176 mm^3. Mann-Whitney Test (nonsinging males vs. nonsinging females), U = 0, P value = 0.00492. Mann-Whitney Test (nonsinging males vs. singing females), U = 8, P value = 0.127. Testosterone-implanted canaries: males 0.478 mm^3, females 0.253 mm^3. Mann-Whitney Test, U = 0, P value = 0.0238. * P value < 0.05; ** P value < 0.01. Each colour-coded dot indicates the measurement from one bird. The boxes indicate the 25th/50th/75th percentiles (bottom/middle/top bar), and the extent of the whiskers indicates the most

extreme values that are within 1.5 times the IQR (interquartile range) of the hinge. T: testosterone; music note: presence of singing behaviour. **B**, Hierarchical clustering showing that phylogenetic relatedness accounts for the most variation in the HVC transcriptomes of females and males of the forest weaver, cordon-bleu, and canary. Among the seven canary groups, the testosterone-treated animals were the least similar to the untreated canaries. **C**, A PCA of the HVC transcriptomes of the seven canary groups distinguished male birds (PC2 < 0) from female birds (PC2 > 0). Each point is colour-coded by group (see B) and represents an HVC sample.

Sex-biased gene expression in the HVC of songbirds

We defined the degree of sex differences in HVC gene expression levels as the number of sex-biased genes whose expression levels showed significant differences between males and females of the same species (male-biased genes showed higher expression levels in males than in females, $\log_2$ (fold change) ≥ 0.5; female-biased genes presented higher expression levels in females than in males, $\log_2$ (fold change) ≤ -0.5). For canary, we quantified the sex differences from five distinct male-to-female comparisons (Figure 2A, comparisons 3 to 7).

Interestingly, we found persistent sex differences in HVC gene expression levels from all the comparisons, regardless of the degree of sex differences in singing behaviour (Figure 2A). The majority (> 90%) of sex-biased genes were lowly biased ($0.5 \leq |\log_2$ (fold change)$| < 1$, Supplementary Figure 5). The comparison of forest weaver (the female birds can sing identical songs to the males; least differences in the HVC volumes between the sexes; Figure 1A) showed the least prominent sex differences in the HVC transcriptomes across all the comparisons performed in this study. Nevertheless, the extent of sex differences in gene expression levels was extensive (> 1,300 sex-biased genes, Figure 2A, comparison 1). The comparison showing the second-lowest degree of sex-differences was the comparison of nonbreeding male and nonbreeding female canaries (> 1,500 sex-biased genes, Figure 2A, comparison 7), both lacking singing behaviour, but HVC volume was male biased (Figure 1A). Testosterone implantation induced singing behaviour and increased the HVC volume of both male and female canaries, although a markedly smaller HVC volume continued to be observed in these females (Figure 1A and Supplementary Figure 1C). Such processes in females are referred to as "masculinization" of the female brain and behaviour (Arnold and Gorski, 1984; Wade, 2001). However, the gene expression levels of the testosterone-stimulated singing female canaries were very different from those of the male canaries administered the same treatment (> 5,000 sex-biased genes, Figure 2A, comparison 4). The comparisons showing the most striking sex differences were the two that included spontaneously singing female canaries. As

revealed in these comparisons, many genes showed markedly different regulation in spontaneously singing female canaries, as reflected by expression levels, compared with breeding singing male canaries (> 8,600 sex-biased genes, comparison 5) and nonbreeding nonsinging male canaries (> 7,500 sex-biased genes, comparison 6, Figure 2A).

Figure 2. Female- and male-biased sex differences in HVC gene expression levels in songbirds.

A, The bar graph summarizes the number of sex-biased genes in the HVC transcriptome identified from each male-to-female comparisons. The phenotypes of the groups being compared in each male-to-female comparison are listed at the bottom of the graph. **B,** The bar graph summarizes the number of autosomal and Z chromosomal genes classified as male- and female-biased genes in the HVC transcriptome based on each male-to-female comparison. Abbreviations: B, breeding; NB, nonbreeding; T, testosterone implantation. The music note indicates the presence or absence of singing behaviour; the asterisk (*)

indicates enrichment for Z chromosomal genes (adjusted $p < 0.05$, Fisher's exact test followed by the Bonferroni correction for multiple testing).

184  For each comparison, we decomposed the sex-biased genes into two classes,
185 autosomal genes and Z sex chromosomal genes, to assess whether the sex-biased genes
186 were mainly concentrated on the sex chromosomes. Annotation for W chromosomal
187 genes was unfortunately not available in our study. We used Fisher's exact test to evaluate
188 whether a particular chromosome was enriched in sex-biased genes. We observed
189 enrichments of Z chromosomal genes among the male-biased genes identified in most of
190 the comparisons (Figure 2B and Supplementary Table 3). This observation was not
191 surprising because males are the homogametic sex (have two copies of Z), whereas
192 females are the heterogametic sex (have one copy of Z and one copy of W) in birds, and
193 the dosage compensation for Z-linked genes is less complete in birds than that in
194 mammals (Itoh et al., 2007; Mank, 2013; Nätt et al., 2014; Uebbing et al., 2015; Wolf and
195 Bryk, 2011). However, one canary comparison (breeding singing males and nonbreeding
196 singing females, Figure 2B, comparison 5) was an exception (Supplementary Table 3). In
197 this comparison, the spontaneously singing female canaries expressed a high number of
198 Z chromosomal genes (260 genes, approximately 24% of all Z chromosomal genes) at
199 higher levels than the breeding singing male canaries, which eliminated the male
200 enrichment of Z chromosomal genes.

201  Sex-biased genes were not only Z chromosomal genes; in fact, sizeable numbers
202 of the autosomal genes were found to be sex-biased genes, particularly female-biased
203 genes (male-biased genes: >40%; female-biased genes: >96%, Figure 2B). Many studies
204 have reported sex differences in the gene expression levels in multiple tissues and animal
205 species (Frésard et al., 2013; Itoh et al., 2007; Nätt et al., 2014; Uebbing et al., 2015; Wolf
206 and Bryk, 2011; Yang et al., 2006). Gene expression levels are generally under tight
207 regulation, and perturbed expression levels might result in functional consequences. For
208 example, several genes, including transcription factors, modulate distinct gene sets
209 depending on their expression levels (Birchler et al., 2001; Doghman et al., 2013; Schulz,
210 2017).

211  We compared the male-biased genes in a pairwise manner across the
212 comparisons and found that the male-biased autosomal genes showed low similarity to
213 each other (Supplementary Figure 6A), whereas the male-biased Z chromosomal genes
214 showed some similarity across the groups (Supplementary Figure 6B). Pairwise
215 comparison of the female-biased genes across groups showed low similarity, and this
216 finding was obtained for both autosomal genes (Supplementary Figure 6C) and Z

chromosomal genes (Supplementary Figure 6D). Fisher's exact test showed that chromosome 2 (comparisons 1, 4, and, 5) and chromosome 3 (comparisons 1, 2, and 4) are hotspots for female-biased genes; the enrichments of these chromosomes were observed in four out of the seven pairwise comparisons (Supplementary Table 3).

Figure 3. Cross-species comparisons of sex-biased genes expressed in the HVC of birds with singing phenotypes shows high species specificity.

Venn diagrams of male-biased (**A**) and female-biased (**B**) autosomal and Z chromosomal genes. The numbers of sex-biased genes are indicated in parentheses. **C,** GO term enrichment analysis predicting the biological functions of sex-biased (autosomal and Z chromosomal) genes. The GO terms were categorized and are colour coded. Bonferroni-adjusted P values are shown by colour scales in the heatmap. Only five GO terms with the lowest Bonferroni-adjusted P values are shown for each set (see Supplementary Table 4 for the complete results). The phenotypes of the groups used for comparisons 1, 2, and 5 are listed at the bottom. Abbreviations: B, breeding; NB, nonbreeding. The music note indicates the presence or absence of singing behaviour

Further examination of the sex-biased genes (comparisons 1, 2, and 5) identified in the groups of naturally singing birds revealed that almost none of the male-biased autosomal genes (Figure 3A) and none of the female-biased Z chromosomal genes (Figure 3B) were shared across the three songbird species. In addition, relatively low numbers of male-biased Z chromosomal genes (Figure 3A) and female-biased autosomal genes (Figure 3B) were shared across the three songbird species. This observation indicates that only a small set of genes were regulated in the same manner (male- or female-biased)

between species even though all birds exhibit singing behaviours. Moreover, a Gene Ontology (GO) enrichment analysis suggested that male- and female-biased genes did not functionally converge to similar pathways in the three studied species (Figure 3C).

The plasma androgen levels substantially alter the HVC transcriptomes of canaries

With the aim of understanding the features that distinguish the HVC transcriptomes in within-species contexts, we performed a principal component analysis (PCA) of the HVC transcriptomic data from the seven canary groups to identify variables that would explain the most variation in the data. By calculating the correlation coefficient between the PCs and the variables (plasma testosterone levels (Supplementary Figure 1), HVC volume (Figure 1A), sex, and singing), we identified the variables that were highly correlated with the most important PCs. PC1 explained 32% of the data variance (Figure 1C and Supplementary Figure 4) and was strongly correlated with the blood plasma androgen concentrations and the presence of singing activity (plasma androgen level: Pearson's $r = -0.57$, Bonferroni-adjusted $p = 1.0 \times 10^{-4}$; singing: Pearson's $r = -0.51$, Bonferroni-adjusted $p = 0.001$, Supplementary Table 2). Sex and HVC volume were strongly correlated with PC2 (sex: Pearson's $r = -0.91$, Bonferroni-adjusted $p = 4.4 \times 10^{-16}$; HVC volume: Pearson's $r = -0.73$, Bonferroni-adjusted $p = 2.1 \times 10^{-7}$), which explained 19% of the variance in the data (Supplementary Table 2). Taken together, the hierarchical clustering and PCA results suggest that although phylogenetic relationships dominate the variation between songbird species, the circulating testosterone levels, the presence of singing activity, and sex identity dominate the HVC gene expression patterns within a single species.

The application of testosterone does not yield an HVC transcriptome that mimics that of natural singing canaries

To investigate the molecular mechanisms underlying the singing phenotype, we performed a differential gene expression analysis of the HVC transcriptomes of the two groups of singing canary females (nonbreeding spontaneously singing females and nonbreeding testosterone-stimulated singing females) against the nonsinging nonbreeding female canaries. Similarly, a differential gene expression analysis of the HVC transcriptomes of the two groups of singing male canaries (breeding singing males and nonbreeding testosterone-stimulated singing males) with nonsinging nonbreeding male canaries was performed. A substantial number of genes were differentially expressed between singing and nonsinging birds (Figure 4A, nonbreeding singing females: 4,125 genes; nonbreeding testosterone-stimulated singing females: 7,702 genes; breeding

264 singing males: 3,359 genes; nonbreeding testosterone-stimulated singing males: 5,106

265 genes). Approximately 65% of the differentially expressed genes found in the naturally

266 singing birds overlapped with those found in the testosterone-treated birds of the same

267 sex (male: 65%; female: 62%). However, both testosterone-treated groups had markedly

268 higher numbers of differentially expressed genes than the naturally singing groups of the

269 same sex (Figure 4A). Thus, although testosterone implantation induced singing in

270 females and males, most testosterone-responsive genes (female: 67%; male: 57%) might

271 not be necessary for singing behaviour *per se* but rather a response to nonphysiological

272 levels of testosterone (Supplementary Figure 1). Alternatively, mechanisms that lead to

273 the first song in life might be very different from those that reinduce singing in animals

274 that sang before (Vellema et al., 2019a). Because male canaries sing regularly starting

275 from approximately 50 days of age (Nottebohm et al., 1986) while most female canaries

276 never sing (Ko et al., 2020), the induction of singing in females by testosterone likely

277 activated genes related to first-time singing, whereas in males, this treatment activated

278 genes related to reinduced singing.

The majority of differentially expressed genes between singing and nonsinging
canaries are sex-specific

281 We examined whether the overlapping gene sets of the female canary signing groups

282 were similar to the overlapping gene sets of the singing males (Figure 4A). The Venn

283 diagram indicated that only approximately 25% (642 of 2569 genes) of the female-

284 specific expressed genes and 29% (642 of 2210 genes) of the male-specific expressed

285 genes were shared by both sexes of singing canaries.

The sex-specific genes show functional overlap

287 To understand the putative biological functions of the female-specific, male-specific, and

288 sex-shared expressed genes in the HVC, we performed a GO term enrichment analysis.

289 Interestingly, the results suggested that the female- and male-specific expressed genes

290 largely overlapped at the functional level (Figure 4B and Supplementary Table 5). GO

291 terms such as nervous system development (GO:0007399), neuron development

292 (GO:0048666), and intracellular signal transduction (GO:0035556) were shared among

293 the sexes. Female-specific GO terms were mainly related to cellular maintenance, such as

294 organonitrogen compound metabolic process (GO:1901564) and phospholipid metabolic

295 process (GO:0006644). In contrast, the male-specific GO terms were axon development

296 (GO:0061564), DNA replication (GO:0006260), cell migration (GO:0016477), and blood

vessel development (GO:0001568). In summary, although only approximately one-fourth to one-third of the sex-specific expressed genes are shared among the sexes in terms of their identities, most of the predicted functions of the sex-specific expressed genes are nevertheless sex-shared in singing canaries.

Figure 4. HVC transcriptomes of naturally singing canaries are dissimilar from testosterone-stimulated singing canaries.

Venn diagrams comparing (**A**) the two female singing groups, (**B**) the two male singing groups, and (**C**) the female- and male-biased genes resulting from comparisons A and B. Note that only approximately

20% of the genes were shared among the singing females and approximately 26% were shared among the singing males. The majority of these overlapping genes were not sex-shared; approximately 75% of female intersections were specific to female groups, whereas 71% of male intersections were specific to male groups. **D**, GO term enrichment analysis of female-specific, male-specific, and sex-shared expressed genes derived from comparisons A to C. Because many putative functions show similarities between male- and female-specific expressed genes, the results suggest functional intraspecies convergence based on sex-specific gene expression in the HVC. See Supplementary Table 5 for the full results of the GO term enrichment analysis.

Testosterone can be converted to 5α-dihydrotestosterone (5α-DHT) and 17β-
estradiol, which activate androgen receptor (AR) and estrogen receptors (ERα and ERβ),
respectively. Both AR (encoded by *AR*) and ERs (ERα encoded by *ESR1* and ERβ encoded
by *ESR2*) are transcription factors that play important roles in the transcription of
numerous genes (Bourdeau et al., 2004; Pihlajamaa et al., 2015; Takayama et al., 2007;
Wilson et al., 2016). Interestingly, *ESR1* was specifically expressed in males, whereas
ESR2 was female-specifically expressed in the HVC of canaries, which suggests that
estrogen receptor paralogues could provide finer-tuned mechanisms for sex-specific
regulation. Empirical results have shown that ERα and ERβ bind to the same estrogen
response element motifs (Zhao et al., 2010) and might functionally overlap in some
tissues. Moreover, ER paralogues show sex differences in expression levels and tissue
specificity (Zhang et al., 2017). In quail, the administration of an agonist specific to ERβ
on embryonic day 7 demasculinizes male sexual behaviour and midbrain nuclei
characteristics in Japanese quails, whereas an agonist specific to ERα does not exert this
effect (Court et al., 2020). The specific roles of each ER paralogue in the adult HVC of male
and female songbirds warrant further investigation. Until now, ERα but not ERβ was
expected to regulate the function of HVC neurons of adult songbirds in addition to the AR
(Frankl-Vilches and Gahr, 2017).

Conclusion

In this study, we investigated the sex differences in the gene expression patterns in the
HVC of three songbird species with different levels of sex-specific singing and several
different song-related phenotypes between male and female canaries. Our inter- and
intraspecies comparisons yielded large-scale transcriptional sex differences regardless of
singing behaviour. Instead, fundamental sex differences in gene expression levels were
found in the HVC, and these differences were highly species-specific. By leveraging
several experimental groups of canaries, we found that the plasma androgen levels and
sex were the major contributors to the variations in the HVC transcriptome. Although

testosterone reliably induced singing in both female and male canaries, testosterone treatment did not alter the transcriptome to imitate that of natural singing birds. Our results suggest that female and male canaries rely on different gene networks for singing behaviour, but the sex-specific gene networks might show functional convergence.

Materials and methods

Animals

Forest weaver (*Ploceus bicolor*) pairs were observed in their breeding territories in eastern South Africa; the songs of the pairs were recorded to ensure that both mates were singing. Subsequently, the animals were caught and sacrificed in accordance with permits issued by the local authorities (Chief Professional Officer for Research at the Natal Parks, Game and Fish Preservation Board, P. O. B. 662, Pietermaritzburg 3200). Blue-capped cordon-bleus (*Uraeginthus cyanocephalus*) and canaries (*Serinus canaria*) were bred at the animal facilities at the Max Planck Institute for Ornithology in Seewiesen, Germany. The procedures used for animal housing and welfare complied with the European directives for the protection of animals used for scientific purposes (2010/63/EU), and the protocols were approved by the Government of Upper Bavaria (AZ No. 55.2-1-54-2532-181-12). Adult canaries and blue-capped cordon-bleus (aged at least 1 year) were housed in pairs under long-day conditions (light:dark = 14:10 hours), and their breeding activities were monitored. The birds (male canaries and cordon-bleus) were sacrificed after singing activity was observed during the breeding season. Nonbreeding canaries were housed pairwise with a 9-hour light:15-hour dark schedule for at least 8 weeks before their singing activity was monitored (generally starting in late September). Nonbreeding male and female canaries were maintained alone in sound-attenuated boxes (70 × 50 × 50 cm), recorded continuously during the song monitoring phase (two weeks) and sacrificed after confirmation of no singing activity. Additional groups of male and female nonbreeding and nonsinging canaries were implanted with testosterone for 2 weeks and sacrificed after observation of singing activity. Water and food were available *ad libitum*. The sex was confirmed by PCR using the P2 and P8 primers for CHD genes (Griffiths et al., 1998) and by visual inspection of the reproduction system after sacrifice. We previously found that six nonbreeding female canaries exhibited singing behaviour during long-term monitoring, and their songs have been described in detail (Ko et al., 2020). In this study, we included the transcriptomes of these birds. All the birds were sacrificed via an overdose of isoflurane followed by decapitation, the body weights were recorded, and the brains were dissected, weighed, snap-frozen on dry ice and stored at -

365 80°C until further use. Supplementary Table 1 summarizes the information of the
366 experimental groups used in this study.

Testosterone implantation

A Silastic™ tube (Dow Corning; 1.47-mm inner diameter, 1.96-mm outer diameter, 0.23-
mm thickness) was cut to a length of 7 mm and loaded with testosterone (86500, Fluka)
as densely as possible. The two ends of the Silastic™ tube were sealed with silicone
elastomer (3140, Dow Corning). After closure, the implants were cleaned with 100%
ethanol to remove testosterone particles and then immersed in ethanol overnight in a
hood to ensure no leakage at either end. Implants with apparent dampness were
discarded. One day prior to the implantation, the implants were incubated in 0.1 M
phosphate buffered saline (PBS) overnight to enable the immediate release of
testosterone upon implantation (Rasika et al., 1994). We started implantation at
approximately 8:30 am (immediately after the light was turned on in the morning), which
resulted in a 20-minute interval between birds based on the scarification time. A small
incision was made on the back of the bird over the pectoral musculature, and one
testosterone implant was placed subcutaneously. The skin was closed by the application
of tissue glue. After 2 weeks, the animals were sacrificed. The testosterone implants were
checked, and this inspection revealed that the implants were all in place and were not
empty at the end of the experiments.

Measurement of the HVC volume

One series of 20-µm sections was mounted on RNase-free Superfrost slides, subjected to
Nissl staining with 0.1% thionin (Sigma-Aldrich), dehydrated, immersed in xylene and
cover-slipped with Roti-Histokitt II mounting medium (Roth). The HVC areas (typically 8-
10 slices) were measured with a Leica DM6000 B microscope connected to a computer-
based image-analysis system (IMAtec). All brains were coded such that the observers
were blind towards any additional information about the sections they measured during
the delineations. The volumes were derived from the summed area measurements
multiplied by the section thickness and the intersection distance.

Radioimmunoassay of plasma testosterone

394 Blood was sampled (< 150 µl) at the time sacrifice. All blood samples were taken between
395 8 and 11 am and were taken within 3 min after caught to avoid the effect of handling
396 (Wingfield et al., 1982). Blood samples were centrifuged (5,000 rpm, 10 min) to separate

the plasma from blood cells. Testosterone metabolites were measured with a radioimmunoassay using a commercial antiserum against testosterone (T3-125, Endocrine Sciences, Tarzana, USA) as previously described (Goymann et al., 2002). Standard curves and sample concentrations were calculated with Immunofit 3.0 (Beckman Inc. Fullerton, CA) using a four-parameter logistic curve fit and corrected for individual recoveries.

The testosterone concentrations were assayed in duplicate in five separate assays. The mean extraction efficiency for plasma testosterone was 85.0 ± 3.9% (mean ± SD, N = 42). The lower detection limits of the testosterone assays were 0.34, 0,35, 0.36, 0.38 and 0.35 pg per tube, and all the samples were above the detection limit. The intra-assay coefficients of variation of a chicken plasma pool were 8.7%, 3.4%, 12.8%, 1.9%, and 4.4%. The interassay coefficient of variation as determined by the variation in the chicken plasma pool between all the assays was 5.1%. Because the testosterone antibody used shows significant cross-reactions with 5α-dihydrotestosterone (44%), our measurement might include a fraction of 5α-DHT.

Microarray procedures and annotation

The birds were killed by an overdose of isoflurane, and their brains were snap-frozen on dry ice. The brains were sectioned sagittally into four series of 40-μm sections and two series of 20-μm sections with a cryostat (Jung CM3000 Leica). The 40-μm sections were mounted on glass slides for subsequent tissue dissection, whereas the thin sections were mounted on RNase-free Superfrost slides for Nissl staining. All sections were stored at -80°C until further processing. For total RNA extraction, the song control nucleus HVC and the visual area of the entopallium were dissected from the abovementioned 40-μm sections under a stereomicroscope (typically 24-32 slices for the HVC and 16-20 slices for the entopallium) and transferred into an Eppendorf tube containing 340 μl of RLT buffer mixture (containing DTT, Qiagen). This dissection procedure using rather thin sections reduces the contamination of HVC tissue with surrounding tissue. RNA was then extracted using the RNeasy® Micro Kit (Qiagen) with the optional DNase digest step. The RNA quality was assessed using the Agilent Model 2100 Bioanalyzer (Agilent Technologies), and the RNA concentrations were assessed using a Nanodrop 1000 spectrometer (Thermo Fisher Scientific). The RNA quality of all samples was good (RIN > 7). The purified total RNA samples (at least 100 ng per sample) were subsequently processed and hybridized using the Ambion WT Expression Kit and the Affymetrix WT Terminal Labelling and Controls Kit. The resulting cDNA was hybridized to the Custom

Affymetrix Gene Chip® MPIO-ZF1s520811 Exon Array, which has been used successfully and validated in cross-species hybridization studies (Dittrich et al., 2014; Frankl-Vilches et al., 2015). The 5.7 million male zebra finch-specific probes spotted on this array correspond to approximately 4,711,133 probe sets and hence to 25,816 transcripts published in public databases (NCBI and Ensembl). We annotated more than 90% of the transcripts to 12,729 human orthologous genes using several publicly available databases (Ensembl, GenBank, UniProt, and DAVID (Benson et al., 2005; Consortium, 2015; Flicek et al., 2014; Huang et al., 2008, 2009; Yates et al., 2016)) and commercial databases (El Dorado, Genomatix, Precigen Bioinformatics Germany GmbH (PBG), Munich, Germany). Hybridization was performed for 16 hours at 45°C and 60 rpm in a GeneChip Hybridization Oven 640 (Affymetrix). The arrays were washed, stained, and scanned using the Affymetrix GeneChip Fluidics Station 450 and the Affymetrix GeneChip scanner 3000 7G. The CEL files were generated using Affymetrix® GeneChip® Command Console® Software (AGCC), and for quality control, the success of individual hybridizations was assessed using Affymetrix® Expression Console™ software.

Differential expression was calculated using ChipInspector software version 21 (El Dorado Database version E28R1306 (Genomatix)). ChipInspector is a single probe-based analysis tool for microarray data that can show increased sensitivity compared with that obtained with conventional probe set-based analyses, such as robust multiarray analysis. ChipInspector consists of four steps: single probe-transcript annotation (ensuring up-to-date annotation), total intensity normalization, SAM (significance analysis of microarrays, adapted to single-probe handling) analysis (Tusher et al., 2001), and transcript identification based on significantly changed probes (Cohen et al., 2008). We set the delta threshold to 0 (to control the false positive rate) and used the groupwise exhaustive comparison tool in ChipInspector. The minimum coverage for each transcript was set to 10 significant probes, and the minimum expression difference was $|\log_2(\text{fold change})| \geq 0.5$. The significantly differentially expressed transcripts obtained were annotated to human orthologous genes as described above. For transcripts belonging to the same genes, the average expression was calculated if all transcripts were regulated in the same manner (e.g., all upregulated or all downregulated). If a gene contains both up- or downregulated transcripts, the transcripts that showed changes in expression in the minority direction were discarded (<40% of the total transcripts), and the average expression was instead calculated from the remaining transcripts (60% or more). Transcripts without human orthologous gene annotation were removed prior to subsequent analyses. We were cautious about possible cross-species bias in hybridization, and differential expression analyses were only performed between two

467   conspecific groups. The microarray data discussed in this publication have been
468   deposited in NCBI's Gene Expression Omnibus (Edgar et al., 2002) and are accessible
469   through the GEO Series accession number GSE83674.

Hierarchical clustering

The normalized gene expression levels (across all transcriptome samples) were
calculated using the "justRMA" function of the R package "affy" v1.66.0 (Gautier et al.,
2004), and the expression levels were further collapsed to gene levels using the
"collapseRows" functions of the R package "WGCNA" v1.69 (Langfelder and Horvath,
2008, 2012). The group-level expression levels were calculated by averaging the gene
expression levels. Spearman's ρ was calculated between samples (or groups) using the
"cor" function, and a Euclidean distance matrix was calculated (dist(1-cor)) and used for
hierarchical clustering analysis (method: complete) with the "hclust" function. This
computation was performed with R v4.0.2 (R_Core_Team, 2020) using the "dendextend"
v1.14.0 package for visualization (Galili, 2015).

Principal component analysis (PCA)

482   PCA (Ringner, 2008) was performed using the normalized and collapsed HVC gene
483   expression data (described in the hierarchical clustering section) from 41 canaries
484   (Supplementary Table 1). The computation was performed using the "pca" function of the
485   R package "pcaMethods" (v1.60.0 (Stacklies et al., 2007), method = svd). The data were
486   centred but not scaled because the expression data had already been normalized.

Fisher's exact testing for chromosomal enrichment

Fisher's exact test was used to test whether sex-biased genes were enriched on a
chromosome by comparing the gene lists of interest to the zebra finch annotation
(https://www.ncbi.nlm.nih.gov/genome/?term=zebra+finch). We used the "fisher.test"
function in R (R_Core_Team, 2020) with the alternative set to "greater" for enrichment.
The male- and female-biased genes identified from each comparison as well as the male-
specific, female-specific, and sex-shared genes were tested separately. The P values were
adjusted using the Bonferroni correction to account for multiple comparisons using
"p.adjust" in R (R_Core_Team, 2020).

Gene Ontology (GO) term enrichment analysis

We used ClueGO v2.2.4, an application built in the Cytoscape environment v3.3.0 (Bindea et al., 2009; Shannon et al., 2003), to predict the putative biological functions of the genes of interest. This software performs GO term enrichment hierarchical analyses and fuses GO terms with similar functions. The enrichment was determined by the right-sided test and corrected using the Bonferroni step-down method considering multiple comparisons.

Data accessibility

The microarray CEL files (GEO Series accession number GSE83674) are available on NCBI's Gene Expression Omnibus. The processed microarray data, plasma androgen levels, HVC volume measures, and all scripts used for analysis and visualization are available in GitHub (https://github.com/maggieMCKO/SongbirdSexDiff).

Authors' contributions

M-CK collected the nonbreeding testosterone-implanted canaries, analysed and visualized the transcriptomic data and was a major contributor to the writing of the manuscript. AB and M-CK performed the RNA extractions and microarray hybridizations. MG and M-CK isolated the tissues for microarray. MG collected the forest weavers and cordon-bleus. MG conceived the study and obtained financial support for the study. M-CK, CF-V and MG drafted the manuscript. All the authors read and approved the final manuscript.

Acknowledgements

We thank Dieter Schmidl for catching the forest weavers, Stefan Leitner for providing the breeding canaries, Roswitha Brighton and David Witkowski for maintaining the breeding colonies of the blue-capped cordon-bleus and the canaries, Doris Walcher for providing data from the breeding male and female canaries as well as the nonbreeding nonsinging male and female canaries, and Wolfgang Goymann and Monika Trappschuh for the RIA of the plasma testosterone. Furthermore, we express our appreciation to Maude Baldwin, Glenn Cockburn, Falk Dittrich, Lisa Gill, Vincent Van Meir, Keren Sadanandan, and Michiel Vellema for their comments on previous versions of this manuscript and to the International Max Planck Research School for Organismal Biology for the training and support provided.

Competing interests

The authors declare no competing financial interests.

Funding

This research was supported by Max-Planck-Gesellschaft to MG. The funder had no role in the study design, data collection and analysis, decision to publish, or preparation of the manuscript.

References

Alward, B.A., Balthazart, J., and Ball, G.F. (2013). Differential effects of global versus local testosterone on singing behavior and its underlying neural substrate. Proc Natl Acad Sci U S A *110*, 19573-19578.

Arnold, A.P., and Gorski, R.A. (1984). Gonadal steroid induction of structural sex differences in the central nervous system. Annu Rev Neurosci *7*, 413-442.

Ball, G.F. (2016). Species variation in the degree of sex differences in brain and behaviour related to birdsong: adaptations and constraints. Philos Trans R Soc Lond B Biol Sci *371*.

Benson, D.A., Karsch-Mizrachi, I., Lipman, D.J., Ostell, J., and Wheeler, D.L. (2005). GenBank. Nucleic Acids Research *33*, D34-D38.

Bernard, D.J., and Ball, G.F. (1997). Photoperiodic Condition Modulates the Effects of Testosterone on Song Control Nuclei Volumes in Male European Starlings. General and Comparative Endocrinology *105*, 276-283.

Bindea, G., Mlecnik, B., Hackl, H., Charoentong, P., Tosolini, M., Kirilovsky, A., Fridman, W.-H., Pagès, F., Trajanoski, Z., and Galon, J. (2009). ClueGO: a Cytoscape plug-in to decipher functionally grouped gene ontology and pathway annotation networks. Bioinformatics *25*, 1091-1093.

Birchler, J.A., Bhadra, U., Bhadra, M.P., and Auger, D.L. (2001). Dosage-Dependent Gene Regulation in Multicellular Eukaryotes: Implications for Dosage Compensation, Aneuploid Syndromes, and Quantitative Traits. Developmental Biology *234*, 275-288.

Bourdeau, V.r., Deschênes, J., Métivier, R.l., Nagai, Y., Nguyen, D., Bretschneider, N., Gannon, F., White, J.H., and Mader, S. (2004). Genome-Wide Identification of High-Affinity Estrogen Response Elements in Human and Mouse. Molecular Endocrinology *18*, 1411-1427.

Brenowitz, E. (1991). Altered perception of species-specific song by female birds after lesions of a forebrain nucleus. Science *251*, 303 - 305.

Brenowitz, E.A., Arnold, A.P., and Levin, R.N. (1985). Neural correlates of female song in tropical duetting birds. Brain Res *343*, 104-112.

Cohen, C.D., Lindenmeyer, M.T., Eichinger, F., Hahn, A., Seifert, M., Moll, A.G., Schmid, H., Kiss, E., Grone, E., Grone, H.J*., et al.* (2008). Improved elucidation of biological processes linked to diabetic nephropathy by single probe-based microarray data analysis. PLoS One *3*, e2937.

Consortium, T.U. (2015). UniProt: a hub for protein information. Nucleic Acids Research *43*, D204-D212.

Court, L., Vandries, L., Balthazart, J., and Cornil, C.A. (2020). Key role of estrogen
receptor beta in the organization of brain and behavior of the Japanese quail. Horm
Behav *125*, 104827.

Del Negro, C., Gahr, M., Leboucher, G., and Kreutzer, M. (1998). The selectivity of sexual
responses to song displays: effects of partial chemical lesion of the HVC in female
canaries. Behavioural Brain Research *96*, 151-159.

Dittrich, F., Ramenda, C., Grillitsch, D., Frankl-Vilches, C., Ko, M.C., Hertel, M., Goymann,
W., Ter Maat, A., and Gahr, M. (2014). Regulatory mechanisms of testosterone-
stimulated song in the sensorimotor nucleus HVC of female songbirds. BMC Neurosci *15*,
128.

Doghman, M., Figueiredo, B.C., Volante, M., Papotti, M., and Lalli, E. (2013). Integrative
analysis of SF-1 transcription factor dosage impact on genome-wide binding and gene
expression regulation. Nucleic Acids Research *41*, 8896-8907.

Edgar, R., Domrachev, M., and Lash, A. (2002). Gene Expression Omnibus: NCBI gene
expression and hybridization array data repository. Nucleic Acids Res *30*, 207 - 210.

Flicek, P., Amode, M.R., Barrell, D., Beal, K., Billis, K., Brent, S., Carvalho-Silva, D.,
Clapham, P., Coates, G., Fitzgerald, S*., et al.* (2014). Ensembl 2014. Nucleic Acids
Research *42*, D749-D755.

Frankl-Vilches, C., and Gahr, M. (2017). Androgen and estrogen sensitivity of bird song: a
comparative view on gene regulatory levels. Journal of Comparative Physiology A.

Frankl-Vilches, C., and Gahr, M. (2018). Androgen and estrogen sensitivity of bird song: a
comparative view on gene regulatory levels. Journal of Comparative Physiology A *204*,
113-126.

Frankl-Vilches, C., Kuhl, H., Werber, M., Klages, S., Kerick, M., Bakker, A., de Oliveira, E.H.,
Reusch, C., Capuano, F., Vowinckel, J*., et al.* (2015). Using the canary genome to decipher
the evolution of hormone-sensitive gene regulation in seasonal singing birds. Genome
Biol *16*, 19.

Frésard, L., Morisson, M., Brun, J.-M., Collin, A., Pain, B., Minvielle, F., and Pitel, F. (2013).
Epigenetics and phenotypic variability: some interesting insights from birds. Genetics
Selection Evolution *45*, 16.

Fusani, L., Metzdorf, R., Hutchison, J., and Gahr, M. (2003). Aromatase inhibition affects
testosterone-induced masculinization of song and the neural song system in female
canaries. J Neurobiol *54*, 370 - 379.

Gahr, M. (2001). Distribution of sex steroid hormone receptors in the avian brain:
Functional implications for neural sex differences and sexual behaviors. Microscopy
Research and Technique *55*, 1-11.

Gahr, M., Metzdorf, R., Schmidl, D., and Wickler, W. (2008). Bi-directional sexual
dimorphisms of the song control nucleus HVC in a songbird with unison song. PLoS One
3, e3073.

Gahr, M., Sonnenschein, E., and Wickler, W. (1998). Sex difference in the size of the
neural song control regions in a dueting songbird with similar song repertoire size of
males and females. Journal of Neuroscience *18*, 1124-1131.

Galili, T. (2015). dendextend: an R package for visualizing, adjusting and comparing
trees of hierarchical clustering. Bioinformatics *31*, 3718-3720.

Gautier, L., Cope, L., Bolstad, B.M., and Irizarry, R.A. (2004). affy--analysis of Affymetrix
GeneChip data at the probe level. Bioinformatics *20*, 307-315.

Geberzahn, N., and Gahr, M. (2011). Undirected (solitary) birdsong in female and male
blue-capped cordon-bleus (Uraeginthus cyanocephalus) and its endocrine correlates.
PLoS One *6*, e26485.

Genomatix (http://www.genomatix.de: Precigen Bioinformatics Germany GmbH
(PBG)).

Goymann, W., Möstl, E., and Gwinner, E. (2002). Non-invasive methods to measure
androgen metabolites in excrements of European stonechats, Saxicola torquata rubicola.
General and Comparative Endocrinology *129*, 80-87.

Griffiths, R., Double, M.C., Orr, K., and Dawson, R.J.G. (1998). A DNA test to sex most
birds. Molecular Ecology *7*, 1071-1075.

Gulledge, C.C., and Deviche, P. (1999). Photoperiod and testosterone independently
affect vocal control region volumes in adolescent male songbirds. Journal of
Neurobiology *36*, 550-558.

Gurney, M.E., and Konishi, M. (1980). Hormone-induced sexual differentiation of brain
and behavior in zebra finches. Science *208*, 1380-1383.

Hahnloser, R.H.R., Kozhevnikov, A.A., and Fee, M.S. (2002). An ultra-sparse code
underliesthe generation of neural sequences in a songbird. Nature *419*, 65-70.

Hall, M.L., Rittenbach, M.R.D., and Vehrencamp, S.L. (2015). Female song and vocal
interactions with males in a neotropical wren. Frontiers in Ecology and Evolution *3*.

Hall, Z.J., MacDougall-Shackleton, S.A., Osorio-Beristain, M., and Murphy, T.G. (2010).
Male bias in the song control system despite female bias in song rate in streak-backed
orioles (Icterus pustulatus). Brain Behav Evol *76*, 168-175.

Hartley, R.S., Chinn, M.S., and Ullrich, N.F.E. (1997). Left Syringeal Dominance in
Testosterone-Treated Female Canaries. Neurobiology of Learning and Memory *67*, 248-
253.

Hartley, R.S., and Suthers, R.A. (1989). Airflow and pressure during canary song: direct
evidence for mini-breaths. Journal of Comparative Physiology A *165*, 15-26.

Heid, P., Güttinger, H.R., and Pröve, E. (1985). The Influence of Castration and
Testosterone Replacement on the Song Architecture of Canaries (Serinus canaria).
Zeitschrift für Tierpsychologie *69*, 224-236.

Herpin, A., and Schartl, M. (2015). Plasticity of gene-regulatory networks controlling sex
determination: of masters, slaves, usual suspects, newcomers, and usurpators. EMBO
Rep *16*, 1260-1274.

Herrick, E.H., and Harris, J.O. (1957). Singing Female Canaries. Science *125*, 1299-1300.

Hoffmann, S., Trost, L., Voigt, C., Leitner, S., Lemazina, A., Sagunsky, H., Abels, M.,
Kollmansperger, S., Maat, A.T., and Gahr, M. (2019). Duets recorded in the wild reveal
that interindividually coordinated motor control enables cooperative behavior. Nat
Commun *10*, 2577.

Huang, D.W., Sherman, B.T., and Lempicki, R.A. (2008). Systematic and integrative
analysis of large gene lists using DAVID bioinformatics resources. Nature Protocols *4*,
44-57.

Huang, D.W., Sherman, B.T., and Lempicki, R.A. (2009). Bioinformatics enrichment tools:
paths toward the comprehensive functional analysis of large gene lists. Nucleic Acids
Research *37*, 1-13.

Immelmann, K. (1968). Zur biologischen Bedeutung des Estrildidengesanges, Vol 109.

Irwin, D.E. (2018). Sex chromosomes and speciation in birds and other ZW systems.
Molecular Ecology *0*.

Itoh, Y., Melamed, E., Yang, X., Kampf, K., Wang, S., Yehya, N., Van Nas, A., Replogle, K.,
Band, M.R., Clayton, D.F.*, et al.* (2007). Dosage compensation is less effective in birds
than in mammals. Journal of Biology *6*, 1-15.

Ko, M.-C., Van Meir, V., Vellema, M., and Gahr, M. (2020). Characteristics of song, brain-
anatomy and blood androgen levels in spontaneously singing female canaries.
Hormones and Behavior *117*, 104614.

Kreutzer, M., Vallet, E., and Doucet, S. (1992). Sexual responses of female canaries at the
onset of song stimuli. Experientia *48*, 679-682.

Lambeth, L.S., Raymond, C.S., Roeszler, K.N., Kuroiwa, A., Nakata, T., Zarkower, D., and
Smith, C.A. (2014). Over-expression of DMRT1 induces the male pathway in embryonic
chicken gonads. Developmental Biology *389*, 160-172.

Langfelder, P., and Horvath, S. (2008). WGCNA: an R package for weighted correlation
network analysis. BMC Bioinformatics *9*, 1-13.

Langfelder, P., and Horvath, S. (2012). Fast R Functions for Robust Correlations and
Hierarchical Clustering. Journal of statistical software *46*, i11.

Leitner, S., Voigt, C., and Gahr, M. (2001a). Seasonal Changes in the Song Pattern of the
Non-Domesticated Island Canary (Serinus Canaria) a Field Study. Behaviour *138*, 885-
904.

Leitner, S., Voigt, C., Garcia-Segura, L., Van't Hof, T., and Gahr, M. (2001b). Seasonal
activation and inactivation of song motor memories in wild canaries is not reflected in
neuroanatomical changes of forebrain song areas. Horm Behav *40*, 160 - 168.

Leonard, S.L. (1939). Induction of Singing in Female Canaries by Injections of Male
Hormone. Experimental Biology and Medicine *41*, 229-230.

Lobato, M., Vellema, M., Gahr, C., Leitão, A., de Lima, S.M.A., Geberzahn, N., and Gahr, M.
(2015). Mismatch in sexual dimorphism of developing song and song control system in
blue-capped cordon-bleus, a songbird species with singing females and males. Frontiers
in Ecology and Evolution *3*, 117.

MacDougall-Shackleton, S.A., and Ball, G.F. (1999). Comparative studies of sex
differences in the song-control system of songbirds. Trends in Neurosciences *22*, 432-
436.

Madison, F.N., Rouse, M.L., Jr., Balthazart, J., and Ball, G.F. (2015). Reversing song
behavior phenotype: Testosterone driven induction of singing and measures of song
quality in adult male and female canaries (Serinus canaria). Gen Comp Endocrinol *215*,
61-75.

Mank, J.E. (2013). Sex chromosome dosage compensation: definitely not for everyone.
Trends in Genetics *29*, 677-683.

Nätt, D., Agnvall, B., and Jensen, P. (2014). Large Sex Differences in Chicken Behavior and
Brain Gene Expression Coincide with Few Differences in Promoter DNA-Methylation.
PLoS ONE *9*, e96376.

Nixdorf, B.E., Davis, S.S., and DeVoogd, T.J. (1989). Morphology of Golgi-impregnated
neurons in hyperstriatum ventralis, pars caudalis in adult male and female canaries. J
Comp Neurol *284*, 337-349.

Nottebohm, F. (1980). Testosterone triggers growth of brain vocal control nuclei in
adult female canaries. Brain Res *189*, 429-436.

Nottebohm, F., and Arnold, A.P. (1976). Sexual dimorphism in vocal control areas of the
songbird brain. Science *194*, 211-213.

Nottebohm, F., Nottebohm, M., Crane, L., and Wingfield, J. (1987). Seasonal changes in
gonadal hormone levels of adult male canaries and their relation to song. Behav Neural
Biol *47*, 197 - 211.

Nottebohm, F., Nottebohm, M.E., and Crane, L. (1986). Developmental and seasonal
changes in canary song and their relation to changes in the anatomy of song-control
nuclei. Behavioral and neural biology *46*, 445-471.

Nottebohm, F., Stokes, T.M., and Leonard, C.M. (1976). Central control of song in the
canary, Serinus canarius. The Journal of Comparative Neurology *165*, 457-486.

Odom, K.J., Hall, M.L., Riebel, K., Omland, K.E., and Langmore, N.E. (2014). Female song is
widespread and ancestral in songbirds. Nat Commun *5*, 3379.

Ota, N., Gahr, M., and Soma, M. (2018). Couples showing off: Audience promotes both
male and female multimodal courtship display in a songbird. Sci Adv *4*, eaat4779.

Pesch, A., and Güttinger, H.-R. (1985). Der Gesang des weiblichen Kanarienvogels.
Journal für Ornithologie *126*, 108-110.

Pihlajamaa, P., Sahu, B., and Jänne, O.A. (2015). Determinants of Receptor- and Tissue-
Specific Actions in Androgen Signaling. Endocrine Reviews *36*, 357-384.

Price, J.J. (2019). Sex differences in song and plumage color do not evolve through sexual
selection alone: new insights from recent research. Journal of Ornithology *160*, 1213-
1219.

Price, J.J., Lanyon, S.M., and Omland, K.E. (2009). Losses of female song with changes
from tropical to temperate breeding in the New World blackbirds. Proceedings of the
Royal Society B: Biological Sciences *276*, 1971-1980.

Price, T.D. (2002). Domesticated Birds as a Model for the Genetics of Speciation by
Sexual Selection. Genetica *116*, 311-327.

Pryke, S., R. (2010). Sex chromosome linkage of mate preference and color signal
maintains assortative mating between interbreeding finch morphs. Evolution *64*, 1301-
1310.

R_Core_Team (2020). R: A language and environment for statistical computing
[Internet]. Vienna, Austria: R Foundation for Statistical Computing; (R Foundation for
Statistical Computing).

Rasika, S., Nottebohm, F., and Alvarez-Buylla, A. (1994). Testosterone increases the
recruitment and/or survival of new high vocal center neurons in adult female canaries.
Proceedings of the National Academy of Sciences *91*, 7854-7858.

Ringner, M. (2008). What is principal component analysis? Nat Biotechnol *26*, 303-304.

Robinson, C.M., Snyder, K.T., and Creanza, N. (2019). Correlated evolution between
repertoire size and song plasticity predicts that sexual selection on song promotes open-
ended learning. Elife *8*.

Sæther, S.A., Sætre, G.-P., Borge, T., Wiley, C., Svedin, N., Andersson, G., Veen, T., Haavie, J.,
Servedio, M.R., Bureš, S.*, et al.* (2007). Sex Chromosome-Linked Species Recognition and
Evolution of Reproductive Isolation in Flycatchers. Science *318*, 95.

Sayers, E.W., Beck, J., Brister, J.R., Bolton, E.E., Canese, K., Comeau, D.C., Funk, K., Ketter,
A., Kim, S., Kimchi, A.*, et al.* (2020). Database resources of the National Center for
Biotechnology Information. Nucleic Acids Res *48*, D9-D16.

Schulz, E.G. (2017). X-chromosome dosage as a modulator of pluripotency, signalling
and differentiation? Philosophical Transactions of the Royal Society B: Biological
Sciences *372*.

Schwabl, H., Dowling, J., Baldassarre, D.T., Gahr, M., Lindsay, W.R., and Webster, M.S.
(2015). Variation in song system anatomy and androgen levels does not correspond to
song characteristics in a tropical songbird. Animal Behaviour *104*, 39-50.

Shannon, P., Markiel, A., Ozier, O., Baliga, N.S., Wang, J.T., Ramage, D., Amin, N.,
Schwikowski, B., and Ideker, T. (2003). Cytoscape: a software environment for
integrated models of biomolecular interaction networks. Genome Res *13*, 2498-2504.

Shoemaker, H.H. (1939). Effect of Testosterone Propionate on Behavior of the Female
Canary. Proceedings of the Society for Experimental Biology and Medicine *41*, 299-302.

Slater, P.J.B., and Mann, N.I. (2004). Why do the females of many bird species sing in the
tropics? Journal of Avian Biology *35*, 289-294.

Smeds, L., Warmuth, V., Bolivar, P., Uebbing, S., Burri, R., Suh, A., Nater, A., Bures, S.,
Garamszegi, L.Z., Hogner, S., *et al.* (2015). Evolutionary analysis of the female-specific
avian W chromosome. Nat Commun *6*, 7330.

Smith, C.A., Roeszler, K.N., Ohnesorg, T., Cummins, D.M., Farlie, P.G., Doran, T.J., and
Sinclair, A.H. (2009). The avian Z-linked gene DMRT1 is required for male sex
determination in the chicken. Nature *461*, 267-271.

Smith, G.T., Brenowitz, E.A., and Wingfield, J.C. (1997). Roles of photoperiod and
testosterone in seasonal plasticity of the avian song control system. J Neurobiol *32*, 426-
442.

Stacklies, W., Redestig, H., Scholz, M., Walther, D., and Selbig, J. (2007). pcaMethods—a
bioconductor package providing PCA methods for incomplete data. Bioinformatics *23*,
1164-1167.

Takayama, K., Kaneshiro, K., Tsutsumi, S., Horie-Inoue, K., Ikeda, K., Urano, T., Ijichi, N.,
Ouchi, Y., Shirahige, K., Aburatani, H., *et al.* (2007). Identification of novel androgen
response genes in prostate cancer cells by coupling chromatin immunoprecipitation and
genomic microarray analysis. Oncogene *26*, 4453.

Tusher, V.G., Tibshirani, R., and Chu, G. (2001). Significance analysis of microarrays
applied to the ionizing radiation response. Proc Natl Acad Sci U S A *98*, 5116-5121.

Uebbing, S., Konzer, A., Xu, L., Backström, N., Brunström, B., Bergquist, J., and Ellegren, H.
(2015). Quantitative Mass Spectrometry Reveals Partial Translational Regulation for
Dosage Compensation in Chicken. Molecular Biology and Evolution.

Vallet, E., Kreutzer, M., and Gahr, M. (1996). Testosterone Induces Sexual Release Quality
in the Song of Female Canaries. Ethology *102*, 617-628.

Van Meir, V., Verhoye, M., Absil, P., Eens, M., Balthazart, J., and Van der Linden, A. (2004).
Differential effects of testosterone on neuronal populations and their connections in a
sensorimotor brain nucleus controlling song production in songbirds: a manganese
enhanced-magnetic resonance imaging study. Neuroimage *21*, 914-923.

Vellema, M., Diales Rocha, M., Bascones, S., Zsebők, S., Dreier, J., Leitner, S., Van der
Linden, A., Brewer, J., and Gahr, M. (2019a). Accelerated redevelopment of vocal skills is
preceded by lasting reorganization of the song motor circuitry. eLife *8*, e43194.

Vellema, M., Rocha, M., Bascones, S., Zsebők, S., Dreier, J., Leitner, S., Van der Linden, A.,
Brewer, J., and Gahr, M. (2019b). Accelerated redevelopment of vocal skills is preceded
by lasting reorganization of the song motor circuitry. eLife *8*, e43194.

Voigt, C., and Leitner, S. (2008). Seasonality in song behaviour revisited: Seasonal and
annual variants and invariants in the song of the domesticated canary (Serinus canaria).
Hormones and Behavior *54*, 373-378.

Wade, J. (2001). Zebra finch sexual differentiation: The aromatization hypothesis
revisited. Microscopy Research and Technique *54*, 354-363.

Wickler, W., and Seibt, U. (1980). Vocal Dueting and the Pair Bond II. Unisono dueting of
the African forest weaver, Symplectes bicolor. Zeitschrift für Tierpsychologie *52*, 217-
226.

Wild, J.M. (2004). Functional Neuroanatomy of the Sensorimotor Control of Singing.
Annals of the New York Academy of Sciences *1016*, 438-462.

Wilson, S., Qi, J., and Filipp, F.V. (2016). Refinement of the androgen response element
based on ChIP-Seq in androgen-insensitive and androgen-responsive prostate cancer
cell lines. Scientific Reports *6*, 32611.

Wingfield, J.C., Smith, J.P., and Farner, D.S. (1982). Endocrine Responses of White-
Crowned Sparrows to Environmental Stress. The Condor *84*, 399-409.

Wolf, J.B., and Bryk, J. (2011). General lack of global dosage compensation in ZZ/ZW
systems? Broadening the perspective with RNA-seq. BMC Genomics *12*, 91.

Yang, X., Schadt, E.E., Wang, S., Wang, H., Arnold, A.P., Ingram-Drake, L., Drake, T.A., and
Lusis, A.J. (2006). Tissue-specific expression and regulation of sexually dimorphic genes
in mice. Genome Res *16*, 995-1004.

Yates, A., Akanni, W., Amode, M.R., Barrell, D., Billis, K., Carvalho-Silva, D., Cummins, C.,
Clapham, P., Fitzgerald, S., Gil, L.*, et al.* (2016). Ensembl 2016. Nucleic Acids Research *44*,
D710-D716.

Zhang, R., Yin, Y., Sun, L., Yan, P., Zhou, Y., Wu, R., and Wu, X. (2017). Molecular cloning of
ESR2 and gene expression analysis of ESR1 and ESR2 in the pituitary gland of the
Chinese alligator (Alligator sinensis) during female reproductive cycle. Gene *623*, 15-23.

Zhao, C., Dahlman-Wright, K., and Gustafsson, J.-Å. (2010). Estrogen Signaling via
Estrogen Receptor β. Journal of Biological Chemistry *285*, 39575-39579.

Supplementary Tables

Supplementary Table 1. Experimental groups and sample sizes.

Species	Group	Sex	Singing	Tissue	Sample size
Forest weaver	Breeding	Female	Yes	HVC	6
		Male	Yes	HVC	6
Blue-capped cordon-bleu	Breeding	Female	Yes	HVC	6
		Male	Yes	HVC	6
Canary	Breeding	Female	No	HVC	6
		Male	Yes	HVC	6
	Nonbreeding	Female	Yes	HVC	6
		Female	No	HVC	5
		Male	No	HVC	6
		Female	Yes	HVC	6

Nonbreeding + testosterone	Male	Yes	HVC	6

826 **Supplementary Table 2. Pearson's correlation analysis of principal components**

827 **and variables (plasma androgen levels, HVC volume, sex, and singing).**

828 **Supplementary Table 3. Fisher's exact tests of sex-biased genes for chromosome**

829 **enrichment.**

830 **Supplementary Table 4. GO term enrichment analysis of sex-biased genes**

831 **identified from the forest weaver (comparison 1), cordon-bleu (comparison 2), and**

832 **canary (comparison 5) comparisons.**

Abbreviations: FWm, forest weaver male-biased genes; CBm, cordon-bleu male-biased genes; Cm, canary male-biased genes; FWf, forest weaver female-biased genes; CBm, cordon-bleu female-biased genes; Cm, canary female-biased genes.

833 **Supplementary Table 5. GO term enrichment analysis of female-specific, male-**

834 **specific and sex-shared genes.**

835 **Supplementary Figures**

836

837 **Supplementary Figure 1. Plasma androgen levels of seven groups of canaries on the**

838 **day of sacrifice.**

A, Canaries in breeding conditions. Singing males 1.76 ng/ml (mean), nonsinging females 59.2 pg/ml. Mann-Whitney Test, U = 6, P value = 0.07. **B,** Canaries in nonbreeding conditions. Nonsinging males 66.7 pg/ml (mean), nonsinging females 47.2 pg/ml, singing females 52.6 pg/ml. Mann-Whitney Test (nonsinging males vs. nonsinging females), U = 9, P value = 0.172. Mann-Whitney Test (nonsinging males vs. singing females), U = 24, P value = 0.377. **C,** Testosterone-implanted nonbreeding canaries. Males 13.5 ng/ml, females 9.31 ng/ml. Mann-Whitney Test, U = 13, P value = 0.485. Testosterone implantation significantly increased the plasma androgen levels of both nonbreeding males and females

Mann-Whitney Test (nonbreeding nonsinging males vs. testosterone-implanted singing males), U = 0, P value = 0.00492. Mann-Whitney Test (nonbreeding nonsinging females vs. testosterone-implanted singing females), U = 0, P value = 0.00217. The plasma androgen levels were higher in the breeding males than in the nonbreeding nonsinging males (Mann-Whitney Test, U = 32, P value = 0.0275). The boxes indicate the 25th/50th/75th percentiles (bottom/middle/top bar), and the extent of the whiskers indicates the most extreme values that are within 1.5 times the IQR (interquartile range) of the hinge. Each colour-coded dot indicates the measurement from one bird. The nonbreeding singing female canaries data were obtained from Ko et al. (2020).

Supplementary Figure 2. Hierarchical clustering of the HVC transcriptomes of 65

birds used in this study.

Hierarchical clustering showed that the HVC transcriptomes were first clustered based on the phylogenetic relationship; among canaries, nonbreeding females implanted with testosterone showed the most distinctive patterns. +T: testosterone implantation.

Supplementary Figure 3. Hierarchical clustering of the entopallium transcriptomes of 65 birds used in this study.

Hierarchical clustering showed that the entopallium transcriptomes were first clustered based on the phylogenetic relationship. The canaries were clustered by sex. +T: testosterone implantation.

Supplementary Figure 4. PCA scree plot showing the percentage of variance explained by each principal component.

Supplementary Figure 5. The majority of sex-biased genes showed low sex bias.

The bar graph summarizes the number of sex-biased genes in the HVC transcriptome identified from each male-to-female comparison. High bias: $|\log_2(\text{fold change})| \geq 2$; moderate bias: $1 \leq |\log_2(\text{fold change})| < 2$; low bias: $0.5 \leq |\log_2(\text{fold change})| < 1$. The phenotypes of the groups being compared in each male-to-female comparison are listed at the bottom of the graph.

Supplementary Figure 6. Pairwise comparisons between autosomal and Z chromosomal sex-biased genes.

A, Autosomal male-biased genes. **B,** Z chromosomal male-biased genes. **C,** Autosomal female-biased genes. **D,** Z chromosomal female-biased genes. The numbers indicated in the matrices are the proportions of identical genes identified from each pair. The phenotypes of the groups used for each comparison are listed at the bottom. B: breeding; NB: nonbreeding; T: testosterone implantation; the music note indicates the presence or absence of singing behaviour.