Highly social pinyon jays, but not less social Clark's nutcrackers, modify

their food-storing behaviour when observed by a heterospecific.

Alizée Vernouillet[1], Dawson Clary[2] & Debbie M. Kelly[1,2]

1. Department of Biological Sciences, University of Manitoba

2. Department of Psychology, University of Manitoba

Author Note

Alizée Vernouillet ORCID: https://orcid.org/0000-0003-2525-2936

Dawson Clary ORCID: https://orcid.org/0000-0002-0152-0484

Debbie M. Kelly ORCID: https://orcid.org/0000-0003-0575-1447

Alizée Vernouillet is now a Postdoctoral Researcher at the Department of

Experimental Psychology at Ghent University.

AV and DMK developed the study; AV conducted the experiments and

analysed the data; AV wrote the manuscript with comments from DC and DMK.

Research support was provided by a Natural Science and Engineering

Research Council Discovery grant (#RGPIN/4944-2017) and Canada Research Chair

to DMK.

The authors are grateful to Iroshini Gunasekara, Thomas Rawliuk, and Nicole

Tongol for helping with video scoring, and to Ben Farrar for his comments.

Data are available as supplementary material. The experiment was not

preregistered.

Correspondence concerning this article should be addressed to Alizée

Vernouillet. Email: alizee.vernouillet@gmail.com

Abstract

Behavioural flexibility can be described as the ability to use information and generalise it across contexts. Social living is thought to favour behavioural flexibility. We used a food-storing (caching) paradigm, during which individuals either ate or cached food under different conditions, to investigate whether they could flexibly adjust their caching behaviour when observed by conspecifics and heterospecifics. We examined the location and number of caches made by two corvid species differing in sociality, highly social pinyon jays (*Gymnorhinus cyanocephalus*) and less social Clark's nutcrackers (*Nucifraga columbiana*). Although pinyon jays cached a similar amount of food across conditions, they allocated more caches to areas less accessible to the observer when the observer spent more time close to the caching locations. Nutcrackers, however, reduced the number of seeds cached when observed by another nutcracker in comparison to when they cached alone, but did not significantly change their caching behaviour when observed by a pinyon jay. The differences in cache protection strategies, and the social cues (e.g., presence and behaviour of an observing bird) that elicit them, may be explained by the species' social organisation. Overall, our results provide insight into understanding how pressures associated with the social environment may influence foraging behaviours.

Keywords: caching, conspecifics, corvids, heterospecifics, nutcrackers, pinyon jays

Highly social pinyon jays, but not less social Clark's nutcrackers, modify their

food-storing behaviour when observed by a heterospecific.

Behavioural flexibility, the ability of an animal to adapt its behaviour to novel

circumstances, is thought to be one key component of complex cognition[1]. Flexible

behaviour may have evolved in species whose environment is highly unpredictable[2].

For example, species living in groups with changing relationships (e.g., fission-fusion)

may display more behavioural flexibility than less social species[2-5], as individuals in

complex social groups need to rapidly adjust their behaviour to dynamic social

environments. In such environments, it would be beneficial for individuals to switch

among behavioural strategies that best fit the current situation, but also to apply

adaptive behaviour to novel contexts.

Corvids are a taxonomic family of birds that are reported to possess many

sophisticated cognitive abilities, some of which rival those of primates[6-8] such as tool

use[9], future planning[10], and mirror self-recognition[11-12]. Corvid species differ in their

social organisations along a spectrum, which makes them ideal candidates for

investigating the relationship between sociality and behavioural flexibility. For

example, pinyon jays (*Gymnorhinus cyanocephalus*) live in flocks of hundreds of

individuals with high fission-fusion dynamics[13-14], whereas Clark's nutcrackers

(*Nucifraga columbiana*) are less social, with a social group consisting of the mating

pair and offspring of the year[13,15]. Hence, if the demands of a complex social life co-

evolve with behavioural and social flexibility, then pinyon jays might be expected to

show a more flexible behavioural repertoire than Clark's nutcrackers.

The socio-cognitive abilities of corvids are often assessed by their food-caching

behaviours. Caching refers to the behaviour of storing food to survive when resources

in the environment are scarce[16]. To protect their food caches from theft, corvids engage

in a suite of *cache protection strategies*, during which individuals flexibly change their

foraging behaviour depending on whether they are being observed. A well-studied

example is the California scrub jay (*Aphelocoma californica*), which supresses caching

in the presence of conspecifics[17], caches in locations that are less visible to the

observer[17], and selectively re-caches food items in new locations after the observing

bird has left[6].

To date, most studies examining cache protection strategies by corvids have focused

on moderately social species (e.g., scrub jays[6,17]) and less-social species (e.g.,

nutcrackers[18]; adult ravens[13]; Eurasian jays[19]). In contrast, few studies have addressed

the challenges encountered by highly social species (but see[20-21]), and no studies have

directly compared the caching behaviour of highly social and less-social species using

the same paradigm. Hence, we examined the caching behaviour of highly social pinyon

jays and of less-social Clark's nutcrackers when in the presence of a conspecific

observer, as well as when in the presence of a heterospecific observer. Displaying cache

protection strategies when observed by a heterospecific may require even greater

cognitive and behavioural flexibility, as the cacher must generalise the threat of losing

their caches to a different species[22] and appropriately modify its caching behaviour. We

also examined whether cachers changed their foraging behaviour based on social cues

provided by the observing bird during caching and pilfering. If living in complex social

structures influences behavioural flexibility, then highly social pinyon jays should be

more likely to generalise their cache protection strategies when faced with novel

heterospecific observers than less-social Clark's nutcrackers, but also that pinyon jays

would adjust their foraging behaviour based on more subtle social cues than Clark's

nutcrackers.

93   **Methods**

Subjects

Pinyon jays and Clark's nutcrackers served as caching birds (herein referred as

"cachers"; $n = 10$ for each species, five female jays and six female nutcrackers) or as

observing birds (herein referred as "observers"; two females for each species). All

individuals were captured as adults from populations around Flagstaff (Arizona, USA)

and were in captivity for approximately seven to ten years prior to this study. Five of

the nutcracker cachers had previous experience with caching paradigms[11,18,23]; the five

other nutcracker cachers, and all ten pinyon jay cachers had previous unrelated

experimental experience (e.g., concept learning[24-25]; exploratory behaviour[26]).

All birds were housed in individual cages (pinyon jays: 51 x 51 x 72 cm,

nutcrackers: 82 x 54 x 76 cm, width x depth x height, respectively), with multiple wood

perches, at the University of Manitoba, Canada. Pinyon jays were housed in a colony

room alongside California scrub jays, whereas nutcrackers were kept in a species-

specific colony room. All colony rooms were maintained at 22°C with a 12:12 day-

night cycle, with light onset at 0700 (Central Daylight time). These temperature and

lighting conditions correspond to the average temperature and photoperiod during

September in Flagstaff (USA), the peak caching season for pinyon jays and Clark's

nutcrackers. Birds were given *ad libitum* water and grit. When not experiencing the

food restriction procedure (see General procedures below), birds were fed a regular diet

consisting of a mixture of parrot pellets, turkey starter, sunflower seeds, mealworms,

peanuts, powder of oyster shells, and the vitamin powder supplement Prime®. Birds

were monitored and weighed daily to ensure a healthy weight during the experiment.

All applicable international, national, and institutional guidelines for the care

and use of animals were followed. Our research protocol was approved by University

of Manitoba's Animal Care Committee (#F2014-037) and complied with the guidelines

set by the Canadian Council on Animal Care.

Caching Apparatus

The caching apparatus was the same as used in Vernouillet et al.[21]. The

experiment was conducted in an experimental room, separated from the colony room.

Individual birds were tested in a cage (123.5 x 63.5 x 74.5 cm, width x depth x height),

divided into two equally-sized compartments. The entire cage was surrounded by white

curtains (Figure S1; also Figure 1 from Vernouillet et al. [21]). A transparent acrylic

divider separated the two compartments, each of which contained a perch. One

compartment served as the "caching compartment", whereas the other served as the

"observing compartment".

Within the caching compartment, cachers were given two plastic ice cube trays

(49.5 x 11 cm, length x width) each with 2 rows of 13 wells filled with sand. Trays were

made visually distinctive by affixing coloured plastic MegaBlocks™ at the base of each

tray. Each cacher received the same pair of distinct trays throughout the study, but the

arrangement of colored blocks differed across subjects. One tray (hereafter referred to

as the "pilfered tray") was positioned parallel and flush against the acrylic cage divider,

whereas the other tray (hereafter referred to as the "safe tray") was positioned on the

opposite wall of the caching compartment (i.e., parallel to the first tray). A food dish

was placed in the caching compartment beside the cage door, between the two trays

(Figure S1). All trials were recorded using an EverFocus® 1/3″ color digital camera

positioned either beside or above the experimental cage and using the motion tracking

program, BiObserve® through Windows XP.

General Procedures

The experimental procedures were the same as Vernouillet et al.[21]. Briefly, each

bird was given a weekly experimental session. Cachers were food deprived for 24h

before the start of the *Caching phase*, by removing all the food from their home cage.

This procedure created food uncertainty and motivated individuals to cache[18,21,27]. The

session started with a 45-min *Caching phase*, during which the cacher was provided

with a dish of 50 pine nuts to cache or consume. The observing compartment was

arranged as per the condition to be completed that session (see Conditions below). This

phase was immediately followed by a 3-min *Pilfering phase*, during which the pilfered

tray was placed in the observing compartment, alongside the divider and in the same

orientation as during the *Caching phase*. The safe tray was placed on a stool outside of

both compartments (remaining visible, but inaccessible to both birds). Upon the

completion of the *Pilfering phase*, all birds were returned to the colony room. After a

one-hour delay, the cacher was given a 5-min *Re-caching phase*, during which both

trays were returned to the cacher, such that they could eat or re-cache (i.e., move a pine

seed to a different location from where it was previously cached) some of the seeds

while alone. A re-cache was thus recorded when a pine seed was found in a different

well of either the same tray or the opposite tray, compared to where it was originally

placed. Upon completion of the *Re-caching phase*, the cacher was returned to the

colony room and provided with a small amount of food to ensure a healthy bodyweight

was maintained, while ensuring the bird would remain motivated to retrieve its caches

the following day. During the *Retrieving phase*, which occurred 24 hours after the

completion of the *Caching phase*, the cacher was placed in the caching compartment

with the safe and pilfered trays, unaltered from the previous *Re-caching phase*. The

cacher was permitted 45 minutes to consume or re-cache the previously cached pine

seeds, during which the observing compartment of the caching cage was empty (i.e.,

retrieval was always conducted while alone). Additional *Retrieving phases* were

administered, on the following day, every three hours if necessary, until the cacher

recovered the entirety of its caches. After each phase, the researcher recorded the

number and location of each pine seed (for more details, see Supplementary Material,

or Vernouillet et al. [21]).

Conditions. After three weeks of baseline trials during which individuals

experienced the exact same procedure as during the experiment in private (similar to

the *Alone* condition), each cacher experienced three blocks of four conditions (*Alone*,

Conspecific, Heterospecific and *Object*), with each condition randomly selected and

not repeated within the same block. Each condition refers to what (if anything) was

present in the observing compartment during the *Caching* and *Pilfering phases*.

Observers did not participate as cachers to exclude potential effects of pilfering

experience on their caching behaviour[28]. The pairing of observers to cacher remained

consistent throughout the experiment (each observer watched five cachers of each

species).

Alone. During the *Caching phase*, the observing compartment remained empty

(Figure 1a). During the *Pilfering phase*, the pilfered tray was placed in the empty

observing compartment for three minutes, and the pine seeds cached in that tray

remained undisturbed. This condition was conducted to assess the baseline caching

behaviour of cachers and was used to assess whether exposure to the experimental

conditions changed the caching behaviour of individuals.

Object. During the *Caching phase*, an inanimate object (a black water bottle,

27.5 x 7.0 cm, height x diameter) was affixed to the perch in the observing compartment

(Figure 1b). During the *Pilfering phase*, the pilfered tray was placed in the observing

compartment with the object for three minutes, whereas the safe tray was placed on a

stool outside of the cage, remaining visible to the cacher. The experimenter removed

pine seeds from the pilfered tray after the *Pilfering phase* (but before the *Re-caching*

phase). Pilfering rate was kept consistent across individuals by removing 33% of pine

seeds cached during the first and second blocks, and 50% of pine seeds cached during

the third block, to resemble natural variation of the pilfering rate in the wild[29]. Both

trays were given back to the cacher during the *Re-caching phase*. This condition served

as a control to determine whether the cacher modified its caching behaviour in response

to cache loss, and when compared with the *Conspecific* and *Heterospecific* conditions,

assessed the importance of social and motion cues in the display of cache protection

strategies.

Conspecific. During the *Caching phase*, a bird of the same species as the cacher

occupied the observing compartment (Figure 1c). During the *Pilfering phase*, the

pilfered tray was placed in the observing compartment with the observer, who was

given three minutes to access the tray in view of the cacher. The safe tray was placed

on an inaccessible stool outside of the cage, but visible to both birds. If the observer did

not pilfer enough seeds according to the pilfering rate described for the Object

condition, additional pine seeds were removed by the experimenter after the *Pilfering*

phase (but before the *Re-caching phase*). Similarly, if the observer pilfered more seeds

than the standardized threshold, the experimenter added caches back in the tray after

the *Pilfering phase* (but before the *Re-caching phase*). In both cases, the experimenter

adjusted the seed number such that all areas of the tray were pilfered equally. This

condition was conducted to assess whether the presence and behaviour of a conspecific

observer influenced the cacher's behaviour.

Heterospecific. This condition was conducted as during the *Conspecific*

condition, with the exception that the bird in the observing compartment was of a

different species than the cacher (Figure 1d). This condition was conducted to assess

whether the presence and the behaviour of a heterospecific observer influenced the

cacher's behaviour.

Behavioural Measures

Dependent variables. During the *Caching phase*, we examined the number of

pine seeds cached (in the trays and externally) and eaten. During the *Re-caching phase*,

we examined the proportion of re-cached seeds. This proportion was calculated by

dividing the number of caches found in new locations (compared to the location of

caches documented at the end of the *Caching phase*) by the total number of pine seeds

remaining after the *Re-caching phase*. All measures were evaluated at a global-level

(i.e., with respect to the total number of pine seeds cached) and at a tray-level, to

determine whether cachers associated the pilfering risk with the pilfered tray, but not

the safe tray. At the tray level, the proportion of re-cached pine seeds corresponds to

the proportion of pine seeds present in the tray at the end of the *Re-caching phase* that

were found in new locations in comparison to the locations documented at the end of

the *Caching phase*.

Behaviour of the observer. From the recorded sessions of the *Conspecific* and

Heterospecific conditions, we measured the duration (in seconds) the observer spent

interacting with the cacher during the *Caching phase* and with the pilfered tray during

the *Pilfering phase*. During the *Caching phase*, the observer was defined as interacting

with the cacher when the observer was standing within the third section of the observing

compartment closest to the divider separating the two compartments. During the

Pilfering phase, the observer was defined as interacting with the pilfered tray when

searching in the tray, retrieving pine seeds, or standing on the tray.

Statistical Analyses

We performed analyses separately for each species. We used a generalized

linear mixed model (GLMM) approach for our analyses, with the number of seeds

cached and eaten during the *Caching phase* and the proportion of seeds re-cached

during the *Pilfering phase* as dependent variables. Condition, duration of time the

observer spent interacting with the cacher during the *Caching phase,* and duration of

time the observer spent interacting with the pilfered tray during the *Pilfering phase*

were included as fixed factors in our models to assess whether the cacher responded to

the observer's behaviour during the *Caching* and *Re-caching phases*. Other fixed

factors included in the models were experimental block and the cacher's sex. We

performed separate analyses on the number of seeds cached in each tray and the

proportion of seeds re-cached in each tray to determine whether the cachers treated the

two trays differently. For these analyses, we included tray identity (pilfered vs. safe) as

an additional fixed factor. Identity of the cacher was included in all models as a random

factor to account for repeated measures taken on each cacher. Residual plots indicated

assumptions of normality were met for the number of caches made during the *Caching*

phase. We used the logarithmic number of pine seeds eaten during the *Caching phase*

and the square root of the proportion of pine seeds that were re-cached during the *Re-*

caching phase to meet the assumptions of normality. Analyses were conducted using R

version 3.3.2[30] with the *lme4*[31], *lsmeans*[32], *psych* [33], and *qpcR*[34] packages.

To assess the fit of each model, we used Akaike's Information Criterion (*AIC*).

Models were considered equivalent when *ΔAIC* < 2.0. We obtained *p*-values for each

fixed factor (i.e. tray, condition, block, sex, amount of time the observer interacted with

the cacher during the *Caching phase*, amount of time the observer interacted with the

pilfered tray during the *Pilfered phase*, block x condition, condition x tray, condition x

block x tray, condition x amount of time the observer interacted with the cacher during

the *Caching phase*, condition x amount of time the observer interacted with the pilfered

tray during the *Pilfered phase*, tray x amount of time the observer interacted with the

cacher during the *Caching phase*, tray x amount of time the observer interacted with

the pilfered tray during the *Pilfered phase*) using likelihood ratio tests of the model with

the fixed factor in question against the model without the fixed factor in question.

Parameter estimation was achieved using residual maximum likelihood. Post-hoc

analyses were conducted using pairwise comparisons. *P*-values of post-hoc analyses

were adjusted using the Tukey method to consider multiple comparisons[32]. Alpha was

set at 0.05 for all statistical analyses.

277   **Results**

Pinyon Jays

Number of pine seeds cached and eaten during the Caching phase. There

were no statistical differences in the overall number of pine seeds cached by pinyon

jays between conditions (*M ± SE*: *Alone*: 14.6 ± 2.4, *Object*: 15.0 ± 2.1, *Conspecific*:

15.6 ± 2.2, *Heterospecific*: 16.2 ± 2.2; $\chi^2_{(3)}$ = 0.579, *p* = 0.901, *ΔAIC* = 6.7; Figure 2;

Table S1). No other fixed factors examined in our analyses explained the total number

of seeds cached during the *Caching phase* by the pinyon jays.

Instead, we found that pinyon jays preferentially allocated their caches in the

safe tray when observed. The number of pine seeds cached in each tray was best

explained by the duration of time the observer spent interacting with the cacher during

the *Caching phase*, regardless of the species of the observer ($\chi^2_{(2)} = 28.978$, $p < 0.001$;

Table S1). As the observer spent more time interacting with the cacher, the cacher

decreased the number of pine seeds cached in the pilfered tray and increased the number

of pine seeds cached in the safe tray (Intercept: 7.50 ± 1.41 pine seeds; Pilfered tray

estimate: -0.18 ± 0.06 pine seeds/min; Safe tray estimate: $+0.24 \pm 0.06$ pine seeds/min;

Figure 3).

There were no statistical differences in the number of pine seeds eaten by pinyon

jays between conditions (see Supplementary Material, Table S2).

Proportion of re-cached pine seeds during the Re-caching phase. There

were no statistical differences in the proportion of re-cached pine seeds across

conditions by pinyon jays (*M ± SE*: *Alone*: 0.24 ± 0.05, *Object*: 0.20 ± 0.04,

Conspecific: 0.28 ± 0.06, *Heterospecific*: 0.29 ± 0.07; $\chi^2_{(3)} = 2.573$, $p = 0.462$, *ΔAIC =*

3.4; Table S3). No other fixed factors considered in our models explained the overall

proportion of re-cached pine seeds during the *Re-caching phase* by pinyon jays (*M ±*

SE: 0.25 ± 0.03; Table S3).

The proportion of re-cached pine seeds in each tray was best explained by the

amount of time the observer spent interacting with the cacher during the *Caching phase*

($\chi^2_{(2)} = 24.315$, $p < 0.001$; Table S3). After the observer spent more time interacting

with the cacher during the *Caching phase*, there was an increase in the proportion of

pine seeds present in the Pilfered tray that were re-cached (i.e., found in new locations

within the Pilfered tray), but not in the Safe tray, during the *Re-caching phase*

(Intercept: 0.378 ± 0.067, Pilfered tray estimate: 0.018 ± 0.006 proportion/min; Safe

tray estimate: -0.000 ± 0.006 proportion/min).

Clark's Nutcrackers

Number of pine seeds cached and eaten during the Caching phase. The

overall number of pine seeds cached by nutcrackers was best explained by conditions

($\chi^2_{(3)} = 11.953$, $p = 0.008$; Table S4; Figure 2). Nutcrackers cached significantly fewer

pine seeds when observed by a conspecific compared to when caching alone ($M \pm SE$:

Alone: 22.0 ± 1.6; *Conspecific*: 16.2 ± 1.8; $t = 3.39$, $p = 0.005$), and tended to cache

fewer pine seeds when observed by a conspecific than when observed by a

heterospecific (i.e., pinyon jay), though this difference was not significant ($M \pm SE$:

Heterospecific: 20.3 ± 1.8; $t = -2.37$, $p = 0.089$). There was no difference in the number

of pine seeds cached when caching alone and when observed by a heterospecific ($t =$

1.02, $p = 0.741$).

The number of pine seeds cached in each tray by the nutcrackers was also best

explained by condition ($\chi^2_{(7)} = 33.847$, $p < 0.001$; Figure 4, Table S4). Nutcrackers

cached more pine seeds in the pilfered tray than in the safe tray during the *Alone* ($M \pm$

SE: Pilfered tray: 13.6 ± 1.6, Safe tray: 8.5 ± 1.1; $t = 3.152$, $p = 0.002$; Figure 4) and

Object conditions ($M \pm SE$: Pilfered tray: 12.4 ± 1.5, Safe tray: 6.1 ± 1.1; $t = 3.867$, $p =$

0.001; Figure 4). During the *Conspecific* condition, nutcrackers showed an opposite

tendency to cache more pine seeds in the safe tray than in the pilfered tray, but this

difference did not reach statistical significance ($M \pm SE$: Pilfered tray: 6.6 ± 1.1, Safe

tray: 9.6 ± 1.3; $t = -3.050$, $p = 0.061$; Figure 4). There were no differences in the number

of pine seeds cached between trays during the *Heterospecific* condition ($M \pm SE$:

Pilfered tray: 10.9 ± 1.4, Safe tray: 9.4 ± 1.3; $t = 0.977$, $p = 0.330$; Figure 4).

There were no statistical differences in the number of pine seeds eaten by

Clark's nutcrackers between conditions (see Supplementary Material; Table S5).

Proportion of pine seeds re-cached during the Re-caching phase. There were

no statistical differences in the proportion of pine seeds re-cached across conditions by

nutcrackers ($M \pm SE$: *Alone*: 0.24 ± 0.05, *Object*: 0.24 ± 0.05, *Conspecific*: 0.23 ± 0.06,

Heterospecific: 0.32 ± 0.11; $\chi^2_{(3)} = 1.337$, $p = 0.721$, $\Delta AIC = 4.9$; Table S4). None of

the factors we considered in our models explained the overall proportion of re-cached

pine seeds by nutcrackers during the *Re-caching phase* ($M \pm SE$: 0.26 ± 0.03; Table

S6).

The best-ranked model explaining the proportion of pine seeds re-cached in

each tray during the *Re-caching phase* included only tray identity ($\chi^2_{(1)} = 27.017$, $p <$

0.001; Table S6). Across all conditions, the proportion of pine seeds present in the

Pilfered tray that were re-cached (i.e., found in new locations within the Pilfered tray)

was greater than the proportion of pine seeds present in the Safe tray that were re-cached

($M \pm SE$: Pilfered tray: 0.43 ± 0.04, Safe tray: 0.19 ± 0.03).

348   **Discussion**

In the presence of a pilfering conspecific, pinyon jays preferentially allocated

their caches in safer locations (see also[21]), whereas Clark's nutcrackers reduced the

number of caches made overall (see also[11,18,23]). The social cues triggering the display

of cache protection strategies differed between species. Pinyon jays' caching

behaviour was influenced by the amount of time an observer spent interacting near the

caching tray, whereas nutcrackers' caching behaviour was influenced by the mere

presence or absence of a conspecific.

In the presence of a pilfering heterospecific, pinyon jays allocated their caches in

safer locations in comparison to when they were caching alone, as they did when

observed by a conspecific. Also, of the caches remaining after the *Re-caching phase*, a

greater proportion of these caches were re-caches (i.e., found in new locations from

what the observer would have witnessed) in the pilfered tray compared to the safe tray.

Both the allocation of caches and the increase in re-caching depended on the duration

of time the observing bird interacted close to the cacher and the caching locations

during the caching session.

We found less evidence that Clark's nutcrackers modified their caching

behaviour when observed by a pinyon jay compared to when they cached alone.

Nutcrackers allocated their caches preferentially in the pilfered tray when they were

caching alone and in the presence of an object, but not when they were caching in

presence of an observing conspecific or heterospecific. This result may indicate that

nutcrackers generally associated the pilfered tray with a higher risk of losing their

caches when observed by another individual. This was also supported by the greater

proportion of the seeds present in the pilfered tray that were moved to a new location

by caching nutcrackers at the end of the re-caching phase than for the safe tray.

However, despite experiencing repeated cache loss to pinyon jay observers to the same

degree as with nutcracker observers, they did not reduce the amount of caches made,

nor did they preferentially allocate their caches in safe locations when caching. This

result may indicate that, unlike pinyon jay cachers, nutcrackers did not view the pinyon

jays as a pilfering threat.

An explanation for the species differences in caching behaviour may be that

observing pinyon jays spent less time interacting with cachers during the *Caching phase*

compared to observing nutcrackers (Table S7). Thus, nutcrackers may have viewed

pinyon jay observers as less threatening than conspecifics based on their behaviour,

despite similar rates of cache loss. Pinyon jays, however, viewed both conspecifics and

heterospecifics as threatening. More studies are required to determine which cues

nutcrackers use to assess the pilfering risk associated to the observer.

The social cues that elicit the display of cache protection behaviours in pinyon

jays and nutcrackers may also align with the demands of social living. Pinyon jays'

caching and re-caching behaviours were influenced by the amount of time an observer

spent interacting near the caching tray, whereas nutcrackers' caching behaviour was

influenced by the mere presence or absence of a conspecific. As highly social pinyon

jays rarely cache in private[14], it may be more advantageous for an individual to adjust

its behaviour only when warranted by the behaviour of the observer, and not simply in

the presence of an observer. On the contrary, less social species such as Clark's

nutcrackers and Eurasian jays (*Garrulus glandarius*) can reduce their caching rate in

the presence of another bird[18,19], since they have more opportunities to cache in private.

Therefore, pinyon jays may use more nuanced social information during caching to

adjust their behaviour than nutcrackers, due to the more restrictive conditions in which

they cache, regardless of whether the observer pilfers the caches afterwards[21]. The

difference in use of social cues may be evidence that pinyon jays have evolved greater

behavioural flexibility compared to nutcrackers.

Social information can be exchanged between conspecifics and heterospecifics

when multiple species share the same territory or resources. For example, some species

are influenced by heterospecifics when responding to alarm calls[36,37], when choosing a

breeding site[38,39], or even when foraging[40]. In the case of our study species, individual

Clark's nutcrackers briefly join flocks of pinyon jays for protection during late summer

and early fall, when both species collect pine seeds to cache for the winter[42]. Both

species' caches are susceptible to pilfering by one another, but also by Steller's jays

(*Cyanocitta stelleri*), another social jay species[43]. Here, we provide experimental

evidence that one corvid species, the pinyon jay, can flexibly use the social information

provided by an observer, regardless of the species, and adjust their caching behaviour

accordingly. To date, only one other study evaluated whether a caching species, the

mountain chickadee (*Poecile gambeli*), can adjust its caching behaviour in the presence

of a pilfering heterospecific, the red-breasted nuthatch (*Sitta canadensis*;[22]). Mountain

chickadees cached preferentially in hidden sites in presence of observing nuthatches

but used both hidden and visible sites when they were alone, suggesting chickadees

could view nuthatches as a threat for their caches. Both pinyon jays and mountain

chickadees have complex social behaviour[44,45], supporting the social living hypotheses

that suggest social living might promote more flexible behaviours, especially in a social

context[2,5].

Our results provide equivocal evidence that social living could promote cognitive

flexibility, at least in corvids[2]. Up until now, most socio-cognitive abilities thought to

be present in corvids have been found in both social and non-social corvid species. For

instance, both social magpies and less-social nutcrackers show evidence of mirror self-

recognition[11,12]. Even in a social context, nutcrackers and more social corvids such as

pinyon jays and azure-winged magpies can both perform complex social interactions

with conspecifics, like cooperatively sharing food[23,46,47]. Based on our findings, one

possibility is that the ability to view a conspecific as a threat might be common to all

corvids, but that only highly social species adjust their caching behaviour to the social

context due to the selection pressures associated with social living[2,5].

Alternatively, the species difference in caching behaviour could also be

ontogenetic, as pinyon jays will likely witness a greater number of caching and pilfering

events between members of their social group, and in more varied social contexts than

nutcrackers. These additional learning opportunities may facilitate greater

generalisation of the cognitive abilities underlying cache protection behaviours. In both

cases, social living could explain the differences in behavioural flexibility between the

two caching species.

In summary, the current study investigated, for the first time, whether corvids

display cache protection strategies in the presence of a heterospecific in a controlled

setting. We found that highly social pinyon jays and less social Clark's nutcrackers

displayed different caching behaviours when observed. Pinyon jays preferentially

allocate cached food in areas less accessible to a conspecific or heterospecific observer,

but primarily when the observer spent more time near a caching tray. Contrarily,

nutcrackers reduced the number of caches they made, but only when observed by

another nutcracker. Overall, these findings suggest that corvids may differ in their use

of social information and in the extent to which they interpret heterospecifics as a threat.

These findings may indicate that the ability to view a conspecific as a threat is shared

among corvids regardless of social organisation. However, highly social species may

display more behavioural flexibility by generalising the use of cache protection

strategies against heterospecifics to a greater extent and by adjusting their caching

behaviour to the social context. Our results provide insight into understanding how

pressures associated with the social environment may influence the caching behaviour

of corvids and support the idea of more social species adjusting their behaviour more

flexibly than less social species. Our findings provide a step further into understanding

how complex cognition has evolved.

References

1. Emery, N. J., & Clayton, N. S. (2004). The mentality of crows: convergent evolution of intelligence in corvids and apes. *Science, 306,* 1903-1907.

2. Bond, A. B., Kamil, A. C., & Balda, R. P. (2007). Serial reversal learning and the evolution of behavioral flexibility in three species of North American corvids (*Gymnorhinus cyanocephalus*, *Nucifraga columbiana*, *Aphelocoma californica*). *Journal of Comparative Psychology, 121*, 372-379.

3. Blumstein, D.T., & Armitage, K. B. (1997). Does sociality drive the evolution of communicative complexity? A comparative test with ground-dwelling sciurid alarm calls. *American Naturalist, 150,* 179-200.

4. Dunbar, R.I.M., & Shultz, S. (2007). Evolution in the social brain. *Science, 317,* 1344.

5. Taborsky, B., & Oliveira, R.F. (2012). Social competence: an evolutionary approach. *Trends in Ecology and Evolution, 27,* 679-688.

6. Emery, N.J., Dally, J.M., & Clayton, N.S. (2004). Western scrub-jays (*Aphelocoma californica*) use cognitive strategies to protect their caches from thieving conspecifics. *Animal Cognition, 7,* 37-43.

7. Seed, A., Emery, N., & Clayton, N. (2009). Intelligence in corvids and apes: a case of convergence evolution? *Ethology, 115,* 401-420.

8. Taylor, A. H. (2014) Corvid cognition. *Wiley Interdisciplinary Reviews: Cognitive Science, 5,* 361-372.

9. Bluff, L.A., Weir, A.A.S., Rutz, C., Wimpenny, J.H., & Kacelnik, A. (2007) Tool-related cognition in New Caledonian crows. *Comparative Cognition & Behavior Reviews, 2,* 1-25.

10. Raby, C.R., Alexis, D.M., Dickinson, A., & Clayton, N.S. (2007). Planning for the future by western scrub-jays. *Nature, 445,* 919-921.

11. Clary, D., & Kelly, D.M. (2016a). Graded mirror self-recognition by Clark's nutcrackers. *Scientific Reports, 6,* 36459.

12. Prior, H., Schwartz, A., & Güntürkün, O. (2008). Mirror-induced behaviour in the magpie (Pica pica): evidence of self-recognition. *PLoS Biology, 6,* e202.

13. Clayton, N.S., & Emery, N.J. (2008) The social life of corvids. *Current Biology, 17,* 652-656.

14. Marzluff, J. M., & R. P. Balda. (1992). The Pinyon Jay: the behavioral ecology of a colonial and cooperative corvid. T & AD Poyser, London, UK.

15. Tomback, D.F. (1998). Clark's Nutcracker (*Nucifraga columbiana*) in *The Birds of North America* (P.G. Rodewald, editor). Cornell Lab of Ornithology, Ithaca, NY, USA.

16. Vander Wall, S. B. (1990) Food hoarding in animals. Chicago: University of Chicago Press.

17. Dally, J.M., Emery, N.J., & Clayton, N.S. (2005). Cache protection strategies by western scrub-jays, *Aphelocoma californica*: Implications for social cognition. *Animal Behaviour, 70,* 1251-1263.

18. Clary, D., & Kelly, D.M. (2011). Cache protection strategies of a non-social food-caching corvid, Clark's nutcracker (*Nucifraga columbiana*). *Animal Cognition, 14,* 735-44.

19. Shaw, R. C., & Clayton, N. S. (2012) Eurasian jays, *Garrulus glandarius*, flexibly switch caching and pilfering tactics in response to social context. *Animal Behaviour, 84,* 1191-1200.

20. Bednekoff, P. A., & Balda, R. P. (1996). Social caching and observational spatial memory in pinyon jays. *Behaviour, 133,* 807-826.

21. Vernouillet, A., Casidsid, H.J.M., & Kelly, D.M. (2020). Conspecific presence, but not pilferage, influences pinyon jays' (*Gymnorhinus cyanocephalus*) caching behavior. *Learning & Behavior.*

22. Pravosudov, V.V. (2008). Mountain chickadees discriminate between potential cache pilferers and non-pilferers. *Proceedings of the Royal Society B: Biological Sciences, 275,* 55-61.

23. Clary, D., & Kelly, D.M. (2016b). Clark's nutcrackers (*Nucifraga columbiana*) flexibly adapt caching behavior to a cooperative context. *Frontiers in Psychology, 7,* 1-11.

24. Wright, A.A., Magnotti, J.F., Katz, J.S., Leonard, K., Vernouillet, A., & Kelly, D.M. (2017). Corvids outperform pigeons and primates in learning a basic concept. *Psychological Science, 28,* 437-444.

25. Vernouillet, A., Leonard, K., Katz, J.S., Magnotti, J.F., Wright, A.A., & Kelly, D.M. (In press). Abstract-concept learning in two species of New World corvids, pinyon jays (*Gymnorhinus cyanocephalus*) and California scrub jays (A*phelocoma californica*). *Journal of Experimental Psychology: Animal Learning and Cognition.*

26. Vernouillet, A., & Kelly, D.M. (2020). Individual exploratory responses are not repeatable across time or contexts for four species of food-storing corvid. *Scientific Reports, 10,* 394.

27. Kamil, A.C., & Balda, R.P. (1985). Cache recovery and spatial memory in Clark's nutcrackers (*Nucifraga columbiana*). *Journal of Experimental Psychology: Animal Behavior Processes, 11,* 95-111.

28. Emery, N.J., & Clayton, N.S. (2001). Effects of experience and social context
on prospective caching strategies by scrub jays. *Nature, 414,* 443-446.

29. Vander Wall, S. B., & Jenkins, S. (2003) Reciprocal pilferage and the evolution
of food-hoarding behavior. *Behavioral Ecology, 14,* 656-667.

30. R Core Team (2013). R: A language and environment for statistical computing.
R Foundation for Statistical Computing, Vienna, Austria.URL http://www.R-
project.org/.

31. Bates, D., Maechler, M., Bolker, B., & Walker, S. (2015). Fitting linear mixed-
effects models using lme4. *Journal of Statistical Software, 67,* 1-48.

32. Lenth, R.V. (2016). Least-squares means: the R package lsmeans. *Journal of*
Statistical Software, 69, 1-33.

33. Revelle, W.R. (2017). Psych: procedures for personality and psychological
research. Software. Northwestern University, Evanston, Illinois, USA.
https://CRAN.R-project.org/package=psych

34. Spiess, A.N. (2018). qpcR: Modelling and analysis of real-time PCR data.
https://CRAN.R-project.org/package=qpcR

35. Bugnyar, T., & Heinrich, B. (2005). Pilfering ravens, *Corvus corax*, adjust their
behavior to social context and identity of competitors. *Animal Cognition, 9,*
369-376.

36. Dutour, M., & Danel, S. (2020). Wild great tits' alarm calls prompt vigilant
behaviours in free-range chickens. *Animal Cognition.*

37. Magrath, R.D., Haff, T.M., Fallow, P.M., & Radford, A.N. (2015).
Eavesdropping on heterospecific alarm calls: from mechanisms to
consequences. *Biological Reviews, 90,* 560-586.

38. Morinay, J., Forsman, J.K., Germain, M., & Doligez, B. (2020). Behavioural
traits modulate the use of heterospecific social information for nest site
selection: experimental evidence from a wild bird population. *Proceedings of*
the Royal Society B, 287, 20200265.

39. Tolvanen, J., Kivelä, S.M., Doligez, B., Morinay, J., Gustafsson, L., Bijma, P.,
Pakanen, V.M., & Forsman, J.T. (2020). Quantitative genetics of the use of
conspecific and heterospecific social cues. *Evolution, 74,* 2332-2347.

40. Romero-Gonzalez, J. E., Solvi, C., & Chittka, L. (2020). Honey bees adjust
colour preferences in response to concurrent social information from
conspecifics and heterospecifics. *Animal Behaviour.*

41. Sridhar, H., & Guttal, V. (2018). Friendship across species borders: factors that
facilitate and constrain heterospecific sociality. *Philosophical Transactions B,*
373, 20170014.

42. Balda, R. P., Bateman, G. C., & Foster, G. F. (1972). Flocking associates of the
Piñon Jay. *Wilson Bulletin, 84,* 60-73.

43. Burnell, K.L., & Tomback, D.F. (1985). Steller's jays steal gray jay caches:
field and laboratory observations. *The Auk, 102,*417-419.

44. Harrap, S., & Quinn, D. (1995). Chickadees, tits, nuthatches and treecreepers.
Princeton, NJ: Princeton University Press.

45. Hogstad, O. (1989). Social organization and dominance behavior in some *Parus*
species. *Wilson Bulletin, 101,* 254–262.

46. Duque, J.F., & Stevens, J.R. (2016). Voluntary food sharing in pinyon jays: The
role of reciprocity and dominance. Animal Behaviour, 122, 135-144.

47. Horn, L., Scheer, C., Bugnyar, T., & Massen, J. M. (2016). Proactive

prosociality in a cooperatively breeding corvid, the azure-winged magpie

(*Cyanopica cyana*). *Biology Letters, 12,* 20160649.

Figure 1. Pictorial representation of the experimental conditions during the *Caching*

phase. The caching compartment contained the cacher (nearest bird, here a pinyon

jay), and the observing compartment **a)** remained empty, contained **b)** an object, **c)** a

conspecific (here, another pinyon jay), or **d)** a heterospecific (here, a Clark's

nutcracker).

Figure 2. Total number of pine seeds cached (± *SE*) by Clark's nutcrackers and

pinyon jays during the *Caching phase* during the Alone, Object, Conspecific, and

Heterospecific conditions. Nutcrackers cached less when in the presence of another

nutcracker (*Conspecific* condition) in comparison to when they were caching alone (*p*

= 0.008). There was no difference in the number of seeds cached by pinyon jays

across conditions (*p* = 0.901).

Figure 3. Number of pine seeds cached by pinyon jays in the pilfered tray and in the

safe tray during the *Caching phase* depended on the amount of time the observing

bird interacted with the caching bird ($p < 0.001$).

Figure 4. Number of pine seeds cached in the safe and in the pilfered trays by caching

nutcrackers in each experimental condition during the *Caching phase* ($p < 0.001$).