**Poly(rC)-binding protein 1 limits hepatitis C virus virion assembly and secretion**

**Sophie E. Cousineau[1], Marylin Rheault[1], and Selena M. Sagan[1,2]#**

[1]Department of Microbiology & Immunology, McGill University, Montreal, QC, Canada

[2]Department of Biochemistry, McGill University, Montreal, QC, Canada

#Corresponding Author: Dr. Selena Sagan, McGill University, 3655 Promenade Sir William

Osler, Room 805, Montréal, QC, H3G 1Y6; email: selena.sagan@mcgill.ca

**Keywords:** Hepatitis C virus (HCV), poly(rC)-binding protein 1 (PCBP1), hnRNP E1,

assembly, virion secretion

**Running title**: PCBP1 limits HCV assembly and secretion

**ABSTRACT**

The hepatitis C virus (HCV) co-opts numerous cellular elements – including proteins, lipids, and microRNAs – to complete its viral life cycle. The cellular RNA-binding protein, poly(rC)-binding protein 1 (PCBP1), was previously reported to bind to the 5′ untranslated region (UTR) of the HCV genome; however, its importance in the viral life cycle has remained unclear. Herein, we sought to clarify the role of PCBP1 in the HCV life cycle. Using the HCV cell culture (HCVcc) system, we found that knockdown of endogenous PCBP1 resulted in an overall decrease in viral RNA accumulation, yet resulted in an increase in extracellular viral titers. To dissect PCBP1's specific role in the HCV life cycle, we carried out assays for viral entry, translation, genome stability, RNA replication, as well as virion assembly and secretion. We found that PCBP1 knockdown did not directly affect viral entry, translation, RNA stability, or RNA replication, but resulted in an overall increase in infectious particle secretion. This increase in virion secretion was evident even when viral RNA synthesis was inhibited, and blocking virus secretion could partially restore the viral RNA accumulation decreased by PCBP1 knockdown. We therefore propose a model where endogenous PCBP1 normally limits virion assembly and secretion, which increases viral RNA accumulation in infected cells by preventing the departure of viral genomes packaged into virions. Overall, our findings improve our understanding of how cellular RNA-binding proteins influence viral genomic RNA utilization during the HCV life cycle.

## INTRODUCTION

Hepatitis C virus (HCV) is an enveloped virus of the *Flaviviridae* family (genus: hepacivirus) that typically causes a persistent liver infection (1). Its ~9.6 kb single-stranded, positive-sense RNA genome contains a single open reading frame flanked by 5´ and 3´ untranslated regions (UTR). A highly structured internal ribosomal entry site (IRES) in the 5´ UTR drives the translation of the viral polyprotein, which is subsequently processed into 10 mature viral proteins: 3 structural proteins (core, E1 and E2 glycoproteins), and 7 non-structural (NS) proteins (p7, NS2, NS3, NS4A, NS4B, NS5A and NS5B) (1, 2). While the structural proteins form the nucleocapsid and viral envelope, the NS3-5B proteins form the viral replicase, required for viral RNA replication (3). The p7, NS2, NS3 and NS5A proteins are also implicated in viral genome packaging and the assembly of viral particles (4-7). As a positive-sense RNA virus, the HCV genome itself must serve as a template for viral translation, genome replication, and packaging; however, the mechanisms that determine which process each viral RNA is engaged in at any given time have not been well defined. Importantly, due to their limited coding capacity, viruses are highly dependent on the molecular machinery of the host cell. While numerous cellular proteins and RNAs have been shown to interact with the HCV genome, their precise role(s) in the viral life cycle often remain poorly defined.

The poly(rC)-binding protein 1 (PCBP1), along with its paralogs hnRNP K and PCBP2, is one of the most abundant cellular RNA-binding proteins with an affinity for poly(rC) (8). These multifunctional proteins can regulate translation and enhance the stability of their cellular mRNA targets, which they interact with through their hnRNP K homologous (KH) domains (9). Notably, all three paralogs have been reported to bind to the HCV 5´ UTR (10-12). However, the degree to which each protein has been studied in the context of HCV infection varies

significantly – while hnRNP K and PCBP2 have been fairly extensively studied and reported to

play markedly different roles in the viral life cycle, the role of PCBP1 in the HCV life cycle has

not been investigated in detail (13, 14). Beyond its interactions with the 5′ UTR, previous reports

suggested that PCBP1 was not necessary for HCV IRES-mediated translation, but that

knockdown of PCBP1 decreases HCV RNA accumulation during infection (15, 16).

Herein, we sought to further clarify the role of PCBP1 in the HCV life cycle. Using cell

culture-adapted strains of HCV (HCVcc), we found that PCBP1 knockdown decreased viral

RNA accumulation, yet led to an increase in virion secretion. By examining individual steps of

the viral life cycle, we ruled out a direct role for PCBP1 in viral entry, translation, genome

stability and viral RNA replication. Further analysis of the assembly step revealed that, similarly

to its paralog hnRNP K, endogenous PCBP1 limits HCV virion assembly and release.

**MATERIALS AND METHODS**

**Cell culture**

Huh-7.5 human hepatoma cells were obtained from Charlie Rice (Rockefeller University) and

maintained in complete media: Dulbecco's Modified Eagle Media (DMEM) supplemented with

inactivated 10% fetal bovine serum (FBS), 2 mM L-glutamine, and 1X MEM non-essential

amino acids. Human embryonic kidney (293T) cells were kindly provided by Martin J. Richer

(McGill University, Montreal, QC, Canada) and were maintained in DMEM supplemented with

10% FBS. All cell lines were maintained at 37°C/5% $CO_2$ and were routinely screened for

mycoplasma contamination.

**Plasmids and viral RNAs**

The pJFH-1$_T$ plasmid, encoding a cell culture-adapted Japanese Fulminant Hepatitis (JFH-1; HCV genotype 2a) with three adaptive mutations that increase viral titers in cell culture, was provided by Rodney Russell (Memorial University of Newfoundland) (17). The pJ6/JFH1 plasmid bears a full-length viral sequence derived from the J6 (structural genes through first transmembrane domain of NS2) and JFH-1 (remaining NS genes) isolates of HCV, and the pJ6/JFH-1 FL RLuc GNN ("RLuc-GNN") viral sequence also includes a *Renilla* luciferase (RLuc) reporter gene inserted between p7 and NS2 and an inactivating GNN mutation within the NS5B RNA polymerase active site (4, 18). The pJ6/JFH-1 mono RLuc-NS2 plasmid ("Δcore-p7") – a truncated version of the *Renilla* reporter virus with a deletion of the structural genes through p7 – was provided by Joyce Wilson (University of Saskatchewan) (19).

To make full-length uncapped viral RNAs, all plasmid templates were linearized and *in vitro* transcribed as previously described (20). The firefly luciferase (FLuc) mRNA was transcribed from the Luciferase T7 Control DNA plasmid (Promega) linearized using *XmnI* and *in vitro* transcribed using the mMessage mMachine T7 Kit (Life Technologies) according to the manufacturer's instructions.

**Generation of infectious HCV stocks**

To generate JFH-1$_T$ stocks, 30 µg of *in vitro* transcribed RNA was transfected into Huh-7.5 cells using the DMRIE-C reagent (Life Technologies) according to the manufacturer's instructions. Four days post-transfection, infectious cell supernatants were passed through a 0.45 µm filter and infectious viral titers were determined by focus-forming unit assay (17). Infectious virus was amplified for two passages through Huh-7.5 cells at a MOI of 0.2 before being used for

infections. To generate J6/JFH-1 stocks, 25 µg of *in vitro* transcribed RNA was electroporated

into 6 x $10^6$ Huh-7.5 cells using a Bio-Rad GenePulser XCell at 270 V, 950 µF, and infinite

resistance in a 4-mm cuvette. Two days post-electroporation, infectious cell supernatants were

passed through a 0.45 µm filter and concentrated at least 10-fold using Amicon Ultra-15-100K

centrifugal filter units (Millipore Sigma) according to the manufacturer's instructions. Viral titers

were determined by focus-forming unit assay. All viral stocks were aliquoted and stored at -80°C

until use.

**Focus-forming unit (FFU) assays**

One day prior to infection, 8-well chamber slides (Lab-Tek) were seeded with 4 x $10^5$ Huh-7.5

cells/well. Infections were performed with 10-fold serial dilutions of viral samples in 100 µL for

4 h, after which the supernatant was replaced with fresh media. Three days post-infection, slides

were fixed in 100% acetone and stained with anti-HCV core antibody (1:100, clone B2,

Anogen), and subsequently with the AlexaFluor-488-conjugated anti-mouse antibody (1:200,

ThermoFisher Scientific) for immunofluorescence analysis. Viral titers are expressed as the

number of focus-forming units (FFU) per mL.

Extracellular virus titers were determined directly from cell supernatants, while

intracellular virus titers were determined after cell pellets were subjected to lysis via four freeze-

thaw cycles, removal of cellular debris via centrifugation, and recovery of virus-containing

supernatants.

**MicroRNAs and siRNA sequences**

siGL3 (siCTRL): 5'-CUUACGCUGAGUACUUCGAUU-3', siGL3* : 5'-

UCGAAGUACUCAGCGUAAGUU-3', miR122$_{p2-8}$ (siCTRL for luciferase experiments): 5'-

UAAUCACAGACAAUGGUGUUUGU-3', miR122$_{p2-8}$*: 5'-

AAACGCCAUUAUCUGUGAGGAUA-3' (21), siPCBP1: 5'-

CUGUGUAAUUUCUGGUCAGUU-3', siPCBP1*: 5'-CUGACCAGAAAUUACACAGUU-3'

(16) were all synthesized by Integrated DNA Technologies.

All microRNA and siRNA duplexes were diluted to a final concentration of 20 μM in

RNA annealing buffer (150 mM HEPES pH 7.4, 500 mM potassium acetate, 10 mM magnesium

acetate), annealed at 37°C for 1 h and stored at -20°C. For all knockdown experiments, 50 nM

siRNA transfections were conducted 2 days prior to infection or electroporation of viral RNAs.

Transfections were conducted using Lipofectamine RNAiMAX (Invitrogen) according to the

manufacturer's instructions with the modification that 20 μL of reagent were used to transfect a

10-cm dish of cells.

**HCV and VSV pseudoparticles (HCVpp and VSVpp)**

HCVpp consisting of a FLuc reporter lentiviral vector pseudotyped with the HCV E1 and E2

glycoprotein (from the H77 isolate, a genotype 1a strain) were a kind gift from John Law

(University of Alberta) (22). To generate lentiviral vectors pseudotyped with the VSV-G

glycoprotein (VSVpp), a 90% confluent 10-cm dish of 293T cells were transfected with 10 μg

pPRIME-FLuc, 5 μg psPAX.2, and 2.5 μg pVSV-G plasmid with 10 μL Lipofectamine 2000

(Invitrogen) diluted in 4 mL Opti-MEM. Media was changed 4, 20, and 28 h post-transfection.

At 48 h post-transfection, the cell culture media was passed through a 0.45 µm filter and stored

at -80°C.

To assay for cell entry, HCVpp and VSVpp were diluted 1/3 in dilution media (1X

DMEM, 3% FBS, 100 IU penicillin and 100 µg/mL streptomycin) with 20 mM HEPES and 4

µg/µL polybrene, and introduced to Huh-7.5 cells by spinoculation at 1,200 rpm for 1 h at room

temperature. The cells were left to recover at 37°C for at least 5 h before the pseudoparticle-

containing media was changed for fresh complete Huh-7.5 media. In parallel, cells seeded in a 6-

well plate were transfected with 1 µg of pPRIME-FLuc plasmid using Lipofectamine 2000

(Invitrogen) according to the manufacturer's instructions. Three days post-spinoculation and

transfection, cells were lysed in passive lysis buffer (Promega) and FLuc activity was assayed

using the Dual Reporter Luciferase kit (Promega).

**Infections**

Three days prior to infection, 10-cm dishes were seeded with $5 \times 10^5$ Huh-7.5 cells, which were

transfected with siRNA duplexes on the following day. On the day of infection, each 10-cm dish

– containing approximately $1 \times 10^6$ cells – was infected with $5 \times 10^4$ FFU of JFH-$1_T$ diluted in 3

mL complete media. Four to five hours post-infection, each infected plate was split into three 10-

cm dishes. Protein, RNA, and virus samples were collected three days post-infection.

**Electroporations**

For each electroporation, 400 µL of resuspended cells ($1.5 \times 10^7$ cell/mL) were mixed with 2 µg

of FLuc mRNA and 5 µg of replicating (Δcore-p7 J6/JFH-1 RNA) or 10 µg GNN J6/JFH-1

RNA, and electroporated in 4-mm cuvettes at 270 V, 950 µF, and infinite resistance, optimized

for the Bio-Rad GenePulser XCell (Bio-Rad). Electroporated cells were resuspended in complete

Huh-7.5 media and transferred to 6-well plates for luciferase assays and protein analysis.

**Inhibition of RNA replication by 2′CMA**

Two days post-siRNA transfection, Huh-7.5 cells were infected with JFH-1$_T$ at a MOI of 0.05.

Four to five hours post-infection, each plate of infected cells was split into 6-well plates. Three

days post-infection, the media on these cells was changed for complete Huh-7.5 media with 5

μM 2′CMA (2′C-methyladenosine, Carbosynth), an HCV NS5B polymerase inhibitor, or DMSO

(vehicle control) (23). Total RNA and intracellular virus samples were collected at 0, 6 and 12 h

post-treatment, while cell culture supernatants were collected 6 and 12 h post-treatment. Protein

samples were collected from untreated plates to assess PCBP1 knockdown efficiency by Western

blot.

**Inhibition of virus secretion by Brefeldin A**

Three days post-infection with JFH-1$_T$, the cell culture media was replaced with complete Huh-

7.5 media with 5 μg/mL BFA (Brefeldin A, Biolegend) or equal volumes of DMSO (vehicle

control) (24). Total RNA, intracellular virus, and extracellular virus samples were collected at 0

and 6 h post-treatment. Protein samples were collected from untreated cells to verify PCBP1

knockdown by Western blot.

**Western blot analysis**

To collect total intracellular protein samples, cells were lysed in RIPA buffer (150 mM sodium

chloride, 1% NP-40, 0.5% sodium deoxycholate, 0.1% SDS, 50 mM Tris pH 8.0), supplemented

with Complete Protease Inhibitor Cocktail (Roche) and frozen at -80°C. Cellular debris was

pelleted by centrifugation at 16,000 x g for 30 min at 4°C, and the supernatant was quantified by

BCA Protein Assay (ThermoScientific). Ten micrograms of sample were loaded onto 10-12%

SDS-PAGE gels. Samples were transferred onto Immobilon-P PVDF membranes (Millipore),

blocked in 5% milk, and incubated overnight with primary antibodies diluted in 5% BSA: rabbit

anti-PCBP1 (clone EPR11055, Abcam ab168378, diluted 1:10,000); rabbit anti-actin (A2066,

Sigma, 1:20,000); mouse anti-HCV core (clone B2, Anogen MO-I40015B, 1:7,500); mouse anti-

JFH-1 NS5A (clone 7B5, BioFront Technologies, 1:10,000). Blots were incubated for 1 hour

with HRP-conjugated secondary antibodies diluted in 5% skim milk: anti-mouse (HAF007, R&D

Systems, 1:25,000); anti-rabbit (111-035-144, Jackson ImmunoResearch Laboratories, 1:50,000)

and visualized using enhanced chemiluminescence (ECL Prime Western Blotting Detection

Reagent, Fisher Scientific).

**RNA isolation and Northern blot analysis**

Total RNA was harvested using TRIzol Reagent (ThermoFisher Scientific) according to the

manufacturer's instructions. Ten micrograms of total RNA were separated on a 1% agarose gel

containing 1X 3-(N-morpholino)propanesulfonic acid (MOPS) buffer and 2.2 M formaldehyde

and transferred to a Zeta-probe membrane (Bio-Rad) by capillary transfer in 20X SSC buffer (3

M NaCl, 0.3 M sodium citrate). Membranes were hybridized in ExpressHyb Hybridization

Buffer (ClonTech) to random-primed $^{32}$P-labeled DNA probes (RadPrime DNA labelling

system, Life Technologies) complementary to HCV (nt 84-374) and γ-actin (nt 685-1171).

Autoradiograph band densities were quantified using Fiji (25).

**RT-qPCR analysis**

The iTaq Universal Probes One-Step kit (Bio-Rad) was used to perform duplex assays probing

for the HCV genome (NS5B-FW primer: 5'-AGACACTCCCCTATCAATTCATGGC-3';

NS5B-RV primer: 5'-GCGTCAAGCCCGTGTAACC-3'; NS5B-FAM probe: 5'-

ATGGGTTCGCATGGTCCTAATGACACAC-3') and the GAPDH loading control (PrimePCR

Probe assay with HEX probe, Bio-Rad). Each 20 µL reaction contained 500 ng of total RNA, 1.5

µL of the HCV primers and probe, and 0.5 µL of the GAPDH primers and probe. RT-PCR

reactions were conducted in a CFX96 Touch Deep Well Real-Time PCR system (Bio-Rad).

Genome copies were calculated using a standard curve and fold-differences in gene expression

were calculated using the $2^{-\Delta\Delta Ct}$ method (26).

**Luciferase assays**

For translation and replication assays, cells were washed in PBS and harvested in 100 µL of 1X

passive lysis buffer (Promega). The Dual-Luciferase Assay Reporter Kit (Promega) was used to

measure both *Renilla* and firefly luciferase activity according to the manufacturer's instructions

with the modification that 25 µL of reagent were used with 10 µL of sample. All samples were

measured in triplicate.

**Data analysis**

Statistical analyses were performed using GraphPad Prism v9 (GraphPad, USA). Statistical

significance was determined by paired t-test to compare results obtained from multiple

experiments, and by two-way ANOVA with Geisser-Greenhouse and Bonferroni corrections

when more than two comparisons were applied at once. To calculate half-lives, a one-step decay

curve using the least-squares regression was used, and error was reported as the asymmetrical

(profile likelihood) 95% confidence interval of the half-life. To calculate virus accumulation and

virus secretion rates, a simple linear regression was performed using the least squares regression

method. The slope and standard error calculated for each regression represents the rate of virus

accumulation or secretion.

**RESULTS**

**PCBP1 plays a role in the HCV life cycle**

Previous reports suggested that the PCBP1 protein directly interacted with the 5′ UTR of the

HCV genome, and a siRNA screen reported that PCBP1 knockdown reduced HCV RNA

accumulation (10, 16). Thus, we sought to further characterize the role of PCBP1 in the HCV life

cycle. Firstly, we assessed how PCBP1 knockdown affected the accumulation of HCVcc using

two viral strains: the cell culture-adapted JFH-1$_T$ strain (which has three adaptive mutations in

E2, p7, and NS2), and the chimeric J6/JFH-1 strain (where the coding region of the core through

to the first transmembrane domain of NS2 comes from the J6 sequence, and the remaining

sequences are derived from JFH-1) (**Figure 1A**) (17, 18). We found that knocking down

endogenous PCBP1 decreased viral protein expression and HCV RNA accumulation of both

strains, resulting in an approximately 2.2-fold and 3.3-fold decrease in JFH-1$_T$ and J6/JFH-1

viral RNA accumulation, respectively (**Figure 1B-D**). Quantification of intracellular virions

revealed no significant differences in JFH-1$_T$ titers between the PCBP1 knockdown and control

conditions; however, we observed an approximately 2.6-fold reduction in intracellular J6/JFH-1

titers during PCBP1 knockdown (**Figure 1E**). Surprisingly, when we quantified extracellular

virions, we observed that PCBP1 knockdown resulted in an overall increase in extracellular titers

of approximately 3.9-fold and 2.6-fold for JFH-1$_T$ and J6/JFH-1, respectively (**Figure 1F**).

Overall, PCBP1 knockdown increased the total quantity of virions produced by 3.7-fold for JFH-

1$_T$ and 2.1-fold for J6/JFH-1 (**Supplementary Figure S1**). To verify that this effect was

specifically due to the decrease in endogenous PCBP1 protein expression, we attempted to

rescue PCBP1 protein levels by ectopically expressing a siRNA-resistant PCBP1 with a N-

terminal FLAG tag (**Supplementary Figure S2**). Although we were unable to ectopically

express PCBP1 to the same level as the endogenous protein, we observed that ectopic PCBP1

expression diminished the impact of endogenous PCBP1 knockdown. Thus, in line with previous

findings, we found that PCBP1 knockdown decreased viral protein expression and intracellular

viral RNA accumulation. However, despite this overall decrease in viral protein and RNA

accumulation, we observed a marked increase in virion production and these effects were

partially reversed by ectopic expression of PCBP1.

**Figure 1. PCBP1 knockdown decreases HCV protein expression and intracellular viral RNA accumulation, but increases secreted virus titers. (A)** Schematic representation of the

HCVcc infectious particles and genomic RNA used in infections. Huh-7.5 cells were transfected

with siPCBP1 or siCTRL two days prior to infection with JFH-1$_T$ or J6/JFH-1 (MOI = 0.05).

Total protein, RNA, and intracellular and extracellular infectious virus were harvested at day 3

post-infection. **(B)** Viral protein expression analysis by Western blot (ns, non-specific band;

arrow, core and NS5A bands). **(C)** Viral RNA accumulation analysis by Northern blot and **(D)**

quantification by RT-qPCR. **(E)** Intracellular and **(F)** extracellular (secreted) virus titers,

quantified by FFU assay. All data are representative of at least three independent biological

replicates, and error bars represent the standard deviation of the mean. Statistical significance

was calculated by paired t-test (ns, not significant; ** $p < 0.01$; *** $p < 0.001$; **** $p < 0.0001$).

**PCBP1 knockdown does not have a direct impact on HCV entry, translation, genome**

**stability or viral RNA replication**

To clarify PCBP1's precise role(s) in the HCV life cycle, we used assays to specifically examine

different steps of the viral life cycle (**Figure 2**). Firstly, we explored whether PCBP1 knockdown

had an effect on viral entry using the HCV pseudoparticle (HCVpp) system. HCVpp are

lentiviral vectors with a firefly luciferase reporter gene pseudotyped with the HCV E1 and E2

glycoproteins (22). HCVpp engage with HCV-specific entry receptors, then enter cells by

clathrin-mediated endocytosis; thus, to account for any changes in clathrin-mediated endocytosis,

we used vesicular stomatitis virus (VSV) pseudoparticles (VSVpp) as a control. In addition, to

verify that PCBP1 knockdown did not affect luciferase reporter gene expression, we assessed

firefly luciferase expression from cells directly transfected with a FLuc reporter plasmid. In all

cases, we found that depletion of endogenous PCBP1 had no impact on luciferase activity

(**Figure 2A**). This suggests that PCBP1 knockdown does not affect FLuc reporter expression,

clathrin-mediated endocytosis, or HCVpp entry.

**Figure 2. PCBP1 knockdown has no direct effect on HCV entry, translation, genome**

**stability or RNA replication. (A)** siRNA-transfected Huh-7.5 cells were spinoculated with

luciferase reporter pseudoparticles expressing the HCV E1/E2 glycoproteins (HCVpp) or the

VSV-G glycoprotein (VSVpp). In parallel, cells were transfected with a firefly luciferase

expression plasmid. Samples were harvested 3 days post-infection/transfection and analyzed by

luciferase assay. **(B)** siRNA-transfected Huh-7.5 cells were electroporated with replication-

defective J6/JFH-1-RLuc-GNN RNAs, and luciferase activity was monitored at several

timepoints in the two days post-electroporation. **(C)** Huh-7.5 cells were electroporated with the

replication-competent Δcore-p7 J6/JFH RLuc RNA, and luciferase activity was monitored at

several timepoints post-electroporation. RLuc values were normalized to the early timepoint

(3h), to control for disparities in electroporation efficiency between experiments. All data are

representative of at least three independent biological replicates and error bars represent the

standard deviation of the mean.

To examine PCBP1's impact on viral translation, we used a full-length J6/JFH-1 RLuc

reporter RNA containing an inactivating mutation in the NS5B polymerase gene (GNN). The

RLuc activity thus served as a direct measure of HCV IRES-mediated translation, and over time,

this signal also served as a proxy measure for viral RNA stability. We found that the PCBP1

knockdown and control conditions had similar RLuc activity at all timepoints (**Figure 2B**).

Moreover, the luciferase signal half-lives were nearly identical, with 2.68 h (95% CI 1.64 – 4.64)

and 2.71 h (95% CI 1.43 – 5.50) for the PCBP1 knockdown and control conditions, respectively.

Thus, PCBP1 knockdown does not directly impact either HCV IRES-mediated translation or

genome stability.

To assess the impact of PCBP1 knockdown on HCV RNA replication, we made use of a

J6/JFH-1 subgenomic replicon RLuc reporter RNA, with a deletion of the structural proteins

(Δcore-p7), which allowed us to assess viral RNA replication in the absence of virion assembly

(**Figure 2C**). We observed no significant differences in subgenomic replicon RNA accumulation

during PCBP1 knockdown compared with control conditions, suggesting that PCBP1

knockdown does not affect viral RNA replication. Interestingly, we also observed no significant

differences in luciferase activity during PCBP1 knockdown and control conditions of a full-

length WT J6/JFH-1 RLuc genomic RNA or of a full-length J6/JFH-1 genomic RNA containing

a GFP insertion in the NS5A gene (NS5A-GFP), which was previously shown to impair virion

assembly without impairing viral RNA replication (**Supplementary Figure S3**) (6, 27). While

this is seemingly in contrast to our findings with HCVcc, where PCBP1 depletion decreased viral

RNA levels (**Figure 1C and D**); it is important to note that all of these reporter viral RNAs are

defective in viral packaging due to the large RLuc reporter gene insertion, which results in >100-

to >1000-fold reductions in infectious virion production (**Supplementary Figure S3**). Thus,

taken together, our data suggests that PCBP1 knockdown does not have a direct effect on viral

RNA replication.

**PCBP1 knockdown limits virus secretion**

As we had found that silencing PCBP1 did not have a significant impact on HCV entry, translation, genome stability or viral RNA replication, we reasoned that PCBP1 was likely to be exerting an effect on virion assembly and/or secretion. To investigate this possibility, we used a nucleoside analog, 2´C-methyladenosine (2´CMA), to block viral RNA synthesis by the HCV NS5B RNA-dependent RNA polymerase and monitored virion production over time (**Figure 3**) (23). Similar to related studies using Zika virus, we reasoned that blocking viral RNA synthesis could allow us to assess if PCBP1 knockdown modulated viral assembly and secretion in the absence of genomic RNA production (28). To this end, we infected cells with JFH-1$_T$ and, three days post-infection, treated cells with 2´CMA or vehicle (DMSO), and monitored viral RNA accumulation as well as intracellular and extracellular virion production over the next 12 hours (**Figure 3A**). In agreement with our previous findings, we observed an overall reduction in JFH-1$_T$ viral RNA accumulation by day 3 post-infection during PCBP1 knockdown (0 h timepoint, **Figure 3B**). In addition, 2´CMA treatment efficiently blocked further viral RNA accumulation, which continued to increase under the vehicle (DMSO) condition (6-12 h timepoints, **Figure 3B**). We observed that intracellular viral titers were initially similar in both PCBP1 knockdown and control conditions, but that infectious virions accumulated at a slightly faster rate during PCBP1 knockdown in DMSO-treated cells (**Figure 3C and D**). In contrast, the 2´CMA treatment induced a similar decrease in intracellular titers during both PCBP1 knockdown and control conditions, resulting in similar rates of intracellular virion accumulation (**Figure 3C and D**). Interestingly, as we had observed previously (**Figure 1F**), PCBP1 knockdown resulted in an increase in extracellular titers which continued upon 2´CMA treatment (**Figure 3E and F**). Despite an overall reduction in extracellular virion production during 2´CMA treatment, we observed that PCBP1 knockdown still resulted in higher extracellular titers and virion secretion rates than the control condition under both 2´CMA and vehicle (DMSO) treatment (**Figure 3E and F**). Taken together, these results suggest that endogenous PCBP1 typically limits the secretion of infectious viral particles during the HCV life cycle.

**Figure 3. PCBP1 knockdown enhances infectious virion secretion even in the absence of RNA replication**. **(A)** Schematic representation of the experimental approach for 2′CMA experiments: two days post-siRNA transfection, Huh-7.5 cells were infected with JFH-1$_T$ (MOI = 0.05). Three days post-infection, the cell culture medium was replaced with media containing

2´CMA or DMSO (vehicle control). Total RNA, intracellular and extracellular virus were

collected at t = 0, 6 and 12 h post-treatment. **(B)** Quantitative RT-PCR analysis after 2´CMA

treatment. **(C)** Intracellular viral titers and **(D)** intracellular virus accumulation rate calculated by

linear regression. **(E)** Extracellular viral titers and **(F)** virus secretion rates calculated by linear

regression. All data are representative of three independent replicates and error bars in (B), (C)

and (E) represent the standard deviation of the mean. Error bars in (D) and (F) represent the

slopes of the linear regressions ± standard error. Statistical significance was calculated by two-

way ANOVA. (ns, not significant; ** $p < 0.01$; **** $p < 0.0001$).

**Blocking virion secretion partially restores HCV RNA accumulation during PCBP1**

**knockdown**

During virion secretion, viral genomes that have been packaged into new infectious particles are

removed from the cell. Since our previous findings suggested that PCBP1 knockdown increases

virion secretion, we wondered if this increased egress of assembled viral particles (packaged

viral genomic RNAs) might explain our initial observation that PCBP1 knockdown led to an

overall reduction in viral RNA accumulation (**Figure 1C-D**). To directly test this, we treated

HCV-infected cells with Brefeldin A (BFA), a protein trafficking inhibitor previously shown to

prevent HCV virion secretion (**Figure 4**) (24). As previously reported, we found that BFA

treatment blocked the secretion of infectious virions, resulting in a sharp decline in extracellular

titers and an increase in intracellular titers relative to untreated cells or to vehicle (DMSO)

treatment (**Figure 4A and B**). Consistent with our prior results, PCBP1 knockdown resulted in a

greater than 2-fold decrease in viral RNA accumulation, while BFA treatment resulted in a

partial rescue of HCV RNA accumulation during PCBP1 knockdown (**Figure 4C**). Thus,

blocking cellular secretory pathways, thereby preventing virion secretion, can partially rescue

intracellular viral RNA accumulation defects resultant from PCBP1 knockdown.

**Figure 4. Blocking HCV secretion partially rescues intracellular RNA accumulation during**
**PCBP1 knockdown.** Three days post-JFH-1$_T$ infection (MOI 0.05), the cell culture medium was
replaced with media containing Brefeldin A (BFA) or DMSO (vehicle control). Total
intracellular RNA, intracellular virus, and extracellular virus were collected at 0 h and 6 h post-
treatment. **(A)** Extracellular (secreted) viral titers and **(B)** intracellular viral titers from untreated
cells (0h) and after 6h of DMSO or BFA treatment, quantified by FFU assay. **(C)** Intracellular
RNA accumulation relative to the siCTRL condition, assessed by RT-qPCR analysis. All data
are representative of three independent biological replicates, and error bars represent the standard
deviation of the mean. Statistical significance was calculated by paired t-test (ns, not significant;
** $p < 0.01$; **** $p < 0.0001$).

**DISCUSSION**

Herein, we investigated the role of PCBP1 in the HCV life cycle. Previous studies had found that

PCBP1 interacts with the HCV 5′ UTR *in vitro* (10, 15), and that PCBP1 silencing inhibited

HCV RNA accumulation in cell culture (16), yet the precise steps of the HCV life cycle

modulated by PCBP1 remained unclear. Our systematic evaluation of the effects of PCBP1

knockdown on individual steps of the viral life cycle revealed that PCBP1 depletion did not

directly affect virus entry, translation, genome stability or viral RNA replication, but rather

resulted in increased infectious virion production in cell culture. Specifically, PCBP1

knockdown significantly increased virion secretion, resulting in the depletion of intracellular

viral RNA. Additionally, inhibiting virion secretion could partially restore HCV RNA

accumulation during PCBP1 knockdown. Thus, our findings suggest a model whereby

endogenous PCBP1 normally limits virion assembly and secretion, such that viral genomes

remain available to engage in the translation and replication steps of the viral life cycle (**Figure**

**5**). As such, PCBP1 depletion increases the proportion of viral genomes engaged in virion

assembly and secretion, which decreases the pool of translating/replicating viral RNAs. We

anticipate that these differences in intracellular viral RNA levels would compound over time,

with PCBP1-replete cells rapidly increasing their pool of intracellular viral RNA through

repeated cycles of translation and RNA replication, while the smaller pool of

translating/replicating RNAs in PCBP1-depleted cells would result in a much slower rate of viral

RNA accumulation over time.

**Figure 5. Model for PCBP1's effects on HCV assembly and secretion.** Under wild-type
conditions, endogenous PCBP1 limits HCV assembly and secretion, and the majority of
intracellular viral RNAs are engaged in translation and viral genome replication. In contrast,
PCBP1 knockdown promotes virion assembly and secretion. This results in a shift in viral RNA
utilization such that an increased proportion of viral RNAs are engaged in viral particle
production, leaving a smaller proportion of viral RNAs to engage in the translating/replicating
pool.

Notably, most of the experiments conducted herein made use of JFH-1$_T$, a cell culture-

adapted strain that produces greater yields of infectious virions in cell culture than the parental

JFH-1 strain (5, 17). To verify that the phenotype we observed was not an artefact of the JFH-1$_T$

strain itself, we validated our findings with J6/JFH-1, a chimeric virus that is widely used in

HCVcc studies, commonly referred to as Jc1 (18). Knockdown of PCBP1 had a similar effect on

JFH-1$_T$ and J6/JFH-1 protein expression, RNA accumulation, and extracellular viral titers;

however, we found that while JFH-1$_T$ intracellular titers were not significantly different from

control condition, J6/JFH-1 intracellular titers were significantly decreased by approximately

2.6-fold (**Figure 1E**). Nonetheless, PCBP1 knockdown still resulted in an overall greater than 2-

fold increase in the total number of infectious virions produced by both strains (**Supplementary**

**Figure S1**). Since an infectious viral particle must have undergone the process of virion

assembly, this suggests that PCBP1 depletion enhances the assembly of infectious virions in

addition to their subsequent secretion. Importantly, this data could reflect pre-existing strain

differences in virion assembly, as J6/JFH-1 has previously been reported to be more efficient

than JFH-1 in virion assembly (29, 30). Specifically, the J6 core protein was shown to localize

closer to the sites of virion assembly on the endoplasmic reticulum (ER) than the JFH-1 core

protein, which accumulates to a greater extent on lipid droplets (29, 30). Furthermore, J6/JFH-1

allocates a greater proportion of its viral genomes to virion assembly than JFH-1, which allocates

more viral RNAs to the translating/replicating pool (29, 30). Thus, it is possible that PCBP1

knockdown promotes both the assembly and secretion of J6/JFH-1 virions, but minor gains in the

already efficient virion assembly step are less apparent than the gains in virion secretion for this

particular strain.

Interestingly, PCBP1 has also been previously implicated in attenuation of antiviral

signalling, specifically in the regulation and turnover of MAVS, a signal transduction protein

directly downstream of RIG-I, which is known to limit HCV infection (31). However, all of our

conclusions were drawn from experiments conducted in the Huh-7.5 cell line, which has well-

documented defects in the RIG-I antiviral signalling pathway (32). Additionally, our own brief

explorations of interferon induction and MAVS turnover did not reveal any significant

differences between PCBP1 knockdown and control conditions (data not shown). Moreover, the

HCV NS3-4A protease also inactivates MAVS during infection to reduce antiviral signaling and

recognition of the viral RNA (33, 34). However, as hepatocytes typically express RIG-I and

MAVS, we cannot rule out the possibility that PCBP1 may also modulate this pathway during

infection *in vivo*.

Finally, PCBP1's closely related paralogs, hnRNP K and PCBP2, have been more

extensively studied during HCV infection. While the hnRNP K protein was shown to restrict

infectious virion production, the PCBP2 protein has been suggested to modulate viral translation

and RNA replication (13, 14). Since the PCBP1 amino acid identity is far more similar to PCBP2

(~80% identity) than hnRNP K (~24% identity), we were initially surprised that our findings for

PCBP1 in HCV infection closely matched those reported for hnRNP K. Yet, while PCBP1 and

PCBP2 have been shown to perform similar functions, they have also been reported to play

distinct roles during poliovirus, vesicular stomatitis virus, and human immunodeficiency virus

infection (35-37). In addition to the similarities we observed between PCBP1 and the effects

previously reported for hnRNP K, our results and conclusions also echo those reported for the

IGF2BP2/YBX-1 complex, METTL3/METTL14 $N$6-methyladenosine (m$^6$A) writers, and the

YTHDF m$^6$A-binding proteins during HCV infection (13, 38, 39). All of these complexes have

been reported to inhibit HCV infectious particle production, with no effect on viral translation or

RNA replication – with the exception of the IGF2BP2/YBX-1 complex, which plays an

additional role in facilitating viral RNA replication (38). It is currently unclear if PCBP1, hnRNP

K, the YBX-1 complex, or $m^6A$ modification of the HCV genome inhibit virion assembly as

components of a common pathway, or through distinct and/or additive mechanisms.

Interestingly, high-throughput affinity capture and proximity ligation studies have found that

PCBP1 interacts with hnRNP K, IGF2BP2, YBX-1, METTL3 and METTL14; although this has

been demonstrated only in non-hepatic cell lines to date (40-42). Should these interactions be

conserved during HCV infection, it seems plausible that PCBP1 may participate in one or more

of these complexes which together inhibit HCV virion assembly and secretion. Future

investigations will thus be needed to reveal whether these proteins function in an overlapping or

distinct manner; and, are likely to further improve our understanding of HCV virion assembly as

well as how this process is regulated by cellular RNA binding proteins.

In conclusion, our results support a model whereby endogenous PCBP1 limits HCV

infectious particle production. By preventing virion assembly and secretion, PCBP1 indirectly

enhances viral RNA accumulation. The model presented herein helps to inform our

understanding of how cellular RNA-binding proteins modulate HCV genomic RNA utilization

during the viral life cycle, specifically as it pertains to virion assembly. While the precise

molecular mechanism(s) employed by PCBP1 to inhibit HCV assembly and egress remain to be

characterized, similar phenotypes reported for hnRNP K, IGF2BP2/YBX-1 and $m^6A$

modification of the HCV genome offer promising leads for future investigations.

**ACKNOWLEDGEMENTS**

We would like to acknowledge Charlie Rice (Rockefeller University) for kindly providing the

Huh-7.5 cells, pJ6/JFH (pJc1), pJ6/JFH FL RLuc WT and GNN plasmids; Rodney Russell

(Memorial University) for providing JFH-1$_T$; Mamata Panigrahi and Joyce Wilson (University of

Saskatchewan) for the pJ6/JFH mono RLuc NS2 ("Δcore-p7") plasmid; Martin J Richer (McGill

University) for the 293T cells; John Law (University of Alberta) for the kind gift of HCVpp. We

are also grateful to Nathan Taylor, Julie Magnus, and Carolina Camargo (McGill University) for

technical support, and to Craig McCormick's graduate journal club (Dalhousie University) for

their critical feedback on a previous iteration of this manuscript.

**FUNDING INFORMATION**

This research was supported by the Canadian Institutes for Health Research (CIHR) [MOP-

136915 and PJT-169214]. S.E.C. was supported by the Canadian Network on Hepatitis C

(CanHepC) training program, as well as a Vanier Canada Graduate Scholarship. M.R. was also

supported by the CanHepC training program. In addition, this research was undertaken, in part,

thanks to the Canada Research Chairs program (S.M.S.).

**AUTHOR CONTRIBUTIONS**

S.E.C. and S.M.S. designed the study; S.E.C. and M.R. performed the experiments and analyzed

the data, and S.E.C and S.M.S. wrote and edited the manuscript.

## REFERENCES

1.  **Tsukiyama-Kohara K**, **Iizuka N**, **Kohara M**, **Nomoto A**. 1992. Internal ribosome entry site within hepatitis C virus RNA. Journal of Virology **66**:1476–1483.

2.  **Lancaster AM**, **Jan E**, **Sarnow P**. 2006. Initiation factor-independent translation mediated by the hepatitis C virus internal ribosome entry site. RNA **12**:894–902.

3.  **Lohmann V**, **Korner F**, **Koch J**, **Herian U**, **Theilmann L**, **Bartenschlager R**. 1999. Replication of subgenomic hepatitis C virus RNAs in a hepatoma cell line. Science **285**:110–113.

4.  **Jones CT**, **Murray CL**, **Eastman DK**, **Tassello J**, **Rice CM**. 2007. Hepatitis C virus p7 and NS2 proteins are essential for production of infectious virus. Journal of Virology **81**:8374–8383.

5.  **Jones DM**, **Atoom AM**, **Zhang X**, **Kottilil S**, **Russell RS**. 2011. A genetic interaction between the core and NS3 proteins of hepatitis C virus is essential for production of infectious virus. Journal of Virology **85**:12351–12361.

6.  **Masaki T**, **Suzuki R**, **Murakami K**, **Aizaki H**, **Ishii K**, **Murayama A**, **Date T**, **Matsuura Y**, **Miyamura T**, **Wakita T**, **Suzuki T**. 2008. Interaction of hepatitis C virus nonstructural protein 5A with core protein is critical for the production of infectious virus particles. Journal of Virology **82**:7964–7976.

7.  **Appel N**, **Zayas M**, **Miller S**, **Krijnse-Locker J**, **Schaller T**, **Friebe P**, **Kallis S**, **Engel U**, **Bartenschlager R**. 2008. Essential role of domain III of nonstructural protein 5A for hepatitis C virus infectious particle assembly. PLoS Pathog **4**:e1000035.

8.  **Matunis MJ**, **Michael WM**, **Dreyfuss G**. 1992. Characterization and primary structure of the poly(C)-binding heterogeneous nuclear ribonucleoprotein complex K protein. Mol Cell Biol **12**:164–171.

9.  **Du Z**, **Lee JK**, **Fenn S**, **Tjhen R**, **Stroud RM**, **James TL**. 2007. X-ray crystallographic and NMR studies of protein-protein and protein-nucleic acid interactions involving the KH domains from human poly(C)-binding protein-2. RNA **13**:1043–1051.

10. **Spångberg K**, **Schwartz S**. 1999. Poly(C)-binding protein interacts with the hepatitis C virus 5' untranslated region **80**:1371–1376.

11. **Flynn RA**, **Martin L**, **Spitale RC**, **Do BT**, **Sagan SM**, **Zarnegar B**, **Qu K**, **Khavari PA**, **Quake SR**, **Sarnow P**, **Chang HY**. 2015. Dissecting noncoding and pathogen RNA-protein interactomes. RNA **21**:135–143.

12. **Fan B**, **Lu K-Y**, **Reymond Sutandy FX**, **Chen Y-W**, **Konan K**, **Zhu H**, **Kao CC**, **Chen C-S**. 2014. A human proteome microarray identifies that the heterogeneous nuclear

558      ribonucleoprotein K (hnRNP K) recognizes the 5' terminal sequence of the hepatitis C
559      virus RNA. Mol Cell Proteomics **13**:84–92.

560    13.    **Poenisch M**, **Metz P**, **Blankenburg H**, **Ruggieri A**, **Lee J-Y**, **Rupp D**, **Rebhan I**,
561            **Diederich K**, **Kaderali L**, **Domingues FS**, **Albrecht M**, **Lohmann V**, **Erfle H**,
562            **Bartenschlager R**. 2015. Identification of HNRNPK as Regulator of Hepatitis C Virus
563            Particle Production. PLoS Pathog **11**:e1004573–21.

564    14.    **Wang L**, **Jeng K-S**, **Lai MMC**. 2011. Poly(C)-binding protein 2 interacts with sequences
565            required for viral replication in the hepatitis C virus (HCV) 5' untranslated region and
566            directs HCV RNA replication through circularizing the viral genome. Journal of Virology
567            **85**:7954–7964.

568    15.    **Choi K**, **Kim JH**, **Li X**, **Paek KY**, **Ha SH**, **Ryu SH**, **Wimmer E**, **Jang SK**. 2004.
569            Identification of cellular proteins enhancing activities of internal ribosomal entry sites by
570            competition with oligodeoxynucleotides. Nucleic Acids Research **32**:1308–1317.

571    16.    **Randall G**, **Panis M**, **Cooper JD**, **Tellinghuisen TL**, **Sukhodolets KE**, **Pfeffer S**,
572            **Landthaler M**, **Landgraf P**, **Kan S**, **Lindenbach BD**, **Chien M**, **Weir DB**, **Russo JJ**, **Ju**
573            **J**, **Brownstein MJ**, **Sheridan R**, **Sander C**, **Zavolan M**, **Tuschl T**, **Rice CM**. 2007.
574            Cellular cofactors affecting hepatitis C virus infection and replication. PNAS **104**:12884–
575            12889.

576    17.    **Russell RS**, **Meunier J-C**, **Takikawa S**, **Faulk K**, **Engle RE**, **Bukh J**, **Purcell RH**,
577            **Emerson SU**. 2008. Advantages of a single-cycle production assay to study cell culture-
578            adaptive mutations of hepatitis C virus. Proc Natl Acad Sci USA **105**:4370–4375.

579    18.    **Pietschmann T**, **Kaul A**, **Koutsoudakis G**, **Shavinskaya A**, **Kallis S**, **Steinmann E**,
580            **Abid K**, **Negro F**, **Dreux M**, **Cosset F-L**, **Bartenschlager R**. 2006. Construction and
581            characterization of infectious intragenotypic and intergenotypic hepatitis C virus chimeras.
582            PNAS **103**:7408–7413.

583    19.    **Panigrahi M**, **Thibault PA**, **Wilson JA**. 2021. miR-122 affects both the initiation and
584            maintenance of Hepatitis C Virus infections. bioRxiv [preprint] 1–43.

585    20.    **Amador-Cañizares Y**, **Bernier A**, **Wilson JA**, **Sagan SM**. 2018. miR-122 does not
586            impact recognition of the HCV genome by innate sensors of RNA but rather protects the
587            5′ end from the cellular pyrophosphatases, DOM3Z and DUSP11. Nucleic Acids Research
588            **46**:5139–5158.

589    21.    **Machlin ES**, **Sagan SM**, **Sarnow P**. 2011. Masking the 5' terminal nucleotides of the
590            hepatitis C virus genome by an unconventional microRNA-target RNA complex. PNAS
591            **108**:3193–3198.

592    22.    **Hsu M**, **Zhang J**, **Flint M**, **Logvinoff C**, **Cheng-Mayer C**, **Rice CM**, **McKeating JA**.
593            2003. Hepatitis C virus glycoproteins mediate pH-dependent cell entry of pseudotyped
594            retroviral particles. PNAS **100**:7271–7276.

23.   **Carroll SS**, **Tomassini JE**, **Bosserman M**, **Getty K**, **Stahlhut MW**, **Eldrup AB**, **Bhat
B**, **Hall D**, **Simcoe AL**, **LaFemina R**, **Rutkowski CA**, **Wolanski B**, **Yang Z**, **Migliaccio
G**, **De Francesco R**, **Kuo LC**, **MacCoss M**, **Olsen DB**. 2003. Inhibition of hepatitis C
virus RNA replication by 2'-modified nucleoside analogs. Journal of Biological Chemistry
**278**:11979–11984.

24.   **Gastaminza P**, **Cheng G**, **Wieland S**, **Zhong J**, **Liao W**, **Chisari FV**. 2008. Cellular
determinants of hepatitis C virus assembly, maturation, degradation, and secretion. Journal
of Virology **82**:2120–2129.

25.   **Schindelin J**, **Arganda-Carreras I**, **Frise E**, **Kaynig V**, **Longair M**, **Pietzsch T**,
**Preibisch S**, **Rueden C**, **Saalfeld S**, **Schmid B**, **Tinevez J-Y**, **White DJ**, **Hartenstein V**,
**Eliceiri K**, **Tomancak P**, **Cardona A**. 2012. Fiji: an open-source platform for biological-
image analysis. Nat Meth **9**:676–682.

26.   **Livak KJ**, **Schmittgen TD**. 2001. Analysis of Relative Gene Expression Data Using
Real-Time Quantitative PCR and the 2−ΔΔCT Method. Methods **25**:402–408.

27.   **Moradpour D**, **Evans MJ**, **Gosert R**, **Yuan Z**, **Blum HE**, **Goff SP**, **Lindenbach BD**,
**Rice CM**. 2004. Insertion of green fluorescent protein into nonstructural protein 5A
allows direct visualization of functional hepatitis C virus replication complexes. Journal of
Virology **78**:7400–7409.

28.   **Taguwa S**, **Yeh M-T**, **Rainbolt TK**, **Nayak A**, **Shao H**, **Gestwicki JE**, **Andino R**,
**Frydman J**. 2019. Zika Virus Dependence on Host Hsp70 Provides a Protective Strategy
against Infection and Disease. CellReports **26**:906–920.e3.

29.   **Boson B**, **Granio O**, **Bartenschlager R**, **Cosset F-L**. 2011. A concerted action of
hepatitis C virus p7 and nonstructural protein 2 regulates core localization at the
endoplasmic reticulum and virus assembly. PLoS Pathog **7**:e1002144.

30.   **Iwanami S**, **Kitagawa K**, **Ohashi H**, **Asai Y**, **Shionoya K**, **Saso W**, **Nishioka K**, **Inaba
H**, **Nakaoka S**, **Wakita T**, **Diekmann O**, **Iwami S**, **Watashi K**. 2020. Should a viral
genome stay in the host cell or leave? A quantitative dynamics study of how hepatitis C
virus deals with this dilemma. PLoS Biol **18**:e3000562–17.

31.   **Zhou X**, **You F**, **Chen H**, **Jiang Z**. 2012. Poly(C)-binding protein 1 (PCBP1) mediates
housekeeping degradation of mitochondrial antiviral signaling (MAVS). Cell Res **22**:717–
727.

32.   **Sumpter R**, **Loo Y-M**, **Foy E**, **Li K**, **Yoneyama M**, **Fujita T**, **Lemon SM**, **Gale M**.
2005. Regulating intracellular antiviral defense and permissiveness to hepatitis C virus
RNA replication through a cellular RNA helicase, RIG-I. Journal of Virology **79**:2689–
2699.

33.    **Meylan E**, **Curran J**, **Hofmann K**, **Moradpour D**, **Binder M**, **Bartenschlager R**,
**Tschopp J**. 2005. Cardif is an adaptor protein in the RIG-I antiviral pathway and is
targeted by hepatitis C virus. Nature **437**:1167–1172.

34.    **Bender S**, **Reuter A**, **Eberle F**, **Einhorn E**, **Binder M**, **Bartenschlager R**. 2015.
Activation of Type I and III Interferon Response by Mitochondrial and Peroxisomal
MAVS and Inhibition by Hepatitis C Virus. PLoS Pathog **11**:e1005264–30.

35.    **Walter BL**, **Parsley TB**, **Ehrenfeld E**, **Semler BL**. 2002. Distinct Poly(rC) Binding
Protein KH Domain Determinants for Poliovirus Translation Initiation and Viral RNA
Replication. Journal of Virology **76**:12008–12022.

36.    **Dinh PX**, **Beura LK**, **Panda D**, **Das A**, **Pattnaik AK**. 2011. Antagonistic effects of
cellular poly(C) binding proteins on vesicular stomatitis virus gene expression. Journal of
Virology, 3rd ed. **85**:9459–9471.

37.    **Woolaway K**, **Asai K**, **Emili A**, **Cochrane A**. 2007. hnRNP E1 and E2 have distinct roles
in modulating HIV-1 gene expression. Retrovirology **4**:28–18.

38.    **Chatel-Chaix L**, **Germain M-A**, **Motorina A**, **Bonneil É**, **Thibault P**, **Baril M**,
**Lamarre D**. 2013. A host YB-1 ribonucleoprotein complex is hijacked by hepatitis C
virus for the control of NS3-dependent particle production. Journal of Virology **87**:11704–
11720.

39.    **Gokhale NS**, **McIntyre ABR**, **McFadden MJ**, **Roder AE**, **Kennedy EM**, **Gandara JA**,
**Hopcraft SE**, **Quicke KM**, **Vazquez C**, **Willer J**, **Ilkayeva OR**, **Law BA**, **Holley CL**,
**Garcia-Blanco MA**, **Evans MJ**, **Suthar MS**, **Bradrick SS**, **Mason CE**, **Horner SM**.
2016. N6-Methyladenosine in Flaviviridae Viral RNA Genomes Regulates Infection. Cell
Host & Microbe **20**:654–665.

40.    **Rosenbluh J**, **Mercer J**, **Shrestha Y**, **Oliver R**, **Tamayo P**, **Doench JG**, **Tirosh I**,
**Piccioni F**, **Hartenian E**, **Horn H**, **Fagbami L**, **Root DE**, **Jaffe J**, **Lage K**, **Boehm JS**,
**Hahn WC**. 2016. Genetic and Proteomic Interrogation of Lower Confidence Candidate
Genes Reveals Signaling Networks in β-Catenin-Active Cancers. Cell Syst **3**:302–316.e4.

41.    **Youn J-Y**, **Dunham WH**, **Hong SJ**, **Knight JDR**, **Bashkurov M**, **Chen GI**, **Bagci H**,
**Rathod B**, **MacLeod G**, **Eng SWM**, **Angers S**, **Morris Q**, **Fabian M**, **Côté J-F**,
**Gingras A-C**. 2018. High-Density Proximity Mapping Reveals the Subcellular
Organization of mRNA-Associated Granules and Bodies. Molecular Cell **69**:517–532.e11.

42.    **Yue Y**, **Liu J**, **Cui X**, **Cao J**, **Luo G**, **Zhang Z**, **Cheng T**, **Gao M**, **Shu X**, **Ma H**, **Wang**
**F**, **Wang X**, **Shen B**, **Wang Y**, **Feng X**, **He C**, **Liu J**. 2018. VIRMA mediates preferential
m(6)A mRNA methylation in 3'UTR and near stop codon and associates with alternative
polyadenylation. Cell Discov **4**:10.

 **SUPPLEMENTARY INFORMATION**

**SUPPLEMENTARY METHODS**

**Plasmids**

To generate the pPCBP1(N)FLAG-puro plasmid, the complete PCBP1 coding sequence was

amplified using PCBP1noAUG-KpnI-FW (5'-AAG CTG GCG GTA CCG GAG ATG CCG

GTG TGA CTG-3') and PCBP1-XhoI-RV (5'-TTT TTC TC GAG CTA GCT GCA CCC CAT

GCC CTT -3') and the high-fidelity Hot Start Q5 DNA polymerase (NEB). The PCR product

was digested with KpnI and XhoI and ligated in a pcDNA3.1(+)-derived plasmid that contained

a 3xFLAG tag located downstream of a human cytomegalovirus (CMV) immediate early

enhancer/promoter and upstream of a bovine growth hormone (bGH) polyadenylation signal; this

insertion left the PCBP1 coding sequence in frame with the upstream 3x FLAG tag. To note, this

pcDNA3.1(+)-derived vector also contained a puromycin resistance gene located downstream of

a phosphoglycerate kinase 1 (PGK) promoter, and upstream of a simian virus 40 (SV40)

polyadenylation signal.

To generate the pFLuc-puro plasmid, the complete Firefly luciferase coding sequence

was amplified using FLuc-NheI-FW (5'-ATC CGC TAG CAT GGA TTA CAA GGA C-3') and

FLuc-XhoI-RV (5'-AAG GTA TCT CGA GTT AGT AAA CAA GAT AAT TGC TCC TAA

AGT A-3') using the high-fidelity Hot Start Q5 DNA polymerase (NEB). The PCR product was

digested with NheI and XhoI and ligated into the same pcDNA3.1(+)-derived plasmid, between

the CMV enhancer/promoter and the bGH polyadenylation signal; this insertion removed the

3xFLAG tag.

The pJ6/JFH1 FL RLuc WT ("RLuc-wt") viral sequence contains include a *Renilla* luciferase

(RLuc) reporter gene inserted between p7 and NS2 (1). The pJ6/JFH-1 FL RLuc-NS5A-GFP

("NS5A-GFP") plasmid contains a GFP insertion between P2390 and L2391 within the NS5A

domain III, as previously described (2). The NS5A-GFP region was subcloned into the pJ6/JFH-

1 FL RLuc WT backbone using the *AvrII* and *XbaI* restriction sites.

To construct a JFH-1$_T$-RLuc viral genome template, a MluI site was created between the

p7 and NS2 genes by PCR amplification of the pJFH-1$_T$ plasmid using JFH1-p7NS2-MluI-FW

(5'-CCT ATG ACG CGT CTG TGC ACG GAC AGA TAG GC-3'), JFH1-p7NS2-MluI-RV (5'-

CAG ACG CGT CAT AGG CAT AAG CCT GCC G-3'), and the high-fidelity Hot Start Q5

DNA polymerase (NEB). The PCR product was digested with MluI and dephosphorylated with

calf intestinal phosphatase (Quick CIP, NEB) and ligated with an insert containing a *Renilla*

luciferase (RLuc) with a C-terminal foot-and-mouth disease virus 2A peptide (FMDV2A), which

had been cut out of the pJ6/JFH WT RLuc plasmid by MluI digestion. The correct orientation of

the insert was screened by digestion with BstBI and NotI and by sequencing. Once a plasmid

with the correct insert orientation was identified, the E1 through NS2 region was cut out using

the BsiWI and NotI unique cut sites and ligated into the original pJFH-1$_T$ plasmid backbone. The

inserted E1 through NS2 region was sequenced to verify that the PCR amplifications had not

introduced any unexpected substitutions.

**Puromycin selection of cells and infection with JFH-1$_T$**

To select cells that would stably express PCBP1(N)FLAG or the FLuc control protein, 15-cm

dishes were seeded with 2 x $10^6$ Huh-7.5 cells one day prior to transfection with 36 µg of

pPCBP1(N)FLAG-puro or pFLuc-puro plasmid and 30 µL Lipofectamine 2000 diluted in 12 mL

Opti-MEM serum-free media (Thermo Fisher Scientific). Two days post-transfection, the media

on all transfected plates and one non-transfected plate was changed for complete Huh-7.5 media

supplemented with 3 µg/mL puromycin (Thermo Fisher Scientific), which was replenished every

two to three days, passaging the cells as necessary. Once all cells in the non-transfected plate

were completely dead (within 7 days of selection), selected cells could be used for infection

experiments. Selected cells were transfected with siRNA duplexes and infected with JFH-1$_T$ at

an MOI of 0.05 as described in the article's main methods; the selection pressure of 3 µg/mL

puromycin was maintained in these cells' media until their infection with JFH-1$_T$, after which

they were kept in puromycin-free media.

**REFERENCES**

1.  **Pietschmann T**, **Kaul A**, **Koutsoudakis G**, **Shavinskaya A**, **Kallis S**, **Steinmann E**, **Abid**
**K**, **Negro F**, **Dreux M**, **Cosset F-L**, **Bartenschlager R**. 2006. Construction and
characterization of infectious intragenotypic and intergenotypic hepatitis C virus chimeras.
PNAS **103**:7408–7413.
725

726  2.  **Moradpour D**, **Evans MJ**, **Gosert R**, **Yuan Z**, **Blum HE**, **Goff SP**, **Lindenbach BD**, **Rice**
727      **CM**. 2004. Insertion of green fluorescent protein into nonstructural protein 5A allows direct
728      visualization of functional hepatitis C virus replication complexes. Journal of Virology
729      **78**:7400–7409.
730

**Supplementary Figure S1. PCBP1 knockdown increases the total quantity of infectious**

**virions produced by the JFH-1$_T$ and J6/JFH-1 strains.** To calculate the total quantity of

infectious virions (FFU) present in a sample, the viral titer (FFU/mL) was multiplied by the total

volume of the sample (mL). These data were derived from the same experiments presented in

Figure 1 and are representative of three independent biological replicates. Error bars represent

the standard deviation of the mean.

**Supplementary Figure S2. Ectopic PCBP1 expression can reduce the impact of endogenous**

**PCBP1 knockdown on extracellular viral titers.** Huh-7.5 cells that were puromycin-selected

to stably express ectopic Firefly luciferase (FLuc, control) or PCBP1 with a N-terminal FLAG

tag (PCBP1(N)FLAG) were transfected with siCTRL or siPCBP1 two days prior to infection

with JFH-1$_T$ (MOI 0.05). Three days post-infection, total intracellular protein, RNA, and

intracellular and extracellular virus were collected. **(A)** Viral protein expression analysis by

Western blot. **(B)** Viral RNA accumulation analysis by Northern blot and **(C)** quantification by

RT-qPCR. **(D)** Intracellular and **(E)** extracellular (secreted) virus titers, quantified by FFU assay.

For (C-E), data from each experiment was normalized to the mean of the cell-matched siCTRL

condition before replicate data were combined. All data are representative of three independent

biological replicates, with the exception of (B) which is an $N = 1$. Error bars represent the

standard deviation of the mean. Statistical significance was calculated by paired t-test (ns, not

significant; *** $p < 0.001$; **** $p < 0.0001$).

**Supplementary Figure S3. PCBP1 has no effect on RNA replication in the absence of efficient virion assembly, and the addition of a RLuc gene reduces the assembly efficiency of HCV genomes.** SiRNA-transfected Huh-7.5 cells were electroporated with 5 μg of (**A**) full-length J6/JFH RLuc WT RNA, or of (**B**) a full-length J6/JFH RLuc with a GFP insertion in the NS5A gene (NS5A-GFP), which had previously been shown to impair virion assembly without impairing viral RNA replication. Luciferase activity was monitored for three days post-electroporation, and RLuc values were normalized to the early timepoint (3h) to control for disparities in electroporation efficiency between experiments. (**C**) Equal amounts (10 μg) of JFH-1$_T$, JFH-1$_T$-RLuc, or J6/JFH-RLuc RNAs were electroporated into Huh-7.5 cells; intracellular and extracellular viruses were collected three days post-electroporation and titered by focus-forming unit assay. Compared with the untagged JFH-1$_T$ RNA electroporations, the extracellular titers were reduced by over 100-fold and 1000-fold for JFH1$_T$-RLuc and J6/JFH-

765    RLuc, respectively; intracellular titers were reduced by more than 200-fold for both luciferase

766    reporter genomes. Data in (A) and (B) are representative of three independent biological

767    replicates; data in (C) is representative of two independent biological replicates, and the limit of

768    detection is indicated. Error bars represent the standard deviation of the mean. Statistical

769    significance was calculated by t-test (** $p < 0.005$, **** $p < 0.0001$)