1

Poly(rC)-binding protein 1 limits hepatitis C virus assembly and egress

Sophie E. Cousineau[1] and Selena M. Sagan[1,2]#

[1]Department of Microbiology & Immunology, McGill University, Montreal, QC, Canada

[2]Department of Biochemistry, McGill University, Montreal, QC, Canada

#Corresponding Author: Dr. Selena Sagan, McGill University, 3655 Promenade Sir William Osler, Room 805, Montréal, QC, H3G 1Y6; email: selena.sagan@mcgill.ca

Keywords: Hepatitis C virus (HCV), poly(rC)-binding protein 1 (PCBP1), hnRNP E1, assembly, egress

Running title: PCBP1 limits HCV assembly and egress

ABSTRACT

The hepatitis C virus (HCV) co-opts a number of cellular elements – including proteins, lipids, and microRNAs – to complete its viral life cycle. The cellular RNA-binding protein poly(rC)-binding protein 1 (PCBP1) had previously been reported to bind the HCV genome 5′ untranslated region (UTR), but its importance in the viral life cycle has remained unclear. Herein, we aimed to clarify the role of PCBP1 in the HCV life cycle. Using the HCV cell culture (HCVcc) system, we found that endogenous PCBP1 knockdown decreased viral RNA accumulation yet increased extracellular virus titers. To dissect PCBP1's specific role in the viral life cycle, we carried out assays for viral entry, translation, genome stability, RNA replication, virion assembly and egress. We found that PCBP1 did not affect viral entry, translation, RNA stability, or RNA replication in the absence of efficient virion assembly. To specifically examine virion assembly and egress, we inhibited viral RNA replication with an RNA-dependent RNA polymerase inhibitor and tracked both intracellular and extracellular viral titers over time. We found that when viral RNA accumulation was inhibited, knockdown of PCBP1 still resulted in an overall increase in HCV particle secretion. We therefore propose a model where endogenous PCBP1 limits virion assembly and egress, thereby indirectly enhancing viral RNA accumulation in infected cells. This model furthers our understanding of how cellular RNA-binding proteins modulate HCV genomic RNA utilization during the viral life cycle.

IMPORTANCE

Hepatitis C virus (HCV) is a positive-sense RNA virus, and as such, its genome must be a template for multiple mutually exclusive steps of the viral life cycle, namely translation, RNA replication, and virion assembly. However, the mechanism(s) that regulate how the viral genome

is used throughout the viral life cycle still remain unclear. A cellular RNA-binding protein –

PCBP1 – had previously been reported to bind the HCV genome, but its precise role in the viral

life cycle was not known. In this study, we found that depleting PCBP1 decreased viral RNA

accumulation but increased virus secretion. We ruled out a role for PCBP1 in virus entry,

translation, genome stability or RNA replication, and demonstrate that PCBP1 knockdown

enhances virus secretion when RNA replication is inhibited. We conclude that PCBP1 normally

prevents virus assembly and egress, which allows more of the viral genomic RNA to be available

for translation and viral RNA replication.

INTRODUCTION

The hepatitis C virus (HCV) is an enveloped virus of the *Flaviviridae* family (genus:

hepacivirus) that typically causes a persistent liver infection (1). Its ~9.6 kb single-stranded,

positive-sense RNA genome contains a single open reading frame flanked by 5′ and 3′

untranslated regions (UTR). A highly structured, type 3 internal ribosomal entry site (IRES) in

the 5′ UTR drives the translation of the viral polyprotein, which is subsequently processed into

10 mature viral proteins: 3 structural proteins (core, E1 and E2 glycoproteins), and 7

nonstructural (NS) proteins (p7, NS2, NS3, NS4A, NS4B, NS5A and NS5B) (2, 3). While the

structural proteins form the nucleocapsid and viral envelope, the NS3-5B form the viral

replicase, required for viral RNA replication (4). The p7, NS2, NS3 and NS5A proteins have also

been implicated in viral genome packaging and the assembly of new virion particles (5-8). As a

positive-sense RNA virus, the HCV genome itself must serve as a template for viral translation,

genome replication, and packaging; however, the mechanisms that determine which process each

viral RNA is engaged in at any given time have not been defined. Due to their limited coding

capacity, viruses are highly dependent on the molecular machinery of the host cell; thus, it is

likely that cellular components participate in regulation of the viral RNA during the HCV life

cycle. While a number of cellular proteins and RNAs have been shown to interact with the HCV

genome, their precise role(s) in the viral life cycle have yet to be defined.

The poly(rC)-binding protein 1 (PCBP1) is one of the three most abundant cellular RNA-

binding proteins with a strong affinity for poly(rC), along with its paralogs hnRNP K and PCBP2

(9). These multifunctional proteins can regulate translation and enhance the stability of their

cellular mRNA targets, which they interact with through their hnRNP K homologous (KH)

domains (10). Notably, all three paralogs have been reported to bind to the HCV 5′ UTR (11-

13). However, the degree to which each protein has been studied in the context of HCV infection

varies significantly – while hnRNP K and PCBP2 have been fairly extensively studied and

reported to play markedly different roles in the viral life cycle, the role of PCBP1 in the HCV

life cycle has not been investigated in detail (14, 15). Beyond its interactions with the 5´ UTR,

previous reports suggested that PCBP1 was not necessary for HCV IRES-mediated translation,

but that knockdown of PCBP1 decreases HCV RNA accumulation during infection (16, 17).

Herein, we sought to clarify the role of PCBP1 in the HCV life cycle. Using a cell

culture-adapted strain of HCV, we found that PCBP1 knockdown decreased viral RNA

accumulation, yet led to an increase in virus secretion. By examining individual steps of the viral

life cycle, we ruled out a role for PCBP1 in viral entry, translation, genome stability and viral

RNA replication. Further analysis of the assembly step revealed that, similarly to its paralog

hnRNP K, endogenous PCBP1 limits HCV virion assembly and release.

MATERIALS AND METHODS

Cell culture

Huh-7.5 human hepatoma cells were obtained from Charlie Rice (Rockefeller University) and

maintained in complete media: Dulbecco's Modified Eagle Media (DMEM) supplemented with

inactivated 10% fetal bovine serum (FBS), 2 mM L-glutamine, and 1X MEM non-essential

amino acids. Human embryonic kidney (293T) cells were kindly provided by Martin J. Richer

(McGill University, Montreal, QC, Canada) and were maintained in DMEM supplemented with

10% FBS. All cell lines were maintained at 37°C/5% CO_2 and were routinely screened for

mycoplasma contamination.

Plasmids and viral RNAs

The pJFH-1$_T$ plasmid, encoding a cell culture-adapted Japanese Fulminant Hepatitis (JFH-1; HCV genotype 2a) with three adaptive mutations that increase viral titers in cell culture, was a gift from Rodney Russell (Memorial University of Newfoundland) (18). Plasmids pJ6/JFH1 FL RLuc WT ("RLuc-wt") and pJ6/JFH-1 FL RLuc GNN ("RLuc-GNN") bear full-length viral sequence derived from the J6 (structural genes) and JFH-1 (NS genes) isolates of HCV, with a *Renilla* luciferase (RLuc) reporter (5). The pJ6/JFH-1 mono RLuc-NS2 plasmid ("Δcore-p7") – a truncated version of the *Renilla* reporter virus with a deletion of the structural genes through p7 – was a gift from Joyce Wilson (University of Saskatchewan) (19). The pJ6/JFH-1 FL RLuc-NS5A-GFP ("NS5A-GFP") was created via overlapping PCR and subcloned using the *AvrII* and *XbaI* restriction sites, as previously described (20).

To make full-length uncapped viral RNAs, all plasmid templates were linearized and *in vitro* transcribed as previously described (21). The firefly luciferase (FLuc) mRNA was transcribed from the Luciferase T7 Control DNA plasmid (Promega) linearized using *XmnI* and *in vitro* transcribed using the mMessage mMachine T7 Kit (Life Technologies) according to the manufacturer's instructions.

Generation of infectious HCV stocks

To generate viral stocks, 30 µg of *in vitro* transcribed JFH-1$_T$ RNA was transfected into Huh-7.5 cells using the DMRIE-C reagent (Life Technologies) according to the manufacturer's instructions. Four days post-transfection, infectious cell supernatants were passed through a 0.45 µm filter and infectious viral titers were determined by focus-forming unit assay (18). Infectious

virus was amplified for two passages through Huh-7.5 cells at a MOI of 0.1. Viral stocks were

aliquoted and stored at -80°C until use.

Focus-forming unit (FFU) assays

One day prior to infection, 8-well chamber slides (Lab-Tek) were seeded with 4 x 10^5 Huh-7.5

cells/well. Infections were performed with 10-fold serial dilutions of viral samples in 100 μL for

4 h, after which the supernatant was replaced with fresh media. Three days post-infection, slides

were fixed in 100% acetone and stained with anti-HCV core antibody (1:100, clone B2,

Anogen), and subsequently with the AlexaFluor-488-conjugated anti-mouse antibody (1:200,

ThermoFisher Scientific) for immunofluorescence analysis. Viral titers are expressed as the

number of focus-forming units (FFU) per mL.

Extracellular virus titers were determined directly from cell supernatants, while

intracellular virus titers were determined after cell pellets were subjected to lysis via four freeze-

thaw cycles, removal of cellular debris via centrifugation, and recovery of virus-containing

supernatants.

MicroRNAs and siRNA sequences

siGL3 (siCTRL): 5'-CUUACGCUGAGUACUUCGAUU-3', siGL3* : 5'-

UCGAAGUACUCAGCGUAAGUU-3', miR122$_{p2-8}$ (siCTRL for luciferase experiments): 5'-

UAAUCACAGACAAUGGUGUUUGU-3', miR122$_{p2-8}$*: 5'-

AAACGCCAUUAUCUGUGAGGAUA-3' (22), siPCBP1: 5'-

CUGUGUAAUUUCUGGUCAGUU-3', siPCBP1*: 5'-CUGACCAGAAAUUACACAGUU-3'

(17) were all synthesized by Integrated DNA Technologies.

All microRNA and siRNA duplexes were diluted to a final concentration of 20 µM in

RNA annealing buffer (150 mM HEPES pH 7.4, 500 mM potassium acetate, 10 mM magnesium

acetate), and annealed at 37°C for 1 h and stored at -20°C. For all knockdown experiments, 50

nM siRNA transfections were conducted 2 days prior to infection or electroporation of viral

RNAs. Transfections were conducted using the Lipofectamine RNAiMAX (Invitrogen)

according to the manufacturer's instructions with the modification that 20 µL of reagent were

used to transfect a 10-cm dish of cells.

HCV and VSV pseudoparticles (HCVpp and VSVpp)

HCVpp consisting of a FLuc reporter lentiviral vector pseudotyped with the HCV E1 and E2

glycoprotein (from the H77, a genotype 1a strain) were a kind gift from John Law (University of

Alberta) (23). To generate lentiviral vectors pseudotyped with the VSV-G glycoprotein

(VSVpp), a 90% confluent 10-cm dish of 293T cells were transfected with 10 µg pPRIME-FLuc,

5 µg psPAX.2, and 2.5 µg pVSV-G plasmid with 10 µL Lipofectamine 2000 (Invitrogen) diluted

in 4 mL Opti-MEM. Media was changed 4, 20, and 28 h post-transfection. At 48 h post-

transfection, the cell culture media was passed through a 0.45 µm filter and stored at -80°C.

To assay for cell entry, HCVpp and VSVpp were diluted 1/3 in dilution media (1X

DMEM, 3% FBS, 100 IU penicillin and 100 µg/mL streptomycin) with 20 mM HEPES and 4

µg/µL polybrene, and then introduced to Huh-7.5 cells by spinoculation at 1,200 rpm for 1 h at

room temperature. The cells were left to recover at 37°C for at least 5 h before the

pseudoparticle-containing media was changed for fresh complete Huh-7.5 media. In parallel,

cells seeded in a 6-well plate were transfected with 1 µg of pPRIME-FLuc plasmid using

Lipofectamine 2000 (Invitrogen) according to the manufacturer's instructions. Three days post-

spinoculation and transfection, cells were lysed in passive lysis buffer (Promega) and FLuc

activity was assayed using the Dual Reporter Luciferase kit (Promega).

Electroporations

For each electroporation, 400 µL of resuspended cells (1.5 x 10^7 cell/mL) were mixed with 2 µg

of FLuc mRNA and 5 µg of replicating (WT, Δcore-p7 or NS5A-GFP J6/JFH-1 RNA) or 10 µg

GNN J6/JFH-1 RNA, and electroporated in 4-mm cuvettes at 270 V, 950 µF, and infinite

resistance, optimized for the Bio-Rad GenePulser XCell (Bio-Rad). Electroporated cells were

resuspended in complete Huh-7.5 media and transferred to 6-well plates for luciferase assays and

protein analysis.

Inhibition of RNA replication by 2´CMA

Two days post-siRNA transfection, Huh-7.5 cells were infected with JFH-1$_T$ at a MOI of 0.05.

Four to five hours post-infection, each plate of infected cells was split into 6-well plates. Three

days post-infection, the media on these cells was changed for complete Huh-7.5 media with 5

µM 2´CMA (2´C-methyladenosine, Carbosynth), an HCV NS5B polymerase inhibitor, or DMSO

control (24). Total RNA and intracellular virus samples were collected at 0, 6 and 12 h post-

treatment, while cell culture supernatants were collected 6 and 12 h post-treatment. Protein

samples were collected from untreated plates to assess PCBP1 knockdown efficiency by Western

blot.

Western blot analysis

To collect total intracellular protein samples, cells were lysed in RIPA buffer (150 mM sodium chloride, 1% NP-40, 0.5% sodium deoxycholate, 0.1% SDS, 50 mM Tris pH 8.0, Complete Protease Inhibitor Cocktail (Roche)) and frozen at -80°C. Cellular debris was pelleted by centrifugation at 16,000 x g for 30 min at 4°C, and the supernatant was quantified by BCA Protein Assay (ThermoScientific). Ten micrograms of sample were loaded onto 10-12% SDS-PAGE gels. Samples were transferred onto Immobilon-P PVDF membranes (Millipore), blocked in 5% milk, and incubated overnight with primary antibodies diluted in 5% BSA: rabbit anti-PCBP1 (clone EPR11055, Abcam ab168378, 1:10,000); rabbit anti-actin (A2066, Sigma, 1:20,000); mouse anti-HCV core (clone B2, Anogen MO-I40015B, diluted 1:7,500); mouse anti-JFH-1 NS5A (clone 7B5, BioFront Technologies, 1:10,000). Blots were incubated for 1 hour with HRP-conjugated secondary antibodies diluted in 5% skim milk: anti-mouse (HAF007, R&D Systems, 1:25,000); anti-rabbit (111-035-144, Jackson ImmunoResearch Laboratories, 1:50,000) and visualized using enhanced chemiluminescence (ECL Prime Western Blotting Detection Reagent, Fisher Scientific).

RNA isolation and Northern blot analysis

Total RNA was harvested using TRIzol Reagent (ThermoFisher Scientific) according to the manufacturer's instructions. Ten micrograms of total RNA were separated on a 1% agarose gel containing 1X 3-(N-morpholino)propanesulfonic acid (MOPS) buffer and 2.2 M formaldehyde and transferred to a Zeta-probe membrane (Bio-Rad) by capillary transfer in 20X SSC buffer (3 M NaCl, 0.3 M sodium citrate). Membranes were hybridized in ExpressHyb Hybridization Buffer (ClonTech) to random-primed [32]P-labeled DNA probes (RadPrime DNA labelling

system, Life Technologies) complementary to HCV (nt 84-374) and γ-actin (nt 685-1171).

Autoradiograph band densities were quantified using Fiji (25).

RT-qPCR analysis

The iTaq Universal Probes One-Step kit (Bio-Rad) was used to perform duplex assays probing

for the HCV genome (NS5B-FW primer: 5'-AGACACTCCCCTATCAATTCATGGC-3';

NS5B-RV primer: 5'-GCGTCAAGCCCGTGTAACC-3'; NS5B-FAM probe: 5'-

ATGGGTTCGCATGGTCCTAATGACACAC-3') and the GAPDH loading control (PrimePCR

Probe assay with HEX probe, Bio-Rad). Each 20 µL reaction contained 500 ng of total RNA, 1.5

µL of the HCV primers and probe, and 0.5 µL of the GAPDH primers and probe. RT-PCR

reactions were conducted in a CFX96 Touch Deep Well Real-Time PCR system (Bio-Rad).

Genome copies were calculated using a standard curve and fold-differences in gene expression

were calculated using the $2^{-\Delta\Delta Ct}$ method (26).

Luciferase assays

For translation and replication assays, cells were washed in PBS and harvested in 100 µL of 1X

passive lysis buffer (Promega). The Dual-Luciferase Assay Reporter Kit (Promega) was used to

measure both *Renilla* and firefly luciferase activity according to the manufacturer's instructions

with the modification that 25 µL of reagent were used with 10 µL of sample. All samples were

measured in triplicate.

Data analysis

Statistical analyses were performed using GraphPad Prism v9 (GraphPad, USA). Statistical

significance was determined by paired t-test to compare results obtained from multiple

experiments, and by two-way ANOVA with Geisser-Greenhouse and Bonferroni corrections

when more than two comparisons were applied at once. To calculate half-lives, a one-step decay

curve using the least-squares regression was used, and error was reported as the asymmetrical

(profile likelihood) 95% confidence interval of the half-life. To calculate virus accumulation and

virus secretion rates, a simple linear regression was performed using the least squares regression

method. The slope and standard error calculated for each regression represents the rate of virus

accumulation or secretion.

RESULTS

PCBP1 plays a role in the HCV life cycle

The PCBP1 protein was previously reported to directly interact with the 5′ UTR of the HCV

genome, and in an siRNA screen PCBP1 knockdown resulted in a reduction in HCV RNA

accumulation (11, 17). Thus, we sought to further characterize the role of PCBP1 in the HCV life

cycle. We began by assessing how PCBP1 knockdown affected the accumulation of cell culture-

derived HCV (HCVcc), using the cell-culture adapted JFH-1$_T$ strain. Compared to the parental

JFH-1 strain, JFH-1$_T$ has three adaptive mutations in the E2, p7 and NS2 coding-region, which

enable it to produce higher viral titers in cell culture (18). We found that knockdown of

endogenous PCBP1 resulted in an approximately 2.2-fold decrease in viral protein and RNA

accumulation in Huh-7.5 cells (**Figure 1A-C and Supplementary Figure 1**). Interestingly,

when we quantified intracellular and extracellular virions, we found that while intracellular titers

were not significantly different between the PCBP1 knockdown and control conditions,

extracellular titers were elevated, with an average increase of approximately 3.90-fold in the

PCBP1 knockdown condition (**Figure D-E**). Thus, in line with previous findings, we found that

PCBP1 knockdown decreased viral protein expression and intracellular viral RNA accumulation.

Yet, despite this overall decrease, we observed an increase in extracellular (secreted) virus titers.

These results imply that endogenous PCBP1 indeed plays a role in the HCV life cycle, although

the precise step(s) influenced by PCBP1 remain unclear.

PCBP1 knockdown has no impact on HCV entry

Firstly, we explored whether PCBP1 knockdown had any effect on virus entry. To this end, we

made use of the HCV pseudoparticle (HCVpp) system, which consists of lentiviral vectors with a

firefly luciferase reporter gene pseudotyped with the HCV E1 and E2 glycoproteins (23). HCVpp

enter cells by clathrin-mediated endocytosis after engaging with HCV-specific entry receptors;

thus, to account for any changes in clathrin-mediated endocytosis, we used a vesicular stomatitis

virus (VSV) pseudoparticle system (VSVpp) as a control. In addition, to verify that PCBP1

knockdown did not affect luciferase reporter gene expression, we assessed firefly luciferase

expression from cells directly transfected with a FLuc reporter plasmid. In all cases, we found

that depleting endogenous PCBP1 had no impact on luciferase activity (**Figure 2**). This suggests

that PCBP1 knockdown does not affect FLuc reporter expression, clathrin-mediated endocytosis,

or the HCV entry process.

PCBP1 knockdown has no impact of HCV translation or genome stability

PCBP1 was previously reported to bind the HCV 5′ UTR, which contains the viral IRES that

drives translation of the viral polyprotein in the absence of most canonical translation initiation

factors (2, 3, 11, 27). Furthermore, PCBP1 has been reported to contribute to the IRES-mediated

translation of some cellular mRNAs such as Bag-1 and *c-myc* (28-30). Thus, it was plausible that

PCBP1's interactions with the viral 5′ UTR could affect viral protein expression by altering

IRES-mediated translation. To assess PCBP1's impact on HCV translation, we used full-length

J6/JFH-1 RLuc reporter RNAs containing an inactivating mutation in the NS5B polymerase gene

(GNN). The RLuc activity thus served as a direct measure of HCV IRES-mediated translation,

and over time, this signal also served as a proxy measure for viral RNA stability. We found that

the siPCBP1 and siCTRL conditions had similar RLuc activity at all timepoints (**Figure 3**).

Moreover, the luciferase signal half-lives were nearly identical, with 2.68 h (95% CI 1.64 – 4.64)

for the siPCBP1 condition and 2.71 h (95% CI 1.43 – 5.50) for the siCTRL condition. Thus,

PCBP1 knockdown does not appear to affect either viral IRES-mediated translation or genome

stability.

PCBP1 knockdown does not affect viral RNA replication

PCBP1 has been reported to promote viral RNA replication by binding to the 5′ UTR of several

RNA viruses (31, 32). Thus, to specifically assess whether PCBP1 knockdown has an effect on

HCV RNA replication, we assessed RLuc expression of replication-competent but assembly-

deficient reporter RNAs (**Figure 4**). This includes the full-length WT J6/JFH-1 RLuc RNA, a

subgenomic J6/JFH-1 replicon RNA containing a deletion of the structural protein genes (Δcore-

p7), and a full-length J6/JFH-1 genome with a GFP insertion in the NS5A gene (NS5A-GFP),

the latter previously shown to impair virion assembly without impairing viral RNA replication

(7, 20). In all cases, we did not observe any significant differences in viral RNA accumulation

between the siPCBP1 and siCTRL conditions (**Figure 4A-C**). This is seemingly in contrast to

our findings using the JFH-1$_T$ system, where PCBP1 depletion decreased viral RNA

accumulation (**Figure 1B-C and Supplementary Figure 1**). However, it is notable that all of the

viral RNAs used in **Figure 4** are defective in viral packaging, including the full-length WT

J6/JFH-1 RLuc RNA (**Figure 4A**), likely due to the large RLuc reporter gene insertion

(**Supplementary Figure 2**). Thus, taken together, our data suggests that PCBP1 has no effect on

the viral RNA replication step of the HCV life cycle.

PCBP1 knockdown limits virus egress

Finally, since knockdown of PCBP1 did not seem to have a significant impact on HCV entry,

translation, genome stability or viral RNA replication, we reasoned that PCBP1 was likely to be

exerting an effect on virion assembly and/or secretion. We had previously established that

PCBP1 knockdown did not have a significant effect on intracellular viral titers yet resulted in an

overall increase in extracellular (secreted) virions (**Figure 1D-E**). Thus, to further investigate

whether PCBP1 plays a role in viral packaging and/or egress, we used a nucleoside analog, 2´C-

methyladenosine (2´CMA), to block viral RNA synthesis by the HCV NS5B RNA-dependent

RNA polymerase and monitored viral titers over time (24). Similar to related studies using Zika

virus, we reasoned that blocking viral RNA synthesis could allow us to assess viral packaging

and egress in the absence of genomic RNA production (33).

To this end, we infected cells with JFH-1$_T$ and, three days post-infection, we replaced the

culture media with media containing 5 μM 2′CMA (or DMSO, as a control) and collected total

RNA, intracellular virions and extracellular virions over the next 12 h (**Figure 5A**). In agreement

with our previous findings, we observed an overall reduction in JFH-1$_T$ viral RNA accumulation

by day 3 post-infection in the siPCBP1 condition (0 h timepoint, **Figure 5B**). In addition, the

2′CMA treatment efficiently blocked viral RNA accumulation, which continued to increase

under the control (DMSO) condition (6-12 h timepoints, **Figure 5B**). Additionally, we observed

similar initial intracellular titers as well as similar 2′CMA-induced decreases in intracellular

titers under both the siPCBP1 and siCTRL conditions (**Figure 5C and D**). In contrast, we

observed a continued rise in extracellular titers as packaged viruses continued to egress out of the

cell, with the PCBP1 knockdown resulting in a >3-fold greater virus secretion rate than the

siCTRL condition during 2′CMA treatment (**Figure 5E and F**). Notably, since inhibiting viral

RNA synthesis equalized the intracellular virus accumulation rates yet failed to equalize the

virus secretion rates, these results suggest that PCBP1 modulates the virion assembly and egress

steps of the HCV life cycle.

DISCUSSION

Herein, we investigated the role of PCBP1 in the HCV life cycle. We found that PCBP1

knockdown did not directly affect virus entry, translation, genome stability or viral RNA

replication; but resulted in an increase in infectious particle secretion. Furthermore, since PCBP1

knockdown does not alter intracellular infectious particle accumulation yet reduces total

intracellular viral RNA, our results suggest that PCBP1 knockdown promotes virion assembly

and egress.

Previous studies had found that PCBP1 interacts with the HCV 5′ UTR *in vitro* (11, 16).

Notably, the full 5′ UTR appeared to be critical for PCBP1 binding, as truncated fragments of

the 5′ UTR were unable to bind to PCBP1 (11, 16). Somewhat more recently, PCBP1 was one of

many host factors evaluated in a siRNA screen targeting cellular proteins predicted to interact

with HCV (17). This screen revealed that PCBP1 knockdown decreased viral RNA accumulation

approximately 2.3-fold, a result consistent with our observation that PCBP1 knockdown

decreased JFH-1$_T$ viral RNA accumulation by approximately 2.2-fold (**Figure 1D**) (17).

However, our systematic evaluation of the effect of PCBP1 knockdown on viral translation and

viral RNA replication revealed that PCBP1 does not modulate either of these steps in the viral

life cycle. Moreover, we did not observe any significant effects on viral entry or genome

stability.

Initially, we were quite puzzled to find that PCBP1 did not affect viral RNA accumulation

of a full-length WT J6/JFH-1 RLuc RNA; however, this RNA generates far fewer infectious

viral particles than the JFH-1$_T$ strain we used for our infection experiments, likely due to the

increased genome size due to the RLuc insertion resulting in reduced packaging efficiency

(**Supplementary Figure S2**) (14). The fact that PCBP1 knockdown failed to have a significant

impact on the luciferase reporter RNA is therefore consistent with a model whereby PCBP1

primarily acts at the level of virion assembly and egress (**Figure 6**). As such, knockdown of

PCBP1 results in an overall decrease in the translating/replicating pool of viral RNAs and a

concomitant increase in extracellular (secreted) virions. Since intracellular viral titers remained

similar in PCBP1 knockdown and control conditions, a greater proportion of the intracellular

viral RNA must represent viral RNA packaged into virions, implying that endogenous PCBP1

normally limits virion assembly. This model is further supported by our finding that 2′CMA

equalized the rate of intracellular virus accumulation yet resulted in a virus secretion rate >3-fold

greater during PCBP1 knockdown, suggesting that endogenous PCBP1 limits virion egress.

Moreover, this model is consistent with the observed decreases in viral protein expression and

negative-strand replicative intermediate accumulation (**Figure 1B and Supplementary Figure**

1), since increasing viral genome packaging into new virions would sequester the RNA from the

translation and RNA replication machinery. Thus, endogenous PCBP1 normally limits viral

assembly and egress, and PCBP1 knockdown therefore indirectly impairs viral protein synthesis

and viral RNA replication by liberating the RNA for virion assembly and egress.

Notably, PCBP1 has been previously implicated in turnover of MAVS, a signal

transduction protein directly downstream of RIG-I, which is known to limit HCV infection (34).

However, all of our conclusions were drawn from experiments conducted in the Huh-7.5 cell

line, which has well-documented defects in the RIG-I antiviral signalling pathway, and our own

brief explorations of interferon induction and MAVS turnover did not reveal any significant

differences between PCBP1 knockdown and control conditions (data not shown) (35). Moreover,

the HCV NS3-4A protease also inactivates MAVS during infection to reduce antiviral signaling

and recognition of the viral RNA (36, 37). However, as hepatocytes typically express RIG-I and

MAVS, we cannot rule out the possibility that PCBP1 may modulate this pathway during

infection *in vivo*.

Finally, PCBP1's closely related paralogs, hnRNP K and PCBP2, have been more

extensively studied during HCV infection. While the hnRNP K protein was shown to restrict

infectious virion production, the PCBP2 protein has been suggested to modulate viral translation

and RNA replication (14, 15). Since the PCBP1 amino acid identity is far more similar to PCBP2

(~80% identity) than hnRNP K (~24% identity), we were initially surprised that our findings for

PCBP1 in HCV infection closely matched those reported for hnRNP K. Yet, while PCBP1 and

PCBP2 have been shown to perform similar functions, they have also been reported to play

distinct roles during poliovirus, VSV, and human immunodeficiency virus infection (31, 38, 39).

In addition to the similarities we observed with PCBP1 and those previously reported with

hnRNP K during HCV infection, our results and conclusions also echo those reported for the

IGF2BP2/YBX-1 complex, METTL3/METTL14 N6-methyladenosine (m^{6}A) writers, and the

YTHDF m^{6}A-binding proteins (14, 40, 41). These have all been reported to inhibit HCV

infectious particle production, with no effect on viral translation or RNA replication – with the

exception of the IGF2BP2/YBX-1 complex, which plays an additional role in facilitating viral

RNA replication (40). It is currently unclear if PCBP1, hnRNP K, the YBX-1 complex or m^{6}A

modifications of the HCV genome inhibit virion assembly as components of a common pathway,

or through distinct and/or additive mechanisms. Interestingly, high-throughput affinity capture

and proximity ligation studies have found that PCBP1 interacts with hnRNP K, IGF2BP2, YBX-

1, METTL3 and METTL14; although this has been demonstrated only in non-hepatic cell lines

to date (42-44). Should these interactions be conserved during HCV infection, it seems plausible

that PCBP1 could participate in these virion assembly inhibition pathways – or, conversely, that

it may have an inhibitory role to promote one pathway over another. Future investigations will

reveal whether these proteins function in an overlapping or a distinct manner; and are likely to

improve our understanding of HCV virion assembly and how this process is regulated by cellular

RNA binding proteins.

Taken together, our results support a model where endogenous PCBP1 limits HCV

infectious particle production. By preventing virion assembly and egress, PCBP1 indirectly

enhances viral RNA accumulation. The model presented herein helps to inform understanding of

how cellular RNA-binding proteins modulate HCV genomic RNA utilization during the viral life

cycle, specifically as it pertains to virion assembly. While the precise molecular mechanism

employed by PCBP1 to inhibit HCV assembly and egress remains to be characterized, similar

phenotypes reported for hnRNP K, IGF2BP2/YBX-1 and m^{6}A modifications of the HCV

genome offer promising leads for future investigations.

ACKNOWLEDGEMENTS

We would like to acknowledge Charlie Rice (Rockefeller University) for kindly providing the

Huh-7.5 cells, pJ6/JFH FL RLuc WT and GNN plasmids; Rodney Russell (Memorial University)

for providing JFH-1$_T$; Mamata Panigrahi and Joyce Wilson (University of Saskatchewan) for the

pJ6/JFH mono RLuc NS2 plasmid; Martin J Richer (McGill University) for the 293T cells; and

John Law (University of Alberta) for supplying the HCVpp used herein. We are also grateful to

Nathan Taylor, Julie Magnus and Carolina Camargo (McGill University) for technical support.

FUNDING INFORMATION

This research was supported by the Canadian Institutes for Health Research (CIHR) [MOP-

136915 and PJT-169214]. S.E.C. was supported by the Canadian Network on Hepatitis C

(CanHepC) training program, as well as a Vanier Canada Graduate Scholarship. In addition, this

research was undertaken, in part, thanks to the Canada Research Chairs program (S.M.S.).

AUTHOR CONTRIBUTIONS

S.E.C. and S.M.S. designed the study; S.E.C. performed the experiments and analyzed the data,

and S.E.C and S.M.S. wrote and edited the manuscript.

REFERENCES

1. **Simmonds P**, **Becher P**, **Bukh J**, **Gould EA**, **Meyers G**, **Monath T**, **Muerhoff S**, **Pletnev A**, **Rico-Hesse R**, **Smith DB**, **Stapleton JT**, **ICTV Report Consortium**. 2017. ICTV Virus Taxonomy Profile: Flaviviridae. J Gen Virol **98**:2–3.

2. **Tsukiyama-Kohara K**, **Iizuka N**, **Kohara M**, **Nomoto A**. 1992. Internal ribosome entry site within hepatitis C virus RNA. Journal of Virology **66**:1476–1483.

3. **Lancaster AM**, **Jan E**, **Sarnow P**. 2006. Initiation factor-independent translation mediated by the hepatitis C virus internal ribosome entry site. RNA **12**:894–902.

4. **Lohmann V**, **Korner F**, **Koch J**, **Herian U**, **Theilmann L**, **Bartenschlager R**. 1999. Replication of subgenomic hepatitis C virus RNAs in a hepatoma cell line. Science **285**:110–113.

5. **Jones CT**, **Murray CL**, **Eastman DK**, **Tassello J**, **Rice CM**. 2007. Hepatitis C virus p7 and NS2 proteins are essential for production of infectious virus. Journal of Virology **81**:8374–8383.

6. **Jones DM**, **Atoom AM**, **Zhang X**, **Kottilil S**, **Russell RS**. 2011. A genetic interaction between the core and NS3 proteins of hepatitis C virus is essential for production of infectious virus. Journal of Virology **85**:12351–12361.

7. **Masaki T**, **Suzuki R**, **Murakami K**, **Aizaki H**, **Ishii K**, **Murayama A**, **Date T**, **Matsuura Y**, **Miyamura T**, **Wakita T**, **Suzuki T**. 2008. Interaction of hepatitis C virus nonstructural protein 5A with core protein is critical for the production of infectious virus particles. Journal of Virology **82**:7964–7976.

8. **Appel N**, **Zayas M**, **Miller S**, **Krijnse-Locker J**, **Schaller T**, **Friebe P**, **Kallis S**, **Engel U**, **Bartenschlager R**. 2008. Essential role of domain III of nonstructural protein 5A for hepatitis C virus infectious particle assembly. PLoS Pathog **4**:e1000035.

9. **Matunis MJ**, **Michael WM**, **Dreyfuss G**. 1992. Characterization and primary structure of the poly(C)-binding heterogeneous nuclear ribonucleoprotein complex K protein. Mol Cell Biol **12**:164–171.

10. **Du Z**, **Lee JK**, **Fenn S**, **Tjhen R**, **Stroud RM**, **James TL**. 2007. X-ray crystallographic and NMR studies of protein-protein and protein-nucleic acid interactions involving the KH domains from human poly(C)-binding protein-2. RNA **13**:1043–1051.

11. **Spångberg K**, **Schwartz S**. 1999. Poly(C)-binding protein interacts with the hepatitis C virus 5' untranslated region **80**:1371–1376.

12. **Flynn RA**, **Martin L**, **Spitale RC**, **Do BT**, **Sagan SM**, **Zarnegar B**, **Qu K**, **Khavari PA**, **Quake SR**, **Sarnow P**, **Chang HY**. 2015. Dissecting noncoding and pathogen RNA-protein interactomes. RNA **21**:135–143.

13. **Fan B**, **Lu K-Y**, **Reymond Sutandy FX**, **Chen Y-W**, **Konan K**, **Zhu H**, **Kao CC**, **Chen**
C-S. 2014. A human proteome microarray identifies that the heterogeneous nuclear
ribonucleoprotein K (hnRNP K) recognizes the 5' terminal sequence of the hepatitis C
virus RNA. Mol Cell Proteomics **13**:84–92.

14. **Poenisch M**, **Metz P**, **Blankenburg H**, **Ruggieri A**, **Lee J-Y**, **Rupp D**, **Rebhan I**,
Diederich K, **Kaderali L**, **Domingues FS**, **Albrecht M**, **Lohmann V**, **Erfle H**,
Bartenschlager R. 2015. Identification of HNRNPK as Regulator of Hepatitis C Virus
Particle Production. PLoS Pathog **11**:e1004573–21.

15. **Wang L**, **Jeng K-S**, **Lai MMC**. 2011. Poly(C)-binding protein 2 interacts with sequences
required for viral replication in the hepatitis C virus (HCV) 5' untranslated region and
directs HCV RNA replication through circularizing the viral genome. Journal of Virology
85:7954–7964.

16. **Choi K**, **Kim JH**, **Li X**, **Paek KY**, **Ha SH**, **Ryu SH**, **Wimmer E**, **Jang SK**. 2004.
Identification of cellular proteins enhancing activities of internal ribosomal entry sites by
competition with oligodeoxynucleotides. Nucleic Acids Research **32**:1308–1317.

17. **Randall G**, **Panis M**, **Cooper JD**, **Tellinghuisen TL**, **Sukhodolets KE**, **Pfeffer S**,
Landthaler M, **Landgraf P**, **Kan S**, **Lindenbach BD**, **Chien M**, **Weir DB**, **Russo JJ**, **Ju**
J, **Brownstein MJ**, **Sheridan R**, **Sander C**, **Zavolan M**, **Tuschl T**, **Rice CM**. 2007.
Cellular cofactors affecting hepatitis C virus infection and replication. PNAS **104**:12884–
12889.

18. **Russell RS**, **Meunier J-C**, **Takikawa S**, **Faulk K**, **Engle RE**, **Bukh J**, **Purcell RH**,
Emerson SU. 2008. Advantages of a single-cycle production assay to study cell culture-
adaptive mutations of hepatitis C virus. Proc Natl Acad Sci USA **105**:4370–4375.

19. **Panigrahi M**, **Thibault PA**, **Wilson JA**. 2021. Chapter 7: miR-122 affects both the early
and late stages of a Hepatitis C Virus infection, pp. 129–165. *In* miR-122-independent
propagation of Hepatitis C Virus: Understanding the underlying mechanism through
characterization of viral replication [dissertation]. Saskatoon: University of Saskatchewan;
2021.

20. **Moradpour D**, **Evans MJ**, **Gosert R**, **Yuan Z**, **Blum HE**, **Goff SP**, **Lindenbach BD**,
Rice CM. 2004. Insertion of green fluorescent protein into nonstructural protein 5A
allows direct visualization of functional hepatitis C virus replication complexes. Journal of
Virology **78**:7400–7409.

21. **Amador-Cañizares Y**, **Bernier A**, **Wilson JA**, **Sagan SM**. 2018. miR-122 does not
impact recognition of the HCV genome by innate sensors of RNA but rather protects the
5′ end from the cellular pyrophosphatases, DOM3Z and DUSP11. Nucleic Acids Research
46:5139–5158.

22.   **Machlin ES**, **Sagan SM**, **Sarnow P**. 2011. Masking the 5' terminal nucleotides of the
499   hepatitis C virus genome by an unconventional microRNA-target RNA complex. PNAS
500   **108**:3193–3198.

501   23.   **Hsu M**, **Zhang J**, **Flint M**, **Logvinoff C**, **Cheng-Mayer C**, **Rice CM**, **McKeating JA**.
502   2003. Hepatitis C virus glycoproteins mediate pH-dependent cell entry of pseudotyped
503   retroviral particles. PNAS **100**:7271–7276.

504   24.   **Carroll SS**, **Tomassini JE**, **Bosserman M**, **Getty K**, **Stahlhut MW**, **Eldrup AB**, **Bhat**
505   **B**, **Hall D**, **Simcoe AL**, **LaFemina R**, **Rutkowski CA**, **Wolanski B**, **Yang Z**, **Migliaccio**
506   **G**, **De Francesco R**, **Kuo LC**, **MacCoss M**, **Olsen DB**. 2003. Inhibition of hepatitis C
507   virus RNA replication by 2'-modified nucleoside analogs. Journal of Biological Chemistry
508   **278**:11979–11984.

509   25.   **Schindelin J**, **Arganda-Carreras I**, **Frise E**, **Kaynig V**, **Longair M**, **Pietzsch T**,
510   **Preibisch S**, **Rueden C**, **Saalfeld S**, **Schmid B**, **Tinevez J-Y**, **White DJ**, **Hartenstein V**,
511   **Eliceiri K**, **Tomancak P**, **Cardona A**. 2012. Fiji: an open-source platform for biological-
512   image analysis. Nat Meth **9**:676–682.

513   26.   **Livak KJ**, **Schmittgen TD**. 2001. Analysis of Relative Gene Expression Data Using
514   Real-Time Quantitative PCR and the $2-\Delta\Delta CT$ Method. Methods **25**:402–408.

515   27.   **Wang C**, **Sarnow P**, **Siddiqui A**. 1993. Translation of Human Hepatitis C Virus RNA in
516   Cultured Cells Is Mediated by an Internal Ribosome-Binding Mechanism. Journal of
517   Virology **67**:3338–3344.

518   28.   **Le Quesne JPC**, **Stoneley M**, **Fraser GA**, **Willis AE**. 2001. Derivation of a structural
519   model for the c-myc IRES. Journal of Molecular Biology **310**:111–126.

520   29.   **Pickering BM**, **Mitchell SA**, **Spriggs KA**, **Stoneley M**, **Willis AE**. 2004. Bag-1 internal
521   ribosome entry segment activity is promoted by structural changes mediated by poly(rC)
522   binding protein 1 and recruitment of polypyrimidine tract binding protein 1. Mol Cell Biol
523   **24**:5595–5605.

524   30.   **Evans JR**, **Mitchell SA**, **Spriggs KA**, **Ostrowski J**, **Bomsztyk K**, **Ostarek D**, **Willis**
525   **AE**. 2003. Members of the poly (rC) binding protein family stimulate the activity of the c-
526   myc internal ribosome entry segment in vitro and in vivo. Oncogene **22**:8012–8020.

527   31.   **Walter BL**, **Parsley TB**, **Ehrenfeld E**, **Semler BL**. 2002. Distinct Poly(rC) Binding
528   Protein KH Domain Determinants for Poliovirus Translation Initiation and Viral RNA
529   Replication. Journal of Virology **76**:12008–12022.

530   32.   **Luo Z**, **Dong X**, **Li Y**, **Zhang Q**, **Kim C**, **Song Y**, **Kang L**, **Liu Y**, **Wu K**, **Wu J**. 2014.
531   PolyC-Binding Protein 1 Interacts with 5′-Untranslated Region of Enterovirus 71 RNA in
532   Membrane-Associated Complex to Facilitate Viral Replication. PLoS ONE **9**:e87491–13.

33.   **Taguwa S**, **Yeh M-T**, **Rainbolt TK**, **Nayak A**, **Shao H**, **Gestwicki JE**, **Andino R**,
Frydman J. 2019. Zika Virus Dependence on Host Hsp70 Provides a Protective Strategy
against Infection and Disease. CellReports **26**:906–920.e3.

34.   **Zhou X**, **You F**, **Chen H**, **Jiang Z**. 2012. Poly(C)-binding protein 1 (PCBP1) mediates
housekeeping degradation of mitochondrial antiviral signaling (MAVS). Cell Res **22**:717–
727.

35.   **Sumpter R**, **Loo Y-M**, **Foy E**, **Li K**, **Yoneyama M**, **Fujita T**, **Lemon SM**, **Gale M**.
2005. Regulating intracellular antiviral defense and permissiveness to hepatitis C virus
RNA replication through a cellular RNA helicase, RIG-I. Journal of Virology **79**:2689–
2699.

36.   **Meylan E**, **Curran J**, **Hofmann K**, **Moradpour D**, **Binder M**, **Bartenschlager R**,
Tschopp J. 2005. Cardif is an adaptor protein in the RIG-I antiviral pathway and is
targeted by hepatitis C virus. Nature **437**:1167–1172.

37.   **Bender S**, **Reuter A**, **Eberle F**, **Einhorn E**, **Binder M**, **Bartenschlager R**. 2015.
Activation of Type I and III Interferon Response by Mitochondrial and Peroxisomal
MAVS and Inhibition by Hepatitis C Virus. PLoS Pathog **11**:e1005264–30.

38.   **Dinh PX**, **Beura LK**, **Panda D**, **Das A**, **Pattnaik AK**. 2011. Antagonistic effects of
cellular poly(C) binding proteins on vesicular stomatitis virus gene expression. Journal of
Virology, 3rd ed. **85**:9459–9471.

39.   **Woolaway K**, **Asai K**, **Emili A**, **Cochrane A**. 2007. hnRNP E1 and E2 have distinct roles
in modulating HIV-1 gene expression. Retrovirology **4**:28–18.

40.   **Chatel-Chaix L**, **Germain M-A**, **Motorina A**, **Bonneil É**, **Thibault P**, **Baril M**,
Lamarre D. 2013. A host YB-1 ribonucleoprotein complex is hijacked by hepatitis C
virus for the control of NS3-dependent particle production. Journal of Virology **87**:11704–
11720.

41.   **Gokhale NS**, **McIntyre ABR**, **McFadden MJ**, **Roder AE**, **Kennedy EM**, **Gandara JA**,
Hopcraft SE, **Quicke KM**, **Vazquez C**, **Willer J**, **Ilkayeva OR**, **Law BA**, **Holley CL**,
Garcia-Blanco MA, **Evans MJ**, **Suthar MS**, **Bradrick SS**, **Mason CE**, **Horner SM**.
2016. N6-Methyladenosine in Flaviviridae Viral RNA Genomes Regulates Infection. Cell
Host & Microbe **20**:654–665.

42.   **Rosenbluh J**, **Mercer J**, **Shrestha Y**, **Oliver R**, **Tamayo P**, **Doench JG**, **Tirosh I**,
Piccioni F, **Hartenian E**, **Horn H**, **Fagbami L**, **Root DE**, **Jaffe J**, **Lage K**, **Boehm JS**,
Hahn WC. 2016. Genetic and Proteomic Interrogation of Lower Confidence Candidate
Genes Reveals Signaling Networks in β-Catenin-Active Cancers. Cell Syst **3**:302–316.e4.

43.   **Youn J-Y**, **Dunham WH**, **Hong SJ**, **Knight JDR**, **Bashkurov M**, **Chen GI**, **Bagci H**,
Rathod B, **MacLeod G**, **Eng SWM**, **Angers S**, **Morris Q**, **Fabian M**, **Côté J-F**,

Gingras A-C. 2018. High-Density Proximity Mapping Reveals the Subcellular
Organization of mRNA-Associated Granules and Bodies. Molecular Cell **69**:517–532.e11.

44.   **Yue Y**, **Liu J**, **Cui X**, **Cao J**, **Luo G**, **Zhang Z**, **Cheng T**, **Gao M**, **Shu X**, **Ma H**, **Wang**
F, **Wang X**, **Shen B**, **Wang Y**, **Feng X**, **He C**, **Liu J**. 2018. VIRMA mediates preferential
m(6)A mRNA methylation in 3'UTR and near stop codon and associates with alternative
polyadenylation. Cell Discov **4**:10.

FIGURE LEGENDS

Figure 1. PCBP1 knockdown decreases viral protein expression and intracellular viral

RNA accumulation, but increases secreted virus titers. (A) Schematic representation of the

HCVcc (JFH-1$_T$) infectious particles and genomic RNA used in infections. Huh-7.5 cells were

transfected with siPCBP1 or siCTRL at day -2, and at day 0 were infected with JFH-1$_T$ (MOI =

0.05). Viral protein, Total RNA and intracellular and extracellular infectious virus were

harvested at day 3 post-infection. **(B)** Viral protein expression analysis by Western blot (ns

indicates non-specific band; and arrow indicates core and NS5A bands). **(C)** Viral RNA

accumulation analysis by Northern blot, and **(D)** quantification by densitometry. **(E)** Intracellular

and **(F)** extracellular (secreted) virus titers, quantified by FFU assay. All data are representative

of three independent replicates, and error bars represent the standard deviation of the mean. P-

values were calculated by paired t-test.

Figure 2. PCBP1 knockdown has no effect on HCV entry. Two days post-siRNA transfection,

cells were spinoculated with luciferase reporter pseudoparticles expressing the HCV E1/E2

glycoproteins (HCVpp) or the VSV-G glycoprotein (VSVpp). In parallel, cells were transfected

with a firefly luciferase expression plasmid. Samples were harvested 3 days post-

infection/transfection, and analyzed by luciferase assay. The HCVpp and pFLuc data is

representative of three independent replicates, while the VSVpp data is representative of two

independent replicates. Error bars represent the standard deviation of the mean.

Figure 3. PCBP1 knockdown has no impact on HCV translation or genome stability. (A)

Schematic representation of the J6/JFH-RLuc-GNN reporter RNA. **(B)** Huh-7.5 cells were

electroporated with J6/JFH-1-RLuc-GNN reporter RNAs and FLuc control mRNA, and

luciferase activity was monitored at several timepoints post-electroporation. All data are

representative of three independent replicates. Error bars on the y-axis represent the standard

deviation of the mean, while error bars on the x-axis indicate the range of sample harvest times

across replicates.

Figure 4. PCBP1 knockdown does not affect viral RNA replication. Two days post-siRNA

transfection, Huh-7.5 cells were electroporated with **(A)** full-length RLuc WT J6/JFH-1 RNA,

(B) Δcore-p7 J6/JFH-1 RLuc RNA, or **(C)** full-length NS5A-GFP J6/JFH-1 RLuc RNA. RLuc

values were normalized to the early timepoint (3h), to control for disparities in electroporation

efficiency between experiments. All data are representative of three independent replicates and

error bars represent the standard deviation of the mean.

Figure 5. Viral particle production after 2´CMA treatment confirms that PCBP1

knockdown enhances virion secretion. **(A)** Schematic representation of the experimental

approach for 2´CMA experiments: two days post-siRNA transfection, Huh-7.5 cells were

infected with JFH-1$_T$ (MOI = 0.05). Three days post-infection, cells were treated with 2´CMA or

DMSO (control). Total RNA, intracellular and extracellular virus was collected at t = 0, 6 and 12

h post-treatment. **(B)** Quantitative RT-PCR analysis after 2´CMA treatment. **(C)** Intracellular

viral titers and **(D**) intracellular virus accumulation rate calculated by linear regression. **(E)**

Extracellular viral titers and **(F)** virus secretion rates calculated by linear regression. All data are

representative of three independent replicates and error bars in (B), (C) and (E) represent the

standard deviation of the mean. Error bars in (D) and (F) represent the slopes of the linear

regressions ± standard error. P-values were calculated by two-way ANOVA.

Figure 6. Model for PCBP1 effects on HCV assembly and egress. Under wild-type

conditions, PCBP1 inhibits HCV assembly and egress, leaving the majority of intracellular viral

RNAs to engage in protein translation and viral genome replication. However, during PCBP1

knockdown, a disinhibition of assembly and egress leads to an overall decrease in the

translating/replicating pool of viral RNAs, and a concomitant increase in viral infectious particle

production.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Wild-type

PCBP1 knockdown

Release

Egress

Assembly

Cell Surface

Golgi Apparatus

ER Lumen

Translating/
Replicating Pool

Figure 6

Supplementary Figure 1. PCBP1 knockdown decreases both positive-sense and negative-sense viral RNA accumulation. Two days post-siRNA transfection, Huh-7.5 cells were infected with JFH-1$_T$ (MOI = 0.05). Total RNA was harvested 3 days post-infection and analysed by strand-specific RT-qPCR. All data is representative of three independent replicates and error bars represent the standard deviation. P-values were calculated by paired t-test.

Supplementary Figure 2. The J6/JFH FL Rluc WT can produce infectious viral particles, but much less efficiently than JFH-1$_T$. Three days post-electroporation of viral RNA, intracellular and extracellular viral titers were harvested and assayed by focus-forming unit assay. Data is representative of one independent replicate measured in duplicate. The limit of detection is indicated (dotted line).