Novel newt regeneration genes regulate Wingless signaling to restore patterning in *Drosophila* eye

Abijeet Singh Mehta[1], Prajakta Deshpande[1], Anuradha Venkatakrishnan Chimata[1], Panagiotis A. Tsonis[1†], Amit Singh[1, 2, 3, 4, 5]*

(1) Department of Biology, University of Dayton, Dayton, OH 45469, USA.

(2) Premedical Program, University of Dayton.

(3) Center for Tissue Regeneration and Engineering at Dayton (TREND), University of Dayton.

(4) The Integrative Science and Engineering Center, University of Dayton, Dayton, OH 45469, USA.

(5) Center for Genomic Advocacy (TCGA), Indiana State University, Terre Haute, IN, USA.

[†]Deceased

Running Title: Newt genes regulate Wg signaling to recue eye mutant phenotype

Key words: *Drosophila*, Eye, Newts, Wingless, Cell death, Growth, Regeneration

*Corresponding author

Amit Singh

Email: asingh1@udayton.edu

Abstract

A fundamental process of regeneration, which varies among animals, recruits conserved signaling pathways to restore missing parts. Only a few animals like newts can repeatedly regenerate lost body parts throughout their lifespan that can be attributed to strategic regulation of conserved signaling pathways by newt's regeneration tool-kit genes. Here we report the use of a genetically tractable *Drosophila* eye model to demonstrate the regeneration potential of a group of unique protein(s) from newt (*Notophthalmus viridescens),* which when ectopically expressed can significantly rescue missing photoreceptor cells in a *Drosophila* eye mutant. These newt proteins with signal peptides motifs exhibit non-cell-autonomous rescue properties and their regeneration potential even extends into later stages of fly development. Ectopic expression of these newt genes can rescue eye mutant phenotype by promoting cell proliferation and blocking cell death. These novel newt genes downregulate the evolutionarily conserved Wingless (Wg)/Wnt signaling pathway to promote rescue. Modulation of Wg/Wnt signaling levels by using antagonists or agonists of Wg/Wnt signaling pathway in eye mutant background where newt gene(s) is ectopically expressed suggests that Wg signaling acts downstream of newt genes. Our data highlights the regeneration potential of novel newt proteins that regulate conserved pathways to trigger a robust regeneration response in *Drosophila* model with weak regeneration capability.

Introduction

A urodele amphibian, newts belongs to *Salamandridae* family (Weisrock et al., 2006). The newts are the only four-legged vertebrates that exhibits a remarkable capability to regenerate tissues like limbs, tail, heart, lens, spinal cord, brain, and retina throughout its life time (Mehta and Singh, 2019a; Sanchez Alvarado and Tsonis, 2006). This regeneration potential/ ability of the newt is due to their ability to reprogram and dedifferentiate the terminally differentiated cells to trigger regeneration response (Mehta and Singh, 2019a). Such an exceptional regeneration ability of newts has been attributed to unique gene(s) that may have evolved to form the regeneration tool box (Bryant et al., 2017; Casco-Robles et al., 2018; Elewa et al., 2017; Evans et al., 2018; Keinath et al., 2015; Kumar et al., 2007; Matsunami et al., 2019; Mehta and Singh, 2019a; Nowoshilow et al., 2018; Sanor et al., 2020; Smith et al., 2009; Smith et al., 2019). Earlier, a newt gene Prod1, which encodes a transmembrane receptor, was found to be critical for maintaining proximo-distal identity (pattern memory) during newt limb regeneration (da Silva et al., 2002; Echeverri and Tanaka, 2005; Kumar et al., 2007; Mehta and Singh, 2019a). Homologs of Prod1 are yet to be identified outside of salamanders (Garza-Garcia et al., 2009). Similarly, the transcript of a novel newt gene *newtic1* was found to be significantly enriched in a subset of erythrocytes which formed an aggregate structure called erythrocyte clump. These *newtic* expressing erythrocytes promote limb regeneration (Casco-Robles et al., 2018). Surprisingly, our understanding of underlying molecular genetic mechanism(s) responsible for promoting lifelong regeneration of missing structures and/or rescue of pattern defects in newts is far from complete. Some strategies to determine their regeneration potential are to introduce these unique genes in genetically tractable model(s) that lack regeneration potential exhibited by newts such as mammals, *Drosophila* etc. Since the genetic machinery is conserved across the species, the *Drosophila* model has been successfully used for such cross species studies to model human disease and determine their underlying molecular genetic mechanism(s) (Hughes et al., 2012; Sarkar et al., 2018; Sarkar et al., 2016; Singh and Irvine, 2012; Tare et al., 2011).

Previously, a novel family of five protein members was identified in a
comprehensive transcriptomic analysis from *Notophthalmus viridescens* (red-spotted
newt). These proteins are expressed in adult tissues and are modulated during lens
regeneration (Looso et al., 2013). The three-dimensional structure models of these
novel newt proteins by *ab initio* methods suggested that these proteins could act as ion
transporters and/or involved in signaling (Mehta et al., 2019). These unique newt genes
were introduced in genetically tractable *Drosophila melanogaster* (fruit fly) model by
transgenic approaches to test their regeneration potential. Such transgenic approaches
were not feasible in *N. viridescens*, therefore, *Drosophila* served as a suitable system to
address these queries. The transcriptomic analysis was performed in the transgenic
Drosophila that are expressing newt genes. Based on these studies, graded expression
of 2775 transcripts was reported in transgenic fly where one of the five newly identified
newt genes was ectopically expressed using the GAL4/UAS binary target system
(Brand and Perrimon, 1993; Mehta et al., 2019). Among these 2775 transcripts, genes
involved in the fundamental developmental processes like cell cycle, apoptosis, and
immune response were highly enriched suggesting that these foreign genes from newt
were able to modulate the expression of *Drosophila* gene(s) (Mehta et al., 2019; Mehta
and Singh, 2019b).

Drosophila, a hemimetabolous insect, serves as a versatile model organism to
study development and disease due to presence of fully sequenced and less redundant
genome, presence of homologs or orthologs for human disease related genes, a shorter
life cycle and economy to maintain fly cultures (Bier, 2005; Lenz et al., 2013; Singh and
Irvine, 2012). *Drosophila* eye has been extensively used to understand the fundamental
process of development and model neurodegenerative disorders (Gogia et al., 2020a;
Kumar, 2018; Sarkar et al., 2018; Sarkar et al., 2016; Singh and Irvine, 2012; Tare et
al., 2011; Yeates et al., 2019). The rich repository of genetic tools available in fly model,
and the fact that eye is dispensable for survival, makes *Drosophila* eye suitable for
screening genes from other animals that can promote growth and regeneration. The
compound eye of the adult fly develops from a monolayer epithelium called the eye-
antennal imaginal disc, which develops from an embryonic ectoderm (Cohen, 1993;
Held, 2002). The imaginal disc, a sac-like structure present inside the larva, comprise of

two different layers: the peripodial membrane (PM) and the disc proper (DP) (Kumar,
2020b). The DP develops into retina whereas the PM of the eye-antennal imaginal disc
contributes to the adult head structures (Haynie and Bryant, 1986; Kumar, 2020b;
Milner et al., 1983). In third instar larval eye imaginal disc, a wave of synchronous
retinal differentiation moves anteriorly from the posterior margin of the eye disc, and is
referred to as Morphogenetic Furrow (MF) (Kumar, 2013, 2020a; Ready et al., 1976).
The progression of MF transforms the undifferentiated cells into differentiated
photoreceptor cells. The adult compound eye comprises of 800 unit eyes or ommatidia
(Kumar, 2013, 2018, 2020a; Ready et al., 1976). The ommatidial cells in the compound
eye are grouped together into two chiral forms, which are arranged in mirror image
symmetry along the Dorso-Ventral (DV) midline called the equator. Ventral domain is
the default state of the entire early eye imaginal primordium (Singh and Choi, 2003),
and onset of the expression of the dorsal selector gene *pannier* (*pnr*), establishes the
dorso-ventral (DV) lineage in the eye (Maurel-Zaffran and Treisman, 2000; Singh and
Choi, 2003; Tare et al., 2013). It has been reported that *Lobe* (*L*) is a cell survival gene
required for ventral eye development and growth, and acts upstream to the Wg
signaling pathway (Singh et al., 2006).

The *wg* gene, a *Drosophila* homolog of Wnt, serves as a ligand, and encodes a
secreted signaling protein that acts as a morphogen (Baker, 1987; Bejsovec, 2013;
Swarup and Verheyen, 2012). The transcriptional effector of the Wg/β-catenin signaling
is *Drosophila* T-cell Factor (dTCF), which upon activation by Armadillo (Arm, human
homolog β-catenin) leads to the transcription of Wg target genes. In the absence of
ligand Wg, a destruction complex comprised of Adenomatous polyposis coli (Apc), and
Axin degrades Arm and thus prevents it to form a complex with dTCF. The pathway is
activated by binding of Wg ligand to its co-receptors Frizzled (Fz) and Arrow (Arr). Upon
activation, Fz binds to Dishevelled (Dsh) and Arr, which binds to Axin and thus
inactivating the destruction complex. This results in the translocation of Arm (β-catenin)
from the cytosol to the nucleus where Arm binds to dTCF and activates transcription of
Wg target genes (Bejsovec, 2006). In the eye-antennal disc, Wg acts as a negative
regulator of the eye development (Ma and Moses, 1995; Treisman and Rubin, 1995). In

early third instar eye-antennal imaginal disc, *wg* expresses on the ventral margin.
Ectopic expression of Wg triggers cell death of photoreceptor cells causing eye defects
(Baker, 1987; Cavodeassi et al., 1999; Singh, 2012; Singh et al., 2006). Previously it
has been shown that Wg pathway was upregulated in developing eye of L^2 background.
This perturbation in Wg pathway causes loss of ventral eye in 100% of flies of L^2/+
background (Gogia et al., 2020a; Singh et al., 2006; Singh et al., 2011). Therefore,
timing and level of *wg* expression in this region is crucial for formation of a normally
patterned eye. Here, we demonstrate using a unique gene pool of newt genes (5 newly
identified newt genes) to rescue *Drosophila L^2* mutant pattern defect*.*
150
Misexpression of these novel newt gene in *Drosophila* downregulate Wg in L^2
mutant background, resulting in significant restoration of missing photoreceptor cells in
the developing *Drosophila* eye. Ectopic expression of these genes restore eye mutant
phenotypes by promoting cell proliferation, and downregulating cell death. Furthermore,
these eye mutant phenotype rescues by newt genes are non-cell-autonomous and are
all along the developmental stages *viz.,* early as well as later time points of
development.
158

Results

We employed *Drosophila* eye model to test the regeneration potential of these five

unique newt genes, *candidate 1* (*C1*), *candidate 2 (C2), candidate 3 (C3), candidate 4*

(C4), and *candidate 5 (C5)*. We isolated the total RNA from newt tail and generated

cDNA to amplify these newt genes, and generated transgenic flies for these five genes

(Figure 1A). We employed GAL4/UAS system (Brand and Perrimon, 1993) to allow

targeted misexpression of these newt transgenes in the fly tissues (Figure 1B). To

confirm if targeted misexpression approach can generate newt proteins in *Drosophila*

(Brand and Perrimon, 1993), we tested localization of the fusion protein. As there are

no antibodies available against these novel newt proteins, so we used V5 epitope

tagged transgenes to generate transgenic flies (Wang et al., 2019). The V5 tag

sequence of about 42 bp long is fused towards the 3' end of the 501 bp long open

reading frame (ORF). We tested all these transgenes using the antibody against V5 tag.

Here we present the expression of one of the transgenes *viz.*, C4 using V5 tag (Figure

S1). Next, to detect if misexpression of these newt genes in *Drosophila* can generate

any developmental defects, we used *tubulin* – GAL4 (*tub* –GAL4) to ubiquitously

misexpress newt genes in flies in the wild-type background (Figure 1B), and found no

defects.

Newt genes can rescue *Drosophila* eye mutants

For our experiments, we used the *Drosophila* eye model as it is not required for the

viability of the fly and it is easy to score the phenotypes. We used *ey*-GAL4 driver

(Hazelett et al., 1998) to misexpress these newt genes in the developing eye imaginal

disc, which develops into an adult eye. The phenotypes of UAS-transgene alone were

tested to rule out any contribution(s) in phenotype due to insertion of the transgene. In

comparison to the control, *ey*-GAL4 (Figure 1C), misexpression of these transgenes *ey*-

GAL4>UAS-*C1* (*ey*>*C1*, Figure 1D),   *ey*-GAL4>UAS-*C2 (ey>C2*, Figure 1E),   *ey*-

GAL4>UAS-*C3* (*ey*>*C3*, Figure 1F),   *ey*-GAL4>UAS-*C4* (*ey*>*C4*, Figure 1G), and *ey*-

GAL4>UAS-*C5* (*ey*>*C5*, Figure 1H),   did not affect the eye size (Figure 1I). To

investigate the regeneration potential of these genes, we used eye mutant(s) such as *L*

mutant which exhibits selective loss of ventral half of the eye (Figure 1J). This mutant

was selected as the penetrance of the phenotype of loss-of-ventral eye is 100%.

Interestingly, targeted misexpression of the newt gene(s) in L^2; *ey*-GAL4 background

[(L^2; *ey>C1*; Figure. 1K), (L^2; *ey>C2*; Figure 1L), (L^2; *ey>C3*; Figure 1M), (L^2; *ey>C4*;

Figure 1N), (L^2; *ey>C5*; Figure 1O)] exhibit significant rescue of the loss-of-ventral eye

phenotype (Figure 1P). For each cross, three independent cultures were established

and two hundred flies for each set (triplicate, 600 flies) were counted to determine

phenotype frequency. The rescue frequency was about 40.83% in C1, 37.7% in C2,

49% in C3, 58.2% in C4, and 38.7% in C5 (Figure S2). Note that the strongest rescues

were observed with C4 transgene (Figure 1P; S2; and Table S1).

Since the eye mutant represent developmental defects in the genetic machinery

involved in early eye development, it raises a question whether the regeneration

potential of these newt genes is restricted to the early time window or even later stages

of eye development. The GMR- GAL4 driver was used to drive expression of

transgenes in the differentiating retinal neurons of larval eye imaginal disc, which

continues all along to the adult eye (Moses and Rubin, 1991). Gain-of-function of *head*

involution defective (*hid*), an executioner caspase of cell death machinery, triggers cell

death (Bergmann et al., 1998; Grether et al., 1995; Hay et al., 1995; White et al., 1994).

Misexpression of *hid* using GMR-GAL4 (GMR>*hid*) results in a "No-eye" or highly

reduced eye phenotype (Figure S3B). However, misexpression of newt transgenes

along with *hid* in the eye (GMR-GAL4>*hid*+*C4*) results in a significant rescue of the "No-

eye" phenotype (Figure S3C). Since these newt genes have signal peptides (Looso et

al., 2013), it is possible that these genes may have non-autonomous effects.

Ectopic expression of newt genes exhibits non-autonomous rescue

To determine if these genes have non-autonomous effect, we misexpressed these

genes within a subset of retinal neuron population in the developing eye field of L^2

mutant and assay their regeneration effect. Misexpression of *C4* using *dorsal-eye -*

GAL4 (*DE - GAL4*) (Figure 1B) directs its expression in the dorsal half of the developing

eye (Morrison and Halder, 2010) (Figure 2A, E). The rationale of this experiment was to

misexpress *C4* in the dorsal half of the eye and test its effect on an eye mutant which

results in the loss of the ventral half of the developing eye (Figure 2). Misexpression of

both GFP (reporter) and newt *C4* in the dorsal domain of the developing eye, which

serves as a positive control (+/+*; DE>GFP+C4)* (Figure 2A, B, B′, B″), exhibits a near

wild-type eye. As a negative control, we tested GFP expression alone in L^2-mutant

background (L^2/+*; DE>GFP)* (Figure 2C, D, D', D"). We found that GFP expression was

restricted to the dorsal half as seen in the wild-type control (Figure 2B, B', B").

Furthermore, we did not see any rescue of *L* mutant phenotype of ventral eye loss

(marked by red dotted boundary, Figure 2B, B′, B″). Misexpression of both GFP+C4 in

the dorsal half of the L^2 mutant eye (L^2/+*; DE>GFP+C4)* exhibits a significant rescue of

the loss-of-ventral eye phenotype as seen in the adult eye, and the eye imaginal disc

(marked by white-dotted boundary, Figure 2E, F, F′, F″). Even though the targeted

misexpression of C4 is restricted to the dorsal half of the developing eye, it was able to

rescue the loss-of-ventral eye phenotype suggesting that newt genes can show non-

cell-autonomous rescue of *Drosophila* eye mutants.

In order to rule out any domain specific restraint, or contribution of a *DE*-GAL4

background, we employed genetic mosaic approach. We generated gain-of-function

clones using Flp out approach (Xu and Rubin, 1993), which result in clones of cells that

misexpress high levels of the transgene and these clones are marked by the GFP

reporter (Figure 2G,H,I,J). We found that the misexpression of these transgenes

rescues the loss-of-ventral eye phenotypes, however these rescues extended beyond

the boundary of the clone. This data validate our prior observation that these newt gene

exhibit non-cell-autonomous rescue due to their secretory nature. Next, we wanted to

investigate the mechanism by which these newt genes can rescue loss-of-ventral eye

phenotype.

Newt genes promote cell proliferation and downregulate cell death

To test the possibility of upregulation of cell proliferation resulting in the rescue of the

missing tissue in the L^2 mutant background, we calculated relative fold change in mitotic

index (MI) of the eye imaginal discs showing rescue of L mutant phenotype and

compared it to the control L^2 mutant (L^2; ey-GAL4) eye disc (Figure 3A-D). Mitotic index

is the ratio between the number of proliferating cells to the rows of ommatidial cells

(Baonza and Freeman, 2005; Vollmer et al., 2016; Yang et al., 2005). The proliferating

cells are marked by anti-phospho-Histone H3 (PH3) marker (Figure 3A-C, A′-C′ and

Figure S4A-F, A′-F′)**.** Phosphorylation of a highly conserved Serine residue (Ser-10) in

the histone H3 tail is considered a crucial event for the onset of mitosis, and represents

completion of cell cycle (Crosio et al., 2002; Kim et al., 2017). Our results show robust

upregulation of about 5.79 ± 1.5 folds increase in the MI when $C4$ is misexpressed in

the L^2-mutant background (L^2; $ey>C4$) with respect to the L^2-mutant (L^2; ey-GAL4) only

(Figure 3D). Similar phenotypes were observed with other transgenes (Figure S4 A-G)

and the MI values were $C3$: 5.18 ± 1.10, $C2$: 4.6 ± 1.7, $C1$: 4.3± 1.4, and $C5$: 4.2 ± 0.9

respectively**.** These results show that the newt gene(s) can induce cell proliferation to

restore the missing photoreceptor cells of L^2-mutant eye disc and thus exhibit significant

rescue of the loss-of-ventral eye phenotype as seen in the eye imaginal disc and the

adult eye (shown in white dotted boundary, Figure 3B). The control where no newt gene

is misexpressed, in the L^2 mutant background did not show any rescue of the loss-of-

ventral eye phenotype due to lack of cell proliferation as evident from PH3 staining

(shown in red dotted boundary, Figure 3A).

During the newt lens regeneration both cell proliferation and apoptosis are seen

(Tsonis et al., 2004), which warranted the need to look at the levels of cell death. It has

been shown that the loss-of-ventral-eye phenotype observed in L^2 mutation is due to

induction of both caspase-dependent and caspase-independent cell death (Singh et al.,

2006). Therefore, we tested if these newt genes can block or downregulate cell death to

rescue L^2 mutant phenotype by using the TUNEL (Terminal deoxynucleotidyl

transferase dUTP nick end labeling) assay (Figure 3E-G, and S4 H-M). TUNEL assay is

based on the detection of fragmented DNA ends, which are characteristic of apoptotic

cells (Sarkissian et al., 2014; White et al., 2001). Our results clearly demonstrate strong

decrease in the number of dying cells (0.28 ± 0.04 fold) when newt gene ($C4$) is

misexpressed under the L^2 mutant background (L^2; ey>C4) (Figure 3F, F', H) relative to

the L^2-mutant strain in which newt gene is not misexpressed (L^2; ey-GAL4) (Figure 3E,

E', H). The positive control (ey-GAL4) shows 0.32 ± 0.04 fold downregulation in the rate

of cell death (Figure 3G, G', H) relative to the L^2 mutant (L^2; ey-GAL4). The value is

significantly lower in comparison to the L^2 mutant (L^2; ey-GAL4), but approximately

equivalent to the L^2 mutant where newt gene is misexpressed (L^2; ey>C4) (Figure 3H).

This suggests that newt gene(s) can recue the L^2 mutant loss-of-ventral eye phenotype

to wild-type by preventing excessive cell death of photoreceptor cells (marked in white

dotted boundary, Figure 3 F, F'). We found similar results when other members of novel

newt gene family were misexpressed (Figure S4 H-O). Downregulation in cell death is

observed in C3: 0.36 ± 0.14, C2: 0.38 ± 0.09, C1: 0.39 ± 0.15, and C5: 0.45 ± 0.22

respectively (Figure S4N). We also tested the role of cell death using Dcp1 (Death

caspase-1) staining (Figure S5). Antibody against cleaved Dcp1 marks the dying cells in

Drosophila (Sarkissian et al., 2014). Dcp1 staining exhibits reduction in the number of

dying cells when newt genes are misexpressed in L^2 mutant background (Figure S5 C-

F), compared to when newt genes are not expressed in L^2 mutant background (Figure

S5 B). These results suggest that newt genes upregulate cell proliferation and

downregulate cell death to significantly rescue the L^2 mutant phenotype of loss-of-

ventral eye.

We further validated the role of cell death by blocking caspase dependent cell

death using baculovirus p35 misexpression (Hay et al., 1994). Even though

misexpression of *p35* in L^2 mutant background (L^2/+; ey>p35) exhibits rescue of loss-of-

ventral eye phenotype, the phenotype strength (relative eye size ratio of 1.42 ± 0.29 and

rescue frequency of 22.7%) of L^2/+; ey>p35 was significantly weaker than phenotype

strength (relative eye size ratio = 2.34 ± 0.51 and rescue frequency of 58.2%) of L^2/+;

ey>C4 (Figure S6 C, D, F, G). This data suggests that the newt genes upregulate cell

proliferation and block cell death as well, but cell proliferation may be playing significant

role in rescuing the missing tissues in *the* L^2 mutant eye. It is therefore important to

understand what signal is triggered by these newt genes to promote the rescue.

Newt genes downregulate *wg* expression in developing eye

We screened for the signaling pathway(s) that are involved in the rescue of L^2 mutant

loss-of-ventral eye phenotype. We identified members of evolutionarily conserved

Wg/Wnt signaling pathway in our high throughput screen (RNA sequencing) where newt

genes were misexpressed in fly tissue (Figure S7). It is reported that L^2 mutant

phenotype can be rescued by downregulating *wg* (Singh et al., 2006). To determine the

involvement of *wg* in promoting rescue of L^2 mutant phenotype by unique newt genes,

we examined expression levels of Wg in larval eye-antennal imaginal disc (Figure 4A-C,

A′-C′, D). Wg expression is seen at antero-lateral margins of the developing eye

imaginal disc (Baker, 1987; Cavodeassi et al., 1999; Singh et al., 2006). In L^2 mutant

robust ectopic expression of Wg is seen at the ventral eye margins (Figure 4B, B′)

(Singh et al., 2006; Singh et al., 2011). Misexpression of newt gene (*C4*) in L^2 mutant

background downregulates the Wg expression at the ventral margin of eye-antennal

imaginal disc (Figure 4C, C′), as evident from the signal intensity plots (integrative

density) in various backgrounds (Figure 4D). The semi-quantitative western blot,

showed that Wg levels increased 3 folds in L^2 mutant as compared to *ey-GAL4* control

whereas misexpressing *C4* in the L^2 mutant background decreased the Wg expression

by 1.6-fold as compared to L^2 mutant (*L^2; ey*-GAL4) (Figure 4 E, F). Similar trends were

seen with other novel newt genes (C1, C2, C3, and C5) when misexpressed in L^2

mutant background (Figure S8 C-F, C′-F′, G, G′). Thus, the newt genes can

downregulate *wg* expression in *Drosophila* L^2 mutant to partially rescue ventral eye-

loss-phenotype.

To further test our hypothesis, we misexpressed *wg* (the target identified in our

screen) along with *C4* transgene in the L^2 mutant background. As control, we

misexpressed *wg* in the developing eye (*ey>wg*), which result in reduced eye phenotype

(Figure 5 A, B) (Lee and Treisman, 2001; Singh et al., 2012a). Misexpressing *C4* in the

wild-type background where *wg* is ectopically induced (*+/+; ey>wg+C4*), rescues the

reduced eye phenotype (Figure 5 C, D) at a frequency of 16.8% (Figure S9 and Table

S1). Misexpression of *wg* in the L^2/+ mutants eye background (*L^2/+; ey>wg*) completely

eliminate the eye field as seen in the eye imaginal disc and the adult eye (Figure 5E, F)

whereas, misexpressing *C4* along with *wg* in the L^2/+ mutant eye disc (L^2/+*; ey>wg; C4*)

partially rescues the loss-of-ventral eye phenotype (Figure 5 G, H) in 21% flies (Figure

S9 and TableS1). Thus, newt genes can even downregulate Wg expression in the wild-

type (+/+*; ey>wg+C4*; Figure 5 C, D) as well as L^2 (L^2/+*; ey>wg; C4*; Figure 5 G, H)

background(s) where *wg* was ectopically co-expressed. *Drosophila* eye to promote

rescue of the loss-of-ventral eye phenotype.

Newt genes negatively regulate Wg/Wnt signaling

To investigate if newt genes regulate expression of *wg* alone or they regulate the Wg

signaling, we tested the antagonists, and agonists of the Wg-signaling pathway. The

rationale was to modulate the levels of Wg signaling and sample its effect on the L^2

mutant phenotype as well as the L^2 mutant phenotype where *C4* is misexpressed (L^2/+*;*

ey> C4) (Figure 6). Increasing the levels of Arm (a vertebrate homolog of β-Catenin, an

agonist of Wg signaling) in the wild-type eye (*ey>arm*) resulted in small eyes (Figure

S10 A, B) and in the L^2/+ background, (L^2/+*; ey>arm)* it results in elimination of the

entire eye field (Figure 6B, C). Misexpressing *C4* along with *arm* in the L^2/+ mutant

background (L^2/+*; ey>arm; C4*) partially rescues the loss-of-ventral eye phenotype

(Figure 6D, E) and the rescue frequency is 22.8% (Figure S11A and Table S1)*.*

Similarly, misexpressing *C4* and *arm* in the wild-type background (+/+*; ey>arm; C4*)

partially rescues the small eye phenotype defect caused by *ey>arm* phenotype (Figure

S10 C, D), and the rescue frequency is 21% (Figure S11A and Table S1). This data

suggests that newt gene can modulate the Wg signaling pathway components to

promote rescue of reduced eye phenotype both in L^2-mutant as well as the wild-type

background(s). However, the rescue frequency obtained by misexpressing newt gene in

addition to ectopically misexpressing *arm* (L^2/+*; ey>arm; C4*, 22.8%) and/or *wg* (L^2/+*;*

ey>wg; C4, 21%) (Figure S9, S11A and Table S1) is lower than the rescue frequency

obtained when only newt gene is misexpressed in the L^2-mutant background (L^2/+*;*

ey>arm; C4, 58.2%). This could be due to severity of loss-of-ventral eye phenotype

caused by the combined effect of L^2 mutation and gain-of-function of *wg*, or *arm.*

Blocking Wg signaling by misexpressing *shaggy (sgg)*, an antagonists of Wg
signaling, in the eye (*ey>sgg,* Figure S10E, F) (Hazelett et al., 1998; Singh et al., 2002;
Singh et al., 2006) or along with *C4* in the wild-type background (*+/+; ey>sgg; C4,*
Figure S10 G, H) does not affect the eye size. Misexpression of *sgg* in the L^2/+ mutants
(*L^2/+; ey>sgg*), significantly rescues the loss-of-ventral eye phenotype in 27.3% of flies
(Figure 6F, G, S11B and Table S1) (Singh et al., 2006). Similarly, misexpressing *C4*
and *sgg* in the L^2/+ mutant eye (*L^2/+; ey>sgg; C4*) rescues the loss-of-ventral eye
phenotype (Figure 6H, I) and rescue frequency increased dramatically from 27.3% to
71.5% (Figure S11B and Table S1). The transcription factor dTCF is the downstream
target of Wg signaling, and is downregulated by misexpression of the N-terminal deleted
dominant-negative form of TCF (dTCF^{DN}) (van de Wetering et al., 1997). Misexpression
of dTCF^{DN} alone in the eye (*ey>dTCF^{DN}*, Figure S10I, J) or dTCF^{DN} with *C4* (*+/+; ey>*
dTCF^{DN}; C4, Figure S10K, L) does not affect the eye size. Misexpressing *dTCF^{DN}* in the
eye of L^2/+ mutants (*L^2/+; ey>dTCF^{DN}*) significantly rescues the loss-of-ventral eye
phenotype in 21.8% of flies (Figure 6J, K, S11B and Table S1) (Singh et al., 2006).
Similarly, misexpressing C4 and *dTCF^{DN}* in the eye of L^2 mutants (*L^2/+; ey> dTCF^{DN};*
C4) rescues the loss-of-ventral eye phenotype (Figure 6L, M). As expected rescue
frequency again dramatically increased from 21.8% to 70.6% (Figure S11B and Table
S1).

Our results demonstrate that misexpression of newt genes when Wg signaling is
downregulated (using antagonists of Wg) in L^2 mutant background (*L^2/+; ey> sgg; C4,*
71.5%) or (*L^2/+; ey> dTCF^{DN}; C4*, 70.6%) significantly increased the rescue frequency
of the L^2 mutant phenotype. The rescue frequency obtained is greater than either
misexpressing candidate gene *(L^2/+; ey>C4,* 58.2%) alone or only misexpressing
negative regulators (*L^2/+; ey> sgg*, 27.3%) or (*L^2/+; ey> dTCF^{DN}*, 21.8%). The converse
phenotypes were seen when Wg signaling was upregulated. This strongly suggest that
newt gene(s) interact with the Wg signaling pathway to promote rescue of L^2 mutant
phenotype.

Porcupine (Porc) is involved in post-translational modification of Wg nascent
protein in the producing cell and thus facilitates its transport outside to the receiving cell.

Wg ligand can bind to its receptor on the receiving cell (Manoukian et al., 1995; Swarup

and Verheyen, 2012) thus modulating downstream pathway components (Figure 6A).

We blocked Wg transport from the producing cell using *porc porc^{RNAi}* and sampled its

effect on L^2 mutant phenotype, and L^2 mutant phenotype where C4 is misexpressed.

Misexpression of *porc^{RNAi}* alone (*ey> porc^{RNAi}*) (Figure S10M, N), and along with C4 in

the wild-type background (*+/+; ey> porc^{RNAi}; C4*) (Figure S10O, P) does not affect the

eye size. Misexpressing *porc^{RNAi}* in the L^2 mutant eye (*L^2/+; ey> porc^{RNAi}*) significantly

rescues the loss-of-ventral eye phenotype in 20.8% of flies (Figure 6N, O, S11B and

Table S1) whereas when *C4* is also misexpressed in the eye of L^2 mutants along with

porc^{RNAi} (*L^2/+; ey> porc^{RNAi}; C4*) rescued the loss-of-ventral eye phenotype (Figure 6P,

Q). The rescue frequency increased dramatically from 20.8% to 68.1% (Figure S11B,

and Table S1). Thus, newt genes can modulate levels of Wg signaling to promote

regeneration response in L^2 eye mutants.

Discussion

Notophthalmus viridescens have enormous genome size ($\sim c \times 10^{10}$ bases), a long reproductive cycle, and have limited genetic tools that makes it difficult to use this model to ascertain the molecular mechanism(s) behind the function of its novel genes (Mehta and Singh, 2019a). One of the strategies can be to introduce these genes into models with array of genetic tools and where regeneration potential is lower as seen in models like *Drosophila, Bombyx mori* etc. (Gopinathan et al., 1998; Kango-Singh et al., 2001; Mehta and Singh, 2019a; Singh et al., 2007). The rationale is to test the efficacy of these novel newt genes in triggering regeneration response in these model systems with lower regeneration potential. We and others have introduced human, plant and other vertebrate genes by transgenic approaches in *Drosophila* (Deshpande et al., 2019; Gogia et al., 2020b; Hughes et al., 2012; Sarkar et al., 2018; Tare et al., 2011). These transgenic flies have been utilized for targeted expression of foreign genes in flies using a targeted misexpression approach (Brand and Perrimon, 1993). The successful misexpression of the unique newt genes (vertebrates) in *Drosophila* using the same targeted misexpression approach has been reported earlier (Mehta et al., 2019; Mehta and Singh, 2019b).

To test the regeneration potential of these novel newt genes, we took a distinctive approach of using *Drosophila* model because the signaling pathways, which are involved in regeneration and/or tissue growth, patterning and development are evolutionarily conserved across the species (Kango-Singh and Singh, 2009; Mehta and Singh, 2019a; Wang and Hu, 2020). Even though humans *(Homo sapiens),* which evolved approx. 200,000 years ago (Stringer, 2016) and are 541 million years apart from *Drosophila* still share 75% of genes with flies (Pandey and Nichols, 2011). Despite the gap of 250 million years between newts and *Drosophila*, our studies suggest that the proteins encoded by unique newt genes are functional in *Drosophila* (Figure 1). Therefore, it is plausible that the pathways that newt genes can modulate in *Drosophila* might share parallels with their mechanism of action in newts. We found that the five newly identified genes from newt, can (i) rescue *L* mutant eye phenotype (early eye mutant) as well as GMR-*hid* GMR-GAL4 (late retinal degeneration) phenotype, (ii)

440   promote cell proliferation, and downregulate cell death to restore loss-of-ventral eye

441   phenotype, and (iii) downregulate Wg signaling (Figure 7). Our studies clearly

442   demonstrate the regeneration potential of these novel newt genes in triggering rescue

443   response in *Drosophila* model along the temporal axis.

Newt genes promote cell proliferation to restore *Drosophila* eye mutant

phenotype

Regeneration involves the formation of a blastema, a proliferating zone of

undifferentiated cells to restore the missing structures (Brockes and Kintner, 1986;

Mehta and Singh, 2019a; Sanchez Alvarado and Tsonis, 2006). Furthermore, a network

of newt gene(s) trigger proliferation response to promote lens regeneration (Eguchi and

Shingai, 1971; Grogg et al., 2005; Looso et al., 2013; Madhavan et al., 2006) A similar

proliferating response was observed when newt genes were misexpressed in the

Drosophila L^2 mutant eye (Figure 3C, and Figure S4). L^2 mutant flies exhibit a loss-of-

ventral eye phenotype, and misexpressing newt genes causes 4.2-5.8 fold increase in

number proliferating cells (Figure 3, 7 and Figure S4), which clearly mimics the classic

mechanism of epimorphic regeneration in the newt lens.

In our study, we found that newt genes can also modulate cell death in

Drosophila developing eye. When newt genes were misexpressed in the L^2 mutant

background (L^2; *ey>newt gene*), the rate of cell death was lower as compared to the L^2

mutant (L^2; *ey*-GAL4) eye (Figure 3E, F, H and Figure S4H-L, N), but minimally higher

compared to the positive control (*ey*-GAL4) (Figure 3F, G, H and Figure S4I-N). A

minimal rate of apoptosis is observed during the newt lens regeneration (Tsonis et al.,

2004). Our data suggest that newt genes hijack the evolutionarily conserved signaling

pathway(s) in *Drosophila* to regulate cell proliferation, and cell death. It is a strong

possibility that the same pathways may promote regeneration in newts.

Newt genes tightly regulate Wg signaling pathway in *Drosophila*

We screened for the various signaling pathways using the candidate approach as well as compared them to our transcriptomic analysis (Mehta et al., 2019). Among the various genetic modifiers of the L^2 mutant phenotype, we found *wg* as one of the possible candidate. Wg, a member of the evolutionarily conserved Wg/Wnt pathway, has been previously reported to promote regeneration, growth, development etc. (Harris et al., 2016; Schubiger et al., 2010; Singh et al., 2012b; Singh et al., 2018; Smith-Bolton et al., 2009; Yokoyama et al., 2007). Misexpresion of newt genes in *Drosophila* wild-type background modulate both agonists of Wg/Wnt signaling like *frizzled (fz),* and *microtubule star,*(*mts*) and antagonists like *Casein Kinase IIβ* (*CKIIβ*), and *sinaH* (Figure S7). This modulation in the levels of both antagonists as well as agonists can be a homeostatic reaction shown by the *Drosophila* to normalize expression levels of Wg/Wnt signaling pathway. As a result, changes in the eye growth after misexpressing newt genes in the wild-type was non-significant when compared to the wild-type control (*ey*-GAL4) (Figure 1). We did not observe any lethality, developmental defects etc.

Interestingly, when newt genes were misexpressed in L^2 mutant background, there was significant downregulation of Wg/Wnt expression (Figures 4, 7, and Figure S8), which resulted in rescue of the loss-of-ventral eye phenotype. It is known that overexpressing Wnt, and its receptor Fz promote lens regeneration in newts, whereas blocking Wnt pathway by expressing Dkk1 inhibits lens induction (Hayashi et al., 2006). Therefore, our study may have parallels with the regeneration mechanism in newts.

Furthermore, we found that when Wg pathway components are modulated in *Drosophila* along with the newt genes, it not only rescue the L^2 mutant loss-of-ventral eye phenotype but can also rescue the eye-loss caused by the activation of the Wg signaling pathway in the wild-type background. Hence, our data strongly suggest that the newt genes can modulate Wg signaling pathway in loss-of-ventral eye background. Interestingly, these newt genes did not show any phenotype when they were

496   misexpressed alone suggesting that regeneration response is triggered only to the cues
497   generated due to tissue damage.

Newt proteins exhibit non-cell-autonomous response

Genetic mosaic experimental approach, which resulted in the generation of clone(s) of
retinal neurons expressing high levels of newt genes within the L^2 mutant eye disc.
These experiments showed that clones expressing high levels of newt genes can
rescue the L^2 mutant phenotype, however the rescue was not limited to the clone itself
and spread into the neighboring cells where newt genes were not misexpressed. This
non-cell-autonomous rescue of the missing tissue is possible only if the newt proteins
are getting secreted from one cell to another and/or newt proteins are acting on the
signaling molecules. These newt proteins have signal peptides (Looso et al., 2013),
which can facilitate its transport from one cell to another. Similarly, we have reported
that these genes interact with Wg (Figure 4), which is a morphogen that can have a long
distance effect in the *Drosophila* eye (Zecca et al., 1996a).

It will be interesting to investigate, (a) if newt genes can regulate Wg signaling
pathway to promote regeneration in other tissues of *Drosophila* e.g. wing, leg etc., and
under the background of other congenital mutations (Schubiger et al., 2010; Smith-
Bolton et al., 2009), (b) if these proteins can rescue/regenerate missing tissues/organs
in other animals with low regenerative potentials. The pathways that have roles in
regeneration, growth, development are evolutionarily conserved throughout the animal
kingdom (Mehta and Singh, 2019a). For example, the Wg/Wnt signaling pathway, is
one such evolutionarily conserved pathway that apart from promoting regeneration in
hydra, planaria, newts, zebrafish, etc. has also been found to promote cell proliferation
during regeneration of mammalian muscles, liver, and bone (Polesskaya et al., 2003;
Sodhi et al., 2005; Zhong et al., 2006). Therefore, it raises the possibility that genetically
engineering genes from the highly regenerative animals that show regeneration in a
precisely controlled fashion might systematically regulate evolutionarily conserved
pathways in a foreign host to promote tissue regeneration and/or cell survival.

Materials and methods

Animals

Handling and operations of *Notophthalmus viridescens* (newts) have been described previously (Mehta et al., 2019; Sousounis et al., 2013). For our studies, *Notophthalmus viridescens* (newts) were purchased from Charles Sullivan Inc. newt farm. All procedures involving animals were approved by the University of Dayton Institutional Animal Care and Use Committee (IACUC; Protocol ID: 011-02). Newts were anesthetized in 0.1% (w/v) ethyl-3-aminobenzoate methanesulfonic acid (MS222; Sigma) dissolved in phosphate buffered saline. The surgical procedures were performed in anesthetized with MS222 newts, and appropriate procedures were undertaken to alleviate pain and distress while working with newts.

Generation of transgenic flies

The design and generation of transgenic flies (UAS-C1, UAS-C2, UAS-C3, UAS-C4, and UAS-C5) used in this study have been described (Mehta et al., 2019). To validate the successful insertion of newt genes in *Drosophila* by targeted misexpression approach, we generated transgenic fly like UAS-C1-5-V5 with 14 aa long V5-epitope sequence tagged towards the 3' end of the C4 ORF. The complete gene fragment (5'-C4-V5-3') was synthesized, and cloned into the XbaI and XhoI restriction sites in pUAST-attB plasmid. The pUAST-attB plasmid containing both the transgene and donor sequence (attB) was co-injected along with ϕC31 integrase mRNA into attP-containing recipient embryos, resulting in site-specific insertion of the transgene at Chromosome III (Fish et al., 2007; Groth et al., 2004). The integration and expression of transgene was verified by the western blots using V5 antibody.

***Drosophila* stocks**

In this study, we used transgenic flies UAS-C1, UAS-C2, UAS-C3, UAS-C4, UAS-C5,
and UAS-C1-5-V5). Other fly stocks used in this study are described in Flybase
(http://flybase.bio.indiana.edu). We used *Tub*-GAL4*/S-T* (Yang et al., 2019)*, L²/CyO;*
ey-GAL4*/ey*-GAL4*, L²/C4; ey*-GAL4*/ey*-GAL4*, UAS-GFP* (Singh et al., 2005; Singh and
Choi, 2003; Singh et al., 2006). *DE*-GAL4*/CyO; UAS-GFP/Tb* (Morrison and Halder,
2010), *UAS-wg-GFP* (Azpiazu and Morata, 1998) , *UAS-arm (Zecca et al., 1996b),*
UAS-sgg (Hazelett et al., 1998)*, UAS-dTCF^{DN}* (van de Wetering et al., 1997) *, UAS-p35*
(Hay et al., 1994).

Targeted misexpression studies

We used the GAL4/UAS system for targeted misexpression studies (Brand and
Perrimon, 1993). All GAL4/UAS crosses were performed at 25°C. In order to
misexpress the UAS-transgenes, we used *tub* –GAL4 (Yang et al., 2019) to ubiquitously
misexpress genes*, ey*-GAL4 (Hazelett et al., 1998) to misexpress transgene in the
developing eye, and *DE*-GAL4 (Morrison and Halder, 2010) that drives transgene in the
dorsal domain of the developing eye.

Genetic mosaic analysis

Clones were generated using the FLP/FRT system of mitotic recombination (Xu and
Rubin, 1993). To generate clones of C4 *in L²*-mutant eye, *yw* hsFLP; Act>y+>GAL4,
UAS-*GFP*/*CyO* females were crossed to *L²/CyO;* UAS-*C4/Tb* males. C4 is expressed in
all the GFP+ cells of eye-antennal imaginal disc.

Immunohistochemistry

Eye-antenna discs were dissected from wandering third instar larvae and stained
following the standard protocol (Singh et al., 2006). The eye imaginal discs were
dissected in cold 1X solution of Phosphate Buffered Saline (PBS) and fixed using 4% p-
formaldehyde (Electron Microscopy Sciences, Catalog no. 15710). Fixed tissues were

washed using PBS-T [0.2% Triton X-100 (Sigma, Catalog no. T8787) in 1XPBS] and

immunostained using primary antibodies: mouse anti-Phospho histone-3 (1:200)

(Sigma-Aldrich); Rabbit anti- *Drosophila* caspase-1 (DCP-1) (1:100) (Sigma-Aldrich);

rat anti-Elav (1:100) (Developmental Studies Hybridoma Bank); mouse anti-Wg (1:100)

(Developmental Studies Hybridoma Bank), and goat anti-Hth (a gift from H. Sun and R.

Mann). Secondary antibodies (Jackson Laboratories) used were goat anti-rat IgG

conjugated with Cy5 (1:200); donkey anti-rat IgG conjugated to FITC (1:200); donkey

anti-rat IgG conjugated to Cy5 (1:200), donkey anti-mouse IgG conjugated to Cy3

(1:200); donkey anti-goat IgG conjugated to Cy3 (1:200), and goat anti-rabbit IgG

conjugated with Cy5 (1:200). The discs were mounted in Vectashield (Vector labs,

H1000) and imaged using Olympus Fluoview 1000 microscope.

Western Blot

In this study, two different western blots were run. For blot 1, protein samples were

prepared from ($n = $ ~10) the wandering third instar larvae of transgenic fly UAS-C4-V5

as experimental sample, and *ey*-GAL4 was used as control. For blot 2, protein samples

were prepared from ($n = $ ~25) the adult heads of *ey*-GAL4, L^2; *ey*-GAL4 and L^2; *ey>C4*.

Protein samples were isolated by standardized protocol (Gogia et al., 2017), ran on

10% gel, and then transferred onto nitrocellulose membranes. For blot 1, the membrane

was then incubated overnight at 4°C with primary antibodies: 1:15000 rabbit anti-V5

antibody (SIGMA, Cat. # V8137), and 1:12000 mouse anti-α-Tubulin antibody (SIGMA,

Cat. # T5168). Followed by 1h incubation with secondary antibodies: horseradish

peroxidase conjugated goat anti–rabbit IgG (1:5000 dilution) secondary antibody, and

goat anti-mouse IgG-HRP (1:5000 dilution) (Santa Cruz Biotechnology, Cat. Sc-2005).

For blot 2, the membrane was then incubated overnight at 4°C with primary antibody

1:12000 mouse anti-α-Tubulin antibody (SIGMA, Cat. # T5168). Followed by

1h incubation with secondary antibody: horseradish peroxidase conjugated goat anti-

mouse IgG-HRP (1:5000 dilution) (Santa Cruz Biotechnology, Cat. Sc-2005). The blot

was then stripped and then incubated overnight with primary antibody 1:1000 mouse

anti-Wingless antibody (DSHB, Cat. # 4D4). Followed by 1h incubation with secondary

antibody 1:5000 horseradish peroxidase conjugated goat anti-mouse IgG-HRP (Santa

Cruz Biotechnology, Cat. Sc-2005). The signal in both the blots were detected using

Super Signal West Dura Extended Duration Substrate (ThermoFisher Scientific, Cat.

#34076). Images were captured using the UVP Imaging System. Wg levels were

quantified and normalized by using Licor Image Studio lite 5.2 software and this was

followed by statistical analysis.

613 TUNEL assay for detection of cell death

Apoptosis was detected by using Terminal deoxynucleotidyl transferase-mediated

dUTP nick-end labeling (TUNEL) assay (Tare et al., 2011; White et al., 2001). TUNEL

assay is based on the detection of fragmented DNA ends, which are characteristic of

apoptotic cells. A deoxynucleotidyl transferase (TdT) is used to add fluorescently

labeled nucleotides to free 3'-OH DNA ends in a template-independent manner. The

modified nucleotides can then be detected by fluorescence microscopy. Eye-antennal

discs, after secondary-antibody staining were blocked in 10% normal goat serum in

phosphate buffered saline with 0.2% Triton X100 (PBT) for an hour. After blocking,

samples were incubated in 0.1 M sodium citrate (Fisher Scientific) and 10% TritonX-100

for 30 minutes in dark at 65^{o}C. Samples were washed and incubated in TUNEL dilution

buffer (Roche Diagnostics, Catalog no. 11966006001) for 10 minutes at room

temperature. Followed by labeling with 50% TUNEL labeling solution for 30 minutes

(Roche Diagnostics, Catalog no. 12156792910) in dark at room temperature, and finally

samples were subjected to TdT enzyme solution (Roche Diagnostics) for 2 hours at

37^{0}C. After which samples were washed with PBS-T and mounted in Vectashield

(Vector labs, Catalog no. H1000).

630 Bright Field imaging

Adult flies were selected and were transferred to -20^{0}C for two hours, followed by

removal of legs and wings. The legs and wings were removed in order to get a clear

view of the compound eye. The Bright field pictures of adult heads were taken using a

Zeiss Apotome Imager Z1 (Wittkorn et al., 2015). Eye size was measured using ImageJ

software tools (http://rsb.info.nih.gov/ij/). To normalize the eye size values and neglect

any variations caused by overall different head size we took ratio of eye size to the head

size for each individual picture. Then relative ratio was calculated for all the treatments,

and the value reported is the relative fold change for each experimental treatment

compared to the control.

Statistical analysis

Statistical analysis was performed using student's t-test for independent samples.

Sample size used for the calculation was five in number (n=5), otherwise specified.

Statistical significance was determined with 95% confidence ($p < 0.05$).

Acknowledgments

We thank Bloomington *Drosophila* Stock Center (BDSC) for *Drosophila* strains, and the Developmental Studies Hybridoma Bank (DSHB) for antibodies. We would like to thank to Katia Del Rio-Tsonis, Labib Rouhana, Madhuri Kango-Singh, Meghana Tare, Neha Gogia, Shilpi Verghese, Pothitos Pitychoutis, Deepika K Sodhi for critical comments on the manuscript. We also thank Justin Kumar, Y. Henry Sun, and Kyung Ok Cho for gift of fly strains and antibodies. Confocal microscopy was supported by core facility at University of Dayton.

Funding resources

A.S. is supported by NIH1R15GM124654-01 from NIH, Schuellein Chair Endowment Fund and STEM Catalyst Grant from the University of Dayton.

References

Azpiazu, N., and Morata, G. (1998). Functional and regulatory interactions between Hox and extradenticle genes. Genes Dev *12*, 261-273.

Baker, N.E. (1987). Molecular cloning of sequences from wingless, a segment polarity gene in Drosophila: the spatial distribution of a transcript in embryos. EMBO J *6*, 1765-1773.

Baonza, A., and Freeman, M. (2005). Control of Cell Proliferation in the Drosophila Eye by Notch Signaling. Developmental Cell *8*, 529-539.

Bejsovec, A. (2006). Flying at the head of the pack: Wnt biology in Drosophila. Oncogene *25*, 7442-7449.

Bejsovec, A. (2013). Wingless/Wnt signaling in Drosophila: the pattern and the pathway. Mol Reprod Dev *80*, 882-894.

Bergmann, A., Agapite, J., McCall, K., and Steller, H. (1998). The Drosophila gene hid is a direct molecular target of Ras-dependent survival signaling. Cell *95*, 331-341.

Bier, E. (2005). Drosophila, the golden bug, emerges as a tool for human genetics. Nat Rev Genet *6*, 9-23.

Brand, A.H., and Perrimon, N. (1993). Targeted gene expression as a means of altering cell fates and generating dominant phenotypes. Development *118*, 401.

Brockes, J.P., and Kintner, C.R. (1986). Glial growth factor and nerve-dependent proliferation in the regeneration blastema of urodele amphibians. Cell *45*, 301-306.

Bryant, D.M., Johnson, K., DiTommaso, T., Tickle, T., Couger, M.B., Payzin-Dogru, D., Lee, T.J., Leigh, N.D., Kuo, T.H., Davis, F.G., *et al.* (2017). A Tissue-Mapped Axolotl De Novo Transcriptome Enables Identification of Limb Regeneration Factors. Cell reports *18*, 762-776.

Casco-Robles, R.M., Watanabe, A., Eto, K., Takeshima, K., Obata, S., Kinoshita, T., Ariizumi, T., Nakatani, K., Nakada, T., Tsonis, P.A., *et al.* (2018). Novel erythrocyte clumps revealed by an orphan gene Newtic1 in circulating blood and regenerating limbs of the adult newt. Sci Rep *8*, 7455-7455.

Cavodeassi, F., Diez Del Corral, R., Campuzano, S., and Dominguez, M. (1999). Compartments and organising boundaries in the Drosophila eye: the role of the homeodomain Iroquois proteins. Development *126*, 4933-4942.

Cohen, S.M. (1993). Imaginal disc development. Paper presented at: Bate, Martinez Arias, 1993.

Crosio, C., Fimia, G.M., Loury, R., Kimura, M., Okano, Y., Zhou, H., Sen, S., Allis, C.D., and Sassone-Corsi, P. (2002). Mitotic phosphorylation of histone H3: spatio-temporal regulation by mammalian Aurora kinases. Molecular and cellular biology *22*, 874-885.

da Silva, S.M., Gates, P.B., and Brockes, J.P. (2002). The Newt Ortholog of CD59 Is Implicated in Proximodistal Identity during Amphibian Limb Regeneration. Developmental Cell *3*, 547-555.

Deshpande, P., Gogia, N., and Singh, A. (2019). Exploring the efficacy of natural products in alleviating Alzheimer's disease. Neural regeneration research *14*, 1321-1329.

Echeverri, K., and Tanaka, E.M. (2005). Proximodistal patterning during limb regeneration. Developmental Biology *279*, 391-401.

Eguchi, G., and Shingai, R. (1971). CELLULAR ANALYSIS ON LOCALIZATION OF LENS FORMING POTENCY IN THE NEWT IRIS EPITHELIUM*,**. Development, Growth & Differentiation *13*, 337-350.

Elewa, A., Wang, H., Talavera-López, C., Joven, A., Brito, G., Kumar, A., Hameed, L.S., Penrad-Mobayed, M., Yao, Z., Zamani, N., *et al.* (2017). Reading and editing the Pleurodeles waltl genome reveals novel features of tetrapod regeneration. Nature Communications *8*, 2286.

Evans, T., Johnson, A.D., and Loose, M. (2018). Virtual Genome Walking across the 32 Gb Ambystoma mexicanum genome; assembling gene models and intronic sequence. Sci Rep *8*, 618.

Fish, M.P., Groth, A.C., Calos, M.P., and Nusse, R. (2007). Creating transgenic Drosophila by microinjecting the site-specific φC31 integrase mRNA and a transgene-containing donor plasmid. Nature Protocols *2*, 2325.

Garza-Garcia, A., Harris, R., Esposito, D., Gates, P.B., and Driscoll, P.C. (2009). Solution structure and
phylogenetics of Prod1, a member of the three-finger protein superfamily implicated in salamander limb
regeneration. PloS one *4*, e7123.
Gogia, N., Puli, O.R., Raj, A., and, and Singh, A. (2020a). Generation of Third Dimension: Axial Patterning
in the Developing Drosophila Eye. In Molecular Genetics of Axial Patterning, Growth and Disease in the
Drosophila Eye, A. Singh, and, and M. Kango-Singh, eds. (Springer NewYork Springer), pp. 53-95.
Gogia, N., Sarkar, A., Mehta, A.S., Ramesh, N., Deshpande, P., Kango-Singh, M., Pandey, U.B., and Singh,
A. (2020b). Inactivation of Hippo and cJun-N-terminal Kinase (JNK) signaling mitigate FUS mediated
neurodegeneration in vivo. Neurobiol Dis *140*, 104837.
Gogia, N., Sarkar, A., and Singh, A. (2017). An Undergraduate Cell Biology Lab: Western
Blotting to Detect Proteins from Drosophila Eye. Drosophila Information Service.
Gopinathan, K.P., Joy, O., and Singh, A. (1998). Developmental aspects of Mulberry and Nonmulberry
Silkworm Species: A comparative study. In "Genome Analysis in Eukaryotes: Developmental and
evolutionary aspects", R.N. Chatterjee, and L. Sanchez, eds. (New York: Springer-Verlag New York,
Incorporated), pp. 59-91.
Grether, M.E., Abrams, J.M., Agapite, J., White, K., and Steller, H. (1995). The head involution defective
gene of Drosophila melanogaster functions in programmed cell death. Genes Dev *9*, 1694-1708.
Grogg, M.W., Call, M.K., Okamoto, M., Vergara, M.N., Del Rio-Tsonis, K., and Tsonis, P.A. (2005). BMP
inhibition-driven regulation of six-3 underlies induction of newt lens regeneration. Nature *438*, 858-862.
Groth, A.C., Fish, M., Nusse, R., and Calos, M.P. (2004). Construction of transgenic Drosophila by using
the site-specific integrase from phage phiC31. Genetics *166*, 1775-1782.
Harris, R.E., Setiawan, L., Saul, J., and Hariharan, I.K. (2016). Localized epigenetic silencing of a damage-
activated WNT enhancer limits regeneration in mature Drosophila imaginal discs. eLife *5*, e11588.
Hay, B.A., Wassarman, D.A., and Rubin, G.M. (1995). Drosophila homologs of baculovirus inhibitor of
apoptosis proteins function to block cell death. Cell *83*, 1253-1262.
Hay, B.A., Wolff, T., and Rubin, G.M. (1994). Expression of baculovirus P35 prevents cell death in
Drosophila. Development *120*, 2121-2129.
Hayashi, T., Mizuno, N., Takada, R., Takada, S., and Kondoh, H. (2006). Determinative role of Wnt signals
in dorsal iris-derived lens regeneration in newt eye. Mech Dev *123*, 793-800.
Haynie, J.L., and Bryant, P.J. (1986). Development of the eye-antenna imaginal disc and morphogenesis
of the adult head in Drosophila melanogaster. J Exp Zool *237*, 293-308.
Hazelett, D., Bourouis, M., Walldorf, U., and Treisman, J. (1998). decapentaplegic and wingless are
regulated by eyes absent and eyegone and interact to direct the pattern of retinal differentiation in the
eye disc. . Development *125*, 3741-3751.
Held, L.I.J. (2002). The eye disc. In Imaginal Disc, L.I. Held, ed. (Cambridge University Press), pp. 197-236.
Hughes, T.T., Allen, A.L., Bardin, J.E., Christian, M.N., Daimon, K., Dozier, K.D., Hansen, C.L., Holcomb,
L.M., and Ahlander, J. (2012). Drosophila as a genetic model for studying pathogenic human viruses.
Virology *423*, 1-5.
Kango-Singh, M., and Singh, A. (2009). Regulation of organ size: Insights from the Drosophila Hippo
signaling pathway. Developmental Dynamics *238*, 1627-1637.
Kango-Singh, M., Singh, A., and Gopinathan, K.P. (2001). The wings of Bombyx mori develop from larval
discs exhibiting an early differentiated state: a preliminary report. J Biosci *26*, 167-177.
Keinath, M.C., Timoshevskiy, V.A., Timoshevskaya, N.Y., Tsonis, P.A., Voss, S.R., and Smith, J.J. (2015).
Initial characterization of the large genome of the salamander Ambystoma mexicanum using shotgun
and laser capture chromosome sequencing. Sci Rep *5*, 16413.
Kim, J.-Y., Jeong, H.S., Chung, T., Kim, M., Lee, J.H., Jung, W.H., and Koo, J.S. (2017). The value of
phosphohistone H3 as a proliferation marker for evaluating invasive breast cancers: A comparative study
with Ki67. Oncotarget *8*, 65064-65076.

Kumar, A., Godwin, J.W., Gates, P.B., Garza-Garcia, A.A., and Brockes, J.P. (2007). Molecular basis for the
nerve dependence of limb regeneration in an adult vertebrate. Science *318*, 772-777.
Kumar, J.P. (2013). Catching the next wave: patterning of the Drosophila eye by the morphogenetic
furrow. In In Molecular genetics of axial patterning, growth and disease in the Drosophila eye, A.a.K.-S.
Singh, Madhuri, ed. (NewYork: Springer), pp. 75-97.
Kumar, J.P. (2018). The fly eye: Through the looking glass. Developmental Dynamics *247*, 111-123.
Kumar, J.P. (2020a). Catching the next wave: patterning of the Drosophila eye by the morphogenetic
furrow. In Molecular Genetics of Axial Patterning, Growth and Disease in the Drosophila Eye, A. Singh,
and, and M. Kango-Singh, eds. (Springer NewYork Heidelberg Dordrecht London: Springer International
Publishing), pp. 97-120.
Kumar, J.P. (2020b). Ghost in the Machine: The Peripodal Membrane. In Molecular Genetics of Axial
Patterning, Growth and Disease in the Drosophila Eye, A. Singh, and, and M. Kango-Singh, eds. (Springer
NewYork Heidelberg Dordrecht London: Springer International Publishing), pp. 121-142.
Lee, J.D., and Treisman, J.E. (2001). The role of Wingless signaling in establishing the anteroposterior and
dorsoventral axes of the eye disc. Development *128*, 1519-1529.
Lenz, S., Karsten, P., Schulz, J.B., and Voigt, A. (2013). Drosophila as a screening tool to study human
neurodegenerative diseases. Journal of Neurochemistry *127*, 453-460.
Looso, M., Preussner, J., Sousounis, K., Bruckskotten, M., Michel, C.S., Lignelli, E., Reinhardt, R., Höffner,
S., Krüger, M., Tsonis, P.A., *et al.* (2013). A de novo assembly of the newt transcriptome combined with
proteomic validation identifies new protein families expressed during tissue regeneration. Genome
biology *14*, R16-R16.
Ma, C., and Moses, K. (1995). Wingless and patched are negative regulators of the morphogenetic
furrow and can affect tissue polarity in the developing Drosophila compound eye. Development *121*,
2279-2289.
Madhavan, M., Haynes, T.L., Frisch, N.C., Call, M.K., Minich, C.M., Tsonis, P.A., and Del Rio-Tsonis, K.
(2006). The role of Pax-6 in lens regeneration. Proceedings of the National Academy of Sciences *103*,
14848.
Manoukian, A.S., Yoffe, K.B., Wilder, E.L., and Perrimon, N. (1995). The porcupine gene is required for
wingless autoregulation in Drosophila. Development *121*, 4037-4044.
Matsunami, M., Suzuki, M., Haramoto, Y., Fukui, A., Inoue, T., Yamaguchi, K., Uchiyama, I., Mori, K.,
Tashiro, K., Ito, Y., *et al.* (2019). A comprehensive reference transcriptome resource for the Iberian
ribbed newt Pleurodeles waltl, an emerging model for developmental and regeneration biology. DNA
Res *26*, 217-229.
Maurel-Zaffran, C., and Treisman, J.E. (2000). pannier acts upstream of wingless to direct dorsal eye disc
development in Drosophila. Development *127*, 1007-1016.
Mehta, A.S., Luz-Madrigal, A., Li, J.-L., Tsonis, P.A., and Singh, A. (2019). Comparative transcriptomic
analysis and structure prediction of novel Newt proteins. PloS one *14*, e0220416.
Mehta, A.S., and Singh, A. (2019a). Insights into regeneration tool box: An animal model approach.
Developmental Biology *453*, 111-129.
Mehta, A.S., and Singh, A. (2019b). Regeneration Data: RNA Sequences [Internet] (University of Dayton:
eCommons).
Milner, M.J., Bleasby, A.J., and Pyott, A. (1983). The role of the peripodial membrane in the
morphogenesis of the eye-antennal disc ofDrosophila melanogaster. Wilehm Roux Arch Dev Biol *192*,
164-170.
Morrison, C.M., and Halder, G. (2010). Characterization of a dorsal-eye Gal4 Line in Drosophila. Genesis
48, 3-7.
Moses, K., and Rubin, G.M. (1991). Glass encodes a site-specific DNA-binding protein that is regulated in
response to positional signals in the developing Drosophila eye. Genes Dev *5*, 583-593.

Nowoshilow, S., Schloissnig, S., Fei, J.-F., Dahl, A., Pang, A.W.C., Pippel, M., Winkler, S., Hastie, A.R.,
Young, G., Roscito, J.G., *et al.* (2018). The axolotl genome and the evolution of key tissue formation
regulators. Nature *554*, 50-55.
Pandey, U.B., and Nichols, C.D. (2011). Human disease models in Drosophila melanogaster and the role
of the fly in therapeutic drug discovery. Pharmacol Rev *63*, 411-436.
Polesskaya, A., Seale, P., and Rudnicki, M.A. (2003). Wnt signaling induces the myogenic specification of
resident CD45+ adult stem cells during muscle regeneration. Cell *113*, 841-852.
Ready, D.F., Hanson, T.E., and Benzer, S. (1976). Development of the Drosophila retina, a
neurocrystalline lattice. Developmental Biology *53*, 217-240.
Sanchez Alvarado, A., and Tsonis, P.A. (2006). Bridging the regeneration gap: genetic insights from
diverse animal models. Nat Rev Genet *7*, 873-884.
Sanor, L.D., Flowers, G.P., and Crews, C.M. (2020). Multiplex CRISPR/Cas screen in regenerating haploid
limbs of chimeric Axolotls. eLife *9*, e48511.
Sarkar, A., Gogia, N., Glenn, N., Singh, A., Jones, G., Powers, N., Srivastava, A., Kango-Singh, M., and
Singh, A. (2018). A soy protein Lunasin can ameliorate amyloid-beta 42 mediated neurodegeneration in
Drosophila eye. Sci Rep *8*, 13545.
Sarkar, A., Irwin, M., Singh, A., Riccetti, M., and Singh, A. (2016). Alzheimer's disease: the silver tsunami
of the 21(st) century. Neural Regen Res *11*, 693-697.
Sarkissian, T., Timmons, A., Arya, R., Abdelwahid, E., and White, K. (2014). Detecting apoptosis in
Drosophila tissues and cells. Methods *68*, 89-96.
Schubiger, M., Sustar, A., and Schubiger, G. (2010). Regeneration and transdetermination: the role of
wingless and its regulation. Developmental biology *347*, 315-324.
Singh, A. (2012). Neurodegeneration- A Means to an End. J Cell Science & Therapy *3*, 10000e10107.
Singh, A., Chan, J., Chern, J.J., and Choi, K.-W. (2005). Genetic Interaction of Lobe
With Its Modifiers in Dorsoventral Patterning and Growth of the Drosophila Eye. Genetics *171*, 169.
Singh, A., and Choi, K.-W. (2003). Initial state of the Drosophila eye before dorsoventral specification is
equivalent to ventral. Development *130*, 6351.
Singh, A., and Irvine, K.D. (2012). Drosophila as a model for understanding development and disease.
Dev Dyn *241*, 1-2.
Singh, A., Kango-Singh, M., Parthasarathy, R., and Gopinathan, K.P. (2007). Larval legs of mulberry
silkworm Bombyx mori are prototypes for the adult legs. Genesis *45*, 169-176.
Singh, A., Kango-Singh, M., and Sun, Y.H. (2002). Eye suppression, a novel function of teashirt, requires
Wingless signaling. Development *129*, 4271-4280.
Singh, A., Shi, X., and Choi, K.W. (2006). Lobe and Serrate are required for cell survival during early eye
development in Drosophila. Development *133*, 4771-4781.
Singh, A., Tare, M., Kango-Singh, M., Son, W.-S., Cho, K.-O., and Choi, K.-W. (2011). Opposing
interactions between homothorax and Lobe define the ventral eye margin of Drosophila eye.
Developmental biology *359*, 199-208.
Singh, A., Tare, M., Puli, O.R., and Kango-Singh, M. (2012a). A glimpse into dorso-ventral patterning of
the Drosophila eye. Developmental dynamics : an official publication of the American Association of
Anatomists *241*, 69-84.
Singh, B.N., Doyle, M.J., Weaver, C.V., Koyano-Nakagawa, N., and Garry, D.J. (2012b). Hedgehog and
Wnt coordinate signaling in myogenic progenitors and regulate limb regeneration. Developmental
biology *371*, 23-34.
Singh, B.N., Weaver, C.V., Garry, M.G., and Garry, D.J. (2018). Hedgehog and Wnt Signaling Pathways
Regulate Tail Regeneration. Stem Cells and Development *27*, 1426-1437.
Smith-Bolton, R.K., Worley, M.I., Kanda, H., and Hariharan, I.K. (2009). Regenerative growth in
Drosophila imaginal discs is regulated by Wingless and Myc. Developmental cell *16*, 797-809.

Smith, J.J., Putta, S., Zhu, W., Pao, G.M., Verma, I.M., Hunter, T., Bryant, S.V., Gardiner, D.M., Harkins,
T.T., and Voss, S.R. (2009). Genic regions of a large salamander genome contain long introns and novel
genes. BMC Genomics *10*, 19-19.

Smith, J.J., Timoshevskaya, N., Timoshevskiy, V.A., Keinath, M.C., Hardy, D., and Voss, S.R. (2019). A
chromosome-scale assembly of the axolotl genome. Genome research *29*, 317-324.

Sodhi, D., Micsenyi, A., Bowen, W.C., Monga, D.K., Talavera, J.C., and Monga, S.P. (2005). Morpholino
oligonucleotide-triggered beta-catenin knockdown compromises normal liver regeneration. J Hepatol
43, 132-141.

Sousounis, K., Michel, C.S., Bruckskotten, M., Maki, N., Borchardt, T., Braun, T., Looso, M., and Tsonis,
P.A. (2013). A microarray analysis of gene expression patterns during early phases of newt lens
regeneration. Molecular vision *19*, 135-145.

Stringer, C. (2016). The origin and evolution of Homo sapiens. Philos Trans R Soc Lond B Biol Sci *371*,
20150237.

Swarup, S., and Verheyen, E.M. (2012). Wnt/Wingless signaling in Drosophila. Cold Spring Harb Perspect
Biol *4*.

Tare, M., Modi, R.M., Nainaparampil, J.J., Puli, O.R., Bedi, S., Fernandez-Funez, P., Kango-Singh, M., and
Singh, A. (2011). Activation of JNK signaling mediates amyloid-ss-dependent cell death. PloS one *6*,
e24361.

Tare, M., Puli, O.R., and, and Singh, A. (2013). Molecular Genetic Mechanisms of Axial Patterning:
Mechanistic Insights into Generation of Axes in the Developing Eye. In Molecular Genetics of Axial
Patterning, Growth and Disease in the Drosophila Eye, A. Singh, and, and M. Kango-Singh, eds. (Springer
NewYork Heidelberg Dordrecht London: Springer), pp. 37-75.

Treisman, J.E., and Rubin, G.M. (1995). wingless inhibits morphogenetic furrow movement in the
Drosophila eye disc. Development *121*, 3519-3527.

Tsonis, P.A., Madhavan, M., Tancous, E.E., and Del Rio-Tsonis, K. (2004). A newt's eye view of lens
regeneration. Int J Dev Biol *48*, 975-980.

van de Wetering, M., Cavallo, R., Dooijes, D., van Beest, M., van Es, J., Loureiro, J., Ypma, A., Hursh, D.,
Jones, T., Bejsovec, A., *et al.* (1997). Armadillo Coactivates Transcription Driven by the Product of the
Drosophila Segment Polarity Gene dTCF. Cell *88*, 789-799.

Vollmer, J., Fried, P., Sánchez-Aragón, M., Lopes, C.S., Casares, F., and Iber, D. (2016). A quantitative
analysis of growth control in the Drosophila eye disc. Development *143*, 1482.

Wang, H., Wang, X., Zhang, K., Wang, Q., Cao, X., Wang, Z., Zhang, S., Li, A., Liu, K., and Fang, Y. (2019).
Rapid depletion of ESCRT protein Vps4 underlies injury-induced autophagic impediment and Wallerian
degeneration. Sci Adv *5*, eaav4971-eaav4971.

Wang, W., and Hu, C.K. (2020). Changes in regeneration-responsive enhancers shape regenerative
capacities in vertebrates. *369*.

Weisrock, D.W., Papenfuss, T.J., Macey, J.R., Litvinchuk, S.N., Polymeni, R., Ugurtas, I.H., Zhao, E.,
Jowkar, H., and Larson, A. (2006). A molecular assessment of phylogenetic relationships and lineage
accumulation rates within the family Salamandridae (Amphibia, Caudata). Mol Phylogenet Evol *41*, 368-
383.

White, K., Grether, M.E., Abrams, J.M., Young, L., Farrell, K., and Steller, H. (1994). Genetic control of
programmed cell death in Drosophila. Science *264*, 677-683.

White, K., Lisi, S., Kurada, P., Franc, N., and Bangs, P. (2001). Methods for studying apoptosis and
phagocytosis of apoptotic cells in Drosophila tissues and cell lines. In Methods in Cell Biology (Academic
Press), pp. 321-338.

Wittkorn, E., Sarkar, A., Garcia, K., Kango-Singh, M., and Singh, A. (2015). The Hippo pathway effector Yki
downregulates Wg signaling to promote retinal differentiation in the Drosophila
eye. Development *142*, 2002.

Xu, T., and Rubin, G.M. (1993). Analysis of genetic mosaics in developing and adult Drosophila tissues.
Development *117*, 1223-1237.
Yang, C.-P., Chen, M.-S., Liaw, G.-J., Chen, S.-F., Chou, G., and Fan, S.-S. (2005). Using Drosophila eye as a
model system to characterize the function of mars gene in cell-cycle regulation. Experimental Cell
Research *307*, 183-193.
Yang, F., Quan, Z., Huang, H., He, M., Liu, X., Cai, T., and Xi, R. (2019). Ovaries absent links dLsd1 to HP1a
for local H3K4 demethylation required for heterochromatic gene silencing. eLife *8*, e40806.
Yeates, C.J., Sarkar, A., Kango-Singh, M., and Singh, A. (2019). Unravelling Alzhiemer's Disease using
Drosophila. In Insights into Human Neurodegeneration: Lessons Learnt from Drosophila, M. Mutsuddi,
and, and A. Mukherjee, eds. (Springer Singapore: Springer), pp. 251-277.
Yokoyama, H., Ogino, H., Stoick-Cooper, C.L., Grainger, R.M., and Moon, R.T. (2007). Wnt/beta-catenin
signaling has an essential role in the initiation of limb regeneration. Developmental biology *306*, 170-
178.
Zecca, M., Basler, K., and Struhl, G. (1996a). Direct and long-range action of a wingless morphogen
gradient. Cell *87*, 833-844.
Zecca, M., Basler, K., and Struhl, G. (1996b). Direct and Long-Range Action of a Wingless Morphogen
Gradient. Cell *87*, 833-844.
Zhong, N., Gersch, R.P., and Hadjiargyrou, M. (2006). Wnt signaling activation during bone regeneration
and the role of Dishevelled in chondrocyte proliferation and differentiation. Bone *39*, 5-16.

Figure legends:

Figure 1. Novel newt genes rescues L^2 mutant's loss-of-ventral eye phenotype.

(A, B) Schematic representation of generating transgenic *Drosophila* carrying novel

newt (*Notophthalmus viridescens*) candidate genes (*C1, C2, C3, C4,* and *C5*).

Bright field image of a (C) wild type adult eye as positive control (*ey*-GAL4), (D-H) adult

eye phenotype when *ey*-Gal4 drives expression of newt transgenes (D) UAS-*C1*

(ey>C1), (E) UAS-*C2* (*ey>C2*), (F) UAS-*C3 (ey>C3)*, (G) UAS-*C4 (ey>C4)*, and (H)

UAS-*C5 (ey>C5)*. (I) The bar graph shows there is no significant (ns) change in eye size

between control and where ey-GAL4 drives expression of newt transgenes (*ey>C1-C5*).

Bright field image of a (J) L^2 mutant (*L^2; ey*-GAL4*)* adult eye where there is preferential

loss-of- ventral eye phenotype, (K-O) L^2 mutant (*L^2; ey*-GAL4*)* background where newt

transgene (K) UAS-*C1 (L^2; ey >C1)*, (L) UAS-*C2 (L^2; ey >C2)*, (M) UAS-*C3 (L^2; ey >C3)*,

(N) UAS-*C4 (L^2;ey>C4)*, and (O) UAS-*C5 (L^2;ey>C5)* are expressed in the developing

eye using ey-GAL4. (P) L^2 mutant adult eye is half the adult eye size of the *ey*-GAL4

which serve as positive control. The bar graph clearly displays the significant increase in

the adult eye size of L^2-mutant background where unique newt genes were

misexpressed compared to the adult eye size of the L^2-mutant only (*L^2; ey*-GAL4).

Eye size was measured using ImageJ software tools (http://rsb.info.nih.gov/ij/). Sample

size used for statistical analysis was five (n=5). Statistical analysis was performed using

student's t-test for independent samples. Statistical significance was determined with

95% confidence (p<0.05). Error bars show standard deviation, and symbols above the

error bar signify as: ns is non-significant p value, * signifies p value <0.05, ** signifies p

value <0.005, *** signifies p value <0.0005 respectively. In this study *ey* is used as a

driver to misexpress *newt genes* in the developing *Drosophila* eye. All the images are

displayed in same polarity as dorsal domain-towards top, and ventral domain-towards

bottom. Scale bar= 100 **μm.**

See also Figure S1, S2, and S3 and Table S1

Figure 2. Misexpression of newt genes exhibits non-cell-autonomous rescue. (A,

B, B′, B″) Misexpression of *C4* in the dorsal half using *DE*-GAL4 driver results in (A)

Adult eye and (B, B′, B″) developing third instar larval eye imaginal disc (as positive

control). Note that Green fluorescence protein (GFP) (green) reporter marks the DE-

GAL4 driver domain and proneural marker Embryonic lethal abnormal visual system

(Elav) (red) marks the retinal fate. (B′) is a single channel confocal image showing GFP

expression, and (B″) is a single channel confocal image showing Elav expression. (C,

D, D′, D″) L^2 mutant eye where there is no misexpression of newt genes (L^2/+*;*

DE>GFP) (as negative control). (C) Adult eye bright field image and (D, D′, D″) third

instar eye disc of L^2 mutant (L^2/+*; DE*>GFP) exhibiting loss-of-ventral eye phenotype

and red dotted line marks the boundary of eye. GFP reporter (green) marks domain of

DE-GAL4 expression domain in L^2 mutant background. . (E, F, F′, F″) Misexpression of

C4 in dorsal eye domain of L^2 mutant (L^2/+*; DE*>GFP; C4) exhibits non-cell autonomous

rescue of loss-of-ventral eye phenotype in (E) Adult eye (white dotted boundary) and

(F′, F′, F″) third instar eye disc. White dotted boundary marks the significantly rescued

loss-of-ventral eye phenotype.   (G-J) Genetic mosaic "Flp out" somatic clones of

(G,G′,G″) newt gene C4 in the developing eye disc wild-type, (H,H′,H″) GFP in L^2

background (L^2/+*; hsflp*>GFP, served as the negative control) and (I,I′,I″ & J,J′,J″) newt

gene *C4* clones in L^2 mutant (L^2/+*; hsflp*>GFP*;C4*) background. All clones are marked

by GFP reporter (green). Note that misexpressing *C4* rescue missing photoreceptor

cells in a non- cell-autonomous manner, as photoreceptor cells rescue marked by Elav

(in red) extends outside the boundary of the clone marked by GFP (in green). (J, J′, J″)

exhibits the magnified view of the *C4* misexpressing clone (green) in L^2/+*; hsflp*>GFP*;*

C4 background. All the images are displayed in same polarity as dorsal domain-towards

top, and ventral domain-towards bottom. Scale bar= 100 **μm,**

Figure 3. Newt genes promote cell proliferation and downregulate cell death to

promote rescue of L^2 mutant phenotype of loss-of-ventral eye. (A-D) PH-3 staining

(blue), as a marker to calculate cell proliferation and mitotic index, in (A) L^2 mutant eye

disc (red dotted boundary marking eye boundary), (B) L^2 mutant eye disc where C4

gene is misexpressed (L^2; *ey*>C4), and (C) Wild type (*ey*-GAL4) eye disc which serve

as control. A′-C′) single channel confocal images demonstrating PH-3 stained nuclei. (B)

White dotted boundary represents the partially rescued loss-of-ventral eye phenotype.

The zone of interest that was considered to count number of dividing cells is marked in

yellow dotted boundary. (D) Bar graph showing relative increase in the mitotic index

(MI) when newt genes are misexpressed in L^2 mutant background. MI is calculated as

ratio of cells dividing (stained by PH-3 in blue) to total rows of ommatidial cells (stained

by Elav in red). There is no significant change in MI between the L^2 mutant, and wild-

type (*ey*-GAL4) eye disc, but comparatively there is robust increase in the MI inL^2;

ey>C4 background. (E-H) Terminal deoxynucleotidyl transferase-mediated dUTP nick-

end labeling (TUNEL) staining as a marker for cell death assay. TUNEL staining

(shown in green) marks the dying nuclei in eye imaginal disc of (E) L^2 mutant, (F) L^2

mutant where C4 gene is misexpressed (L^2; *ey*>C4), and (G) Wild type (*ey*-GAL4) eye

disc. Red dotted boundary represents ventral-eye-loss area, and white dotted boundary

represent the rescued loss-of-ventral eye region. (E′-G′)The tunnel positive cells are

counted within the zone of interest marked within yellow dotted boundary as shown in

single filter confocal images. (H) bar graph showing relative decrease in the cell death

when newt gene is misexpressed under the L^2-mutant background. The downregulation

in cell death is comparable to the cell death happening in the normal wild type

background

Statistical analysis was performed using student's t-test for independent samples.

Sample size used for the calculation was five in number (n=5). Statistical significance

was determined with 95% confidence (p<0.05). All bar graphs show relative fold change

in cell proliferation and cell death as average between 5 samples. Error bars show

standard deviation, and symbols above the error bar ** signify p value <0.005,   ***

signify p value <0.0005, and ns signify p value > 0.05 respectively. All the images are

displayed in same polarity as dorsa domain-towards top, and ventral domain-towards

bottom. Scale bar= 100 **μm**

1006

Figure 4. Newt genes downregulate Wg expression to rescue loss-of-ventral eye

phenotype.   (A-D) Immunohistochemistry showing Wg modulation. (A-C) Eye disc

showing Wg staining in green, and Elav in red (A′-C′) are single channel confocal

images showing Wg expression. The green window is the area of interest that is utilized

to calculate (D, D′) integrative density (intensity plot). The bar graph represent the Wg

intensity at the ventral margin (within the green box) in (A, A′) wild type fly (*ey*-GAL4),

(B, B′) L^2-mutant fly, (C, C′) Misexpression of newt gene (*C4*) in the L^2 mutation

background. Note that *C4* misexpression downregulates Wg expression. (E, F) Western

blot (WB) to semi quantitate modulation of Wg expression levels. (E) The blot shows the

expression of Wg protein (C4-V5) in *Drosophila melanogaster.* The samples were

loaded in the following sequence: Lane 1- *ey*-GAL4, Lane 2- L^2; *ey*-GAL4, Lane 3- L^2;

ey>C4. Alpha tubulin is used as a loading control. Molecular weight of Wg is 46kD, and

alpha tubulin is 55 kD. The samples were treated with anti-Wg antibody, and anti-α

tubulin antibody. (F) Bar graph representing as a relative Wg level, is a measure of

signal intensity of the bands, which clearly demonstrates that newt genes when

misexpressed under the L^2-mutant background downregulates Wg levels. Statistical

analysis was performed using student's t-test for independent samples. Statistical

significance was determined with 95% confidence ($p<0.05$). All bar graphs show

integrative density as a scale to measure Wg intensity for each sample represented as

the average between 5. Error bars show standard deviation, and symbols above the

error bar * signify p value <0.05, and ** signify p value <0.005 respectively. All the

images are displayed in same polarity as dorsal domain-towards top, and ventral

domain-towards bottom. Scale bar= 100 **μm**

1030

1031   **See also <u>Figure S7</u>, <u>S8</u>**

1032

1033   **Figure 5. Newt gene regulates ectopically misexpressed Wg both under the wild**

1034   **type background, and under the L^2-mutant background.**

1035

1036   (A-H) Wg is ectopically misexpressed in the *Drosophila* developing eye. (A,B) wild type

1037   background (*+/+; ey>wg-GFP*). (C, D) Newt gene and Wg are misexpressed under the

1038   wild type background (*+/+; ey>wg-GFP;C4*) showing increase in the eye size compared

1039   to (*+/+; ey>wg-GFP*). Comparatively represented as both (A, C) bright field adult eye

picture, and (B, D) third instar eye-antennal imaginal disc confocal image. (E, F) Wg is misexpressed under the L^2–mutant background (L^2/+; ey>wg-GFP) that lack newt gene. L^2–mutant small eye phenotype becomes more severe. (G, H) Newt gene and Wg are misexpressed under the L^2-mutant background (L^2/+; ey>wg-GFP; C4). There is a significant rescue in the eye-loss phenotype. Comparatively represented as both (E, G) bright field adult eye picture, and (F, H) third instar eye-antennal imaginal disc confocal image. The GFP as reporter is in green that signifies ectopic misexpression of Wg in the developing eye. Elav staining is shown in red. All the images are displayed in same polarity as dorsal domain-towards top, and ventral domain-towards the bottom. Scale bar= 100 μm

See also Figure S9

Figure 6. Modulating positive and negative regulators of Wg affect *L* mutant phenotype. (A-Q) Schematic presentation of Wg signaling pathway showing various members of the canonical pathway. (B-E) Activating Wg signaling in L^2 mutant eye background by misexpression of (B, C) *arm* alone (L^2/+; ey>arm), (D, E) *arm* and *C4* (L^2/+; ey>arm; C4). Note that *C4* misexpression with *arm* (L^2/+; ey>arm; C4) in the eye rescues the L^2/+; ey>arm reduced eye phenotype. (F-M) Blocking Wg signaling in L^2 mutant eye background by misexpression of (F, G) *sgg* alone (L^2/+; ey>sgg), (J, K) dTCF^{DN} alone (L^2/+; ey> dTCF^{DN}) results in rescue of L^2 loss-of-ventral eye phenotype. However, in L^2 mutant background misexpression of *C4* along with (H, I) *sgg* (L^2/+; ey>sgg; C4) and (L, M) dTCF^{DN} (L^2/+; ey> dTCF^{DN}; C4) significantly rescue the L^2/+; ey>sgg and L^2/+; ey> dTCF^{DN} partial loss-of-ventral eye phenotype.

(N-Q) Blocking transport of Wg morphogen in L^2 mutant eye background by misexpression of (N, O) porc^{RNAi} alone (L^2/+; ey> porc^{RNAi}), (D, E) porc^{RNAi} and C4 (L^2/+; ey> porc^{RNAi}; C4). Note that L^2/+; ey> porc^{RNAi} exhibits weak rescue of loss-of-ventral eye phenotype whereas *C4* misexpression with *sgg* (L^2/+; ey> porc^{RNAi}; C4) in the eye rescues the L^2/+; ey> porc^{RNAi} small eye phenotype. All the images are displayed in same polarity as dorsal domain-towards the top, and ventral domain-towards the bottom. Scale bar= 100 μm

1071

1072 **See also Figure S10, and S11**

1073

Figure 7. Model of newt genes function during development. These novel newt
genes (C1-C5) can negatively regulate evolutionarily conserved Wg /Wnt signaling
pathway in *Drosophila*, promote cell proliferation, and block cell death, which together
results in the rescue of L^2 mutant loss-of-ventral eye phenotype. This regenerative
ability of these novel newt genes extend from early eye development to late larval eye
and adult eye.

1080

1081

Mehta *et al.*, 2021, Figure 1

Mehta *et al.*, 2021, Figure 2

Mehta *et al.,* 2021, Figure 3

Mehta *et al.*, 2021, Figure 4

Mehta *et al.*, 2021, Figure 6