September 3, 2021

Collagen polarization promotes epithelial elongation by stimulating locoregional cell proliferation.

Hiroko Katsuno-Kambe[1], Jessica L. Teo[1], Robert J. Ju[1], James E. Hudson[2], Samantha J. Stehbens[1] and Alpha S. Yap[1,3]

[1]Division of Cell and Developmental Biology, Institute for Molecular Bioscience, The University of Queensland; and
[2]QIMR Berghofer Medical Research Institute, Brisbane, Australia

[3]Correspondence: a.yap@uq.edu.au

Keywords: Epithelia, elongation, tubulogenesis, collagen I, symmetry breaking, anisotropy

Abstract

Epithelial networks are commonly generated by processes where multicellular aggregates elongate and branch. Here we focus on understanding cellular mechanisms for elongation, using an organotypic culture system as a model of mammary epithelial anlage. Isotropic cell aggregates broke symmetry and slowly elongated when transplanted into collagen 1 gels. The elongating regions of aggregates displayed enhanced cell proliferation that was necessary for elongation to occur. Strikingly, this loco-regional increase in cell proliferation occurred where collagen 1 fibrils reorganized into bundles which were polarized with the elongating aggregates. Applying external stretch as a cell-independent way to reorganize the ECM, we found that collagen polarization stimulated regional cell proliferation to precipitate symmetry-breaking and elongation. This required β1-integrin and ERK signaling. We propose that collagen polarization supports epithelial anlagen elongation by stimulating loco-regional cell proliferation. This could provide a long-lasting structural memory of the initial axis that is generated when anlage break symmetry.

Introduction

Branched tubules represent one of the archetypal modes of epithelial organization (Iruela-Arispe and Beitel, 2013; Varner and Nelson, 2014). In organs such as the mammary gland and lungs, networks of hollow epithelial tubes mediate the physiological exchange of gases, nutrients and solutes between the body and its external environment (Chung and Andrew, 2008). These definitive networks are established by a complex morphogenetic process, where tubules grow outwards

from their precursors until they are instructed to branch, after which outgrowth
continues until the network is completed (Affolter et al., 2009; Andrew and Ewald,
2010). The branching and elongation of multicellular aggregates can thus be
considered as fundamental processes in the generation of tubular networks.

Interestingly, different organs also use distinct strategies for tubulogenesis.
For example, in the trachea, lumens appear early and accompany the growth of
tubules (Schottenfeld et al., 2010), whereas in the salivary and mammary glands
tubules first begin as non-polarized cellular aggregates (or anlage), which then
elongate as solid, multicellular cords before eventually forming lumens (Bastidas-
Ponce et al., 2017; Nerger and Nelson, 2018; Tucker, 2007). In the present study we
use the mammary epithelium as a model to analyse what guides the elongation of
multicellular anlage.

Tubulogenesis is a highly regulated phenomenon. The decision to branch is
recognized to be a critical checkpoint that is controlled by developmental signals and
cell-cell and cell-ECM interactions (Goodwin and Nelson, 2020). The elongation of
tubule precursors is also thought to be a regulated process controlled by receptor
tyrosine kinases and other signaling pathways (Costantini and Kopan, 2010;
Gjorevski and Nelson, 2010; Sternlicht et al., 2006). However, elongation occurs
over hours to days, time scales that are much longer than the underlying cellular
processes and the signaling pathways that guide them. This raises the question of
whether there may be mechanisms that can help guide elongation over longer time
scales, effectively serving as a bridge between the rapidity of cell signaling and the
slow progression of macroscopic anlage elongation. In this study, we show how
collagen 1 within the extracellular matrix (ECM) can provide such a bridge by
stimulating cell proliferation.

The ECM comprises complex mixtures of proteins, glycosaminoglycans and
glycoconjugates that fill the extracellular spaces of tissues and organs (Frantz et al.,
2010). One of the main components of ECM is collagen, particularly type 1 collagen
(also called collagen 1), which is often the dominant form during epithelial
tubulogenesis (Graham et al., 1988; Keely et al., 1995; Llacua et al., 2018; Nakanishi
et al., 1986; Simon-Assmann et al., 1995). Type 1 collagen exerts a diverse range of
effects that can potentially influence epithelial elongation. Fibrillar collagen helps
scaffold other molecules, such other ECM proteins and growth factors (Kanematsu
et al., 2004; Wipff and Hinz, 2008). Adhesion between cells and ECM allows the
chemical and mechanical properties of the ECM to regulate cell signaling and gene
regulation (Shi et al., 2011). These cell-ECM adhesions also allow cell-based forces
to reorganize the ECM. In particular, cells can rearrange collagen fibrils to influence
cell migration (Buchmann et al., 2021; Gjorevski et al., 2015; Guo et al., 2012; Shi et
al., 2014).

Importantly, fibrillar components of the ECM, such as collagen 1, are relatively
long-lived (Price and Spiro, 1977; Verzijl et al., 2000), making them attractive
candidates to bridge time scales during the elongation process. Indeed,
rearrangement of collagen has been implicated in patterning epithelial branching
(Brownfield et al., 2013; Guo *et al.*, 2012; Harunaga et al., 2011; Ingman et al., 2006;

Patel et al., 2006). However, it is difficult to elucidate the specific contribution for
collagen in the complex environment of an organ, where there are many additional
contributions from other cell types and various chemotactic signals. Therefore, in this
study we used three-dimensional (3D) organotypic cultures to test how the
extracellular matrix (ECM) regulates epithelial elongation. We report that a collagen
1 matrix induces the elongation of mammary epithelial anlage. This is accompanied
by the polarization of the matrix itself, an event that sustains anlagen elongation by
stimulating cell proliferation.
97
98   **Results**
99
100 **An experimental system to capture symmetry-breaking and elongation of**
101 **epithelial aggregates.**
102
To model the elongation of epithelial anlage, we induced multicellular aggregates to
break symmetry by manipulating the ECM environment in which the cells were
grown. When MCF10A cells are grown on Matrigel[TM] substrates and in media
supplemented with soluble Matrigel[TM] (Debnath et al., 2003), they proliferate from
single isolated cells to form a multicellular aggregate, which then polarizes and
clears the central mass of apoptotic cells, generating a central lumen in a process
known as cavitation (Figure 1-figure supplement 1A). In contrast, when isolated cells
were embedded in a gel of Type 1 collagen, which is the major ECM component in
the stromal environment during mammary tubulogenesis (Schedin and Keely, 2011),
they proliferated to form elongated, solid cords of non-polarized cells (Figure 1-figure
supplement 1B) (Krause et al., 2008). This indicated that some properties of the
collagen 1 environment might provide an instructive cue for elongation.
In order to capture the process by which isotropic aggregates broke symmetry
and then elongated, cells were first seeded in Matrigel[TM] until proliferation arrested at
10 days (Figure 1A); aggregates were then isolated and embedded into collagen 1
(acid solubilized Rat Type I collagen, 37□ for 30 minutes). Lumens were apparent in
some Matrigel-embedded aggregates, but most remained as solid spheres,
resembling anlage. Live-cell imaging revealed that the transplanted aggregates
displayed small jiggling motions for several hours, then spontaneously broke
symmetry and elongated to form cord-like structures similar to those formed when
cultured in collagen from the outset (Figure 1B and Video 1). These elongating
aggregates were generally solid, with no evident lumens (Figure 2A). Commonly,
elongation began with a group of cells that protruded away from the more spherical
original aggregate (Figure 1C,E). In contrast, when aggregates were transplanted
from Matrigel[TM] back into Matrigel[TM], they grew slightly but did not elongate (Figure
1-figure supplement 1C). Therefore, transplantation into a collagen I matrix
effectively caused isotropic MCF10A aggregates to break symmetry and elongate.
130
131 **Cell proliferation drives anlagen elongation.**
132

Multiple cellular processes have been implicated in epithelial elongation (Andrew and
Ewald, 2010; Economou et al., 2013; Keller, 2002). We first examined cell
proliferation, given the evident increase in the size of aggregates that elongated.
Staining for Ki-67, a nonhistone nuclear protein commonly used as a marker of
proliferating cells (Soliman and Yussif, 2016), showed clearly evident proliferation in
aggregates that had elongated 8 d after transplantation into collagen (Figure 1C,D).
These elongating aggregates commonly consisted of elongated extensions as well
as a rounded region that marked the original aggregate (Figure 1B). Strikingly, Ki-67-
positive cells were more frequent in the elongating parts of the aggregates (41.00 ±
4.55 % of all cells) compared with the non-elongating parts of the aggregates (18.86
± 3.05 %) (Figure 1C,D). Live-cell imaging of cells expressing NLS-mCherry also
showed that the proportion of nuclei which divided was higher in the elongating than
the non-elongating areas within aggregates (Figure 1E,F). Thus, elongation was
associated with loco-regional differences in cell proliferation, which was enhanced in
those parts of the aggregates that elongated.
More detailed inspection revealed that increased proliferation was first evident
in aggregates two days after transplantation (Figure 1G) and appeared to precede
the onset of symmetry-breaking (Figure 1H). To quantitate aggregate elongation we
calculated a symmetry ratio, i.e., the ratio of the maximum length (L_1) and width (L_2)
of the aggregates. Completely round aggregates will have a symmetry ratio of 1,
whereas for elongated aggregates L_1/L_2 will be >1 (Figure 2B). The symmetry ratio
first increased at Day 3 (Figure 1H), approximately 24 h after an increase in
proliferation was detected (Figure 1G). To test if any loco-regional differences in
proliferation were to be found at these early stages of symmetry breaking, we
analysed aggregates that had broken symmetry (defined experimentally as a
symmetry ratio > 1.5) after 3 days of culture. We compared the proportion of cells
that were Ki67-positive in the elongating regions of aggregates with the proportion to
be found in the non-elongating areas of the same aggregates (Figure 1I). Ki67
positivity was two-fold greater in the elongating areas (Figure 1I), suggesting that
loco-regional differences in proliferation were established early in the elongation
process.
Importantly, elongation was inhibited when proliferation was blocked using the
cell cycle inhibitors mitomycin C and aphidicolin (Figure 1J). In control cultures the
proportion of Ki-67-positive cells increased progressively after transplantation into
collagen (from 2.28 ± 0.85 % at Day 1 to 26.63 ± 3.34 % at Day 8). However, cell
proliferation was significantly reduced by mitomycin C (3.88 ± 1.18 %, Day 8) and
aphidicolin (11.54 ± 3.55 %, Day 8) (Figure 1K). Aggregate elongation was also
significantly reduced by both mitotic inhibitors (Figure 1J). Whereas the symmetry
ratio increased as control anlage elongated, this was blocked by both mitomycin C
and aphidicolin (Figure 1L). Formally, cell proliferation might have been necessary to
initiate symmetry-breaking and cell elongation or ongoing proliferation might also
have served to sustain elongation. To pursue this, we added mitomycin C or
aphidicolin after the first signs of symmetry breaking were evident (at 3 days after
transplantation). This reduced elongation by ~ 3-fold (Figure 1M). Therefore, cell

proliferation was required for elongation, with the capacity to contribute early in the
symmetry-breaking process and also later to sustain elongation.
We then used live cell imaging of labelled nuclei to evaluate other processes
implicated in epithelial elongation. Cells within the elongating regions tended to
divide along the axis of elongation (Figure 1-figure supplement 2A), as revealed by
comparing the axis of cell division and the principal axis of the region. The angle
difference between these axes was smaller in elongating areas compared with non-
elongating areas within the same aggregate (Figure 1-figure supplement 2A). This
suggested that polarized cell division accompanied enhanced proliferation during
aggregate elongation (Gong et al., 2004; Keller, 2006).
Tracking also revealed that cells were motile within elongating aggregates
(Figure 1-figure supplement 2B). This was evident in rounded aggregates that had
not broken symmetry and as well as in aggregates that had elongated (Figure 1-
figure supplement 2C). When we examined regional differences within aggregates
that had undergone elongation, we found that the speeds of migration were identical
in the parts that were elongating, compared with the non-elongating regions of the
aggregates (Figure 1-figure supplement 2D). However, the straightness of the tracks,
used an index of the persistence of migration, was slightly greater in the elongating
areas than in the non-elongating areas (Figure 1-figure supplement 2E). Furthermore,
cells within regions of elongation appeared to orient better with the axis of the
aggregate than did cells found in non-elongating areas. We measured this by
comparing the orientation of the tracks with the principal axis of the aggregates
(track displacement angle, Figure 1-figure supplement 2F). The track displacement
angle was less in the elongating areas than in the non-elongating areas.
Inhibition of proliferation with either mitomycin C or aphidicolin did not affect
cell migration speeds, in either the rounded or elongated areas (Figure 1-figure
supplement 2C). However, blocking proliferation slightly reduced track straightness,
implying a reduction in persistence (Figure 1-figure supplement 2G) and increased
the track displacement angle (Figure 1-figure supplement 2H). Therefore, the
apparently orderly migration of cells was compromised by blocking cell proliferation.
Together, these results suggest that proliferation was essential for the elongation
process and may also have influenced aspects of directional migration that would be
predicted to reinforce the elongation process.
210
211 **Collagen I condenses around epithelial aggregates that break symmetry.**
212
To gain further insight into the process of elongation, we sought to identify changes
in the ECM during this process. Consistent with the change in matrix environment
associated with the transplantation process, re-embedded aggregates lost laminin V
expression over time (Figure 2-figure supplement 1A). They expressed fibronectin
throughout the experiments without any evident regional differences in the
elongating areas of aggregates (Figure 2-figure supplement 1B).
Then we visualized Collagen 1 by labelling with the collagen-binding peptide
CNA35 (mCherry-CNA35; (Krahn et al., 2006)), mixed with soluble Collagen I before

the incorporation of MCF10A aggregates (Figure 2A). Collagen I, in contrast to the
basement membrane-like Matrigel^{TM}, is composed of fibrillar polymers that can guide
cell movement and whose organization is influenced by the application of cellular
forces (Brownfield *et al.*, 2013; Gjorevski *et al.*, 2015; Piotrowski-Daspit et al., 2017).
Aggregate elongation appeared to coincide with change in fibril organization.
Whereas fibrils appeared isotropic in acellular gels, collagen condensed around the
aggregates that had begun to elongate, forming dense bands that extended away
from the cells into the gel. However, it was seldom possible to visualize individual
fibrils with the low magnification, long working distance lenses that were required to
visualize aggregates within the gels.

Therefore, we measured the coherency (or co-orientation) of gels with
Orientation J, which calculates the local orientation and isotropy for each pixel in an
image based on the structure tensor for that pixel (see Methods for
details;(Rezakhaniha et al., 2012)). We interpret coherency as reflecting collagen
bundling and condensation (which we shall call "bundling" for short), such as has
been observed elsewhere during elongation (Brownfield *et al.*, 2013; Buchmann *et
al.*, 2021; Gjorevski *et al.*, 2015). Then, we compared collagen coherency with
aggregate shape, as measured by the symmetry ratio (Figure 2B), after 8 days
culture, when elongation was established. Overall, collagen coherency increased
with the increase in symmetry ratio (Figure 2C), implying that the degree of collagen
bundling increased with aggregate elongation. Furthermore, closer examination
around elongating aggregates showed that collagen coherency was greater in
regions proximate to the elongating parts of the aggregates compared with the non-
elongating parts (Figure 2D). In contrast, aggregates that remained spherical
showed no regional differences in collagen organization. Similarly, earlier studies
showed that collagen reorganized ahead of the tips of elongating aggregates
(Brownfield *et al.*, 2013; Gjorevski *et al.*, 2015). This suggested that collagen
became increasingly bundled and condensed where aggregates were elongating.

Increased collagen bundling was also evident at the early stages of elongation.
Because aggregates varied in the timing of when they broke symmetry, we
examined aggregates by live imaging in the first 3 days of the assays. We then
subdivided these based on their symmetry ratio, and compared the coherency of
collagen 1 around the elongating areas of aggregates with that around the non-
elongating areas (i.e. those that remained rounded, expressed as a fold-difference).
This showed that collagen 1 coherency increased around the elongating areas as
they broke symmetry (Figure 2E). Thus, from the early stages of symmetry breaking
collagen 1 bundling increased around the sites where aggregates elongated.

Collagen bundling required cell proliferation, as coherency was reduced by
treatment with either mitomycin C or aphidicolin (Figure 2F,G). Collagen bundling
was also reduced when we inhibited cellular contractility with the myosin antagonist,
blebbistatin, or the Rho kinase (ROCK) inhibitor, Y27632 (Figure 2H,I), consistent
with reports that cell contractility can condense collagen (Brownfield *et al.,* 2013;
Buchmann *et al.*, 2021; Gjorevski *et al.*, 2015). In contrast, collagen condensation
persisted around aggregates treated with the Rac1 GEF inhibitor NSC23766, which

did not affect aggregate elongation or the speed with which cells moved within the aggregates (Figure 2-figure supplement 2). This suggested that aggregates exerted forces to reorganize their local collagen environment as they broke symmetry.

Collagen polarity coaligns with cell aggregates during elongation.

We then asked if higher order organization in the collagen 1 gels was altered as aggregates elongated. For this, we used Orientation J to extract the principal axis of orientation in the gel, as a measure of its polarity, around elongating or non-elongating parts of an aggregate. Then we compared these gel orientations with the axis of elongation of the cell aggregate. First, we assembled histograms of gel orientation, setting the principal axis of elongation in the whole aggregate as 0^{o} (Figure 2J). The gels around non-elongating areas of the aggregates showed a broad distribution of orientations relative to the axis of the aggregate (Figure 2J). In contrast, the gel around the elongating regions of aggregates tended to orient with the principal axis of the aggregates. This suggested that the gel preferentially co-aligned with the aggregates around the regions of elongation. This notion was reinforced by comparing the angle differences between the principal axis (polarity) of the gel and of the aggregates. In this analysis, a decrease in the difference in angles between these two axes indicates an increase in their co-alignment. We found that the angle differences between the axes of collagen polarity and aggregate elongation was significantly smaller around the elongating regions than around the non-elongating regions of the aggregates (Figure 2K). Thus, the collagen matrix became polarized and co-aligned with the aggregates around regions of elongation.

Collagen polarization stimulates cell proliferation to induce aggregate elongation.

This led us to wonder if polarization of the collagen could itself influence the process of aggregate elongation. To test this, we developed a strategy that applied exogenous stretch to polarize the matrix independently of cell-generated forces. In this protocol, a collagen gel was created within a ring-shaped polydimethylsiloxane (PDMS) frame, then stretched uniaxially by expanding a stretcher inserted into the hole of the ring for 4 h (Figure 3-figure supplement 1A,B). Second harmonic imaging confirmed that collagen fibers were isotropically distributed in the unstretched gels and became more coherent (bundled) and more polarized immediately after stretching (Figure 3-figure supplement 1C,D), with their principal axes oriented in the direction of stretch (Figure 3-figure supplement 1E). Moreover, polarization of the collagen fibrils could be preserved for at least 7 days after the application of stretch by re-embedding the collagen rings in a larger collagen gel (Figure 3A-C, Figure 3-figure supplement 1B). In contrast, when gels were allowed to float in medium after stretching (Figure 3D, Figure 3-figure supplement 1B), gel coherency (Figure 3E) and polarity (Figure 3F) reverted to that of unstretched gels. Thus, reembedding

allowed us to sustain collagen polarization for a prolonged period after the initial
stretching.

We then transplanted spherical cell aggregates from Matrigel^{TM} into the
collagen rings and applied our stretching protocol. Interestingly, the aggregates did
not become elongated during the period when stretch was actively applied. The
symmetry ratios of aggregates immediately after stretching were identical to those in
unstretched aggregates (Figure 3-figure supplement 2A,B). Instead, aggregates
broke symmetry many hours after the period of active stretching. However,
symmetry breaking occurred earlier when stretched gels were allowed to retain
collagen polarization by re-embedding (35.14 ± 3.28 h, Figure 3G,H) compared with
aggregates that were allowed to break symmetry spontaneously in non-stretched
gels (68.40 ± 5.88 h, p<0.001). Moreover, the degree of elongation was also
enhanced once it had begun. The symmetry ratios of aggregates embedded in
stretch-polarized gels were significantly greater than those seen with native gels
(Figure 3I). Therefore, polarization of the collagen gel could stimulate the cell
aggregates to break symmetry.

Stretch-polarized aggregates also elongated preferentially along the axis of
the exogenous stretch, whereas aggregates in non-stretched gels elongated in a
random direction (Figure 3G,J, Figure 3-figure supplement 2C,D; Video 2). The
angle difference between collagen fiber polarity and aggregate elongation was
significantly smaller in stretch-polarized gels than in unstretched control gels (Figure
3K). Of note, collagen gel reorganization was preserved in the stretch-polarized gels
(Figure 3L,M), even in the presence of cellular aggregates, and this was oriented in
the direction of the original stretch. This suggested that in this assay the elongating
aggregates were following the polarity of the gel itself. In contrast, similar intensity of
fibronectin staining was seen in stretch-polarized aggregates compared with controls,
suggesting that fibronectin deposition was not stimulated by the stretch stimulation
(Figure 3-figure supplement 2E).

Importantly, the impact of stretch on the aggregates was disrupted when
collagen polarization was reversed. The proportion of aggregates that elongated
after stretching was reduced when the gels were floated rather than being
reembedded (Figure 4A) and their length of elongation was reduced (Figure 4B). Of
note, cell-based forces could not overcome the external forces acting on the gel as
the reversal of collagen polarization occurred even when aggregates were
incorporated into the gel (Figure 4C). This implied that collagen polarization may
have been responsible for allowing external stretch to promote aggregate elongation.

To confirm that these results were due to a critical role of collagen polarization,
rather than as-yet-unknown impacts of stretch upon the cells, we used collagenase
to extract aggregates from gels immediately after their 4 h stretch, then transplanted
them into unstretched gels where the collagen fibers oriented randomly (Figure 4D).
We reasoned that if elongation were due to a cell-intrinsic mechanism that bore the
memory of the stretch, then this should be preserved even after cells were
transplanted into a naïve gel. As noted earlier, aggregate elongation occurred earlier
when stretched gels were reembedded. However, this effect was lost when cells

were removed from the stretched gel immediately after stretching and transplanted into a naïve isotropic gel (Figure 4E). Together, these results indicate that collagen polarization can direct aggregate elongation, to accelerate its initiation and orient the direction of symmetry-breaking.

Collagen matrix polarization stimulates cell proliferation for anlagen elongation.

Since cell proliferation was a major driver of spontaneous aggregate elongation, we then asked if it was altered when the collagen gel was polarized by stretch. Indeed, we found that stretch-polarization stimulated cell proliferation (Figure 5A). An increase in the proportion of Ki67-positive cells was evident at day 1 after stretch-polarization, whereas it did not increase till day 2 in native gels. Moreover, the proportion of Ki-67 cells was consistently greater in stretch-polarized gels during the early phase of culture (day 1-3) as well as at the end of our experiments (day 8). Cell proliferation mediated the enhanced elongation in the stretched gels, as both the length (Figure 5B) and symmetry ratio (Figure 5C) of stretch-stimulated aggregates were reduced by mitomycin C and aphidicolin.

However, proliferation was not increased if gels were allowed to float, rather than being re-embedded, after stretching. Floating the gels reduced the proportion of proliferating cells in the aggregates overall (Figure 5D,E) as well as those specifically within the area of elongation (Figure 5F), compared with gels that had been re-embedded to preserve collagen polarization. This implied that it was sustained polarization of the gel, rather than simply transfer into the collagen environment, that could stimulate cell proliferation to elongate aggregates.

Polarized collagen promotes cell proliferation via ERK pathway.

Extracellular signal-regulated kinase (ERK1/2) is one of the major pathways that stimulates cell proliferation in mammary epithelial cells (Moreno-Layseca and Streuli, 2014; Streuli and Akhtar, 2009; Walker and Assoian, 2005). To evaluate its possible role in aggregate elongation, we expressed the ERK/KTR-mClover biosensor, which translocates out of the nucleus when ERK is activated (de la Cova et al., 2017). We confirmed the action of the sensor, as it accumulated in the nuclei of monolayer cultures that were treated with the ERK inhibitor FR180204 (50 μM) (Figure 6-figure supplement 1A). Using this sensor in 3D cultures, we found that the proportion of ERK-activated cells was significantly greater in the elongating regions of aggregates that had spontaneously broken symmetry, than in the non-elongating areas (Figure 6A, B). This was especially apparent in cells at the surface of aggregates that were in contact with the collagen gel (Figure 6C). In contrast, although Hippo pathway signaling has been implicated in breaking the symmetry of epithelial organoids (Serra et al., 2019), we did not detect any changes in nuclear Yap1 staining that might indicate that this pathway was being activated in our experiments (Figure 6-

395 figure supplement 1B,C). This suggested that ERK 1/2 signaling might be a
396 candidate for collagen polarization to stimulate cell proliferation.
397    This notion was supported by finding that FR180204 (50 µM) reduced cell
398 proliferation in response to stretch-polarization (Figure 6D,E) and decreased the
399 proportion of aggregates that elongated (Figure 6F). As well, the length (Figure 6G)
400 and asymmetric morphology (Figure 6H) were reduced in those aggregates that did
401 elongate. Thus, ERK1/2 appeared to be critical for collagen polarization to stimulate
402 cell proliferation for aggregate elongation.
403
404 **Integrins are necessary for polarized collagen to stimulate elongation**.
405
406 Integrins are the major ECM receptors in the mammary gland and other epithelia,
407 and they control diverse aspects of cellular function, including cell proliferation
408 (Miranti and Brugge, 2002; Wozniak et al., 2003). We therefore asked if integrin
409 signaling might have stimulated cell proliferation and aggregate elongation in
410 response to collagen polarization (Figure 7A).
411    We focused on the β1-integrins which have been implicated in regulating cell
proliferation during mammary gland development (Li et al., 2005). This class of
integrins contains a number of collagen 1 receptors, including α2β1-integrin (Heino,
2000; Käpylä et al., 2000) which was expressed in our MCF10A cells (Figure 7B).
Then we used the inhibitory AIIB2 antibody to block β1-integrin during stretch-
polarization experiments. mAb AIIB2 (15 µg/ml) was added as aggregates were
transplanted into the collagen gel rings and replenished after 7 days. mAb AIIB2
blocked the induction of proliferation in stretch-polarized gels, the proportion of Ki67-
positive cells being significantly reduced in AIIB2-treated cultures compared with
controls (Figure 7C,D). This was accompanied by inhibition of elongation. After 3
days incubation, 49.19 ± 1.61 % aggregates started to elongate in controls, whereas
less than 20 % of aggregates elongated when β1-integrins were inhibited (AIIB2:
17.86 ± 4.78 %) (Figure 7E). Moreover, even when mAb AIIB2-treated aggregates
eventually elongated upon prolonged culture (comparing the proportion of elongated
cells at 7 days with that at 3 days), these aggregates remained shorter (Figure 7F)
and less asymmetric (Figure 7G) than in the absence of AIIB2. Finally, we asked if
the ERK response required β1-integrins. Indeed, we found that the ability of stretch-
polarization to stimulate ERK signaling at early (day 2) and later stages (day 7) was
reduced when β1-integrins were blocked with mAb AIIB2 (Figure 7H,I). This implied
that a β1-integrin-ERK pathway was responsible for stimulating epithelial
proliferation.
432
433 **Discussion**
434
435 These results lead us to conclude that a polarized collagen 1 matrix can promote
436 epithelial elongation by stimulating locoregional cell proliferation. We infer this
437 because: 1) Loco-regional cell proliferation was a striking feature of MCF10A
438 elongation in our experiments, whose loss was not compensated for by other

morphogenetic processes. 2) The pattern of proliferation within aggregates was critically influenced by the polarized organization of the surrounding collagen 1 matrix. Together, we suggest that these reveal an interplay between collagen polarization and cell proliferation that can guide epithelial elongation (Figure 8).

Cell proliferation was a distinguishing feature of the elongation process in our MCF10A model. Increased proliferation was observed in elongating aggregates and, indeed, began to increase before elongation was detected. Furthermore, proliferation was greater in the regions of the aggregates that underwent elongation compared to those areas that did not. This suggested that a loco-regional increase in cell proliferation might be important for the elongation process, a notion that was supported when elongation was inhibited by blocking proliferation. Elongation was reduced when mitomycin C and aphidicolin were added from the beginning of the assays, consistent with earlier evidence that inhibiting proliferation early in the culture process blocked branching morphogenesis in mouse mammary organoids (Ewald et al., 2008). We also found that elongation was compromised if proliferation was inhibited after the elongation process had begun, a later dependence that was not seen in organoids (Huebner et al., 2016). Differences in cell system may therefore influence the impact of cell proliferation on elongation. Nonetheless, our data strongly suggest that loco-regional stimulation of cell proliferation was required for epithelial elongation in our model.

This raised the question of how cell proliferation might be preferentially increased within specific regions of MCF10A aggregates. Several of our observations implicate polarized reorganization of the collagen 1 matrix in this specification process. First, loco-regional polarization of the collagen 1 matrix accompanied aggregate elongation when cell aggregates spontaneously broke symmetry. This reorganization was distinguished by increased coherency of the matrix, consistent with an increase in collagen bundling, as well as polarization of the gel so that it became oriented with the axis of the elongating aggregate. Matrix reorganization concentrated where the aggregates were elongating and proliferation, as has been reported earlier (Brownfield *et al.*, 2013; Buchmann *et al.*, 2021; Gjorevski *et al.*, 2015). Second, aggregate elongation was stimulated when the gel was polarized by applying an external stretch. Similarly, collagen 1 orientation has been reported to mammary organoid branching (Brownfield *et al.*, 2013). But the stimulatory effect on elongation was lost if the stretched gels were allowed to lose their polarization or if aggregates were transplanted into a naïve, unstretched gel. This implied that the cells were responding to the altered organization of the gel, rather than simply exposure to collagen 1. Consistent with what we had observed when aggregates spontaneously broke symmetry, stretch-induced polarization of gels stimulated cell proliferation and this was necessary for the accelerated elongation process to occur. This appeared to be mediated by a β1-integrin and ERK-dependent pathway.

What property of the polarized collagen 1 was being recognized by the cells to enhance their proliferation? The polarization of collagen networks can have complex effects on cells. Increasing fibril alignment in acellular collagen gels was reported to

increase fibre stiffness and decrease pore size (Riching et al., 2014; Taufalele et al.,
2019), even in the absence of externally applied forces. Moreover, in single cell
cultures aligned collagen fibres were associated with larger focal adhesions than
non-aligned fibrils (Doyle and Yamada, 2016). One possibility, then, is that integrin
signaling was responding to the increased stiffness of the polarized collagen gels.
This is consistent with the well-characterized role for integrins to sense changes in
matrix stiffness (Giannone and Sheetz, 2006). As well, collagen 1 can bind to, and
sequester, a variety of growth factors. Whether this reservoir can be released when
cells apply tension to collagen is an interesting question for further consideration
(Wipff and Hinz, 2008).

Cell migration is a key driver of branching morphogenesis (Gjorevski *et al.*,
2015; Huebner *et al.*, 2016) and earlier experiments reported that the application of
external stretch to collagen-embedded aggregates could promote elongation through
cell migration (Brownfield *et al.*, 2013). We, too, observed cell migration in our
aggregates and, although their speeds were not different, migrating cells within
elongating parts of the aggregates tended to align with the axis of elongation and
move slightly more persistently than cells moving within the non-elongating regions
of the aggregates. These features of collective migration might be expected to
contribute to aggregate elongation.

Interestingly, blocking cell proliferation decreased the apparent persistence of
the migrating cells and their alignment with the principal axes of the aggregates.
Therefore, rather than collective migration being enhanced to compensate for the
decrease in proliferation, cell proliferation appeared to support these elongation-
facilitating aspects of cell migration. One possible explanation is that persistence and
directional alignment were being guided by the reorganized collagen which, in effect,
served to create a 3-dimensional micropattern around the aggregates. Studies in 2-
dimensional systems have shown that cells can orient their patterns of migration
when collagen networks condense and form bundles (Mohammed et al., 2020; Wang
et al., 2018). As collagen reorganization was reduced by inhibiting proliferation,
mitomycin C and aphidicolin might have affected cell migration indirectly through
matrix organization. Such micropatterning might also influence other aspects of cell
behaviour during elongation. Micropatterning experiments using 2-dimensional
substrata have shown that anisotropic confinement can orient patterns of cell division
(Thery et al., 2005), potentially by orienting tensile stresses (Campinho et al., 2013;
Legoff et al., 2013), as would also be predicted to accompany stiffer, condensed
collagen bundles. Consistent with this, we observed that cells within the elongating
areas appeared to orient their divisions with the axis of elongation, an effect which
would also be predicted to enhance elongation. These observations reinforce the
notion that interplay between cell proliferation and collagen (re)organization is a key
contributor to epithelial elongation.

The reductionist model that we used in these experiments allowed us to focus
on testing how collagen organization could promote epithelial elongation. How might
the results from this system operate in the more complex environment of the tissue?
One important factor to consider is the basement membrane, which can separate the

epithelial cell compartment from collagen 1 in the stroma. However, earlier studies reported that the basement membrane thins substantially and is remodelled in regions of epithelial elongation, such as the tips of mammary or salivary gland buds (Harunaga et al., 2014; Silberstein and Daniel, 1982; Williams and Daniel, 1983), potentially allowing elongating regions of epithelia to engage with collagen 1. Evaluating how cell proliferation, stromal collagen organization are coordinated with other components of the ECM in physiological models of developing glands will be an important question for future research.

In conclusion, we propose the following working model (Figure 8): Regional polarization of the collagen 1 matrix begins in response to anisotropies in force that aggregates exert upon their ECM via their integrin adhesions (Figure 7A). Cell-based forces exerted through integrins can apply strain on collagen fibrils (Brownfield *et al.*, 2013; Buchmann *et al.*, 2021; Gjorevski *et al.*, 2015; Hall et al., 2016) and, consistent with this, collagen reorganization in our experiments was compromised by inhibiting cellular contractility. However, compression of collagen by increasing cell numbers or cell movements could also have contributed (Buchmann *et al.*, 2021). By implication, collagen polarization will be greater around the parts of the aggregate that are generating more force. The consequent loco-regional collagen polarization then stimulates further cell proliferation in the adjacent parts of the aggregate to sustain elongation. It was interesting to note that collagen reorganization was compromised when cell proliferation was blocked from the outset of the assays. Possibly, subtle differences in cell proliferation may have contributed to generating initial anisotropies in force to reorganize the gels. If so, we speculate that the increase in cell proliferation may have contributed to further collagen reorganization, with the capacity to develop a feedback system that might help sustain aggregate elongation. This would allow polarized collagen networks to effectively provide a structural memory of the initial axis of symmetry breaking, i.e., a relatively long-lived spatial cue that directs further elongation of the epithelial anlagen.

Acknowledgements

We thank Selwin Wu for advice and all our lab colleagues for their feedback and support. The authors were supported by the National Health and Medical Research Council of Australia (Fellowship 1136592 and GNT1123816, 1140090 to ASY), Australian Research Council (DP19010287, 190102230 to AY and FT190100516 to SS), a Snow Medical Fellowship to JH, and a postdoctoral fellowship from The Uehara Memorial Foundation to HK.

Materials and Methods

Cell culture and lentivirus infection
MCF10A human mammary epithelial cells were cultured in DMEM/F12 medium supplemented with 5% horse serum, 10 µg/ml insulin, 0.5 µg/ml hydrocortisone, 100

ng/ml cholera toxin, 20 ng/ml EGF, 100 units/ml Penicillin and 100 units/ml
Streptomycin as previously described (Debnath et al., 2002).
HEK-293T cells were transfected using Lipofectamine 2000 (Invitrogen) for lentiviral
expression vector pLL5.0 and third generation packaging constructs pMDLg/pRRE,
RSV-Rev and pMD.G. Third generation packaging constructs were kindly provided
by Prof. James Bear (UNC Chapel Hill, North Carolina, USA). After transducing
MCF10A cells with either pLL5.0-EGFP-HRasC20, pLL5.0-NLS-mCherry or pLenti-
ERK/KTR-mClover lentivirus construct, we isolated highly-expressing cells using
Fluorescence-activated cell sorting (Influx Cell sorter; Cytopeia).
581
3D Matrigel and collagen culture
MCF10A aggregates and acini were cultured on 100% growth factor reduced
Matrigel (FAL354230; Corning) and overlayed with Matrigel including medium in 8
well chamber coverglass (0030742036; Eppendorf) as previously described
(Debnath *et al.*, 2002; Debnath *et al.*, 2003). In brief, single isolated MCF10A cells
were mixed with an assay medium containing 2% Matrigel, 5 ng/ml EGF and seeded
on a solidified layer of 100% Matrigel at 10000 cells/well. After seeding cells on
Matrigel, medium was changed every 4 days.
In collagen culture, single isolated MCF10A cells were mixed with collagen gel at
20000 cells/well (FAL354236; Corning). This mixture was neutralized with NaOH and
HEPES buffer and collagen concentration was adjusted to 1.0 mg/ml with culture
medium on ice. The pH of collagen solution was checked by litmus paper. Collagen
solution with cells was seeded in 8 well chamber coverglass and solidified at 37 □
for 30 min.
596
Transplantation of MCF10A aggregates
MCF10A aggregates cultured on Matrigel for 10 days were washed with PBS and
incubated with cold cell recovery solution (FAL354253; Corning) for 30 min on ice.
Aggregates were collected into a tube containing the cell recovery solution, spun
down at 1200 rpm, and washed with cold PBS. PBS-washed aggregates were
resuspended in MCF10A culture medium and mixed with collagen solution
neutralized with NaOH and HEPES buffer. Collagen gel solution with aggregates
were solidified at 37 □ for 30 min.
Aggregates embedded in collagen gel were isolated from gel by dissolving the gel
with collagenase (C2799; Sigma). Gels were washed with PBS and incubated in 20
µg/ml collagenase in Hanks' Balanced Salt Solution (H8264; Sigma) at 37 □ for 30
min to dissolve the collagen gel. Aggregates were collected with incubated solution
into the tube and mixed with DMEM/F12 medium which contains 20% horse serum.
Solution was spun down at 1200 rpm and pellet was washed with PBS. PBS-washed
aggregates were resuspended in culture medium and re-embedded in collagen gel.
612
Plasmids
pLL5.0-NLS-mCherry was constructed in previous study (Leerberg et al., 2014).
pLent-ERK/KTR-mClover was obtained from addgene (#59150). pLL5.0-EGFP-
HRasC20 was generated by insertion of EGFP-HRasC20 fragment which was
amplified from pEGFP-F (#6074-1; Clontech) by PCR into pLL5.0 vector. pLL5.0-
EGFP (gift from Prof. James Bear; UNC Chapel Hill, North Carolina, USA) was
digested with EcoRI and SbfI to remove EGFP, and then EGFP-C20 fragment was
inserted and ligated by In-Fusion cloning kit (638910; Clontech).
621

Antibodies and inhibitors

Primary antibodies used for immunocytochemistry in this study were rabbit anti-Ki67 (ab15580; Abcam), mouse anti-laminin V (MAB19562; Chemicon), rat anti-E-cadherin (131900; Invitrogen), mouse anti-GM130 (610822; BD), rabbit anti-fibronectin (F3648; Sigma) and rabbit anti-YAP1 (4912; Cell Signaling Technology). F-actin was stained with AlexaFluor 488-, 594-, 647-phalloidin (Invitrogen). Primary antibodies used for immunoblot were rabbit anti-integrin α2 (ab181548; Abcam), mouse anti-integrin β1 (610467; BD Transduction Laboratories) and rabbit anti-p44/42 MAPK (Erk1/2) (9102; Cell Signaling Technology). AIIB2 antibody for blocking integrin β 1 was purchased from Developmental Studies Hybridoma Bank and treated at 15 µg/ml in this study.

Proliferation inhibitors mitomycin C (M7949; Sigma) and aphidicolin (178273; Merck) were used at 10 µM and 2 µM respectively. Myosin II inhibitor blebbistatin (203390; Sigma) and ROCK inhibitor Y-27632 (688000; Sigma) were used at 25 µM and 30 µM respectively. ERK1/2 inhibitor FR180204 (SC-203945; Santa Cruz) was used at 50 µM. Except for mitomycin C, cell aggregates were treated with the inhibitors after transplantation into collagen gels. Inhibitors were then replenished when medium was changed and left in the collagen culture until cell aggregates were ready to be fixed. For mitomycin C wash out experiments, cell aggregates were treated with mitomycin C for 1 hour, then washed 3 times with PBS and left to recover in MCF10A culture medium.

Immunocytochemistry and microscopy

3D cultured cells in gel were fixed with 4% paraformaldehyde in cytoskeleton stabilization buffer (10mM PIPES at pH 6.8, 100mM KCl, 300 mM sucrose, 2 mM EGTA and 2 mM $MgCl_2$) at room temperature for 30 min, followed by a treatment for 30 min with 0.5% TritonX-100 and 10% goat serum in PBS for 1 hour at room temperature. Then they were stained with primary antibodies for overnight at 4 ☐. Subsequently, cells were washed with PBS and incubated with secondary antibodies with phalloidin and DAPI for 1 hour at room temperature.

Confocal images were acquired with an upright Meta laser scanning confocal microscope (LSM710; Zeiss) equipped with plan-Apochromat 20x 0.8NA or 40x 1.3NA objectives (Zeiss) and zen2012 software (Zeiss). The fluorescent images of collagen fibrils probed with mCherry-CNA35 were acquired by inverted microscope (Ti2; Nikon) equipped with Dragonfly spinning disc (Andor), by using plan-Apo 10x 0.45NA, 20x 0.75NA or 40x 0.95NA dry objectives (Nikon) and Fusion software (Andor).

Fluorescent Images of the SHG signal from collagen-1 were collected with an inverted confocal microscope (LSM710; Zeiss) equipped with multiphoton laser by using 860 nm excitation with SHG signal obtained with 690 nm bandpass filter. A 40x 1.3NA or 63x 1.40NA plan-apochromat oil objectives (Zeiss) were used to obtain SHG signals.

Fluorescence and phase contrast live images of elongating aggregates were acquired with an inverted fluorescence microscope (IX81; Olympus) equipped with CCD-camera (Hamamatsu) and an incubation box (Clear State Solutions) maintained at 37☐ and 5% CO_2 with gas controller (OkoLab), using plan-Apo 10x 0.4NA objective (Olympus) and CellSens software (Olympus).

Collagen gel labeling and stretching

Collagen fibrils were labelled with mCherry-CNA35 by mixing with purified mCherry-
CNA35 protein at 2 µM before gelling. Gels were solidified at 37 □ for 30 min.
PDMS gel frame and stretchers for gel stretching were kindly gifted from Dr. James
Hudson (QIMR Berghofer, Queensland, Australia). 1.5 mg/ml collagen gels were
solidified as a ring-shape in PDMS frame. PDMS stretchers were inserted into the
hole of the gel and expanded, and incubated for 4 hours in culture medium.
Stretched gels were released and then floated in medium or re-embedded in
collagen gel. For re-embedding, stretched gels were put in 1.5 mg/ml collagen gel
and solidified at at 37 □ for 30 min.
680
CNA35-mcherry protein purification
Protein expression vector pET28a-mCherry-CNA35 was obtained from Addgene
(#61607). Transformed E.coli (BL21) was cultured in 400 ml LB, and induced protein
expression with isopropyl β-D-1-tthiogalactopyranoside for 20 hours at 25□. Cultured
bacteria were spun down, and collected bacteria pellet was sonicated. The lysate
was centrifuged and the resulting supernatant was run through a column filled with
N-NTA His-band resin (Millipore). Bound protein was eluted and then dialyzed for
overnight in PBS at 4□. Endotoxin was removed by endotoxin removal columns
(88274; Thermo) following with manufacturer's protocol.
690
Quantitative analysis of collagen fibril alignment and aggregates elongation
Fluorescent confocal images of collagen fibrils acquired SHG microcopy or CNA35
probes were used for collagen fibril analysis. We analysed collagen organization in
these images using Orientation J which calculates the local orientation and isotropy
for each pixel in an image based on the structure tensor for that pixel (Rezakhaniha
et al., 2012). The structure tensor is evaluated for each pixel of the given image by
calculating the spatial partial derivatives by using (a cubic B-spline) interpolation.
The local orientation and isotropy for each pixel are computed based on the
eigenvalues and eigenvectors of the structure tensor. We characterized three
features in the organization of collagen fibrils orientations (Clemons et al., 2018): i)
The coherency of fibrils, defined as co-orientation in the same direction, as a
measure of bundling. ii) Isotropy, the distribution of fibril orientation in the field of
analysis; and iii) Polarization, defined as the principal axis of fibril orientation in
anisotropic gels, relative to a reference axis. 50 X 50 um or 100 X 100 um size of
ROI were selected from SHG images or CNA35 probed images respectively and
used for analysis. Elongation axis of aggregates were measured by the angle tool in
ImageJ, and then we calculated the angle difference between principal axis of
collagen fibrils or stretching axis. The fold difference of gel coherency was measured
by dividing the coherency of fibrils in the elongating area by the non-elongating area.
The symmetry ratio of aggregates was measured by dividing the longest length by
widest width of aggregates.
712
Quantitative analysis of proliferating cells and nuclear division angle
The number of cells co-stained with DAPI and anti-Ki67 antibody were counted by
Imaris software (Bitplane). The number of Ki67 positive cells were divided by the
total number of nucleus stained with DAPI to calculate percentage of Ki67 positive
cells. For live imaging, cells expressed with NLS-mCherry to count their number. The
frequency and orientation of cell division were analyzed from the time lapse images
by ImageJ. We set the elongation axis of aggregate as a reference and measured
the angle difference between elongation axis and dividing axis of nucleus.

Analysis of nuclear tracking

To obtain the migration speed of cells, we tracked individual cell nuclei using the Spot function in the Imaris software (Huebner *et al.*, 2016). The fluorescent images of cells expressing NLS-mCherry were acquired every 10 minutes for at least 50 hours. To obtain the track displacement angle, we first calculated the displacement angle of nuclei from the displacement of X and Y axis between first and last position, and then measured the angle difference from elongating angle of aggregates. Track straightness was calculated from track displacement by track length. Inhibitors were treated 1 hour before the imaging.

Quantitative analysis of ERK/KTR biosensor

ERK activity was judged by the location of mClover fluorescent signal in individual cells. Briefly, fluorescent tagged KTRs translocate between nuclei and cytoplasm depends on kinase activity. When ERK activity is high, KTR-mClover should localize in cytoplasm (de la Cova *et al.*, 2017). Aggregates expressing ERK/KTR-mClover were transplanted into collagen gel, and were co-stained with phalloidin and DAPI after fixation. To judge the delocalization of KTR-mClover, we used line intensity scan in single plane images and then manually counted ERK active cells.

Statistical analysis

Significance was determined by unpaired Student's t test and one-way ANOVA by using GraphPad Prism 8 (GraphPad software).

References

Affolter, M., Zeller, R., and Caussinus, E. (2009). Tissue remodelling through branching morphogenesis. Nat Rev Mol Cell Biol *10*, 831-842. 10.1038/nrm2797.

Andrew, D.J., and Ewald, A.J. (2010). Morphogenesis of epithelial tubes: Insights into tube formation, elongation, and elaboration. Dev Biol *341*, 34-55. 10.1016/j.ydbio.2009.09.024.

Bastidas-Ponce, A., Scheibner, K., Lickert, H., and Bakhti, M. (2017). Cellular and molecular mechanisms coordinating pancreas development. Development *144*, 2873-2888. 10.1242/dev.140756.

Brownfield, D.G., Venugopalan, G., Lo, A., Mori, H., Tanner, K., Fletcher, D.A., and Bissell, M.J. (2013). Patterned collagen fibers orient branching mammary epithelium through distinct signaling modules. Curr Biol *23*, 703-709. 10.1016/j.cub.2013.03.032.

Buchmann, B., Engelbrecht, L.K., Fernandez, P., Hutterer, F.P., Raich, M.K., Scheel, C.H., and Bausch, A.R. (2021). Mechanical plasticity of collagen directs branch elongation in human mammary gland organoids. Nat Commun *12*, 2759. 10.1038/s41467-021-22988-2.

Campinho, P., Behrndt, M., Ranft, J., Risler, T., Minc, N., and Heisenberg, C.P. (2013). Tension-oriented cell divisions limit anisotropic tissue tension in epithelial spreading during zebrafish epiboly. Nat Cell Biol *15*, 1405-1414. 10.1038/ncb2869.

Chung, S., and Andrew, D.J. (2008). The formation of epithelial tubes. Journal of Cell Science *121*, 3501-3504. 10.1242/jcs.037887.

Clemons, T.D., Bradshaw, M., Toshniwal, P., Chaudhari, N., Stevenson, A.W., Lynch, J., Fear, M.W., Wood, F.M., and Iyer, K.S. (2018). Coherency image analysis to quantify collagen architecture: implications in scar assessment. Rsc Adv *8*, 9661-9669. 10.1039/c7ra12693j.

Costantini, F., and Kopan, R. (2010). Patterning a complex organ: branching morphogenesis
and nephron segmentation in kidney development. Dev Cell *18*, 698-712.
10.1016/j.devcel.2010.04.008.
de la Cova, C., Townley, R., Regot, S., and Greenwald, I. (2017). A Real-Time Biosensor for
ERK Activity Reveals Signaling Dynamics during C. elegans Cell Fate Specification. Dev Cell *42*,
542-553 e544. 10.1016/j.devcel.2017.07.014.
Debnath, J., Mills, K.R., Collins, N.L., Reginato, M.J., Muthuswamy, S.K., and Brugge, J.S.
(2002). The role of apoptosis in creating and maintaining luminal space within normal and
oncogene-expressing mammary acini. Cell *111*, 29-40.
Debnath, J., Muthuswamy, S.K., and Brugge, J.S. (2003). Morphogenesis and oncogenesis of
MCF-10A mammary epithelial acini grown in three-dimensional basement membrane
cultures. Methods *30*, 256-268.
Doyle, A.D., and Yamada, K.M. (2016). Mechanosensing via cell-matrix adhesions in 3D
microenvironments. Exp Cell Res *343*, 60-66. 10.1016/j.yexcr.2015.10.033.
Economou, A.D., Brock, L.J., Cobourne, M.T., and Green, J.B. (2013). Whole population cell
analysis of a landmark-rich mammalian epithelium reveals multiple elongation mechanisms.
Development *140*, 4740-4750. 10.1242/dev.096545.
Ewald, A.J., Brenot, A., Duong, M., Chan, B.S., and Werb, Z. (2008). Collective epithelial
migration and cell rearrangements drive mammary branching morphogenesis. Dev Cell *14*,
570-581. 10.1016/i.devcel.2008.03.003.
Frantz, C., Stewart, K.M., and Weaver, V.M. (2010). The extracellular matrix at a glance. J
Cell Sci *123*, 4195-4200. 10.1242/jcs.023820.
Giannone, G., and Sheetz, M.P. (2006). Substrate rigidity and force define form through
tyrosine phosphatase and kinase pathways. Trends in Cell Biology *16*, 213-223.
https://doi.org/10.1016/j.tcb.2006.02.005.
Gjorevski, N., and Nelson, C.M. (2010). Branch formation during organ development. Wiley
Interdiscip Rev Syst Biol Med *2*, 734-741. 10.1002/wsbm.96.
Gjorevski, N., Piotrowski, A.S., Varner, V.D., and Nelson, C.M. (2015). Dynamic tensile forces
drive collective cell migration through three-dimensional extracellular matrices. Sci Rep *5*,
11458. 10.1038/srep11458.
Gong, Y., Mo, C., and Fraser, S.E. (2004). Planar cell polarity signalling controls cell division
orientation during zebrafish gastrulation. Nature *430*, 689-693. 10.1038/nature02796.
Goodwin, K., and Nelson, C.M. (2020). Branching morphogenesis. Development *147*.
10.1242/dev.184499.
Graham, M.F., Diegelmann, R.F., Elson, C.O., Lindblad, W.J., Gotschalk, N., Gay, S., and Gay,
R. (1988). Collagen content and types in the intestinal strictures of Crohn's disease.
Gastroenterology *94*, 257-265. 10.1016/0016-5085(88)90411-8.
Guo, C.L., Ouyang, M., Yu, J.Y., Maslov, J., Price, A., and Shen, C.Y. (2012). Long-range
mechanical force enables self-assembly of epithelial tubular patterns. Proc Natl Acad Sci U S
A *109*, 5576-5582. 10.1073/pnas.1114781109.
Hall, M.S., Alisafaei, F., Ban, E., Feng, X., Hui, C.-Y., Shenoy, V.B., and Wu, M. (2016). Fibrous
nonlinear elasticity enables positive mechanical feedback between cells and ECMs.
Proceedings of the National Academy of Sciences *113*, 14043-14048.
10.1073/pnas.1613058113.
Harunaga, J., Hsu, J.C., and Yamada, K.M. (2011). Dynamics of salivary gland morphogenesis.
J Dent Res *90*, 1070-1077. 10.1177/0022034511405330.

Harunaga, J.S., Doyle, A.D., and Yamada, K.M. (2014). Local and global dynamics of the
basement membrane during branching morphogenesis require protease activity and
actomyosin contractility. Dev Biol *394*, 197-205. 10.1016/j.ydbio.2014.08.014.
Heino, J. (2000). The collagen receptor integrins have distinct ligand recognition and
signaling functions. Matrix Biology *19*, 319-323. https://doi.org/10.1016/S0945-
053X(00)00076-7.
Huebner, R.J., Neumann, N.M., and Ewald, A.J. (2016). Mammary epithelial tubes elongate
through MAPK-dependent coordination of cell migration. Development *143*, 983-993.
10.1242/dev.127944.
Ingman, W.V., Wyckoff, J., Gouon-Evans, V., Condeelis, J., and Pollard, J.W. (2006).
Macrophages promote collagen fibrillogenesis around terminal end buds of the developing
mammary gland. Dev Dyn *235*, 3222-3229. 10.1002/dvdy.20972.
Iruela-Arispe, M.L., and Beitel, G.J. (2013). Tubulogenesis. Development *140*, 2851-2855.
10.1242/dev.070680.
Kanematsu, A., Marui, A., Yamamoto, S., Ozeki, M., Hirano, Y., Yamamoto, M., Ogawa, O.,
Komeda, M., and Tabata, Y. (2004). Type I collagen can function as a reservoir of basic
fibroblast growth factor. Journal of Controlled Release *99*, 281-292.
https://doi.org/10.1016/j.jconrel.2004.07.008.
Käpylä, J., Ivaska, J., Riikonen, R., Nykvist, P., Pentikäinen, O., Johnson, M., and Heino, J.
(2000). Integrin alpha(2)I domain recognizes type I and type IV collagens by different
mechanisms. J Biol Chem *275*, 3348-3354. 10.1074/jbc.275.5.3348.
Keely, P.J., Wu, J.E., and Santoro, S.A. (1995). The spatial and temporal expression of the
alpha 2 beta 1 integrin and its ligands, collagen I, collagen IV, and laminin, suggest important
roles in mouse mammary morphogenesis. Differentiation *59*, 1-13. 10.1046/j.1432-
0436.1995.5910001.x.
Keller, R. (2002). Shaping the vertebrate body plan by polarized embryonic cell movements.
Science *298*, 1950-1954.
Keller, R. (2006). Mechanisms of elongation in embryogenesis. Development *133*, 2291-
2302. 10.1242/dev.02406.
Krahn, K.N., Bouten, C.V., van Tuijl, S., van Zandvoort, M.A., and Merkx, M. (2006).
Fluorescently labeled collagen binding proteins allow specific visualization of collagen in
tissues and live cell culture. Anal Biochem *350*, 177-185. 10.1016/j.ab.2006.01.013.
Krause, S., Maffini, M.V., Soto, A.M., and Sonnenschein, C. (2008). A novel 3D in vitro culture
model to study stromal-epithelial interactions in the mammary gland. Tissue Eng Part C
Methods *14*, 261-271. 10.1089/ten.tec.2008.0030.
Leerberg, J.M., Gomez, G.A., Verma, S., Moussa, E.J., Wu, S.K., Priya, R., Hoffman, B.D.,
Grashoff, C., Schwartz, M.A., and Yap, A.S. (2014). Tension-sensitive actin assembly supports
contractility at the epithelial zonula adherens. Curr Biol *24*, 1689-1699.
10.1016/j.cub.2014.06.028.
Legoff, L., Rouault, H., and Lecuit, T. (2013). A global pattern of mechanical stress polarizes
cell divisions and cell shape in the growing Drosophila wing disc. Development *140*, 4051-
4059. 10.1242/dev.090878.
Li, N., Zhang, Y., Naylor, M.J., Schatzmann, F., Maurer, F., Wintermantel, T., Schuetz, G.,
Mueller, U., Streuli, C.H., and Hynes, N.E. (2005). β1 integrins regulate mammary gland
proliferation and maintain the integrity of mammary alveoli. The EMBO Journal *24*, 1942-
1953. https://doi.org/10.1038/sj.emboj.7600674.

Llacua, L.A., Faas, M.M., and de Vos, P. (2018). Extracellular matrix molecules and their
potential contribution to the function of transplanted pancreatic islets. Diabetologia *61*,
1261-1272. 10.1007/s00125-017-4524-8.
Miranti, C.K., and Brugge, J.S. (2002). Sensing the environment: a historical perspective on
integrin signal transduction. Nature Cell Biology *4*, E83-E90. 10.1038/ncb0402-e83.
Mohammed, D., Pardon, G., Versaevel, M., Bruyère, C., Alaimo, L., Luciano, M., Vercruysse,
E., Pruitt, B.L., and Gabriele, S. (2020). Producing Collagen Micro-stripes with Aligned Fibers
for Cell Migration Assays. Cellular and Molecular Bioengineering *13*, 87-98. 10.1007/s12195-
019-00600-4.
Moreno-Layseca, P., and Streuli, C.H. (2014). Signalling pathways linking integrins with cell
cycle progression. Matrix Biology *34*, 144-153.
https://doi.org/10.1016/j.matbio.2013.10.011.
Nakanishi, Y., Sugiura, F., Kishi, J., and Hayakawa, T. (1986). Collagenase inhibitor stimulates
cleft formation during early morphogenesis of mouse salivary gland. Dev Biol *113*, 201-206.
10.1016/0012-1606(86)90122-3.
Nerger, B.A., and Nelson, C.M. (2018). 3D culture models for studying branching
morphogenesis in the mammary gland and mammalian lung. Biomaterials.
10.1016/j.biomaterials.2018.08.043.
Patel, V.N., Rebustini, I.T., and Hoffman, M.P. (2006). Salivary gland branching
morphogenesis. Differentiation *74*, 349-364. 10.1111/j.1432-0436.2006.00088.x.
Piotrowski-Daspit, A.S., Nerger, B.A., Wolf, A.E., Sundaresan, S., and Nelson, C.M. (2017).
Dynamics of Tissue-Induced Alignment of Fibrous Extracellular Matrix. Biophys J *113*, 702-
713. 10.1016/j.bpj.2017.06.046.
Price, R.G., and Spiro, R.G. (1977). Studies on the metabolism of the renal glomerular
basement membrane. Turnover measurements in the rat with the use of radiolabeled
amino acids. J Biol Chem *252*, 8597-8602.
Rezakhaniha, R., Agianniotis, A., Schrauwen, J.T., Griffa, A., Sage, D., Bouten, C.V., van de
Vosse, F.N., Unser, M., and Stergiopulos, N. (2012). Experimental investigation of collagen
waviness and orientation in the arterial adventitia using confocal laser scanning microscopy.
Biomech Model Mechanobiol *11*, 461-473. 10.1007/s10237-011-0325-z.
Riching, K.M., Cox, B.L., Salick, M.R., Pehlke, C., Riching, A.S., Ponik, S.M., Bass, B.R., Crone,
W.C., Jiang, Y., Weaver, A.M., et al. (2014). 3D collagen alignment limits protrusions to
enhance breast cancer cell persistence. Biophys J *107*, 2546-2558.
10.1016/j.bpj.2014.10.035.
Schedin, P., and Keely, P.J. (2011). Mammary gland ECM remodeling, stiffness, and
mechanosignaling in normal development and tumor progression. Cold Spring Harb
Perspect Biol *3*, a003228. 10.1101/cshperspect.a003228.
Schottenfeld, J., Song, Y., and Ghabrial, A.S. (2010). Tube continued: morphogenesis of the
Drosophila tracheal system. Curr Opin Cell Biol *22*, 633-639.
https://doi.org/10.1016/j.ceb.2010.07.016.
Serra, D., Mayr, U., Boni, A., Lukonin, I., Rempfler, M., Challet Meylan, L., Stadler, M.B.,
Strnad, P., Papasaikas, P., Vischi, D., et al. (2019). Self-organization and symmetry breaking
in intestinal organoid development. Nature *569*, 66-72. 10.1038/s41586-019-1146-y.
Shi, M., Zhu, J., Wang, R., Chen, X., Mi, L., Walz, T., and Springer, T.A. (2011). Latent TGF-
beta structure and activation. Nature *474*, 343-349. 10.1038/nature10152.

Shi, Q., Ghosh, R.P., Engelke, H., Rycroft, C.H., Cassereau, L., Sethian, J.A., Weaver, V.M., and
Liphardt, J.T. (2014). Rapid disorganization of mechanically interacting systems of mammary
acini. Proc Natl Acad Sci U S A *111*, 658-663. 10.1073/pnas.1311312110.
Silberstein, G.B., and Daniel, C.W. (1982). Glycosaminoglycans in the basal lamina and
extracellular matrix of the developing mouse mammary duct. Dev Biol *90*, 215-222.
10.1016/0012-1606(82)90228-7.
Simon-Assmann, P., Kedinger, M., De Arcangelis, A., Rousseau, V., and Simo, P. (1995).
Extracellular matrix components in intestinal development. Experientia *51*, 883-900.
10.1007/BF01921739.
Soliman, N.A., and Yussif, S.M. (2016). Ki-67 as a prognostic marker according to breast
cancer molecular subtype. Cancer Biol Med *13*, 496-504. 10.20892/j.issn.2095-
3941.2016.0066.
Sternlicht, M.D., Kouros-Mehr, H., Lu, P., and Werb, Z. (2006). Hormonal and local control of
mammary branching morphogenesis. Differentiation *74*, 365-381. 10.1111/j.1432-
0436.2006.00105.x.
Streuli, Charles H., and Akhtar, N. (2009). Signal co-operation between integrins and other
receptor systems. Biochemical Journal *418*, 491-506. 10.1042/bj20081948.
Taufalele, P.V., VanderBurgh, J.A., Munoz, A., Zanotelli, M.R., and Reinhart-King, C.A. (2019).
Fiber alignment drives changes in architectural and mechanical features in collagen matrices.
PLoS One *14*, e0216537. 10.1371/journal.pone.0216537.
Thery, M., Racine, V., Pepin, A., Piel, M., Chen, Y., Sibarita, J.B., and Bornens, M. (2005). The
extracellular matrix guides the orientation of the cell division axis. Nat Cell Biol *7*, 947-953.
Tucker, A.S. (2007). Salivary gland development. Semin Cell Dev Biol *18*, 237-244.
10.1016/j.semcdb.2007.01.006.
Varner, V.D., and Nelson, C.M. (2014). Cellular and physical mechanisms of branching
morphogenesis. Development *141*, 2750-2759. 10.1242/dev.104794.
Verzijl, N., DeGroot, J., Thorpe, S.R., Bank, R.A., Shaw, J.N., Lyons, T.J., Bijlsma, J.W., Lafeber,
F.P., Baynes, J.W., and TeKoppele, J.M. (2000). Effect of collagen turnover on the
accumulation of advanced glycation end products. J Biol Chem *275*, 39027-39031.
10.1074/jbc.M006700200.
Walker, J.L., and Assoian, R.K. (2005). Integrin-dependent signal transduction regulating
cyclin D1 expression and G1 phase cell cycle progression. Cancer and Metastasis Reviews *24*,
383-393. 10.1007/s10555-005-5130-7.
Wang, W.Y., Pearson, A.T., Kutys, M.L., Choi, C.K., Wozniak, M.A., Baker, B.M., and Chen, C.S.
(2018). Extracellular matrix alignment dictates the organization of focal adhesions and
directs uniaxial cell migration. APL Bioeng *2*, 046107. 10.1063/1.5052239.
Williams, J.M., and Daniel, C.W. (1983). Mammary ductal elongation: differentiation of
myoepithelium and basal lamina during branching morphogenesis. Dev Biol *97*, 274-290.
10.1016/0012-1606(83)90086-6.
Wipff, P.J., and Hinz, B. (2008). Integrins and the activation of latent transforming growth
factor beta1 - an intimate relationship. Eur J Cell Biol *87*, 601-615.
10.1016/j.ejcb.2008.01.012.
Wozniak , M.A., Desai , R., Solski , P.A., Der , C.J., and Keely , P.J. (2003). ROCK-generated
contractility regulates breast epithelial cell differentiation in response to the physical
properties of a three-dimensional collagen matrix. Journal of Cell Biology *163*, 583-595.
10.1083/jcb.200305010.
952

Figure legends

Figure 1. Type 1 collagen induces elongation of MCF10A anlage via cell proliferation.

(A) Cartoon: transplantation of MCF10A cell aggregates from Matrigel^{TM} into collagen gel. Single isolated cells were cultured on Matrigel^{TM} and overlayed with Matrigel^{TM}-containing medium. Matrigel^{TM} was washed out after 10 days and the aggregates re-embedded into type-1 collagen gel.

(B) Time lapse images of MCF10A aggregates after transfer into a collagen gel. Red arrowheads: elongation from aggregates.

(C) Fluorescence image of an elongated aggregate cultured for 8 days after transfer into collagen gel and stained with anti-Ki67 antibody (green), phalloidin (magenta) and DAPI (blue). Blue parenthesis: non-elongating area, red parenthesis: elongating area.

(D) Percentage of Ki67 positive cells in elongating and non-elongating areas of the elongated aggregates. (n = 15 aggregates).

(E) Time lapse images of elongating aggregates expressing NLS-mCherry. Yellow arrowheads: dividing nucleus.

(F) Frequency of cell division in elongating and non-elongating regions of aggregates. (n = 12 aggregates).

(G) Percentage of Ki67-positive cells in aggregates cultured for 0-3 days. (n = 102 aggregates).

(H) Symmetry ratio of aggregates cultured for 0-3 days. (n = 102 aggregates).

(I) Percentage of Ki67 positive cells in elongating area and non-elongating areas of aggregates which broke symmetry (defined as symmetry ratio >1.5) in the first 3 days of culture. (n = 18 aggregates).

(J) MCF10A aggregates co-stained with anti-Ki67 antibody (green) and DAPI (blue). Aggregates were cultured for 0 day (i) or 8 days (ii-iv) after treatment with vehicle (ii), mitomycin C (iii) or aphidicolin (iv).

(K) Percentage of Ki67 positive cells in aggregates cultured for 1 day or 8 days with or without mitomycin C or aphidicolin. (n = 63 aggregates).

(L) Symmetry ratio of aggregates cultured for 1 day or 8 days with or without mitomycin C or aphidicolin. (n = 237 aggregates).

(M) Effect of delayed inhibition of proliferation on aggregate elongation. Aggregates were cultured for 3 days before treatment with mitomycin C or aphidicolin. Data are fold change of elongation in control and drug-treated cultures (n = 57 aggregates).

All data are means ± SEM, *$P<0.05$, **$P<0.01$, ***$P<0.001$. Data in (D, F, I) were analyzed by unpaired Student's t-test. Data in (G, H, K-M) were analyzed with one-way ANOVA Tukey's multiple comparisons test.

The following figure supplements are available for figure 1:

Figure supplement 1. 3D morphology of MCF10A cells.

Figure supplement 2. Cell division and motility during aggregate elongation.

Figure 2. Mammary cell aggregates polarize the ECM as they elongate.

(A) Fluorescent image of collagen fiber alignment and elongated aggregates
expressing the cell membrane marker GFP-HRasC20. Collagen fibers were labeled
with mCherry-CNA35 peptide (magenta).
(B) Cartoon of symmetry ratio of aggregates (i) and gel coherency (ii).
(C) Scatter plot of aggregate symmetry ratio and collagen fiber coherency. (n = 75
aggregates).
(D) Regional analysis of collagen coherency around aggregates. Cartoon illustrates
the approach: for elongating aggregates coherency was measured in regions of
interest (ROIs) placed both at the tips of elongations and proximate to their non-
rounded areas. For rounded aggregates, ROIs were placed orthogonally. Regional
differences in coherency were measured as the fold difference, measured around
rounded aggregates (n = 19 aggregates) and elongated aggregates (n = 57
aggregates). Elongated aggregates were defined as symmetry ratio > 1.5. Double
headed arrow: elongating axis.
(E) Fold difference of collagen fiber coherency around aggregates measured at early
stages of elongation (first 3 days of culture) subdivided based on symmetry ratio
(n = 98 aggregates).
(F) Fluorescent images of collagen fibers labeled with mCherry-CNA35 (magenta)
and aggregates expressing GFP-HRasC20 cultured for 8 days with mitomycin C or
aphidicolin.
(G) Coherency of collagen fibers surrounding the aggregates treated with mitomycin
C and aphidicolin. (n = 40 aggregates).
(H) Fluorescent images of collagen fiber alignment with aggregates expressing GFP-
HRasC20 treated with blebbistatin or Y-27632 for 8 days. Collagen fibrils were
labeled with CNA35-mCherry (magenta).
(I) Symmetry ratio of aggregates cultured for 8 days with or without blebbistatin or Y-
27632. (n = 88 aggregates).
(J) Distribution of collagen fiber orientation surrounding elongated aggregates (n =
22 aggregates) or non-elongated aggregates (n = 13 aggregates).
(K) Difference between elongation axis of aggregates and average angle of collagen
fibers in non-elongating area or elongating area. (n = 57 aggregates).
All data are means ± SEM, ns, not significant, *P<0.05, **P<0.01, ***P<0.001. Data
in
(D, E, G, I) were analyzed with one-way ANOVA Tukey's multiple comparisons test.
Data in (K) were analyzed by unpaired Student's *t*-test.
The following figure supplements are available for figure 2:
Figure supplement 1. Immunofluorescent staining of ECM proteins in the
aggregates.
Figure supplement 2. Effect of Rac1 GEF inhibitor on MCF10A aggregate
elongation.
1038
1039 **Figure 3. Collagen polarization induces mammary aggregate elongation.**

(A) Second harmonic generation (SHG) images of collagen fibers in the gel with or without stretching and incubated for 7 days after re-embedding in gel. Double head arrow: stretching axis.

(B) Coherency of collagen fiber in the gel with or without stretching. (n = 24 positions in multiple gels).

(C) Distribution of collagen fiber orientation in the gel with or without stretching. 0^o is defined as the axis of stretch. (N = 3 independent experiments).

(D) SHG images of collagen fiber floated for 7 days with or without after stretching. Double head arrow: stretching axis.

(E) Coherency of collagen fiber in 7 days floated gel with or without stretching. (n = 46 positions in multiple gels).

(F) Distribution of collagen fiber orientation in gels that had been allowed to float (7 days) with or without prior stretching. (N = 3 independent experiments).

(G) Time lapse images of aggregates embedded in stretched gel. Double head arrow: stretching axis.

(H) Initiation time of aggregate elongation in the gel with or without stretching. (-Stretch: n = 25 aggregates, +Stretch: n = 71 aggregates).

(I) Symmetry ratio of aggregates in the early phase of culture (0-3 days) with or without stretching. (-Stretch, day 0: n = 12, day 1: n = 12, day 2: n = 24, day 3: n = 24, +Stretch, day 0: n = 21, day 1: n = 32, day 2: n = 17, day 3: n = 22).

(J) Distribution of elongation axes of aggregates in the gel with or without stretching. (- Stretch: n = 112 aggregates, +Stretch: n = 115 aggregates).

(K) Difference between elongating axis of aggregates and average angle of collagen fibers in the gel with or without stretching. (- Stretch: n = 21 aggregates, +Stretch: n = 102 aggregates).

(L) Fluorescence image of aggregates cultured for 5 days after gel stretching and co-stained with phalloidin (green) and DAPI (blue) in stretched gel. Collagen fibrils were labeled with CNA35-mCherry (magenta).

(M) Coherency of collagen fibers surrounding elongated aggregates in the gel with or without stretching. (- Stretch: n = 21 aggregates, +Stretch: n = 102 aggregates).

All data are means ± SEM; ns, not significant, **$P<0.01$, ***$P<0.001$. Data in (B, E, H, K, M) were analyzed by unpaired Student's *t*-test. Data in (I) was analyzed with one-way ANOVA Tukey's multiple comparisons test.

The following figure supplements are available for Figure 3:

Figure supplement 1. External gel stretching aligns collagen fiber.

Figure supplement 2. MCF10A aggregates elongate along the gel stretching axis.

Figure 4. Collagen polarization must be sustained to stimulate mammary aggregate elongation.

(A) Population of elongated aggregates in the re-embedded or floated gel. (N = 3 independent experiments).

(B) The length of elongated aggregates in the re-embedded or floated gel. (N = 3 independent experiments).

(C) Coherency of collagen fibers surrounding the aggregates in gels that were re-
embedded after stretch or floated for 7 days after stretching. (n = 68 aggregates).
(D) Cartoon of aggregate from stretched gel into normal collagen gel. Aggregates
were isolated from stretched gel by collagenase and re-embedded in naïve gel.
(E) Initiation time for aggregate elongation in non-stretched control gels, stretched
gels that had been re-embedded to preserve collagen polarization (embed) and after
cells were extracted and transferred into non-stretched gels (transfer) (n = 83
aggregates).
All data are means ± SEM; ns, not significant, **P<0.01, ***P<0.001. Data in (A-C)
were analyzed by unpaired Student's t-test. Data in (E) was analyzed with one-way
ANOVA Tukey's multiple comparisons test.
1094
Figure 5. Collagen polarization induces cell proliferation for aggregate
elongation.
(A) Time course of cell proliferation within aggregates in control gels (- stretch) or
after stretching (+ stretch). Data are percentage of cells that were Ki67 positive (n =
206 aggregates).
(B,C) Length (B) and (C) symmetry ratio of aggregates in stretched gel incubated
with mitomycin C or aphidicolin for 8 days. (n = 214 aggregates)
(D) Fluorescence images of elongated aggregates cultured for 7 days in the re-
embedded or floated gel after stretching. Aggregates were co-stained with anti-Ki67
antibody (green), Phalloidin (red) and DAPI (blue). Collagen fibers were labeled with
mCherry-CNA35 (magenta).
(E) Percentage of Ki67 positive cells in the aggregates cultured for 7 days in the re-
embedded or floated gel after stretching. (n = 56 aggregates).
(F) Percentage of Ki67 positive cells in the elongating area of aggregates in the re-
embedded or floated gel. (n = 47 aggregates).
All data are means ± SEM; ns, not significant, **P<0.01, ***P<0.001. Data in (E, F)
were analyzed by unpaired Student's t-test. Data in (A, B, C) were analyzed with
one-way ANOVA Tukey's multiple comparisons test.
1113
Figure 6. Polarized collagen promotes cell proliferation via the ERK pathway.
(A) Fluorescent images of aggregates expressing ERK/KTR-mClover biosensor
cultured for 7 days and stained with phalloidin (magenta) and DAPI (blue).
(B) Percentage of ERK active cells in elongating areas and non-elongating areas of
the aggregates. (n = 30 aggregates).
(C) Percentage of ERK active cells at the surface in elongating and non-elongating
areas of aggregates. (n = 30 aggregates).
(D) Fluorescent images of aggregates cultured for 7 days treated with FR180207
after gel stretching. Aggregates were co-stained with anti-Ki67 antibody (green),
Phalloidin (red) and DAPI (blue). Collagen fibers were labeled with mCherry-CNA35
(magenta).
(E) Percentage of Ki67-positive cells in aggregates incubated with FR180207 for
7days after stretching. (n = 24 aggregates).

(F) Effect of inhibiting ERK on stretch-induced aggregate elongation. Proportion of elongated aggregates in stretched gel incubated with FR180207 for 3 days and 7 days. (N = 3 independent experiments).

(G) Length and (H) symmetry ratio of elongated aggregates incubated with FR180207 for 7 days. (n = 166 aggregates).

All data are means ± SEM; ns, not significant, **P<0.01, ***P<0.001. Data in (B, C, E, -H) were analyzed by unpaired Student's *t*-test.

The following figure supplement is available for figure 6:

Figure supplement 1. ERK biosensor and YAP1 localization in MCF10A cells.

Figure 7. Integrins are necessary for polarized collagen to stimulate elongation.

(A) Schematic of potential integrin-ERK pathway that mediates the effect of collagen polarization on cell proliferation.

(B) Immunoblot of integrin α2, β1 and ERK1/2 protein levels in MCF10A cell lysate.

(C) Fluorescent images of aggregates cultured for 7 days treated with AIIB2 antibody after gel stretching. Aggregates were co-stained with anti-Ki67 antibody (green), Phalloidin (red) and DAPI (blue). Collagen fibers were labeled with mCherry-CNA35 (magenta).

(D) Percentage of Ki67 positive cells in the aggregates incubated with AIIB2 antibody for 7 days after stretching the gels. (n = 35 aggregates).

(E) Proportion of elongated aggregates in stretched gel incubated with AIIB2 antibody for 3 days and 7 days. (N = 3 independent experiments).

(F) Length and (G) symmetry ratio of elongated aggregates incubated with AIIB2 antibody for 7 days. (n = 134 aggregates).

(H) Fluorescent images of aggregates expressed with ERK/KTR-mClover and incubated with IgG (control) or AIIB2 antibody for 2 days after gel stretching. Aggregates were co-stained with phalloidin (red) and DAPI (blue). Collagen fibers were labeled with mCherry-CNA35 (magenta).

(I) Percentage of ERK active cells in aggregates incubated with AIIB2 antibody for 2 days or 7 days in stretched gel. (n = 78 aggregates).

All data are means ± SEM; ns, not significant, **P<0.01, ***P<0.001. Data in (D-G, I) were analyzed by unpaired Student's *t*-test.

Figure 8. Model of collagen polarization as a structural memory for epithelial anlage elongation.

(i) Initially isotropic epithelia anlage exert isotropic patterns of force on a non-polarized collagen 1 gel.

(ii) Initial anisotropies in force associated with symmetry-breaking of the aggregate exert strain on collagen fibrils leading to bundling and polarization.

(iii) The polarized collagen matrix provides a structural memory, that promotes regional cell proliferation to direct further elongation of the anlage.

Supplemental Figure legends

Figure 1-figure supplement 1. 3D morphology of MCF10A cells.

(A) MCF10A cells cultured in Matrigel^{TM} for 7 days (left) and 31 days (right). Day 7 aggregate was co-stained with anti-E-cadherin antibody (green), anti-GM130 antibody (red), phalloidin (magenta) and DAPI (blue). Day 31 acinus was co-stained with anti-Laminin V antibody (green), anti-E-cadherin antibody (red) and DAPI (blue).

(B) MCF10A cells cultured in Type 1 collagen gel for 3 days (left) and 10 days (right). Day 3 MFC10A cells expressing GFP-HRasC20 were co-stained with anti-GM130 antibody (red), phalloidin (magenta) and DAPI (blue). Day 10 MCF10A cells were co-stained with phalloidin (magenta) and DAPI (blue).

(C) MCF10A aggregate cultured for 7 days after transplantation from Matrigel^{TM} back into Matrigel^{TM}. Aggregate was co-stained with anti-Ki67 antibody (green), phalloidin (red) and DAPI (blue).

Figure 1-figure supplement 2. Cell division and motility during aggregate elongation.

(A) Average angle of cell division. Principal axis of aggregate is 0°. (n = 12 aggregates).

(B) Nuclear tracking during aggregate elongation. Tails show tracking path for each cell in the last 120 min of the movie.

(C) Speed of cell motility based on nuclear tracking in rounded and elongated aggregates treated with mitomycin C or aphidicolin. (n = 137 movies).

(D) Cell speed, (E) Track straightness and (F) track displacement angle in elongating aggregates. (n = 18 movies).

(G) Track straightness and (H) track displacement angle of cells in aggregates treated with mitomycin C or aphidicolin (n = 60 movies).

All data are means ± SEM; ns, not significant, *$P<0.05$, ***$P<0.001$. Data in (D-F) were analyzed by unpaired Student's *t*-test. Data in (C, G, H) were analyzed with one-way ANOVA Tukey's multiple comparisons test.

Figure 2-figure supplement 1. Immunofluorescent staining of ECM proteins in the aggregates.

(A) MCF10A aggregates transferred from Matrigel^{TM} into collagen gel at day 0 (left) and at day 4 (right). Day 0 aggregate was co-stained with anti-Laminin V antibody (green), anti-E-cadherin antibody (red), and DAPI (blue). Day 4 aggregate was co-stained with anti-Laminin V antibody (green), anti-E-cadherin antibody (red), phalloidin (magenta), and DAPI (blue). *: Y-Z slice image of yellow line.

(B) MCF10A aggregates transferred from Matrigel^{TM} into collagen gel at day 1 (left) and at day 7 (right). Aggregates were co-stained with anti-fibronectin antibody (green) and DAPI (blue).

Figure 2-figure supplement 2. Rac1 GEF inhibitor does not affect MCF10A aggregate elongation.

(A) Fluorescent images of collagen fiber alignment with aggregates expressing GFP-HRasC20 treated with NSC23766 for 8 days. Collagen fibers were labeled with mCherry-CNA35 (magenta).

(B) Speed of cell movement based on nuclear tracking in rounded and elongated aggregates treated with NSC23766. (n = 82 movies)

(C) Effect of delayed inhibition of proliferation on aggregate elongation. Aggregates were cultured for 3 days before treatment with NSC23766. Data are fold change of elongation in control and drug-treated cultures (n = 40 aggregates).

All data are means ± SEM; ns, not significant, *$P<0.05$, **$P<0.01$. Data in (B,C) were analyzed with one-way ANOVA Tukey's multiple comparisons test. Data in (D) was analyzed by unpaired Student's *t*-test.

Figure 3-figure supplement 1. External gel stretching aligns collagen fiber.

(A) PDMS gel frame and stretching device.

(B) Schematic image of stretching experiment. Gel was stretched for 4 hours and then re-embedded in gel or floated in culture medium for 7 days.

(C) Second harmonic generation microscopy images of collagen gel with or without stretch. Double head arrow: stretching axis.

(D) Coherency of collagen fibers in gel and (E) distribution of fiber orientation. (N = 4 independent experiments)

All data are means ± SEM; **$P<0.01$, ***$P<0.001$. Data was analyzed by unpaired Student's *t*-test.

Figure 3-figure supplement 2. MCF10A aggregates elongate along the gel stretching axis.

(A) MCF10A aggregates in the gel after 4 hours stretching. Aggregate was co-stained with Phalloidin (green) and DAPI (blue). Collagen fibers were labeled with mCherry-CNA35 (magenta).

(B) Symmetry ratio of aggregates with or without gel stretching for 4 hours (n = 274 aggregates).

(C) Schematic image of measurement of elongation angle.

(D) The average angle of elongated aggregates in non-stretched gel and stretched gel. (n = 227 aggregates).

(E) MCF10A aggregates cultured for 1 day in non-stretched and stretched gels. Aggregates were co-stained with anti-fibronectin antibody (green) and DAPI (blue). Collagen fibrils were labeled with CNA35-mCherry (red).

All data are means ± SEM; ns, not significant, ***$P<0.001$. Data were analyzed by unpaired Student's *t*-test.

Figure 6-figure supplement 1. ERK biosensor and YAP1 localization in MCF10A cells.

(A) Time lapse images of MCF10A monolayer cells that express ERK/KTR-mClover biosensor treated with ERK inhibitor (FR180204).

(B) YAP1 localization in rounded MCF10A aggregate cultured for 1 day after transfer into collagen gel.

(C) YAP1 localization in elongated MCF10A aggregate cultured for 8 days after transfer into collagen gel.

Aggregates in (B,C) were co-stained with anti-YAP1 antibody (green), phalloidin (magenta) and DAPI (blue).

Video 1. Time lapse images of MCF10A aggregates after transferred into collagen gel.

Images were taken every 10 minutes.

Video 2. Time lapse images of aggregates embedded in stretched gel.

Images were taken every 1 hour.

A

E-cadherin / GM130 / F-actin / DAPI

day 7

20 µm

Laminin V / E-cadherin / DAPI

Lumen

day 31

50 µm

B

Membrane / GM130 / F-actin / DAPI

day 3

20 µm

F-actin / DAPI

day 10

200 µm

C

Ki67 / F-actin / DAPI   Matrigel

day7 after transplant

50 µm

A

Gel frame

9 mm

3 mm

Stretcher

7 mm

3 mm

B

Ring shape gel

Stretch
4 hours

Stretcher

Re-embed

Collagen gel

Float

Medium

C

- Stretch

+ Stretch

10 μm

10 μm

Stretching axis   0°

D

Gel coherency (%)

**

- Stretch   + Stretch

E

Collagen fibril orientation

Fibrils (%)

— - Stretch
— + Stretch

Angle (°)

A +FR180204

00:00 ERK/KTR-mClover
00:02
00:04
00:06
00:08
00:10
00:40
50 μm

B YAP1 / F-actin / DAPI   day 1   YAP1 / F-actin
50 μm

C YAP1 / F-actin / DAPI   day 8
100 μm

YAP1 / F-actin