Combined effects of salinity and temperature on grazing activity of invasive

Gammarus tigrinus on charophytes

Authors: Berthold[1, 2, *], M.; Porsche[3], C.; Hofmann[3], A., Nowak[3], P.

[1] – Applied Ecology and Phycology, University of Rostock, Rostock, Germany

[2] – Phytoplankton Ecophysiology, Mount Allison University, Sackville, Canada

[3] – Aquatic Ecology, University of Rostock, Rostock, Germany

* – corresponding author: maximilian.berthold@uni-rostock.de

Abstract

Charophytes are a refuge for zooplankton and stabilize sediments, but they are also a food

source for various animal species (water birds, fishes, invertebrates). This grazing pressure

may be an important factor prohibiting macrophyte re-colonization in eutrophic coastal

waters. Especially the introduction of new species, like *Gammarus tigrinus*, into the Baltic

Sea led to yet not understood changes in the food web. Furthermore, the occurrence of native

and invasive species is determined by their capacity to acclimatize to changing abiotic factors.

In this study we investigated the influence of temperature and salinity on the grazing pressure

of *Gammarus tigrinus* on two charophyte species: *Chara tomentosa* and *Chara aspera*. The

grazing experiments were conducted in a full factorial design with the factors salinity (3 – 13

PSU), temperature (5 – 30 °C), and charophyte species. Grazing rates were determined as

mass deviation within 48 hours considering simultaneous growth of the charophytes.

Experimental results were further used to determine gross macrophyte biomass within the

ecosystem, using field data. Gammarids preferred *C. aspera* before *C. tomentosa*, but both

were grazed within their respective growth optima. The potential grazing peak of 24 °C is not

yet reached in these coastal waters. Furthermore, calculation of gross production revealed that

macrophytes could show up to 60% and 15% more biomass without grazing in deeper light-

limiting, and shallow, not light limited water depths, respectively. The impact of food web on

coastal water restoration should therefore be equally considered, as the nutrient control.

Keywords: charophytes, grazing, *Gammarus tigrinus*, temperature, salinity

Introduction

Anthropogenic-induced eutrophication affected coastal water bodies worldwide, through

increased growth of phytoplankton mediated by elevated nutrient inputs from river, diffuse

run-off and precipitation. This increased blooming of phytoplankton subsequently increased

turbidity within the water column and led to a loss of macrophyte cover (González Sagrario et

al. 2005, Barker et al. 2008). Such primary production shifts from benthic to pelagic zone

were described as alternative stable states, and can be very stable especially in shallow aquatic

systems. This loss of submerged vegetation had several implications on ecosystem

functioning and services, e.g. , altered species composition (Suikkanen et al. 2013) and

possibility of potentially harmful algal blooms (Paerl and Paul 2012). To counteract these

developments, nutrient inputs as one eutrophication-causing factor were reduced (Gulati and

Van Donk 2002, Nausch et al. 2011). However, these reductions worked only partly and, in

some cases, macrophytes need a long time frame to re-colonize (Sand-Jensen et al. 2017).

One theory emphasizes, that the current phytoplankton dominated state has strong own

feedback mechanisms, like increasing attenuation, that inhibit a return to pre-eutrophication

conditions (Scheffer et al. 1997). Furthermore, Global Change will affect water temperatures

worldwide, as well as altering run-off from land, leading to possibly different salinity

conditions within the next decades (Schindler 2006). The Baltic Sea is such a system prone to

eutrophication and Global Change, due to already increased sear surface temperatures,

compared to other oceans worldwide (Reusch et al. 2018). The Baltic is subject to large

phytoplankton blooms since decades, with the riparian states working to reduce the impact of

nutrient-run-off to the sea (Janssen et al. 2004, HELCOM 2018). Similar, the coastal water

bodies of the Baltic Sea were impacted by eutrophication and shifts on primary producer

dominance, i.e. from submerged macrophytes to phytoplankton (Schiewer 2007). Even

though nutrient inputs declined in the Baltic and its adjacent coastal water bodies through the

last decades, most coastal water bodies (CWB) remain eutrophied (Berthold et al. 2018a).

However, there were also hypotheses that not only nutrients, but also top-down control (i.e.

grazing) may affect re-mesotrophication and macrophyte recolonization (Körner and Dugdale

2003, Östman et al. 2016).

Besides (accidental) grazing from native species (Körner and Dugdale 2003), there is also an

increasing grazing pressure of invasive species introduced through, e.g. ballast water of ships.

One of these species is the gammarid *Gammarus tigrinus* (Sexton 1939), a successful new

species introduced from North America to European waters (Berezina 2007). This gammarid

was described to colonize shallow waters, especially within reed, and soft bottom systems of

the Baltic Sea (Daunys and Zettler 2006, Kotta et al. 2011). Amid Global Change, invasive

species can alter known food web interactions and may counteract re-colonization of former

native primary producers (Puntila 2016).

Charophytes receive attention as indicator species for oligotrophic water bodies (e.g. Melzer

1999), and were listed as 'endangered species' in Germany. However, this view has been

relativized by the fact that during a Germany-wide survey, an increased number of charophyte

populations were reported at strongly eutrophic sites (Doege et al. 2016). Blümel et al. (2002)

found charophyte species in the Darss-Zingst lagoon system (DZLS) even with high nutrient

levels and turbidity (Selig et al. 2009). An explanation was offered by Schubert et al. (2018),

who concluded that many charophyte species are "pioneer plants" reacting particularly

sensitive to the competitive pressure of higher plants. Shading in particular reduces the

competition with other species. This distribution pattern means that charophytes can be found

in shallow water areas even under highly eutrophic conditions, if the level of light exposure

and the water level stability admits it. Nonetheless, charophytes are also under pressure in

other parts of the Baltic Sea. Comparisons of historical and recent distributions in the northern

Baltic found that *C. aspera* and *C. tomentosa* were found 8 and 86% times less in the most

recent transect, respectively (Pitkänen et al. 2013).

Salinity constitutes another factor in brackish water, where the increased salinity prevents the

growth of macrophyte species (Lindner 1972). This means that regardless of the degree of

trophism, a niche is created in which charophytes can be established. It was assumed that

charophytes have several turbidity-reducing traits, like e.g. dissolved nutrient competition

with phytoplankton, refuge function for zooplankton and young fish, or reducing of

resuspension of particulate matter (Kufel and Kufel 2002). Therefore, re-colonization with

those species could create feedback mechanism, favoring a macrophyte-dominated stable

state. Charophytes at the southern Baltic Sea coast diminished in some CWB already in the

1930ies (Schlungbaum et al. 2000), and later in adjacent CWB as well (Blümel et al. 2002).

Even though nutrient concentrations were reduced to a comparable level between today and

the 1930ies, no dense macrophyte cover was able to form in e.g. the Darss-Zingst lagoon

system (DZLS) (Gessner 1957, Berthold et al. 2018a, Paar et al. 2021). Indeed, the maximum

depth distribution of charophytes has not increased significantly in the DZLS during the last

years, and charophytes are still found only in very shallow waters, where light is not limiting

(Piepho 2017). The DZLS can be considered as model ecosystem for the southern Baltic Sea,

and is monitored for decades (see Schiewer 2007). The process of sudden alternative stable

state shifts were observed in the 1930s in the innermost part, and in the 1970s in the outer

parts (Gessner 1957, Schiewer 2007, Berthold 2016). This state-shift caused a major decline

in submerged vegetation cover (Blümel et al. 2002). Restoration measures starting thirty years

ago, lowered the nutrient inputs by up to 90% for Phosphorus and 60% for Nitrogen (Nausch

et al. 2011, LUNG 2013). However, the macrophyte covers were not able to recover until

today.

The question evolves, if re-colonization of macrophytes is additionally hampered by grazing

pressure of (newly introduced) mesograzers, and how this pressure changes throughout the

year. *Gammarus tigrinus* and charophytes occur within the same depth distribution, and

grazing can potentially happen. It was questioned what for effects the variable temperature,

and salinities in lagoons have on grazing behavior on charophytes. Therefore, a full-factorial

design was chosen with the variables water temperature and salinity at four levels. This

design would also allow to discuss future developments within CWB at e.g. rising water

temperatures, or reduced salinities at increasing precipitation events. Shallow CWB are likely

to be more affected by such events due to their relatively lower water column and close

coupling with the adjacent land. Therefore, we combined the laboratory determined grazing

potentials with field-determined elemental budgets of submerged macrophytes of the DZLS.

The data sets were further used to calculate potential nutrient flows from the macrophytes to

the water column, therefore putting macrophytes as nutrient mediators between the sediment

and higher trophic levels. It was hypothesized that there is either a combination of factors

(abiotic and biotic) that amplify mass loss on charophytes, or that some abiotic factors quench

each other, e.g. lower salinity amplifies/buffers effects of higher temperatures. Those mass

losses may depend on the acclimation potential of the local population.

Material and Methods

Experimental design

We chose a full factorial design with three factors, for which one factor relates to species and includes two levels (*Chara aspera* and *C. tomentosa*), and the other two factors each with four levels are water temperature (9, 15, 24, and 30 °C), and salinity (3, 5, 7, 13). These salinity levels represent the salinity gradient from the Western belt Sea to the Gulf of Finland, covering most of the Baltic Sea area (HELCOM 2010). The water temperature levels represent the blooming period of shallow coastal waters (winter to summer +1 – +22 °C, Schiewer 2007), with 30 ° C as prospective endpoint of possible charophyte growth. Charophytes were cultured from intact plants extracted from the DZLS. The DZLS is formed by four consecutive shallow lagoons, and shows a salinity and trophic gradient from the main river inflows Recknitz and Barthe (inner and middle part, salinity of 1 – 6, eutrophic) to the open Baltic Sea (salinity of 10 – 12, mesotrophic, Schumann et al. 2006). Salinity varies around 6 to 8 at the respective sampling locations in the Bodstedter Bodden (54°22'30,5" N, 12°34'14,9" E) and the Grabow (54°22'2,3" N, 12°48'27" E) (Schumann et al. 2006). During the experiment, two charophyte species were used, *Chara tomentosa* and *Chara aspera*. Both species are common in the Baltic Sea, with *C. aspera* covering the widest salinity range among charophytes (Blindow 2000). Along the west coast of Sweden, *C. aspera* is regularly found at sites with salinities up to 15 (temporarily up to 20), but it is also common in freshwater (Hasslow 1931, Blindow 2000). *Chara tomentosa* is among the largest charophyte species and can be found in freshwaters all over the world. Only in the Baltic Sea, *C. tomentosa* also occurs in brackish water, with a salinity range between 0-7.5 (Björkman 1947, Torn et al. 2003). Gammarids were derived from the Aquaculture section in Born of the State Research Centre for Agriculture and Fishery Mecklenburg-Vorpommern. This research facility is located at the DZLS and uses brackish water from the lagoon with a salinity of 4 –

6. Gammarids were identified by morphology and genetic analyses (see below).

Charophytes were cultured in beakers (vol. 500 ml), containing sediment containers (vol. 64

ml) with acid-rinsed, phosphorus-enriched sediment (6 g P kg^{-1} sediment) according to

Wüstenberg et al. (2011). Mineral growth media was prepared to culture charophytes (Pohl et

al. 1987, Wüstenberg et al. 2011), but was adjusted to the respective salinities by adding

marine salt (Instant Ocean, Aquarium Systems, Sarrebourg). Salinity was checked with a

salinometer (Multiline P4, WTW). Each of the 16 combinations were replicated eight times

with gammarids and without gammarids present (control) (Fig. 1). Gammarids were pre-

acclimatized to experimental conditions by culturing them at the respective temperature and

salinity combinations. Ten to 20 individuals were simultaneously cultured per temperature-

salinity combination, aerating all cultures. Charophytes were added as food supplement, as

well as resting surface for the gammarids.

<u>Field sampling and analyses</u>

In addition, we used elemental and biomass composition of submerged macrophytes from the

DZLS, to compare the experimental derived-grazing rates to the actual ecosystem. Field

sampling was conducted in the Bodstedter Bodden and the Grabow in 2014, as part of the

BACOSA project. Submerged macrophytes were sampled throughout the year, first in 54x54

cm plots, and later in the year in 25×25 cm plots at nine different depths. Here, three sampling

locations were pooled into three depth zones, according to water depth and proximity to the

reed belt. Water depths for both stations varied with water levels in the respective lagoons and

were shallow (30 – 60 cm), intermediate (60 – 90 cm), and deep (90 – 120 cm). The pooling

became necessary, as macrophytes grew patchy at each sampling station. Sampled

macrophytes were separated on the genus level. Whole plants, including roots, were wet-

weighed and then dried at 60 °C for 24 h. Afterwards, plants dry mass (DM) were re-weighed

again to calculate water content. Dried plant material was ground with a ball mill. The powder

was treated with 1M HCl to remove inorganic carbon residues and then weighed in tin-

container (1×0.5 cm). These samples were analyzed for their carbon and nitrogen content

using an elemental analyzer (varioEL III). The analyzer was calibrated using acetanilide (~5

mg per sample).

Aliquots of dried plant material were weighed in crucibles and combusted at 550 °C for 4 h.

The ash was re-weighed to calculate ash-free dry mass. Ashes were used to determine total

phosphorus content (mg P g^{-1} dry mass), using an acid persulfate extraction (see Berthold et

al. 2015).

Charophyte growth experiment

Growth responses of *C. tomentosa* and *C. aspera* to the different culture conditions were

followed. Fresh mass was determined before and at the end of the experiment by blotting

them dry with a paper towel. Experiments lasted 48 to 72 hours and were simultaneously the

gammarid-free controls of the grazing experiment. The differences in mass were used to

calculate short-term developments and compensate for possible growth in grazing

experiments (see below).

Gammarid grazing experiments

For grazing experiments, charophytes were wet weighed and set into beakers filled with

sediment and growth media of the respective salinity. Gammarids were photographed before

and after the experiment to determine possible increases in length. One charophyte and five

gammarids were set together in one beaker. Each replicate run lasted for 48 hours and in four

cases over the weekend, i.e. for 72 hours. Replicate runs for each *Chara* species were

performed over a period of four weeks, with two replicates of each respective temperature-

salinity combination at a time. Charophytes were wet weighed after each replicate run to

calculate mass differences during the replicate run. Dead gammarids were counted at the end

of each replicate run and replaced for the next replicate run. If possible, the same gammarids

were used throughout all replicate runs. Equations 1 – 3 were used to extract grazing rates of

gammarids from growth rates of charophytes during the experiments. Grazing rates were

treated with an error-propagation to account for experimental variability.

Equation 1 subtract the mean effect of pure growth of the control (C) from the treatment

which also includes grazing (T):

$$G_i = T_i - \bar{C} \ \{i = 1..n$$

199   Equation 2 standardizes the grazing effect from prior formula (**z**-score):

$$Gz_i = \frac{(G_i - \bar{G})}{\sigma_G} \ \{i: 1..n$$

200   Equation 3 calculates the mean of pure grazing and include its error caused by the experiment:

$$Gf_i = \bar{G} + Gz_i \cdot \sqrt{|(\sigma_T^2 - \sigma_C^2)|} \ \{i: 1..n$$

T = treatment, C = control, G= detrended grazing, G_z = standardized grazing, G_f = real

grazing with included experimental error, n = replicates

<u>Data analyses</u>

We further used the experimental grazing data to calculate grazing rates based on salinity and

water temperature at two sites of the DZLS. The data set failed to show homogeneity of the

residuals, and was therefore Box-Cox power transformed (*bcnPower* function, R package

"*car*", Fox and Weisberg 2019). Finally, general linear models (GLM, *glm* function, R

package "stats", R Core Team 2019) and generalized additive models (GAM, *gam* function, R

package *"mcgv",* Wood 2011), both with Gaussian distributional assumptions, were

compared by the Akaike Information Criterion (AIC, R package "stats", R Core Team 2019)

and the Bayesian Information Criterion (BIC, R package *"stats",* R Core Team 2019),

choosing the model with low AIC and BIC values. We chose the GAM (with knots, k = 16) as

the best model to reflect the experimental data (see Results). Recent biomass and individual

numbers of gammarids were modelled according to described numbers within the DZLS and

nearby lagoons (Zettler 1995, M. Paar personal communication). We assumed that *Gammarus*

tigrinus populations followed the same seasonal pattern as described in Chambers (1977). We

used again a GAM model to calculate the seasonal population development of *Gammarus*

tigrinus, normalized the predicted seasonal data to have a maximum of 1. Then, we used this

normalized prediction to calculate population densities with a maximum of 400 Ind m^{-2}

(Zettler 1995, M. Paar personal communication, supplement figure S3). To calculate seasonal

grazing rates within the DZLS, these population numbers were combined with the individual

grazing rates of the temperature-salinity model with field data from the two sites of the DZLS

(Equation 4). We used a fixed grazing preference of 20% macrophyte share in gammarid diets

(Pellan et al. 2016). Furthermore, we used the GAM to predict the charophyte development as

function of sampling date (k = 8) at two sites of the DZLS. We calculated daily biomass

increase and subsequently added gammarid grazing rates to determine the potential gross

biomass production per site (see Results and Discussion below).

Equation 4 calculates the seasonal grazing rates per square meter and day for specific sites:

$$Graz_t = G_{GAM_t}\{T_t; S_t\} \cdot \delta_t \cdot 0.2 \quad \{t: 1..n$$

229   t = day of the year, $Graz_t$ = grazing at day t (mg·m^{-2}·d^{-1}), G_{GAMt} = modelled grazing rate of one

230   individual at time t with given temperature (T) and salinity (S) from specific site at time t, δ_t =

231   population density at time t (individuals·m^{-2}), 0.2 = reflects the 20 % fixed grazing preference,

232   n = length of the year in days

 <u>Genetic analyses</u>

At the end of the grazing experiment, 69 gammarids were determined as *Gammarus tigrinus*

following Zettler and Zettler (2017) and preserved in 80 % ethanol. To verify the

morphological analyses, a genetic approach was conducted on 15 individuals. Total genomic

DNA was extracted using a silica spin column procedure with the DNeasy Blood and Tissue

Kit (Qiagen, Hilden, Germany) following the protocol provided by the manufacturer. Partial

sequences of the COI gene were amplified with the universal primers LCO 1490 and HCO

2198 (Folmer et al. 1994).

PCR amplifications were performed with a denaturation step for 60 s at 94°C, followed by 5

cycles of: 60 s at 94°C, 90 s at 45°C and 60 s at 72°C, 35 cycles of: 60 s at 94°C, 90 s at 51°C

and 60 s at 72°C, and completed with 5 min at 72°C as a final extension step. PCR was

performed in a 30 µl reaction volume with a Taq PCR Master Mix (Qiagen, Hilden,

Germany) consisting 2.5 mM MgCl2, and 0.5 pmol of each primer (final concentration). The

PCR products were extracted from agarose gels according to the protocol of the Biometra-

innuPrep Gel Extraction Kit (Analytik Jena, Jena, Germany), and were sequenced directly

using a 3130×L Genetic Analyzer (Applied Biosystems, NY, USA) with sequencing primers

identical to the primers that were used for the PCR reaction. Obtained sequences were quality

controlled and aligned via the BIOEDIT software (Hall 1999). The COI gene is widely used

for inter- and intraspecific diversification questions concerning amphipod taxa. For the

taxonomic determination addition GenBank sequences of *Gammarus tigrinus* (see

Supplement Table S1), *G. duebeni* Liljeborg, 1852 (EU421779), *G. locusta* Linnaeus, 1758

(KT209211), *G. oceanicus* Segerstråle, 1947 (GQ341809), *G. pulex* Linnaeus, 1758

(MN400977), *G. roeseli* Gervais, 1835 (EF570337) *G. salinus* Spooner, 1947 (KT208533),

and *G. zaddachi* Sexton, 1912 (KU845083) were used for the phylogenetic analyses.

In order to estimate the evolutionary divergence between haplotypes, pair-wise uncorrected p-

distances and the number of substitutions were conducted using MEGA version X (Kumar et

al. 2018). To uncover phylogenetic relationships, a Maximum likelihood tree was constructed

using MEGA version X (Kumar et al. 2018) on the basis of the Kimura 2-parameter model.

Branch support for the nodes was calculated from 1000 bootstrap replicates.

To study the distribution of mtDNA sequence diversity, a haplotype network was constructed

using the Median-joining algorithm (Bandelt et al. 1999) implemented in PopART (Leigh and

Bryant 2015). The distribution basemap was created with QGIS.org software

(http://www.qgis.org) and modified in CorelDRAW (Corel Corporation, Ottawa, Ontario,

Canada).

Results

Genetic data

The present study analysed 15 specimens of *Gammarus tigrinus* collected at the DZLS with

sequence data from previous studies (see Table S1) to validate morphological species

determination and identify their genetic haplotypes. Based on the 407 bp region of the COI,

the 15 analysed individuals could be assigned to the northern species of *G. tigrinus*. The

phylogenetic analyses revealed that the individuals of this study belonged to two different,

very closely related haplotypes: N4a & N4e (Fig 2&3). Both haplotypes differed by a single

base pair substitution and were identical with *G. tigrinus* specimens from Europe (N4a&N4e),

the British Isles (N4e-1) and North Amerika (N4a-4). Both haplotypes were sympatrically

present in the DZLS and were observed in both fresh and brackish water habitats. Haplotype

N4e-1 comprised 4 individuals of the DZLS and was already identified in previous studies in

the Baltic Sea and in freshwater habitats in Northern Ireland. 11 DZLS individuals

corresponded to haplotype N4a-4, which has been described at the Baltic Sea coast, but also

in North America and moreover it was found in the northern Rhine (freshwater habitat).

Growth and survival rate

Starting biomasses of *C. aspera* in control and experimental beaker ranged from 50 to up to

300 mg, with the most macrophytes being within the range of 100 – 175 mg (70%). After two

days, biomass distribution of *C. aspera* in control beacons stayed relatively the same, but

smaller and larger biomass proportions increased. Biomass of *C. aspera* decreased with

increasing temperature at salinities of 3 and 13. Contrary, the biomass stayed almost constant

at all temperatures at salinities of 5. A small possible biomass increase was detected at

temperatures 15 and 24 degrees at salinities of 7, and a strong decrease at 30 degrees for

almost all salinities (Supplement Figure S1).

Starting biomass of *C. tomentosa* in control beacons ranged from 75 to up to 600 mg, and

40% of all macrophytes were within the range of 100 – 125 mg at the beginning. Contrary to

C. aspera, biomass of *C. tomentosa* in experimental beacons was close to a normal

distribution with 50% of all macrophytes being within the range 175 – 225 mg at the

beginning. The biomass distribution of *C. tomentosa* in control beacons stayed almost the

same, but with a trend to smaller biomasses, indicating biomass loss even without grazing.

Biomass of *C. tomentosa* stayed very constant at almost every temperature-salinity

combination. There was a possible growth at temperatures of 15 to 24 degrees at salinities of

7 (Supplement Figure 1). In total, these results indicate an overall stable experimental

approach.

Two thirds of the time (65%) all gammarids survived the incubation period at all temperature-

salinity-macrophyte combinations. Additionally, in 30% of the time, only one gammarid died

during incubation. These results indicate overall stable culture conditions and the capability of

these gammarids to survive a wide range of temperature-salinity gradients.

<u>Grazing rate</u>

Growth-corrected grazing ranged between 1 to 10 mg DM per individual and day on *C.*

aspera. Grazing rates peaked at 24 °C, except for salinity concentrations of 5, where it

probably peaked at 30 °C. This grazing rate was contrary to all other salinities, were grazing

rates at 30 °C were lower compared to 24 °C. Furthermore, growth of *C. aspera* was less

affected of high temperatures at salinity of 5, compared to other salinities (see Figure 4 and

Supplement Figure S2). Growth-corrected grazing ranged between 1 to 5 mg DM per

individual and day on *C. tomentosa*. Grazing rates peaked at 24 °C for all salinity levels and

at 15 °C for salinities of 7. Interestingly, grazing rates of gammarids coincided with growth

optima of the charophytes in this study. These results may indicate a grazing preference on

actively growing charophytes. In general, grazing rates on *C. aspera* were up to two times

higher compared to *C. tomentosa*. The grazing rate considering both charophytes to be

present, would have been up to 5.5 mg Ind^{-1} d^{-1}, with a general larger temperature-salinity

range of highest observed grazing.

<u>Field data results and model</u>

Individual daily grazing rates within the system would range from 0.1 up to 5 mg Ind^{-1} d^{-1},

when applying the predicted GAM output on field observations of temperature and salinity

(Figure 5). Grazing rates would peak during summer and be lowest at winter at the two

sampling stations of the DZLS (see Figure 5). The here used individuals of *G. tigrinus* grazed

therefore twice as much at the experimental 24 °C, as would be expected from their initial

harvesting season in April and May, indicating acclimation processes. Interestingly, grazing

rates differ between both stations, even though the difference in salinity is only 1 – 2 PSU.

The yearly potential grazing rate for *G. tigrinus* on charophytes ranges from 37.2 to 42.5 g m^{-}

^{2}, considering regional salinity and temperature patterns within the DZLS. These values are

based on the assumptions of a maximum of 400 Ind m^{-2} and a food preference of 20% for

macrophytes compared to other available food sources (see discussion below).

Charophyte biomass was highest in the shallow areas, close to the reed belt at both sampling

(Figure 6). Biomass in the shallow zone peaked in July in the less eutrophic Grabow, and in

October in the highly eutrophic Bodstedter Bodden. Contrary, charophyte biomass was higher

in the Bodstedter Bodden at intermediate distances/water depths compared to the Grabow.

However, Grabow biomass peaked during summer, whereas biomass in the Bodstedter

Bodden became depressed during that time. There were no regular biomasses of charophytes

in the deeper water areas at both sampling locations, indicating a possible light limitation

from shallow to deep.

There were no significant differences for the C:N and C:P ratios along the depth distribution,

or between stations (see pooled values in Fig 7). Interestingly, charophytes showed either a

weak seasonal trend in elemental composition (C:N), or the ratio decreased (C:P). Lowest C:P

ratios were found from July to October. Contrary, co-occurring tracheophytes increased their

C:N ratio during summer, as well as their C:P ratios. These findings indicate that charophytes

had probably a higher protein content per biomass compared to tracheophytes.

Discussion

Grazing rates in our experiment and in the field depend on several factors, like the population

structure of gammarids, available food and food quality, occurrence/absence of (competing)

predators, and ultimately habitat choice and niche occupation.

Populations of *G. tigrinus* were described to turn-over at least three times per year, with a

peak of large animals (6 – 9 mm) in spring, and a concomitant shift to smaller animals in late

summer (2 – 4 mm) (Chambers 1977). We used for all our experiments animals in the size

range of 6 – 12 mm, i.e. large adults. This size class is only dominant by up to 60% until May,

and is almost completely replaced by smaller animals from June to September (Chambers

1977). This pre-selection had likely an effect on the described grazing rates, as metabolic

rates of *G. tigrinus* increase with lower weight in form of a power function (Normant et al.

2007). However, the gammarids in our study weight from 10 – 90 mg wet mass Ind^{-1} (data not

shown), whereas the gammarids of Normant et al. were caught in August, and weighed only 5

– 25 mg wet weight Ind^{-1}. We therefore calculated with the provided power function, that our

animals showed even lower specific metabolic rates of 0.25 – 1.5 mW g^{-1} wet mass, compared

to 1 – 3.5 mW g^{-1} wet mass in smaller gammarids (Normant et al. 2007). This reduced

metabolic rate had likely an impact on the grazing rates we found in this study, as grazing for

metabolic upkeep is lower in large adult animals. Our rates are therefore only representative

for springtime, if similar population dynamics are assumed for our system like described in

Chambers (1977).

To our knowledge, published studies on seasonal population dynamics of *G. tigrinus* are

currently missing in the system studied here, as elsewhere in the Baltic Sea. However, recent

field studies found 5 to 2200 Ind m^{-2} (gammarids on genus level) in the DZLS and the

adjacent Westrügensche lagoon system. Largest, but fewest animals were found early in the

year (22 – 28 mg Ind^{-1}), with decreasing mass, but increasing abundance in summer (3.5 mg

Ind^{-1}) (M. Paar, personal communication). These patterns point to the same seasonal

development as described in Chambers (1977). If we assume that *G. tigrinus* constitutes 20%

of the local gammarid population in the DZLS (Zettler 2001), we get a population size of up

to 450 Ind m^{-2}. These numbers are in agreement with first described abundances of *G. tigrinus*

of 300 – 500 Ind m^{-2} in the phytal and reed belt of the DZLS (Zettler 1995), or 30 – 110 Ind

m^{-2} in the Neva estuary (Baltic Sea, Berezina 2007).

Nonetheless, springtime is a crucial period for charophyte sprouting, as the active growth of

for example *C. tomentosa* takes place during a relatively short period at the beginning of

summer (Torn et al. 2006). Furthermore, young sprouted charophytes show a higher

elemental content per biomass, than for example tracheophytes (Volkmann 2016, this study),

making them a preferred grazing opportunity (Kraufvelin et al. 2006). *G. locusta* follows a

preference for high value plant material and selectively feeds on more nutrient-rich

macrophytes, especially periphyton, brown and green algae (Kraufvelin et al. 2006). If *G.*

tigrinus shows similar grazing preferences, periphyton from tracheophytes would be the first

choice followed by charophytes in our system. Periphyton (Sanudo-Wilhelmy et al. 2004) and

possibly charophytes (Kufel and Kufel 2002) can take up fast amounts of nutrient pulses that

are likely to happen close to the wetland in this area (Berthold et al. 2018b). The charophytes

in our study area showed higher C:N and N:P ratios, then tracheophytes, making them more

likely to gammarid grazing as high-value food. Furthermore, this pre-selection of high-value

food may explain the differences in grazing rates on *C. tomentosa* vs. *C. aspera* found in this

study. *C. tomentosa* can show higher C:N ratios (44:1) than *C. aspera* (27:1), making it

maybe a less favorable food. Furthermore, *C. aspera* showed in our study signs of early

decay, contrary to *C. tomentosa* (see trends in growth control beacons, supplement figure S1).

Such beginning decay can weaken *C. aspera* and make it more susceptible to grazing

(Buchsbaum et al. 1991), explaining the deviation of grazing rates between both charophytes

found in this study. Tracheophytes in our and adjacent lagoons can show high colonization

rates by epiphytes (Paar et al. 2021), offering an additional plant food source to charophytes.

Furthermore, food preferences can change throughout the year depending on gammarid life

stage and temperature (Felten et al. 2008, Pellan et al. 2016). Plant biomass becomes

increasingly important, when temperatures increase and can reach up to 20 – 30% of total

food consumed (Pellan et al. 2016). Detritus, plant litter, or animal matter, like chironomides,

are usually important food sources during winter time (Felten et al. 2008, Pellan et al. 2016).

Sediment organic content can be as high as 20% in sediments of the DZLS (Berthold et al.

2018c). Sediment detritus was not an alternative source in our experiments, as we used only

inorganic sand. Plant litter occurs abundantly within the reed belt during winter time in the

DZLS (Karstens et al. 2016). Chironodmides biomass can be as high as 40 g m^{-3} and is

positively impacted by the occurrence of another invasive species, *Marenzelaeria viridis*

(Zettler 1995, Schiewer 2007). Functional feeding in gammarids relies furthermore on

population composition and food availability (Kelly et al. 2002). Our experimental results

show therefore only a potential grazing rate, as we did not represent the variety and

abundance of alternative food sources from within the ecosystem. However, our suggested

weighed grazing rates of only 20% macrophyte biomass may still be a realistic assumption

(see Figure 6).

Furthermore, grazing rates depend on niche occupation and competition with native and other

invasive species. *G. tigrinus* can probably outcompete the native *G. salinus*, if both depend on

P. littoralis in the northern Baltic Sea (Orav-Kotta et al. 2009). Both gammarids had the

potential to exceed daily macrophyte production during summer (Orav-Kotta et al. 2009).

Comparably, we found that macrophyte biomass could have been at least 25% higher during

summer, without the impact of *G. tigrinus* grazing. However, these results are only valid, if

G. tigrinus outcompeted other gammarids in our study area. Otherwise, grazing rates on

charopytes could be even higher.

Additionally, microhabitats, like macrophyte patches, are important for native species to co-

exist with invasive species (Korpinen and Westerbom 2009). Nonetheless, recent studies

showed that *G. tigrinus* would compete with native species, either directly through food

choices (Orav-Kotta et al. 2009), or grazing on native juveniles (Jänes et al. 2015). *G. tigrinus*

may be more tolerant against heat waves and sub-oxic conditions, than native species (Lenz et

al. 2011). These broader temperature acclimation may help to permanently introduce *G.*

tigrinus within the Baltic Sea, as sea surface temperature there are already higher, then in

other ocean parts (Reusch et al. 2018). Furthermore, endurance of sub-oxic conditions is an

advantage in eutrophic coastal waters of the Baltic, as redox conditions within the reed belt

can change fast (Karstens et al. 2015), and an increase of nutrients is challenging submerged

macrophytes in the coastal water bodies of the southern Baltic Sea (Paar et al. 2021), stressing

habitats of native gammarids. Contrary, an *in silico* study assumed that *G. tigrinus* has

actually a narrower niche than native gammarids in the norther Baltic (Herkül et al. 2016).

These results are somewhat contradictory to other studies in this area (Korpinen and

Westerbom 2009, Kotta et al. 2011, Lenz et al. 2011), and may have not considered the

impact of haplotype-diversity, that means locally adapted populations of *G. tigrinus*. Different

studies revealed high genetic variability of *G. tigrinus* in the Baltic Sea (this study, Kelly et

al. 2006) and Paiva et al. (2018) indicated that the invasive species may have different

tolerance and different limits to salinities due spatially varying selection among populations.

The patchy macrophyte stands found in the DZLS, and in other eutrophic water bodies, may

affect the occurrence of *G. tigrinus*, and therefore the grazing rates, especially in intermediate

and deeper waters. Palaemonid prawns put a higher grazing pressure on *G. tigrinus* in

unvegetated mesocosms, then in vegetated ones (Kuprijanov et al. 2015). It is likely that *G.*

tigrinus stays close to the reed belt, as described for this lagoon and other coastal waters

(Zettler 1995, 2001). This spatial distribution would further hamper charophyte

recolonization, as the shallow areas in the DZLS are the only spots, where light availability is

high enough to support charophyte biomass (see Figure 6, Piepho (2017). Furthermore,

elemental concentrations of reed, macrophytes and sediment is highest close to the land

(Berthold et al. 2018b). Grazing on macrophytes can impact nutrient sequestration rates by

increasing grazing-mediated release rates, therefore slowing down restoration measures to

reduce eutrophication.

Acknowledgements

The authors thank the State Agency for Environment, Nature Conservation and Geology
Mecklenburg-Vorpommern for providing monitoring data and the Aquaculture section in
Born of the State Research Centre for Agriculture and Fishery Mecklenburg-Vorpommern for
providing gammarids. We also want to thank Dr. Ralf Bastrop for providing the respective
primers, and Dr. Martin Paar for providing us with recent data on gammarid distribution in
the southern Baltic Sea.

References

Bandelt, H. J., P. Forster, and A. Rohl. 1999. Median-joining networks for inferring
    intraspecific phylogenies. Molecular Biology and Evolution 16:37–48.

Barker, T., K. Hatton, M. O'Connor, L. Connor, and B. Moss. 2008. Effects of nitrate load on
    submerged plant biomass and species richness: results of a mesocosm experiment.
    Fundamental and Applied Limnology / Archiv für Hydrobiologie 173:89–100.

Berezina, N. A. 2007. Expansion of the North American amphipod Gammarus tigrinus
    Sexton, 1939 to the Neva Estuary (easternmost Baltic Sea). Oceanologia 49:129–135.

Berthold, M. 2016. The influence of phosphorus on the eutrophication process in the Darß-
    Zingst Bodden chain. University of Rostock.

Berthold, M., U. Karsten, M. von Weber, A. Bachor, and R. Schumann. 2018a. Phytoplankton
    can bypass nutrient reductions in eutrophic coastal water bodies. Ambio 47 (1):146–158.

Berthold, M., S. Karstens, U. Buczko, and R. Schumann. 2018b. Potential export of soluble
    reactive phosphorus from a coastal wetland in a cold-temperate lagoon system: Buffer
    capacities of macrophytes and impact on phytoplankton. Science of the Total
    Environment 616–617:46–54.

Berthold, M., D. Zimmer, V. Reiff, and R. Schumann. 2018c. Phosphorus Contents Re-visited

After 40 Years in Muddy and Sandy Sediments of a Temperate Lagoon System.

Frontiers in Marine Science 5:1–14.

Berthold, M., D. Zimmer, and R. Schumann. 2015. A simplified method for total phosphorus

digestion with potassium persulphate at sub-boiling temperatures in different

environmental samples. Rostocker Meeresbiologische Beiträge 25:7–25.

Björkman, S. O. 1947. On the distribution of Chara tomentosa L. round the Baltic and some

remarks on its specific epithet. Botaniska Notiser 1947:157–170.

Blindow, I. 2000. Distribution of charophytes along the Swedish coastin relation to salinity

and eutrophication. International Review of Hydrobiology 85:707–717.

Blümel, C., A. Domin, J. C. Krause, M. Schubert, U. Schiewer, and H. Schubert. 2002. Der

historische Makrophytenbewuchs der inneren Gewässer der deutschen Ostseeküste.

Rostocker Meeresbiologische Beiträge 10:5–111.

Buchsbaum, R., I. Valiela, T. Swain, M. Dzierzeski, and S. Allen. 1991. Available and

refractory nitrogen in detritus of coastal vascular plants and macroalgae. Marine Ecology

Progress Series 72:131–143.

Chambers, M. R. 1977. The population ecology of Gammarus tigrinus (sexton) in the reed

beds of the Tjeukemeer. Hydrobiologia 53:155–164.

Daunys, D., and M. L. Zettler. 2006. Invasion of the North Amrican Amphipod (Gammarus

tigrinus Sexton, 1939) into the Curonian Lagoon, South-Eastern Baltic Sea. Acta

Zoologica Lituanica 16:20–26.

Doege, A., K. van de Weyer, R. Becker, and H. Schubert. 2016. Bioindikation mit Characeen.

Pages 97–137 Armleuchteralgen. Springer Berlin Heidelberg, Berlin, Heidelberg.

Felten, V., G. Tixier, F. Guérold, V. De Crespin De Billy, O. Dangles, F. Guerold, De Crespin

V. De Billy, O. Dangles, F. Guérold, V. De Crespin De Billy, and O. Dangles. 2008.

Quantification of diet variability in a stream amphipod: Implications for ecosystem

functioning. Fundamental and Applied Limnology / Archiv für Hydrobiologie 170:303–

313.

Folmer, O., M. Black, W. Hoeh, R. Lutz, and R. Vrijenhoek. 1994. DNA primers for

amplification of mitochondrial cytochrome c oxidase subunit I from diverse metazoan

invertebrates. Molecular marine biology and biotechnology 3:294–299.

Fox, J., and S. Weisberg. 2019. An {R} Companion to Applied Regression. Third. Sage,

Thousand Oaks {CA}.

Gessner, F. 1957. Meer und Strand. Page (F. Gessner, Ed.). Second edition. VEB Deutscher

Verlag der Wissenschaften, Berlin.

González Sagrario, M. A., E. Jeppesen, J. Gomà, M. Søndergaard, J. P. Jensen, T. Lauridsen,

and F. Landkildehus. 2005. Does high nitrogen loading prevent clear-water conditions in

shallow lakes at moderately high phosphorus concentrations? Freshwater Biology 50:27–

41.

Gulati, R. D., and E. Van Donk. 2002. Lakes in the Netherlands, their origin, eutrophication

and restoration: State-of-the-art review. Hydrobiologia 478:73–106.

Hasslow, O. J. 1931. Sveriges characéer. Botaniska Notiser:63–136.

HELCOM. 2010. Ecosystem Health of the Baltic sea 2003-2007: HELCOM Initial Holistic

Assessment. Baltic Sea Environment Proceedings 122:68.

HELCOM. 2018. Sources and pathways of nutrients to the Baltic Sea. Baltic Sea

Environmental Proceedings 153:48.

Herkül, K., V. Lauringson, and J. Kotta. 2016. Specialization among amphipods: The invasive

Gammarus tigrinus has narrower niche space compared to native gammarids. Ecosphere

7:1–16.

Jänes, H., J. Kotta, and K. Herkül. 2015. High fecundity and predation pressure of the

invasive Gammarus tigrinus cause decline of indigenous gammarids. Estuarine, Coastal

and Shelf Science 165:185–189.

Janssen, F., T. Neumann, and M. Schmidt. 2004. Inter-annual variability in cyanobacteria

blooms in the Baltic Sea controlled by wintertime hydrographic conditions. Marine

Ecology Progress Series 275:59–68.

Karstens, S., U. Buczko, and S. Glatzel. 2015. Phosphorus storage and mobilization in coastal

Phragmites wetlands: Influence of local-scale hydrodynamics. Estuarine, Coastal and

Shelf Science 164:124–133.

Karstens, S., U. Buczko, G. Jurasinski, R. Peticzka, and S. Glatzel. 2016. Impact of adjacent

land use on coastal wetland sediments. Science of the Total Environment 550:337–348.

Kelly, D. W., J. T. A. Dick, and W. I. Montgomery. 2002. The functional role of Gammarus

(Crustacea, Amphipoda): Shredders, predators, or both? Hydrobiologia 485:199–203.

Kelly, D. W., J. R. Muirhead, D. D. Heath, and H. J. Macisaac. 2006. Contrasting patterns in

genetic diversity following multiple invasions of fresh and brackish waters. Molecular

Ecology 15:3641–3653.

Körner, S., and T. Dugdale. 2003. Is roach herbivory preventing re-colonization of submerged

macrophytes in a shallow lake? Hydrobiologia 506–509:497–501.

Korpinen, S., and M. Westerbom. 2009. Microhabitat segregation of the amphipod genus

Gammarus (Crustacea: Amphipoda) in the Northern Baltic Sea. Marine Biology

157:361–370.

Kotta, J., H. Orav-Kotta, K. Herkül, and I. Kotta. 2011. Habitat choice of the invasive

Gammarus tigrinus and the native *Gammarus salinus* indicates weak interspecific

competition. Boreal Environment Research 16:64–72.

Kraufvelin, P., S. Salovius, H. Christie, F. E. Moy, R. Karez, and M. F. Pedersen. 2006.

Eutrophication-induced changes in benthic algae affect the behaviour and fitness of the

marine amphipod Gammarus locusta. Aquatic Botany 84:199–209.

Kufel, L., and I. Kufel. 2002. Chara beds acting as nutrient sinks in shallow lakes—a review.

Aquatic Botany 72:249–260.

Kumar, S., G. Stecher, M. Li, C. Knyaz, and K. Tamura. 2018. MEGA X: Molecular

evolutionary genetics analysis across computing platforms. Molecular Biology and

Evolution 35:1547–1549.

Kuprijanov, I., J. Kotta, V. Lauringson, and K. Herkül. 2015. Trophic interactions between

native and alien palaemonid prawns and an alien gammarid in a brackish water

ecosystem. Proceedings of the Estonian Academy of Sciences 64:518–524.

Leigh, J. W., and D. Bryant. 2015. POPART: Full-feature software for haplotype network

construction. Methods in Ecology and Evolution 6:1110–1116.

Lenz, M., B. A. P. P. da Gama, N. V. Gerner, J. Gobin, F. Gröner, A. Harry, S. R. Jenkins, P.

Kraufvelin, C. Mummelthei, J. Sareyka, E. A. Xavier, and M. Wahl. 2011. Non-native

marine invertebrates are more tolerant towards environmental stress than taxonomically

related native species: results from a globally replicated study. Environmental Research

111:943–52.

Lindner, A. 1972. Soziologisch-ökologische Untersuchungen an der submersen Vegetation in

der Boddenkette südlich des Darß und Zingst. University of Rostock.

LUNG. 2013. Zur Entwicklung und zum Stand der Nährstoffbelastung der Küstengewässer

Mecklenburg-Vorpommerns. Berichte zur Gewässergüte. Güstrow.

Melzer, A. 1999. Aquatic macrophytes as tools for lake management. Hydrobiologia

395/396:181–190.

Nausch, G., A. Bachor, T. Petenati, J. Voss, and M. Von Weber. 2011. Nutrients in the

German coastal waters of the Baltic Sea and adjacent areas (in German). Meeresumwelt

Aktuell Nord- und Ostsee:1–16.

Normant, M., M. Feike, A. Szaniawska, and G. Graf. 2007. Adaptation of Gammarus tigrinus

Sexton, 1939 to new environments-Some metabolic investigations. Thermochimica Acta

458:107–111.

Orav-Kotta, H., J. Kotta, K. Herkül, I. Kotta, and T. Paalme. 2009. Seasonal variability in the

grazing potential of the invasive amphipod Gammarus tigrinus and the native amphipod

Gammarus salinus (Amphipoda: Crustacea) in the northern Baltic Sea. Biological

Invasions 11:597–608.

Östman, Ö., J. Eklöf, B. K. Eriksson, J. Olsson, P. O. Moksnes, and U. Bergström. 2016. Top-

down control as important as nutrient enrichment for eutrophication effects in North

Atlantic coastal ecosystems. Journal of Applied Ecology 53:1138–1147.

Paar, M., M. Berthold, R. Schumann, S. Dahlke, and I. Blindow. 2021. Seasonal Variation in

Biomass and Production of the Macrophytobenthos in two Lagoons in the Southern

Baltic Sea. Frontiers in Earth Science 8:1–16.

Paerl, H. W., and V. J. Paul. 2012. Climate change: Links to global expansion of harmful

cyanobacteria. Water Research 46:1349–1363.

Paiva, F., A. Barco, Y. Chen, A. Mirzajani, F. T. Chan, V. Lauringson, M. Baltazar-Soares,

A. Zhan, S. A. Bailey, J. Javidpour, and E. Briski. 2018. Is salinity an obstacle for

biological invasions? Global Change Biology 24:2708–2720.

Pellan, L., V. Médoc, D. Renault, T. Spataro, and C. Piscart. 2016. Feeding choice and

predation pressure of two invasive gammarids, Gammarus tigrinus and Dikerogammarus

villosus, under increasing temperature. Hydrobiologia 781:43–54.

Piepho, M. 2017. Assessing maximum depth distribution, vegetated area, and production of

submerged macrophytes in shallow, turbid coastal lagoons of the southern Baltic Sea.

Hydrobiologia 794:303–316.

Pitkänen, H., M. Peuraniemi, M. Westerbom, M. Kilpi, and M. Von Numers. 2013. Long-

term changes in distribution and frequency of aquatic vascular plants and charophytes in

an estuary in the Baltic Sea. Annales Botanici Fennici 50:1–54.

Pohl, P., M. K. Ohlhase, S. K. Rautwurst, and K. Laus-Kinnerk Baasch. 1987. An inexpensive

inorganic medium for the mass cultivation of freshwater microalgae. Phytochemistry

26:1657–1659.

Puntila, R. 2016. Trophic Interactions and Impacts of Non-indigenous Species in Baltic Sea

Coastal Ecosystems. University of Helsinki.

R Core Team. 2019. R: A Language and Environment for Statistical Computing. Vienna,

Austria.

Reusch, T. B. H., J. Dierking, H. C. Andersson, E. Bonsdorff, J. Carstensen, M. Casini, M.

Czajkowski, B. Hasler, K. Hinsby, K. Hyytiäinen, K. Johannesson, S. Jomaa, V.

Jormalainen, H. Kuosa, S. Kurland, L. Laikre, B. R. MacKenzie, P. Margonski, F.

Melzner, D. Oesterwind, H. Ojaveer, J. C. Refsgaard, A. Sandström, G. Schwarz, K.

Tonderski, M. Winder, and M. Zandersen. 2018. The Baltic Sea as a time machine for

the future coastal ocean. Science Advances 4.

Sand-Jensen, K., H. H. Bruun, and L. Baastrup-Spohr. 2017. Decade-long time delays in

nutrient and plant species dynamics during eutrophication and re-oligotrophication of

Lake Fure 1900–2015. Journal of Ecology 105:690–700.

Sanudo-Wilhelmy, S. A., A. Tovar-Sanchez, F. Fu, D. G. Capone, E. J. Carpenter, and D. A.

Hutchins. 2004. The impact of surface-adsorbed phosphorus on phytoplankton Redfield

stoichiometry. Nature 432:897–901.

Scheffer, M., S. Rinaldi, and L. R. Mur. 1997. On the dominance of filamentous blue-green

algae in shallow lakes. Ecology 78:272–282.

Schiewer, U. 2007. Darß-Zingst Boddens, Northern Rügener Boddens and Schlei. Pages 35–

86 *in* U. Schiewer, M. M. Caldwell, G. Heldmaier, R. B. Jackson, O. L. Lange, H. A.

Mooney, E.-D. Schulze, and U. Sommer, editors. Ecology of Baltic coastal waters. First

edition. Springer, Berlin, Heidelberg.

Schindler, D. W. 2006. Recent advances in the understanding and management of

eutrophication. Limnology and Oceanography 51:356–363.

Schlungbaum, G., H. Baudler, M. Krech, and B. Kwiatkowski. 2000. Die Darß-Zingster

Bodden - eine Studie. Page (G. Schlungbaum, H. Baudler, M. Krech, and B.

Kwiatkowski, Eds.). Second edition. Landesamt für Umwelt, Naturschutz und Geologie

Mecklenburg-Vorpommern, Güstrow.

Schubert, H., I. Blindow, N. C. Bueno, M. T. Casanova, M. Pełechaty, and A. Pukacz. 2018.

Ecology of charophytes – permanent pioneers and ecosystem engineers. Perspectives in

Phycology 5:61–74.

Schumann, R., H. Baudler, Ä. Glass, K. Dümcke, and U. Karsten. 2006. Long-term

observation on salinity dynamics in a tideless shallow coastal lagoon of the Southern

Baltic Sea coast and their biological relevance. Journal of Marine Systems 60:330–344.

Selig, U., T. Steinhardt, and H. Schubert. 2009. Interannual variability of submerged

vegetation in a brackish coastal lagoon on the southern Baltic Sea. Ekologia Bratislava

28:412–423.

Suikkanen, S., S. Pulina, J. Engström-Öst, M. Lehtiniemi, S. Lehtinen, and A. Brutemark.

2013. Climate Change and Eutrophication Induced Shifts in Northern Summer Plankton

Communities. PLoS ONE 8:1–10.

Torn, K., G. Martin, and R. Munsterhjelm. 2003. Chara tomentosa L. 1753. Pages 131–141 *in*

H. Schubert and I. Blindow, editors. Charophytes of the Baltic Sea. Gantner, Ruggel,

Königstein, Germany.

Torn, K., G. Martin, and T. Paalme. 2006. Seasonal changes in biomass, elongation growth

and primary production rate of Chara tomentosa in the NE Baltic Sea. Annales Botanici

Fennici 43:276–283.

Volkmann, C. 2016. The impact of organic material for macrophytes in coastal waters of the

Baltic Sea. University of Rostock.

Wickham, H. 2016. ggplot2: Elegant Graphics for Data Analysis. Springer-Verlag New York.

Wood, S. N. 2011. Fast stable restricted maximum likelihood and marginal likelihood

estimation of semiparametric generalized linear models. Journal of the Royal Statistical

Society (B) 73:3–36.

Wüstenberg, A., Y. Pörs, and R. Ehwald. 2011. Culturing of stoneworts and submersed

angiosperms with phosphate uptake exclusively from an artificial sediment. Freshwater

Biology 56:1531–1539.

Zettler, M. L. 1995. Erstnachweis von Gammarus tigrinus Sexton, 1939 (Crustacea:

Amphipoda) in der Darss-Zingster-Boddenkette und seine derzeitige Verbreitung an der

deutschen Ostsekueste. Arch. Freunde Naturg. Mecklb. XXXIV:137–141.

Zettler, M. L. 2001. Some malacostracan crustacean assemblages in the southern and western

Baltic Sea. Rostock. Meersebiolog. Beitr. 9:127–143.

Zettler, M. L., and A. Zettler. 2017. Marine and freshwater Amphipoda from the Baltic Sea

and adjacent territories. 2. Auflage. Conchbooks, Hackenheim.

<u>Figures</u>

  Figure 1 Flow scheme of the full factorial experimental design for temperature, and salinity

  with *Chara aspera* and *C. tomentosa*, with and without *Gammarus tigrinus*.

675
676   Figure 2 Phylogeny of *Gammarus tigrinus* COI haplotypes inferred by using the Maximum

677   Likelihood method (Tamura-Nei model +I+Γ) with bootstrap values above the branches

678   (1000 replicates). The clade numbers are identical to Kelly et al. (2006) and the coloured

679   boxes indicate haplotypes of clade N4. Haplotypes N4a and N4e comprised the individuals

680   from the DZLS.

Figure 3 Distribution of *Gammarus tigrinus* populations of clade N4 identified by COI

network analysis. Pie charts indicated haplotype frequencies in European (A) and North

American (B) populations.

Figure 4 Generalized additive model for grazing rates of *Gammarus tigrinus* on *Chara aspera*

(top left), *Chara tomentosa* (top right) and on both Chara species (bottom) in respect to

different salinity and temperature levels.

Figure 5: Seasonal grazing rates of *Gammarus tigrinus* in the Grabow and Bodstedter Bodden

over the years of 2000 to 2018, whereas individual grazing rates (points) were modelled

depending on temperature and salinity conditions at the respective time point in the water

body (left). The loss by grazing per square meter were calculated as the product of the left

side model and the seasonal pattern of *Gammarus tigrinus* (Supplement Figure S3) with a

maximum abundance of 400 individuals per square meter (right).

Figure 6 Charophyte fresh mass in gram per square meter at two sampling locations of the

Darss-Zingst lagoon system along a water depth/distance to reed belt transect. Shallow depth

(30 – 60 cm) and close to reed belt, intermediate depth (60 – 90 cm) and distance to reed belt,

deep depth (90 – 120 cm) and farthest distance to reed belt. Net biomass represents model-

fitted values (generalized additive model) of observed charophyte biomass. Green points

represent observed charophyte biomass at the respective day. Gross biomass represents the

sum of net biomass plus biomass possibly lost to *Gammarus tigrinus* grazing.

Figure 7 Elemental ratio of Carbon to Nitrogen (C:N) and Carbon to Phosphorus (C:P) of

macrophyte biomass (separated by order) at two stations of the Darss-Zingst lagoon system

across all sampling stations (Bodstedter Bodden and Grabow) and sampling depths in 2014.

Please note the different scaling. Points represent measured elemental ratios, lines a local

regression (LOESS – locally estimated scatter smoothing), and the ribbons the 95%

confidence intervals (Wickham 2016).

 Supplement

714 Supplement Table 1: Sample list of specimens used for the phylogenetic tree and the

715 determination of haplotypes. Indicated are the localities, if available the salinity of the

716 sampling site, the identified haplotypes and the accession numbers (F=freshwater sites).

Country	Site	Lat., Lon.	Sal.	haplotype	AccessionNo	Ref.
Denmark	Bornholm, Hovedstaden, Baltic Sea, Svaneke	55.1329, 15.152		N1	MK403734	Baltazar-Soares et al. (2017)
	Bornholm, Hovedstaden, Ostersoen stream, Nexo	55.0569, 15.127		N1, N4a	MK403733, MK403736	Baltazar-Soares et al. (2017)
Estonia	Liu	58.2744, 24.2528	4.7	N1, N4a, N4e	KU845009-KU845015, KU845017-KU845021, KU845023-KU845027	Paiva et al. 2018
	Pärnu	58.3748, 24.4734	4.28	N1, N4a, N4d, N4e	KU845029, KU845030, KU845033, KU845035-KU845038, KU845040-KU845043, KU845046, KU845049, KU845050	Paiva et al. 2018
Finland	Turku	60.4, 22.2	4.4	N4a, N4d, N4e	DQ523177, DQ523178, DQ523183	Kelly et al. 2006
Germany	**Born, DZLS**	**54.379434, 12.524819**	**5.5**	**N4a, N4e**	**MW509071-MW509085**	**this study**
	Dierhagen lagoon	54.2, 12.3	1.5	N4a, N4e	DQ523178, DQ523183	Kelly et al. 2006
	Travemümde	53.9655, 10.8851	10	N1	KU844994-KU845006	Paiva et al. 2018
	North Rhine	51.7328950, 7.1774890	F	N1, N4a, N4c	KT075215, KT075216, KT075218-KT075220	Grabner et al. 2015
Netherlands	Buiten-IJ, Amsterdamm	52.369028, 4.990722	F	N4d	EF570348	Hou et al. 2007
	Lake Gouwzee,	52.4, 5.0	F	N4a, N4c	DQ300227, DQ523183	Kelly et al. 2006
North Ireland	Bann river	54.8, -6.4	F	N4a, N4b, N4e	DQ523178, DQ523181, DQ523183	Kelly et al. 2006

	Lough Neagh,	54.7, -6.5	F	N4a, N4d, N4e	DQ523177, DQ523178, DQ523183	Kelly et al. 2006
Poland	Brody, Oder river	52.0, 15.4	F	N4a, N4d, N4e	DQ523177, DQ523178, DQ523183	Kelly et al. 2006
	Bytom, Oder river	51.7, 15.8	F	N4a, N4d, N4e	DQ523177, DQ523178, DQ523183	Kelly et al. 2006
	Gdansk Bay, Puck	54.723394, 18.416633		N4a, N4e	GQ341858-GQ341861	Costa et al. 2009
	Vistula lagoon	54.3, 19.7	4.5	N1, N4a	DQ300212, DQ523183	Kelly et al. 2006
Canada	New Brunswick	47.037389, -65.163250		N1, N2	FJ581678, FJ581679	Radulovici et al. 2009
	Nova Scotia	45.631983, -61.958414		N2	FJ581680-FJ581683	Radulovici et al. 2009
	Quebec	47.149100, -70.521900		N1, N3	FJ581684-FJ581690	Radulovici et al. 2009
	St. John estuary	45.3, -66.2	4.1	N1	DQ300250, DQ300251	Kelly et al. 2006
	St. Lawrence, d/s Quebec, Cap Brule	46.9, -70.5	6.5	N3	DQ300211	Kelly et al. 2006
USA	Berrys creek, New Hampshire	43.0, -70.7	16	N1	DQ300208-DQ300210	Kelly et al. 2006
	Chesapeake Bay, Virginia	37.45, -76.67		N4b	DQ300219, DQ300220, DQ523180, DQ523181	Kelly et al. 2006
	Delaware estuary, Deemers Beach, Delaware	39.6, -75.5	5	N3, N4a, N4b	DQ523181-DQ523184, DQ523189	Kelly et al. 2006
	Delaware estuary, Delaware	39.57, -75.58		N3, N4a	DQ300221, DQ300222, DQ300225	Kelly et al. 2006
	Elizabeth estuary, Virginia	36.7, -76.2	10.2	N3, N4a, N4c	DQ300224, DQ300226, DQ300227-DQ300231, DQ523183, DQ523186-DQ523188, DQ523190-DQ523195	Kelly et al. 2006

Hudson estuary, New York	40.9, -73.8	7.8	N3	DQ523179	Kelly et al. 2006
Maryland, Rhode River, SERC Education Dock	38.885, -76.542		N4a, N4b	KU905729, KU905737, KU905742, KU905952	unpublished, direct submission
Maryland, Thoroughfare Island, Potomac River, Chesapeake Bay	38.5731, -77.175		N4b	MH235818	Baltazar-Soares et al. (2017)
Neuse River: d/s New Bern, N. Carolina	35.1, -77.0	14.2	Southern Species	DQ300240-DQ300244	Kelly et al. 2006
Pawcatuck estuary, Rhode Island	41.3, -71.8	11.2	N2	DQ300245-DQ300249	Kelly et al. 2006

Figure S1 Growth of *Chara aspera* (A) and *Chara tomentosa* (B) in beakers without

gammarids after 48 hours (n = 8).

Figure S2 Growth-corrected grazing rates of *Gammarus tigrinus* on *Chara aspera* (A) and

Chara tomentosa (B) in beakers after 48 hours (n = 8).

Figure S3 Model of seasonal pattern of *Gammarus tigrinus* calculated with population

densities from Chambers (1977) and modified to have a maximum of 400 individuals per

square meter based on values of Zettler (1995) and of M. Paar (personal communication).

728