Desalination and temperature increase will shift seasonal grazing patterns of

invasive *Gammarus tigrinus* on charophytes

Authors: Berthold[1, 2,*], M.; Porsche[3], C.; Hofmann[3], A., Nowak[3], P.

[1] – Applied Ecology and Phycology, University of Rostock, Rostock, Germany

[2] – Phytoplankton Ecophysiology, Mount Allison University, Sackville, Canada

[3] – Aquatic Ecology, University of Rostock, Rostock, Germany

* – corresponding author: maximilian.berthold@uni-rostock.de

Abstract

Charophytes are a refuge for zooplankton and stabilize sediments, but they are also a food

source for various animal species (water birds, fishes, invertebrates). Especially the

introduction of new species, like *Gammarus tigrinus*, into the Baltic Sea led to yet not

understood changes in the food web. Furthermore, future projections point to increased water

temperatures at lowered salinity levels affecting species capacity to acclimatize to changing

abiotic factors. In this study we investigated the influence of temperature and salinity on the

grazing pressure of *Gammarus tigrinus* on two charophyte species: *Chara aspera* and *Chara*

tomentosa. The grazing experiments were conducted in a full factorial design with the factors

salinity ($3 – 13$ g kg^{-1}), temperature ($5 – 30$ °C), and charophyte species. Grazing rates were

determined as mass deviation within 48 hours considering biomass changes in the presence

and absence of gammarids. Grazing rate were further used to calculate charophyte losses in

two coastal lagoons with different salinity concentrations for recent and future time periods.

The potential grazing peak of about 24 °C is not yet reached in these coastal waters but may

be reached in the near future as shown by our future projection results. However, the

temperature increase, and desalination will cause a shift in seasonal individual grazing

patterns from summer to spring and autumn. Desalination and temperature increase can lead

to a shift in optimal habitats for *G. tigrinus* in the future.

Keywords: charophytes, grazing, *Gammarus tigrinus*, temperature, salinity, climate change

Introduction

Global climate change imposes several interacting effects on aquatic organisms. Besides the

increase of water temperature, there may be regionally altered hydrological flow regimes that

alter timing and amount of freshwater input into coastal waters. Native species get stressed by

this development and ecosystems may be more susceptible to invasive species colonization.

The Baltic Sea is such a system prone to eutrophication and global climate change, due to

already increased sea surface temperatures, compared to other oceans worldwide (Reusch et

al. 2018). The Baltic Sea is subject to large phytoplankton blooms since decades, with the

riparian states working to reduce the impact of nutrient-run-off to the sea (Janssen et al. 2004,

HELCOM 2018). Similar, the coastal water bodies of the Baltic Sea were impacted by

eutrophication and shifts on primary producer dominance, i.e. from submerged macrophytes

to phytoplankton (Schiewer 2007).

Charophytes are widely distributed in fresh and brackish water ecosystems (e.g. Schubert and

Blindow 2003) and act as keystone organisms in shallow water ecosystems of the Baltic Sea

because of their high biomasses (Lindner 1972). During the last decades, charophytes

received attention as indicator species for oligotrophic water bodies, but also as pioneer

species in post-eutrophied water bodies (e.g. Melzer 1999, Schubert and Blindow 2003).

These traits make charophytes interesting foundation species as habitats are colonized

regardless of trophic status (Schubert et al. 2018). Algae of the Characeaen family are

currently listed as 'endangered species' in several European areas (Stewart and Church 1992,

Blazencic et al. 2005, Gärdenfors 2010, Auderset Joye and Schwarzer 2012) and they are also

under pressure in the Baltic Sea. Comparisons of historical and recent distributions in the

northern Baltic found that *C. aspera* and *C. tomentosa* were found 8 and 86% times less in the

most recent transect, respectively (Pitkänen et al. 2013). Besides eutrophication related issues,

salinity and/or temperature changes may influence successful recolonization in the future.

Salinity changes in brackish water are such a factor, where increased salinity prevents the

growth of macrophyte species (Lindner 1972). It was assumed that charophytes have several

turbidity-reducing traits, like through for example lowering of resuspension of particulate

matter, and direct (dissolved nutrients) and indirect (refuge for zooplankton) competition with

phytoplankton (Kufel and Kufel 2002). Therefore, recolonization with those species could

create feedback mechanism, favoring a macrophyte-dominated stable state.

However, restoration measures may be hampered by global climate change and the effects of

desalination and increasing water temperature. Furthermore, there are hypotheses that not

only abiotic parameters, but also top-down control (i.e. grazing) may affect the restoration of

the pristine trophic state and macrophyte recolonization (Körner and Dugdale 2003, Östman

et al. 2016). Besides grazing from native species (Körner and Dugdale 2003), there is also an

increasing grazing pressure of invasive species introduced through, e.g. ballast water of ships.

One of these species is the gammarid *Gammarus tigrinus* (Sexton 1939), a successful new

species introduced from North America to European waters (Berezina 2007). This gammarid

was described to colonize shallow waters, especially within reed, and soft bottom systems of

the Baltic Sea (Daunys and Zettler 2006, Kotta et al. 2011). Amid global climate change,

invasive species can alter known food web interactions and may counteract recolonization of

former native primary producers (Puntila 2016). However, even these species are subject to

global climate change and changes in the abiotic environment may support these invasive

species even further (Kelley 2014).

The question evolves, if recolonization of macrophytes is not only hampered by abiotic

factors, like temperature and salinity, but additionally by grazing pressure of (newly

introduced) grazers. This grazing pressure likely changes throughout the year and may result

in high-stress situations, where pressures of salinity and temperature amplify with grazing.

Gammarus tigrinus and charophytes occur within the same depth distribution, and grazing

can potentially happen. It was hypothesized that there is either a combination of factors

(abiotic and biotic) that amplify mass loss on charophytes, or that some abiotic factors quench

each other, e.g. lower salinity amplifies/buffers effects of higher temperatures. We tested the

impact of salinity and temperature as well as their interactions on biomass change of

charophytes in the presence and absence of grazers. These results also allow to discuss future

developments within coastal water bodies at e.g. rising water temperatures, or desalination.

Shallow coastal water bodies are likely to be more affected by such events due to their

relatively shallow water column and close coupling with the adjacent land.

We chose the Darss-Zingst lagoon system (DZLS) as model ecosystem for the southern Baltic

Sea, due to its salinity and trophic gradient and decade-long monitoring record (see Schiewer

2007). Even though nutrient concentrations were reduced to a comparable level between

today and the 1930ies, no dense macrophyte cover was able to form in DZLS (Gessner 1957,

Berthold et al. 2018a, Paar et al. 2021). Indeed, the maximum depth distribution of

charophytes has not increased significantly in the DZLS during the last years, and

charophytes are found only in very shallow waters, where light is not limiting (Piepho 2017).

To account for this development, we combined the laboratory determined grazing potentials

with field-based growth patterns of submerged macrophytes of the DZLS, and projected them

with future salinity and temperature developments in these waters.

Material and Methods

Experimental design

We chose a full factorial design with three factors (Figure S1), for which one factor relates to

species and includes two levels (*Chara aspera* and *C. tomentosa*), and the other two factors

each with four levels are water temperature (9, 15, 24, and 30 °C), and salinity (3, 5, 7, 13 g

kg^{-1}). These salinity levels represent the salinity gradient from the Western belt Sea to the

Gulf of Finland, covering most of the Baltic Sea area (HELCOM 2010). The water

temperature levels represent the blooming period of shallow coastal waters (+1 °C in winter to

+22 °C in summer, Schiewer 2007), with 30 °C as prospective endpoint of possible

charophyte growth.

The DZLS is formed by four consecutive shallow lagoons, and shows a salinity and trophic

gradient from the main river inflows Recknitz and Barthe (inner and middle part, salinity of 1

– 6 g kg^{-1}, eutrophic) to the open Baltic Sea (salinity of 10 – 12 g kg^{-1}, mesotrophic,

Schumann et al. 2006). Salinity varies from 4 – 6 g kg^{-1} at the respective sampling locations

in the Bodstedter Bodden (54°22'30,5" N, 12°34'14,9" E) and around 6 – 8 g kg^{-1} in the

Grabow (54°22'2,3" N, 12°48'27" E) (Schumann et al. 2006). During the experiment, two

charophyte species were used, *Chara aspera* and *Chara tomentosa*. Both species are common

in the Baltic Sea, with *C. aspera* covering the widest salinity range among charophytes

(Blindow 2000). Along the west coast of Sweden, *C. aspera* is regularly found at sites with

salinities up to 15 g kg^{-1} (temporarily up to 20 g kg^{-1}), but it is also common in freshwater

(Hasslow 1931, Blindow 2000). *C. tomentosa* is among the largest charophyte species and can

be found in freshwaters all over the world. Only in the Baltic Sea, *C. tomentosa* also occurs in

brackish water, with a salinity range between 0 – 7.5 g kg^{-1} (Björkman 1947, Torn et al.

2003). Gammarids were derived from the Aquaculture section in Born of the State Research

Centre for Agriculture and Fishery Mecklenburg-Vorpommern. This research facility is

located at the DZLS and uses brackish water from the lagoon with a salinity of 4 – 6 g kg^{-1}.

Gammarids were identified by morphology and genetic analyses (see supplement).

Charophytes were extracted as intact plants from the DZLS. Charophytes were then cultured

in beakers (vol. 500 ml) following the culture design described by Wüstenberg et al. (2011).

Each beaker contained sediment containers (vol. 64 ml) with acid-rinsed, phosphorus-

enriched sediment (6 g P kg^{-1} sediment). Mineral growth media was prepared to culture

charophytes (Pohl et al. 1987, Wüstenberg et al. 2011), but was adjusted to the respective

salinities by adding marine salt (Instant Ocean, Aquarium Systems, Sarrebourg). Salinity was

checked with a salinometer (Multiline P4, WTW). Each of the 16 combinations were

replicated eight times with gammarids (= grazing treatment) and in absence of gammarids (=

control) (Fig. S1). Gammarids were pre-acclimatized to experimental conditions by culturing

them at the respective temperature and salinity combinations. Ten to 20 individuals were

simultaneously cultured per temperature-salinity combination, aerating all cultures.

Charophytes were added as food supplement, as well as resting surface for the gammarids.

Charophytes were wet weighed and set into beakers filled with sediment and growth media of

the respective salinity. Controls consisted of one charophyte, whereas grazing treatments

consisted of one charophyte and five gammarids in one beaker. Control and grazing

treatments were exposed to light intensities between 80 and 120 µmol photons m^{-2} s^{-1} using

fluorescent tubes (Phillips, TLD 36W/950). Each replicate run lasted for about 48 hours.

Replicates were run in blocks over a period of four weeks per *Chara* species. Two out of the

eight replicates per salinity-temperature combinations were conducted simultaneously to

spread and randomize replicate runs over the four week period. According to previous studies

on growth of charophyte (Nowak et al. 2017) we did not expect significant growth responses

of *C. aspera* and *C. tomentosa* over the course of two days. However, it was expected to find

significant mass deviations when gammarids were present.

Charophytes were wet weighed after each replicate run to calculate mass differences. Dead

gammarids were counted at the end of each replicate run and replaced for the next replicate

run. If possible, the same gammarids were used throughout all replicate runs. Equations 1 – 3

were used to extract grazing rates of gammarids from growth rates of charophytes during the

experiments. Grazing rates were treated with an error-propagation to account for experimental

variability.

Equation 1 subtract the mean effect of biomass change of the control (C) from the treatment

which also includes grazing (T):

$$G_i = T_i - \bar{C} \ \{i = 1..n$$

153   Equation 2 standardizes the grazing effect from prior formula (**z**-score):

$$Gz_i = \frac{(G_i - \bar{G})}{\sigma_G} \ \{i: 1..n$$

154   Equation 3 calculates the mean of pure grazing and include its error caused by the experiment:

$$Gf_i = \bar{G} + Gz_i \cdot \sqrt{|(\sigma_T^2 - \sigma_C^2)|} \ \{i: 1..n$$

T = treatment, C = control, G= detrended grazing, G_z = standardized grazing, G_f = grazing

with included experimental error, n = replicates

<u>Field sampling and analyses</u>

In addition, we used elemental and biomass composition of submerged macrophytes from the

DZLS, to compare the experimental derived grazing rates to the actual ecosystem. Field

sampling was conducted in the Bodstedter Bodden and the Grabow in 2014, as part of the

BACOSA project. Interestingly, the Bodstedter Bodden has on average a lower salinity of 2 g

kg^{-1}, then the Grabow, coinciding with future desalination projections (Neumann and

Friedland 2011). This difference may be used as model validation, if grazing rates of the

present Bodstedter Bodden represent future grazing rates in the Grabow. Submerged

macrophytes were sampled throughout the year, first in 54x54 cm plots, and in 25×25 cm

plots later during the year at nine different depths. Here, three sampling locations were pooled

into three depth zones, according to water depth and proximity to the reed belt. Water depths

for both stations varied with water levels in the respective lagoons and were shallow (30 – 60

cm), intermediate (60 – 90 cm), and deep (90 – 120 cm). Sampled macrophytes were

separated on the genus level. Whole plants, including roots, were wet-weighed and then dried

at 60 °C for 24 h. Afterwards, dry mass (DM) was re-weighed again to calculate water

content. Dried plant material was ground with a ball mill. The powder was treated with 1M

HCl to remove inorganic carbon residues and then weighed in tin-container (1 × 0.5 cm).

These samples were analyzed for their carbon and nitrogen content using an elemental

analyzer (varioEL III). The analyzer was calibrated using acetanilide (~5 mg per sample).

Aliquots of dried plant material were weighed in crucibles and combusted at 550 °C for 4 h.

The ash was re-weighed to calculate ash-free dry mass. Ashes were used to determine total

phosphorus content (mg P g^{-1} dry mass), using an acid persulfate extraction (see Berthold et

al. 2015).

<u>Data analyses</u>

We further used the experimental grazing data to calculate grazing rates based on salinity and

water temperature at two sites of the DZLS. The data set failed to show homogeneity of the

residuals, and was therefore Box-Cox power transformed (*bcnPower* function, R package

"*car*", Fox and Weisberg 2019). Finally, general linear models (GLM, *glm* function, R

package "stats", R Core Team 2019) and generalized additive models (GAM, *gam* function, R

package *"mcgv",* Wood 2011), both with Gaussian distributional assumptions, were

compared by the Akaike Information Criterion (AIC, R package "stats", R Core Team 2019)

and the Bayesian Information Criterion (BIC, R package *"stats",* R Core Team 2019),

choosing the model with low AIC and BIC values. We chose the GAM (with knots, k = 16) as

the best model to reflect the experimental data (see Results). To our knowledge, recent studies

on seasonal population dynamics of *G. tigrinus* are currently missing in our study system, as

elsewhere in the Baltic Sea. However, recent field studies found 5 to 2200 Ind m^{-2}

(gammarids on genus level) in the DZLS and the adjacent Western Rügen lagoon system.

Largest, but fewest animals were found early in the year (22 – 28 mg Ind^{-1}), with decreasing

mass, but increasing abundance in summer (3.5 mg Ind^{-1}) (M. Paar, personal communication).

These patterns point to the same seasonal development as described in Chambers (1977). If

we assume that *G. tigrinus* constitutes 20% of the local gammarid population in the DZLS

(Zettler 2001), we get a population size of up to 450 Ind m^{-2}. These numbers are in agreement

with first described abundances of *G. tigrinus* of 300 – 500 Ind m^{-2} in the phytal and reed belt

of the DZLS (Zettler 1995), or 30 – 110 Ind m^{-2} in the Neva estuary (Baltic Sea, Berezina

2007).

We used again a GAM model to calculate the seasonal population development of *Gammarus*

tigrinus, normalized the predicted seasonal data to have a maximum of 1. Then, we used this

normalized prediction to calculate population densities with a maximum of 400 Ind m^{-2}

(Zettler 1995, M. Paar personal communication, Fig. S6). These population numbers were

combined with the individual grazing rates of the temperature-salinity model with field data

from the two sites of the DZLS (Equation 4) to calculate seasonal grazing rates. We used a

fixed grazing preference of 20% macrophyte share in gammarid diets, that means grazing

rates were multiplied by a factor of 0.2 (Pellan et al. 2016). Furthermore, we used the GAM to

predict the charophyte development as function of sampling date (k = 8) at two sites of the

DZLS. We calculated daily biomass increase and subsequently added gammarid grazing rates

to determine the potential gross biomass production per site (see Results and Discussion

below).

Equation 4 calculates the seasonal grazing rates per square meter and day for specific sites:

$$Graz_t = G_{GAM_t}\{T_t; S_t\} \cdot \delta_t \cdot 0.2 \quad \{t: 1..n$$

t = day of the year, $Graz_t$ = grazing at day t (mg·m^{-2}·d^{-1}), G_{GAMt} = modelled grazing rate of one

individual at time t with given temperature (T) and salinity (S) from specific site at time t, δ_t =

population density at time t (individuals·m^{-2}), 0.2 = reflects the 20 % fixed grazing preference,

n = length of the year in days

Results

<u>Growth and survival rate</u>

Starting biomasses of *C. aspera* in control and treatment beakers ranged from 50 to up to 300

mg, with the most macrophytes being within the range of 100 – 175 mg (70 %). After two

days, biomass distribution of *C. aspera* in control beakers stayed relatively the same, but

smaller and larger biomass proportions increased. Biomass of *C. aspera* decreased with

increasing temperature at all salinities, but strongest at salinities > 7 g kg^{-1}. Contrary, biomass

stayed almost constant at lower temperature, even at higher salinities (Supplement Table S2

and Figure S4).

Starting biomass of *C. tomentosa* in control beakers ranged from 75 to up to 600 mg, and 40

% of all macrophytes were within the range of 100 – 125 mg at the beginning. Contrary to *C.*

aspera, biomass of *C. tomentosa* in treatment beakers was close to a normal distribution with

50 % of all macrophytes being within the range 175 – 225 mg at the beginning. The biomass

distribution of *C. tomentosa* in control beakers stayed almost the same, but with a trend to

smaller biomasses, indicating minor biomass losses even without grazing. Biomass of *C.*

tomentosa stayed very constant at almost every temperature-salinity combination. There was a

possible growth at temperatures of 15 to 24 °C at salinities of 7 g kg^{-1} (Supplement Figure

S4). If results of both charophyte control beakers were combined, the area with stable biomass

over two days expands from 10 – 25 °C and salinities of 7 – 13 g kg^{-1}. In total, these results

indicate an overall stable experimental approach, as significant growth was not expected

within two days.

Two thirds of the time (65 %) all gammarids survived at all temperature-salinity-macrophyte

combinations in treatment beakers. Additionally, 30 % of the time, only one gammarid died

during incubation. These results indicate overall stable culture conditions and the capability of

these gammarids to survive a wide range of temperature-salinity gradients.

Gammarid grazing on charophytes

Biomass in treatment beakers decreased in all combinations and beakers. Differences to

control beakers were always significant, regardless of charophyte species, or treatment (see

Supplement table S2). Grazing rates ranged between 1 to 10 mg FM Ind^{-1} gammarid d^{-1} on C.

$aspera$, already accounting for biomass changes in control beakers. Grazing rates peaked at

24 °C and tended to be higher with increasing salinities. Grazing ranged between 1 to 5 mg

FM Ind^{-1} gammarid d^{-1} on $C.\ tomentosa$. Grazing rates peaked again at 24 °C for all salinity

levels but with a tendency to slightly lower salinity compared to grazing rates on $C.\ aspera$.

Interestingly, highest grazing rates per charophyte species fell into temperature ranges where

charophytes already showed highest loss rates in control beakers. In general, grazing rates on

$C.\ aspera$ were up to two times higher compared to $C.\ tomentosa$. Pooled grazing rates of

both charophyte species were on average up to 5.5 mg Ind^{-1} d^{-1}.

Field data results and model

Seasonal daily grazing rates within the ecosystem would range from 0.1 up to 5 mg FM Ind^{-1}

d^{-1}, when applying the grazing model on field observations of temperature and salinity of

Grabow and Bodstedter Bodden (Figure 5). Grazing rates would peak during summer and be

lowest at winter at the two sampling stations of the DZLS (see Figure 5). $Gammarus\ tigrinus$

was originally collected in April/May, where grazing rates are not at its peak. Nonetheless,

gammarids grazed twice as much at 24 °C in our treatments, as would be expected from their

initial collection time in April and May. These findings indicate acclimation processes.

Interestingly, the modelled grazing rates differed significantly between both stations ($p <$

0.001, paired Mann-Whitney U-Test), even though the difference in salinity are only 2 g kg^{-1}.

Charophyte biomass was highest in the shallow areas, close to the reed belt at both sampling

sites (Figure 6). Biomass in the shallow zone peaked in July in the less eutrophic Grabow, and

in October in the highly eutrophic Bodstedter Bodden. Contrary, charophyte biomass was

higher in the Bodstedter Bodden at intermediate distances/water depths compared to the

Grabow. However, Grabow biomass peaked during summer, whereas biomass in the

Bodstedter Bodden became depressed during that time. Charophytes were only found in very

little biomasses in the deeper water areas at both sampling locations, indicating a possible

light limitation from shallow to deep.

Charophyte biomass would have been up to 15 – 50% higher at shallow and intermediate

depths, if macrophyte growth is corrected for seasonal gammarid grazing. Additionally,

gammarid grazing would follow charophyte biomass peaks at the respective stations. The

yearly potential grazing rate for *G. tigrinus* on charophytes ranges from 37.2 to 42.5 g FM m^{-}

^{2}, considering regional salinity and temperature patterns within the DZLS.

<u>Future projections on desalination and temperature increase in the period of 2050 – 2100</u>

The projected grazing rates vary significantly from today's grazing rates ($p < 0.001$, paired

Mann-Whitney U-Test), when considering future desalination and temperature increase in this

region (van der Linden and Mitchell n.d., Neumann and Friedland 2011). Individual grazing

rates would increase in spring and autumn at both stations, but grazing rates in the Grabow

being at least twice as high as in the Bodstedter Bodden. Furthermore, individual grazing rates

would probably drop during summer, as water temperatures will fall out of optimum ranges

for *G. tigrinus*. These differences in future grazing rates are caused by earlier warmer

temperatures, and lower salinities, increasing future individual grazing pressure in the

Grabow. However, this double peak is not apparent, when the seasonal population structure is

included. Areal grazing rates would not be different during spring, and even lower during

summer. However, grazing rates would strongly increase during autumn at both stations. This

peak is caused by the late population peak of gammarids, and the increased temperature in

future projections. This grazing pattern would likely affect charophytes in the turbid

Bodstedter Bodden, as they showed highest biomass during autumn.

Discussion

Charophyte growth and gammarid grazing rates in our experiment and in the field depend on

changes in the abiotic environment. However, grazing rates are simultaneously influenced by

available food sources for gammarids, occurrence of competing species and predators, or

ultimately habitat choice and niche occupation.

In this study, desalination and temperature increase experiments lasted only for two days.

Charophytes of control beakers showed only little, if any biomass increase, and cannot be

used for long-term growth projections. There are complex interactions between abiotic factors

(e.g. salinity, temperature, light climate) that influence charophyte growth and these effects

depend on species-specific characteristics, local environmental conditions, and probably

locally adapted sub-populations (Blindow et al. 2009, Auderset Joye and Rey-Boissezon

2015, Rojo et al. 2017). Distinct *C. aspera* populations can show different growth optima

depending on the habitat (freshwater and brackish), which is translated by changing

photophysiological characteristics at varying salinities (Blindow et al. 2003, Blindow and

Schütte 2007). Likewise, water temperature shapes charophyte populations, which may result

in local biomass decreases of *C. aspera*, *C. tomentosa* and *C. vulgaris* if temperature increases

more than 2 °C (Auderset Joye and Rey-Boissezon 2015, Choudhury et al. 2019). Even our

short-term experiments may confirm these results, as both charophyte species lost most

biomass at temperatures above average summer habitat temperatures (> 24 °C). However,

interactions of abiotic factors do not necessarily induce synergistic effects in charophytes

(Rojo et al. 2017, Puche et al. 2018). Our short-term growth controls showed synergistic

effects on charophyte growth rates for different temperature and salinity conditions (Fig S4).

Further studies are needed to clarify long term charophyte changes in the face of increasing

water temperatures and decreasing salinities, especially among locally adapted sub-

populations.

Several physiological reactions, like survivability or oxygen consumption of *G. tigrinus* on

salinity and temperature changes have already been described in the literature (Dorgelo 1973,

1974, Koop and Grieshaber 2000). *Gammarus tigrinus* survives within a broad salinity-

temperature spectrum, where only full marine or freshwater conditions show highest

mortalities. Furthermore, potential synergetic were observed with higher temperatures

buffering higher salinities, with survival rates 10-times lower at 5 °C, compared to 25 °C

(Dorgelo 1974).

Grazing rates in our study suggest similar findings, as lowest temperatures strongly reduced

grazing rates, regardless of salinity. There was also a tendency to higher grazing rates, at

higher salinity/temperature combinations, compared to high salinity/low temperatures,

indicating a grazing optima at higher temperatures. Ion regulation in *G. tigrinus* happens fast,

and can regulate sudden high ionic stress to a variety of ions (Na, Cl, K) without

compensation losses. However, hypoosmotic, that means low salinity conditions, cause

increased NO_3 inflow into *G. tigrinus* hemolymph, probably causing decreased survivability

(Koop and Grieshaber 2000). The growth media for charophytes in our experiment has

elevated NO_3 concentrations (3 mM KNO_3, Pohl et al. 1987). This elevated NO_3

concentrations may have caused lower grazing rates at lower salinities in our study.

Besides abiotic factors, population structure influenced the grazing rates of gammarids.

Populations of *G. tigrinus* were described to turn-over at least three times per year, with a

peak of large animals (6 – 9 mm) in spring, and a concomitant shift to smaller animals in late

summer (2 – 4 mm) (Chambers 1977). We used animals in the size range of 6 – 12 mm, i.e.

large adults for all of our experiments. This size class is only dominant by up to 60% until

May, and is almost completely replaced by smaller animals from June to September

(Chambers 1977). This pre-selection had likely an effect on the described grazing rates, as

metabolic rates of *G. tigrinus* increase with lower weight in form of a power function

(Normant et al. 2007). Gammarids in our study weighed from 10 – 90 mg wet mass Ind^{-1}

(data not shown), whereas the gammarids of Normant et al. (2007) were caught in August,

and weighed only 5 – 25 mg wet weight Ind^{-1}. Animals of our study should therefore showed

lower specific metabolic rates of 0.25 – 1.5 mW g^{-1} wet mass, compared to 1 – 3.5 mW g^{-1}

wet mass in smaller gammarids (Normant et al. 2007). This reduced metabolic rate had likely

an impact on the grazing rates we found in this study, as grazing for metabolic upkeep is

lower in large adult animals.

The results of this study indicate that the individual grazing rates of *G. tigrinus* will change

over the next 50 to 100 years. During that time, the optimal properties of temperature and

salinity for *G. tigrinus* will increase in spring and autumn months and decrease during

summer months.

Springtime is a crucial period for charophyte sprouting, as the active growth of for example

C. tomentosa takes place during a relatively short period at the beginning of summer (Torn et

al. 2006). Furthermore, young sprouted charophytes show a higher elemental content per

biomass, than for example tracheophytes (Volkmann 2016, this study), making them a

preferred grazing opportunity (Kraufvelin et al. 2006). *Gammarus locusta* follows a

preference for high value plant material and selectively feeds on more nutrient-rich

macrophytes, especially periphyton, brown and green algae (Kraufvelin et al. 2006). If *G.*

tigrinus shows similar grazing preferences, periphyton from tracheophytes would be the first

choice followed by charophytes in the DZLS. Periphyton (Sanudo-Wilhelmy et al. 2004) and

possibly charophytes (Kufel and Kufel 2002) can take up large amounts of dissolved nutrient

that occur frequently through diffuse run-off (Berthold et al. 2018b). Tracheophytes in the

DZLS and adjacent lagoons can show high colonization rates by epiphytes (Paar et al. 2021),

offering an additional plant food source to gammarids. Nonetheless, charophytes in the DZSL

showed higher C:N and N:P ratios, then tracheophytes, making them more likely to gammarid

grazing as high-value food (see Fig S7). Furthermore, this pre-selection of high-value food

may explain the differences in grazing rates on *C. tomentosa* vs. *C. aspera* found in this study.

C. tomentosa can show higher C:N ratios (44:1) than *C. aspera* (27:1), making it maybe a less

favorable food source. Furthermore, *C. aspera* showed in our study signs of early decay at

higher temperatures, contrary to *C. tomentosa* (see trends in control beakers, Figure S4). Such

beginning decay can weaken *C. aspera* and make it more susceptible to grazing (Buchsbaum

et al. 1991), explaining the deviation of grazing rates between both charophytes along the

temperature gradient found in this study.

The drop of grazing rates during future summer conditions is probably caused by below-

optimum water temperatures in combination with lowered salinities. The impact on future

summer impact grazing is more pronounced in Bodstedter Bodden, then in the Grabow. This

difference is probably caused by even lower salinity concentrations in the Bodstedter Bodden,

indicating that future desalination can lead to shifts of preferred habitats.

Food preferences can change throughout the year depending on gammarid life stage and

temperature (Felten et al. 2008, Pellan et al. 2016). Plant biomass becomes increasingly

important, when temperatures increase and can reach up to 20 – 30% of total food consumed

(Pellan et al. 2016). Grazing on charophytes can happen earlier in the year, and lasts longer

during autumn. This effect does not come into effect in our future projection model, as the

current modelled population densities of gammarids are lowest in spring. Nonetheless,

prolonged grazing may hamper charophytes sprouting in spring (*C. aspera*) and lower

biomass of overwintering species (*C. tomentosa*) (Torn et al. 2006, Blindow et al. 2016).

Functional feeding in gammarids relies furthermore on population composition and food

availability (Kelly et al. 2002). Other food sources like detritus, plant litter, or animal matter,

like chironomides, are usually important food sources during winter time (Felten et al. 2008,

Pellan et al. 2016). Sediment organic content can be as high as 20% in sediments of the DZLS

(Berthold et al. 2018c). Sediment detritus was not an alternative source in our experiments, as

we used only inorganic sand. Plant litter occurs abundantly within the reed belt during winter

time in the DZLS (Karstens et al. 2016). Our experimental results show therefore only a

potential grazing rate, as we did not represent the variety and abundance of alternative food

sources from within the ecosystem.

Furthermore, grazing rates depend on niche occupation and competition with native and other

invasive species. *Gammarus tigrinus* can probably outcompete the native *G. salinus*, if both

depend on *Pylaiella littoralis* in the northern Baltic Sea (Orav-Kotta et al. 2009). Both

gammarids had the potential to exceed daily macrophyte production during summer (Orav-

Kotta et al. 2009).

G. tigrinus may be more tolerant against heat waves and sub-oxic conditions, than native

species (Lenz et al. 2011). These broader temperature acclimation abilities may help to

permanently introduce *G. tigrinus* within the Baltic Sea, as sea surface temperature are

already higher there, then in other ocean parts (Reusch et al. 2018). Furthermore, endurance

of sub-oxic conditions is an advantage in eutrophic coastal waters of the Baltic, as redox

conditions within the reed belt can change fast (Karstens et al. 2015), and an increase of

nutrients is challenging submerged macrophytes in the coastal water bodies of the southern

Baltic Sea (Paar et al. 2021), stressing habitats of native gammarids. Contrary, an *in silico*

study assumed that *G. tigrinus* has actually a narrower niche than native gammarids in the

northern Baltic Sea (Herkül et al. 2016). These results are somewhat contradictory to other

studies in this area (Korpinen and Westerbom 2009, Kotta et al. 2011, Lenz et al. 2011), and

may have not considered the impact of haplotype-diversity (Baltazar-Soares et al. 2017), that

means locally adapted populations of *G. tigrinus*. Different studies revealed high genetic

variability of *G. tigrinus* in the Baltic Sea (Supplement Table 1, Figure S2, Kelly et al. 2006

and Paiva et al. 2018) indicated that the invasive species may have different tolerance and

different limits to salinities due spatially varying selection among populations.

The patchy macrophyte stands found in the DZLS, and in other eutrophic water bodies, may

affect the occurrence of *G. tigrinus*, and therefore the grazing rates, especially in intermediate

and deeper waters. Palaemonid prawns put a higher grazing pressure on *G. tigrinus* in

unvegetated mesocosms, then in vegetated ones (Kuprijanov et al. 2015). It is likely that *G.*

tigrinus stays close to the reed belt, as described for this lagoon and other coastal waters

(Zettler 1995, 2001). This spatial distribution would further hamper charophyte

recolonization, as the shallow areas in the DZLS are the only spots, where light availability is

high enough to support charophyte biomass (see Figure S6, Piepho 2017).

Acknowledgements

The authors thank the State Agency for Environment, Nature Conservation and Geology

Mecklenburg-Vorpommern for providing monitoring data and the Aquaculture section in

Born of the State Research Centre for Agriculture and Fishery Mecklenburg-Vorpommern for

providing gammarids. We also want to thank Dr. Ralf Bastrop for providing the respective

primers, and Dr. Martin Paar for providing us with recent data on gammarid distribution in

the southern Baltic Sea.

References

Auderset Joye, D., and A. Rey-Boissezon. 2015. Will charophyte species increase or decrease

their distribution in a changing climate? Aquatic Botany 120:73–83.

Auderset Joye, D., and A. Schwarzer. 2012. Red List Characeae. Threatened Species in

Switzerland. Status 2010.

Baltazar-Soares, M., F. Paiva, Y. Chen, A. Zhan, and E. Briski. 2017. Diversity and

distribution of genetic variation in gammarids: Comparing patterns between invasive and

non-invasive species. Ecology and Evolution 7:7687–7698.

Bandelt, H. J., P. Forster, and A. Rohl. 1999. Median-joining networks for inferring

intraspecific phylogenies. Molecular Biology and Evolution 16:37–48.

Berezina, N. A. 2007. Expansion of the North American amphipod Gammarus tigrinus

Sexton, 1939 to the Neva Estuary (easternmost Baltic Sea). Oceanologia 49:129–135.

Berthold, M., U. Karsten, M. von Weber, A. Bachor, and R. Schumann. 2018a. Phytoplankton

can bypass nutrient reductions in eutrophic coastal water bodies. Ambio 47 (1):146–158.

Berthold, M., S. Karstens, U. Buczko, and R. Schumann. 2018b. Potential export of soluble

reactive phosphorus from a coastal wetland in a cold-temperate lagoon system: Buffer

capacities of macrophytes and impact on phytoplankton. Science of the Total

Environment 616–617:46–54.

Berthold, M., D. Zimmer, V. Reiff, and R. Schumann. 2018c. Phosphorus Contents Re-visited

After 40 Years in Muddy and Sandy Sediments of a Temperate Lagoon System.

Frontiers in Marine Science 5:1–14.

Berthold, M., D. Zimmer, and R. Schumann. 2015. A simplified method for total phosphorus

digestion with potassium persulphate at sub-boiling temperatures in different

environmental samples. Rostocker Meeresbiologische Beiträge 25:7–25.

Björkman, S. O. 1947. On the distribution of Chara tomentosa L. round the Baltic and some

remarks on its specific epithet. Botaniska Notiser 1947:157–170.

Blazencic, J., B. Stevanovic, Z. Blazencic, and V. Stevanovic. 2005. Red Data List of

Charophytes in the Balkans. Biodiversity & Conservation 15:3445–3457.

Blindow, I. 2000. Distribution of charophytes along the Swedish coastin relation to salinity

and eutrophication. International Review of Hydrobiology 85:707–717.

Blindow, I., S. Dahlke, A. Dewart, S. Flügge, M. Hendreschke, A. Kerkow, and J. Meyer.

2016. Long-term and interannual changes and viable diaspore reservoir of submerged

macrophytes in a shallow brackish water bay of the southern Baltic Sea - influence of

eutrophication and climate. Hydrobiologia:1–16.

Blindow, I., J. Dietrich, N. Möllmann, and H. Schubert. 2003. Growth, photosynthesis and

fertility of Chara aspera under different light and salinity conditions. Aquatic Botany

76:213–234.

Blindow, I., N. Möllmann, M. G. Boegle, and M. Schütte. 2009. Reproductive isolation in

Chara aspera populations. Aquatic Botany 91:224–230.

Blindow, I., and M. Schütte. 2007. Elongation and mat formation of Chara aspera under

different light and salinity conditions. Hydrobiologia 584:69–76.

Buchsbaum, R., I. Valiela, T. Swain, M. Dzierzeski, and S. Allen. 1991. Available and

refractory nitrogen in detritus of coastal vascular plants and macroalgae. Marine Ecology

Progress Series 72:131–143.

Chambers, M. R. 1977. The population ecology of Gammarus tigrinus (sexton) in the reed

beds of the Tjeukemeer. Hydrobiologia 53:155–164.

Choudhury, M. I., P. Urrutia-Cordero, H. Zhang, M. K. Ekvall, L. R. Medeiros, and L.-A.

Hansson. 2019. Charophytes collapse beyond a critical warming and brownification

threshold in shallow lake systems. Science of The Total Environment 661:148–154.

Daunys, D., and M. L. Zettler. 2006. Invasion of the North Amrican Amphipod (Gammarus

tigrinus Sexton, 1939) into the Curonian Lagoon, South-Eastern Baltic Sea. Acta

Zoologica Lituanica 16:20–26.

Dorgelo, J. 1973. Comparative ecophysiology of gammarids (crustacea: amphipoda) from

marine, brackish and fresh-water habitats exposed to the influence of salinity-

temperature combinations. III. Oxygen uptake. Netherlands Journal of Sea Research

7:253–266.

Dorgelo, J. 1974. Comparative ecophysiology of gammarids (Crustacea: Amphipoda) from

marine, brackish and fresh-water habitats, exposed to the influence of salinity-

temperature combinations - I. Effect on survival. Hydrobiological Bulletin 8:90–108.

Felten, V., G. Tixier, F. Guérold, V. De Crespin De Billy, O. Dangles, F. Guerold, De Crespin

V. De Billy, O. Dangles, F. Guérold, V. De Crespin De Billy, and O. Dangles. 2008.

Quantification of diet variability in a stream amphipod: Implications for ecosystem

functioning. Fundamental and Applied Limnology / Archiv für Hydrobiologie 170:303–

313.

Folmer, O., M. Black, W. Hoeh, R. Lutz, and R. Vrijenhoek. 1994. DNA primers for

amplification of mitochondrial cytochrome c oxidase subunit I from diverse metazoan

invertebrates. Molecular marine biology and biotechnology 3:294–299.

Fox, J., and S. Weisberg. 2019. An {R} Companion to Applied Regression. Third. Sage,

Thousand Oaks {CA}.

Gärdenfors, U. 2010. 2010 Rödlistade arten I Sverige (The 2010 Red List of Swedish

Species). ArtDatabanken. Uppsala.

Gessner, F. 1957. Meer und Strand. Page (F. Gessner, Ed.). Second edition. VEB Deutscher

Verlag der Wissenschaften, Berlin.

Grabner, D. S., A. M. Weigand, F. Leese, C. Winking, D. Hering, R. Tollrian, and B. Sures.

2015. Invaders, natives and their enemies: distribution patterns of amphipods and their

microsporidian parasites in the Ruhr Metropolis, Germany. Parasites and Vectors 8:1–

15.

Hasslow, O. J. 1931. Sveriges characéer. Botaniska Notiser:63–136.

HELCOM. 2010. Ecosystem Health of the Baltic sea 2003-2007: HELCOM Initial Holistic

Assessment. Baltic Sea Environment Proceedings 122:68.

HELCOM. 2018. Sources and pathways of nutrients to the Baltic Sea. Baltic Sea

Environmental Proceedings 153:48.

Herkül, K., V. Lauringson, and J. Kotta. 2016. Specialization among amphipods: The invasive

Gammarus tigrinus has narrower niche space compared to native gammarids. Ecosphere

7:1–16.

Janssen, F., T. Neumann, and M. Schmidt. 2004. Inter-annual variability in cyanobacteria

blooms in the Baltic Sea controlled by wintertime hydrographic conditions. Marine

Ecology Progress Series 275:59–68.

Karstens, S., U. Buczko, and S. Glatzel. 2015. Phosphorus storage and mobilization in coastal

Phragmites wetlands: Influence of local-scale hydrodynamics. Estuarine, Coastal and

Shelf Science 164:124–133.

Karstens, S., U. Buczko, G. Jurasinski, R. Peticzka, and S. Glatzel. 2016. Impact of adjacent

land use on coastal wetland sediments. Science of the Total Environment 550:337–348.

Kelley, A. L. 2014. The role thermal physiology plays in species invasion. Conservation

Physiology 2:1–14.

Kelly, D. W., J. T. A. Dick, and W. I. Montgomery. 2002. The functional role of Gammarus

(Crustacea, Amphipoda): Shredders, predators, or both? Hydrobiologia 485:199–203.

Kelly, D. W., J. R. Muirhead, D. D. Heath, and H. J. Macisaac. 2006. Contrasting patterns in

genetic diversity following multiple invasions of fresh and brackish waters. Molecular

Ecology 15:3641–3653.

Koop, J. H. E., and M. K. Grieshaber. 2000. The role of ion regulation in the control of the

distribution of Gammarus tigrinus (Sexton) in salt-polluted rivers. Journal of

Comparative Physiology - B Biochemical, Systemic, and Environmental Physiology

170:75–83.

Körner, S., and T. Dugdale. 2003. Is roach herbivory preventing re-colonization of submerged

macrophytes in a shallow lake? Hydrobiologia 506–509:497–501.

Korpinen, S., and M. Westerbom. 2009. Microhabitat segregation of the amphipod genus

Gammarus (Crustacea: Amphipoda) in the Northern Baltic Sea. Marine Biology

157:361–370.

Kotta, J., H. Orav-Kotta, K. Herkül, and I. Kotta. 2011. Habitat choice of the invasive

Gammarus tigrinus and the native *Gammarus salinus* indicates weak interspecific

competition. Boreal Environment Research 16:64–72.

Kraufvelin, P., S. Salovius, H. Christie, F. E. Moy, R. Karez, and M. F. Pedersen. 2006.

Eutrophication-induced changes in benthic algae affect the behaviour and fitness of the

marine amphipod Gammarus locusta. Aquatic Botany 84:199–209.

Kufel, L., and I. Kufel. 2002. Chara beds acting as nutrient sinks in shallow lakes—a review.

Aquatic Botany 72:249–260.

Kumar, S., G. Stecher, M. Li, C. Knyaz, and K. Tamura. 2018. MEGA X: Molecular

evolutionary genetics analysis across computing platforms. Molecular Biology and

Evolution 35:1547–1549.

Kuprijanov, I., J. Kotta, V. Lauringson, and K. Herkül. 2015. Trophic interactions between

native and alien palaemonid prawns and an alien gammarid in a brackish water

ecosystem. Proceedings of the Estonian Academy of Sciences 64:518–524.

Leigh, J. W., and D. Bryant. 2015. POPART: Full-feature software for haplotype network

construction. Methods in Ecology and Evolution 6:1110–1116.

Lenz, M., B. A. P. P. da Gama, N. V. Gerner, J. Gobin, F. Gröner, A. Harry, S. R. Jenkins, P.

Kraufvelin, C. Mummelthei, J. Sareyka, E. A. Xavier, and M. Wahl. 2011. Non-native

marine invertebrates are more tolerant towards environmental stress than taxonomically

related native species: results from a globally replicated study. Environmental Research

111:943–52.

van der Linden, P., and J. F. B. Mitchell. (n.d.). ENSEMBLES: Climate Change and its

Impacts: Summary of research and results from the ENSEMBLES project. FitzRoy

Road, Exeter EX1 3PB, UK.

Lindner, A. 1972. Soziologisch-ökologische Untersuchungen an der submersen Vegetation in

der Boddenkette südlich des Darß und Zingst. University of Rostock.

Melzer, A. 1999. Aquatic macrophytes as tools for lake management. Hydrobiologia

395/396:181–190.

Neumann, T., and R. Friedland. 2011. Climate Change Impacts on the Baltic Sea. Pages 23–

32 *in* G. Schernewski, J. Hofstede, and T. Neumann, editors. Global Change and Baltic Coastal Zones. Springer.

Normant, M., M. Feike, A. Szaniawska, and G. Graf. 2007. Adaptation of Gammarus tigrinus Sexton, 1939 to new environments-Some metabolic investigations. Thermochimica Acta 458:107–111.

Orav-Kotta, H., J. Kotta, K. Herkül, I. Kotta, and T. Paalme. 2009. Seasonal variability in the grazing potential of the invasive amphipod Gammarus tigrinus and the native amphipod Gammarus salinus (Amphipoda: Crustacea) in the northern Baltic Sea. Biological Invasions 11:597–608.

Östman, Ö., J. Eklöf, B. K. Eriksson, J. Olsson, P. O. Moksnes, and U. Bergström. 2016. Top-down control as important as nutrient enrichment for eutrophication effects in North Atlantic coastal ecosystems. Journal of Applied Ecology 53:1138–1147.

Paar, M., M. Berthold, R. Schumann, S. Dahlke, and I. Blindow. 2021. Seasonal Variation in Biomass and Production of the Macrophytobenthos in two Lagoons in the Southern Baltic Sea. Frontiers in Earth Science 8:1–16.

Paiva, F., A. Barco, Y. Chen, A. Mirzajani, F. T. Chan, V. Lauringson, M. Baltazar-Soares, A. Zhan, S. A. Bailey, J. Javidpour, and E. Briski. 2018. Is salinity an obstacle for biological invasions? Global Change Biology 24:2708–2720.

Pellan, L., V. Médoc, D. Renault, T. Spataro, and C. Piscart. 2016. Feeding choice and predation pressure of two invasive gammarids, Gammarus tigrinus and Dikerogammarus villosus, under increasing temperature. Hydrobiologia 781:43–54.

Piepho, M. 2017. Assessing maximum depth distribution, vegetated area, and production of submerged macrophytes in shallow, turbid coastal lagoons of the southern Baltic Sea.

Hydrobiologia 794:303–316.

Pitkänen, H., M. Peuraniemi, M. Westerbom, M. Kilpi, and M. Von Numers. 2013. Long-
term changes in distribution and frequency of aquatic vascular plants and charophytes in
an estuary in the Baltic Sea. Annales Botanici Fennici 50:1–54.

Pohl, P., M. K. Ohlhase, S. K. Rautwurst, and K. Laus-Kinnerk Baasch. 1987. An inexpensive
inorganic medium for the mass cultivation of freshwater microalgae. Phytochemistry
26:1657–1659.

Puche, E., S. Sánchez-Carrillo, M. Álvarez-Cobelas, A. Pukacz, M. A. Rodrigo, and C. Rojo.
2018. Effects of overabundant nitrate and warmer temperatures on charophytes: The
roles of plasticity and local adaptation. Aquatic Botany 146:15–22.

Puntila, R. 2016. Trophic Interactions and Impacts of Non-indigenous Species in Baltic Sea
Coastal Ecosystems. University of Helsinki.

R Core Team. 2019. R: A Language and Environment for Statistical Computing. Vienna,
Austria.

Radulovici, A. E., B. Sainte-Marie, and F. Dufresne. 2009. DNA barcoding of marine
crustaceans from the Estuary and Gulf of St Lawrence: A regional-scale approach.
Molecular Ecology Resources 9:181–187.

Reusch, T. B. H., J. Dierking, H. C. Andersson, E. Bonsdorff, J. Carstensen, M. Casini, M.
Czajkowski, B. Hasler, K. Hinsby, K. Hyytiäinen, K. Johannesson, S. Jomaa, V.
Jormalainen, H. Kuosa, S. Kurland, L. Laikre, B. R. MacKenzie, P. Margonski, F.
Melzner, D. Oesterwind, H. Ojaveer, J. C. Refsgaard, A. Sandström, G. Schwarz, K.
Tonderski, M. Winder, and M. Zandersen. 2018. The Baltic Sea as a time machine for
the future coastal ocean. Science Advances 4:eaar8195.

Rojo, C., M. Carramiñana, D. Cócera, G. P. Roberts, E. Puche, S. Calero, and M. A. Rodrigo. 2017. Different responses of coexisting Chara species to foreseeable Mediterranean temperature and salinity increases. Aquatic Botany in press.

Sanudo-Wilhelmy, S. A., A. Tovar-Sanchez, F. Fu, D. G. Capone, E. J. Carpenter, and D. A. Hutchins. 2004. The impact of surface-adsorbed phosphorus on phytoplankton Redfield stoichiometry. Nature 432:897–901.

Schiewer, U. 2007. Darß-Zingst Boddens, Northern Rügener Boddens and Schlei. Pages 35–86 *in* U. Schiewer, M. M. Caldwell, G. Heldmaier, R. B. Jackson, O. L. Lange, H. A. Mooney, E.-D. Schulze, and U. Sommer, editors. Ecology of Baltic coastal waters. First edition. Springer, Berlin, Heidelberg.

Schubert, H., and I. Blindow, editors. 2003. Charophytes of the Baltic Sea. Gantner, Ruggel, Königstein and Germany.

Schubert, H., I. Blindow, N. C. Bueno, M. T. Casanova, M. Pełechaty, and A. Pukacz. 2018. Ecology of charophytes – permanent pioneers and ecosystem engineers. Perspectives in Phycology 5:61–74.

Schumann, R., H. Baudler, Ä. Glass, K. Dümcke, and U. Karsten. 2006. Long-term observation on salinity dynamics in a tideless shallow coastal lagoon of the Southern Baltic Sea coast and their biological relevance. Journal of Marine Systems 60:330–344.

Stewart, N. F., and J. M. Church. 1992. Red data books of Britain and Ireland-- Stoneworts. {Joint Nature Conservation Committee} and {Office of Public Works} and Plantlife, Peterborough.

Torn, K., G. Martin, and R. Munsterhjelm. 2003. Chara tomentosa L. 1753. Pages 131–141 *in* H. Schubert and I. Blindow, editors. Charophytes of the Baltic Sea. Gantner, Ruggel,

Königstein, Germany.

Torn, K., G. Martin, and T. Paalme. 2006. Seasonal changes in biomass, elongation growth

and primary production rate of Chara tomentosa in the NE Baltic Sea. Annales Botanici

Fennici 43:276–283.

Volkmann, C. 2016. The impact of organic material for macrophytes in coastal waters of the

Baltic Sea. University of Rostock.

Wickham, H. 2016. ggplot2: Elegant Graphics for Data Analysis. Springer-Verlag New York.

Wood, S. N. 2011. Fast stable restricted maximum likelihood and marginal likelihood

estimation of semiparametric generalized linear models. Journal of the Royal Statistical

Society (B) 73:3–36.

Wüstenberg, A., Y. Pörs, and R. Ehwald. 2011. Culturing of stoneworts and submersed

angiosperms with phosphate uptake exclusively from an artificial sediment. Freshwater

Biology 56:1531–1539.

Zettler, M. L. 1995. Erstnachweis von Gammarus tigrinus Sexton, 1939 (Crustacea:

Amphipoda) in der Darss-Zingster-Boddenkette und seine derzeitige Verbreitung an der

deutschen Ostsekueste. Arch. Freunde Naturg. Mecklb. XXXIV:137–141.

Zettler, M. L. 2001. Some malacostracan crustacean assemblages in the southern and western

Baltic Sea. Rostock. Meersebiolog. Beitr. 9:127–143.

Zettler, M. L., and A. Zettler. 2017. Marine and freshwater Amphipoda from the Baltic Sea

and adjacent territories. 2. Auflage. Conchbooks, Hackenheim.

 <u>Figures</u>

Figure 1: Seasonal losses of *Chara* sp. biomass by grazing of *Gammarus tigrinus* in the

Bodstedter Bodden (upper panel) and Grabow (lower panel) over the years of 2000 to 2018.

The losses of *Chara* sp. biomass by individual grazing (points) were modelled (generalized

additive model) depending on temperature and salinity conditions at the respective time point

in the water body (left). Grazing per square meter (right) was calculated as the product of the

individual grazing model (left) and the seasonal population development of *Gammarus*

tigrinus (Supplement Figure S5) with a maximum abundance of 400 individuals per square

meter. The solid line represents the mean and the dotted line the 95 % confidence interval.

Figure 2 Charophyte fresh mass in gram per square meter at two sampling locations of the

Darss-Zingst lagoon system along a water depth/distance to reed belt transect. Shallow depth

(30 – 60 cm) and close to reed belt, intermediate depth (60 – 90 cm) and distance to reed belt,

deep depth (90 – 120 cm) and farthest distance to reed belt. Net biomass represents model-

fitted values (generalized additive model) of observed charophyte biomass. Green points

represent observed charophyte biomass at the respective day. Gross biomass represents the

sum of net biomass plus biomass possibly lost by abiotic influenced seasonally resolved

Gammarus tigrinus grazing.

Figure 3: Future projection of seasonal losses of *Chara* sp. biomass by grazing of *Gammarus*

tigrinus in the Bodstedter Bodden (upper panel) and Grabow (lower panel) as difference to

present day grazing. Projections were calculated with a delta change approach of recent data

with projected changes. Data from 2000 to 2018 was modified in temperature (+2.75 K) and

salinity (-1.75 g kg^{-1}) as described in Neumann and Freidland (2011) and van der Linden, P.

and Mitchell, J. F. B. (2009). Losses of *Chara* sp. biomass by individual grazing were

modelled depending on the modified temperature and salinity conditions at the respective

time point in the water body (left). Grazing per square meter (right) was calculated as the

product of the individual grazing model (left) and the seasonal population development of

Gammarus tigrinus (Supplement Figure S5) with a maximum abundance of 400 individuals

per square meter. The solid line represents the mean and the dotted line the 95 % confidence

interval.

Tables

Table 1: Sum of losses of *Chara* sp. biomass by individual grazing or by grazing at one
square meter with respect to the seasonal population pattern of *Gammarus tigrinus*
(Supplement Figure S5) with a maximum abundance of 400 individuals per square meter.
These were calculated for the Bodstedter Bodden and the Grabow as well as for the recent
data and the future projection.

CWB	base	recent [mg a^{-1}]	projected [mg a^{-1}]	$\Delta_{(proj.-rec.)}$ [mg a^{-1}]	Δ / rec. [%]
Bodstedter Bodden	ind.	672,6	687,3	14,7	2,2
Grabow	ind.	768,3	847,9	79,6	10,4
Bodstedter Bodden	m²	36132,5	35860,5	-272	-0,8
Grabow	m²	41014,9	44230,3	3215,5	7,8

Supplement

<u>Genetic analyses</u>

At the end of the grazing experiment, 69 gammarids were determined as *Gammarus tigrinus*

following Zettler and Zettler (2017) and preserved in 80 % ethanol. To verify the

morphological analyses, a genetic approach was conducted on 15 individuals. Total genomic

DNA was extracted using a silica spin column procedure with the DNeasy Blood and Tissue

Kit (Qiagen, Hilden, Germany) following the protocol provided by the manufacturer. Partial

sequences of the COI gene were amplified with the universal primers LCO 1490 and HCO

2198 (Folmer et al. 1994).

PCR amplifications were performed with a denaturation step for 60 s at 94°C, followed by 5

cycles of: 60 s at 94°C, 90 s at 45°C and 60 s at 72°C, 35 cycles of: 60 s at 94°C, 90 s at 51°C

and 60 s at 72°C, and completed with 5 min at 72°C as a final extension step. PCR was

performed in a 30 µl reaction volume with a Taq PCR Master Mix (Qiagen, Hilden,

Germany) consisting 2.5 mM MgCl2, and 0.5 pmol of each primer (final concentration). The

PCR products were extracted from agarose gels according to the protocol of the Biometra-

innuPrep Gel Extraction Kit (Analytik Jena, Jena, Germany), and were sequenced directly

using a 3130×L Genetic Analyzer (Applied Biosystems, NY, USA) with sequencing primers

identical to the primers that were used for the PCR reaction. Obtained sequences were quality

controlled and aligned via the BIOEDIT software (Hall 1999). The COI gene is widely used

for inter- and intraspecific diversification questions concerning amphipod taxa. For the

taxonomic determination addition GenBank sequences of *Gammarus tigrinus* (see

Supplement Table S1), *G. duebeni* Liljeborg, 1852 (EU421779), *G. locusta* Linnaeus, 1758

(KT209211), *G. oceanicus* Segerstråle, 1947 (GQ341809), *G. pulex* Linnaeus, 1758

(MN400977), *G. roeseli* Gervais, 1835 (EF570337) *G. salinus* Spooner, 1947 (KT208533),

and *G. zaddachi* Sexton, 1912 (KU845083) were used for the phylogenetic analyses.

In order to estimate the evolutionary divergence between haplotypes, pair-wise uncorrected p-

distances and the number of substitutions were conducted using MEGA version X (Kumar et

al. 2018). To uncover phylogenetic relationships, a Maximum likelihood tree was constructed

using MEGA version X (Kumar et al. 2018) on the basis of the Kimura 2-parameter model.

Branch support for the nodes was calculated from 1000 bootstrap replicates.

To study the distribution of mtDNA sequence diversity, a haplotype network was constructed

using the Median-joining algorithm (Bandelt et al. 1999) implemented in PopART (Leigh and

Bryant 2015). The distribution basemap was created with QGIS.org software

(http://www.qgis.org) and modified in CorelDRAW (Corel Corporation, Ottawa, Ontario,

Canada).

Supplement Table 1: Sample list of specimens used for the phylogenetic tree and the
determination of haplotypes. Indicated are the localities, if available the salinity of the
sampling site, the identified haplotypes and the accession numbers (F=freshwater sites).

Country	Site	Lat., Lon.	Sal.	haplotype	AccessionNo	Ref.
Denmark	Bornholm, Hovedstaden, Baltic Sea, Svaneke	55.1329, 15.152		N1	MK403734	Baltazar-Soares et al. (2017)
	Bornholm, Hovedstaden, Ostersoen stream, Nexo	55.0569, 15.127		N1, N4a	MK403733, MK403736	Baltazar-Soares et al. (2017)
Estonia	Liu	58.2744, 24.2528	4.7	N1, N4a, N4e	KU845009-KU845015, KU845017-KU845021, KU845023-KU845027	Paiva et al. 2018
	Pärnu	58.3748, 24.4734	4.28	N1, N4a, N4d, N4e	KU845029, KU845030, KU845033, KU845035-KU845038, KU845040-KU845043, KU845046, KU845049, KU845050	(Paiva et al. 2018)
Finland	Turku	60.4,	4.4	N4a, N4d,	DQ523177, DQ523178,	Kelly et al. 2006

		22.2		N4e	DQ523183	
Germany	**Born, DZLS**	**54.379434, 12.524819**	**5.5**	**N4a, N4e**	**MW509071-MW509085**	**this study**
	Dierhagen lagoon	54.2, 12.3	1.5	N4a, N4e	DQ523178, DQ523183	Kelly et al. 2006
	Travemümde	53.9655, 10.8851	10	N1	KU844994-KU845006	Paiva et al. 2018
	North Rhine	51.7328950, 7.1774890	F	N1, N4a, N4c	KT075215, KT075216, KT075218-KT075220	(Grabner et al. 2015)
Netherlands	Buiten-IJ, Amsterdamm	52.369028, 4.990722	F	N4d	EF570348	Hou et al. 2007
	Lake Gouwzee,	52.4, 5.0	F	N4a, N4c	DQ300227, DQ523183	Kelly et al. 2006
North Ireland	Bann river	54.8, -6.4	F	N4a, N4b, N4e	DQ523178, DQ523181, DQ523183	Kelly et al. 2006
	Lough Neagh,	54.7, -6.5	F	N4a, N4d, N4e	DQ523177, DQ523178, DQ523183	Kelly et al. 2006
Poland	Brody, Oder river	52.0, 15.4	F	N4a, N4d, N4e	DQ523177, DQ523178, DQ523183	Kelly et al. 2006
	Bytom, Oder river	51.7, 15.8	F	N4a, N4d, N4e	DQ523177, DQ523178, DQ523183	Kelly et al. 2006
	Gdansk Bay, Puck	54.723394, 18.416633		N4a, N4e	GQ341858-GQ341861	Costa et al. 2009
	Vistula lagoon	54.3, 19.7	4.5	N1, N4a	DQ300212, DQ523183	Kelly et al. 2006
Canada	New Brunswick	47.037389, -65.163250		N1, N2	FJ581678, FJ581679	(Radulovici et al. 2009)
	Nova Scotia	45.631983, -61.958414		N2	FJ581680-FJ581683	Radulovici et al. 2009
	Quebec	47.149100, -70.521900		N1, N3	FJ581684-FJ581690	Radulovici et al. 2009

	St. John estuary	45.3, -66.2	4.1	N1	DQ300250, DQ300251	(Kelly et al. 2006)
	St. Lawrence, d/s Quebec, Cap Brule	46.9, -70.5	6.5	N3	DQ300211	Kelly et al. 2006
USA	Berrys creek, New Hampshire	43.0, -70.7	16	N1	DQ300208-DQ300210	Kelly et al. 2006
	Chesapeake Bay, Virginia	37.45, -76.67		N4b	DQ300219, DQ300220, DQ523180, DQ523181	Kelly et al. 2006
	Delaware estuary, Deemers Beach, Delaware	39.6, -75.5	5	N3,N4a, N4b	DQ523181-DQ523184, DQ523189	Kelly et al. 2006
	Delaware estuary, Delaware	39.57, -75.58		N3, N4a	DQ300221, DQ300222, DQ300225	Kelly et al. 2006
	Elizabeth estuary, Virginia	36.7, -76.2	10.2	N3, N4a, N4c	DQ300224, DQ300226, DQ300227-DQ300231, DQ523183, DQ523186-DQ523188, DQ523190-DQ523195	Kelly et al. 2006
	Hudson estuary, New York	40.9, -73.8	7.8	N3	DQ523179	Kelly et al. 2006
	Maryland, Rhode River, SERC Education Dock	38.885, -76.542		N4a, N4b	KU905729, KU905737, KU905742, KU905952	unpublished, direct submission
	Maryland, Thoroughfare Island, Potomac River, Chesapeake Bay	38.5731, -77.175		N4b	MH235818	(Baltazar-Soares et al. 2017)
	Neuse River: d/s New Bern, N. Carolina	35.1, -77.0	14.2	Southern Species	DQ300240-DQ300244	Kelly et al. 2006
	Pawcatuck estuary, Rhode Island	41.3, -71.8	11.2	N2	DQ300245-DQ300249	Kelly et al. 2006

746   Supplement Table 2: Differences of biomass change between control group (without

747   G. tigrinus) and treatment group (with G. tigrinus) with respect to species, Salinity (S) and

748   temperature (T). Furthermore the differences between the groups were statistical analyzed and

749   the p-value calculated with the non-parametrical Mann-Whitney U-Test.

Species	S [g·kg⁻¹]	T [°C]	N	Median Control	Median Treatment	Δ Median (T-C)	p-value (U-Test)
Chara aspera	3	9	8	-0,7	-11,0	-10,3	0,01*
Chara aspera	3	15	8	-0,7	-13,7	-13,0	0,01*
Chara aspera	3	24	8	-7,3	-26,0	-18,7	0,03*
Chara aspera	3	30	8	-15,3	-11,8	3,5	0,60
Chara aspera	5	9	8	-0,5	-9,0	-8,5	0,01*
Chara aspera	5	15	8	-2,9	-15,2	-12,2	0,01*
Chara aspera	5	24	8	-2,3	-20,1	-17,8	0,01*
Chara aspera	5	30	8	-3,3	-28,5	-25,2	0,00*
Chara aspera	7	9	8	0,2	-7,6	-7,8	0,12
Chara aspera	7	15	8	0,4	-18,5	-18,8	0,01*
Chara aspera	7	24	8	-2,2	-33,6	-31,3	0,00*
Chara aspera	7	30	8	-12,9	-34,9	-22,0	0,03*
Chara aspera	13	9	8	0,9	-5,9	-6,8	0,02*
Chara aspera	13	15	8	0,2	-13,5	-13,7	0,02*
Chara aspera	13	24	8	-4,2	-35,3	-31,1	0,01*
Chara aspera	13	30	8	-14,3	-34,0	-19,8	0,02*
Chara tomentosa	3	9	8	-0,7	-2,6	-1,9	0,17
Chara tomentosa	3	15	8	-2,8	-7,3	-4,5	0,12
Chara tomentosa	3	24	8	-2,0	-17,3	-15,4	0,01*
Chara tomentosa	3	30	8	-4,7	-14,5	-9,7	0,04*
Chara tomentosa	5	9	8	0,9	-5,5	-6,4	0,04*
Chara tomentosa	5	15	8	-0,3	-9,0	-8,7	0,14
Chara tomentosa	5	24	8	-3,6	-8,2	-4,6	0,03*
Chara tomentosa	5	30	8	-3,3	-11,1	-7,9	0,03*
Chara tomentosa	7	9	8	-1,7	-6,0	-4,3	0,12
Chara tomentosa	7	15	8	3,2	-6,5	-9,7	0,01*
Chara tomentosa	7	24	8	2,0	-17,0	-19,0	0,04*
Chara tomentosa	7	30	8	-7,0	-9,6	-2,6	0,35
Chara tomentosa	13	9	8	-0,9	-5,7	-4,8	0,00*
Chara tomentosa	13	15	8	-0,1	-7,5	-7,4	0,25
Chara tomentosa	13	24	8	-1,5	-6,7	-5,2	0,02*
Chara tomentosa	13	30	8	-4,0	-13,7	-9,7	0,02*
Chara sp.	3	9	16	-0,7	-7,5	-6,8	0,01*
Chara sp.	3	15	16	-1,7	-8,6	-6,9	0,00*
Chara sp.	3	24	16	-3,1	-17,7	-14,6	0,00*
Chara sp.	3	30	16	-6,1	-14,0	-7,8	0,03*
Chara sp.	5	9	16	0,2	-8,3	-8,4	0,00*
Chara sp.	5	15	16	-1,7	-10,4	-8,7	0,00*
Chara sp.	5	24	16	-2,6	-14,9	-12,4	0,00*
Chara sp.	5	30	16	-3,3	-15,5	-12,2	0,00*
Chara sp.	7	9	16	-1,1	-6,2	-5,1	0,03*
Chara sp.	7	15	16	1,8	-8,3	-10,0	0,00*
Chara sp.	7	24	16	1,2	-22,9	-24,2	0,00*

Chara sp.	7	30	16	-10,7	-20,4	-9,6	0,04*
Chara sp.	13	9	16	0,6	-5,9	-6,6	0,00*
Chara sp.	13	15	16	0,0	-9,9	-9,9	0,01*
Chara sp.	13	24	16	-2,3	-29,1	-26,9	0,00*
Chara sp.	13	30	16	-8,1	-20,0	-11,8	0,01*

Figure S1 Flow scheme of the full factorial experimental design for temperature, and salinity

with *Chara aspera* and *C. tomentosa*, with and without *Gammarus tigrinus*.

Figure S2 (A) Phylogeny of *Gammarus tigrinus* COI haplotypes inferred by using the

Maximum Likelihood method (Tamura-Nei model +I+Γ) with bootstrap values above the

branches (1000 replicates). The clade numbers are identical to Kelly et al. (2006) and the

coloured boxes indicate haplotypes of clade N4. Haplotypes N4a and N4e comprised the

individuals from the DZLS. (B & C) Distribution of *Gammarus tigrinus* haplotypes of clade

N4 identified by COI network analysis. Pie charts indicated haplotype frequencies in

European (B) and North American (C) populations. More information on haplotypes can be

found in Supplement Table S1.

Figure S3 Generalized additive model for change in biomass of *Chara aspera* (top left),

Chara tomentosa (top right) and for both Chara species (bottom) in respect to different

salinity and temperature levels.

Figure S4 Generalized additive model for loss of biomass caused by growth corrected grazing

of an individual of *Gammarus tigrinus* (G.t.) on *Chara aspera* (top left), *Chara tomentosa*

(top right) and on both Chara species (bottom) in respect to different salinity and temperature

levels.

Figure S5 Model of seasonal pattern of *Gammarus tigrinus* calculated with population

densities from Chambers (1977) and modified to have a maximum of 400 individuals per

square meter based on values of Zettler (1995) and of M. Paar (personal communication).

<u>Elemental composition</u>

There were no significant differences for the C:N and C:P ratios along the depth distribution,

or between stations (see pooled values Figure S6). Charophytes showed either a weak

seasonal trend in elemental composition (C:N), or the ratio decreased (C:P). Lowest C:P

ratios were found from July to October. Contrary, co-occurring tracheophytes increased their

C:N ratio during summer, as well as their C:P ratios. These findings indicate that charophytes

had probably a higher protein content per biomass compared to tracheophytes.

Figure S6 Elemental ratio of Carbon to Nitrogen (C:N) and Carbon to Phosphorus (C:P) of

macrophyte biomass (separated by order) at two stations of the Darss-Zingst lagoon system

across all sampling stations (Bodstedter Bodden and Grabow) and sampling depths in 2014.

Please note the different scaling. Points represent measured elemental ratios, lines a local

regression (LOESS – locally estimated scatter smoothing), and the ribbons the 95%

confidence intervals (Wickham 2016).

793