Disruption of the *Pseudomonas aeruginosa* Tat system perturbs PQS-dependent quorum sensing and biofilm maturation through loss of the Rieske cytochrome *bc₁* sub-unit

*Eliza Ye-Chen Soh[1], *Frances Smith[1], Maxime Rémi Gimenez[2], Liang Yang[3†], Rebecca Munk Vejborg[4], Matthew Fletcher[1], Nigel Halliday[1], Sophie Bleves[2], Stephan Heeb[1], Miguel Camara[1], Michael Givskov[3,4], Kim R. Hardie[1], Tim Tolker-Nielsen[4,] Bérengère Ize[2] and *Paul Williams[1]*

[1]Biodiscovery Institute, National Biofilms Innovation Centre and School of Life Sciences, University of Nottingham, Nottingham, U.K.
[2]Laboratoire d'Ingénierie des Systèmes Macromoléculaires (LISM-UMR7255), Institut de Microbiologie de la Méditerranée, CNRS and Aix-Marseille Univ. 31 Chemin Joseph Aiguier, CS 70071, 13402 Marseille cedex 09, France.
[3]Singapore Centre for Environmental Life Sciences Engineering, Nanyang Technological University, 60 Nanyang Drive, Singapore 637551.
[4]Costerton Biofilm Center, Department of Immunology and Microbiology, Faculty of Health Sciences, University of Copenhagen, 2200 Copenhagen N, Denmark.

*joint first authors

†Present Address: School of Medicine, Southern University of Science and Technology, Shenzhen 518055, Guangdong, China

Running Head: Tat, quorum sensing and biofilms in *P. aeruginosa*

Keywords: quorum sensing; biofilms; twin arginine translocation system (Tat) tobramycin; *Pseudomonas aeruginosa*; *Pseudomonas* quinolone signal; PQS; 2-heptyl-3-hydroxy-4-quinolone; cytochrome bc₁ Rieske sub-unit

Abstract

Extracellular DNA (eDNA) is a major constituent of the extracellular matrix of *Pseudomonas aeruginosa* biofilms and its release is regulated via the pseudomonas quinolone signal (PQS) dependent quorum sensing (QS). By screening a *P. aeruginosa* transposon library to identify factors required for DNA release, mutants with insertions in the twin-arginine translocation (Tat) pathway were identified as exhibiting reduced eDNA release, and defective biofilm architecture with enhanced susceptibility to tobramycin. *P. aeruginosa tat* mutants showed substantial reductions in pyocyanin, rhamnolipid and membrane vesicle (MV) production consistent with perturbation of 2-heptyl-3-hydroxy-4-quinolone (PQS) dependent QS as demonstrated by changes in *pqsA* expression and 2-alkyl-4-quinolone (AQ) production. Provision of exogenous PQS to the *tat* mutants did not return *pqsA*, *rhlA* or *phzA1* expression or pyocyanin production to wild type levels. However, transformation of the *tat* mutants with the AQ-independent *pqs* effector *pqsE* restored *phzA1* expression and pyocyanin production. Since mutation or inhibition of Tat prevented PQS-driven auto-induction, we sought to identify the Tat secretion substrate responsible. A *pqsA::lux* fusion was introduced into each of 34 validated *P. aeruginosa* Tat substrate deletion mutants. Analysis of each mutant for reduced bioluminescence revealed that the signalling defect was associated with the Rieske iron-sulfur subunit of the cytochrome bc_1 complex. In common with the parent strain, a Rieske mutant exhibited defective PQS signalling, AQ production, *rhlA* expression and eDNA release that could be restored by genetic complementation. Thus, lack of the Rieske sub-unit export is clearly responsible for the Tat-mediated perturbation of PQS-dependent QS, the loss of virulence factor production, biofilm eDNA and the tobramycin tolerance of *P. aeruginosa* biofilms.

Author Summary

Pseudomonas aeruginosa is a highly adaptable human pathogen responsible for causing chronic biofilm-associated infections. Biofilms are highly refractory to host defences and antibiotics and thus difficult to eradicate. The biofilm extracellular matrix incorporates extracellular DNA (eDNA). This stabilizes biofilm architecture and helps confer tolerance to antibiotics. Since mechanisms that control eDNA release are not well understood, we screened a *P. aeruginosa* mutant bank for strains with defects in eDNA release and discovered a role for the twin-arginine translocation (Tat) pathway that exports folded proteins across the cytoplasmic membrane. Perturbation of the Tat pathway resulted in defective biofilms susceptible to antibiotic treatment as a consequence of perturbed pseudomonas quinolone (PQS) signalling. This resulted in the failure to produce or release biofilm components including eDNA, phenazines and rhamnolipids as well as microvesicles. Furthermore, we discovered that perturbation of PQS signalling was a consequence of the inability of *tat* mutants to translocate the Rieske subunit of the cytochrome bc_1 complex involved in electron transfer and energy transduction. Given the importance of PQS signalling and the Tat system to virulence and biofilm maturation in *P. aeruginosa*, our findings underline the potential of the Tat system as a drug target for novel antimicrobial agents.

Introduction

Pseudomonas aeruginosa is an opportunistic pathogen that causes a wide range of human infections including lung, urinary tract and wound, bacteremia and infections associated with medical devices [1]. It is notorious for its tolerance to antimicrobial agents, a property that is largely a consequence of its ability to form biofilm communities [1,2]. Bacterial exoproducts including cell surface appendages, extracellular polymeric substances, biosurfactants and secondary metabolites all contribute to *P. aeruginosa* biofilm formation and maturation [3-7].

Apart from exopolysaccharides such as Psl, Pel and alginate, the extracellular polymeric matrix of *P. aeruginosa* biofilms incorporates proteins, rhamnolipids, membrane vesicles (MVs) and extracellular DNA (eDNA) [5,8-10]. Rhamnolipid biosurfactants are required during the initial stages of micro-colony formation and are involved in the migration-dependent formation of the caps of mushroom-shaped micro-colonies formed in flow-cell grown biofilms [10]. They also aid the maintenance of channels between multicellular structures within biofilms and contribute to biofilm dispersal [5]. With respect to the biofilm micro-colonies that characteristically form in flow-chambers fed with glucose minimal medium, eDNA is present at high concentrations in the outer layers of microcolonies in young biofilms. However, in mature biofilms, eDNA is primarily located in the stalks at the borders between micro-colony caps and stalks [8].

The release of eDNA occurs via the lysis of a sub-population of bacterial cells [10-14]. It is involved in attachment, aggregation and stabilization of biofilm microcolonies. eDNA can act as a nutrient source, chelate metal cations and confer tolerance to antibiotics such as the polymyxins and aminoglycosides [10,12,13]. eDNA also binds other biopolymers (exopolysaccharides and proteins) stabilizing biofilm architecture and conferring protection against adverse chemical and physical

challenges [12,13]. By intercalating with eDNA, secondary metabolites such as

phenazines enhance biofilm integrity [12,15]. Pyocyanin for example can contribute

to DNA release through the formation of reactive oxygen species such as hydrogen

peroxide that damage cell membranes [12]. Although the mechanism(s) responsible

for eDNA release has not been fully elucidated both eDNA and MVs can be generated

via explosive cell lysis mediated via a cryptic prophage endolysin encoded within the

R- and F-pyocin gene clusters [14].

In *P. aeruginosa*, rhamnolipids and pyocyanin production, eDNA and MV release, and

hence biofilm development, are all controlled by quorum sensing (QS) [1,8,16].

Consequently, *P. aeruginosa* mutants with defects in this cell-to-cell communication

system form aberrant, flat undifferentiated biofilms [10]. In *P. aeruginosa*, the QS

regulatory network consists of a hierarchical cascade incorporating the overlapping

las, *rhl* and *pqs* pathways that employ *N*-acylhomoserine lactones (AHLs) and 2-

alkyl-4-quinolones (AQs) as signal molecules [1,16,17].   All three QS systems

contain auto-induction loops whereby activation of a dedicated transcriptional

regulator by the cognate QS signal molecule induces expression of the target

synthase such that QS signal molecule production can be rapidly amplified to

promote co-ordination of gene expression at the population level.

P. aeruginosa produces a diverse family of AQs and AQ *N*-oxides [18,19] of which

2-heptyl-3-hydroxy-4-quinolone (the *Pseudomonas* Quinolone Signal, PQS) and its

immediate precursor 2-heptyl-4-hydroxyquinoline (HHQ) are most closely

associated with PQS signalling [17]. Most of the genes required for AQ biosynthesis

(*pqsABCDE*) and response (*pqsR;* also called *mvfR*) are located at the same genetic

locus although *pqsH* and *pqsL* are distally located [17]. PqsA catalyses the formation

of anthraniloyl-CoA that is condensed with malonyl-CoA by PqsD to form 2-

aminobenzoylacetyl-CoA (2-ABA-CoA) [20,21]. The latter is converted to 2-

aminobenzoylacetate (2-ABA) via the thioesterase functionality of PqsE [22]. The

PqsBC heterodimer condenses 2-ABA with octanoyl-coA to generate HHQ [23,24].

PQS is formed through the oxidation of HHQ by PqsH [25] while formation of the AQ

N-oxides such as 2-heptyl-4-hydroxyquinoline *N*-oxide (HQNO) requires the

alternative mono-oxygenase PqsL [18]. The PqsE protein has dual functions; while

it is not essential for AQ biosynthesis, it is required for the AQ-independent

production of several factors that contribute to biofilm maturation including

pyocyanin, rhamnolipids and lectin A [26]. While activation of RhlR-dependent

genes depends on PqsE [27], the AQ-independent, thioesterase-independent

mechanism by which PqsE acts has not yet been elucidated [17,22].

The *pqs* system is subject to positive autoinduction, since the LysR-type

transcriptional regulator PqsR (MvfR), binds to the promoter region of *pqsABCDE*

(P*pqsA*) triggering transcription once activated by HHQ or PQS [28-30] (**Fig. 1**).

Therefore, by analogy with other QS systems, HHQ and PQS can both act as

autoinducers by generating a positive feedback loop that accelerates their

biosynthesis. However, in contrast to HHQ which only regulates the *pqsABCDE*

operon [17], PQS is a ferric iron chelator [31] that not only drives AQ biosynthesis

via PqsR but also the expression of genes involved in the iron-starvation response

and virulence factor production *via* PqsR-dependent and PqsR-independent

pathways [17]. In addition, PQS can act as a cell-sensitizing pro-oxidant [32] and

is required for MV production via a direct physical interaction with lipopolysaccharide

(LPS) within the outer membrane [33]. The packaging of PQS within MVs also

provides a means for trafficking this hydrophobic QS signal within

a *P. aeruginosa* population [34].

With respect to biofilm development, the *pqs* system is of particular interest because

pqs biosynthetic mutants fail to produce eDNA, rhamnolipids, pyocyanin and MVs,

and form thin defective biofilms containing little eDNA [8, 27,33,35]. The mechanism

involved in PQS-mediated DNA-release in biofilms is not understood but has been

suggested to be linked to phage induction causing cell lysis [8,36-40]. Although
explosive cell lysis releases eDNA in biofilms and generates MVs through
vesicularization of shattered membrane fragments, *pqsA* mutants are not defective
for explosive lysis [14] and therefore this phenomenon is unlikely to account for PQS-
dependent eDNA release.

In the present study we sought to identify additional factors involved in eDNA release
by screening a transposon (Tn) mutant library for eDNA-release defective mutants.
Apart from *pqs* biosynthetic mutants, we obtained Tn insertion mutants within the
twin-arginine translocation (Tat) pathway that exhibited reduced levels of eDNA
release, fail to produce rhamnolipids or pyocyanin and form defective, eDNA- poor,
antibiotic susceptible biofilms. Since mutation or deletion of *tat* resulted in altered
AQ production, reduced pyocyanin, rhamnolipid and MVs, and as the *tat* mutants
were refractory to exogenously supplied PQS, the aberrant biofilm phenotype
observed could be accounted for by perturbation of PQS autoinduction. By screening
a library of *P. aeruginosa* Tat substrate mutants, we identified the Rieske sub-unit of
the cytochrome bc_1 complex as being required for *pqs*-dependent QS and hence
eDNA release and biofilm maturation.

Results

Transposon mutagenesis screen for P. aeruginosa mutants exhibiting
reduced DNA release

To identify *P. aeruginosa* genes that contribute to eDNA release, a mariner
transposon (Tn) mutant library was generated in strain PAO1. Approximately 10,000
mutants grown in microtitre plates were assayed for reduced eDNA release using
propidium iodide (PI) to quantify eDNA because it is unable to penetrate live bacteria
and because its fluorescence is enhanced 30-fold on binding DNA [41]. From the

initial screen, 84 Tn insertion mutants were selected and re-screened to eliminate

strains with double Tn insertions and to confirm their eDNA phenotype. For 34 of

the remaining mutants, the regions flanking each Tn insertion were sequenced and

the corresponding genes identified. For many of the eDNA-release deficient mutants,

the Tn insertions were located within genes required for AQ biosynthesis (*pqsC* and

pqsH) or regulation (*pqsR*) (**Fig. 2**) These data confirm our previous work that first

uncovered a role for PQS signalling in eDNA release [8].   Apart from the *pqs*

mutants, two mutants were obtained with insertions in the *tatA* and *tatB* genes

respectively (**Fig. 2**). These code for components of the twin-arginine translocation

(Tat) system that mediates the export of folded proteins and was originally named

with respect to the presence of an Arg-Arg motif in the signal sequence of Tat-

exported products also called Tat substrates [42,43]. In *P. aeruginosa* diverse

proteins involved in phosphate and iron metabolism, virulence and energy

transduction are exported to the periplasm, or secreted via the Tat export system

and *tat* mutants exhibit pleiotropic phenotypes [44,45]. **Fig. 2** shows that genetic

complementation of the *P. aeruginosa tatA* mutant with a plasmid-borne copy

restored eDNA release.

The Tat pathway contributes to biofilm development in P. aeruginosa

Since eDNA makes an important contribution to biofilm development and

architecture [10,12,13], biofilm formation by the *P. aeruginosa tatA* mutant in flow-

chambers was investigated. After 4-days growth, the *P. aeruginosa* wild-type and

complemented *tatA*/pTatA mutant formed biofilms with mushroom-shaped

structures whereas the *tatA* mutant formed thin, flat biofilms (**Fig. 3A**). Ethidium

bromide staining of the latter revealed that they contained much less eDNA than the

structured biofilms formed by the wild-type and complemented *tatA* mutant strain

(**Fig. 3A**). Consistent with this biofilm phenotype, exposure of the biofilms formed

by each of the three strains to tobramycin showed that the *tatA* mutant biofilm was

more sensitive to tobramycin than either the wild-type or *tatA*/pTatA mutant biofilm

(**Fig. 3B**).

P. aeruginosa tatA mutants are defective in the production of rhamnolipids,

pyocyanin and MVs

Since rhamnolipids, pyocyanin and MVs are all important components of

P. aeruginosa biofilms, their production was quantified in the *tatA* Tn insertion

mutant and in a Δ*tatABC* deletion mutant. **Fig. 4 (A and B)** shows that both *tat*

mutants produce substantially less pyocyanin and rhamnolipid than the parent or

tatA complemented strain. Furthermore, MV levels (**Fig. 4C**) were reduced by ~50%

in the *tatA* mutant compared with the wild type and could be restored by genetic

complementation.

Inactivation of the tat pathway by mutation or small molecule-mediated

inhibition perturbs PQS signalling

The reductions in eDNA release, rhamnolipids, pyocyanin and MVs noted in the *tat*

mutant as well as its biofilm phenotype are similar to those observed for

P. aeruginosa strains with mutations in *pqs* genes such as *pqsA*, the first gene in

the AQ biosynthetic pathway (see **Figs. 2** and **Fig 4** and [8, 40]). These data

suggested that the *tat* mutant biofilm phenotype was likely, at least in part, to be

due to a defect in PQS signalling.

To investigate the impact of the *tatA* mutation on the expression of *pqsA,* a

CTX::*pqsA′-lux* fusion was introduced into the chromosomal CTX site of both the

wild type and *tatA* mutant. **Fig. 5A** shows that *pqsA* expression in the *tatA* mutant

is reduced ~4 fold compared with the wild type strain and restored by genetic

complementation of the mutant. In agreement with these data, the Tat inhibitor,

Bayer 11-7082, identified by Vasil *et al* [46] reduced *pqsA* expression in the wild type PAO1 strain by ~4 fold consistent with the reduction noted for the CTX::*pqsA-lux* fusion in the *tatA* mutant (**Fig. 5B**). Bayer-11 7082 had no effect on growth or light output in *P. aeruginosa* expressing the *lux* genes from a derepressed *lac* promoter (**S1 Fig.**). In addition, the concentrations of PQS in whole culture extracts of *P. aeruginosa* after growth in LB medium as determined by LC-MS/MS was respectively ~56% lower in the *tatA* mutant compared with the wild type and complemented *tat* mutant (**Fig. 5C**). Since *pqsA* expression and hence AQ production is also PqsR/MvfR-dependent, we compared the expression of *pqsR* in the Δ*tatABC* with the parent strain and found no difference (**S2 Fig.**).

Since mutation of *tatA* results in reduced *pqsA* expression, it is possible that the auto-induction of PQS signalling via PqsR/MvfR is compromised. To uncouple the autoinduction of AQ production, the *pqsABCD* genes were introduced into the *P. aeruginosa* Δ*pqsA* and Δ*pqsA*Δ*tatA* mutants respectively on a plasmid (pBBRMCS5::*pqsABCD*) that constitutively expresses the *pqsABCD* genes [47]. **S3 Fig.** shows that PQS, HHQ and HQNO are present in the culture medium of the Δ*pqsA* mutant transformed with pBBRMCS5::*pqsABCD*. However neither the cell free supernatant nor whole cells of the *tatA* Δ*pqsA* double mutant transformed with pBBRMCS5::*pqsABCD* contained or accumulated intracellular AQs (**S4 Fig.**) suggesting that in the absence of autoinduction in the *tat* mutant background. *P. aeruginosa* is unable to synthesize AQs.

***Exogenous PQS does not restore PQS signalling in a P. aeruginosa* tatA mutant**

QS systems are characteristically autoinducible such that exogenous provision of the cognate signal molecule induces expression of the signal synthase and hence downstream target genes [48]. When the *tatA* mutant was provided with exogenous

PQS, eDNA release did not increase (**Fig. 2**). To investigate this finding further,

either PQS or HHQ was exogenously supplied to the wild type, *tatA* mutant or the

complemented *tatA* mutant strains carrying chromosomal *pqsA'-lux* fusions. The

data presented in **Fig. 6A** show that the response of the *tatA* mutant to PQS or HHQ

respectively at 5 or 20 µM with respect to *pqsA* expression was at least 2-fold lower

than the controls. Since both wild type and *tatA* mutant still produce AQs

endogenously, the experiments were repeated in the *P. aeruginosa* Δ*pqsA* and *tatA*

Δ*pqsA* mutants since no AQs are produced in these genetic backgrounds. **Fig. 6B**

shows that the response to both PQS and HHQ is substantially reduced (e.g. ~4 fold

at 5 µM PQS) in the absence of *tatA* in the *P. aeruginosa tatA* Δ*pqsA* double mutant.

To determine the consequences of perturbed PQS signalling on the expression of the

rhamnolipid (*rhlA*) and pyocyanin biosynthetic genes (*P. aeruginosa* has two, almost

identical redundant 7 gene phenazine biosynthetic operons termed *phzA1-G1* and

phzA2-G2; [49]), the corresponding miniCTX-*lux* promoter fusions for *rhlA* and

phzA1 respectively were constructed and introduced onto the chromosomes of the

wild type, Δ*pqsA* and *tatA* Δ*pqsA* mutants respectively. **Fig. 7A** shows the

expression profiles of *rhlA'-lux* as a function of time. Both the wild type and Δ*pqsA*

mutant show an ~2 fold increase in *rhlA* expression when supplied with exogenous

PQS (20 µM) and share similar profiles over the growth curve. In contrast, the *rhlA'-*

lux fusion in the *tatA* mutant does not show the same expression profile or response

to exogenous PQS as the wild type and Δ*pqsA* mutant strains. The *rhlA'-lux*

expression profile in the *tatA* mutant supplied with exogenous PQS is however

clearly restored when the mutation is complemented by pTatA (**Fig. 7A**).

AQ-dependent QS is required for *phzA1* expression [27,49]. Exogenous PQS

increased *phzA1'-lux* expression by ~4 fold in both wild type and Δ*pqsA* mutant

backgrounds (**Fig. 7B**). However, the *tatA ΔpqsA* double mutant responded

comparatively poorly to PQS (**Fig. 7B**).

313
314
315 ***PqsE restores pyocyanin in the tat mutants***
316
317

Although PqsE is not essential for AQ biosynthesis, it is required for the production

of pyocyanin, rhamnolipids and biofilm maturation and its function is independent

of PQS, HHQ or PqsR [27,50]. Consequently, the expression of certain PQS

signalling-dependent exoproducts such as pyocyanin can be restored in a *pqsA*

negative (and hence AQ-negative) mutant by expressing *pqsE*. To determine

whether expression of *pqsE* alone could restore *phzA1* expression and pyocyanin

production in the *tat* mutants, *pqsE* was introduced on pUCP18. **Fig. 7B** shows that

phzA1 expression can be restored in the *tatA ΔpqsA* mutant in the absence of AQs

while **Fig. 8** confirms that plasmid-borne *pqsE* restores pyocyanin production.

The reduction in rhamnolipid production in the *tat* mutants does not

account for the perturbation of *PQS* signalling

In *P. aeruginosa* biofilms, rhamnolipids provide protective shielding against

neutrophils [51,52] and contribute to the effectiveness of PQS signalling by

enhancing the solubility and bioactivity of PQS [53]. In **Fig. 4B**, we show that

rhamnolipid production is substantially reduced in the *P. aeruginosa tat* mutant

backgrounds. To determine whether the perturbation of PQS signalling in the *tat*

mutants is a consequence of reduced rhamnolipid production, we investigated the

impact of exogenous rhamnolipids on *pqsA* expression. **S5 Fig.** shows that the

addition of purified rhamnolipids (10 or 50 μg/ml) to the *tatA ΔpqsA* mutant with or

without PQS (40 μM) had little effect on *pqsA* expression.

Identification of the Tat substrate responsible for perturbation of *PQS* signalling

Recently Gimenez et al [45] experimentally validated the Tat-mediated export of 34 *P. aeruginosa* gene products predicted to have Tat signal peptides. To determine which of the exported Tat substrates was responsible for perturbation of PQS signalling, allelic replacement mutants were constructed in *P. aeruginosa* strain PA14 for each substrate. Before introducing the *pqsA'-lux* fusion onto the chromosomal CTX site of each Tat substrate mutant, we first confirmed that PQS signalling in a PA14 Δ*tatABC* mutant was perturbed in a similar manner to that observed for PAO1, the genetic background used so far in the study (**Fig. 9A**). Determination of the maximum expression of *pqsA'-lux* for each of the 34 Tat substrate mutants (**Fig. 9A**) revealed that although a number of mutants exhibited reduced light output, the greatest reduction was observed for deletion of PA14_57570 (equivalent to PA4431 in PAO1, here designated *petA* following the nomenclature of orthologues described in other species). This gene codes an iron-sulfur cluster protein, the Rieske subunit of the cytochrome bc_1 complex.

To confirm that the Δ*petA* mutant in common with the *P. aeruginosa tat* mutants exhibited similar defects as a consequence of perturbed PQS signalling, we compared their AQ production (**Fig. 9B**), *rhlA* expression, and eDNA release in biofilms (**S6 Fig.** and **S7 Fig.**). Comparison of PQS, HHQ and HQNO production in PA14 with the Δ*petA* mutant and the strain with a *cis* complementation of the *petA* mutation at the CTX site confirmed the loss and increase in AQ production (**Fig. 9B**). In *P. aeruginosa* PA14, the *rhlA* expression profiles for the parent and Δ*tatABC* deletion mutant in the absence or presence of exogenously supplied PQS were similar to those of strain PAO1 (compare **S6 Fig.** with **Fig. 7A**). Confocal microscopy images of biofilm formation by *P. aeruginosa* PA14 under static growth conditions shows that in common with the Δ*tatABC* mutant, *petA* mutant biofilms lack the eDNA content observed in the wild type and complemented mutant (**S7**

Fig.) demonstrating that PetA is the Tat substrate required for PQS-dependent eDNA release.

Discussion

Extracellular DNA is one of the most abundant components of the extracellular matrix of bacterial biofilms [10-14]. However, the detailed mechanism(s) by which eDNA is released is not fully understood but likely to involve multiple pathways. *P. aeruginosa pqsL* and *pqsA* mutants release more or less eDNA respectively, consistent with the levels of PQS they produce [8]. Furthermore, the biofilms formed by *pqsA* mutants are thin and flat and contain little eDNA consistent with *pqs* signalling playing an important regulatory role in biofilm maturation [8]. To gain further insights into eDNA release, a *P. aeruginosa* transposon mutant library was screened and two groups of mutants exhibiting reduced eDNA release were identified. The first of these contained Tn insertions in genes involved in *pqs* signalling including *pqsC*, *pqsH* and *pqsR*. Mutations in *pqsC* or *pqsR* in common with *pqsA* both abrogate AQ biosynthesis while *pqsH* mutants are unable to produce 3-hydroxy-AQs such as PQS but maintain production of AQs and AQ N-oxides such as HHQ and HQNO respectively [17, 54]. These data suggest that eDNA release is likely to require PQS/HHQ rather than the effector protein PqsE. This is because although *pqsE* mutants in common with *pqsA* mutants form poor biofilms, *pqsE pqsA* double mutants could not be complemented by *pqsE* alone to restore biofilm development [26]. This contrasts with pyocyanin and lectin A for example, as production of both can be restored by PqsE in the absence of AQ biosynthesis [26].

The transposon insertions in the second group of *P. aeruginosa* eDNA release mutants obtained were located in *tatA* and *tatB*, the first two genes in the *tatABC* operon that codes for the twin-arginine translocase secretion pathway [42,43]. The Tat secretion system was originally discovered in the thylakoid membranes of plant

chloroplasts and subsequently identified in bacteria and archaea [42]. Tat exports folded proteins out of the cytoplasm and across the cytoplasmic membrane in an ATP-independent manner [42]. In bacteria such as *E. coli* and *P. aeruginosa* proteins exported by the Tat system are most frequently terminally localized in the periplasm, but some (e.g. phospholipase C) can be transported across the outer membrane by the Xcp type II secretion system, thereby becoming extracellular. Tat signal peptides harbor a conserved S/TRRxFLK consensus motif, where the twin arginine is invariant and in the vast majority of cases essential for export [45]. Using prediction algorithms for Tat substrates, 44 putative *P. aeruginosa* PA14 Tat signal peptides have been identified, and of these, 34 confirmed experimentally using a novel amidase-based Tat reporter [45]. These include phospholipases and proteins involved in pyoverdine-mediated iron-uptake, respiration, osmotic stress defence, motility, and biofilm formation [44, 45]. However, none of these are known to be involved in AQ biosynthesis, transport or *pqs* signal transduction.

In our flow-chamber grown biofilms, clear differences between the *P. aeruginosa* PAO1 wild type and *tatA* mutant biofilms were apparent. While the wild type formed characteristic mushroom-shaped structures under these conditions, the *tatA* mutant formed flat, thin biofilms. In addition, eDNA was observed primarily in the stalks of the mushroom-shaped structures in the wild-type biofilms, whereas the *tatA* mutant biofilms contained no stalks and little extracellular DNA. Moreover, the biofilms formed by the *tatA* mutant were more sensitive to tobramycin treatment than wild-type biofilms, presumably in part due to the inability of the *tatA* mutant to release DNA and form differentiated multicellular structures. Indeed, DNA binds positively charged antibiotics and can act as a 'shield' against aminoglycosides such as tobramycin [53]. In *P. aeruginosa* biofilms, exogenously provided DNA integrates into *P. aeruginosa* biofilms increasing their tolerance toward aminoglycosides [55]. The formation of flat, eDNA deficient, tobramycin-sensitive biofilms and the reduction in rhamnolipid, pyocyanin and MV production by the *P. aeruginosa tatA*

mutant are consistent with the *pqsA* mutant biofilm phenotype [8] and imply the

existence of a link between Tat export and *pqs* signalling. Since transcriptomic

studies of PQS signalling in *P. aeruginosa* have not provided any evidence that the

Tat export system is QS controlled [17,26], we considered it likely that mutation of

the *tat* genes resulted in perturbed PQS signalling. This observation could also

account, at least in part, for the reduced virulence of *P. aeruginosa tat* mutants in a

rat pulmonary infection model [44].

To explore this hypothesis, the expression of *pqsA* and the concentrations of PQS

and HHQ produced were compared in *P. aeruginosa* PAO1 wild type, *tatA* and

complemented *tatA* mutant strains. Similar results were also obtained for

P. aeruginosa strain PA14. Disruption of the *tat* operon had a pronounced effect on

the expression of *pqsA,* with a nearly 3–fold reduction in *pqsA* expression. Moreover,

addition of the Tat inhibitor, BAY 11-7082 [46] also resulted in a comparable

reduction in *pqsA* expression. LC-MS/MS quantification of PQS levels in early

stationary phase after 8 h growth, showed a ~50% reduction, thus correlating with

the *pqsA′-lux* promoter fusion assays. Although PQS and HHQ both serve as PqsR

ligands for driving the positive feedback loop for *pqsABCDE* expression that is central

to AQ signalling [28-30], exogenous provision of neither PQS nor HHQ could fully

restore *pqsA* expression in the *tatA* mutant. Since this mutant still produces AQs,

albeit at a reduced level, a *pqsA* mutation was introduced into the *tatA* mutant

background to examine the response to exogenous AQs in an AQ-negative genetic

background (*tatA* Δ*pqsA*). The data obtained confirmed that the *P. aeruginosa tatA*

Δ*pqsA* mutant responded only weakly to exogenously added PQS or HHQ, albeit to

a much lesser extent than the isogenic Δ*pqsA* mutant with an intact *tat* operon.

These data suggest that disruption of *tat* impairs the ability of *P. aeruginosa* to fully

induce AQ-dependent QS. The consequences of *tat*-dependent perturbation of *pqs*

signalling are clearly evident in the modified expression profile of the *rhlA′-lux* fusion

in the *tatA* mutant with or without exogenously supplied PQS. Similar results were

also obtained for *phzA1* expression in that the reduction in expression observed in

a *tatA* Δ*pqsA* mutant background was not restored to wild type levels by exogenous

PQS.

Since pyocyanin can be produced in the absence of PQS by ectopic expression of

pqsE [26], it was possible that the *tat* phenotype is either a consequence of the

inability of PQS/HHQ to activate the *pqsABCDE* operon via PqsR to generate

sufficient PqsE protein or because the activity of PqsE depends on a functional Tat

system. To investigate the impact of PqsE on pyocyanin production in the PAO1 *tatA*

mutants, a plasmid borne copy of *pqsE* was introduced into the *tat* mutants.

Pyocyanin production was fully restored suggesting that the *tat* mutant phenotype

is not due to the inability of PqsE to function but rather a failure of the *pqs* auto-

induction circuitry to produce sufficient PQS/HHQ to efficiently activate *pqsABCDE*

transcription and hence *pqsE* expression.

AQs such as PQS are extremely hydrophobic and virtually insoluble in aqueous

environments. Hence PQS signalling can be enhanced by increasing the solubility

and delivery of PQS. Calfee *et al* [53] showed that rhamnolipids increase the

solubility and bioactivity of PQS in *P. aeruginosa* with respect to *lasB* expression.

Since rhamnolipid production is substantially reduced in the *P. aeruginosa tat*

mutants, we added back purified rhamnolipids to the *tatA* Δ*pqsA* mutant. However,

no effect on *pqsA* expression was observed indicating that the PQS signalling defect

in the *tat* mutants is not simply due to the loss of rhamnolipid production and an

inability to solubilize PQS.

In *Escherichia coli,* inactivation of *tat* leads to a characteristic cell envelope defect

because of the mis-localization of two amidases involved in cell wall metabolism.

This results in the formation of bacterial chains, leakage of periplasmic proteins and

enhanced susceptibility to detergents [56]. Since PQS is a signal molecule that must

be transported across the cell envelope in both directions, it was possible that an

envelope defect in *P. aeruginosa tat* mutants impacts on the PQS signalling auto-

induction cascade. However, Ball *et al* [57] showed that *P. aeruginosa tat* mutants

do not show the same Tat-dependent envelope defects found in *E. coli.* In addition,

we found no evidence for intracellular accumulation of AQs indicative of a transport

defect.

To uncover the mechanistic link between Tat and PQS signalling, we sought to

determine whether mutation of *tat* resulted in the inability to export a specific Tat

substrate, the loss of which resulted in perturbation of PQS autoinduction and hence

the expression of PQS-dependent genes and the formation of defective biofilms. By

screening 34 validated Tat substrate mutants for reduced *pqsA* expression, we

identified the *petA* gene as responsible for the defect in *pqs*-dependent QS. This

gene codes for the Rieske protein, an iron-sulfur cluster protein sub-unit component

of the cytochrome bc_1 complex involved in electron transfer and respiration under

aerobic conditions and also essential for enabling *P. aeruginosa* to grow under

anaerobic conditions in the presence of nitrite [58]. Self-poisoning of cytochrome

bc_1 occurs via HQNO which is produced via the same biosynthetic pathway as PQS

(**Fig. 1**). HQNO binds to the quinone reduction (Qi) site of the respiratory

cytochrome bc_1 complex [59]. This results in the generation of reactive oxygen

species that cause *P. aeruginosa* cell death and autolysis favouring biofilm formation

and antibiotic tolerance. Rieske subunit and cytochrome b_1 mutants do not undergo

autolysis and are insensitive to exogenous HQNO [59]. Similarly, *P. aeruginosa pqsL*

mutants that are unable to produce HQNO also fail to undergo autolysis [59]. Our

data also show that in common with the *tat* mutants, the *petA* mutant produces

reduced levels of HQNO (as well as PQS and HHQ). This suggests that lack of eDNA

in the *tat* mutant biofilms is because the cells do not undergo limited autolysis as

they lack the self-poisoning mechanism that depends on both an intact cytochrome

bc_1 and sufficient HQNO. Thus, lack of the Rieske sub-unit export is clearly

responsible for the Tat-mediated perturbation of PQS-dependent QS, the loss of

virulence factor production, biofilm eDNA and the tobramycin tolerance of

P. aeruginosa biofilms. Given the importance of PQS signalling and the Tat system

to virulence and biofilm maturation in *P. aeruginosa*, our findings underline the

potential of the Tat system as a drug target for novel antimicrobial agents.

<h1 align="center">Materials and Methods</h1>

Bacterial strains and growth conditions

The *P. aeruginosa* and *E. coli* strains used are listed in **S1 Table** and were grown in

LB or ABTG [8] at 37^{o}C unless otherwise stated. *P. aeruginosa* biofilms were

cultivated at 30^{o}C in flow-chambers irrigated with FAB medium [60] supplemented

with 0.3 mM glucose or in M9 medium with succinate (for static biofilms). Selective

media were supplemented with ampicillin (Ap; 100 µg ml^{-1}), gentamicin (Gm; 60 µg

ml^{-1}), or streptomycin (Sm; 100 µg ml^{-1}). PQS, HHQ and 2-heptyl-4-

hydroxyquinoline N-oxide (HQNO) were synthesized and characterized in house as

described before [19,30] and dissolved in methanol before being added to growth

media at the appropriate concentration.

Mutant construction, screening and validation

The *P. aeruginosa* PAO1 Tn mutant library was constructed using the Mariner

transposon vector pBT20 as previously described [61]. Transconjugants carrying Tn

insertions were picked from the selective plates and inoculated into microtiter plates

containing ABTG medium [62] supplemented with propidium iodide (PI) and the

level of red fluorescence quantified. Mutants producing reduced levels of eDNA were

selected and the sequences flanking the Tn insertion identified by arbitrary PCR

essentially as described by Friedman and Kolter [63] but using the specific primers

TnM1 and TnM2 (**S2 Table**). DNA Sequencing was performed by Macrogen, Seoul,

Korea with primer TnMseq (**S2 Table**). An in-frame *P. aeruginosa tatABC* deletion

mutant was generated by allelic exchange using the oligonucleotide primers Tat3D-

UF, Tat3D-UR, Tat3D-DF and Tat3D-DR (**S2 Table**) to introduce the up- and down-

stream regions of the *tatABC* locus into the suicide vector pME3087 to generate

pME3087::*tatABC* (**S1 Table**). The latter was introduced into *P. aeruginosa* via

conjugation with *E. coli* S17-1 λpir followed by enrichment for tetracycline-sensitive

cells as described by Ye *et al* [64]. The Δ*tatABC* deletion in *P. aeruginosa* PAO1 was

confirmed by PCR and sequence analysis. To generate the PA14 Δ*petA* mutant,

500 bp upstream and downstream of the gene were amplified using respectively

primers S.PA4431upFor/S.PA4431upRev and S.PA4431downFor/S.PA4431downRev

listed in **S2 Table**. The PCR product was cloned in pKNG101 suicide vector by one-

step sequence and ligation-independent cloning (SLIC) [65] which was then

sequenced. The resulting plasmid, pKNGΔ*petA*, maintained in the *E. coli* CC118λpir

strain, was then mobilized in *P. aeruginosa* strains. The mutant, in which the double

recombination events occurred, was confirmed by PCR analysis. A similar strategy

was used to construct allelic replacement mutants for each of 34 validated Tat

substrates [45]. These strains are summarized in **S3 Table** and their validation will

be described in detail elsewhere. For the generation of the *petA* cis-complemented

strain, PA14Δ*petA::(CTX1*::*petA)*, the *petA* genes along with a 500 bp fragment

corresponding to the putative promoter region for the *petA* gene were PCR amplified

using S-4431CTXFor/S-4431CTXRev and cloned by SLIC into the mini-CTX1 vector

yielding pminiCTX1-*petA*. Transfer of this plasmid in *P. aeruginosa* Δ*petA* strain was

carried out by triparental mating using the conjugative properties of the helper

plasmid pRK2013. The recombinant clones containing the mini-CTX inserted at the

attB locus on the *P. aeruginosa* genome were selected on tetracycline-containing

PIA generating PA14Δ*petA::(attB::petA)*.

Construction of a *tatA* complementation plasmid

The *tatA* gene was amplified by PCR using primers F*tatA* and R*tatA* (**S2 Table**),

introduced into pUCP22 (**S1 Table**) and electroporated into the *P. aeruginosa tatA*

and *tatA ΔpqsA* mutants. Transformants were selected on LB plates containing 200

µg ml^{-1} carbenicillin

Bioluminescent reporter gene fusion assays

To investigate the impact of the *tat* mutation and Tat inhibitor Bayer 11-7082 [46]

on PQS signalling, transcriptional fusions between the promoter regions of *pqsA*,

pqsR, *rhlA*, *phzA1 and phzA2* and the *luxCDABE* operon were constructed using the

miniCTX-*lux* plasmid as previously described [17]. In addition, a constitutively

bioluminescent reporter using a miniCTX::*tac-luxCDABE* promoter fusion was

constructed as a control for Bayer 11-7082. Bioluminescence as a function of

bacterial growth was quantified in 96 well plates using a combined luminometer-

spectrometer.

Cultivation of biofilms

Biofilms were grown in flow-chambers with individual channel dimensions of 1 x 4 x

40 mm as described previously [66]. One hour after inoculation, with bacteria, the

growth medium flow (0.2 mm/s corresponding to laminar flow with a Reynolds

number of 0.02) was started. When required, eDNA in biofilms was stained with 1

µM ethidium bromide prior to microscopy. Tobramycin (10 µM) was added to the

biofilm medium after 4 days of cultivation. After 24 h of tobramycin treatment,

propidium iodide (10 µM) was added to the flow cells to visualize the dead cells via

confocal laser scanning microscopy. *P. aeruginosa* PA14 and *petA* mutant biofilms

were grown under static conditions over 48 h at 37 °C on glass slides (Ibidi)

incorporating 300 µl chambers. After 48 h incubation, spent medium was removed

and the biofilm eDNA stained with YoYo-1 (40 µM).

Microscopy and image processing of flow cell biofilms

All images of flow-chamber-grown and static biofilms were captured with a confocal

laser scanning microscope (CLSM) equipped with detectors and filter sets for

monitoring green fluorescent protein, Syto9, propidium iodide, and ethidium

bromide. Images were obtained using a 63x/1.4 objective or a 40x/1.3 objective.

Simulated 3-D images and sections were generated using the IMARIS software

package (Bitplane AG, Zürich, Switzerland).

AQ, pyocyanin, rhamnolipid and MV analysis

The AQs (PQS, HHQ and HQNO) were quantified by LC-MS/MS after extracting cell

free supernatants or whole bacterial cells in triplicate with acidified ethyl acetate or

methanol respectively as described by Ortori *et al* [19]. Pyocyanin was extracted

with chloroform and quantified spectrophotometrically [26]. Rhamnolipids were

quantified indirectly using the orcinol method [35]. For PQS solubilization

experiments, rhamnolipids were purified as described by Muller *et al* [67]. MVs were

harvested by ultracentrifugation, the pellets resuspended in 10mM HEPES buffer and

the lipid content quantified using FM4-64 essentially as described previously [68].

MV production was normalized by dividing the lipid fluorescence units by CFU values

determined by dilution plating. Assays were performed in triplicate at least twice.

Statistical Analysis.

Significance for differences between wild type and isogenic mutants was determined

by two-tailed *t*-tests where ****$p < 0.001$, ***$p < 0.001$, **$p < 0.01$, and *$p <$

0.05 and n.s., not significant.

S1 Fig. The Tat inhibitor Bayer 11-7082 has no effect on light output at 20 or 40 μM from *P. aeruginosa* CTX:*ptac'-luxCDABE* chromosomal reporter fusion. The co-solvent DMSO, had no effect at 0.8% on the *lux* reporter fusion either. Data are presented as maximal light output as a function of growth (RLU/OD$_{600}$). Experiments were repeated in triplicate at least twice.

S2 Fig. Deletion of the *P. aeruginosa tatABC* genes does not influence *pqsR* expression. The data show that there are no differences in the expression of a chromosomal CTX::*pqsR'-luxCDABE* fusion in the *P. aeruginosa* wild type compared with the Δ*tatABC* mutant Data are presented as maximal light output as a function of growth (RLU/OD$_{600}$). Experiments were repeated in triplicate at least twice.

S3 Fig. AQ biosynthesis is not restored in a *tatA* Δ*pqsA* double mutant by plasmid-borne *pqsABCD* in the absence of autoinduction. Semi-quantitative analysis by LC-MS/MS of PQS, HHQ and HQNO production by *P. aeruginosa pqsA* and *tatA* Δ*pqsA* mutants respectively without or with the *pqsABCD* biosynthetic genes provided *in trans* via pBBRMCS::*pqsABCD.* Experiments were repeated in triplicate at least twice.

S4 Fig.   HHQ and PQS do not accumulate intracellularly a *tatA* Δ*pqsA* double mutant harboring the   plasmid-borne *pqsABCD* genes in the absence of autoinduction*.* Semi-quantitative analysis by LC-MS/MS of PQS (**A**) and HHQ (**B**) extracted from whole cells of *P. aeruginosa* wild type and the *tatA* Δ*pqsA* mutant without or with the *pqsABCD* biosynthetic genes provided via pBBRMCS::*pqsABCD.* Cells were harvested at 8 h and 16 h respectively*.* Experiments were repeated in triplicate

S5 Fig. Exogenous rhamnolipids do not enhance PQS-dependent expression of

pqsA in a *tatA ΔpqsA* mutant*.* PQS (40 μM) was added with or without purified

rhamnolipids (50 μg/ml) to a *pqsA* mutant or a *tatA ΔpqsA* mutant carrying

chromosomal *pqsA'-lux* fusions. Maximal light output as a function of growth

(RLU/OD$_{600}$) is presented. Experiments were repeated in triplicate at least twice.

S6 Fig Rhamnolipid biosynthesis gene *rhlA* shows altered expression profiles in *P.*

aeruginosa PA14 *ΔtatABC* and *ΔpetA* mutants compared with wild type and fail to

respond to exogenous PQS. Bioluminescence from a chromosomal *rhlA'-lux* fusion

as a function of growth (RLU/OD) over time when introduced into (**A**) the PA14 wild

type, (**B**) *ΔtatABC* and (**C**) *ΔpetA* mutants in the absence or presence of exogenous

PQS (20 μM).

S7 Fig Deletion of *petA* in *P. aeruginosa* PA14 results in a reduction in the eDNA

content of biofilms.   Biofilms of wild type PA14, the *ΔtatABC* and *ΔpetA* mutants

and the genetically complemented PA14 mutant (*ΔpetA*::(CTX::*petA*)) were grown

cultured statically and stained for eDNA with YOYO-1. (**A**) CLSM images and (**B**)

eDNA quantification. Experiments were repeated in triplicate at least twice. ***p <

0.001, **p < 0.01.

S1 Table Strains and plasmids used in this study

S2 Table Oligonucleotide primers used in this study

S3 Table *P. aeruginosa* PA14 Tat substrate mutants used in this study

Acknowledgements

We thank Alex Truman for synthesis of AQ standards. This work was funded via

grants to PW and MC from the Biotechnology and Biological Sciences Research

Council, UK (BB/F014392/1; www.bbsrc.ac.uk), the Medical Research Council

U.K.(MR/N501852/1; www.mrc.ac.uk), the Wellcome Trust (103884/Z/14/Z;

www.welcome.ac.uk), the European Union FP7 collaborative action grant

(NABATIVI, Ref. 223670; http://cordis.europa.eu/ project/rcn/90762_en.html).

MC, PW and KH are partly funded by the National Biofilms Innovation Centre (NBIC)

which is an Innovation and Knowledge Centre funded by the BBSRC, InnovateUK

and Hartree Centre (Award Number BB/R012415/1). FA was funded via a Wellcome

Trust Doctoral Training Programme (108876/Z/15/Z). BI was funded by a grant

from Vaincre la Mucoviscidose and the Grégory Lemarchal

associations (RF20140501138) and recurrent funding from the CNRS and Aix-

Marseille Université. LY, MG and TTN were funded by grants from the Danish

Strategic Research Council, the Danish Council for Independent Research, the Novo

Nordisk Foundation and the Lundbeck Foundation. MRG was supported by a PhD

studentship and ATER position from Aix-Marseille Université.

References
691

1. Moradali MF, Ghods S, Rehm BH. *Pseudomonas aeruginosa* Lifestyle: A

paradigm for adaptation, survival, and persistence. Front Cell Infect Microbiol.

2017; 7:39-42. https://doi: 10.3389/fcimb.2017.00039. PMID: 28261568

2. Costerton JW, Stewart PS, Greenberg EP. Bacterial biofilms: a common cause of

persistent infections. Science. 1999; 284:1318-22.

https://doi: 10.1126/science.284.5418.1318. PMID: 10334980.

3. Klausen M, Aaes-Jørgensen A, Molin S, Tolker-Nielsen T. Involvement of

bacterial migration in the development of complex multicellular structures in

Pseudomonas aeruginosa biofilms. Mol Microbiol. 2003; 50:61-8.

https://doi: 10.1046/j.1365-2958.2003.03677.x. PMID: 14507363.

4. Klausen M, Heydorn A, Ragas P, Lambertsen L, Aaes-Jørgensen A, Molin S,

Tolker-Nielsen T. Biofilm formation by *Pseudomonas aeruginosa* wild type, flagella

and type IV pili mutants. Mol Microbiol. 2003; 48:1511-24.

https://doi: 10.1046/j.1365-2958.2003.03525.x. PMID: 12791135.

5. Pamp SJ, Tolker-Nielsen T. Multiple roles of biosurfactants in structural biofilm

development by *Pseudomonas aeruginosa*. J Bacteriol. 2007; 189:2531-39.

https://doi: 10.1128/JB.01515-06. PMID: 17220224

6. Harmsen M, Yang L, Pamp SJ, Tolker-Nielsen T. An update on *Pseudomonas*

aeruginosa biofilm formation, tolerance, and dispersal. FEMS Immunol Med

Microbiol. 2010; 59:253-68. https://doi: 10.1111/j.1574-695X.2010.00690.x.

PMID: 20497222.

7. Yang L, Hu Y, Liu Y, Zhang J, Ulstrup J, Molin S. Distinct roles of extracellular

polymeric substances in *Pseudomonas aeruginosa* biofilm development. Environ

Microbiol. 2011; 13:1705-17. https://doi: 10.1111/j.1462-2920.2011.02503.x.

PMID: 21605307.

8. Allesen-Holm M, Barken KB, Yang L, Klausen M, Webb JS, Kjelleberg S, Molin S,

Givskov M, Tolker-Nielsen T. A characterization of DNA release in *Pseudomonas*

aeruginosa cultures and biofilms. Mol Microbiol. 2006; 59:1114-28.

https://doi: 10.1111/j.1365-2958.2005.05008.x. PMID: 16430688.

9. Tashiro Y, Uchiyama H, Nomura N. Multifunctional membrane vesicles in

Pseudomonas aeruginosa. Environ Microbiol. 2012; 14:1349-62.

https://doi: 10.1111/j.1462-2920.2011.02632.x. PMID: 22103313.

10. Tolker-Nielsen T. Biofilm Development. Microbiol Spectr. 2015 3:MB-0001-

2014. https://doi: 10.1128/microbiolspec.MB-0001-2014. PMID: 26104692.

11. Okshevsky M, Meyer RL. The role of extracellular DNA in the establishment,

maintenance and perpetuation of bacterial biofilms. Crit Rev Microbiol.

2015;41:341-52. https://doi: 10.3109/1040841X.2013.841639. PMID: 24303798.

12. Das T, Sehar S, Manefield M. The roles of extracellular DNA in the structural

integrity of extracellular polymeric substance and bacterial biofilm development.

Environ Microbiol Rep. 2013 5:778-86.

https://doi: 10.1111/1758-2229.12085. PMID: 24249286.

13. Wilton M, Charron-Mazenod L, Moore R, Lewenza S. Extracellular DNA acidifies

biofilms and induces aminoglycoside resistance in *Pseudomonas aeruginosa*.

Antimicrob Agents Chemother. 2015; 60:544-53.

https://doi: 10.1128/AAC.01650-15. PMID: 26552982

14. Turnbull L, Toyofuku M, Hynen AL, Kurosawa M, Pessi G, Petty NK, Osvath SR,

Cárcamo-Oyarce G, Gloag ES, Shimoni R, Omasits U, Ito S, Yap X, Monahan LG,

Cavaliere R, Ahrens CH, Charles IG, Nomura N, Eberl L, Whitchurch CB. Explosive

cell lysis as a mechanism for the biogenesis of bacterial membrane vesicles and

biofilms. Nat Commun. 2016; 7:11220.

https://doi: 10.1038/ncomms11220. PMID: 27075392.

15. Das T, Kutty SK, Tavallaie R, Ibugo AI, Panchompoo J, Sehar S, Aldous L,

Yeung AW, Thomas SR, Kumar N, Gooding JJ, Manefield M. Phenazine virulence

factor binding to extracellular DNA is important for *Pseudomonas aeruginosa*

biofilm formation. Sci Rep. 2015; 5:8398. https://doi: 10.1038/srep08398. PMID:

25669133.

16. Williams P, Cámara M. Quorum sensing and environmental adaptation in

Pseudomonas aeruginosa: a tale of regulatory networks and multifunctional signal

molecules. Curr Opin Microbiol. 2009; 12:182-91.

https://doi: 10.1016/j.mib.2009.01.005. PMID: 19249239.

17. Rampioni G, Falcone M, Heeb S, Frangipani E, Fletcher MP, Dubern JF, Visca P,

Leoni L, Cámara M, Williams P. Unravelling the genome-wide contributions of

specific 2-alkyl-4-quinolones and PqsE to quorum sensing in *Pseudomonas*

aeruginosa. PLoS Pathog. 2016; 12:e1006029.

https://doi: 10.1371/journal.ppat.1006029. PMID: 27851827.

18. Lépine F, Milot S, Déziel E, He J, Rahme LG. Electrospray/mass spectrometric

identification and analysis of 4-hydroxy-2-alkylquinolines (HAQs) produced by

Pseudomonas aeruginosa. J Am Soc Mass Spectrom. 2004; 15:862-9. https://doi:

10.1016/j.jasms.2004.02.012. PMID: 15144975.

19. Ortori CA, Dubern JF, Chhabra SR, Cámara M, Hardie K, Williams P, Barrett

DA. Simultaneous quantitative profiling of *N*-acyl-L-homoserine lactone and 2-

alkyl-4(1*H*)-quinolone families of quorum-sensing signaling molecules using LC-

MS/MS. Anal Bioanal Chem. 2011; 399:839-50.

https://doi: 10.1007/s00216-010-4341-0. PMID: 21046079.

20. Coleman JP, Hudson LL, McKnight SL, Farrow JM 3rd, Calfee MW, Lindsey CA,
Pesci EC. *Pseudomonas aeruginosa* PqsA is an anthranilate-coenzyme A ligase. J
Bacteriol. 2008; 190:1247-55. https://doi: 10.1128/JB.01140-07.
PMID: 18083812
789
790 **21**. Zhang YM, Frank MW, Zhu K, Mayasundari A, Rock CO. PqsD is responsible for
791 the synthesis of 2,4-dihydroxyquinoline, an extracellular metabolite produced by
792 *Pseudomonas aeruginosa*. J Biol Chem. 2008; 283:28788-94.
793 https://doi: 10.1074/jbc.M804555200. PMID: 18728009.
794
795 **22**. Drees SL, Fetzner S. PqsE of *Pseudomonas aeruginosa* acts as pathway-
796 specific thioesterase in the biosynthesis of alkylquinolone signaling molecules.
797 Chem Biol. 2015; 22:611-8. https://doi: 10.1016/j.chembiol.2015.04.012. PMID:
798 25960261.
799
800 **23**. Dulcey CE, Dekimpe V, Fauvelle DA, Milot S, Groleau MC, Doucet N, Rahme
801 LG, Lépine F, Déziel E. The end of an old hypothesis: the pseudomonas signaling
802 molecules 4-hydroxy-2-alkylquinolines derive from fatty acids, not 3-keto fatty
803 acids. Chem Biol. 2013; 20:1481-91.
804 https://doi: 10.1016/j.chembiol.2013.09.021. PMID: 24239007
805
806 **24**. Drees SL, Li C, Prasetya F, Saleem M, Dreveny I, Williams P, Hennecke U,
807 Emsley J, Fetzner S. PqsBC, a condensing enzyme in the biosynthesis of the
808 *Pseudomonas aeruginosa* quinolone signal: crystal structure, inhibition, and
809 reaction mechanism. J Biol Chem. 2016; 291:6610-24.
810 https://doi: 10.1074/jbc.M115.708453. PMID: 26811339;
811
812 **25**. Schertzer JW, Brown SA, Whiteley M. Oxygen levels rapidly modulate
813 *Pseudomonas aeruginosa* social behaviours via substrate limitation of PqsH. Mol
814 Microbiol. 2010; 77:1527-38.
815 https://doi: 10.1111/j.1365-2958.2010.07303.x. PubMed PMID: 20662781.
816
817 **26**. Rampioni G, Pustelny C, Fletcher MP, Wright VJ, Bruce M, Rumbaugh KP, Heeb
818 S, Cámara M, Williams P. Transcriptomic analysis reveals a global alkyl-quinolone-
819 independent regulatory role for PqsE in facilitating the environmental adaptation of
820 *Pseudomonas aeruginosa* to plant and animal hosts. Environ Microbiol. 2010;
821 12:1659-73. https://doi: 10.1111/j.1462-2920.2010.02214.x. PMID: 20406282.
822

27. Groleau, MC, de Oliveira Pereira, T, Dekimpe V, Déziel E. PqsE Is Essential

for RhlR-Dependent Quorum Sensing Regulation in *Pseudomonas aeruginosa*

mSystems. 2020; 5:e00194-20;

https://doi: 10.1128/mSystems.00194-20. PMID: 32457239

28. Wade DS, Calfee MW, Rocha ER, Ling EA, Engstrom E, Coleman JP, Pesci EC.

Regulation of pseudomonas quinolone signal synthesis in *Pseudomonas*

aeruginosa. J Bacteriol. 2005;187:4372-80.

https://doi: 10.1128/JB.187.13.4372-4380.2005. PMID: 15968046.

29. Xiao G, Déziel E, He J, Lépine F, Lesic B, Castonguay MH, Milot S, Tampakaki

AP, Stachel SE, Rahme LG. MvfR, a key *Pseudomonas aeruginosa* pathogenicity

LTTR-class regulatory protein, has dual ligands. Mol Microbiol. 2006; 62:1689-99.

https://doi: 10.1111/j.1365-2958.2006.05462.x. PMID: 17083468.

30. Ilangovan A, Fletcher M, Rampioni G, Pustelny C, Rumbaugh K, Heeb S,

Cámara M, Truman A, Chhabra SR, Emsley J, Williams P. Structural basis for

native agonist and synthetic inhibitor recognition by the *Pseudomonas aeruginosa*

quorum sensing regulator PqsR (MvfR). PLoS Pathog. 2013; 9:e1003508.

https://doi: 10.1371/journal.ppat.1003508. PMID: 23935486.

31. Diggle SP, Matthijs S, Wright VJ, Fletcher MP, Chhabra SR, Lamont IL, Kong X,

Hider RC, Cornelis P, Cámara M, Williams P. The *Pseudomonas aeruginosa* 4-

quinolone signal molecules HHQ and PQS play multifunctional roles in quorum

sensing and iron entrapment. Chem Biol. 2007;14:87-96.

https://doi: 10.1016/j.chembiol.2006.11.014 PMID: 17254955.

32. Häussler S, Becker T. The pseudomonas quinolone signal (PQS) balances life

and death in *Pseudomonas aeruginosa* populations. PLoS Pathog. 2008;

4:e1000166. https://doi: 10.1371/journal.ppat.1000166. PMID: 18818733;

33. Mashburn-Warren L, Howe J, Garidel P, Richter W, Steiniger F, Roessle M,

Brandenburg K, Whiteley M. Interaction of quorum signals with outer membrane

lipids: insights into prokaryotic membrane vesicle formation. Mol Microbiol. 2008

69:491-502.

https://doi: 10.1111/j.1365-2958.2008.06302.x. PMID: 18630345.

34. Mashburn LM, Whiteley M. Membrane vesicles traffic signals and facilitate

group activities in a prokaryote. Nature. 2005; 437:422-5.

https://doi: 10.1038/nature03925 PMID: 16163359

35. Pustelny C, Albers A, Büldt-Karentzopoulos K, Parschat K, Chhabra SR,

Cámara M, Williams P, Fetzner S. Dioxygenase-mediated quenching of quinolone-

dependent quorum sensing in *Pseudomonas aeruginosa*. Chem Biol. 2009;

16:1259-67. https://doi: 10.1016/j.chembiol.2009.11.013. PMID: 20064436.

36. D'Argenio DA, Calfee MW, Rainey PB, Pesci EC. Autolysis and autoaggregation

in *Pseudomonas aeruginosa* colony morphology mutants. J Bacteriol. 2002;

184:6481-9. https://doi: 10.1128/jb.184.23.6481-6489.2002. PMID: 12426335.

37. Webb JS, Thompson LS, James S, Charlton T, Tolker-Nielsen T, Koch B,

Givskov M, Kjelleberg S. Cell death in *Pseudomonas aeruginosa* biofilm

development. J Bacteriol. 2003; 185:4585-92.

https://doi: 10.1128/jb.185.15.4585-4592.2003 PMID: 12867469

38. Spoering AL, Gilmore MS. Quorum sensing and DNA release in bacterial

biofilms. Curr Opin Microbiol. 2006; 9:133-7.

https://doi: 10.1016/j.mib.2006.02.004. PMID: 16529982.

39. Yang L, Barken KB, Skindersoe ME, Christensen AB, Givskov M, Tolker-Nielsen

T. Effects of iron on DNA release and biofilm development by *Pseudomonas*

aeruginosa. Microbiology. 2007;153:1318-28.

https://doi: 10.1099/mic.0.2006/004911-0. PMID: 17464046.

40. Yang L, Nilsson M, Gjermansen M, Givskov M, Tolker-Nielsen T. Pyoverdine

and PQS mediated subpopulation interactions involved in *Pseudomonas aeruginosa*

biofilm formation. Mol Microbiol. 2009;74:1380-92. https://doi: 10.1111/j.1365-

2958.2009.06934.x. PMID: 19889094.

41. Suzuki T, Fujikura K, Higashiyama T, Takata K. DNA staining for fluorescence

and laser confocal microscopy. J Histochem Cytochem. 1997; 45:49-53.

https://doi: 10.1177/002215549704500107. PMID: 9010468.

42. Patel R, Smith SM, Robinson C. Protein transport by the bacterial Tat pathway.

Biochim Biophys Acta. 2014; 1843:1620-8.

https://doi: 10.1016/j.bbamcr.2014.02.013. PMID: 24583120.

43. Costa TR, Felisberto-Rodrigues C, Meir A, Prevost MS, Redzej A, Trokter M,

Waksman G. Secretion systems in Gram-negative bacteria: structural and

mechanistic insights. Nat Rev Microbiol. 2015; 13:343-59.

https://doi: 10.1038/nrmicro3456. PMID: 25978706.

44. Ochsner UA, Snyder A, Vasil AI, Vasil ML. Effects of the twin-arginine

translocase on secretion of virulence factors, stress response, and pathogenesis.

Proc Natl Acad Sci U S A. 2002; 99:8312-7.

https://doi: 10.1073/pnas.082238299. PMID: 12034867.

45. Gimenez MR, Chandra G, Van Overvelt P, Voulhoux R, Ize, B, Bleves, S.

Genome wide identification and experimental validation of *Pseudomonas*

aeruginosa Tat substrates. Sci Rep 2018; 11950.

https://doi: 10.1038/s41598-018-30393-x. PMID: 30093651

46. Vasil ML, Tomaras AP, Pritchard AE. Identification and evaluation of twin-

arginine translocase inhibitors. Antimicrob Agents Chemother. 2012; 56:6223-34.

https://doi: 10.1128/AAC.01575-12. PMID: 23006747

47.Niewerth H, Bergander K, Chhabra SR, Williams P, Fetzner S. Synthesis and

biotransformation of 2-alkyl-4(1H)-quinolones by recombinant *Pseudomonas*

putida KT2440. Appl Microbiol Biotechnol. 2011; 91:1399-408.

https://doi: 10.1007/s00253-011-3378-0. PMID: 21670979.

48. Williams P, Winzer K, Chan WC, Cámara M. Look who's talking:

communication and quorum sensing in the bacterial world. Philos Trans R Soc Lond

B Biol Sci. 2007; 362:1119-34.

https://doi: 10.1098/rstb.2007.2039. PMID: 17360280;

49. Recinos DA, Sekedat MD, Hernandez A, Cohen TS, Sakhtah H, Prince AS,

Price-Whelan A, Dietrich LE. Redundant phenazine operons in *Pseudomonas*

aeruginosa exhibit environment-dependent expression and differential roles in

pathogenicity. Proc Natl Acad Sci U S A. 2012; 109:19420-5.

https://doi: 10.1073/pnas.1213901109. PMID: 23129634

50. Farrow JM 3rd, Sund ZM, Ellison ML, Wade DS, Coleman JP, Pesci EC. PqsE
functions independently of PqsR-Pseudomonas quinolone signal and enhances the
rhl quorum-sensing system. J Bacteriol. 2008; 190:7043-51.
https://doi: 10.1128/JB.00753-08. PMID: 18776012
939

51. Van Gennip M, Christensen LD, Alhede M, Phipps R, Jensen PØ, Christophersen
L, Pamp SJ, Moser C, Mikkelsen PJ, Koh AY, Tolker-Nielsen T, Pier GB, Høiby N,
Givskov M, Bjarnsholt T. Inactivation of the *rhlA* gene in *Pseudomonas aeruginosa*
prevents rhamnolipid production, disabling the protection against
polymorphonuclear leukocytes. APMIS. 2009 117:537-46.
https://doi: 10.1111/j.1600-0463.2009.02466.x. PMID: 19594494
946

947 **52**. Alhede M, Bjarnsholt T, Jensen PØ, Phipps RK, Moser C, Christophersen L,
948 Christensen LD, van Gennip M, Parsek M, Høiby N, Rasmussen TB, Givskov M.
949 *Pseudomonas aeruginosa* recognizes and responds aggressively to the presence of
950 polymorphonuclear leukocytes. Microbiology (Reading). 2009 155:3500-3508.
951 https://doi: 10.1099/mic.0.031443-0. PMID: 19643762.
952

953 **53**. Calfee MW, Shelton JG, McCubrey JA, Pesci EC. Solubility and bioactivity of the
954 Pseudomonas quinolone signal are increased by a *Pseudomonas aeruginosa*-
955 produced surfactant. Infect Immun. 2005; 73:878-82.
956 https://doi: 10.1128/IAI.73.2.878-882.2005. PMID: 15664929.
957

958 **54**. Déziel E, Lépine F, Milot S, He J, Mindrinos MN, Tompkins RG, Rahme LG.
959 Analysis of *Pseudomonas aeruginosa* 4-hydroxy-2-alkylquinolines (HAQs) reveals a
960 role for 4-hydroxy-2-heptylquinoline in cell-to-cell communication. Proc Natl Acad
961 Sci U S A. 2004; 101:1339-44.
962 https://doi: 10.1073/pnas.0307694100. PMID: 14739337
963

964 **55**. Chiang WC, Nilsson M, Jensen PØ, Høiby N, Nielsen TE, Givskov M, Tolker-
965 Nielsen T. Extracellular DNA shields against aminoglycosides in *Pseudomonas*
966 *aeruginosa* biofilms. Antimicrob Agents Chemother. 2013; 57:2352-61.
967 https://doi: 10.1128/AAC.00001-13. PMID: 23478967
968

969 **56**. Ize B, Stanley NR, Buchanan G, Palmer T. Role of the *Escherichia coli* Tat
970 pathway in outer membrane integrity. Mol Microbiol. 2003; 48:1183-93.
971 https://doi: 10.1046/j.1365-2958.2003.03504.x.PMID: 12787348.
972

57. Ball G, Antelmann H, Imbert PR, Gimenez MR, Voulhoux R, Ize B. Contribution
of the twin arginine translocation system to the exoproteome of *Pseudomonas*
aeruginosa. Sci Rep. 2016; 6:27675.
https://doi: 10.1038/srep27675. PMID: 27279369

58. Hasegawa N, Arai H, Igarashi Y. Need for Cytochrome bc 1 Complex for
dissimilatory nitrite reduction of *Pseudomonas aeruginosa*. Biosci Biotechnol
Biochem. 2003 67:121-126.
https://doi: 10.1271/bbb.67.121 PMID: 12619683.

59. Hazan R, Que YA, Maura D, Strobel B, Majcherczyk PA, Hopper LR, Wilbur DJ,
Hreha TN, Barquera B, Rahme LG. Auto poisoning of the respiratory chain by a
quorum-sensing-regulated molecule favors biofilm formation and antibiotic
tolerance. Curr Biol. 2016; 26:195-206.
https://doi: 10.1016/j.cub.2015.11.056. PMID: 26776731

60. Heydorn A, Nielsen AT, Hentzer M, Sternberg C, Givskov M, Ersbøll BK, Molin
S. Quantification of biofilm structures by the novel computer program COMSTAT.
Microbiology. 2000;146:2395-407
https://doi: 10.1099/00221287-146-10-2395. PMID: 11021916.

61. Kulasekara HD, Ventre I, Kulasekara BR, Lazdunski A, Filloux A, Lory S. A
novel two-component system controls the expression of *Pseudomonas aeruginosa*
fimbrial cup genes. Mol Microbiol. 2005 Jan;55(2):368-80.
https://doi: 10.1111/j.1365-2958.2004.04402.x. PMID: 15659157.

62. Chua SL, Tan SY, Rybtke MT, Chen Y, Rice SA, Kjelleberg S, Tolker-Nielsen T,
Yang L, Givskov M. Bis-(3'-5')-cyclic dimeric GMP regulates antimicrobial peptide
resistance in *Pseudomonas aeruginosa*. Antimicrob Agents Chemother. 2013;
57:2066-75. https://doi: 10.1128/AAC.02499-12. PMID: 23403434.

1003

1004 **63**. Friedman L, Kolter R. Genes involved in matrix formation in *Pseudomonas*
1005 *aeruginosa* PA14 biofilms. Mol Microbiol. 2004; 51:675-90.
1006 https://doi: 10.1046/j.1365-2958.2003.03877.x. PMID: 14731271.

1007

1008 **64**. Ye RWl, Haas D, Ka JO, Krishnapillai V, Zimmermann A, Baird C, Tiedje JM.
1009 Anaerobic activation of the entire denitrification pathway in *Pseudomonas*

1010   *aeruginosa* requires Anr, an analog of Fnr. J Bacteriol. 1995; 177:3606-9.

1011   https//:doi: 10.1128/jb.177.12.3606-3609.1995. PMID: 7768875

1012

1013   **65**. Jeong JY, Yim HS, Ryu JY, Lee HS, Lee JH, Seen DS, Kang SG. One-step

1014   sequence- and ligation-independent cloning as a rapid and versatile cloning

1015   method for functional genomics studies. Appl Environ Microbiol. 2012; 78:5440-3.

1016   https://doi: 10.1128/AEM.00844-12. PMID: 22610439.

1017

1018   **66**. Crusz SA, Popat R, Rybtke MT, Cámara M, Givskov M, Tolker-Nielsen T, Diggle

1019   SP, Williams P. Bursting the bubble on bacterial biofilms: a flow cell methodology.

1020   Biofouling. 2012; 28:835-42.

1021   https://doi: 10.1080/08927014.2012.716044. PMID: 22877233

1022

1023   **67**. Müller MM, Hörmann B, Syldatk C, Hausmann R. *Pseudomonas aeruginosa*

1024   PAO1 as a model for rhamnolipid production in bioreactor systems. Appl Microbiol

1025   Biotechnol. 2010; 87:167-74.

1026   https://doi: 10.1007/s00253-010-2513-7. PMID: 20217074.

1027

1028   **68**. McBroom AJ, Johnson AP, Vemulapalli S, Kuehn MJ. Outer membrane vesicle

1029   production by *Escherichia coli* is independent of membrane instability. J Bacteriol.

1030   2006; 188:5385-92. https://doi: 10.1128/JB.00498-06. PMID: 16855227.

1031

1032
1033

Figure Legends

1035

1036

1037

1038

1039

Fig 1. The PQS signalling pathway in *P. aeruginosa*. The PqsABCDE proteins synthesize HHQ, which is converted to PQS by PqsH and also HQNO in conjunction with PqsL. Both HHQ and PQS are released by the cells into the extracellular environment and are taken back up by neighboring cells. Autoinduction occurs when either HHQ or PQS binds to PqsR and amplifies expression of the *pqsABCDE* operon. The *pqsE* gene product has dual functions contributing to AQ biosynthesis as a thioesterase and also via an AQ-independent, thioesterase-independent mechanism to e.g. pyocyanin, rhamnolipid, and lectin production and to biofilm maturation. The conversion of HHQ to PQS confers additional functionalities since PQS unlike HHQ induces microvesicle formation and is a potent iron chelator.

1050

1051

1052

1053

1054

1055

1056

1057

1058

1059

1060

1061

1062

1063

1064

Fig 2. Transposon mutant screen for *P. aeruginosa* strains defective for eDNA
release. *P. aeruginosa* wild-type and mutant strains were grown for 24 h in 96 well
microtiter plates containing ABTG medium, after which the relative levels of eDNA
in the cultures were determined using a PI binding assay. The means and standard
deviations of eight replicates are shown.

1070

Fig 3. *P. aeruginosa tat* mutants form defective biofilms with increased susceptibility to tobramycin. CLSM images showing four-day-old biofilms formed in flow chambers of the *gfp*-tagged *P. aeruginosa* wild-type (a), *tatA* mutant (b) and genetically complemented *tatA* mutant (c). In **(A)** biofilms were stained for total biomass with Syto9 (green) and for eDNA with ethidium bromide (red). In **(B)** biofilms were treated with tobramycin and the medium was supplemented with propidium iodide prior to CLSM such that dead cells appear red while live cells appear green. Each panel shows one horizontal optical section two flanking vertical optical sections. Bars, 20 µm.

1090
1091
Fig 4. Production of pyocyanin (**A**), rhamnolipids (**B**) and MVs (**C**) are reduced in
P. aeruginosa tat mutants. (**A**) Pyocyanin levels are shown in the *P. aeruginosa* wild
type, Δ*pqsA*, *tatA*, and *tatABC* deletion mutants and the *tatA* mutant complemented
with plasmid-borne *tatA*. The impact of exogenous PQS (40 µM) on the *tatA* and
Δ*tatABC* mutants is also shown. Insert panel shows the absence of green pigment
in the *tatA* mutant compared with the wild type and complemented *tatA* mutant. (**B**)
Rhamnolipid production in the *tatA* and Δ*tatABC* mutants compared with the wild
type, Δ*pqsA* mutant and *tatA* complemented with plasmid-borne *tatA*. (**C**)
Comparison of MV production in the *tatA* mutant, complemented *tatA* mutant and in
a double *tatA* Δ*pqsA* mutant compared with the wild type strain. Experiments were
repeated in triplicate at least twice. ***p < 0.001, **p < 0.01; n.s. not significant.
1103

1104

1105
1106
Fig 5. Mutation of *tatA* or exposure to the Tat inhibitor Bayer 11-7082 inhibits *pqsA*
expression and AQ production. (**A**) Mutation of *tatA* or (**B**) treatment with Bayer 11-
7082 supplied at either 20 µM or 40 µM reduces the maximal expression of a
P. aeruginosa PAO1 chromosomal *pqsA'-lux* promoter fusion without affecting
growth. (**C**) LC-MS/MS analysis of PQS production by *P. aeruginosa* PAO1 wild type
compared with the *tatA* mutant and complemented *tatA* mutant. Experiments were
repeated in triplicate at least twice. ***p < 0.001; n.s. not significant.

1114
1115

Fig 6. Exogenous AQs do not restore *pqsA* expression in *P. aeruginosa tatA* (A) or
tatA ΔpqsA (B) mutants. Exogenous PQS or HHQ was added at 5 µM or 20 µM to
(**A**) wild type, the *tatA* mutant and complemented *tatA* mutant or (**B**) Δ*pqsA*, Δ*pqsA*
tatA or *tatA* Δ*pqsA* mutant complemented with *tatA*. Maximal light output from the
chromosomal *pqsA'-lux* fusion was recorded as a function of growth (RLU/OD$_{600}$).
Experiments were repeated in triplicate at least twice. ***p < 0.001, **p < 0.01,
and *p < 0.05; n.s. not significant.

1129
1130
1131
Fig 7. Rhamnolipid (*rhlA*) and pyocyanin biosynthetic (*phzA1*) genes show altered
expression profiles in *P. aeruginosa tat* mutants and fail to respond to exogenous
PQS. (**A**) Light output from a chromosomal *rhlA'-lux* fusion as a function of growth
(RLU/OD) over time when introduced into the wild type, *pqsA* and *tatA* mutants or
complemented *tatA* mutant in the absence or presence of exogenous PQS (20 µM).
(**B**) Maximal light output from a chromosomal *phzA1'-lux* fusion as a function of
growth (RLU/OD$_{600}$) when introduced into the wild type, *pqsA*, or *pqsA tatA* mutants
in the absence or presence of exogenous PQS (10 or 40 µM) or plasmid-borne *pqsE*
or the pUCP18 vector control. Experiments were repeated in triplicate at least twice.
***p < 0.001, **p < 0.01, and *p < 0.05; n.s. not significant.
1142
1143

Fig 8. PqsE restores pyocyanin production in *P. aeruginosa tat* mutants. Production of pyocyanin by the wild type compared with the *tatA* and Δ*tatABC* mutants transformed with plasmid-borne *pqsE* or the empty vector. Experiments were repeated in triplicate at least twice. ***p < 0.001,

1158

1159
1160
Fig 9. Tat substrate screen for *P. aeruginosa* PA14 mutants with defects in *pqs*
signalling uncovers a role for the cytochrome *bc₁* Rieske sub-unit. (**A**) Comparison
of peak *pqsA* expression in each of 34 Tat substrate mutants (**Table S3**)
transformed with a CTX::*pqsA'-lux* fusion compared with the PA14 wild type (white
bar) and the Δ*tatABC* mutant (black bar). The bars represent mutants where *pqsA*
expression is higher (green bar) or lower (grey bar) or the same (grey bar) as the
wild type. PA14 mutant 27 (pink bar) has the lowest *pqsA* expression and carries a
deletion in PA14-57570 (*petA*), the Rieske subunit of cytochrome *bc₁*. (**B**) Production
of PQS, HHQ and HQNO after 16 h growth by *P. aeruginosa* PA14, the Δ*tatABC* and
Δ*petA* mutants and the genetically complemented PA14 mutant
(Δ*petA*::(CTX::*petA*)). ***p < 0.001, **p < 0.01, and *p < 0.05; n.s. not
significant.

1173
1174