Title: BMP signaling is necessary and sufficient for osteoarthritis and a target for disease modifying therapy

AUTHORS:

Akrit Pran Jaswal[1], Anke J. Roelofs[3,#], Amaresh Kumar Singh[1,4,#], Bhupendra Kumar[1], Anna H.K. Riemen[3], Hui Wang[3], Sadaf Ashraf[3], Sanap Vaibhav Nanasaheb[1], Sayeda Fauzia Iqbal[1], Nitin Agnihotri[1], Cosimo De Bari[3], Amitabha Bandyopadhyay[1, 2] *

AFFILIATIONS:
[1] Department of Biological Sciences and Bioengineering, Indian Institute of Technology Kanpur, Kanpur-208016, Uttar Pradesh, India and
[2] The Mehta Family Centre for Engineering in Medicine, Indian Institute of Technology Kanpur, Kanpur, Uttar Pradesh, India.
[3] Arthritis & Regenerative Medicine Laboratory, Institute of Medical Sciences, University of Aberdeen, Aberdeen AB25 2ZD, UK
[4] Department of Zoology, Banaras Hindu University, Varanasi-221005, Uttar Pradesh, India

# These two authors contributed equally

*Corresponding Author: Amitabha Bandyopadhyay

Email address: abandopa@iitk.ac.in

KEYWORDS: BMP, Osteoarthritis, articular cartilage, local inhibition, LDN-193189, post onset

AUTHOR CONTRIBUTIONS: A.B., A.P.J. designed the experiments and A.P.J., A.K.S., B.K., S.V.N. and S.F.I. conducted experiments, collected and analyzed data. A.P.J., A.K.S. and B.K. prepared the manuscript; N.A. conducted the cell-based LDN-193189 assay. A.B., C.D.B., A.J.R. edited the manuscript along with A.P.J.; A.K.S. provided the data for inflammation response studies and mechanistic data including Smurf expression analysis; A.J.R. and A.H.K.R. collected and analysed human cartilage samples; H.W. and S.A. performed the scoring for osteoarthritis.

ABSTRACT:

Osteoarthritis (OA) is among the leading causes of disability across the world. Presently no effective therapy of OA is available as neither the molecular mechanism of disease pathology nor the development and maintenance of articular cartilage is well understood. During OA, articular cartilage undergoes cellular and molecular changes reminiscent of transient cartilage, which is the embryonic precursor of endochondral bone. During endochondral ossification, a precise spatio-temporally regulated WNT-BMP signaling interplay dictates differentiation of a common progenitor pool to either articular or transient cartilage fate in adjacent domains. In the embryonic context, Wnt signaling promotes articular cartilage fate while transient cartilage differentiation is critically BMP signaling dependent. Moreover, any ectopic activation of BMP signaling leads to ectopic transient cartilage differentiation at the expense of articular cartilage. In this study, we show that BMP signaling is sufficient and necessary for the pathogenesis of OA by ectopically activating BMP signaling and depleting BMP ligands in adult mice articular cartilage, respectively. Analysis of human osteoarthritic specimens demonstrated association between OA and ectopic BMP signaling in the articular cartilage. Local inhibition of BMP signaling using a potent pharmacological inhibitor LDN-193189, when administered prophylactically, resulted in delayed onset and reduced severity of OA in mice. Additionally, the same treatment afforded protection against cartilage degeneration post onset of OA in a surgical model of OA in mice. Therefore, inhibiting BMP signaling and consequent block of transient cartilage differentiation within the cells of the joint cartilage can be a possible avenue for developing a disease modifying therapy for OA.

Introduction

Osteoarthritis (OA) is a painful, debilitating musculoskeletal disorder with profound socioeconomic burden and the primary cause of locomotive disability affecting millions of people worldwide (Vos et al., 2012; Cross et al., 2014; Palazzo et al., 2016). The alarmingly increasing prevalence of OA is exacerbated further as no disease modifying therapy exists to manage OA except for symptomatic treatment with anti-inflammatory drugs or surgical intervention in late stage disease (Chen et al., 2017). It is imperative therefore, to discern the molecular basis of pathogenesis of OA in order to develop a disease modifying therapy.

Articular cartilage, the tissue affected in OA, is a lubricated, avascular, alymphatic, aneural tissue that lines the ends of the bones at the joints. During OA, the joint surface undergoes a slew of changes characterised by loss of cartilage proteoglycans, hypertrophy of chondrocytes, angiogenesis, osteophyte formation, and ultimately failure of joint function (Chen et al., 2017). The cellular and molecular changes of the joint cartilage during the onset and progression of OA closely resemble the steps of endochondral ossification, the developmental process by which long bones form within cartilage anlagen.

During endochondral ossification, most of the initial cartilage mass in an appendicular skeletal element is replaced by newly forming bone except for the cartilage at the termini. The cartilage that is replaced by bone is referred to as the transient cartilage while the cartilage at the terminal ends is referred to as the joint or articular or permanent cartilage (Kozhemyakina et al., 2015). During transient cartilage differentiation, type II collagen (Col2a1) expressing cartilage cells undergo a series of changes. These cells undergo pre-hypertrophic differentiation wherein they express Indian hedgehog (IHH), subsequently the transition from pre-hypertrophy to hypertrophy is marked by the expression of type X collagen (ColX). The hypertrophic cells are infiltrated by blood vessels. This is followed by matrix remodelling, where enzymes viz. MMP-13 and ADAMTS-5, degrade the existing collagen matrix and a new matrix, rich in type I collagen (ColI), is synthesised and bone formation is accomplished (Ortega et al., 2004; Long and Ornitz, 2013). Ray et.al. demonstrated that within a developing appendicular skeletal element there is a zone of Col2a1 expressing bipotential proliferating cell population referred to as the Distal Proliferative Zone (DPZ). The DPZ cells under the influence of BMP signaling undergo transient cartilage differentiation whereas when exposed to WNT signaling undergo joint cartilage differentiation (Ray et al., 2015). Some of the molecules involved in transient cartilage differentiation viz. MMP-13, ADAMTS-5, and VEGF-A are reported to be associated and/or necessary for the pathogenesis of OA (Lark et al., 1997; Dahlberg et al., 2000; Neuhold et al., 2001; Enomoto et al., 2003; Glasson et al., 2005; Song et al., 2007; Barranco, 2014; He et al., 2014; Ko et al., 2017).

Previous literature suggests that ectopic activation of BMP signaling in developing cartilage or presumptive joint site, either by overexpression of BMP ligands (Duprez et al., 1996; Ray et al., 2015) or misexpression of constitutively active BMP receptors (Zou et al., 1997), results in transient cartilage differentiation at the expense of joint cartilage. In addition, immobilisation of developing embryonic limbs leads to ectopic differentiation of transient cartilage at the cost of articular cartilage (Kahn et al., 2009; Singh et al., 2018). Recently, it was shown that immobilization leads to ectopic upregulation of BMP signaling within the sub-articular cartilage domain where cartilage precursors are normally exposed only to WNT signaling (Rolfe et al., 2014; Singh et al., 2018).

Taken together, we hypothesized that BMP signaling induced transient cartilage differentiation within adult articular cartilage domain is the molecular basis of the pathogenesis of OA. In this manuscript we tested this hypothesis with conditional gain- and loss-of-function mouse mutants of BMP signaling in conjunction with a surgically induced model of OA. Our study in mice is also supported by data obtained from osteoarthritic human cartilage specimens, wherein we found evidence of active BMP signaling in the joint cartilage. Further, our data suggests that *in situ* pharmacological inhibition of BMP signaling in the synovial joint may serve as an effective disease modifying therapy for OA.

RESULTS:

1. Ectopic activation of BMP signaling in adult joint cartilage is sufficient to induce the development of an OA-like phenotype in mice

To examine whether ectopic activation of BMP signaling in the articular cartilage is sufficient to induce osteoarthritic changes in adult mouse we generated *pMes-caBmpr1a; Col2a1-Cre-ERT2* mouse strain, where BMP signaling can be ectopically activated in adult articular cartilage upon administration of tamoxifen (Fig.1A). We activated BMP signaling in postnatal cartilage at P56 by injecting tamoxifen intraperitoneally (*referred to as induction from hereon*) to *pMes-caBmpr1a; Col2a1-Cre-ERT2* mice. Adult mice articular cartilage is devoid of active BMP signaling (Fig.1B). However, within seven days of mis-expression of *caBmpr1a* in adult mouse cartilage ectopic activation of canonical BMP signaling, as assessed by immunoreactivity towards phosphorylated SMAD1/5/8 (pSMAD1/5/8), is observed and it peaks after two weeks (Fig.1B′-B‴). Subsequently, markers of transient cartilage differentiation were also observed in the articular cartilage domain in the same sequence as observed during endochondral ossification. Expression of IHH, which marks prehypertrophic state of cartilage, was observed within 7 days of induction and concurrent with the first detection of pSMAD1/5/8 immunoreactivity. IHH expression is downregulated by the 14th day after induction (Fig.1C-C″). ColX expression, which marks cartilage hypertrophy, was observed 14 days post induction and the maximum extent of hypertrophy was at 28 days post induction (Fig. 1D-D‴). Embryonic (Mankin, 1962; Ray et al., 2015) as well as adult articular cartilage cells (Mankin, 1963) are proliferation deficient while transient cartilage cells are proliferative (Ray et al., 2015). In our experiments we observed that along with other makers of transient cartilage differentiation, proliferation was also stimulated in the adult mice articular cartilage upon induction. Increased BrdU uptake was observed in the adult joint cartilage 7 days after induction which was more apparent on the 14th day post induction (Fig.1E-E″).

At 60 days post induction, an upregulation of matrix remodelling enzyme MMP-13 was also observed in articular cartilage cells (Fig.1F-F′). Moreover, ColI, marking osteogenic differentiation, was also found to be expressed in multiple zones of articular cartilage 60 days post activation of BMP signaling (Fig.1G-G′). Six months post induction, widespread upregulation of MMP-13 and ColI was observed throughout all zones of articular cartilage suggesting terminal osteogenic differentiation had occurred (Fig. 1F″ and Fig. 1G″). Focal loss of proteoglycan staining in multiple zones of articular cartilage was also observed (Fig. 1H-1H′).

Further, we observed the appearance of NITEGE immunoreactivity in the adult articular cartilage 6 months post induction (Fig. 1I-1I′). Besides the molecular signatures, the most conspicuous feature of osteoarthritic articular cartilage is the formation of osteophytes in the joint, which is otherwise devoid of such features. Micro CT imaging of hind limbs revealed extensive osteophyte (indicated by black arrow) formation upon ectopic activation of *Bmpr1a* in articular cartilage (Fig.1J). Taken together, these observations strongly suggest that ectopic activation of BMP signaling is sufficient to induce the development of an OA like phenotype in adult mice.

2. BMP signaling induced transient cartilage differentiation is necessary for the pathogenesis of OA

Next, we investigated the necessity of BMP signaling in the development of osteoarthritic phenotype. It has been previously reported that levels of BMP-2 ligands are elevated in synovial fluid from OA patients and BMP receptor localisation is associated with OA severity (Nakase et al., 2003; Schmal et al., 2012). However, the presence of active BMP signaling *in situ* osteoarthritic cartilage has not been reported. To induce OA in mice we used Anterior Cruciate Ligament Transection (ACLT), an established model for surgical induction of experimental OA (Hayami et al., 2006; Glasson et al., 2007). We chose to analyse the knee articular cartilage every week after ACLT for the appearance of active BMP signaling. As opposed to sham operated knees (Fig. S1A) or 7 days post ACLT (Fig. S1B) we observed increased pSMAD1/5/8 immunoreactivity in specimens from 14 days post ACLT (Fig. S1B′) which persisted till 56 days post-ACLT (Fig. S1B″, Fig. S1B‴ and Fig. 2B′). In keeping with our observation in the context of ectopic activation of BMP signaling, we observed increased uptake of BrdU in the articular cartilage of mice post ACLT (Fig. S1D-D‴) in contrast to the sham-operated group (Fig. S1C). In order to prevent activation of BMP signaling post ACLT, we used a previously described *Bmp2/4* double conditional knockout strain (Bandyopadhyay et al., 2006). We crossed this strain with *Col2a1-Cre-ERT2* to create *Bmp2^{c/c}; Bmp4^{c/c}; Col2a1-Cre-ERT2* strain. This allowed us to deplete BMP ligands in the articular cartilage in a temporally controlled manner through tamoxifen injection. We injected tamoxifen intraperitoneally in P56 mice. ACLT was performed at P120 (Fig. S2A and Fig. 2A).

As expected, we could not detect any upregulation of BMP signaling in the articular cartilage of *Bmp2/4* depleted animals when subjected to ACLT (Fig. S2B″ and Fig. 2B″). Distribution and abundance of ColII was found to be normal in *Bmp2/4* depleted animals even after 56 days of ACLT (Fig. S2C-C″ and Fig. 2C-C″). Cartilage hypertrophy, as assessed by ColX immunoreactivity (Fig. 2D-D″) as well as expression of MMP-13 (Fig. 2E-E″), a key matrix remodelling enzyme, were significantly elevated after 56 days of ACLT (Fig. 2D′ and Fig. 2E′). However, depletion of *Bmp2/4* protected against extensive upregulation and allowed the maintenance of a level of ColX (Fig. 2D″) and MMP-13 (Fig. 2E″) almost comparable to that of sham (Fig. 2D and Fig. 2E). We also noted that while cartilage loss of proteoglycan staining was observed in ACLT specimens, these changes were minimum post depletion of BMP ligands (Fig. 2F-F″). The emergence of neo-epitope NITEGE resulting from aggrecan degradation was also considerably reduced in *Bmp2/4* depleted articular cartilage (compare, Fig. 2G″ with Fig. 2G′). Similar observations were made 28 days post ACLT as well (Fig. S2G-S2G″). Moreover, ColI expression, which is a marker for bone differentiation (or

ossification), was also found to be at levels significantly lower in articular cartilage post ACLT in BMP depleted tissues as compared to vehicle injected ACLT control animals (compare, Fig. 2H″ with Fig. 2H′). Structural examination using micro-computed tomography (µCT) revealed that in "ACLT + Vehicle" group, extensive damage to articular surfaces (roughness) as well as osteophyte (marked by red arrows) formation was observed (Fig.2I′). However, upon depletion of BMP ligands, the severity and extent of these changes were significantly attenuated (Fig. 2I″), and cartilage quality (smoothness) in this group appeared comparable to that in sham operated group (Fig.2I) suggesting cartilage protection was afforded.

To ascertain the clinical relevance of these findings we examined both osteoarthritic and non-osteoarthritic human articular cartilage. Immunoreactivity of pSMAD1/5/8 was observed in all zones of osteoarthritic cartilage from patients who underwent arthroplasty (Fig. 2 J″, 2J‴), while human cartilage from a donor without known history of OA showed minimal detectable pSMAD1/5/8 immunoreactivity (Fig. 2K″, 2K‴). There was no pSMAD1/5/8 immunoreactivity in osteoarthritic cartilage treated with phosphatase (Fig. 2 J′, 2K′).

3. Local pharmacological inhibition of BMP signaling halts the progression of osteoarthritic changes

We next investigated whether inhibiting BMP signaling locally in articular joints can retard the progress of osteoarthritic changes after surgical induction of OA. For this purpose we used a well characterized BMP signaling inhibitor, LDN-193189, a derivative of dorsomorphin (Yu et al., 2008; Hao et al., 2010; Sanvitale et al., 2013). LDN-193189 activity was assayed using BRITER (BMP Responsive Immortalized Reporter) cell line (Yadav et al., 2012) (please see Materials and Methods). As low as 100nM concentration of LDN-193189 significantly inhibited BMP signaling in BRITER cell line (Fig. S3).

Considering possible dilution and volume loss of LDN-19389 during the injection, we used 10 µM (in 3% w/v 2-hydroxypropyl- β-cyclodextrin in PBS) of LDN-193189 for intra-articular injection to locally inhibit BMP signaling. We devised a regime where we performed ACLT on wild type mice at P120 and injected LDN-193189 into the knee joint intra-articular space for seven consecutive days starting on the 14th day post-surgery and harvested 28 days post-surgery (Fig. 3A).

We observed that local inhibition of BMP signaling severely abrogates OA-associated changes in articular cartilage post transection of ACL. No immunoreactivity to pSMAD1/5/8 was observed in articular cartilage of LDN-treated set post ACLT (Fig. 3B″) and the sham operated knee cartilage (Fig.3B), while pSMAD1/5/8 positive cells were found in zones of articular cartilage in vehicle administered ACLT knee joints (Fig.3B′). The level and distribution of ColII was native like in LDN-193189 treated knee joints (Fig.3C″) as compared to ACLT control set (Fig.3C′) suggesting reduced depletion of ColII post-surgery and protection of cartilage. Hypertrophy of cartilage cells was found to be restricted to the calcified zones and minimal ColX immunostaining was observed in the articular cartilage of LDN-treated ACLT knees (Fig. 3D″) similar to sham set (Fig.3D), while vehicle injected animals showed extensive hypertrophy throughout the cartilage matrix (Fig.3D′). Similarly, MMP-13 levels were found to be significantly reduced in articular cartilage upon intra-articular administration of LDN-193189 (Fig.3E″) whereas in vehicle injected knee joints, a global upregulation of MMP-13

was observed throughout multiple zones of articular cartilage (Fig.3E′). Further, we observed that articular cartilage in LDN-193189 treated group was devoid of any immunoreactivity to NITEGE (Fig.3F″) similar to what was observed in the sham set (Fig.3F), suggesting lesser aggrecanolysis by aggrecanolytic enzymes while abundant NITEGE expression was observed in vehicle treated specimens (Fig.3F′). Proteoglycan depletion and cartilage damage was observed to be minimal in tibial surface of LDN-193189 injected (Fig.3G″) as compared to vehicle injected controls (Fig. 3G′) and cartilage integrity was found to be comparable to sham operated knees (Fig. 3G). To examine the status of terminal transient cartilage differentiation following completion of LDN-193189 regime, we carried out immunostaining for ColI. As opposed to vehicle treated ACLT group (Fig.3H′), we found that very few cells in articular cartilage of LDN-193189 treated joints expressed ColI (Fig.3H″). The structural parameters were examined by μCT imaging. We observed that articular cartilage quality was maintained in terms of minimal erosion, osteophytes formation and less pronounced surface changes upon local inhibition of BMP signaling by using LDN-193189 intra-articular injection (Fig. 3I-I″, red arrows mark osteophytes). Moreover, OARSI scoring of cartilage revealed a significantly attenuated osteoarthritic-like phenotype in LDN-193189 treated group as compared to the vehicle-treated ACLT group (Fig.3J).

Taken together, these findings suggest that *in situ* inhibition of BMP signaling in articular cartilage is sufficient to prevent the phenotypic and molecular changes associated with development and progression of OA in a surgically induced osteoarthritic mouse model.

4. Inhibition of BMP signaling post onset of OA attenuates disease severity

In situ inhibition of BMP signaling before the onset of OA following ACL transection in mice retards the progression of OA. However, in a clinical setting, patients report to the clinic after the disease has set in. We therefore investigated if local inhibition of BMP signaling can mitigate the severity of osteoarthritic changes even after the disease has set in. For this purpose, seven consecutive intra-articular LDN-193189 injections were administered starting on post-surgery day 35 and finishing on post-surgery day 42. The knees were harvested at post-surgery day 56 (Fig.4A).

In contrast to vehicle treated knee joints, where ColX positive cells could be observed throughout articular cartilage (Fig. 4B′) in LDN-193189 treated samples, hypertrophic cells were found restricted only to a few upper zone cells of articular cartilage (Fig. 4B″). The level of MMP-13 was also observed to be markedly less in the articular cartilage of LDN-193189 administered knees (Fig. 4C″), as compared to the vehicle control ACLT group (Fig. 4C′) where a high level of MMP-13 expression was observed throughout. Moreover, NITEGE neo-epitope was absent from the middle or deeper zones of articular cartilage in LDN-193189 administered knees (Fig. 4D″) as opposed to vehicle treated ACLT set (Fig. 4D′) where such cells were found throughout the depth of articular cartilage. Proteoglycan staining was found to be more intense in certain areas of LDN-193189 treated knee joints (Fig. 4E″) which did not coincide with any known marker of hypertrophy or transient cartilage differentiation while overt signs of OA were observed in vehicle treated ACLT knee cartilage (Fig. 4E′). Similarly, while ColI positive cells were found in vehicle treated ACLT set (Fig. 4F′), such cells were not observed in the LDN-193189 treated knee specimens (Fig. 4F″) or in the sham operated group (Fig. 4F). Evaluation of the structural parameters using μCT imaging revealed that cartilage

surface erosion was reduced in LDN-193189 treated set as compared to vehicle injected knees (Fig. 4G-4G″, red arrow marks osteophytes). The OARSI scores in LDN-193189-treated group were significantly lower than that for ACLT group, even though administration of LDN-193189 was performed after the onset of disease (Fig.4H). It should be noted though that less protection of cartilage was afforded, as judged by the OARSI severity scores, to the knee joints treated with LDN-193189 post-onset of OA compared to when knee joints were treated with LDN-193189 pre-onset of OA (compare Fig. 3J and Fig. 4H).

We have observed that intra-articular administration of LDN-193189 provides protection against OA like changes at least for 14 days post injection (Fig. 4). Next, we wanted to investigate the potential for clinical translatability of LDN-193189 or similar molecule as a disease modifying agent. We examined whether LDN-193189 can confer longer term protection against surgically induced OA by emulating a clinic-like regimen of minimum dosage and maximum efficacy over extended durations of time. Our data (Fig. S2) as well as existing literature (Kamekura et al., 2005) suggest that molecular changes associated with OA are apparent within 28 days of ACLT. Hence, we conducted ACLT at P120, injected LDN-193189 intra-articularly on PS28, PS30 and PS32 and harvested the knee joint 56 days later at PS84. As compared to vehicle control, in the LDN-193189 injected specimen, ColII expression (compare Fig. 5B with Fig. 5B‴) as well as cartilage specific proteoglycan content, as assessed by Safranin O staining (compare Fig. 5D with Fig. 5D‴), were largely preserved. Also, ColX immunoreactivity was significantly lower in LDN-193189 treated specimen (compare Fig. 5C with Fig. 5C‴). This set of data suggests that even after onset of surgically induced OA blocking BMP signaling pathway locally can offer protection for at least 56 days in mice.

5. Mechanistic insight into the pathogenesis of OA from a developmental biology perspective

Recently, Singh *et al.*, demonstrated that immobilisation of chick or mouse embryos results in transient cartilage differentiation at the expense of articular cartilage differentiation which is associated with ectopic activation of BMP signaling (Singh et al., 2018). Further, this study also demonstrated that this ectopic activation is associated with a concurrent downregulation of expression of SMURF1, an intracellular inhibitor of BMP signaling pathway (Singh et al., 2018). We detected a downregulation of SMURF1 (Fig. 5E-E″) expression in the mouse articular cartilage 30 or 60 days post ACLT. This suggests that the molecular mechanism of articular cartilage maintenance via mechanical regulation is conserved between embryonic and postnatal stages and likely involved in pathologies such as OA.

6. Effect of local inhibition of BMP signaling on inflammatory responses in a surgically induced osteoarthritic mouse model

We performed an analysis for candidate inflammatory response molecules which are known to be involved in the development of osteoarthritis (Tetlow et al., 2001; Kobayashi et al., 2005). IL-1β immunostaining showed that while there was an increased immunoreactivity to IL-1β in the vehicle-treated group (Fig. 6A′), minimal reactivity could be detected in LDN-193189 treated mice (Fig. 6A″), comparable to that observed in the sham-operated group (Fig.6A). Similarly, we performed immunostaining for iNOS, which is a known key player in the development of OA. iNOS was detected in the subchondral bone of sham-operated mice (Fig.

6B) and upon ACLT, iNOS was also detected in some of the articular cartilage cells of vehicle treated mice (Fig. 6B′) while no such cell could be detected in the LDN-193189 treated group (Fig. 6B″). Immunoreactivity for NF-kB was detected in the articular cartilage of the vehicle treated ACLT group (Fig. 6C′), while it could not be detected in either the LDN-193189 treated group or in the sham operated group (Fig. 6C″ and 6C, respectively). We also investigated TNF-α immunoreactivity in osteoarthritic cartilage upon treatment with LDN-193189 and found that TNF-α immunoreactivity was observed in the ACLT group injected with vehicle (Fig. 6D′, white arrow), while the LDN-193189 treated group displayed minimal immunoreactivity (Fig. 6D″), similar to that in sham-operated mice where TNF-α could be detected in subchondral bone (Fig.6D).

Discussion:

Our findings show that there exists striking resemblance between the molecular changes associated with the onset of osteoarthritic changes in the articular cartilage and the molecular signature associated with endochondral ossification. Moreover, the temporal order of the expression of molecular markers post ACLT induced OA was found to be strikingly similar to what is observed with transient cartilage differentiation during endochondral ossification. Inhibition of transient cartilage differentiation via blocking IHH signaling has been reported to inhibit and attenuate the severity of osteoarthritic phenotype post ACLT (Lin et al., 2009; Zhou et al., 2014a; Zhou et al., 2014b). However, so far, no IHH signaling inhibitor has been approved for clinical use. This could be because Ihh loss-of-function, during embryonic development, results only in a temporary delay in hypertrophic differentiation – an essential step of endochondral ossification, which eventually gets restored and even accelerated at postnatal stages in mice (Vortkamp et al., 1996; Mak et al., 2008). Taken together, these studies give a crucial hint that blocking transient cartilage differentiation is a viable strategy to manage osteoarthritic changes in the articular cartilage. This hypothesis is in line with what has been suggested earlier in the literature (Sandell and Aigner, 2001; Pitsillides and Beier, 2011; Gelse et al., 2012).

BMP signaling is known to play a critical role in transient cartilage differentiation. Previously, intraperitoneal administration of a BMP signaling inhibitor, LDN-193189, has been shown to reverse the phenotype associated with *Fibrodysplasia ossificans progressiva,* a condition where progressive heterotopic ossification of muscle is observed upon injury, due to constitutive activation of BMP signaling (Yu et al., 2008). In this study, we found that activation of BMP signaling is both necessary and sufficient for the pathogenesis of OA in mice. The necessity of BMP signaling in the onset of osteoarthritis like changes in the articular cartilage has been demonstrated using both genetic and pharmacological means whereas sufficiency has been demonstrated using genetic means. Further, analysis of patient samples suggests an association between osteoarthritis and activation of BMP signaling in the articular cartilage cells. Since we have used *Col2a1-Cre-ERT2* mediated recombination as the means to activate expression of caBMPRIA we cannot rule out the possibility that BMP signaling has also been activated in the growth plate cartilage of adult mice and the molecular and cellular changes observed is partly due to activated BMP signaling in the growth plate cartilage. However, in all our analyses (Fig. 1) the changes were first observed in the superficial layers of articular cartilage suggesting that the changes observed were primarily due to ectopic activation of BMP signaling in the articular cartilage.

Interestingly, we also observed a surge in proliferation in articular cartilage cells, as assessed by enhanced BrdU uptake, post ACLT or activation of BMP signaling. Our data suggests that articular cartilage cells, originally having low regenerative potential and proliferative capacity, display a regenerative response upon ACLT or upregulation of BMP signaling. However, this leads to an altered tissue microenvironment that promotes transient cartilage differentiation at the expense of articular cartilage. Thus, instead of healing by regeneration it further promotes the disease condition. Prophylactically blocking BMP signaling *in situ* using LDN-193189 led to an attenuation in the severity of osteoarthritic phenotype following surgical induction in mice. Further, our investigation suggests that administration of LDN-193189 after the onset of OA not only halts the progression of OA but also an intense toluidine blue stained cartilage tissue appears which is negative for transient cartilage markers, suggesting that new cartilage formation takes place.

Finally, it has to be acknowledged that while transient cartilage differentiation could be involved in the initiation of the disease, the inflammation associated with OA can determine the severity and course of disease progression (Goldring and Otero, 2011; Sokolove and Lepus, 2013). However, despite a large body of existing literature exploring the role of inflammation in the pathology of OA, there is no clear demonstration of the hierarchy between onset of inflammation and transient cartilage differentiation. Our data suggest that pharmacologically blocking BMP signaling in surgically induced OA knees prevents activation of inflammatory response as well (Fig. 6). Our data suggest that *in situ* inhibition of BMP signaling, and consequently transient cartilage differentiation, may be a potent means of disease modifying therapy for osteoarthritis.

Materials and Methods

Generation of mice lines

All animals were housed, bred and maintained in Central Experimental Animal Facility (CEAF) of Indian Institute of Technology Kanpur, India. All experiments were performed in accordance with the guidelines of the Institutional Animal Ethics Committee (IAEC) as well as under the aegis of the Centre for Purpose of Control and Supervision of Experiments on Animals (CPCSEA), Government of India under protocols IITK/IAEC/2013/1002; IITK/IAEC/2013/1015; and IITK/IAEC/2013/1040. We obtained *B6By/J* wild type mice, *TgCol2a1-Cre-ERT2* (Nakamura et al., 2006) and *ROSA26 mT/mG (Muzumdar et al., 2007)* strains from Jackson Laboratories, USA; *pMes-caBmpr1a* mice as gift from Prof. YiPing Chen at Tulane University, USA; *Bmp2$^{c/c}$*; *Bmp4$^{c/c}$* mice from Prof. Clifford Tabin at Harvard Medical School, USA. For BMP signaling gain-of-function, *pMes-caBmpr1a* mice were crossed with *TgCol2a1-Cre-ERT2* mice to generate *pMes-caBmpr1a; TgCol2a1-Cre-ERT2*. *Bmp2$^{c/c}$; Bmp4$^{c/c}$ (Bandyopadhyay et al., 2006)* animals were crossed with *TgCol2a1-Cre-*

ERT2 to generate *Bmp2$^{c/c}$; Bmp4$^{c/c}$*; *TgCol2a1-Cre-ERT2* for BMP loss-of-function mutation. Mice were genotyped for the presence of *caBmpr1a* using PCR with primers 5′-CGAAGATATGCGTGAGGTTGTGT-3′ and 5′- TGGGCGAGAGGGGAAAGAC - 3′ while *TgCol2a1-Cre-ERT2* transgene was genotyped using primer pair 5′-CACTGCGGGCTCTACTTCAT-3′ and 5′- ACCAGCAGCACTTTTGGAAG- 3′. *ROSA26 mT/mG* was genotyped using primers 5′-CTCTGCTGCCTCCTGGCTTCT-3′ and 5′-TCAATG GGCGGGGGTCGTT - 3′. *Bmp2$^{c/c}$* was genotyped using 5′-GTGTGGTCCACCGCATCAC-3′ and 5′- GGCAGACATTGTATCTCTAGG - 3′ and *Bmp4$^{c/c}$* was genotyped using 5′-AGACTCTTTAGTGAGCATTTTCAAC-3′ and 5′-AGCCCAATTTCCACAACTTC - 3′ primer pairs.

Anterior Cruciate Ligament Transection (ACLT)

All ACLT surgeries were conducted at P120. Unless specifically mentioned, ACLT was conducted on *B6By/J* wild type male mice. Animals were subjected to ACLT at P120 and harvested at post ACLT day 07, 14, 21, 28, and 56 for molecular and histological analysis. SMURF1 immunoreactivity was analyzed after one and two months of ACLT in wild type mice. All animals were anesthetized using isoflurane during ACLT surgery following standard protocol (Glasson et al., 2007). ACLT was performed in left hind limb of all mice. Patella was dislocated medially to expose femorotibial joint cavity and ACL was transected using a micro-surgical blade under the microscope. After ACL transection, patella was repositioned and quadriceps femoris muscle was sutured to prevent its dislocation. In the sham group, the joint cavity and ACL were exposed but left intact and sutured. Surgical incision was sealed using Vicryl 5-0 Absorbable suture. Povidone-Iodine solution was applied over the sutured skin to prevent microbial infections.

BMP signaling gain-of-function

pMes-caBmpr1a;TgCol2a1-Cre-ERT2 male mice were injected with three consecutive doses of tamoxifen (Sigma-Aldrich, 2.5 mg/25 g body weight) starting from P56 to induce Cre mediated recombination and harvested at post TAM injection day 7,14, 21, 28, 56 and 6 months for molecular and histological analysis.

BMP signaling loss-of-function

All animals used for BMP signaling loss-of-function were homozygous males for the conditional alleles of *Bmp2* and *Bmp4*. Three doses of tamoxifen (Sigma-Aldrich, 2.5 mg/25 g body weight) were injected in *Bmp2$^{c/c}$; Bmp4$^{c/c}$; TgCol2a1-Cre-ERT2* mice intraperitoneally starting from P56 for three consecutive days to activate Cre-mediated recombination to deplete BMP2 and BMP4 in adult cartilage. pSMAD1/5/8 immunoreactivity was used to assess the status of BMP signaling before/after ACLT and/or with/without TAM-induced recombination.

Local inhibition of BMP signaling was achieved through administration of 6µl solution of 10µM LDN-193189, prepared in 3% 2-(hydroxypropyl)-β-cyclodextrin (w/v in PBS) in the knee joint cavity. Local inhibition of BMP signaling was done at two different time points. In one set, LDN-193189 was injected daily from 14th to 21st day post ACLT. In the other set LDN-193189 was administered daily from 35th to 42nd day post ACLT. For the experiment conducted to assess the dosage required and the duration of protection offered by LDN-193189 ACLT was conducted on P120 and LDN-193189 was injected

To observe the persistent effect of LDN-193189 over a prolonged period as well as to ascertain the optimum number of doses needed for providing long term protection following ACLT, LDN-193189 was injected on post-surgery day 28 (single dose) or post-surgery days 28 and 30 (two doses) or post-surgery day 28, 30 and 32 (three doses). Tissues were harvested at post ACLT day 84 for further analysis. In all experiments the vehicle control animals were injected with 3% 2-(hydroxypropyl)-β-cyclodextrin (w/v in PBS).

Cell proliferation analysis

Cell proliferation status was assessed using BrdU incorporation assay following intraperitoneal delivery of 100mg BrdU/10ml PBS/kg of body weight for three consecutive days prior to the day of harvesting tissues.

LDN-193189 activity assay

BRITER (BMP Responsive Immortalized Reporter) cell line (Yadav et al., 2012) were used to determine optimum concentration of LDN-193189 required to inhibit BMP signaling.

Human sample collection

Osteoarthritic cartilage from 6 patients and non-osteoarthritic cartilage from one patient undergoing knee joint excision for malignancy at a site not involving sampled area, were

obtained after informed consent and in accordance with the relevant guidelines and regulations, with approval from the NHS Grampian Biorepository Tissue Bank Committee, UK.

Processing of tissue, histology and immunohistochemistry

Post harvesting, tissues were fixed in 4% paraformaldehyde for 16h at 4°C and then washed in PBS for 2h. The tissues were decalcified in 14% EDTA solution (pH 7.4) for 28 days with regular changes of EDTA solution at least once in a week. Decalcified tissues were embedded in paraffin wax and sectioned at 5 μm thickness.

For immunohistochemistry, sections were treated with xylene and then rehydrated using descending alcohol gradient. Antigen retrieval was done using citrate buffer except for collagens. ColI, ColII and ColX antigens were retrieved by treatment of acidic solution of 0.05% pepsin at 37ºC for 25 minutes. Tissue sections were washed with PBS and treated with 3% H_2O_2 for 15 minutes to block endogenous peroxidase activity. After blocking in 5% HINGS (Heat-inactivated Goat Serum) solution for an hour, tissue sections were incubated with primary antibodies against Type X Collagen (DSHB, Cat.no. X-AC9-s, 1:20), Type II Collagen (DSHB, Cat.no. CIIC1-s, 1:20), NITEGE (Abcam, Cat.no. ab3773,1:20), SMURF1 (Santa Cruz Biotechnology,Cat.no.SC100616,1:100), NF-κB (CST,Cat.no.D14E12,1:400), IL-1β (CST, Cat.no.3A6,1:100), MMP-13 (Santa Cruz Biotechnology, Cat.no.SC30073,1:100), pSMAD1/5/8 (CST Cat.no.13820S, 1:100), GFP (Invitrogen, Cat no.A6455, 1:100), BrdU (Sigma, Cat.no. B8434), iNOS (Abcam, Cat. no.3523, 1:50), and TNF-α (Santa Cruz Biotechnology, Cat.no. SC-52746, 1:100) for 16 hours at 4ºC. Tissue sections were washed with PBST and incubated with fluorescence labelled secondary antibodies for overnight at 4ºC. Secondary antibody against BrdU was conjugated with HRP (horseradish peroxidase) and detected by DAB (3,3 Diaminobenzidine).

Histological analysis was performed using Safranin O and Toluidine blue staining following standard protocols. For Safranin O/ Fast green staining, rehydrated tissue sections were treated with haematoxylin solution for 10 minutes, washed in running tap water, and treated with 0.5% Fast Green solution for 2 minutes. Thereafter, sections were dipped in 1% acetic acid and finally stained with 0.5% Safranin O solution for 2 minutes. Stained sections were dehydrated with 95% and 100% alcohol respectively, cleared with Xylene and mounted in DPX. Toluidine blue staining was performed on rehydrated tissue sections with 0.05% toluidine blue solution and washed in tap water. Slides were air dried, treated with xylene and mounted in DPX.

Human cartilage sections were stained by immunohistochemistry for pSmad1/5/8 using standard protocols (Roelofs and De Bari, 2019). Briefly, antigen retrieval was performed by incubation for 1 hour at 65ºC in antigen unmasking citrate buffer solution (pH 6, Vector Laboratories, UK), after which sections were incubated with phosphatase (CST Lambda Phosphatase kit, New England Biolabs, #P0753) according to manufacturer's instructions, or incubated with buffer only. Sections were then incubated with an antibody against pSmad1/5/8 (Millipore, cat no. ab3848, at 1:50 (10 µg/ml)) followed by biotinylated goat anti-rabbit IgG (Vector Labs, cat no. BA-1000, at 1:200). Stained sections were imaged using Zeiss Axioskop 40 (Zeiss) with Progress XT Core 5 colour digital camera and ProgRes CapturePro 2.9.0.1 software (JenOptik, Germany).

The OARSI scores were calculated using the recommended guidelines for assessment of osteoarthritic severity in small animals (mice). OARSI scoring was performed by researchers who were not involved in the animal experiments and who were blinded to the treatment groups. Data was then unblinded and analysed by the researchers carrying out the experiments. All images were taken using Leica DM 5000B compound microscope and processed using manufacturer provided software suite(s).

Micro-Computed Tomography (µCT)

Fixed tissue were taken in 5ml microfuge tube in hydrated condition and imaged using high resolution µCT (Skyscan 1172). Images were reconstructed and analyzed using NRecon v1.6 and CTAn 1.16.8.0, respectively. 3-D volumetric projections were generated using CTVol 2.0 software. The scanner was set to operate at a voltage of 50 kVp, a current of 200 µA and the resolution of the images was 5.86 µm per pixel. µCT for BMP signaling gain-of-function specimens was performed, processed and analyzed at the Institutional facility of Western University, Ontario, Canada using eXplore Locus GE scanner. The resolution of the image is 40 µm per pixel and 3-D isosurface projection was generated using Parallax Innovations MicroView 2.5.0- rc15 (2.5.0-3557).

Statistics

Graph Pad Prism 8 software was used to ascertain statistical significance of the OARSI histopathological scoring data. One-way ANOVA with post-hoc analysis (Dunnett's test) was

performed to calculate the statistical significance and interdependence between different experimental groups. The results were plotted using Graph Pad Prism 8 software. The error bars represent Standard Deviation (S.D.).

Acknowledgements

We are immensely grateful to Prof. YiPing Chen at Tulane University, USA for the gift of mouse strains. We thank Prof. Frank Beier of Western University, Ontario, Canada for teaching APJ the method of ACL transection. We sincerely thank Shuchi Arora and Ankita Jena for their critical comments on the manuscript and Niveda Udaykumar for help with statistical analysis. We thank Mr. Naresh Gupta for assistance with mouse experiments.

• **Funding**

This work was supported by grants from the Department of Biotechnology, India (DBT) BT/PR17362/MED/30/1648/2017 and BT/IN/DENMARK/08/JD/2016 to A.B.; Versus Arthritis Grants 19667 and 21156 to CDB and AJR, Fellowships to APJ, BK, and SFI are supported by fellowships from the Ministry of Education, Govt. of India. Fellowship to AKS was supported by Science and Engineering Research Board, Govt. of India. APJ travelled to Western University Canada with Shastri Research Student Fellowship (SRSF, 2015-'16). A.H.K.R. was supported by the Wellcome Trust through the Scottish Translational Medicine and Therapeutics Initiative (Grant No. WT 085664).

Competing Interests:

The authors declare the following competing interests:

The use of BMP inhibitors as locally administered agents using sustained drug delivery vehicle(s) has been submitted for patent via Indian patent application number – **201911044840**.

The inventor(s) are:

1. Dr. Amitabha Bandyopadhyay–IIT Kanpur, India Mr. Akrit Pran Jaswal - IIT Kanpur, India.
2. Dr. Praveen Vemula – Institute for Stem Cell Biology and Regenerative Medicine, India.
3. Dr. Manohar Mahato - Institute for Stem Cell Biology and Regenerative Medicine, India.

Legends to Figures

Fig. 1. Ectopic activation of BMP signaling in adult joint cartilage is sufficient to induce OA development.
(A) Schematic for generation of *pMes-caBmpr1a; Col2a1-Cre-ERT2* mice and mis-expression of constitutively active *Bmpr1a* in the adult cartilage by injecting tamoxifen (TAM)

intraperitoneally at P56. (**B-D′′′**) Longitudinal sections through the adult knee joints of vehicle control (B-D) and mice 7 days (B′-D′), 14 days (B′′-D′′), 28 days (B′′′, D′′′) post induction by TAM injection. Immunoreactivity for pSMAD1/5/8 (B-B′′′), IHH (C-C′′) and ColX (D-D′′′). (**E-E′′**) BrdU incorporation after 7 days (E′) and 14 days (E′′) of TAM injection. (**E**) Vehicle control. Insets display the higher magnification view of the marked regions in (E-E′′). (**F-G′′**) Immunoreactivity for MMP-13 (F-F′′) and ColI (G-G′′) in vehicle control (F, G) and TAM injected knee joints at 60 days (F′, G′) and 180 days (F′′-G′′) post induction. (**H-H′**) Toluidine blue staining in vehicle control (H) and TAM injected knee joints at 180 days (H′) post induction. (**I-I′**) Immunoreactivity for NITEGE in vehicle control (I) and TAM injected knee joints at 180 days (I′) post induction. (**J**) 3-D rendering of μCT scan at 40μm resolution in wildtype (WT) control, vehicle control and TAM injected knee joint at 180 days post induction, black arrows show osteophytes. n=4 per group. Scale bar = 100μm.

Fig. 2. BMP signaling induced transient cartilage differentiation is necessary for the pathogenesis of OA.

(**A**) Schematic representation depicting the generation of *Bmp2^c/c^; Bmp4^c/c^; Col2a1-Cre-ERT2* and the regimen for depletion of BMP signaling by administration of tamoxifen followed by ACLT. (**B-H′′**) Longitudinal sections through the knee joints of sham (B-H), "ACLT + vehicle" control (B′-H′) and "BMP depletion + ACLT" (B′′-H′′) mice at 56 days post-surgery (PS56). Immunoreactivity for pSMAD1/5/8 (B-B′′), ColII (C-C′′), ColX (D-D′′), MMP-13 (E-E′′), NITEGE (G-G′′) and ColI (H-H′′) levels. (F-F′′) Safranin O staining. (**I-I′′**) 3-D rendering of μCT at PS56. Red arrowheads indicate osteophytes, surface roughness, and damage. n=4 per time point per group. Scale bar = 100μm. (**J-K′′′**) Histological sections of knee articular cartilage from OA patients (n=6) (J-J′′′), and a patient without known history of knee OA (n=1) (K-K′′′). (J, K) Safranin O/Fast Green staining of OA (J) and normal (K) cartilage. Immunoreactivity for pSMAD1/5/8 with (J′, K′) or without phosphatase pre-treatment to verify antibody specificity (J′′, J′′′, K′′, K′′′), of OA (J′-J′′′) and normal (K′-K′′′) cartilage. (J′′′, K′′′) Higher magnification view of the marked regions in J′′ and K′′. Scale bar = 50μm.

The panels where *Bmp2/4* depleted animals were subjected to ACLT are marked as "BMP depletion + ACLT". Vehicle injected animals were used as genotype controls ("ACLT + Vehicle". "Sham" refers to *Bmp2^c/c^; Bmp4^c/c^; Col2a1-Cre-ERT2* animals which underwent sham surgery without ACLT.

Fig. 3. Local pharmacological inhibition of BMP signaling halts OA progression in mice.

(**A**) Schematic for local inhibition of BMP signaling using LDN-193189 in surgically induced OA in wildtype mice. (**B-H′′**) Longitudinal sections through the knee joints of sham (B-H), "ACLT + vehicle" control (B′-H′) and "ACLT + LDN-193189" (B′′-H′′) mice at 28 days post-surgery (PS28). Immunoreactivity for pSMAD1/5/8 (B-B′′), ColII (C-C′′), ColX (D-D′′), MMP-13 (E-E′′), NITEGE (F-F′′) and ColI (H-H′′) levels. (G-G′′) Safranin O staining. (**I-I′′**) 3-D rendering of μCT scan at resolution of 5.86 μm per pixel in sham (I), "ACLT + vehicle" control (I′) and "ACLT + LDN-193189" injected (I′′) knee joint at PS28. Red arrows show osteophytes. (**J**) OARSI severity score for "ACLT + vehicle" vs "ACLT + LDN-193189" and "ACLT + vehicle" vs Sham. Scale bar = 100μm, n=4 per group. **** p < 0.0001.

Fig. 4. Inhibition of BMP signaling post onset of surgically induced OA attenuates the severity of OA associated changes.
(**A**) Schematic for local inhibition of BMP signaling, using of LDN-193189, in the knee joint of wildtype mouse post-surgical onset of OA. (**B-F″**) Longitudinal sections through the knee joints of sham (B-F), "ACLT + vehicle" control (B′-F′) and "ACLT + LDN-193189" (B″-F″) mice at 56 days post-surgery (PS56). Immunoreactivity for ColX (B-B″), MMP-13 (C-C″), NITEGE (D-D″) and ColI (F-F″) levels. White dashed lines mark the bone ends of femur and tibia. (E-E″) Toluidine blue staining. (**G-G″**) 3-D rendering of μCT scan at resolution of 5.86 μm per pixel in sham, "ACLT + vehicle" control and "ACLT + LDN-193189" injected knee joint at PS56. Red arrows mark osteophytes. (**H**) OARSI severity score for "ACLT + vehicle" vs "ACLT + LDN-193189", *p = 0.0135, "ACLT + vehicle" vs Sham, ***p = 0.0007. Scale bar = 100μm, n=3 per group,

Fig. 5. Local inhibition of BMP signaling post-onset of surgically induced OA attenuates the severity of OA associated changes for longer duration.

Schematic for local inhibition of BMP signaling, using of LDN-193189, in the knee joint of wildtype mouse post-surgical onset of OA. (**A**). Longitudinal sections through the knee joints of ACLT induced OA mice (**B-D‴**). "ACLT + vehicle" control (B, C and D), "ACLT + LDN-193189 one dose" (B′, C′ and D′), "ACLT + LDN-193189 two doses" (B″, C″ and D″), "ACLT + LDN-193189 three doses" (B‴, C‴ and D‴) mice at 84 days post ACLT. Immunoreactivity for ColII (B- B‴), ColX (C- C‴) and Safranin O staining (D- D‴). Immunoreactivity for SMURF1 in control (E), post ACLT 30 days (E′) and 60 days (E″). Scale bar = 100μm, n=3 per group.

Fig. 6. Effect of local inhibition of BMP signaling on inflammatory responses in a surgically induced osteoarthritic mouse model.
(**A-D″**) Longitudinal sections through the knee joints of sham (A-D), "ACLT + vehicle" control (A′-D′) and "ACLT + LDN-193189" (A″-D″) mice at 56 days post-surgery (PS56). Immunoreactivity for IL-1β (A-A″), iNOS (B-B″), NF-κB (C-C″) and TNF-α (D-D″) levels. White arrows indicate TNF-α positive cells. n=4 per group. Scale bar = 100μm.

Fig. S1. Molecular characterization of mouse knee joint post ACLT
(**A-D‴**) Longitudinal sections through the knee joints of sham (A, C), 7 days (B, D), 14 days (B′, D′), 21 days (B″, D″), 28 days (B‴, D‴) post ACLT. (**A-B‴**) Immunoreactivity for pSMAD1/5/8. (**C-D″**) BrdU uptake. Insets display the higher magnification view of the marked regions.

Fig. S2. BMP signaling induced transient cartilage differentiation is necessary for the pathogenesis of OA.
(**A**) Schematic representation depicting the generation of *Bmp2^{c/c}; Bmp4^{c/c}; Col2a1-Cre-ERT2* and the regimen for depletion of BMP signaling by administration of tamoxifen followed by ACLT. (**B-G″**) Longitudinal sections through the knee joints of sham (B-G), "ACLT + vehicle control" (B′-G′) and "BMP depletion + ACLT" (B″-G″) mice at 28 days post-surgery (PS28).

Immunoreactivity for pSMAD1/5/8 (B-B″), ColII (C-C″), ColX (D-D″), MMP-13 (E-E″), NITEGE (G-G″). (**F-F″**) Safranin O staining. n=4 per group. Scale bar = 100μm.

The panels where *Bmp2/4* depleted animals were subjected to ACLT are marked as "BMP depletion + ACLT". Vehicle injected animals were used as genotype controls ("ACLT + Vehicle").

Fig. S3. Determination of optimum LDN-193189 concentration to inhibit BMP signaling using BRITER (BMP Responsive Immortalized Reporter) cell line.
Normalized relative luciferase activity after 3h of LDN-193189 treatment with different concentration (100nM, 200nM, 400nM) to inhibit BMP signaling in BRITER cell line (Fig.S3).

References

Bandyopadhyay, A., Tsuji, K., Cox, K., Harfe, B. D., Rosen, V. and Tabin, C. J. (2006) 'Genetic analysis of the roles of BMP2, BMP4, and BMP7 in limb patterning and skeletogenesis', *PLoS Genet* 2(12): e216.
Barranco, C. (2014) 'Osteoarthritis: Animal data show VEGF blocker inhibits post-traumatic OA', *Nat Rev Rheumatol* 10(11): 638.
Chen, D., Shen, J., Zhao, W., Wang, T., Han, L., Hamilton, J. L. and Im, H. J. (2017) 'Osteoarthritis: toward a comprehensive understanding of pathological mechanism', *Bone Res* 5: 16044.
Cross, M., Smith, E., Hoy, D., Nolte, S., Ackerman, I., Fransen, M., Bridgett, L., Williams, S., Guillemin, F., Hill, C. L. et al. (2014) 'The global burden of hip and knee osteoarthritis: estimates from the global burden of disease 2010 study', *Ann Rheum Dis* 73(7): 1323-30.
Dahlberg, L., Billinghurst, R. C., Manner, P., Nelson, F., Webb, G., Ionescu, M., Reiner, A., Tanzer, M., Zukor, D., Chen, J. et al. (2000) 'Selective enhancement of collagenase-mediated cleavage of resident type II collagen in cultured osteoarthritic cartilage and arrest with a synthetic inhibitor that spares collagenase 1 (matrix metalloproteinase 1)', *Arthritis Rheum* 43(3): 673-82.
Duprez, D., Bell, E. J., Richardson, M. K., Archer, C. W., Wolpert, L., Brickell, P. M. and Francis-West, P. H. (1996) 'Overexpression of BMP-2 and BMP-4 alters the size and shape of developing skeletal elements in the chick limb', *Mech Dev* 57(2): 145-57.
Enomoto, H., Inoki, I., Komiya, K., Shiomi, T., Ikeda, E., Obata, K., Matsumoto, H., Toyama, Y. and Okada, Y. (2003) 'Vascular endothelial growth factor isoforms and their receptors are expressed in human osteoarthritic cartilage', *Am J Pathol* 162(1): 171-81.
Gelse, K., Ekici, A. B., Cipa, F., Swoboda, B., Carl, H. D., Olk, A., Hennig, F. F. and Klinger, P. (2012) 'Molecular differentiation between osteophytic and articular cartilage--clues for a transient and permanent chondrocyte phenotype', *Osteoarthritis Cartilage* 20(2): 162-71.

Glasson, S. S., Askew, R., Sheppard, B., Carito, B., Blanchet, T., Ma, H. L., Flannery, C. R., Peluso, D., Kanki, K., Yang, Z. et al. (2005) 'Deletion of active ADAMTS5 prevents cartilage degradation in a murine model of osteoarthritis', *Nature* 434(7033): 644-8.

Glasson, S. S., Blanchet, T. J. and Morris, E. A. (2007) 'The surgical destabilization of the medial meniscus (DMM) model of osteoarthritis in the 129/SvEv mouse', *Osteoarthritis Cartilage* 15(9): 1061-9.

Goldring, M. B. and Otero, M. (2011) 'Inflammation in osteoarthritis', *Curr Opin Rheumatol* 23(5): 471-8.

Hao, J., Ho, J. N., Lewis, J. A., Karim, K. A., Daniels, R. N., Gentry, P. R., Hopkins, C. R., Lindsley, C. W. and Hong, C. C. (2010) 'In vivo structure-activity relationship study of dorsomorphin analogues identifies selective VEGF and BMP inhibitors', *ACS Chem Biol* 5(2): 245-53.

Hayami, T., Pickarski, M., Zhuo, Y., Wesolowski, G. A., Rodan, G. A. and Duong, L. T. (2006) 'Characterization of articular cartilage and subchondral bone changes in the rat anterior cruciate ligament transection and meniscectomized models of osteoarthritis', *Bone* 38(2): 234-43.

He, Y., Siebuhr, A. S., Brandt-Hansen, N. U., Wang, J., Su, D., Zheng, Q., Simonsen, O., Petersen, K. K., Arendt-Nielsen, L., Eskehave, T. et al. (2014) 'Type X collagen levels are elevated in serum from human osteoarthritis patients and associated with biomarkers of cartilage degradation and inflammation', *BMC Musculoskelet Disord* 15: 309.

Kahn, J., Shwartz, Y., Blitz, E., Krief, S., Sharir, A., Breitel, D. A., Rattenbach, R., Relaix, F., Maire, P., Rountree, R. B. et al. (2009) 'Muscle contraction is necessary to maintain joint progenitor cell fate', *Dev Cell* 16(5): 734-43.

Kamekura, S., Hoshi, K., Shimoaka, T., Chung, U., Chikuda, H., Yamada, T., Uchida, M., Ogata, N., Seichi, A., Nakamura, K. et al. (2005) 'Osteoarthritis development in novel experimental mouse models induced by knee joint instability', *Osteoarthritis Cartilage* 13(7): 632-41.

Ko, J. Y., Lee, M. S., Lian, W. S., Weng, W. T., Sun, Y. C., Chen, Y. S. and Wang, F. S. (2017) 'MicroRNA-29a Counteracts Synovitis in Knee Osteoarthritis Pathogenesis by Targeting VEGF', *Sci Rep* 7(1): 3584.

Kobayashi, M., Squires, G. R., Mousa, A., Tanzer, M., Zukor, D. J., Antoniou, J., Feige, U. and Poole, A. R. (2005) 'Role of interleukin-1 and tumor necrosis factor alpha in matrix degradation of human osteoarthritic cartilage', *Arthritis Rheum* 52(1): 128-35.

Kozhemyakina, E., Lassar, A. B. and Zelzer, E. (2015) 'A pathway to bone: signaling molecules and transcription factors involved in chondrocyte development and maturation', *Development* 142(5): 817-31.

Lark, M. W., Bayne, E. K., Flanagan, J., Harper, C. F., Hoerrner, L. A., Hutchinson, N. I., Singer, II, Donatelli, S. A., Weidner, J. R., Williams, H. R. et al. (1997) 'Aggrecan degradation in human cartilage. Evidence for both matrix metalloproteinase and aggrecanase activity in normal, osteoarthritic, and rheumatoid joints', *J Clin Invest* 100(1): 93-106.

Lin, A. C., Seeto, B. L., Bartoszko, J. M., Khoury, M. A., Whetstone, H., Ho, L., Hsu, C., Ali, S. A. and Alman, B. A. (2009) 'Modulating hedgehog signaling can attenuate the severity of osteoarthritis', *Nat Med* 15(12): 1421-5.

Long, F. and Ornitz, D. M. (2013) 'Development of the endochondral skeleton', *Cold Spring Harb Perspect Biol* 5(1): a008334.

Mak, K. K., Kronenberg, H. M., Chuang, P. T., Mackem, S. and Yang, Y. (2008) 'Indian hedgehog signals independently of PTHrP to promote chondrocyte hypertrophy', *Development* 135(11): 1947-56.

Mankin, H. J. (1962) 'Localization of Tritiated Thymidine in Articular Cartilage of Rabbits: I. Growth In Immature Cartilage', *JBJS* 44(4): 682-688.

Mankin, H. J. (1963) 'Localization of Tritiated Thymidine in Articular Cartilage of Rabbits: III. MATURE ARTICULAR CARTILAGE', *JBJS* 45(3): 529-540.

Muzumdar, M. D., Tasic, B., Miyamichi, K., Li, L. and Luo, L. (2007) 'A global double-fluorescent Cre reporter mouse', *Genesis* 45(9): 593-605.

Nakamura, E., Nguyen, M. T. and Mackem, S. (2006) 'Kinetics of tamoxifen-regulated Cre activity in mice using a cartilage-specific CreER(T) to assay temporal activity windows along the proximodistal limb skeleton', *Dev Dyn* 235(9): 2603-12.

Nakase, T., Miyaji, T., Tomita, T., Kaneko, M., Kuriyama, K., Myoui, A., Sugamoto, K., Ochi, T. and Yoshikawa, H. (2003) 'Localization of bone morphogenetic protein-2 in human osteoarthritic cartilage and osteophyte', *Osteoarthritis Cartilage* 11(4): 278-84.

Neuhold, L. A., Killar, L., Zhao, W., Sung, M. L., Warner, L., Kulik, J., Turner, J., Wu, W., Billinghurst, C., Meijers, T. et al. (2001) 'Postnatal expression in hyaline cartilage of constitutively active human collagenase-3 (MMP-13) induces osteoarthritis in mice', *J Clin Invest* 107(1): 35-44.

Ortega, N., Behonick, D. J. and Werb, Z. (2004) 'Matrix remodeling during endochondral ossification', *Trends Cell Biol* 14(2): 86-93.

Palazzo, C., Nguyen, C., Lefevre-Colau, M. M., Rannou, F. and Poiraudeau, S. (2016) 'Risk factors and burden of osteoarthritis', *Ann Phys Rehabil Med* 59(3): 134-138.

Pitsillides, A. A. and Beier, F. (2011) 'Cartilage biology in osteoarthritis--lessons from developmental biology', *Nat Rev Rheumatol* 7(11): 654-63.

Ray, A., Singh, P. N., Sohaskey, M. L., Harland, R. M. and Bandyopadhyay, A. (2015) 'Precise spatial restriction of BMP signaling is essential for articular cartilage differentiation', *Development* 142(6): 1169-79.

Roelofs, A. J. and De Bari, C. (2019) 'Immunostaining of Skeletal Tissues', *Methods Mol Biol* 1914: 437-450.

Rolfe, R. A., Nowlan, N. C., Kenny, E. M., Cormican, P., Morris, D. W., Prendergast, P. J., Kelly, D. and Murphy, P. (2014) 'Identification of mechanosensitive genes during skeletal development: alteration of genes associated with cytoskeletal rearrangement and cell signalling pathways', *BMC Genomics* 15: 48.

Sandell, L. J. and Aigner, T. (2001) 'Articular cartilage and changes in arthritis. An introduction: cell biology of osteoarthritis', *Arthritis Res* 3(2): 107-13.

Sanvitale, C. E., Kerr, G., Chaikuad, A., Ramel, M. C., Mohedas, A. H., Reichert, S., Wang, Y., Triffitt, J. T., Cuny, G. D., Yu, P. B. et al. (2013) 'A new class of small molecule inhibitor of BMP signaling', *PLoS One* 8(4): e62721.

Schmal, H., Pilz, I. H., Mehlhorn, A. T., Dovi-Akue, D., Kirchhoff, C., Sudkamp, N. P., Gerlach, U. and Niemeyer, P. (2012) 'Expression of BMP-receptor type 1A correlates with progress of osteoarthritis in human knee joints with focal cartilage lesions', *Cytotherapy* 14(7): 868-76.

Singh, P. N. P., Shea, C. A., Sonker, S. K., Rolfe, R. A., Ray, A., Kumar, S., Gupta, P., Murphy, P. and Bandyopadhyay, A. (2018) 'Precise spatial restriction of BMP signaling in developing joints is perturbed upon loss of embryo movement', *Development* 145(5).

Sokolove, J. and Lepus, C. M. (2013) 'Role of inflammation in the pathogenesis of osteoarthritis: latest findings and interpretations', *Ther Adv Musculoskelet Dis* 5(2): 77-94.

Song, R. H., Tortorella, M. D., Malfait, A. M., Alston, J. T., Yang, Z., Arner, E. C. and Griggs, D. W. (2007) 'Aggrecan degradation in human articular cartilage explants is mediated by both ADAMTS-4 and ADAMTS-5', *Arthritis Rheum* 56(2): 575-85.

Tetlow, L. C., Adlam, D. J. and Woolley, D. E. (2001) 'Matrix metalloproteinase and proinflammatory cytokine production by chondrocytes of human osteoarthritic cartilage: associations with degenerative changes', *Arthritis Rheum* 44(3): 585-94.

Vortkamp, A., Lee, K., Lanske, B., Segre, G. V., Kronenberg, H. M. and Tabin, C. J. (1996) 'Regulation of rate of cartilage differentiation by Indian hedgehog and PTH-related protein', *Science* 273(5275): 613-22.

Vos, T. Flaxman, A. D. Naghavi, M. Lozano, R. Michaud, C. Ezzati, M. Shibuya, K. Salomon, J. A. Abdalla, S. Aboyans, V. et al. (2012) 'Years lived with disability (YLDs) for 1160 sequelae of 289 diseases and injuries 1990-2010: a systematic analysis for the Global Burden of Disease Study 2010', *Lancet* 380(9859): 2163-96.

Yadav, P. S., Prashar, P. and Bandyopadhyay, A. (2012) 'BRITER: a BMP responsive osteoblast reporter cell line', *PLoS One* 7(5): e37134.

Yu, P. B., Deng, D. Y., Lai, C. S., Hong, C. C., Cuny, G. D., Bouxsein, M. L., Hong, D. W., McManus, P. M., Katagiri, T., Sachidanandan, C. et al. (2008) 'BMP type I receptor inhibition reduces heterotopic [corrected] ossification', *Nat Med* 14(12): 1363-9.

Zhou, J., Chen, Q., Lanske, B., Fleming, B. C., Terek, R., Wei, X., Zhang, G., Wang, S., Li, K. and Wei, L. (2014a) 'Disrupting the Indian hedgehog signaling pathway in vivo attenuates surgically induced osteoarthritis progression in Col2a1-CreERT2; Ihhfl/fl mice', *Arthritis Res Ther* 16(1): R11.

Zhou, J., Wei, X. and Wei, L. (2014b) 'Indian Hedgehog, a critical modulator in osteoarthritis, could be a potential therapeutic target for attenuating cartilage degeneration disease', *Connect Tissue Res* 55(4): 257-61.

Zou, H., Wieser, R., Massague, J. and Niswander, L. (1997) 'Distinct roles of type I bone morphogenetic protein receptors in the formation and differentiation of cartilage', *Genes Dev* 11(17): 2191-203.

Fig. 1

Fig. 2

Fig. 3

Fig. 4

Vehicle control	LDN-193189-1 dose	LDN-193189-2 doses	LDN-193189-3 doses

ColII

B   B'   B''   B'''

ColX

C   C'   C''   C'''

Safranin O

D   D'   D''   D'''

Control	Post ACLT- 30 days	Post ACLT - 60 days

SMURF1

E   E'   E''

Fig. 5

Fig. 6

Days post ACLT

| Control | 7 days | 14 days | 21 days | 28 days |

pSMAD1/5/8

BrdU

Fig. S1

Fig. S2

Fig. S3