C$_4$-dicarboxylate metabolons: Interaction of C$_4$-dicarboxylate transporters of *Escherichia coli* with cytosolic enzymes and regulators

Christopher Schubert and Gottfried Unden

Institute for Molecular Physiology (IMP), Microbiology and Wine Research; Johannes Gutenberg-University, Mainz, Germany

Address for correspondence:

Gottfried Unden

Institute for Molecular Physiology (IMP), Microbiology and Wine Research

Hanns-Dieter-Hüsch-Weg 17, BZII

55128 Mainz, Germany

unden@uni-mainz.de

Phone +49-6131-3923550

Abstract Metabolons represent the structural organization of proteins for metabolic or regulatory pathways. Here the interaction of enzymes fumarase FumB and aspartase AspA with the C4-DC transporters DcuA and DcuB of *Escherichia coli* was tested by a bacterial two-hybrid (BACTH) assay *in situ*, or by co-chromatography (mSPINE). DcuB interacted strongly with FumB and AspA, and DcuA with AspA. The *fumB-dcuB* and the *dcuA-aspA* genes encoding the respective proteins are known for their colocalization on the genome and the production of co-transcripts. The data consistently suggest the formation of DcuB/FumB, DcuB/AspA and DcuA/AspA metabolons in fumarate respiration for the uptake of L-malate, or L-aspartate, conversion to fumarate and excretion of succinate after reduction. The DcuA/AspA metabolon catalyzes L-Asp uptake and fumarate excretion in concerted action also to provide ammonia for nitrogen assimilation. The aerobic C4-DC transporter DctA interacted with the regulator EIIA^{Glc} of the *E. coli* glucose phosphotransferase system. It is suggested that EIIA^{Glc} inhibits C4-DC uptake by DctA in the presence of the preferred substrate glucose.

Introduction

Versatile bacteria such as *Escherichia coli* co-ordinate metabolism and cellular physiology on various levels including gene expression and post-translational regulation for optimal adaption to current growth conditions. For stabilization and improvement of linear pathways, the enzymes and proteins can be organized in metabolons. Metabolons represent weak complexes of enzymes catalyzing consecutive reactions in order to transfer intermediates between the enzymes of pathways. Channeling prevents the release of labile or toxic intermediates, supports higher local concentration of the intermediates and prevents drifting to other pathways. This organization is able to increase metabolic efficiency and provide an opportunity to control the flux. High-throughput approaches have been used to identify complexes of weakly interacting proteins, or 'interactomes', in microorganisms, including proteins linking membrane processes to cytoplasmic reactions (Butland *et al.* 2005; Hu *et al.* 2009). However, the general approaches provide limited information on complexes when partner proteins are present in low concentrations or are membrane-bound. Thus, established regulatory complexes of the membrane-integral C4-dicarboxylic acids (C4-DC) sensor kinase DcuS with the coregulatory C4-DC carriers DctA, DcuB or DauA (Kleefeld *et al.* 2009b; Witan *et al.* 2012; Karinou *et al.* 2013; Unden *et al.* 2016) and others were not identified.

Alternatively, (Moraes and Reithmeier 2012) suggested an alternative approach for identifying gene clusters of *E. coli* for metabolons consisting of membrane integral transporters together with cytosolic enzymes for the subsequent pathway. The overview (Moraes and Reithmeier 2012) lists polycistronic operons encoding on the one hand transporters for the uptake of sugars, sugar derivatives, amino acids, C4-DC, and other substrates, and on the other hand, enzymes or regulators for subsequent pathways. For some systems interaction of the proteins has been demonstrated, such as for AmtB-GlnK (ammonia transporter and PII-type regulator GlnK) (van Heeswijk *et al.* 1996; Coutts *et al.* 2002). Remarkably, for most of the genetically co-localized systems interaction of the transporter/enzyme pairs is hypothetical.

The list of potential metabolons (Moraes and Reithmeier 2012) also includes the operons *fumB-dcuB* encoding fumarase FumB and the C4-dicarboxylate transporters DcuB, *dcuA-aspA* encoding the C4-dicarboxylate transporters DcuA and aspartase AspA (Six *et al.* 1994; Golby *et al.* 1998b), and *ttdAB-ttdT* encoding tartrate dehydratase TtdAB and the L-tartrate transporter TtdT (Kim and Unden 2007). Both *dcuA* and *dcuB* are transcribed independently from *fumB* and *aspA*, respectively, but also as *fumB-dcuB* and *dcuA-aspA* cotranscripts (Golby *et al.* 1998b). DcuB is produced only under anaerobic conditions (Zientz *et al.* 1998b; Golby *et al.* 1999) and catalyzes the uptake of L-malate which is then dehydrated by FumB to fumarate (Engel *et al.* 1994; Six *et al.* 1994) (Fig. 1A). Fumarate is reduced to succinate by fumarate reductase in fumarate respiration. Succinate is not further catabolized anaerobically and excreted by DcuB via a L-malate/succinate antiport. DcuB also catalyzes the uptake of fumarate which is used directly in fumarate respiration. In the same way, L-Asp is a substrate for transport by DcuB and used after deamination by AspA to fumarate for fumarate respiration. In any case, the succinate is excreted by C4-DC/succinate antiport with C4-DC indicating any of the C4-DCs L-malate, fumarate, or L-aspartate (Engel *et al.* 1994; Six *et al.* 1994; Janausch *et al.* 2002; Schubert *et al.* 2021).

DcuA on the other hand is expressed constitutively under aerobic and anaerobic conditions (Golby *et al.* 1998b). DcuA is able to substitute DcuB in anaerobic growth (Six *et al.* 1994) and catalyze an C4-DC/succinate antiport for fumarate respiration (Fig. 1A). L-Asp, fumarate or L-malate are accepted as the C4-DCs, but with a clear preference for L-Asp over L-malate or fumarate (Strecker *et al.* 2018). In addition to its role in catabolism and fumarate respiration, DcuA catalyzes the uptake of L-aspartate for anabolism (Strecker *et al.* 2018; Schubert *et al.* 2020). Under these conditions (Fig. 1B) L-aspartate is deaminated in bacteria by AspA to

fumarate and ammonia, which can saturate the complete nitrogen requirement for growth. The uptake of L-Asp depends on DcuA and is coupled to the excretion of fumarate (and some L-malate). The combined action of DcuA and AspA results in the net uptake of ammonia, which is able to supply all the nitrogen for cell synthesis without consuming the carbon, provided that other carbon sources are available.

L-Tartrate that is taken up by TtdT for tartrate fermentation (Fig. 1A), has to be dehydrated by dehydratase TtdAB to oxaloacetate (Reaney *et al.* 1993; Kim and Unden 2007). After conversion to fumarate, it is used in fumarate respiration, and the succinate is excreted by TtdT in antiport against L-tartrate.

In the metabolic pathways, DcuA, DcuB and TtdT are closely linked to AspA in case of DcuA and DcuB, to FumB in the case of DcuB, and to TtdAB in the case of TtdT. To test the metabolon hypothesis for the C4-dicarboxylate metabolism, interaction of DcuA and DcuB with AspA and FumB was tested. Interaction was tested *in vivo* by the adenylate cyclase-based bacterial two-hybrid (BACTH) system which has been established as a reliable assay for protein-protein interactions involving membrane-integral proteins, or *in vitro* by co-purification of the proteins.

Fig. 1: Pathways of (A) fumarate respiration of *E. coli* using fumarate, L-malate, and L-aspartate as the electron acceptors, (B) L-tartrate fermentation and (C) nitrogen assimilation from L-aspartate. For details on (A) and (B) see (Unden *et al.* 2016) and on (C) (Schubert *et al.* 2020). Abbreviations: AspA, aspartase, DcuA, C4-DC transporter DcuA; DcuB, C4-DC transporter DcuB, ET, electron transport; FrdABCD, fumarate reductase; Fum, fumarate; FumB, fumarase B; L-Asp, L-aspartate; L-Mal, L-malate; MKH$_2$, menaquinol; Succ, succinate; TtdT, L-tartrate transporter.

Results and Discussion

DcuB interacts with AspA and FumB.

The adenylate cyclase-based bacterial two hybrid (BACTH) system (Karimova *et al.* 1998; Karimova *et al.* 2001) was used to investigate the interaction of AspA or of FumB with DcuB. The BACTH system relies on the restoration of *Bordetella pertussis* adenylate cyclase (AC) that has been separated genetically in two domains T18 and T25. When the domains are fused to interacting proteins, local vicinity of the domains allows restoration of AC activity due to high flexibility of domain arrangement of *B. pertussis* AC. The activity is tested in a reporter strain deficient of the *E. coli* AC CyaA by expression of the cAMP-CRP dependent β-galactosidase LacZ. The N- and C-termini of DcuB (similar to those of DcuA and DcuC) are located in the periplasm, which excludes the use of the BACTH system under standard conditions since cAMP-CRP resides in the cytoplasm. Therefore, the T25 fragment of the AC was fused for the interaction assays into a cytosolic loop of DcuB (or the other Dcu transporters) (Bauer *et al.* 2011; Witan *et al.* 2012; Wörner *et al.* 2016; Strecker *et al.* 2018). DcuB with the sandwich (SW) T25 domain (DcuB$_{SWT25}$) is membrane-integral and active in co-regulating DcuS (Wörner *et al.* 2016) allowing interaction studies. Bacteria producing the pairs DcuB$_{SWT25}$ with FumB$_{T18}$, and DcuB$_{SWT25}$ with AspA$_{T18}$ showed high β-galactosidase activity (Fig. 2) which exceeded the negative control represented by the non-interacting proteins $_{T18}$AspA and $_{T25}$Zip. The interaction was independent of the presence of transport substrate molecules L-malate and L-aspartate (not shown). On the other hand, bacteria producing the sandwich construct DcuC$_{SWT25}$ and FumB$_{T18}$, or AspA$_{T18}$ respectively, showed only low β-galactosidase activity, which exceeded the background activity only slightly (Fig. 2). DcuC serves as a C4-DC efflux transporter (Zientz *et al.* 1996; Zientz *et al.* 1999; Janausch *et al.* 2002) for succinate derived from fermentation metabolism and not from externally supplied C4-DCs. The data indicates specific protein-protein interaction of DcuB with FumB and with AspA which is not observed for the DcuC efflux transporter that has a different metabolic task.

Fig. 2: Interaction of DcuB and DcuC with AspA and FumB tested in the BACTH system. *E. coli* BTH101(Δ*cyaA*) was co-transformed pairwise with plasmids encoding T25 sandwich constructs with DcuB (DcuB_{SWT25}) or DcuC (DcuC_{SWT25}) and fusions of FumB or AspA to T18 (FumB_{T18}, and AspA_{T18}). The combinations are shown on the x-axis. The leucine zipper pair _{T18}Zip and _{T25}Zip are applied as positive control (Karimova et al. 1998; Karimova et al. 2001). The pair _{T18}Zip/_{T25}AspA represents a negative control for background β-galactosidase activity. The corresponding plasmids are derivatives of pUT18 (FumB_{T18 and} AspA_{T18}) and pKNT25 (DcuB_{SWT25}) (Table 1). Bacteria were grown anaerobically in LB medium. β-Galactosidase activities were quantified in Miller-Units MU (Miller 1992).

DcuA interacts with AspA.

The role of DcuA in L-Asp uptake for fumarate respiration in catabolism, and as a source of nitrogen suggests a close metabolic link to aspartase AspA. C- and N-termini of DcuA are located in the periplasm (Golby *et al.* 1998a) which prevents their use for fusion to the T18 and T25 AC domains for the BACTH system. Insertion of the T25 domain into a cytosolic loop of DcuA transporter in a sandwich fusion (DcuA_{SWT25}) produced a fusion protein in analogy to DcuB_{SWT25}. The fusion protein was located in the membrane but was not active in transport (Strecker *et al.* 2018) or any other biological activity, which makes its application in the BACTH assay questionable. In an alternative approach, DcuA was fused to the bacterial alkaline phosphatase PhoA. PhoA-tagged membrane protein like DcuB often retain biological activity (Bauer *et al.* 2011) and the tag can be used for immunodetection with PhoA antisera. In the

experiment, *E. coli* strain C43 was co-transformed with plasmids encoding AspA-Strep and DcuA-PhoA fusions. Proteins were simultaneously overexpressed and the bacteria were treated with formaldehyde after expression to crosslink proteins that are located in close proximity (Sutherland *et al.* 2008). Detergent extracts of the membranes were then applied to a Strep-Tactin column. After a wash step, the specifically bound proteins were eluted with desthiobiotin. A column treated with the extracts from bacteria that produced DcuA-PhoA and AspA-Strep eluted DcuA-PhoA and AspA-Strep from the column upon addition of desthiobiotin (Fig. 3). DcuA-PhoA was missing, however, in the eluate when either AspA-Strep or DcuA-PhoA were not produced by the bacteria. The experiment demonstrates that interaction with AspA-Strep is required to retard DcuA to the Strep-Tactin column. In addition, binding of DcuA to AspA-Strep depended on formaldehyde cross-linking in the cell homogenate (not shown), supporting the notion that the interaction is too weak to survive column chromatography without stabilization by crosslinking.

Fig. 3: Copurification of DcuA-PhoA with AspA-Strep on a Strep-Tactin column. Strep-tagged AspA (AspA-Strep, encoded by pMW3049) and PhoA-tagged DcuA (DcuA-PhoA, encoded by pMW577) were expressed individually or co-expressed in *E. coli* C43. For protein crosslinking, bacterial cell suspensions were incubated with formaldehyde (0.6 % w/v) for 15 min: Then the bacteria were homogenized in a French press cell and treated with lauryldimethylamine oxide (0.05% v/v). The cleared cell homogenate was applied to a Strep-Tactin column. After washing, the bound proteins were eluted with buffer containing desthiobiotin (2.5 mM). Samples were subjected to SDS-PAGE and Western blotting. Western blots were probed with antibodies against PhoA (upper panel) and Strep-tag (lower panel). Protein standards are shown on the right-hand side. The masses of AspA-Strep and DcuA-PhoA calculated from amino acid composition are 56.0 kDa and 95.2 kDa, respectively.

Function of DcuA/AspA, DcuB/FumB and DcuB/AspA metabolons?

The experiments show specific interaction of the C4-DC transporters DcuA with AspA, and of DcuB with FumB and AspA (Fig. 4). Under anaerobic conditions fumarate represents the substrate for fumarate respiration. DcuB is a major carrier for C4-DC/succinate antiport in fumarate respiration (Six *et al.* 1994; Golby *et al.* 1998b; Janausch *et al.* 2002; Unden *et al.*

2016). The major substrate for DcuB is L-malate, but due to the high levels of L-aspartate in the intestine of animals (Bertin *et al.* 2018; Schubert *et al.* 2021), L-Asp is also transported by DcuB (Six *et al.* 1994; Golby *et al.* 1998b). The interaction of DcuB with AspA and FumB suggests a coordination of the C4-DC uptake with the conversion to fumarate, and the excretion of the succinate after fumarate reduction, which indicates in the presence of DcuB/FumB and DcuB/AspA metabolons (Fig. 4A). A very similar situation applies to the role for a DcuA/AspA metabolon in fumarate respiration (Fig.4A). Overall, DcuB/FumB, DcuB/AspA and DcuA/AspA metabolons are consistently indicated for fumarate respiration by interaction of the proteins, the colocalization of the corresponding genes in the genome and the physiological relevance of a coordination (Fig. 4A). The metabolons await, however, direct proof of metabolic coupling and biochemical backgrounds of the coupling.

Fig. 4: Tentative scheme for the DcuA/AspA, DcuB/AspA and DcuB/FumB complexes and the coordinated reactions in fumarate respiration and nitrogen assimilation from L-Asp. Abbreviations as in Fig. 1.

When L-Asp functions as the nitrogen source for bacterial growth, it can be used as the sole source of nitrogen (Strecker *et al.* 2018). The ammonia released by AspA feeds the route for nitrogen assimilation via glutamine synthetase GS and glutamine 2-oxoglutarate amino transferase GOGAT (Strecker *et al.* 2018). The deaminating activity of AspA is further

stimulated by the regulator GlnB (or PII), which responds to the nitrogen status of the bacteria (Schubert *et al.* 2020). In consequence, from the L-Asp only the ammonia used under conditions of nitrogen assimilation, and fumarate with some L-malate is released by DcuA in near stoichiometric amounts to L-aspartate uptake (Strecker *et al.* 2018). Thus, coordination of the pathway with DcuA and AspA in the DcuA/AspA metabolon (Fig. 4B) is also highly indicated by genetic colocalization of the genes, protein interaction and the co-ordination of the L-Asp/Fum antiport to the ammonia supply.

Interaction of DctA with the regulator EIIA^{Glc} of the phosphotransferase system PTS.

Bacteria producing the aerobic C4-DC transporter DctA and the protein EIIA^{Glc} from the glucose phosphotransferase system were tested for interaction using the bacterial two-hybrid system BACTH. DctA represents the main transporter for the uptake of C4-DCs under aerobic condition, and EIIA^{Glc} a component of the glucose-specific phosphotransferase system (PTS) of *E. coli*. EIIA^{Glc} transfers the phosphoryl group to the glucose transporter EIIBC (Deutscher *et al.* 2006). EIIA^{Glc} has in addition a regulatory role in glucose repression by transmitting the status for glucose availability to the AC CyaA and to some transporters for the uptake of carbon substrates. The N-and C-termini of DctA have cytoplasmic location (Witan *et al.* 2012), and fusions of DctA to the T18 and T25 fragments of *B. pertussis* AC were produced in all combinations, as well as to the N- and C-termini of EIIA^{Glc}. DctA and EIIA^{Glc} fusions showed high β-galactosidase activity at the same level as the positive control or higher than the background activity, indicating an interaction of DctA with EIIA^{Glc} (Fig. 5). Although interaction of DctA by EIIA^{Glc} is strong and evident, the role and physiological consequence is not clear. The interaction is suggested to inhibit DctA and C4-DC uptake in the presence of the preferred substrate glucose (i.e. by dephosphorylated EIIA^{Glc}) and to exclude C4-DCs as alternative substrates from the cell. Inhibition of C4-DC uptake does not result in inducer exclusion, since C$_4$-dicarboxylates are sensed in the periplasm by the two-component system DcuS-DcuR (Pappalardo *et al.* 2003; Unden *et al.* 2016). The supposed model suggesting 'substrate exclusion' is at variance with classical inducer exclusion, exemplified by the inhibition of lactose permease LacY or the maltose ABC transporter (Dills *et al.* 1982; Misko *et al.* 1987; Dean *et al.* 1990). Classical inducer exclusion inhibits uptake of substrates that serve also as inducers in the cytoplasm, thereby inhibiting the induction of alternative metabolic pathways (Deutscher *et al.* 2006). The interaction between DctA and EIIA^{Glc} indicates a further level of

glucose regulation on DctA, in addition to the transcriptional regulation of *dctA* by cAMP-CRP (Davies *et al.* 1999).

Fig. 5: DctA-EIIA^Glc interaction *in vivo* using the BACTH system. *E. coli* BTH101(ΔcyaA) was co-transformed pairwise with plasmids encoding fusions of T25 to DctA/EIIA^Glc (_T25DctA) and fusions of T18 to DctA/EIIA^Glc (_T18DctA). The combinations are shown on the x-axis. The leucine zipper pair _T18Zip and _T25Zip are applied as positive control (Karimova *et al.* 1998; Karimova *et al.* 2001), the pair _T18Zip/_T25DctA as the negative control for background β-galactosidase activity. The corresponding plasmids are derivatives of pUT18C (_T18DctA), pUT18 (DctA_T18), pKT25 (_T25DctA), and pKNT25 (DctA_T25) (Tab. 1). The strains were grown aerobically in LB medium at 30°C to the exponential phase (OD_578 = 0.5). β-Galactosidase activities were quantified in Miller-Units (MU).

Materials and Methods

Bacterial strains and growth conditions. The *E. coli* K12 strains and the plasmids are listed in Table 1. All molecular methods, including cloning, DNA isolation, and manipulations were performed according to standard procedures (Jones and Gunsalus 1987; Sambrook *et al.* 1989; Miller 1992; Zientz *et al.* 1998a; Kleefeld *et al.* 2009a). The oligonucleotide primers are shown in Table 2. Bacteria were grown aerobically and anaerobically at 37°C in lysogeny broth (LB).

β-Galactosidase assay. Cells for the BACTH measurements were grown anaerobically in LB medium. Interactions in the BACTH system were measured in terms of the β-galactosidase activity (Miller 1992). BACTH experiments were conducted as described previously (Monzel *et*

al. 2013) with slight modifications (Wörner *et al.* 2016). Activities are the mean of at least two independent experiments and four replicates each.

Membrane protein interaction (mSPINE). The mSPINE assay for protein interactions was performed as described previously (Graf *et al.* 2014; Wörner *et al.* 2016). *E. coli* strain C43 was co-transformed with plasmids, pMW3049 and pMW577, which encoded the fusion proteins AspA-Strep and DcuA-PhoA, respectively. The bacteria were grown aerobically in 400 ml LB medium to an OD_{578} of 0.5. Protein expression was induced for 3 h with 100 µM L-arabinose and 100 µM anhydrotetracycline (AHT). To crosslink the proteins *in vivo*, the bacteria were incubated with formaldehyde (0.6% w/v) for 15 min at 37°C with shaking. Next, the bacteria were harvested with centrifugation ($11300 \times g$ for 10 min), and the sediment was washed in Tris-HCl buffer (pH 8, 50 mM). For Strep-Tactin purification, cells were disrupted with a French Press (1260 psi) in a buffer that contained lauryldimethylamine oxide (LDAO,0.05% (v/v)). The cell homogenates were clarified by centrifugation ($39100 \times g$ for 30 min at 4°C) and subjected to Strep-Tactin chromatography, 10% SDS-PAGE, and Western blotting. Western blots were probed with anti-Strep and anti-PhoA antisera (IBA Lifesciences, Sigma-Aldrich) (Scheu *et al.* 2010; Graf *et al.* 2014; Wörner *et al.* 2016).

SDS-PAGE, Western Blot, and Immunostaining. SDS-PAGE and Western blots were performed according to published procedures (Graf *et al.* 2014; Wörner *et al.* 2016). Immunostaining was performed with horse-radish peroxidase (HRP)-coupled anti-His-HRP, anti-Strep-HRP, and anti-IgG-mouse-HRP polyclonal antiserum (Sigma-Aldrich) or with anti-PhoA antibodies (produced in mouse, Sigma-Aldrich). For visualization chemiluminescent substrate HRP (Merck Millipore) was used and the blots exposed on X-ray film (Advansta).

Acknowledgments. We are grateful to Dr. A. Strecker (Mainz) for preparing the plasmids with the DcuA_SWT25 and DcuC_SWT25 constructs. Financial support from Deutsche Forschungsgemeinschaft (grant UN 49/19-1) is gratefully acknowledged.

References

Bauer, J., Fritsch, M.J., Palmer, T. and Unden, G. 2011 Topology and accessibility of the transmembrane helices and the sensory site in the bifunctional transporter DcuB of Escherichia coli. *Biochemistry* **50**, 5925–5938.

Bertin, Y., Segura, A., Jubelin, G., Dunière, L., Durand, A. and Forano, E. 2018 Aspartate metabolism is involved in the maintenance of enterohaemorrhagic Escherichia coli O157: H7 in bovine intestinal content. *Environmental microbiology* **20**, 4473–4485.

Butland, G., Peregrín-Alvarez, J.M., Li, J., Yang, W., Yang, X., Starostine, A., Richards, D., Beattie, B., Krogan, N. and Davey, M. 2005 Interaction network containing conserved and essential protein complexes in Escherichia coli. *Nature* **433**, 531–537.

Coutts, G., Thomas, G., Blakey, D. and Merrick, M. 2002 Membrane sequestration of the signal transduction protein GlnK by the ammonium transporter AmtB. *EMBO J* **21**, 536–545.

Davies, S.J., Golby, P., Omrani, D., Broad, S.A., Harrington, V.L., Guest, J.R., Kelly, D.J. and Andrews, S.C. 1999 Inactivation and regulation of the aerobic C4-dicarboxylate transport (dctA) gene ofEscherichia coli. *Journal of Bacteriology* **181**, 5624–5635.

Dean, D.A., Reizer, J., Nikaido, H. and Saier Jr, M.H. 1990 Regulation of the maltose transport system of Escherichia coli by the glucose-specific enzyme III of the phosphoenolpyruvate-sugar phosphotransferase system. Characterization of inducer exclusion-resistant mutants and reconstitution of inducer exclusion in proteoliposomes. *J. Biol. Chem.* **265**, 21005–21010.

Deutscher, J., Francke, C. and Postma, P.W. 2006 How phosphotransferase system-related protein phosphorylation regulates carbohydrate metabolism in bacteria. *Microbiology and Molecular Biology Reviews* **70**, 939–1031.

Dills, S.S., Schmidt, M.R. and Saier Jr, M.H. 1982 Regulation of Lactose Transport by the Phosphoenolpyruvate-Sugar Phosphotransferase System in Membrane Vesicles of Escherichia coli. *Journal of cellular biochemistry* **18**, 239–244.

Engel, P., KRÄMER, R. and Unden, G. 1994 Transport of C4-dicarboxylates by anaerobically grown Escherichia coli. *European Journal of Biochemistry* **222**, 605–614.

Golby, P., Davies, S., Kelly, D.J., Guest, J.R. and Andrews, S.C. 1999 Identification and characterization of a two-component sensor-kinase and response-regulator system (DcuS-DcuR) controlling gene expression in response to C4-dicarboxylates in Escherichia coli. *Journal of Bacteriology* **181**, 1238–1248.

Golby, P., Kelly, D.J., Guest, J.R. and Andrews, S.C. 1998a Topological analysis of DcuA, an anaerobic C4-dicarboxylate transporter of Escherichia coli. *Journal of Bacteriology* **180**, 4821–4827.

Golby, P., Kelly, D.J., Guest, J.R. and Andrews, S.C. 1998b Transcriptional Regulation and Organization of the *dcuA* and *dcuB* Genes, Encoding Homologous Anaerobic C4-Dicarboxylate Transporters in *Escherichia coli*. *Journal of Bacteriology* **180**, 6586–6596.

Graf, S., Schmieden, D., Tschauner, K., Hunke, S. and Unden, G. 2014 The sensor kinase DctS forms a tripartite sensor unit with DctB and DctA for sensing C4-dicarboxylates in Bacillus subtilis. *J. Bacteriol.* **196**, 1084–1093.

Guzman, L.M., Belin, D., Carson, M.J. and Beckwith, J. 1995 Tight regulation, modulation, and high-level expression by vectors containing the arabinose PBAD promoter. *J. Bacteriol.* **177**, 4121–4130.

Hu, P., Janga, S.C., Babu, M., Díaz-Mejía, J.J., Butland, G., Yang, W., Pogoutse, O., Guo, X., Phanse, S. and Wong, P. 2009 Global functional atlas of Escherichia coli encompassing previously uncharacterized proteins. *PLoS Biol* **7**, e1000096.

Janausch, I.G., Zientz, E., Tran, Q.H., Kröger, A. and Unden, G. 2002 C 4-dicarboxylate carriers and sensors in bacteria. *Biochimica et Biophysica Acta (BBA)-Bioenergetics* **1553**, 39–56.

Jones, H.M. and Gunsalus, R.P. 1987 Regulation of Escherichia coli fumarate reductase (frdABCD) operon expression by respiratory electron acceptors and the fnr gene product. *J. Bacteriol.* **169**, 3340–3349.

Karimova, G., Pidoux, J., Ullmann, A. and Ladant, D. 1998 A bacterial two-hybrid system based on a reconstituted signal transduction pathway. *Proceedings of the National Academy of Sciences* **95**, 5752–5756.

Karimova, G., Ullmann, A. and Ladant, D. 2001 Protein-protein interaction between Bacillus stearothermophilus tyrosyl-tRNA synthetase subdomains revealed by a bacterial two-hybrid system. *Journal of molecular microbiology and biotechnology* **3**, 73–82.

Karinou, E., Compton, E.L.R., Morel, M. and Javelle, A. 2013 The Escherichia coli SLC26 homologue YchM (DauA) is a C4-dicarboxylic acid transporter. *Molecular Microbiology* **87**, 623–640.

Kim, O.B. and Unden, G. 2007 The L-tartrate/succinate antiporter TtdT (YgjE) of L-tartrate fermentation in Escherichia coli. *J. Bacteriol.* **189**, 1597–1603.

Kleefeld, A., Ackermann, B., Bauer, J., Kramer, J. and Unden, G. 2009a The fumarate/succinate antiporter DcuB of Escherichia coli is a bifunctional protein with sites for regulation of DcuS-dependent gene expression. *The Journal of biological chemistry* **284**, 265–275.

Kleefeld, A., Ackermann, B., Bauer, J., Krämer, J. and Unden, G. 2009b The fumarate/succinate antiporter DcuB of Escherichia coli is a bifunctional protein with sites for regulation of DcuS dependent gene expression. *Journal of Biological Chemistry* **284**, 265–275.

Miller, J.H. 1992 A] short course in bacterial genetics: a laboratory manual and handbook for Escherichia coli and related bacteria.

Miroux, B. and Walker, J.E. 1996 Over-production of proteins in Escherichia coli: mutant hosts that allow synthesis of some membrane proteins and globular proteins at high levels. *Journal of Molecular Biology* **260**, 289–298.

Misko, T.P., Mitchell, W.J., Meadow, N.D. and Roseman, S. 1987 Sugar transport by the bacterial phosphotransferase system. Reconstitution of inducer exclusion in Salmonella typhimurium membrane vesicles. *J. Biol. Chem.* **262**, 16261–16266.

Monzel, C., Degreif-Dünnwald, P., Gröpper, C., Griesinger, C. and Unden, G. 2013 The cytoplasmic PASC domain of the sensor kinase DcuS of Escherichia coli: role in signal transduction, dimer formation, and DctA interaction. *MicrobiologyOpen* **2**, 912–927.

Moraes, T.F. and Reithmeier, R.A.F. 2012 Membrane transport metabolons. *Biochimica et Biophysica Acta (BBA)-Biomembranes* **1818**, 2687–2706.

Pappalardo, L., Janausch, I.G., Vijayan, V., Zientz, E., Junker, J., Peti, W., Zweckstetter, M., Unden, G. and Griesinger, C. 2003 The NMR structure of the sensory domain of the membranous two-component fumarate sensor (histidine protein kinase) DcuS of Escherichia coli. *J. Biol. Chem.* **278**, 39185–39188.

Reaney, S.K., Begg, C., Bungard, S.J. and Guest, J.R. 1993 Identification of the L-tartrate dehydratase genes (ttdA and ttdB) of Escherichia coli and evolutionary relationship with the class I fumarase genes. *Microbiology (Reading, Engl)* **139**, 1523–1530.

Sambrook, J., Fritsch, E.F. and Maniatis, T. 1989 Molecular cloning: a laboratory manual/J. Sambrook, EF Fritsch, T. Maniatis.

Scheu, P.D., Liao, Y.-F., Bauer, J., Kneuper, H., Basche, T., Unden, G. and Erker, W. 2010 Oligomeric sensor kinase DcuS in the membrane of Escherichia coli and in proteoliposomes: chemical cross-linking and FRET spectroscopy. *Journal of Bacteriology* **192**, 3474–3483.

Schubert, C., Winter, M.G., Ebert-Jung, A., Nagel-Wolfrum, K., Kierszniowska, S., Schramm, T., Link, H., Winter, S.E. and Unden, G. 2021 C4-dicarboxylates and L-aspartate utilization by Escherichia coli K-12 in the mouse intestine: L-aspartate as a major substrate for fumarate respiration and as a nitrogen source. *Submitted*.

Schubert, C., Zedler, S., Strecker, A. and Unden, G. 2020 L-Aspartate as a high-quality nitrogen source in Escherichia coli: Regulation of L-aspartase by the nitrogen regulatory system and interaction of L-aspartase with GlnB. *Molecular Microbiology*.

Six, S., Andrews, S.C., Unden, G. and Guest, J.R. 1994 *Escherichia coli* Possesses Two Homologous Anaerobic C4-Dicarboxylate Membrane Transporters (DcuA and DcuB) Distinct from the Aerobic Dicarboxylate Transport System (Dct). *Journal of Bacteriology* **176**, 6470–6478.

Strecker, A. 2018 Der C4-Dicarboxylat-Metabolismus von Escherichia coli Die Rolle von DcuA und von DcuS im Signaltransfer. PhD thesis, Johannes Gutenberg-University, Mainz, Germany.

Strecker, A., Schubert, C., Zedler, S., Steinmetz, P. and Unden, G. 2018 DcuA of aerobically grown Escherichia coli serves as a nitrogen shuttle (L-aspartate/fumarate) for nitrogen uptake. *Mol Microbiol* **109**, 801–811.

Sutherland, B.W., Toews, J. and Kast, J. 2008 Utility of formaldehyde cross-linking and mass spectrometry in the study of protein-protein interactions. *J Mass Spectrom* **43**, 699–715.

Unden, G., Strecker, A., Kleefeld, A. and Kim, O.B. 2016 C4-Dicarboxylate Utilization in Aerobic and Anaerobic Growth. *EcoSal Plus* **7**.

van Heeswijk, W.C., Hoving, S., Molenaar, D., Stegeman, B., Kahn, D. and Westerhoff, H.V. 1996 An alternative PII protein in the regulation of glutamine synthetase in Escherichia coli. *Molecular Microbiology* **21**, 133–146.

Witan, J., Bauer, J., Wittig, I., Steinmetz, P.A., Erker, W. and Unden, G. 2012 Interaction of the Escherichia coli transporter DctA with the sensor kinase DcuS: presence of functional DctA/DcuS sensor units. *Molecular Microbiology* **85**, 846–861.

Wörner, S., Strecker, A., Monzel, C., Zeltner, M., Witan, J., Ebert-Jung, A. and Unden, G. 2016 Conversion of the sensor kinase DcuS of Escherichia coli of the DcuB/DcuS sensor complex to the C4-dicarboxylate responsive form by the transporter DcuB. *Environmental microbiology* **18**, 4920–4930.

Zientz, E., Bongaerts, J. and Unden, G. 1998a Fumarate regulation of gene expression in Escherichia coli by the DcuSR (dcuSR genes) two-component regulatory system. *Journal of Bacteriology* **180**, 5421–5425.

Zientz, E., Bongaerts, J. and Unden, G. 1998b Fumarate Regulation of Gene Expression in *Escherichia coli* by the DcuSR (*dcuSR* Genes) Two-Component Regulatory System. *Journal of Bacteriology* **180**, 5421–5425.

Zientz, E., Janausch, I.G., Six, S. and Unden, G. 1999 Functioning of DcuC as the C4-dicarboxylate carrier during glucose fermentation by Escherichia coli. *Journal of Bacteriology* **181**, 3716–3720.

Zientz, E., Six, S. and Unden, G. 1996 Identification of a Third Secondary Carrier (DcuC) for Anaerobic C4-Dicarboxylate Transport in *Escherichia coli*: Roles of the Three Dcu Carriers in Uptake and Exchange. *Journal of Bacteriology* **178**, 7241–7247.

Table 1. Strains of *E. coli* and plasmids used

Strain	Genotype	Reference
C43(DE3)	Mutant of BL21(DE3), for expression of membrane proteins	(Miroux and Walker 1996)
BTH101	F- *cya-99, araD139, galE15, galK16, rpsL1 (Strr), hsdR2, mcrA1, mcrB1*	(Karimova *et al.* 1998)
Plasmid	**Genotype**	**Reference**
pKT25	C-Terminal T25 protein fusion plasmid, pSU40 derivative, KanR	(Karimova *et al.* 2001)
pKNT25	N-Terminal T25 protein fusion plasmid, pSU40 derivative, KanR	(Karimova *et al.* 2001)
pUT18	N-Terminal T18 protein fusion plasmid, pUC19 derivative, AmpR	(Karimova *et al.* 2001)
pUT18C	C-Terminal T18 protein fusion plasmid, pUC19 derivative, AmpR	(Karimova *et al.* 2001)
pKT25-Zip	$_{T25}$Zip expression plasmid, pKT25 derivative, KanR	(Karimova *et al.* 1998)
pUT18C-Zip	$_{T18}$Zip expression plasmid, pUT18C derivative, AmpR	(Karimova *et al.* 1998)
pMW2367	AspA$_{T18}$ expression plasmid, pUT18 derivative, AmpR	This study
pMW2368	$_{T18}$AspA expression plasmid, pUT18C derivative, AmpR	This study
pMW3086	FumB$_{T18}$ expression plasmid, pUT18 derivative, AmpR	This study
pMW2787	$_{T25}$EIIA^{Glc} expression plasmid, pKT25 derivative, KanR	This study
pMW2788	EIIA^{Glc}$_{T25}$ expression plasmid, pKNT25 derivative, KanR	This study
pMW2781	DctA$_{T18}$ expression plasmid, pUT18 derivative, AmpR	(Witan *et al.* 2012)
pMW2784	DctA $_{T25}$ expression plasmid, pKNT25 derivative, KanR	(Witan *et al.* 2012)
pMW2376	Expression of DcuC$_{1-229}$-T25-DcuC$_{230-461}$, ('DcuC$_{SWT25}$') pKNT25 derivative, KanR	(Strecker 2018)
pMW1028	Expression of DcuB1-211-T25-213-446 ('DcuB$_{SWT25}$'), pKNT25 derivative, KanR	(Wörner *et al.* 2016)
pASK-IBA3plus	Expression plasmid with tet promoter and C-terminal Strep-tag, AmpR	IBA, Göttingen
pMW3049	pASK-IBA3plus with *aspA-Strep*, AmpR	(Schubert *et al.* 2020)

pBAD18	Expression vector; pBR322 *ori*, arabinose-inducible pBAD promoter, KanR	(Guzman *et al.* 1995)
pMW577	pBAD18 with *dcuA-phoA*, KanR	(Bauer *et al.* 2011)

Table 2: Oligonucleotide primers. Primer for PCR amplification for AspA, FumB and EIIA. Restriction sites and amino acid exchange positions are printed in red.

Primer	Sequence (5' --> 3')	Amino acid exchange/restriction site
BACTH experiments		
BACTH_pKT25_aspA_for	CGCGTCTAGAAGCCATGTCAAACAAC	XbaI
BACTH_pKT25_aspA_rev	CGTTCCCGGGTCTGTTCGCTTTC	SmaI
BACTH_pKT25_fumB_for	CATCATGGATCCGATGTCAAACAAACCC	BamHI
BACTH_pKT25_fumB_rev	GTAGTAGGTACCTTCTTAGTGCAGTTCG	KpnI
BACTH_pKT25_EIIA_for	GCGCGCGGATCCGATGGGTTTGTTCGATAAAC	BamHI
BACTH_pKT25_EIIA_rev	GCTAGCGGTACCGGCGACTTCTTGATGC	KpnI