Circular RNA repertoires are associated with evolutionarily young transposable elements

Franziska Gruhl[1,2‡], **Peggy Janich**[2,3§], **Henrik Kaessmann**[4*], **David Gatfield**[2*]

*For correspondence:
h.kaessmann@zmbh.uni-heidelberg.de (HK); david.gatfield@unil.ch (DG)

Present address: [†]SIB Swiss Institute of Bioinformatics, Lausanne, Switzerland; [‡]Krebsforschung Schweiz, Bern, Switzerland

[1]SIB Swiss Institute of Bioinformatics, Lausanne, Switzerland; [2]Center for Integrative Genomics, University of Lausanne, Lausanne, Switzerland; [3]Krebsforschung Schweiz, CH-3001 Bern, Switzerland; [4]Center for Molecular Biology of Heidelberg University (ZMBH), DKFZ-ZMBH Alliance, Heidelberg, Germany

Abstract Circular RNAs (circRNAs) are found across eukaryotes and can function in post-transcriptional gene regulation. Their biogenesis through a circle-forming backsplicing reaction is facilitated by reverse-complementary repetitive sequences promoting pre-mRNA folding. Orthologous genes from which circRNAs arise, overall contain more strongly conserved splice sites and exons than other genes, yet it remains unclear to what extent this conservation reflects purifying selection acting on the circRNAs themselves. Our analyses of circRNA repertoires from five species representing three mammalian lineages (marsupials, eutherians: rodents, primates) reveal that surprisingly few circRNAs arise from orthologous exonic loci across all species. Even the circRNAs from orthologous loci are associated with young, recently active and species-specific transposable elements, rather than with common, ancient transposon integration events. These observations suggest that many circRNAs emerged convergently during evolution – as a byproduct of splicing in orthologs prone to transposon insertion. Overall, our findings argue against widespread functional circRNA conservation.

Introduction

First described more than forty years ago, circular RNAs (circRNAs) were originally perceived as a curiosity of gene expression, yet they have gained significant prominence over the last decade (reviewed in *Kristensen et al.* (*2019*); *Patop et al.* (*2019*)). Large-scale sequencing efforts have led to the identification of thousands of individual circRNAs with specific expression patterns and, in some cases, specific functions (*Conn et al., 2015*; *Du et al., 2016*; *Hansen et al., 2013*; *Piwecka et al., 2017*). CircRNA biogenesis involves so-called "backsplicing", in which an exon's 3' splice site is ligated onto an upstream 5' splice site of an exon on the same RNA molecule (rather than down-stream, as in conventional splicing). Backsplicing occurs co-transcriptionally and is guided by the

canonical splicing machinery (*Guo et al., 2014*; *Ashwal-Fluss et al., 2014*; *Starke et al., 2015*). It can be facilitated by complementary, repetitive sequences in the flanking introns (*Dubin et al., 1995*; *Jeck et al., 2013*; *Ashwal-Fluss et al., 2014*; *Zhang et al., 2014*; *Liang and Wilusz, 2014*; *Ivanov et al., 2015*). Through intramolecular base-pairing and folding, the resulting hairpin-like structures can augment backsplicing over the competing, regular forward-splicing reaction. Backsplicing seems to be rather inefficient in most cases, as judged by the low circRNA expression levels found in many tissues. For example, it has been estimated that about 60% of circRNAs exhibit expression levels of less than 1 FPKM (fragments per kilobase per million reads mapped) – a commonly applied cut-off below which genes are usually considered to not be robustly expressed (*Guo et al., 2014*). Due to their circular structure, circRNAs are protected from the activity of cellular exonucleases, which is thought to favour their accumulation to detectable steady-state levels and, together with the cell's proliferation history, presumably contributes to their complex spatiotemporal expression patterns (*Alhasan et al., 2015*; *Memczak et al., 2013*; *Bachmayr-Heyda et al., 2015*). Overall higher circRNA abundances have been reported for neuronal tissues (*Westholm et al., 2014*; *Gruner et al., 2016*; *Rybak-Wolf et al., 2015*) and during ageing (*Gruner et al., 2016*; *Xu et al., 2018*; *Cortés-López et al., 2018*).

All eukaryotes (protists, fungi, plants, animals) produce circRNAs (*Wang et al., 2014*). Moreover, it has been reported that circRNAs are frequently generated from orthologous genomic regions across species such as mouse, pig and human (*Rybak-Wolf et al., 2015*; *Venøet al., 2015*), and that their splice sites have elevated conservation scores (*You et al., 2015*). In these studies, circRNA coordinates were transferred between species to identify "conserved" circRNAs. However, the analyses did not distinguish between potential selective constraints actually acting on the circRNAs themselves, from those preserving canonical splicing features of genes in which they are formed (termed "parental genes" in the following). Moreover, even though long introns containing reverse complement sequences (RVCs) appear to be a conserved feature of circRNA parental genes (*Zhang et al., 2014*; *Rybak-Wolf et al., 2015*), the rapid evolutionary changes occurring on the actual repeat sequences present a considerable obstacle to a thorough evolutionary understanding. Finally, concrete examples for experimentally validated, functionally conserved circRNAs are still rather scarce. At least in part, the reason may lie in the difficulty to specifically target circular vs. linear transcript isoforms in loss-of-function experiments; only recently, novel dedicated tools for such experiments have been developed (*Li et al., 2020*). Currently, however, the prevalence of functional circRNA conservation remains overall unclear.

Here, we set out to investigate the origins and evolution of circRNAs; to this end, we generated a comprehensive set of circRNA-enriched RNA sequencing (RNA-seq) data from five mammalian species and three organs. Our analyses unveil that circRNAs are typically generated from a distinct class of genes that share characteristic structural and sequence features. Notably, we discovered that circRNAs are flanked by species-specific and recently active transposable elements (TEs). Our findings support a model according to which the integration of TEs is preferred in introns of genes with similar genomic properties, thus facilitating circRNA formation as a byproduct of splicing around the same exons of orthologous genes across different species. Together, our work suggests that most circRNAs - even when occurring in orthologs of multiple species and com-

74 prising the same exons - may nevertheless not trace back to common ancestral circRNAs but have

75 rather emerged convergently during evolution, facilitated by independent TE insertion events.

Results

A comprehensive circRNA dataset across five mammalian species

To explore the origins and evolution of circRNAs, we generated paired-end RNA-seq data for three

organs (liver, cerebellum, testis) in five species (grey short-tailed opossum, mouse, rat, rhesus

macaque, human) representing three mammalian lineages with different divergence times (mar-

supials; eutherians: rodents, primates) (**Figure 1A**). For optimal cross-species comparability, all

organ samples originated from young, sexually mature male individuals; we used biological trip-

licates (**Supplementary Table 1**), with the exception of human liver (single sample) and rhesus

macaque cerebellum (duplicates). From the RNA extracted from each sample, we generated two

types of libraries; that is, with and without prior treatment of the RNA with the exoribonuclease

RNase R. This strategy allowed us to enrich for circRNAs (in libraries with RNase R treatment) and

to calculate the actual enrichment factors (from the ratio with/without RNase R treatment). Using

a custom pipeline that took into account RNase R enrichment and other factors to remove likely

false-positives and low expression noise (see **Material and Methods** and **Supplementary Table**

2), we then identified circRNAs from backsplice junction (BSJ) reads, estimated circRNA steady-state

abundances, and reconstructed their isoforms (**Supplementary Table 3**, **Figure 1-Figure supple-**

ment 1, **Figure 1-Figure supplement 2**).

In total, following rigorous filtering, we identified 1,535 circRNAs in opossum, 1,484 in mouse,

2,038 in rat, 3,300 in rhesus macaque, and 4,491 circRNAs in human, with overall higher numbers in

cerebellum, followed by testis and liver (**Figure 1A**, **Supplementary Table 4**). Identified circRNAs

were generally small in size, overlapped with protein-coding exons, frequently detectable only in

one of the tissues, and were flanked by long introns (**Figure 1-Figure supplement 3**).

The identification of circRNA heterogeneity and hotspot frequency is determined by sequencing depth and detection thresholds

Many genes give rise to multiple, distinct circRNAs (*Venøet al., 2015*). Such "circRNA hotspots" are

of interest as they may be enriched for genomic features that drive circRNA biogenesis. A previ-

ous study defined hotspots as genomic loci that produced at least ten structurally different, yet

overlapping circRNAs (*Venøet al., 2015*). Reaching a specific number of detectable circRNA species

for a given locus (e.g., ten distinct circRNAs, as in the cited example) is likely strongly dependent

on overall sequencing depth and on the CPM (counts per million) detection cut-off that is applied.

We therefore compared circRNA hotspots identified at different CPM values (0.1, 0.05 and 0.01

CPM); moreover, to capture in a comprehensive fashion the phenomenon that multiple circRNAs

can be generated from a gene, we considered genomic loci already as hotspots if they produced

a minimum of two different, overlapping circRNAs at the applied CPM threshold. As expected, the

number of hotspots – and the number of individual circRNAs that they give rise to – depend on the

chosen CPM threshold (**Figure 1B** for human and rhesus macaque data; **Figure 1-Figure supple-**

ment 4 for other species). Thus, at 0.1 CPM only 16-27% of all detected circRNA-generating loci are

A: Dataset and detected circRNAs

B: CircRNA hotspot loci by CPM

C: % of circRNAs and hotspots in multiple tissues

D: CircRNA expression strength in hotspots

E: UCSC genome browser view for *Kansl1l* hotspot in rat

classified as hotspots. Decreasing the stringency to 0.01 CPM increases the proportion of hotspot loci to 32-45%. At the same time, the fraction of circRNAs that originate from hotspots (rather than from non-hotspot loci) increases from 34-49% (0.1 CPM) to 59-76% (0.01 CPM), and the number of circRNAs per hotspot increases from 2 to 6. Together, these analyses show that with lower CPM thresholds, the number of distinct circRNAs that become detectable per locus increases substantially; the number of detectable individual circRNA-generating loci increases as well, yet this effect is overall smaller. Furthermore, we observed that in many cases the same hotspots produces circRNAs across multiple organs (**Figure 1C**), with typically one predominant circRNA expressed per organ (**Figure 1D**). The *Kansl1l* hotspot locus is a representative example: it is a hotspot in rat, where it produces 6 different circRNAs (**Figure 1E**). It is also a hotspot in all other species and produces 8, 5, 7, and 6 different circRNAs in opossum, mouse, rhesus macaque and human, respectively (data

Figure 1. Study design, samples, datasets and characterisation of circRNA properties and hotspots. A: Phylogenetic tree of species analysed in this study and detected circRNAs. CircRNAs were identified and analysed in five mammalian species (opossum, mouse, rat, rhesus macaque, human) and three organs (liver, cerebellum, testis). Each sample was split and one half treated with RNase R to enrich BSJs. A dataset of high confidence circRNAs was established, based on the enrichment of BSJs in RNase R-treated over untreated samples. To the right of the panel, the total number of circRNAs for each species in liver (brown), cerebellum (green) and testis (blue) is shown. B: CircRNA hotspot loci by CPM (human and rhesus macaque). The graph shows, in grey, the proportion (%) of circRNA loci that qualify as hotspots and, in purple, the proportion (%) of circRNAs that originate from such hotspots, at three different CPM thresholds (0.01, 0.05, 0.1). The average number of circRNAs per hotspot is indicated above the purple bars. C. Number of circRNA hotspot loci found in multiple tissues. The graph shows the proportion (%) of circRNAs (light grey) and of hotspots (dark grey) that are present in at least two tissues. D. Contribution of top-1 and top-2 expressed circRNAs to overall circRNA expression from hotspots. The plot shows the contribution (%) that the two most highly expressed circRNAs (indicated as top-1 and top-2) make to the total circRNA expression from a given hotspot. For each plot, the median is indicated with a grey point. E. Example of the *Kansl1l* hotspot in rat. The proportion (%) for each detected circRNA within the hotspot and tissue (cerebellum = green, testis = blue) are shown. The strongest circRNA is indicated by an asterisk. rnCircRNA-819 is expressed in testis and cerebellum.

Figure 1–Figure supplement 1. Overview of the reconstruction pipeline.

Figure 1–Figure supplement 2. Mapping summary of RNA-seq reads.

Figure 1–Figure supplement 3. General circRNA properties.

Figure 1–Figure supplement 4. CircRNA hotspot loci by CPM (opossum, mouse, rat).

---

not shown).

Overall, we concluded that the expression levels of many circRNAs are low. Increasing the sen-

sitivity of detection (i.e., lowering CPM thresholds) led to a substantial gain in the detectability of

additional, low-expressed circRNA species, but less so of additional circRNA-generating genomic

loci. These findings raised the question whether many of the circRNAs that can be identified re-

flected a form of gene expression noise that occurred preferentially at hotspot loci, rather than

functional transcriptome diversity.

CircRNAs formed in orthologous loci across species preferentially comprise consti-

tutive exons

We therefore sought to assess the selective preservation – and hence potential functionality – of

circRNAs. For each gene, we first collapsed circRNA coordinates to identify the maximal genomic

locus from which circRNAs can be produced (**Figure 2A**). In total, we annotated 5,428 circRNA loci

across all species (**Figure 2A**). The majority of loci are species-specific (4,103 loci; corresponding to

75.6% of all annotated loci); there are only comparatively few instances where circRNAs arise from

orthologous loci in the different species (i.e., from loci that share orthologous exons in correspond-

ing 1:1 orthologous genes; **Figure 2A**). For example, only 260 orthologous loci (4.8% of all loci) give

rise to circRNAs in all five species (**Figure 2A**). A considerable proportion of these shared loci also

correspond to circRNA hotspots (opossum: 28.0%, mouse: 43.6%, rat: 53.0%, rhesus macaque:

46.2%, human: 61.6%; calculated from hotspot counts in **Figure 1B** and loci counts in **Figure 2A**).

Thus, despite applying circRNA enrichment strategies for library preparation and lenient thresh-

olds for computational identification, the number of potentially conserved orthologous circRNAs

is surprisingly low. At first sight, this outcome is at odds with previous reports of higher circRNA

conservation that were, however, frequently based on more restricted cross-species datasets (e.g.

comparison human-mouse in *Rybak-Wolf et al.* (*2015*)). Further analyses confirmed that also in

148 our datasets, it was the use of additional evolutionary species that drove the strong reduction in
149 potentially conserved circRNA candidates – see for example how the addition of the rat or of rhesus
150 macaque datasets affect the human-mouse comparison (**Figure 2-Figure supplement 1B**).

151 We next analysed the properties of circRNA exons and started with phastCons scores, which are
152 based on multiple alignments and known phylogenies and describe conservation levels at single-
153 nucleotide resolution (*Siepel et al., 2005*). To assess whether circRNA exons were distinct from
154 non-circRNA exons in their conservation levels, we calculated phastCons scores for different exon
155 types (circRNA exons, non-circRNA exons, UTR exons). CircRNA exons showed higher phastCons
156 scores than exons from the same genes that were not spliced into circRNAs (**Figure 2B**). This would
157 be the expected outcome if purifying selection acted on functionally conserved circRNAs. How-
158 ever, other mechanisms may be relevant as well; constitutive exons, for example, generally exhibit
159 higher conservation scores than alternative exons (*Modrek and Lee, 2003*; *Ermakova et al., 2006*).
160 We thus analysed exon features in more detail. First, the comparison of phastCons scores between
161 exons of non-parental genes, parental genes and circRNAs revealed that parental genes were *per*
162 *se* highly conserved (**Figure 2B**): 85-95% of the observed median differences between circRNA ex-
163 ons and non-parental genes could be explained by the parental gene itself. Next, we compared the
164 usage of parental gene exons across organs (**Figure 2C**). We observed that circRNA exons are more
165 frequently used in isoforms expressed in multiple organs than non-circRNA parental gene exons.
166 Finally, we analysed the sequence composition at the splice sites, which revealed that GC ampli-
167 tudes (i.e., the differences in GC content at the intron-exon boundary) are significantly higher for
168 circRNA-internal exons than for parental gene exons that were located outside of circRNAs (**Figure
169 2D**).

170 Collectively, these observations (i.e., increased phastCons scores, expression in multiple tissues,
171 increased GC amplitudes) prompt the question whether the exon properties associated with circR-
172 NAs actually reflect at their core an enrichment for constitutive exons. Under this scenario, the sup-
173 posed high conservation of circRNAs may not be directly associated with the circRNAs themselves,
174 but with constitutive exons that the circRNAs contain. Thus, even many of the circRNAs "shared"
175 across species might actually not be homologous. That is, rather than reflecting (divergent) evolu-
176 tion from common ancestral circRNAs (**Figure 2E, left panel**), they may frequently have emerged
177 independently (convergently) during evolution in the lineages leading to the different species, thus
178 potentially representing "analogous" transcriptional traits (**Figure 2E, right panel**).

179 **CircRNA parental genes are associated with low GC content and high sequence**
180 **repetiveness**

181 To explore whether convergent evolution played a role in the origination of circRNAs, we set out to
182 identify possible structural and/or functional characteristics that may establish a specific genomic
183 environment (a "parental gene niche") that would potentially favour analogous circRNA production.
184 To this end, we compared GC content and sequence repetitiveness of circRNA parental vs. non-
185 parental genes.

186 GC content is an important genomic sequence characteristic associated with distinct patterns
187 of gene structure, splicing and function (*Amit et al., 2012*). We realised that the increased GC am-

A: Overlap of collapsed circRNA loci

* #overlapping   * #species-specific

B: PhastCons score by exon type

C: Tissue frequency of exon types

D: Splice site amplitude

E: Alternative models for the evolution of overlapping circRNA loci

divergent evolution

circRNA in species A   homologous   circRNA in species B

circRNA originated in common ancestor

-> circRNAs loci overlap, because they evolved from a common ancestor

convergent evolution

circRNA in species A   analogous   circRNA in species B

circRNA originated independently in species A and B

-> circRNA loci overlap, because of similar genomic constraints

---

188   plitudes at circRNA exon-intron boundaries (see above, **Figure 2D**) were mainly caused by a local

189   decrease of intronic GC content rather than by an increase in exonic GC content (**Supplementary**

190   **Table 5**, **Figure 2-Figure supplement 2**). We subsequently explored the hypothesis that GC con-

191   tent could serve to discriminate parental from non-parental genes and grouped all genes into five

192   categories from low (L) to high (H) GC content (isochores; L1 <37%, L2 37-42%, H1 42-47%, H2 47-

Figure 2. Evolutionary properties of circRNAs. A: CircRNA loci overlap between species. Upper panel: Schematic representation of the orthology definition used in our study. CircRNAs were collapsed for each gene, and coordinates were lifted across species. Lower panel: Number of circRNA loci that are species-specific (red) or circRNAs that arise from orthologous exonic loci of 1:1 orthologous genes (i.e., circRNAs sharing 1:1 orthologous exons) across lineages (purple) are counted. We note that in the literature, other circRNA "orthology" definitions can be found, too. For example, assigning circRNA orthology simply based on parental gene orthology implies calling also those circRNAs "orthologous" that do not share any orthologous exons, which directly argues against the notion of circRNA homology; that is, a common evolutionary origin (see **Figure 2-Figure supplement 1A**). Overall, the orthology considerations we applied largely follow the ideas sketched out in *Patop et al.* (*2019*). B: Distribution of phastCons scores for different exon types. PhastCons scores were calculated for each exon using the conservation files provided by ensembl. PhastCons scores for non-parental exons (grey), exons in parental genes, but outside of the circRNA (pink) and circRNA exons (purple) are plotted. The difference between circRNA exons and non-parental exons that can be explained by parental non-circRNA exons is indicated above the plot. C: Mean tissue frequency of different exon types in parental genes. The frequency of UTR exons (grey), non-UTR exons outside of the circRNA (pink) and circRNA exons (purple) that occur in one, two or three tissues was calculated for each parental gene. D: Distribution of splice site amplitudes for different exon types. Distribution of median splice site GC amplitude (log2-transformed) is plotted for different exon types (np = non-parental, po = parental, but outside of circRNA, pi = parental and inside circRNA). Red vertical bars indicate values at which exon and intron GC content would be equal E: Different evolutionary models explaining the origins of overlapping circRNA loci.

Figure 2–Figure supplement 1. CircRNA loci overlap between species.

Figure 2–Figure supplement 2. Amplitude correlations.

52% and H3 >52% GC content) (**Figure 3A**). Non-parental genes displayed a unimodal distribution in

the two rodents (peak in H1), were generally GC-poor in opossum (peak in L1), and showed a more

complex isochore structure in rhesus macaque and human (peaks in L2 and H3), in agreement with

previous findings (*Galtier and Mouchiroud, 1998*; *Mikkelsen et al., 2007*). Notably, circRNA parental

genes showed a distinctly different distribution than non-parental genes and a consistent pattern

across all five species, with the majority of genes (82-94% depending on species) distributing to the

GC-low gene groups, L1 and L2 (**Figure 3A**).

We next analysed intron repetitiveness – a structural feature that has previously been associ-

ated with circRNA biogenesis. We used megaBLAST to align all annotated coding genes with them-

selves in order to identify regions of complementarity in the sense and antisense orientations of

the gene (reverse complement sequences, RVCs) (*Ivanov et al., 2015*). We then compared the level

of self-complementarity between parental and non-parental genes within the same GC isochore (of

note, self-complementarity generally shows negative correlations with GC-content). This analysis

revealed more pronounced self-complementarity for parental genes than for non-parental genes

(**Figure 3B**).

CircRNA parental genes may also show an association with specific functional properties. Using

data from three human cell studies (*Steinberg et al., 2015*; *Pai et al., 2012*; *Koren et al., 2012*), our

analyses revealed that circRNA parental genes are biased towards early replicating genes, showed

higher steady-state expression levels, and are characterised by increased haploinsufficiency scores

(**Figure 3-Figure supplement 1**). Collectively, we conclude that circRNA parental genes exhibit not

only distinct structural features (low GC content, high repetitiveness), but also specific functional

properties associated with important roles in human cells.

A: GC content of parental genes

B: Complementarity in coding genes

C: GC content vs. exon count

D: PhastCons score vs. RVCs

E: Model of circRNA niche

circRNAs are present in orthologous parental genes with similar properties

repeat age in circRNA niche to distinguish between models of divergent and convergent evolution

divergent evolution

common repeats between shared circRNA loci

convergent evolution

species-specific repeats between shared circRNA loci

Figure 3. Characterisation of circRNA parental gene properties. A: GC content of parental genes. Coding genes were classified into L1-H3 based on their GC content, separately for non-parental (grey) and parental genes (purple). The percentage of parental genes in L1-L2 (opossum, mouse, rat) and L1-H1 (rhesus macaque, human) is indicated above the respective graphs. B: Complementarity in coding genes. Each coding gene was aligned to itself in sense and antisense orientation using megaBLAST. The proportion of each gene involved in an alignment was calculated and plotted against its isochore. C-D: Examples of parental gene predictors for linear regression models. A generalised linear model (GLM) was fitted to predict the probability of the murine coding gene to be parental, whereby x- and y-axis represent the strongest predictors. Colour and size of the discs correspond to the p-values obtained for 500 genes randomly chosen from all mouse coding genes used in the GLM. E. Model of circRNA niche.

Figure 3–Figure supplement 1. Replication time, gene expression steady-state levels and GHIS of human parental genes.

Figure 3–Figure supplement 2. Distribution of prediction values for non-parental and parental circRNA genes.

Figure 3–Figure supplement 3. Properties of 'functional circRNAs' from literature.

Figure 3–Figure supplement 4. Validation of parental gene GLM on Werfel *et al.* dataset.

Figure 3–Figure supplement 5. Properties of highly expressed circRNAs.

Among the multiple predictors of circRNA parental genes, low GC content distinguishes circRNA hotspots

²¹⁷ The above analyses established characteristic sequence, conservation and functional features for
²¹⁸ circRNA parental genes. Using linear regression analyses, we next determined which of these prop-
²¹⁹ erties represented the main predictor(s). We used parental vs. non-parental gene as the response
²²⁰ variable of the model, and several plausible explanatory variables. These were: GC content; exon
²²¹ and transcript counts; genomic length; number of repeat fragments in sense/antisense; expres-
²²² sion level; phastCons score; tissue specificity index. After training the model on a data subset
²²³ (80%), circRNA parental gene predictions were carried out on the remainder of the dataset (20%)
²²⁴ (see **Material and Methods**). Notably, predictions occurred with high precision (accuracy 72-79%,
²²⁵ sensitivity of 75%, specificity 71-79% across all species) and uncovered several significantly asso-
²²⁶ ciated features (**Table 1**, **Supplementary Table 6**, **Figure 3-Figure supplement 2**). Consistently
²²⁷ for all species, the main parental gene predictors are low GC content (log-odds ratio -1.84 to -0.72)
²²⁸ and increased number of exons in the gene (log-odds ratio 0.30 to 0.45). Furthermore, features
²²⁹ positively associated with circRNA production are increased genomic length (log-odds ratio 0.17 to
²³⁰ 0.26), increased proportion of reverse-complementary areas (repeat fragments) within the gene
²³¹ (log-odds ratio 0.20 to 0.59), increased expression levels (log-odds ratio 0.25 to 0.38) and higher
²³² phastCons scores (log-odds ratio 0.45 to 0.58) (**Table 1**, **Figure 3C-D**, **Supplementary Table 6**).
²³³ Notably, parental genes of previously reported functional human circRNAs – e.g., circHipk3 (*Zheng*
²³⁴ *et al., 2016*) and circMbnl1 (*Ashwal-Fluss et al., 2014*) that sequester miRNAs and proteins, respec-
²³⁵ tively – obtain high prediction values in our model and share the above specific properties (**Figure**
²³⁶ **3-Figure supplement 3**). In addition, the identified circRNA parental gene predictors were not re-
²³⁷ stricted to our datasets but could be determined from independent circRNA data as well. Thus, the
²³⁸ analysis of mouse and human heart tissue data (*Werfel et al., 2016*) – on which our linear regres-
²³⁹ sion models predicted parental genes with comparable accuracy (74%), sensitivity (75%) and speci-
²⁴⁰ ficity (74%) – revealed that circRNA parental genes were low in GC content, exon-rich, and showed
²⁴¹ enrichment for repeats (**Figure 3-Figure supplement 4**). In conclusion, the identified properties

[242] likely represent generic characteristics of circRNA parental genes that are suitable to distinguish

[243] them from non-parental genes.

Table 1. A generalised linear model was fitted to predict the probability of coding genes to be a parental gene ($n_{opossum}$=18,807, n_{mouse}=22,015, n_{rat}=11,654, n_{rhesus}=21,891, n_{human}=21,744). The model was trained on 80% of the data (scaled values, cross-validation, 1000 repetitions). Only the best predictors were kept and then used to predict probabilities for the remaining 20% of data points (validation set, shown in table). Genomic length, number of exons and GC content are based on the respective ensembl annotations; number of repeats in antisense and sense orientation to the gene was estimated using the RepeatMasker annotation, phastCons scores taken from UCSC (not available for opossum and rhesus macaque) and expression levels and the tissue specificity index based on (*Brawand et al., 2011*). An overview of all log-odds ratios and p-values calculated in the validation set of each species is provided in the table, further details can be found in **Supplementary Table 6**. *Abbreviations: md = opossum, mm = mouse, rn = rat, rm = rhesus macaque, hs = human. Significance levels: '***' < 0.001, '**' < 0.01, '*' < 0.05, 'ns' >= 0.05.*

Predictor	Log-odds range (significance)	Species with significant predictor
Genomic gene length (bp)	rn: 0.26 (***) rm: 0.17 (***) hs: 0.26 (***) md, mm: ns	rn, rm, hs
Number of exons	md: 0.45 (***) mm: 0.38 (***) rn: 0.30 (***) rm: 0.42 (***) hs: 0.32 (***)	md, mm, rn, rm, hs
GC content	md: -1.84 (***) mm: -1.09 (***) rn: -0.72 (***) rm: -1.44 (***) hs: -1.42 (***)	md, mm, rn, rm, hs
Repeat fragments (antisense)	md: 0.28 (**) mm: 0.20 (**) rm: 0.59 (***) rn, hs: ns	md, mm, rm
Repeat fragments (sense)	hs: 0.58 (***) md, mm, rn, rm: ns	hs
PhastCons scores	mm: 0.58 (***) rn: 0.51 (***) hs: 0.45 (***)	mm, rn, hs
Mean expression levels	md: 0.34 (**) rm: 0.38 (***) hs: 0.25 (**) mm, rn: ns	md, rm, hs
Tissue specificity index	md, mm, rn, rm, hs: ns	-

Many circRNAs are formed from circRNA hotspots (**Figure 1C**). We therefore asked whether among the features that our regression analysis identified for parental genes, some would be suitable to further distinguish hotspots. First, we assessed whether hotspots were more likely to be shared between species than parental genes that produced only a single circRNA isoform. The applied regression model indeed detected a positive correlation between the probability of a parental gene being a hotspot and having orthologous parental genes across multiple species (**Supplementary Table 7**); moreover, log-odds ratios increased with the distance and number of species across which the hotspot was shared (e.g., mouse: 0.29 for shared within rodents, 0.67 for shared with eutherian species and 0.72 for shared within therian species). We next interrogated whether any particular feature would be able to specify circRNA hotspots among parental genes. A single factor, low GC content, emerged as a consistent predictor for circRNA hotspots among all circRNA-generating loci (**Supplementary Table 8**). As expected, the predictive power was lower than that of the previous models, which were designed to discriminate parental vs. non-parental genes and which had identified low GC content as well. These findings imply that hotspots emerge across species in orthologous loci that offer similarly favourable conditions for circRNA formation, most importantly low GC content. The increased number of circRNAs that become detectable when CPM thresholds are lowered (see above, **Figure 1C**) is also in agreement with the sporadic formation of different circRNAs whenever genomic circumstances allow for it. Overall, our observations suggest that differences between hotspot and non-hotspot loci, or between high and low abundance circRNAs, are quantitative rather than qualitative in nature. Thus, the comparison of high vs. low expression circRNAs (based on 90% expression quantile; below = low, above = high expression) indicated the same set of properties, albeit amplified, in the highly expressed circRNAs (**Supplementary Table 9**). Parental genes of highly expressed circRNAs in opossum, rhesus macaque and human yielded higher prediction values in our generalised linear model, which was consistently driven by low GC content (**Supplementary Table 9**). High expression circRNAs were also more likely to be expressed in all three tissues (**Figure 3-Figure supplement 5A**) and to originate from a hotspot (**Figure 3-Figure supplement 5B**), and they were more often shared across multiple species (**Figure 3-Figure supplement 5C**, **Supplementary Table 10**).

Collectively, our analyses thus reveal that circRNA parental genes are characterised by a set of distinct features: low GC content, increased genomic length and number of exons, higher expression levels and increased phastCons scores (**Figure 3E**). These features were detected independently across species, suggesting the presence of a unique, syntenic genomic niche in which circRNAs can be produced ("circRNA niche"). While helpful to understand the genomic context of circRNA production, these findings do not yet allow us to distinguish between the two alternative models of divergent and convergent circRNA evolution (**Figure 2E**). To elucidate the evolutionary trajectory and timeline underlying the emergence of the circRNAs, we sought to scrutinize the identified feature "complementarity and repetitiveness" of the circRNA niche. Previous studies have associated repetitiveness with an over-representation of small TEs – such as primate Alu elements or the murine B1 elements – in circRNA-flanking introns; these TEs may facilitate circRNA formation by providing RVCs that are the basis for intramolecular base-pairing of nascent RNA molecules (*Ivanov et al., 2015*; *Jeck et al., 2013*; *Zhang et al., 2014*; *Wilusz, 2015*; *Liang and Wilusz, 2014*). In-

terestingly, while the biogenesis of human circRNAs has so far been mainly associated with the
primate-specific (i.e., evolutionarily young) Alu elements, a recent study has highlighted several
circRNAs that rely on the presence of the more ancient, mammalian MIR elements (*Yoshimoto*
et al., 2020). A comprehensive understanding of the evolutionary age of TEs in circRNA-flanking
introns could thus provide important insights into the modes of circRNA emergence: the presence
of common (i.e., old) repeats would point towards divergent evolution of circRNAs from a common
circRNA ancestor, whereas an over-representation of species-specific (i.e., recent) repeats would
support the notion of convergent circRNA evolution (**Figure 3E**).

CircRNA flanking introns are enriched in species-specific TEs

Using our cross-species datasets, we investigated the properties and composition of the repeat
landscape relevant for circRNA biogenesis – features that have remained poorly characterised so
far. As a first step, we generated for each species a background set of "control introns" from non-
circRNA genes that were matched to the circRNA flanking introns in terms of length distribution and
GC content. We then compared the abundance of different repeat families within the two intron
groups. In all species, TEs belonging to the class of Short Interspersed Nuclear Elements (SINEs) are
enriched within the circRNA flanking introns as compared to the control introns. Remarkably, the
resulting TE enrichment profiles were exquisitely lineage-specific, and even largely species-specific
(**Figure 4A**). In mouse, for instance, the order of enrichment is from the B1 class of rodent-specific B
elements (strongest enrichment and highest frequency of >7.5 TEs per flanking intron) to B2 and B4
SINEs. In rat, B1 (strong enrichment, yet less frequent than in mouse) is followed by ID (Identifier)
elements, which are a family of small TEs characterised by a recent, strong amplification history
in the rat lineage (*Kim et al., 1994*; *Kim and Deininger, 1996*); B2 and B4 SINEs only followed in 3rd
and 4th position. In rhesus macaque and human, Alu elements are the most frequent and strongly
enriched TEs (around 14 TEs per intron), consistent with the known strong amplification history in
the common primate ancestor (reviewed in *Batzer and Deininger* (*2002*)) (**Figure 4A**). The opossum
genome is known for its high number of TEs, many of which may have undergone a very species-
specific amplification pattern (*Mikkelsen et al., 2007*). This is reflected in the distinct opossum
enrichment profile (**Figure 4-Figure supplement 1**).

As pointed out above, TEs are relevant for circRNA formation because they can provide RVCs
for the intramolecular base-pairing of nascent RNA molecules (*Ivanov et al., 2015*; *Jeck et al., 2013*;
Zhang et al., 2014; *Wilusz, 2015*; *Liang and Wilusz, 2014*). Pre-mRNA folding into a hairpin with a
paired stem (formed by the flanking introns via the dimerised RVCs) and an unpaired loop region
(carrying the future circRNA) leads to a configuration that brings backsplice donor and acceptor
sites into close proximity, thus facilitating circRNA formation. In order to serve as efficient RVCs via
this mechanism, TEs likely need to fulfil certain criteria. Thus, the dimerisation potential is expected
to depend on TE identity, frequency, and position. In the simplest case, two integration events
involving the same TE (in reverse orientation) will lead to an extended RVC stretch. Yet also different
transposons belonging to the same TE family will show a certain degree of sequence similarity
that depends on their phylogenetic distance; sequence differences that have evolved are likely to
compromise the base-pairing potential. To account for such effects, we sought to calculate the

A: Enrichment of transposable elements in flanking introns

B: Top–5 dimer contribution

C: Repeat phylogeny, mouse

D: TE phylogentic age, mouse

E: MFE dimers, mouse

F: TE pairing score (age + MFE), mouse

Figure 4. Analysis of the repeat landscape of circRNA parental genes. A: Enrichment of TEs in flanking introns for mouse, rat, rhesus macaque and human. The number of TEs was quantified in both intron groups (circRNA flanking introns and length- and GC-matched control introns). Enrichment of TEs is represented by colour from high (dark purple) to low (grey). The red numbers next to the TE name indicate the top-3 enriched TEs in each species. Enrichment was assessed using a Wilcoxon Signed Rank Test; p-values are indicated at the bottom of each plot. B: Top-5 dimer contribution. The graph shows the proportion of top-5 dimers (purple) vs. other, remaining dimers (white) to all predicted dimers in flanking introns. Top-5 dimers thus account for 78, 50, 55, 43 and 38% of all dimers in opossum, mouse, rat, rhesus and human, respectively. C: Phylogeny of mouse TEs. Clustal-alignment based on consensus sequences of TEs. Most recent TEs are highlighted. D: PCA for phylogenetic age of mouse TE families. PCA is based on the clustal-alignment distance matrix for the reference sequences of all major SINE families in mouse with the MIR family used as an outgroup. TEs present in the top-5 dimers are labelled. E: PCA based on binding affinity of mouse TE families. PCA is based on the minimal free energy (MFE) for all major SINE families in mouse with the MIR family used as an outgroup. TEs present in the top-5 dimers are labelled. F: PCA for TE pairing score of mouse dimers. PCA is based on a merged and normalised score, taking into account binding strength of the dimer structure (= MFE) and phylogenetic distance. Absolute frequency of TEs is visualised by circle size. TEs present in the five most frequent dimers (top-5) are highlighted by blue lines connecting the two TEs engaged in a dimer (most frequent dimer in dark blue = rank 1). If the dimer is composed of the same TE family members, the blue line loops back to the TE (= blue circle).

Figure 4–Figure supplement 1. Enrichment of transposable elements in flanking introns for opossum.

Figure 4–Figure supplement 2. PCA and phylogeny of opossum, rat, rhesus macaque and human repeat dimers.

Given the high abundance of young, still active transposons in the respective genomes (**Figure 4A**),
we suspected that simply basing our further analyses of dimerisation potential on phylogenetic dis-
tance between different TEs would not provide sufficient resolution. Indeed, as shown for mouse
(**Figure 4C-D**), phylogenetic age separates large subgroups, but not TEs of the same family whose
sequences have diverged by relatively few nucleotides. By contrast, classification by binding affini-
ties creates more precise, smaller subgroups that lack, however, the information on phylogenetic
age (**Figure 4E**). Therefore, we combined both age and binding affinity information into an overall
"pairing score" (see **Material and Methods**). Principal component analysis (PCA) showed that this
measure efficiently separated different TE families and individual family members, with PC1 and
PC2 explaining approximately 76% of observed variance (**Figure 4F**; **Figure 4-Figure supplement**
2). Importantly, this analysis suggests that the most frequently occurring dimers (top-5 dimers are
depicted with blue connecting lines in **Figure 4F**) are formed by recently active TE family members.
In mouse, an illustrative example are the dimers formed by the B1_Mm, B1_Mus1 and B1_Mus2
elements (**Figure 4F**), which are among the most recent (and still active) TEs in this species (**Figure**
4C). Across species, our analyses allowed for the same conclusions. For example, the dominant
dimers in rat were the recently amplified ID elements, and not the more abundant (yet older in
their amplification history) B1 family of TEs (**Figure 4-Figure supplement 2B**) (*Kim et al., 1994*;
Kim and Deininger, 1996). In opossum, the most prominent dimers consisted of opossum-specific
SINE1 elements, which are similar to the Alu elements in primates, but possess an independent
origin (**Figure 4-Figure supplement 2A**) (*Gu et al., 2007*). Finally, within the primate lineage, the
dimer composition was more uniform, probably due to the high amplification rate of the AluS sub-
family (>650,000 copies) in the common ancestor of Old World monkeys and the relatively recent
divergence time of macaque and human (**Figure 4-Figure supplement 2C-D**) (*Deininger, 2011*).

In conclusion, the above analyses of RVCs revealed that dimer-forming sequences in circRNA
flanking introns were most frequently composed of recent, and often currently still active, TEs.
Therefore, the dimer repertoires were specific to the lineages (marsupials, rodents, primates) and/or

(as most clearly visible within the rodent lineage) even species-specific.

Flanking introns of shared circRNA loci are enriched in evolutionarily young TEs

We next compared the dimer composition of introns from shared vs. species-specific circRNA loci.
We reasoned that in the case of shared circRNA loci that have evolved from a common, ancestral
circRNA, we would detect evidence for evolutionarily older TE integration events and shared dimers
as compared to species-specific, younger circRNA loci. For our analysis, we took into account the
frequency, enrichment and age of the TEs and, moreover, their degradation rate (milliDiv; see
below) and the minimal free energy (MFE) of the dimer structure.

First, we analysed the dimer composition of flanking introns in shared and species-specific
circRNA loci. We extracted the top-100 most and least frequent dimers of all circRNA loci, and
compared their enrichment factors and mean age (categorised for simplicity into four groups: 1 =
species-specific, 2 = lineage-specific, 3 = eutherian, 4 = therian) across the two groups of parental
genes (shared and species-specific). The analysis revealed that the most frequent dimers are con-
sistently formed by the youngest elements in both groups of genes, and that the frequency dis-
tribution of the top-100 dimers was significantly different between species (see **Figure 5A** for rat
and human; other species in **Figure 5-Figure supplement 1**). In rat, for instance, all top-5 dimers
are composed of repeats from the youngest ID family members; in human, dimers involving AluY
elements are strongly enriched (**Figure 5A**). On average, most dimers occur at least once or twice
per shared circRNA gene, corresponding to a 1.4- to 2.1-fold enrichment in comparison to species-
specific circRNA loci (**Supplementary Table 11**). Conceivably, the multiple resulting dimerisation
possibilities could act cumulatively to position circRNA exons for backsplicing. Furthermore, we
observed that many RVCs overlapped each other, so that one repeat in one RVC could dimerise
with different repeats in multiple other RVCs. Due to the increased frequency of young repeat el-
ements in shared circRNA loci, these "co-pairing possibilities" further increase the number of pos-
sible dimers that can be formed (**Figure 5-Figure supplement 2**). A representative example for
a shared circRNA-generating locus with its complex dimer interaction landscape, involving young
species-specific repeats, is the *Akt3* locus (**Figure 5B**). Thus, although *Akt3* circRNAs are shared
between human (upper panel), mouse (middle panel) and opossum (lower panel), the dimer land-
scapes are entirely specifies-specific (see top-5 dimers that are highlighted in the figure).

The above observations suggest that circRNA-producing genes act as "transposon sinks" that
are prone to insertions of active repeats. Continuously attracting new transposons could con-
tribute to the mechanism that sustains backsplicing and underlies reproducible circRNA expres-
sion levels. Moreover, through the recurring addition of new functional repeats, new dimerisation
potential would be generated that could make older TEs redundant and allow them to rapidly de-
grade, thus explaining why ancient TE integration events are no longer detectable. If a circRNA
is functionally important for the organism, especially the young, dimerisation-competent repeats
may evolve under purifying selection and maintain their pairing ability. We therefore reasoned
that low degradation rates in young dimers of shared circRNA loci could hint at functionality. We
followed up this idea by analysing the degradation rates of repeats based on their milliDiv values.
Briefly, the RepeatMasker annotations (*Smit et al., 2013*) (http://repeatmasker.org; see **Material**

A: Dimer enrichment (shared vs. species-specific circRNA loci)

Rat

Human

B: Examples of repeat landscape

Human, *AKT3* (chr1)

Mouse, *Akt3* (chr1)

Opossum, *AKT* (chr2)

C: MilliDivs and MFE for top-5 dimers and their repeats, human

milliDivs

MFE

Figure 5. Repeat analysis and dimer potential of shared and species-specific parental genes A: Dimer enrichment in shared vs. species-specific repeats in rat and human (see **Figure 5-Figure supplement 1** for other species). The frequency (number of detected dimers in a given parental gene), log2-enrichment (shared vs. species-specific) and mean age (defined as whether repeats are species-specific: age = 1, lineage-specific: age = 2, eutherian: age = 3, therian: age = 4) of the top-100 most frequent and least frequent dimers in parental genes with shared and species-specific circRNA loci in rat and human were analysed. The frequency is plotted on the x- and y-axis, point size reflects the age and point colour the enrichment (blue = decrease, red = increase). Based on the comparison between shared and species-specific dimers (using a Wilcoxon Signed Rank Test), the top-5 dimers defined by frequency and enrichment are highlighted and labelled in red. B: Species-specific dimer landscape for the *Akt3* gene in human, mouse and opossum. UCSC genome browser view for the parental gene, circRNAs and top-5 dimers (as defined in panel B). Start and stop positions of each dimer are connected via an arc. Dimers are grouped by composition represented by different colours, the number of collapsed dimers is indicated to the right-side of the dimer group. Only dimers that start before and stop after a circRNAs are shown as these are potentially those that can contribute to the hairpin structure. The human *Akt3* gene possesses two circRNA clusters. For better visualisation, only the upstream cluster is shown. C: Degradation rates (MilliDivs) and minimal free energy (MFE) for top-5 dimers in human. MilliDiv values for all repeats composing the top-5 dimers (defined by their presence in all parental genes) were compared between parental genes of species-specific (red) and shared (blue) circRNA loci in human (see **Figure 5-Figure supplement 3 for other species**). A Wilcoxon Signed Rank Test was used to compare dimers between parental genes with shared and species-specific circRNA loci, with p-values plotted above the boxplots. MFE values were compared between the least degraded dimers in parental genes of species-specific (red) and shared (blue) circRNA loci. MFE values were calculated using the genomic sequences of all top-5 dimers. For each parental gene, the least degraded dimer (based on its mean milliDiv value) was then chosen which let to a strong enrichment of only a subset of the top-5 dimers (in this case AluSx+AluY and AluSx1+AluY). If enough observations for a statistical test were present, the two distributions (shared/species-specific) where compared using a Student's t-Test and plotted as violin plots with p-values above the plot.

Figure 5–Figure supplement 1. Contribution of species-specific repeats to the formation of shared circRNA loci.

Figure 5–Figure supplement 2. Repeat interaction landscape in shared vs. species-specific circRNA loci.

Figure 5–Figure supplement 3. MilliDivs and MFE for dimers in shared and species-specific circRNA loci.

and Methods) provide a quantification of how many "base mismatches in parts per thousand"
have occurred between each specific repeat copy in its genomic context and the repeat reference
sequence. This deviation from the consensus sequence is expressed as the milliDiv value. Thus, a
high milliDiv value implies that a repeat is strongly degraded, typically due to its age (the older the
repeat, the more time its sequence has had to diverge). Low milliDiv values suggest that the repeat
is younger (i.e., it had less time to accumulate mutations) or that purifying selection prevented the
accumulation of mutations.

Following this rationale, we determined in each species the degradation rates for the repeats
forming the top-5 dimers. Comparing their milliDiv values in shared vs. species-specific parental
genes revealed no significant differences in any of the species (**Figure5C – left panel**, **Figure 5-**
Figure supplement 3 – left panel). Because degradation rates alone may not fully capture the
actual decline in pairing strength within a dimer (e.g., compensatory changes and dimer length
are not/poorly accounted for), we further analysed actual binding energies. To this end, we se-
lected the least-degraded dimer for every parental gene in both groups (shared/species-specific)
and calculated the minimal free energies (MFEs) of dimer formation. We detected no difference
between the groups, suggesting that dimers of shared circRNA loci are not subject to a specific se-
lection pressure, but degrade identically to dimers in species-specific circRNA loci (**Figure 5C – right**
panel, **Figure 5-Figure supplement 3 – right panel)**. Furthermore, we observed that dimers com-
prising "intermediate age" repeats (i.e. B1_Mur2, B1_Mur3, B1_Mur4, present in Muridae) could be
found in the species-specific "least-degraded" dimers, yet they were absent from the shared group,

which rather contained the top-1/top-2 most enriched and youngest dimers (e.g. AluSx+AluY and AluSx1+AluY in human **Figure 5C**; ID_Rn1+ID_Rn1 and ID_Rn1+ID_Rn2 in rat) (**Figure 5C**, **Figure 5-Figure supplement 3C**).

Taken together, we conclude that circRNAs are preferentially formed from loci that have attracted transposons in recent evolutionary history. Even in the case of shared circRNA loci the actual repeat landscapes, dimer predictions, transposon ages and degradation rates, as well as RVC pairing energies, are most consistent with the model that circRNAs are analogous features that have been formed by convergent evolution, rather than homologous features originating from a common circRNA ancestor.

Discussion

Different mechanistic scenarios to explain the origins and evolution of circRNAs have been considered in the field (reviewed in *Patop et al.* (*2019*)). In our study, we have investigated this topic through the analysis of novel, dedicated cross-species datasets. Notably, we propose that many circRNAs have not evolved from common, ancestral circRNA loci, but have emerged independently through convergent evolution, most likely driven by structural commonalities of their parental genes. Thus, the modelling of parental genes uncovered features that are associated with circRNA biogenesis, in support of the concept of a "circRNA niche" in which circRNAs are more likely to be generated: genetic loci giving rise to circRNAs are generally long, exon-rich and located in genomic regions of low GC content. In the case of orthologous parental genes, these structural characteristics are shared as well, and they have led to shared integration biases for transposons, i.e. to shared, genomic "TE hotspots".

It is well established that intronic TE insertions are critical for circRNA biogenesis as they provide reverse-complementary sequences for intramolecular pre-mRNA folding via TE dimers, giving rise to the secondary structures that facilitate productive backsplicing. Important new insights that our study provides on circRNA evolution come from the deep analysis of the transposon landscapes, including the TE identities, their ages, degradation rates and dimerisation potentials. Thus, because the actual TEs predicted as most relevant for dimerisation are mostly not shared across species and are evolutionarily young, we propose that the resulting circRNAs are evolutionarily young as well. In line with this interpretation, circRNAs from orthologous genes frequently do not involve exactly the same 5' and 3' backsplice sites and thus do not encompass precisely the same orthologous exons, but show partial exon overlap across species (see **Figure 2-Figure supplement 1**). These findings all argue for a model of convergent evolution at shared circRNA loci, with circRNAs and TEs co-evolving in a species-specific and dynamic manner.

Our model provides an explanation for how circRNAs can arise from orthologous exonic loci across species even if they themselves are not homologous (i.e., they do not stem from common evolutionary precursors that emerged in common ancestors). Importantly, if most circRNAs are evolutionarily young, then, by extension, it is overall rather unlikely that they fulfil crucial functions. This idea is in agreement with the generally low expression levels of circRNAs that have been reported and with accumulation patterns that are frequently tissue-specific and confined to post-mitotic cells (*Guo et al., 2014*; *Westholm et al., 2014*). Importantly, these and other main con-

clusions of our study overlap with those of two independent manuscripts (with complementary data and analyses) that have appeared in press (*Xu and Zhang, 2021*) and as a publication preprint (*Santos-Rodriguez et al., 2021*), respectively, while we were preparing the revised version of our manuscript.

Why is it frequently the same (orthologous) genes that produce circRNAs, and why do the circRNA hotspots often overlap between species, i.e. they share common exons? A plausible explanation lies in how TE integration is tolerated. Briefly, intronic TE integration in the vicinity of an intron-exon boundary will likely alter local GC content. For example, GC-rich SINE elements integrating close to a splice site would locally increase intronic GC and thereby decrease the GC amplitude at the intron-exon boundary. Especially in GC-low environments, this can interfere with the intron-defined mechanism of splicing and cause mis-splicing (*Amit et al., 2012*). By contrast, TE integration close to a very strong splice site with a strong GC amplitude – as typically found in canonical exons – would have lower impact. Hence, it would be tolerated better than integration close to alternative exons, whose GC amplitudes are less pronounced. Indeed, our analyses show that circRNA exons are typically canonical exons with strong GC amplitudes. While at first sight, circRNA exons thus appear to be endowed with rather specific, evolutionarily relevant properties - most notably with increased phastCons scores - it is probable that these are a mere consequence of a higher tolerance for TE integration in introns flanking canonical exons.

Many additional characteristics associated with circRNAs – identified in this study or previously by others – can be linked to how the impact of TEs on splicing and transcript integrity is likely to be tolerated. Depending on the site of TE integration, potentially hazardous "transcript noise" will arise, and these instances will be subject to purifying selection. In particular, TE integration into exons (changing the coding sequence) or directly into splice sites (affecting splicing patterns) will lead to erroneous transcripts (*Zhang et al., 2011*). Thus, the probability that an integration event is tolerated, will be overall lower in short and compact genes as compared to genes with long introns; of note, long genes are also GC-poor (*Zhu et al., 2009*). These characteristics overlap precisely with those that we identify for circRNAs, which are also frequently generated from GC-poor genes with long introns, complex gene structures, and that contain many TEs.

An interesting feature – not analysed in our study, but previously associated with circRNAs – is RNA editing. In particular, introns bracketing circRNAs are enriched in A-to-I RNA editing events, and the RNA-editing enzyme ADAR1 has been reported as a specific regulator of circRNA expression (*Ivanov et al., 2015*; *Rybak-Wolf et al., 2015*). However, A-to-I editing is also a well-known defense mechanism that has evolved to suppress TE amplification. For example, A-to-I RNA editing is associated with intronic Alu elements to inhibit Alu dimers (*Lev-Maor et al., 2008*; *Athanasiadis et al., 2004*). Therefore, it is quite likely that associations between RNA editing and circRNA abundances are a secondary effect from the primary purpose of A-to-I editing, namely the inhibition of Alu amplification. A similar case can be made for DNA methylation that interferes with TE amplification (*Yoder et al., 1997*) and has been linked to circRNA production (*Enuka et al., 2016*). Or, in the case of N^6-methyladenosine (m^{6}A), it has recently been proposed that this highly prevalent RNA modification is also involved in dynamically regulating circRNA abundances (*Zhou et al., 2017*; *Park et al., 2019*; *Di Timoteo et al., 2020*). Yet the link of circRNAs to m^{6}A, which is known to influence

498 many steps of mRNA metabolism (reviewed in *Zaccara et al.* (*2019*); *Lee et al.* (*2020*)), may simply
499 reflect the general targeting of erroneous transcripts for degradation.

500   In summary, our evolutionary data and the above considerations lead us to conclude that many
501 circRNAs are likely a form of transcript noise - or, more precisely, of mis-splicing - that is provoked
502 by TE integration into parental genes. This conclusion is in full agreement with the observation that
503 in rat neurons, there is a direct correspondence between the pharmacological inhibition of canon-
504 ical splicing and increased circRNA formation, preferentially affecting circRNAs with long introns
505 and many transposons/RVCs (*Wang et al., 2019*). Altogether, these conclusions make it likely that
506 the majority of circRNAs do not have specific molecular functions, although functional circRNAs
507 have arisen during evolution, as demonstrated in several studies (e.g. *Hansen et al.* (*2013*); *Conn*
508 *et al.* (*2015*); *Du et al.* (*2016*)), presumably from initially non-functional (noise) variants whose emer-
509 gence was facilitated by the aforementioned mechanisms. During this process, a functional cir-
510 cRNA may ultimately even become independent from the original RVC-based regulation. Evolving
511 from a sequence-based backsplice mechanism to a protein-based one (i.e., relying on RNA-binding
512 proteins, RBPs) could render regulation more versatile and more controllable. Indeed, RBPs have
513 emerged as important regulators of several circRNAs (see e.g. *Ashwal-Fluss et al.* (*2014*); *Conn*
514 *et al.* (*2015*); *Okholm et al.* (*2020*)). The functions of circRNAs seem to be diverse and may often in-
515 volve the positive or negative regulation of their own parental genes at different expression layers
516 (transcription/splicing, translation, post-translational modification) through various mechanisms
517 (e.g., competition with linear mRNA splicing, microRNA sponge effects, mRNA traps) (*Shao et al.,*
518 *2021*). For several of these functional roles, the exact exons/exon portions that form the circRNA,
519 or which elements in the flanking introns drive the process, may not be important, but rather the
520 general maintenance of circularization at a locus during evolution. In this way, diverting mRNA
521 output to non-functional, dead-end circular transcripts could for example represent a mechanism
522 to limit parental gene expression or to control genes that have transformed into transposon sinks.

523   Finally, we would like to note that circRNAs have emerged as reliable disease biomarkers (*Mem-*
524 *czak et al., 2015*; *Bahn et al., 2015*), and their utility for such predictive purposes is not diminished
525 by our conclusion that most circRNAs are unlikely to fulfil direct functions – on the contrary. Even
526 if an altered circRNA profile will likely not indicate causal involvement in a disease, it could hint at
527 misregulated transcription or splicing of the parental gene, at a novel TE integration event, or at
528 problems with RNA editing or methylation machineries. The careful analysis of the circRNA land-
529 scape may thus teach us about factors contributing to diseases in a causal fashion even if many or
530 perhaps most circRNAs may not be functional but rather represent transcript noise.

531 ## Material and Methods

532 ## Data deposition, programmes and working environment

Table 2. Overview of external programmes.

Programme	Version
Blast	2.2.29+
BEDTools	2.17.0
Bowtie2	2.1.0
Clustal Omega	1.2.4
Cufflinks	2.1.1
FastQC	0.10.1
Mcl	14.137
R	3.0 and 3.1
Ruby	2.0 and 2.1
SAMTools	0.1.19
TopHat2	2.0.11
ViennaRNA	2.1.8

533 The raw data and processed data files discussed in this publication have been deposited in NCBI's
534 Gene Expression Omnibus (*Edgar et al., 2002*) and are accessible through the GEO Series accession
535 number GSE162152. All scripts used to produce the main figures and tables of this publication
536 have been deposited in the Git Repository circRNA_paperScripts. This Git repository also holds
537 information on how to run the scripts, and links to the underlying data files for the main figures.
538 The custom pipeline developed for the circRNA identification can be found in the Git Repository
539 ncSplice_circRNAdetection.

540 ## Library preparation and sequencing
541 We used 5 µg of RNA per sample as starting material for all libraries. For each biological replicate
542 (= *tissue X* of *Animal 1* of a given species) two samples were taken: sample 1 was left untreated,
543 sample 2 was treated with 20 U RNase R (Epicentre/Illumina, Cat. No. RNR07250) for 1 h at 37°C to
544 degrade linear RNAs, followed by RNA purification with the RNA Clean & Concentrator-5 kit (Zymo
545 Research) according to the manufacturer's protocol. Paired-end sequencing libraries were pre-
546 pared from the purified RNA with the Illumina TruSeq Stranded Total RNA kit with Ribo-Zero Gold
547 according to the protocol with the following modifications to select larger fragments: 1.) Instead of
548 the recommended 8 min at 68°C for fragmentation, we incubated samples for only 4 min at 68°C
549 to increase the fragment size; 2.) In the final PCR clean-up after enrichment of the DNA fragments,
550 we changed the 1:1 ratio of DNA to AMPure XP Beads to a 0.7:1 ratio to select for binding of larger
551 fragments. Libraries were analysed on the fragment analyzer for their quality and sequenced with
552 the Illumina HiSeq 2500 platform (multiplexed, 100 cycles, paired-end, read length 100 nt).

Identification and quantification of circRNAs

Mapping of RNA-seq data

The ensembl annotations for opossum (monDom5), mouse (mm10), rat (rn5), rhesus macaque (rheMac2) and human (hg38) were downloaded from Ensembl to build transcriptome indexes for mapping with TopHat2. TopHat2 was run with default settings and the *–mate-inner-dist* and *–mate-std-dev* options set to 50 and 200 respectively. The mate-inner-distance parameter was estimated based on the fragment analyzer report.

Table 3. Ensembl genome versions and annotation files for each species.

Species	Genome	Annotation
Opossum	monDom5	ensembl release 75, feb 2014
Mouse	mm10	ensembl release 75, feb 2014
Rat	rn5	ensembl release 75, feb 2014
Rhesus macaque	rheMac2	ensembl release 77, oct 2014
Human	hg38	ensembl release 77, oct 2014

Analysis of unmapped reads

We developed a custom pipeline to detect circRNAs (**Figure1-Figure supplement 1**), which performs the following steps: Unmapped reads with a phred quality value of at least 25 are used to generate 20 bp anchor pairs from the terminal 3' and 5'-ends of the read. Anchors are remapped with bowtie2 on the reference genome. Mapped anchor pairs are filtered for 1) being on the same chromosome, 2) being on the same strand and 3) for having a genomic mapping distance to each other of a maximum of 100 kb. Next, anchors are extended upstream and downstream of their mapping locus. They are kept if pairs are extendable to the full read length. During this procedure a maximum of two mismatches is allowed. For paired-end sequencing reads, the mate read not mapping to the backsplice junction can often be mapped to the reference genome without any problem. However, it will be classified as "unmapped read" (because its mate read mapping to the backsplice junction was not identified by the standard procedure). Next, all unpaired reads are thus selected from the accepted_hits.bam file generated by TopHat2 (singletons) and assessed for whether the mate read (second read of the paired-end sequencing read) of the anchor pair mapped between the backsplice coordinates. All anchor pairs for which 1) the mate did not map between the genomic backsplice coordinates, 2) the mate mapped to another backsplice junction or 3) the extension procedure could not reveal a clear breakpoint are removed. Based on the remaining candidates, a backsplice index is built with bowtie2 and all reads are remapped on this index to increase the read coverage by detecting reads that cover the BSJ with less than 20 bp, but at least 8 bp. Candidate reads that were used to build the backsplice index and now mapped to another backsplice junction are removed. Upon this procedure, the pipeline provides a first list of backsplice junctions. The set of scripts, which performs the identification of putative BSJs, as well as a short description of how to run the pipeline are deposited in the Git Repository nc-

Splice_circRNAdetection.

Trimming of overlapping reads

Due to small DNA repeats, some reads are extendable to more than the original read length. There-
fore, overlapping reads were trimmed based on a set of canonical and non-canonical splice sites.
For the donor site GT, GC, AT, CT were used and for the acceptor splice site AG and AC. The trim-
ming is part of our custom pipeline described above, and the step will be performed automatically
if the scripts are run.

Generation of high confidence circRNA candidates from the comparison of RNase R-
treated vs. -untreated samples

The detection of circRNAs relies on the identification of BSJs. These are, however, often only cov-
ered by a low number of reads, which carries considerable risk of mistaking biological or techni-
cal noise for a real circRNA event. Their circular structure makes circRNAs resistant to RNase R
treatment - a feature that is not generally expected for spurious RNA molecules that are linear
but may nevertheless resemble BSJs. We therefore compared BSJs between RNase R-treated and
-untreated samples and determined whether BSJs detected in an untreated sample are enriched
in the RNase R-treated sample. To generate a high-confidence dataset of circRNA candidates from
the comparison of untreated and treated samples (**Figure 1-Figure supplement 1**), we applied the
following filtering steps (please also consult **Supplementary Table 2** for a step-by-step description
of filtering outcomes, using the mouse samples as an example.)

Filtering step 1 - mapping consistency of read pairs. When mapping paired-end sequencing
data, both reads should ideally map to the genome (paired-end = "pe"). However, in some cases
one of the mate reads cannot be mapped due to the complexity of the genomic locus. These reads
are reported as "singletons" ("se"). For each potential BSJ, we thus analysed the mapping behaviour
of both read mates. BSJs for which read pairs in the untreated and RNase R-treated sample of the
same biological replicate mapped both either in "pe" or "se" mode were kept; BSJs for which for
example a read pair mapped in "pe" mode in the untreated biological sample, but in "se" mode in
the RNase R-treated sample of the same biological replicate (and vise versa) were considered weak
candidates and removed. This filtering step removed approximately 1% of the total, unique BSJs
detected (**Supplementary Table 2**).

Filtering step 2 - presence of a BSJ in untreated samples. We hypothesized that for circRNAs
to be functionally important, they should generally be expressed at levels that are high enough to
make them detectable in the normal samples, i.e. without RNase R treatment. We thus removed
all BSJs which were only present in RNase R-treated samples, but undetectable in any of the un-
treated, biological replicates (cut-off for absence/presence = minimum one read mapping to BSJ).
This filtering step removed approximately 75% of the initially detected BSJs (**Supplementary Table
2**).

Filtering step 3 - enrichment after RNase R treatment. RNase R treatment leads to the
enrichment of BSJs in the total number of detected junctions due to the preferential degradation
of linear RNAs. To calculate the enrichment factor, BSJs were normalised by the size factor (as

described in **Material and Methods**, section *Reconstruction of circRNA isoforms*) of each sample

and the mean normalised count was calculated for each condition (untreated and RNase R-treated).

Next, the log2-enrichment for RNase R-treated vs. -untreated samples was calculated. All BSJs for

which the log2-enrichment was below 1.5 were removed. This filtering step removed another 15%

of the originally detected unique BSJs (**Supplementary Table 2**).

Filtering step 4 - minimum expression levels. CPM (counts per million) values for BSJs were

calculated for each tissue as follows:

$$counts = \frac{counts\_rep1 + counts\_rep2 + counts\_rep3}{3}$$

$$total\,Mapped\,Reads = \frac{mapped\,Reads\_rep1 + mapped\,Reads\_rep2 + mapped\,Reads\_rep3}{3}$$

$$CPM = \frac{counts \cdot 10^6}{total\,Mapped\,Reads}$$

All BSJs with at least 0.05 CPM were kept. These loci were considered strong circRNA candidates

and used for all subsequent analyses. After this final filtering step, less than 1% of the original BSJs

are left (**Supplementary Table 2**).

Manual filtering steps

We observed several genomic loci in rhesus macaque and human that were highly enriched in

reads for putative BSJs (no such problem was detected for opossum, mouse and rat). Manual

inspection in the UCSC genome browser indicated that these loci are highly repetitive. The detected

BSJs from these regions probably do not reflect BSJs, but instead issues in the mapping procedure.

These candidates were thus removed manually; the concerned regions are:

Table 4. Removed regions during mapping.

species	tissue	chromosome	start	stop	strand
rhesus macaque	testis	7	164261343	164283671	+
rhesus macaque	testis	7	22010814	22092409	-
rhesus macaque	testis	19	52240850	52288425	-
rhesus macaque	testis	19	59790996	59834798	+
rhesus macaque	testis	19	59790996	59847609	+
human	testis	2	178535731	178600667	+
human	testis	7	66429678	66490107	-
human	testis	9	97185441	97211487	-
human	testis	12	97492460	97561047	+
human	testis	14	100913431	100949596	+
human	testis	18	21765771	21849388	+

638 All following analyses were conducted with the circRNA candidates that remained after this step.

639 Reconstruction of circRNA isoforms

640 To reconstruct the exon structure of circRNA transcripts in each tissue, we made use of the junction

641 enrichment in RNase R treated samples. To normalise junction reads across libraries, the size

642 factors based on the geometric mean of common junctions in untreated and treated samples were

643 calculated as

$$geometric\_mean = \left(\prod x \right)^{\frac{1}{length(x)}}$$

$$size\_factor = median \left(\frac{x}{geometric\_mean} \right)$$

with x being a vector containing the number of reads per junction. We then compared read cover-

age for junctions outside and inside the BSJ for each gene and used the log2-change of junctions

outside the backsplice junction to construct the expected background distribution of change in

junction coverage upon RNase R treatment. The observed coverage change of junctions inside the

backsplice was then compared to the expected change in the background distribution and junc-

tions with a log2-change outside the 90% confidence interval were assigned as circRNA junctions;

a loose cut-off was chosen, because involved junctions can show a decrease in coverage if their lin-

ear isoform was present at high levels before (degradation levels of linear isoforms do not correlate

with the enrichment levels of circRNAs). Next, we reconstructed a splicing graph for each circRNA

candidate, in which network nodes are exons connected by splice junctions (edges) (*Heber et al.,*

2002). Connections between nodes are weighted by the coverage in the RNase R treated samples.

The resulting network graph is directed (because of the known circRNA start and stop coordinates),

acyclic (because splicing always proceeds in one direction), weighted and relatively small. We used

a simple breadth-first-search algorithm to traverse the graph and to define the strength for each

possible isoform by its mean coverage. Only the strongest isoform was considered for all subse-

quent analyses.

660 Reconstruction and expression quantification of linear mRNAs

We reconstructed linear isoforms based on the pipeline provided by *Trapnell et al.* (*2012*) (Cufflinks

+ Cuffcompare + Cuffnorm). Expression levels were quantified based on fragments per million

mapped reads (FPKM). Cufflinks was run per tissue and annotation files were merged across tissues

with Cuffcompare. Expression was quantified with Cuffnorm based on the merged annotation file.

All programs were run with default settings. FPKM values were normalised across species and

tissues using a median scaling approach as described in *Brawand et al.* (*2011*).

667 **Identification of shared circRNA loci between species**

668 Definition and identification of shared circRNA loci

Shared circRNA loci were defined on three different levels depending on whether the "parental

gene", the "circRNA locus" in the gene or the "start/stop exons" overlapped between species (see

Figure 2A and **Figure 2-Figure supplement 1A**). Overall considerations of this kind have recently

also been outlined in *Patop et al.* (*2019*).

Level 1 - Parental genes: One-to-one (1:1) therian orthologous genes were defined between
opossum, mouse, rat, rhesus macaque and human using the Ensembl orthology annotation (con-
fidence intervals 0 and 1, restricted to clear one-to-one orthologs). The same procedure was per-
formed to retrieve the 1:1 orthologous genes for the eutherians (mouse, rat, rhesus macaque,
human), for rodents (mouse, rat) and primates (rhesus macaque, human). Shared circRNA loci be-
tween species were assessed by counting the number of 1:1 orthologous parental genes between
the five species. The analysis was restricted to protein-coding genes.

Level 2 - circRNA locus: To identify shared circRNA loci, all circRNA exon coordinates from a given
gene were collapsed into a single transcript using the *bedtools merge* option from the BEDTools
toolset with default options. Next, we used liftOver to compare exons from the collapsed transcript
between species. The minimal ratio of bases that need to overlap for each exon was set to 0.5 (-
minMatch=0.5). Collapsed transcripts were defined as overlapping between different species if they
shared at least one exon, independent of the exon length.

Level 3 - start/stop exon: To identify circRNAs sharing the same first and last exon between
species, we lifted exons coordinates between species (same settings as described above, *liftOver,*
-minMatch=0.5). The circRNA was then defined as "shared", if both exons were annotated as start
and stop exons in the respective circRNAs of the given species. Note, that this definition only
requires an overlap for start and stop exons, internal circRNA exons may differ.

Given that only circRNAs that comprise corresponding (1:1 orthologous exons) in different
species might at least potentially and reasonably considered to be homologous (i.e., might have
originated from evolutionary precursors in common ancestors) and the Level 3 definition might
require strong evolutionary conservation of splice sites (i.e., with this stringent definition many
shared loci may be missed), we decided to use the level 2 definition (circRNA locus) for the analy-
ses presented in the main text, while we still provide the results for the Level 1 and 3 definitions
in the supplement (**Figure 2-Figure supplement 1A**). Importantly, defining shared circRNA loci at
this level allows us to also compare circRNA hostspots which have been defined using a similar
classification strategy.

Clustering of circRNA loci between species

Based on the species set in which shared circRNA loci were found, we categorised circRNAs in the
following groups: species-specific, rodent, primate, eutherian and therian circRNAs. To be part of
the rodent or primate group, the circRNA has to be expressed in both species of the lineage. To
be part of the eutherian group, the circRNA has to be expressed in three species out of the four
species mouse, rat, rhesus macaque and human. To be part of the therian group, the circRNA
needs to be expressed in opossum and in three out of the four other species. Species-specific
circRNAs are either present in one species or do not match any of the other four categories. The
usage of multiple species for defining shared loci, allowed to define "mammalian circRNAs" with
high confidence (**Figure 2-Figure supplement 1B**). To define the different groups, we used the
cluster algorithm MCL (***Enright et al., 2002***; ***Dongen, 2000***). MCL is frequently used to reconstruct
orthology clusters based on blast results. It requires input in *abc* format (file: *species.abc*), in which
a corresponds to event a, *b* to event b and a numeric value *c* that provides information on the

connection strength between event a and b (e.g. blast p-value). If no p-values are available as in

this analysis, the connection strength can be set to 1. MCL was run with a cluster granularity of 2

(*option -I*).

$ mcxload -abc species.abc –stream-mirror -o species.mci -write-tab species.tab

$ mcl species.mci -I 2

$ mcxdump -icl out.species.mci.I20 -tabr species.tab -o dump.species.mci.I20

PhastCons scores

Codings exons were selected based on the attribute "transcript_biotype = protein_coding" in the gtf

annotation file of the respective species and labelled as circRNA exons if they were in our circRNA

annotation. Exons were further classified into UTR-exons and non-UTR exons using the ensembl

field "feature = exon" or "feature = UTR". Since conservation scores are generally lower for UTR-

exons (*Pollard et al., 2010*), any exon labelled as UTR-exon was removed from further analyses to

avoid bias when comparing circRNA and non-circRNA exons. Genomic coordinates of the remain-

ing exons were collapsed using the *merge* command from the BEDtools toolset (*bedtools merge

input_file -nms -scores collapse*) to obtain a list of unique genomic loci. PhastCons scores for all

exon types were calculated using the conservation scores provided by the UCSC genome browser

(mouse: phastCons scores based on alignment for 60 placental genomes; rat: phastCons scores

based on alignment for 13 vertebrate genomes; human: phastCons scores based on alignment

for 99 vertebrate genomes). For each gene type (parental or non-parental), the median phastCons

score was calculated for each exon type within the gene (if non-parental: median of all exons; if

parental: median of exons contained in the circRNA and median of exons outside of the circRNA).

Tissue specificity of exon types

Using the DEXseq package (from HTSeq 0.6.1), reads mapping on coding exons of the parental

genes were counted. The exon-bins defined by DEXseq (filtered for bins >=10 nt) were then mapped

and translated onto the different exon types: UTR-exons of parental genes, exons of parental genes

that are not in a circRNA, circRNA exons. For each exon type, an FPKM value based on the exon

length and sequencing depth of the library was calculated.

$$FPKM = \frac{counts\_for\_exon\_type \cdot 10^9}{exon\_type\_length / sequencing\_depth}$$

Exons were labelled as expressed in a tissue, if the calculated FPKM was at least 1. The maximum

number of tissues in which each exon occurred was plotted separately for UTR-exons, exons out-

side the circRNA and contained in it.

GC amplitude

The ensembl annotation for each species was used to retrieve the different known transcripts in

each coding gene. For each splice site, the GC amplitude was calculated using the last 250 intronic

bp and the first 50 exonic bp (several values for the last *n* intronic bp and the first *m* exonic bp were tested beforehand, the 250:50 ratio was chosen, because it gave the strongest signal). Splice sites were distinguished by their relative position to the circRNA (flanking, inside or outside). A one-tailed and paired Mann-Whitney U test was used to assess the difference in GC amplitude between circRNA-related splice sites and others.

Definition of highly expressed circRNAs

For each species and tissues, circRNAs were grouped into lowly expressed and highly expressed circRNAs based on whether they were found below or above the 90% expression quantile of the respective tissue. Candidates from different tissues were then merged to obtain a unique list of highly expressed circRNAs for each species.

Parental gene analysis

GC content of exons and intron

The ensembl annotation for each species was used to retrieve the different known transcripts in each coding gene. Transcripts were collapsed per-gene to define the exonic and intronic parts. Introns and exons were distinguished by their relative position to the circRNA (flanking, inside or outside). The GC content was calculated based on the genomic DNA sequence. On a per-gene level, the median GC content for each exon and intron type was used for further analyses. Differences between the GC content were assessed with a one-tailed Mann-Whitney U test.

Gene self-complementarity

The genomic sequence of each coding gene (first to last exon) was aligned against itself in sense and antisense orientation using megaBLAST with the following call:

$ blastn -query seq.fa -subject seq.fa -task dc-megablast -word_size 12 -outfmt "6 qseqid qstart qend sseqid sstart send sstrand length pident nident mismatch bitscore evalue" > blast.out

The resulting alignments were filtered for being purely intronic (no overlap with any exon). The fraction of self-complementarity was calculated as the summed length of all alignments in a gene divided by its length (first to last exon).

Generalised linear models

All linear models were developed in the R environment. The presence of multicollinearity between predictors was assessed using the *vif()* function from the R package *car* (version 3.0.3) to calculate the variance inflation factor. Predictors were scaled to be able to compare them with each other using the *scale()* function as provided in the R environment.

For parental genes, the dataset was split into training (80%) and validation set (20%). To find the strongest predictors, we used the R package *bestglm* (version 0.37). Each model was fitted on the complete dataset using the command *bestglm()* with the information criteria set to "CV" (CV = cross validation) and the number of repetitions *t = 1000*. The model family was set to "binomial" as

we were merely interested in predicting the presence (1) or absence (0) of a parental gene. Signifi-

cant predictors were then used to report log-odds ratios and significance levels for the validation

set using the default *glm()* function of the R environment. Log-odds ratios, standard errors and

confidence intervals were standardised using the *beta()* function from the *reghelper* R package (ver-

sion 1.0.0) and are reported together with their p-values in **Supplementary Table 6**. The same

approach was used to predict which parental genes are likely to be a circRNA hotspot with the only

difference that the underlying data was filtered for parental genes. All parental genes were then

analysed for the presence (1) or absence (0) of a hotspot. Log-odds ratios, standard errors and

confidence intervals are reported together with their p-values in **Supplementary Table 8**.

For the correlation of hotspot presence across the number of species, a generalised linear

model was applied using the categorical predictors "lineage" (= circRNA loci shared within rodents

or primates), "eutherian" (= circRNA loci shared within rodents and primates) and "therian" (= cir-

cRNA loci shared within opossum, rodents and primates). Log-odds ratios, standard errors and

confidence intervals were standardised using the *beta()* function from the *reghelper* R package (ver-

sion 1.0.0) and are reported together with their p-values in **Supplementary Table 7**.

Comparison to human and mouse circRNA heart dataset

The circRNA annotations for human and mouse heart as provided by *Werfel et al.* (*2016*) were,

based on the parental gene ID, merged with our circRNA annotations. Prediction values for parental

genes were calculated using the same general linear regression models as described above (sec-

tion *Generalised linear models* in **Material and Methods**) with genomic length, number of exons,

GC content, expression levels, reverse complements (RVCs) and phastCons scores as predictors.

Prediction values were received from the model and compared between parental genes predicted

by our and the Werfel dataset as well as between the predictors in non-parental and parental genes

of the Werfel dataset (**Figure 3-Figure supplement 4**).

Integration of external studies

(1) Replication time

Values for the replication time were used as provided in *Koren et al.* (*2012*). Coordinates of the dif-

ferent replication domains were intersected with the coordinates of coding genes using BEDtools

(*bedtools merge -f 1*). The mean replication time of each gene was used for subsequent analyses.

(2) Gene expression steady-state levels

Gene expression steady-state levels and decay rates were used as provided in Table S1 of *Pai et al.*

(*2012*).

(3) GHIS

Genome-wide haploinsufficiency scores for each gene were used as provided in Supplementary

Table S2 of *Steinberg et al.* (*2015*).

822 **Repeat analyses**

823 Generation of length- and GC-matched background dataset

824 Flanking introns were grouped into a matrix of i columns and j rows representing different genomic

825 lengths and GC content; i and j were calculated in the following way:

$$i = seq(from = quantile(GCcontent, 0.05), to = quantile(GCcontent, 0.95), by = 0.01)$$

$$j = seq(from = quantile(length, 0.05), to = quantile(length, 0.95), by = 1000)$$

826 Flanking introns were sorted into the matrix based on their GC content and length. A second matrix

827 with the same properties was created containing all introns of coding genes. From the latter, a

828 submatrix was sampled with the same length and GC distribution as the matrix for flanking introns.

829 The length distribution and GC distribution of the sampled introns reflect the distributions for the

830 flanking introns as assessed by a Fisher's t Test that was non-significant.

831 Repeat definition

832 The RepeatMasker annotation for full and nested repeats were downloaded for all genomes using

833 the UCSC Table browser (tracks "RepeatMasker" and "Interrupted Rpts") and the two files merged.

834 Nested repeats were included, because it was shown that small repetitive regions are sufficient to

835 trigger base pairing necessary for backsplicing (*Liang and Wilusz, 2014*; *Kramer et al., 2015*). For

836 rhesus macaque, the repeat annotation was only available for the rheMac3 genome. RVC coordi-

837 nates were thus lifted from rheMac2 to rheMac3 (*liftOver, -minMatch=0.5*), which led to a significant

838 drop of overlapping repeats and RVCs in comparison to the other species (only ~20% of RVCs could

839 be intersected with an annotated repeat). The complete list of full and nested repeats was then

840 intersected (*bedtools merge -f1*) with the above defined list of background and flanking introns for

841 further analyses.

842 Identification of repeat dimers

843 The complementary regions (RVCs) that were defined with megaBLAST as described above, were

844 intersected with the coordinates of individual repeats from the RepeatMasker annotation. To be

845 counted, a repeat had to overlap with at least 50% of its length with the region of complementarity

846 (*bedtools merge -f 0.5*). As RVCs can contain several repeats, the "strongest" dimer was selected

847 based on the number of overlapping base pairs (= longest overlapping dimer).

848 We observed that the same genomic repeat can often be present in multiple RVCs. Assuming

849 that repeats are unlikely to form multiple active dimers in the genome at the same given time point,

850 we decided to correct dimer frequency for this "co-counting" to not inflate our numbers and bias

851 subsequent analyses (see also **Figure 5-Figure supplement 2**). We calculated an overestimation

852 factor based on the number of possible interactions each repeat had. Dimer frequency was then

853 calculated as;

$$overestimation\_factor = \frac{co-counts_{\text{Repeat 1}} + co-counts_{\text{Repeat 2}}}{2}$$

$$dimer\_count_{\text{correct}} = \frac{dimer\_count}{overestimation\_factor}$$

The "dimer list" obtained from this analysis for each species was further ranked according to the absolute frequency of each dimer. The proportion of the top-5 dimer frequency to all detected dimers, was calculated based on this list ($n_{\text{top-5}}$ / $n_{\text{all\_dimers}}$).

Pairing scores of repeat dimers

Pairing scores for each TE class (based on the TE reference sequence) were defined by taking into account the (1) phylogenetic distance to other repeat families in the same species and (2) its binding affinity (the Minimal Free Energy = MFE of the dimer structure) to those repeats. We decided to not include the absolute TE frequency into the pairing score, because it is a function of the TE's age, its amplification and degradation rates. Simulating the interplay between these three components is not in scope of this study, and the integration of the frequency into the pairing score creates more noise as tested via PCA analyses (variance explained drops by 10%).

(1) Phylogenetic distance

TE reference sequences were obtained from Repbase (*Bao et al., 2015*) and translated into fasta-format for alignment (*reference_sequences.fa*). Alignments were then generated with Clustal Omega (v1.2.4) (*Sievers et al., 2011*) using the following settings:

$ clustalo -i reference_sequences.fa –distmat-out = repeats.mat –guidetree-out = repeats.dnd –full

The resulting distance matrix for the alignment was used for the calculation of the pairing score. Visualisation of the distance matrix (**Figure 4C**, **Figure 4-Figure supplement 2**) was performed using the standard R functions *dist(method="euclidian")* and *hclust(method="ward.D2")*. Since several TE classes evolved independently from each other, the plot was manually modified to remove connections or to add additional information on the TE's origin from literature.

(2) Binding affinity

To estimate the binding affinity of individual TE dimers, the free energy of the secondary structure of the respective TE dimers was calculated with the RNAcofold function from the ViennaRNA Package:

$ RNAcofold -a -d2 < dimerSequence.fa

with *dimerSequence.fa* containing the two TE reference sequences from which the dimer is composed. The resulting MFE values were used to calculate the pairing score.

889 (3) Final pairing score

890 To generate the final pairing score, values from the distance matrix and the binding affinity were

891 standardised (separately from each other) to values between 0 and 1:

$$f(x) = \frac{x - min(v)}{max(v) - min(v)}$$

892 with *x* being the pairing affinity/dimer frequency and *minv* and *maxv* the minimal and maximal

893 observed value in the distribution. The standardised values for the binding affinity and dimer fre-

894 quency were then summed up (= pairing score) and classified by PCA using the R environment:

895

896 *$ pca <- prcomp(score, center=TRUE, scale.=FALSE)*

897

898 PC1 and PC2 were used for subsequent plotting with the absolute frequency of dimers represented

899 by the size of the data points (**Figure 4D-F**, **Figure 4-Figure supplement 2**).

900 Dimer composition in shared and species-specific circRNA loci

901 Dimers were sorted by their frequency in all parental genes and the 100 most and least frequent

902 dimers were selected to be analysed for their enrichment in shared vs. species-specific circRNA loci.

903 The two dimer frequency distributions were compared using a Wilcoxon Signed Rank Test. Dimer

904 age was defined on whether the repeat family originated in a given species (= rank 1), lineage (=

905 rank 2), in all eutherian species of this study (rank 3) or all therian species (rank 4). Since a dimer is

906 composed of two repeats, the 'mean dimer age' based on the rank value was taken. Based on this

907 analysis, the top-5 most frequent and enriched dimers were then defined.

908 Calculation of TE degradation levels

909 We analysed repeat degradation levels for all TEs present in the top-5 dimers of each species. Re-

910 peatMasker annotations were downloaded from the UCSC Table browser for all genomes (see

911 **Material and Methods**, section *Repeat definition*). The milliDiv values for each TE were retrieved

912 from this annotation for full and nested repeats. All indivudal TEs where then grouped as "species-

913 specific" or "shared" based on whether the circRNA parental gene produced species-specific or

914 shared circRNA loci. Significance levels for milliDiv differences between the TE groups were as-

915 sessed with a simple Mann-Whitney U test.

916 Binding affinity of dimers

917 The binding affinity of dimers was calculated with the RNAcofold function from the ViennaRNA

918 Package:

919

920 *$ RNAcofold -a -d2 < dimerSequence.fa*

921

with *dimerSequence.fa* containing the two TE genomic sequences from which the dimer is com-
posed. To reduce calculation time for human and opossum, the analysis was restricted to the re-
spective top-5 dimers (see section *Dimer composition in shared vs. species-specific circRNA loci*). For
each gene of the two groups (shared/species-specific), the least degraded dimer based on its mean
milliDiv value was chosen. Filtering based on the least degraded dimer, let to a strong enrichment
of only a subset of the top-5 dimers in each species. If enough observations for a statistical test
were present, the two distributions (shared/species-specific) where compared using a Student's
t-Test.

Supplementary Data

Supplementary Tables and Figures

932 Supplementary Tables and Figures are available as an attachment to this document.

Supplementary Files

Supplementary File 1: CircRNA annotation file for opossum. A gtf-file with all circRNA transcripts
including the transcript and exon coordinates.

Supplementary File 2: CircRNA annotation file for mouse. A gtf-file with all circRNA transcripts
including the transcript and exon coordinates.

Supplementary File 3: CircRNA annotation file for rat. A gtf-file with all circRNA transcripts includ-
ing the transcript and exon coordinates.

Supplementary File 4: CircRNA annotation file for rhesus macaque. A gtf-file with all circRNA tran-
scripts including the transcript and exon coordinates.

Supplementary File 5: CircRNA annotation file for human. A gtf-file with all circRNA transcripts
including the transcript and exon coordinates.

All gtf-files have been uploaded to the UCSC genome browser and can be viewed here:

Opossum: http://genome.ucsc.edu/s/Frenzchen/monDom5%20circRNA%20annotation

Mouse; http://genome.ucsc.edu/s/Frenzchen/mm10%20circRNA%20annotation

Rat: http://genome.ucsc.edu/s/Frenzchen/rn5%20circRNA%20annotation

Rhesus macaque: http://genome.ucsc.edu/s/Frenzchen/rheMac2%20circRNA%20annotation

Human: http://genome.ucsc.edu/s/Frenzchen/hg38%20circRNA%20annotation

Ethics statement

The human post-mortem samples were provided by the NICHD Brain and Tissue Bank for Develop-
mental Disorders at the University of Maryland (USA). They originated from individuals with diverse
causes of death that, given the information available, were not associated with the organ sampled.
Written consent for the use of human tissues for research was obtained from all donors or their
next of kin by this tissue bank. The use of these samples was approved by an ERC Ethics Screening
panel (associated with H.K.'s ERC Consolidator Grant 615253, OntoTransEvol), and, in addition, by
the local ethics committee in Lausanne (authorization 504/12). The rhesus macaque samples were

provided by the Suzhou Experimental Animal Center (China); the Biomedical Research Ethics Com-

mittee of Shanghai Institutes for Biological Sciences reviewed the use and care of the animals in the

research project (approval ID: ER-SIBS-260802P). All rhesus macaques used in this study suffered

sudden deaths for reasons other than their participation in this study and without any relation to

the organ sampled. Mouse samples were collected by the Kaessmann lab at the Center for Integra-

tive Genomics in Lausanne. Rat samples were kindly provided by Carmen Sandi, EPFL, Lausanne.

Opossum samples were kindly provided by Peter Giere, Museum für Naturkunde, Berlin. All ani-

mal procedures were performed in compliance with national and international ethical guidelines

and regulations for the care and use of laboratory animals and were approved by the local ani-

mal welfare authorities (Vaud Cantonal Veterinary office, Berlin State Office of Health and Social

Affairs). The use of all animal samples was approved by an ERC Ethics Screening panel (associated

with H.K.'s ERC Consolidator Grant 615253, OntoTransEvol).

Acknowledgments

We thank the Lausanne Genomics Technologies Facility for high throughput sequencing support;

Jean Halbert, Delphine Valloton and Angelica Liechti for opossum, mouse and rat tissue dissec-

tion and RNA extractions; Philipp Khaitovich for providing human and rhesus macaque samples;

Bulak Arpat and Thomas O. Auer for discussions on the manuscript; and Ioannis Xenarios for dis-

cussion and support with IT-infrastructure and data archiving. This research was supported by

grants from the Swiss National Science Foundation to D.G. (NCCR RNA & Disease and individual

grant 179190) and from the European Research Council to H.K. (242597, SexGenTransEvolution;

and 615253, OntoTransEvol). F.G. was supported by the SIB PhD Fellowship granted by the Swiss

Institute of Bioinformatics and the Fondation Leenaards. P.J. was supported by Human Frontiers

Science Program long-term fellowship LT000158/2013-L.

Author contributions

983 Contributions to this publication are distributed as follows: Study design: F.G., D.G., H.K. and P.J.;

984 Experimental work: F.G. and P.J.; Bioinformatics data analyses: F.G.; Paper manuscript and discus-

985 sion: F.G., D.G. and H.K.

Competing interests

987 No competing interests.

References

Alhasan AA, Izuogu OG, Al-Balool HH, Steyn JS, Evans A, Colzani M, Ghevaert C, Mountford JC, Marenah L, Elliott DJ, Santibanez-Koref M, Jackson MS. Circular RNA enrichment in platelets is a signature of transcriptome degradation. Blood. 2015 dec; http://eutils.ncbi.nlm.nih.gov/entrez/eutils/elink.fcgi?dbfrom=pubmed&id=26660425&retmode=ref&cmd=prlinks, doi: 10.1182/blood-2015-06-649434.

Amit M, Donyo M, Hollander D, Goren A, Kim E, Gelfman S, Lev-Maor G, Burstein D, Schwartz S, Postolsky B, Pupko T, Ast G. Differential GC content between exons and introns establishes distinct strategies of splice-site recognition. Cell reports. 2012 may; 1(5):543–556. http://dx.doi.org/10.1016/j.celrep.2012.03.013, doi: 10.1016/j.celrep.2012.03.013.

Ashwal-Fluss R, Meyer M, Pamudurti NR, Ivanov A. circRNA Biogenesis Competes with Pre-mRNA Splicing. Molecular cell. 2014; .

Athanasiadis A, Rich A, Maas S. Widespread A-to-I RNA editing of Alu-containing mRNAs in the human transcriptome. PLoS Biology. 2004 dec; 2(12):e391. http://dx.doi.org/10.1371/journal.pbio.0020391, doi: 10.1371/journal.pbio.0020391.

Bachmayr-Heyda A, Reiner AT, Auer K, Sukhbaatar N, Aust S, Bachleitner-Hofmann T, Mesteri I, Grunt TW, Zeillinger R, Pils D. Correlation of circular RNA abundance with proliferation–exemplified with colorectal and ovarian cancer, idiopathic lung fibrosis, and normal human tissues. Scientific reports. 2015; .

Bahn JH, Zhang Q, Li F, Chan TM, Lin X, Kim Y, Wong DTW, Xiao X. The landscape of microRNA, Piwi-interacting RNA, and circular RNA in human saliva. Clinical Chemistry. 2015 jan; 61(1):221–230. http://dx.doi.org/10.1373/clinchem.2014.230433, doi: 10.1373/clinchem.2014.230433.

Bao W, Kojima KK, Kohany O. Repbase Update, a database of repetitive elements in eukaryotic genomes. Mobile DNA. 2015 jun; 6:11. http://dx.doi.org/10.1186/s13100-015-0041-9, doi: 10.1186/s13100-015-0041-9.

Batzer MA, Deininger PL. Alu repeats and human genomic diversity. Nature Reviews Genetics. 2002 may; 3(5):370–379. http://dx.doi.org/10.1038/nrg798, doi: 10.1038/nrg798.

Brawand D, Soumillon M, Necsulea A, Julien P, Csárdi G, Harrigan P, Weier M, Liechti A, Aximu-Petri A, Kircher M, Albert FW, Zeller U, Khaitovich P, Grützner F, Bergmann S, Nielsen R, Pääbo S, Kaessmann H. The evolution of gene expression levels in mammalian organs. Nature. 2011 oct; 478(7369):343–348. http://dx.doi.org/10.1038/nature10532, doi: 10.1038/nature10532.

Conn SJ, Pillman KA, Toubia J, Conn VM, Salmanidis M, Phillips CA, Roslan S, Schreiber AW, Gregory PA, Goodall GJ. The RNA binding protein quaking regulates formation of circRNAs. Cell. 2015 mar; 160(6):1125–1134. http://dx.doi.org/10.1016/j.cell.2015.02.014, doi: 10.1016/j.cell.2015.02.014.

Cortés-López M, Gruner MR, Cooper DA, Gruner HN, Voda AI, van der Linden AM, Miura P. Global accumulation of circRNAs during aging in Caenorhabditis elegans. BMC Genomics. 2018 jan; 19(1):8. http://dx.doi.org/10.1186/s12864-017-4386-y, doi: 10.1186/s12864-017-4386-y.

Deininger P. Alu elements: know the SINEs. Genome biology. 2011 Dec; 12(12):236. http://eutils.ncbi.nlm.nih.gov/entrez/eutils/elink.fcgi?dbfrom=pubmed&id=22204421&retmode=ref&cmd=prlinks, doi: 10.1186/gb-2011-12-12-236.

Di Timoteo G, Dattilo D, Centrón-Broco A, Colantoni A, Guarnacci M, Rossi F, Incarnato D, Oliviero S, Fatica A, Morlando M, Bozzoni I. Modulation of circRNA Metabolism by m(6)A Modification. Cell reports. 2020 May; 31(6):107641. http://eutils.ncbi.nlm.nih.gov/entrez/eutils/elink.fcgi?dbfrom=pubmed&id=32402287&retmode=ref&cmd=prlinks, doi: 10.1016/j.celrep.2020.107641.

Dongen S. Performance criteria for graph clustering and Markov cluster experiments. . 2000 May; http://dl.acm.org/citation.cfm?id=868979.

Du WW, Yang W, Liu E, Yang Z, Dhaliwal P, Yang BB. Foxo3 circular RNA retards cell cycle progression via forming ternary complexes with p21 and CDK2. Nucleic Acids Research. 2016 apr; 44(6):2846–2858. http://dx.doi.org/10.1093/nar/gkw027, doi: 10.1093/nar/gkw027.

Dubin RA, Kazmi MA, Ostrer H. Inverted repeats are necessary for circularization of the mouse testis Sry transcript. Gene. 1995 dec; 167(1-2):245–248. https://www.ncbi.nlm.nih.gov/pubmed/8566785.

Edgar R, Domrachev M, Lash AE. Gene Expression Omnibus: NCBI gene expression and hybridization array data repository. Nucleic Acids Research. 2002 jan; 30(1):207–210. http://dx.doi.org/10.1093/nar/30.1.207, doi: 10.1093/nar/30.1.207.

Enright AJ, Van Dongen S, Ouzounis CA. An efficient algorithm for large-scale detection of protein families. Nucleic Acids Research. 2002 apr; 30(7):1575–1584. http://dx.doi.org/10.1093/nar/30.7.1575, doi: 10.1093/nar/30.7.1575.

Enuka Y, Lauriola M, Feldman ME, Sas-Chen A, Ulitsky I, Yarden Y. Circular RNAs are long-lived and display only minimal early alterations in response to a growth factor. Nucleic Acids Research. 2016 feb; 44(3):1370–1383. http://dx.doi.org/10.1093/nar/gkv1367, doi: 10.1093/nar/gkv1367.

Ermakova EO, Nurtdinov RN, Gelfand MS. Fast rate of evolution in alternatively spliced coding regions of mammalian genes. BMC Genomics. 2006 apr; 7:84. http://dx.doi.org/10.1186/1471-2164-7-84, doi: 10.1186/1471-2164-7-84.

Galtier N, Mouchiroud D. Isochore evolution in mammals: a human-like ancestral structure. Genetics. 1998 dec; 150(4):1577–1584. https://www.ncbi.nlm.nih.gov/pubmed/9832533.

Gruner H, Cortés-López M, Cooper DA, Bauer M, Miura P. CircRNA accumulation in the aging mouse brain. Scientific Reports. 2016 dec; 6:38907. http://dx.doi.org/10.1038/srep38907, doi: 10.1038/srep38907.

Gu W, Ray DA, Walker JA, Barnes EW, Gentles AJ, Samollow PB, Jurka J, Batzer MA, Pollock DD. SINEs, evolution and genome structure in the opossum. Gene. 2007 jul; 396(1):46–58. http://dx.doi.org/10.1016/j.gene.2007.02.028, doi: 10.1016/j.gene.2007.02.028.

Guo JU, Agarwal V, Guo H, Bartel DP. Expanded identification and characterization of mammalian circular RNAs. Genome Biology. 2014 jul; 15(7):409. http://dx.doi.org/10.1186/s13059-014-0409-z, doi: 10.1186/s13059-014-0409-z.

Hansen TB, Jensen TI, Clausen BH, Bramsen JB, Finsen B, Damgaard CK, Kjems J. Natural RNA circles function as efficient microRNA sponges. Nature. 2013 mar; 495(7441):384–388. http://dx.doi.org/10.1038/nature11993, doi: 10.1038/nature11993.

Heber S, Alekseyev M, Sze SH, Tang H, Pevzner PA. Splicing graphs and EST assembly problem. Bioinformatics (Oxford, England). 2002; 18 Suppl 1:S181–8. http://eutils.ncbi.nlm.nih.gov/entrez/eutils/elink.fcgi?dbfrom=pubmed&id=12169546&retmode=ref&cmd=prlinks, doi: 10.1093/bioinformatics/18.suppl_1.s181.

Ivanov A, Memczak S, Wyler E, Torti F, Porath HT, Orejuela MR, Piechotta M, Levanon EY, Landthaler M, Dieterich C, Rajewsky N. Analysis of intron sequences reveals hallmarks of circular RNA biogenesis in animals. Cell reports. 2015 jan; 10(2):170–177. http://dx.doi.org/10.1016/j.celrep.2014.12.019, doi: 10.1016/j.celrep.2014.12.019.

Jeck WR, Sorrentino JA, Wang K, Slevin MK, Burd CE, Liu J, Marzluff WF, Sharpless NE. Circular RNAs are abundant, conserved, and associated with ALU repeats. RNA (New York). 2013 feb; 19(2):141–157. http://dx.doi.org/10.1261/rna.035667.112, doi: 10.1261/rna.035667.112.

Kim J, Deininger PL. Recent amplification of rat ID sequences. Journal of Molecular Biology. 1996 aug; 261(3):322–327. http://dx.doi.org/10.1006/jmbi.1996.0464, doi: 10.1006/jmbi.1996.0464.

Kim J, Martignetti JA, Shen MR, Brosius J, Deininger P. Rodent BC1 RNA gene as a master gene for ID element amplification. Proceedings of the National Academy of Sciences of the United States of America. 1994 apr; 91(9):3607–3611. http://dx.doi.org/10.1073/pnas.91.9.3607, doi: 10.1073/pnas.91.9.3607.

Koren A, Polak P, Nemesh J, Michaelson JJ, Sebat J, Sunyaev SR, McCarroll SA. Differential relationship of DNA replication timing to different forms of human mutation and variation. American Journal of Human Genetics. 2012 dec; 91(6):1033–1040. http://dx.doi.org/10.1016/j.ajhg.2012.10.018, doi: 10.1016/j.ajhg.2012.10.018.

Kramer MC, Liang D, Tatomer DC, Gold B, March ZM, Cherry S, Wilusz JE. Combinatorial control of Drosophila circular RNA expression by intronic repeats, hnRNPs, and SR proteins. Genes & Development. 2015 oct; 29(20):2168–2182. http://dx.doi.org/10.1101/gad.270421.115, doi: 10.1101/gad.270421.115.

Kristensen LS, Andersen MS, Stagsted LVW, Ebbesen KK, Hansen TB, Kjems J. The biogenesis, biology and characterization of circular RNAs. Nature Reviews Genetics. 2019 aug; 20(11):675–691. http://dx.doi.org/10.1038/s41576-019-0158-7, doi: 10.1038/s41576-019-0158-7.

Lee Y, Choe J, Park OH, Kim YK. Molecular Mechanisms Driving mRNA Degradation by m(6)A Modification. Trends in genetics : TIG. 2020 Mar; 36(3):177–188. http://eutils.ncbi.nlm.nih.gov/entrez/eutils/elink.fcgi?dbfrom=pubmed&id=31964509&retmode=ref&cmd=prlinks, doi: 10.1016/j.tig.2019.12.007.

Lev-Maor G, Ram O, Kim E, Sela N, Goren A, Levanon EY, Ast G. Intronic Alus influence alternative splicing. PLoS Genetics. 2008 sep; 4(9):e1000204. http://dx.doi.org/10.1371/journal.pgen.1000204, doi: 10.1371/journal.pgen.1000204.

Li S, Li X, Xue W, Zhang L, Yang LZ, Cao SM, Lei YN, Liu CX, Guo SK, Shan L, Wu M, Tao X, Zhang JL, Gao X, Zhang J, Wei J, Li J, Yang L, Chen LL. Screening for functional circular RNAs using the CRISPR–Cas13 system. Nature Methods. 2020; https://doi.org/10.1038/s41592-020-01011-4.

Liang D, Wilusz JE. Short intronic repeat sequences facilitate circular RNA production. Genes & Development. 2014 oct; 28(20):2233–2247. http://dx.doi.org/10.1101/gad.251926.114, doi: 10.1101/gad.251926.114.

Memczak S, Jens M, Elefsinioti A, Torti F, Krueger J, Rybak A, Maier L, Mackowiak SD, Gregersen LH, Munschauer M, Loewer A, Ziebold U, Landthaler M, Kocks C, le Noble F, Rajewsky N. Circular RNAs are a large class of animal RNAs with regulatory potency. Nature. 2013 mar; 495(7441):333–338. http://dx.doi.org/10.1038/nature11928, doi: 10.1038/nature11928.

Memczak S, Papavasileiou P, Peters O, Rajewsky N. Identification and characterization of circular rnas as a new class of putative biomarkers in human blood. Plos One. 2015 oct; 10(10):e0141214. http://dx.doi.org/10.1371/journal.pone.0141214, doi: 10.1371/journal.pone.0141214.

Mikkelsen TS, Wakefield MJ, Aken B, Amemiya CT, Chang JL, Duke S, Garber M, Gentles AJ, Goodstadt L, Heger A, Jurka J, Kamal M, Mauceli E, Searle SMJ, Sharpe T, Baker ML, Batzer MA, Benos PV, Belov K, Clamp M, et al. Genome of the marsupial Monodelphis domestica reveals innovation in non-coding sequences. Nature. 2007 may; 447(7141):167–177. http://dx.doi.org/10.1038/nature05805, doi: 10.1038/nature05805.

Modrek B, Lee CJ. Alternative splicing in the human, mouse and rat genomes is associated with an increased frequency of exon creation and/or loss. Nature Genetics. 2003 jun; 34(2):177–180. http://dx.doi.org/10.1038/ng1159, doi: 10.1038/ng1159.

Okholm TLH, Sathe S, Park SS, Kamstrup AB, Rasmussen AM, Shankar A, Chua ZM, Fristrup N, Nielsen MM, Vang S, Dyrskjøt L, Aigner S, Damgaard CK, Yeo GW, Pedersen JS. Transcriptome-wide profiles of circular RNA and RNA-binding protein interactions reveal effects on circular RNA biogenesis and cancer pathway expression. Genome Medicine. 2020; 12(1):112. https://doi.org/10.1186/s13073-020-00812-8.

Pai AA, Cain CE, Mizrahi-Man O, De Leon S, Lewellen N, Veyrieras JB, Degner JF, Gaffney DJ, Pickrell JK, Stephens M, Pritchard JK, Gilad Y. The contribution of RNA decay quantitative trait loci to inter-individual variation in steady-state gene expression levels. PLoS Genetics. 2012 oct; 8(10):e1003000. http://dx.doi.org/10.1371/journal.pgen.1003000, doi: 10.1371/journal.pgen.1003000.

Park OH, Ha H, Lee Y, Boo SH, Kwon DH, Song HK, Kim YK. Endoribonucleolytic Cleavage of m(6)A-Containing RNAs by RNase P/MRP Complex. Molecular cell. 2019 May; 74(3):494–507.e8. http://eutils.ncbi.nlm.nih.gov/entrez/eutils/elink.fcgi?dbfrom=pubmed&id=30930054&retmode=ref&cmd=prlinks, doi: 10.1016/j.molcel.2019.02.034.

Patop IL, Wüst S, Kadener S. Past, present, and future of circRNAs. The EMBO Journal. 2019; 38(16):e100836. https://www.embopress.org/doi/abs/10.15252/embj.2018100836, doi: https://doi.org/10.15252/embj.2018100836.

Piwecka M, Glažar P, Hernandez-Miranda LR, Memczak S, Wolf SA, Rybak-Wolf A, Filipchyk A, Klironomos F, Cerda Jara CA, Fenske P, Trimbuch T, Zywitza V, Plass M, Schreyer L, Ayoub S, Kocks C, Kühn R, Rosenmund C, Birchmeier C, Rajewsky N. Loss of a mammalian circular RNA locus causes miRNA deregulation and affects brain function. Science. 2017 sep; 357(6357). http://dx.doi.org/10.1126/science.aam8526, doi: 10.1126/science.aam8526.

Pollard KS, Hubisz MJ, Rosenbloom KR, Siepel A. Detection of nonneutral substitution rates on mammalian phylogenies. Genome Research. 2010 jan; 20(1):110–121. http://dx.doi.org/10.1101/gr.097857.109, doi: 10.1101/gr.097857.109.

Rybak-Wolf A, Stottmeister C, Glažar P, Jens M, Pino N, Giusti S, Hanan M, Behm M, Bartok O, Ashwal-Fluss R, Herzog M, Schreyer L, Papavasileiou P, Ivanov A, Öhman M, Refojo D, Kadener S, Rajewsky N. Circular RNAs in the mammalian brain are highly abundant, conserved, and dynamically expressed. Molecular Cell. 2015 jun; 58(5):870–885. http://dx.doi.org/10.1016/j.molcel.2015.03.027, doi: 10.1016/j.molcel.2015.03.027.

Santos-Rodriguez G, Voineagu I, Weatheritt RJ. Evolutionary dynamics of circular RNAs in primates. bioRxiv. 2021; https://www.biorxiv.org/content/early/2021/05/01/2021.05.01.442284, doi: 10.1101/2021.05.01.442284.

Shao T, Pan Yh, Xiong Xd. Circular RNA: an important player with multiple facets to regulate its parental gene expression. Molecular Therapy - Nucleic Acids. 2021 Mar; 23:369–376.

Siepel A, Bejerano G, Pedersen JS, Hinrichs AS, Hou M, Rosenbloom K, Clawson H, Spieth J, Hillier LW, Richards S, Weinstock GM, Wilson RK, Gibbs RA, Kent WJ, Miller W, Haussler D. Evolutionarily conserved elements in vertebrate, insect, worm, and yeast genomes. Genome Research. 2005 aug; 15(8):1034–1050. http://dx.doi.org/10.1101/gr.3715005, doi: 10.1101/gr.3715005.

Sievers F, Wilm A, Dineen D, Gibson TJ, Karplus K, Li W, Lopez R, McWilliam H, Remmert M, Söding J, Thompson JD, Higgins DG. Fast, scalable generation of high-quality protein multiple sequence alignments using Clustal Omega. Molecular Systems Biology. 2011 oct; 7:539. http://dx.doi.org/10.1038/msb.2011.75, doi: 10.1038/msb.2011.75.

Smit A, Hubley R, Green P. RepeatMasker Open-4.0, 2013-2015. . 2013; http://www.repeatmasker.org.

Starke S, Jost I, Rossbach O, Schneider T, Schreiner S, Hung LH, Bindereif A. Exon circularization requires canonical splice signals. Cell reports. 2015; .

Steinberg J, Honti F, Meader S, Webber C. Haploinsufficiency predictions without study bias. Nucleic Acids Research. 2015 sep; 43(15):e101. http://dx.doi.org/10.1093/nar/gkv474, doi: 10.1093/nar/gkv474.

Trapnell C, Roberts A, Goff L, Pertea G, Kim D, Kelley DR, Pimentel H, Salzberg SL, Rinn JL, Pachter L. Differential gene and transcript expression analysis of RNA-seq experiments with TopHat and Cufflinks. Nature Protocols. 2012 mar; 7(3):562–578. http://dx.doi.org/10.1038/nprot.2012.016, doi: 10.1038/nprot.2012.016.

VenøMT, Hansen TB, VenøST, Clausen BH, Grebing M, Finsen B, Holm IE, Kjems J. Spatio-temporal regulation of circular RNA expression during porcine embryonic brain development. Genome Biology. 2015 nov; 16:245. http://dx.doi.org/10.1186/s13059-015-0801-3, doi: 10.1186/s13059-015-0801-3.

Wang M, Hou J, Müller-McNicoll M, Chen W, Schuman EM. Long and Repeat-Rich Intronic Sequences Favor Circular RNA Formation under Conditions of Reduced Spliceosome Activity. iScience. 2019 oct; 20:237–247. http://dx.doi.org/10.1016/j.isci.2019.08.058, doi: 10.1016/j.isci.2019.08.058.

Wang PL, Bao Y, Yee MC, Barrett SP, Hogan GJ, Olsen MN, Dinneny JR, Brown PO, Salzman J. Circular RNA is expressed across the eukaryotic tree of life. Plos One. 2014 mar; 9(6):e90859. http://dx.doi.org/10.1371/journal.pone.0090859, doi: 10.1371/journal.pone.0090859.

Werfel S, Nothjunge S, Schwarzmayr T, Strom TM, Meitinger T, Engelhardt S. Characterization of circular RNAs in human, mouse and rat hearts. Journal of Molecular and Cellular Cardiology. 2016 jul; 98:103–107. http://dx.doi.org/10.1016/j.yjmcc.2016.07.007, doi: 10.1016/j.yjmcc.2016.07.007.

Westholm JO, Miura P, Olson S, Shenker S, Joseph B, Sanfilippo P, Celniker SE, Graveley BR, Lai EC. Genome-wide analysis of drosophila circular RNAs reveals their structural and sequence properties and age-dependent neural accumulation. Cell reports. 2014 dec; 9(5):1966–1980. http://dx.doi.org/10.1016/j.celrep.2014.10.062, doi: 10.1016/j.celrep.2014.10.062.

Wilusz JE. Repetitive elements regulate circular RNA biogenesis. Mobile genetic elements. 2015 jun; 5(3):1–7. http://dx.doi.org/10.1080/{2159256X}.2015.1045682, doi: 10.1080/{2159256X}.2015.1045682.

Xu C, Zhang J. Mammalian circular RNAs result largely from splicing errors. Cell Reports. 2021; 36(4):109439. https://www.sciencedirect.com/science/article/pii/S2211124721008561, doi: https://doi.org/10.1016/j.celrep.2021.109439.

Xu K, Chen D, Wang Z, Ma J, Zhou J, Chen N, Lv L, Zheng Y, Hu X, Zhang Y, Li J. Annotation and functional clustering of circRNA expression in rhesus macaque brain during aging. Cell discovery. 2018 sep; 4:48. http://www.nature.com/articles/s41421-018-0050-1, doi: 10.1038/s41421-018-0050-1.

Yoder JA, Walsh CP, Bestor TH. Cytosine methylation and the ecology of intragenomic parasites. Trends in Genetics. 1997 aug; 13(8):335–340. http://dx.doi.org/10.1016/s0168-9525(97)01181-5, doi: 10.1016/s0168-9525(97)01181-5.

Yoshimoto R, Rahimi K, Hansen TB, Kjems J, Mayeda A. Biosynthesis of Circular RNA ciRS-7/CDR1as Is Mediated by Mammalian-wide Interspersed Repeats. iScience. 2020; 23(7):101345. http://www.sciencedirect.com/science/article/pii/S2589004220305320, doi: https://doi.org/10.1016/j.isci.2020.101345.

You X, Vlatkovic I, Babic A, Will T, Epstein I, Tushev G, Akbalik G, Wang M, Glock C, Quedenau C, Wang X, Hou J, Liu H, Sun W, Sambandan S, Chen T, Schuman EM, Chen W. Neural circular RNAs are derived from synaptic genes and regulated by development and plasticity. Nature Neuroscience. 2015 apr; 18(4):603–610. http://dx.doi.org/10.1038/nn.3975, doi: 10.1038/nn.3975.

Zaccara S, Ries RJ, Jaffrey SR. Reading, writing and erasing mRNA methylation. Nature reviews Molecular cell biology. 2019; 20(10):608–624. https://doi.org/10.1038/s41580-019-0168-5.

Zhang XO, Wang HB, Zhang Y, Lu X, Chen LL, Yang L. Complementary sequence-mediated exon circularization. Cell. 2014 sep; 159(1):134–147. http://dx.doi.org/10.1016/j.cell.2014.09.001, doi: 10.1016/j.cell.2014.09.001.

Zhang Y, Romanish MT, Mager DL. Distributions of transposable elements reveal hazardous zones in mammalian introns. PLoS Computational Biology. 2011 may; 7(5):e1002046. http://dx.doi.org/10.1371/journal.pcbi.1002046, doi: 10.1371/journal.pcbi.1002046.

Zheng Q, Bao C, Guo W, Li S, Chen J, Chen B, Luo Y, Lyu D, Li Y, Shi G, Liang L, Gu J, He X, Huang S. Circular RNA
profiling reveals an abundant circHIPK3 that regulates cell growth by sponging multiple miRNAs : Nature
Communications. Nature communications. 2016; 7:11215. http://eutils.ncbi.nlm.nih.gov/entrez/eutils/elink.
fcgi?dbfrom=pubmed&id=27050392&retmode=ref&cmd=prlinks, doi: 10.1038/ncomms11215.

Zhou C, Molinie B, Daneshvar K, Pondick JV, Wang J, Van Wittenberghe N, Xing Y, Giallourakis CC, Mullen AC.
Genome-Wide Maps of m6A circRNAs Identify Widespread and Cell-Type-Specific Methylation Patterns that
Are Distinct from mRNAs. Cell reports. 2017 aug; 20(9):2262–2276. http://dx.doi.org/10.1016/j.celrep.2017.
08.027, doi: 10.1016/j.celrep.2017.08.027.

Zhu L, Zhang Y, Zhang W, Yang S, Chen JQ, Tian D. Patterns of exon-intron architecture variation of genes
in eukaryotic genomes. BMC Genomics. 2009 jan; 10:47. http://dx.doi.org/10.1186/1471-2164-10-47, doi:
10.1186/1471-2164-10-47.

Figure 1–Figure supplement 1.

Figure 1–Figure supplement 2.

Figure 1–Figure supplement 3.

Figure 1–Figure supplement 4.

Figure 2–Figure supplement 1.

Figure 2–Figure supplement 2.

Figure 3–Figure supplement 1.

Figure 3–Figure supplement 2.

Figure 3–Figure supplement 3.

Figure 3–Figure supplement 4.

Figure 3–Figure supplement 5.

Figure 4–Figure supplement 1.

Figure 4–Figure supplement 2.

Figure 5–Figure supplement 1.

Figure 5–Figure supplement 2.

Figure 5–Figure supplement 3.

Supplementary Tables and Figures

Circular RNA repertoires are associated with evolutionarily young transposable elements

Franziska Gruhl [1,2], Peggy Janich [1,3], Henrik Kaessmann [4*], David Gatfield [1*]

[1] Center for Integrative Genomics, University of Lausanne, Lausanne, Switzerland
[2] SIB, Swiss Institute of Bioinformatics, Lausanne, Switzerland
[3] Krebsforschung Schweiz, CH-3001, Bern, Switzerland
[4] Center for Molecular Biology of Heidelberg University (ZMBH), DKFZ-ZMBH Alliance, Heidelberg, Germany

Supplementary Table 1: Sample overview.

Supplementary Table 1. Summary of organism, tissue, age and sex for each sample; last column shows the RNA Quality Number (RQN) for the extracted RNA.

Species	Tissue	Age	Sex	RQN
Opossum	Cerebellum	21 months	male	7.3
Opossum	Cerebellum	19.5 months	male	8.9
Opossum	Cerebellum	15.5 months	male	6.8
Opossum	Liver	15.5 months	male	9.3
Opossum	Liver	21 months	male	8.6
Opossum	Liver	13 months	male	9
Opossum	Testis	21 months	male	8.9
Opossum	Testis	13 months	male	8.5
Opossum	Testis	15.5 months	male	8.9
Mouse	Cerebellum	9 weeks	male	7.1
Mouse	Cerebellum	9 weeks	male	7.4
Mouse	Cerebellum	9 weeks	male	7
Mouse	Liver	9 weeks	male	7.9
Mouse	Liver	9 weeks	male	7.6
Mouse	Liver	9 weeks	male	8.5
Mouse	Testis	9 weeks	male	8.4
Mouse	Testis	9 weeks	male	8.2
Mouse	Testis	9 weeks	male	8.4
Rat	Cerebellum	16 weeks	male	7.2
Rat	Cerebellum	16 weeks	male	7.5
Rat	Cerebellum	16 weeks	male	7.7
Rat	Liver	16 weeks	male	7.2
Rat	Liver	16 weeks	male	7.9
Rat	Liver	16 weeks	male	7.8
Rat	Testis	16 weeks	male	7.7
Rat	Testis	16 weeks	male	8.8

Rat	Testis	16 weeks	male	7.8
Rhesus macaque	Cerebellum	8 years	male	8.5
Rhesus macaque	Cerebellum	9 years	male	7.7
Rhesus macaque	Liver	8 years	male	8.6
Rhesus macaque	Liver	9 years	male	8.2
Rhesus macaque	Liver	9 years	male	8.6
Rhesus macaque	Testis	8 years	male	9.5
Rhesus macaque	Testis	9 years	male	9.1
Rhesus macaque	Testis	8 years	male	8.8
Human	Liver	64 years	male	7.5
Human	Cerebellum	29 years	male	8.2
Human	Cerebellum	41 years	male	8.6
Human	Cerebellum	25 years	male	8.3
Human	Testis	21 years	male	7.8
Human	Testis	41 years	male	6.9
Human	Testis	22 years	male	6.9

Supplementary Table 2: Filtering steps and reduction of circRNAs candidates during the identification pipeline.

Supplementary Table 2. Description of the different filtering steps applied to generate a high confidence circRNA dataset based on the comparison of untreated and RNase R-treated samples. The number of unique BSJs left after each filtering step is shown for each tissue (see **Material and Methods,** section *Generation of high confidence circRNA candidates from the comparison of RNase R-treated vs. -untreated samples*); mouse was chosen as representative example.

	Liver	Cerebellum	Testis
After read mapping, the lists of BSJs in untreated and RNase R treated was merged for each biological replicate keeping all BSJs that were detected in either the untreated or the RNase R-treated sample. The total number of unique BSJs in each biological replicate is shown together with the number of unique BSJs in the untreated and RNase R-treated biological replicate.			
Biological replicate 1 (untreated | RNAse R)	24,474 (4,483 | 20,674)	55,455 (15,409 | 45,454)	47,794 (9,491 | 42,362)
Biological replicate 2 (untreated | RNAse R)	26,575 (4,788 | 22,602)	52,229 (13,724 | 48,322)	36,843 (9,427 | 30,590)
Biological replicate 3 (untreated | RNAse R)	23,699 (5,111 | 19,357)	68,154 (18,510 | 56,725)	40,907 (6,063 | 37,347)
Filtering step 1 When mapping paired-end sequencing data, both reads should ideally map to the genome (paired-end = "pe"). However, sometimes one of the mate reads cannot be mapped due to the complexity of the genomic locus. These reads are reported as "singletons" ("se"). We only kept BSJs for which both read mates mapped consistently either in "pe" or "se" mode (see **Material and Methods** for more details). The number of BSJs in each sample, which remain after filtering step 1, are indicated.			
Biological replicate 1 (% kept after filtering step 1)	24,373 (99.59%)	54,840 (98.89%)	47,416 (99.21%)
Biological replicate 2 (% kept after filtering step 1)	26,502 (99.73%)	51,725 (99.00%)	36,439 (98.90%)
Biological replicate 3 (% kept after filtering step 1)	23,568 (99.57%)	67,370 (98.85%)	40,544 (99.11%)
Total number of unique BSJs across all samples (untreated and RNase R-treated)	66,405	137,615	94,831
Filtering step 2 We assume that to have some kind of potential function, circRNAs need to be present in normal conditions. We thus removed all BSJs which were only present in RNase R treated samples and could not be detected in any of the untreated, biological replicates. The number of unique BSJs, which remain after filtering step 2, are indicated.			
Total number of unique BSJs across all samples (% kept from total, unique BSJs after filtering step 2)	13,084 (19.70%)	37,086 (26.95%)	20,358 (21.47%)

Filtering step 3		
Next, BSJs were normalized by the size factor of each sample (see **Material and Methods**) and the mean, normalised count was calculated for each condition (untreated and RNase R treated). Next, the log2-enrichment for RNase R-treated vs. -untreated samples was calculated. All BSJs for which the log2-enrichment was below 1.5 were removed. The number of BSJs in all untreated samples, which remain after filtering step 3, are indicated.		
Total number of unique BSJs across all samples (% kept from total, unique BSJs after filtering step 3)	1,914 (2.88%)	8,139 (5.91%)	6,381 (6.73%)
Filtering step 4		
The mean RPM value for each BSJ across untreated replicates was calculated. All BSJs with at least 0.05 were kept. These loci were considered strong circRNA candidates and used for all subsequent analyses. The final number of circRNAs, which remain after filtering step 4, are indicated.		
Total number of unique BSJs across all samples = final circRNA candidates (% kept from total, unique BSJs after filtering step 4)	87 (0.13%)	1,054 (0.77%)	523 (0.55%)

Supplementary Table 3: Detected back splice junctions (BSJs) across samples.

Supplementary Table 3. Table summarises the total number of detected BSJs after the filtering step in each species. The percentage of BSJs that are unique to one, two, three or more than three samples of the same species is shown.

Species	Total BSJs	1 replicate	2 replicates	3 replicates	>= 4 replicates
Opossum	76,739	84.74	8.05	4.28	2.93
Mouse	67,249	83.45	9.23	4.73	2.59
Rat	72,855	85.43	7.73	3.88	2.96
Rhesus macaque	100,270	79.29	9.79	4.83	6.09
Human	68,400	79.86	10.71	6.54	2.9

Supplementary Table 4: Total number of circRNAs in different species and tissues.

Supplementary Table 4. Indicated is the total number of different circRNAs that were annotated in each of the tissues across all species.

Species	Liver	Cerebellum	Testis
Opossum	129	417	1229
Mouse	87	1054	523
Rat	114	996	1192
Rhesus macaque	601	2132	1367
Human	765	2994	1761

Supplementary Table 5: Mean amplitude correlations.

Supplementary Table 5. Spearman's rank correlation for the GC amplitude and GC content of introns and exons are calculated for each isochore and species. The mean correlation between the GC amplitude and GC content of introns and exons is shown for different splice sites relative to the circRNA.

Position	Amplitude ~ Intron	Amplitude ~ Exon
Non-parental	-0.42	0.31
Outside of circRNA	-0.44	0.16
Inside of circRNA	-0.48	0.40

Supplementary Table 6: GLM summary for presence of parental genes.

Supplementary Table 6. A generalised linear model was fitted to predict the probability of coding genes to be a parental gene ($n_{opossum}$ = 18,807, n_{mouse} = 22,015, n_{rat} = 11,654, n_{rhesus} = 21,891, n_{human} = 21,744). The model was trained on 80% of the data (scaled values, cross-validation, 1000 repetitions, shown in rows labeled as "prediction"). Only the best predictors were kept and then used to predict probabilities for the remaining 20% of data points (validation set, shown in rows labeled as "validation"). Log-odds ratios, standard error and 95% confidence intervals (CI) for the validation set have been (beta) standardised.

Predictor	Coefficient	Std. error	Lower CI	Upper CI	p-value	Species	Dataset
as.rvc	0.4282	0.0318	0.3658	0.4906	2.93E-41	opossum	prediction
exon_count	0.3267	0.0309	0.2661	0.3872	3.51E-26	opossum	prediction
mean_brawand	0.3314	0.0484	0.2367	0.4263	7.28E-12	opossum	prediction
percentage_gc_content	-1.9481	0.1133	-2.1751	-1.7307	3.24E-66	opossum	prediction
as.rvc	0.2571	0.0307	0.1963	0.3168	5.54E-17	mouse	prediction
exon_count	0.3831	0.0318	0.3206	0.4454	2.14E-33	mouse	prediction
percentage_gc_content	-0.8193	0.058	-0.9341	-0.7068	2.44E-45	mouse	prediction
phastcons	0.5777	0.0607	0.4613	0.6993	1.71E-21	mouse	prediction
exon_count	0.2199	0.0357	0.1495	0.2895	6.91E-10	rat	prediction
genomic_length	0.2624	0.0325	0.1985	0.3263	7.36E-16	rat	prediction
mean_cpm	0.2696	0.0489	0.174	0.3658	3.58E-08	rat	prediction
percentage_gc_content	-0.5576	0.0601	-0.6763	-0.4408	1.68E-20	rat	prediction
phastcons	0.6314	0.0797	0.4802	0.793	2.35E-15	rat	prediction
ss.rvc	0.158	0.0416	0.0737	0.2373	0.000148111	rat	prediction
as.rvc	0.5653	0.0333	0.5001	0.6306	1.23E-64	rhesus	prediction
exon_count	0.3766	0.029	0.3197	0.4335	1.84E-38	rhesus	prediction
genomic_length	0.2506	0.026	0.2001	0.3022	6.36E-22	rhesus	prediction
mean_brawand	0.3162	0.0366	0.2446	0.3879	5.12E-18	rhesus	prediction
percentage_gc_content	-1.3246	0.0586	-1.4412	-1.2114	4.06E-113	rhesus	prediction
exon_count	0.3848	0.0291	0.3279	0.4419	5.10E-40	human	prediction

genomic_length	0.1772	0.0254	0.1279	0.2274	2.87E-12	human	prediction
mean_brawand	0.2675	0.0359	0.197	0.3378	9.71E-14	human	prediction
percentage_gc_content	-1.333	0.056	-1.4442	-1.2247	2.04E-125	human	prediction
phastcons	0.3218	0.0349	0.2538	0.3906	2.91E-20	human	prediction
ss.rvc	0.6142	0.0328	0.55	0.6787	3.25E-78	human	prediction
exon_count	0.4473	0.0646	0.3206	0.574	4.49E-12	opossum	validation
percentage_gc_content	-1.8437	0.2168	-2.2686	-1.4188	1.82E-17	opossum	validation
mean_brawand	0.343	0.0961	0.1547	0.5313	0.000357262	opossum	validation
as.rvc	0.284	0.0656	0.1554	0.4127	1.51E-05	opossum	validation
exon_count	0.3757	0.0682	0.242	0.5095	3.65E-08	mouse	validation
percentage_gc_content	-1.0861	0.1291	-1.3391	-0.8331	3.96E-17	mouse	validation
as.rvc	0.1967	0.063	0.0732	0.3202	0.001801116	mouse	validation
phastcons	0.5802	0.1226	0.3398	0.8205	2.24E-06	mouse	validation
genomic_length	0.2603	0.0727	0.1179	0.4027	0.000340157	rat	validation
exon_count	0.296	0.0732	0.1526	0.4395	5.24E-05	rat	validation
percentage_gc_content	-0.7197	0.1252	-0.9651	-0.4743	9.02E-09	rat	validation
mean_cpm	0.1467	0.0982	-0.0458	0.3392	0.135228403	rat	validation
ss.rvc	0.0848	0.0873	-0.0863	0.2559	0.33133768	rat	validation
phastcons	0.5127	0.1478	0.223	0.8024	0.00052204	rat	validation
genomic_length	0.1716	0.0491	0.0754	0.2678	0.000474304	rhesus	validation
exon_count	0.415	0.0595	0.2984	0.5315	3.02E-12	rhesus	validation
percentage_gc_content	-1.4385	0.121	-1.6757	-1.2013	1.39E-32	rhesus	validation
mean_brawand	0.3781	0.0722	0.2366	0.5197	1.64E-07	rhesus	validation
as.rvc	0.5888	0.0652	0.461	0.7165	1.67E-19	rhesus	validation
genomic_length	0.2624	0.0557	0.1533	0.3716	2.46E-06	human	validation

exon_count	0.3209	0.0613	0.2007	0.4411	1.67E-07	human	validation
percentage_gc_content	-1.4173	0.1224	-1.6572	-1.1774	5.37E-31	human	validation
mean_brawand	0.2475	0.0773	0.096	0.3989	0.001363255	human	validation
ss.rvc	0.5809	0.0692	0.4453	0.7166	4.76E-17	human	validation
phastcons	0.453	0.0763	0.3034	0.6025	2.89E-09	human	validation

Supplementary Table 7: GLM summary for "sharedness" of hotspots.

Supplementary Table 7. A generalised linear model was fitted to predict the probability of a hotspot to be present across multiple species ($n_{opossum}$ = 872, n_{mouse} = 848, n_{rat} = 665, n_{rhesus} = 1,682, n_{human} = 2,022). Reported log-odds ratios, standard error and 95% confidence intervals (CI) are (beta) standardised.

Predictor	Coefficient	Std. error	Lower CI	Upper CI	p-value	Species
therian	0.4283	0.0796	0.2723	0.5843	7.40E-08	opossum
rodents	0.2883	0.0909	0.11	0.4665	0.001525767	mouse
eutherian	0.6723	0.0981	0.4801	0.8646	7.10E-12	mouse
therian	0.7228	0.0882	0.5499	0.8956	2.49E-16	mouse
rodents	0.2048	0.0954	0.0178	0.3918	0.031813121	rat
eutherian	0.5835	0.0997	0.3881	0.779	4.87E-09	rat
therian	0.7539	0.0916	0.5744	0.9335	1.88E-16	rat
primates	0.4241	0.0617	0.3032	0.545	6.07E-12	rhesus
eutherian	0.5736	0.0577	0.4606	0.6867	2.59E-23	rhesus
therian	0.4952	0.0563	0.3848	0.6056	1.49E-18	rhesus
primates	0.4065	0.0506	0.3073	0.5056	9.12E-16	human
eutherian	0.4564	0.0492	0.36	0.5527	1.65E-20	human
therian	0.6161	0.051	0.5162	0.7161	1.35E-33	human

Supplementary Table 8: GLM summary for circRNA hotspots among parental genes.

Supplementary Table 8. A generalised linear model was fitted to predict the probability of circRNA hotspots among parental genes; parental genes were filtered for circRNAs that were either species-specific or occurred in orthologous loci across therian species ($n_{opossum}$ = 869, n_{mouse} = 503, n_{rat} = 425, n_{rhesus} = 912, n_{human} = 1,213). The model was trained on 80% of the data (scaled values, cross-validation, 1000 repetitions, shown in rows labeled as "prediction"). Only the best predictors were kept and then used to predict probabilities for the remaining 20% of data points (validation set, shown in rows labeled as "validation"). Log-odds ratios, standard error and 95% confidence intervals (CI) for the validation set have been (beta) standardised.

Predictor	Coefficient	Std. error	Lower CI	Upper CI	p-value	Species	Dataset
percentage_gc_content	-1.27	0.3557	-2.0031	-0.6096	0.000357104	opossum	prediction
percentage_gc_content	-0.5314	0.2027	-0.9434	-0.1466	0.008758284	mouse	prediction
percentage_gc_content	-0.5665	0.1901	-0.9536	-0.2066	0.00287308	rat	prediction
percentage_gc_content	-0.3979	0.1552	-0.7119	-0.1024	0.01035429	rhesus	prediction
as.rvc	0.3618	0.0882	0.1896	0.5359	4.12E-05	human	prediction
percentage_gc_content	-0.9583	0.1558	-1.2734	-0.6622	7.63E-10	human	prediction
percentage_gc_content	-1.438	0.4137	-2.2489	-0.6271	0.000509099	opossum	validation
percentage_gc_content	-0.4325	0.2781	-0.9776	0.1126	0.119942469	mouse	validation
percentage_gc_content	-0.643	0.3373	-1.3042	0.0182	0.056634202	rat	validation
percentage_gc_content	-0.4345	0.198	-0.8226	-0.0463	0.028234012	rhesus	validation
percentage_gc_content	-0.4319	0.1693	-0.7636	-0.1001	0.010729656	human	validation
as.rvc	0.2547	0.1477	-0.0347	0.5441	0.084501745	human	validation

Supplementary Table 9: Analysis of highly expressed circRNAs.

Supplementary Table 9. Highly expressed circRNAs were defined as the circRNAs present in the 90% expression quantile of a tissue in a species. Per species, the circRNAs in the 90% expression quantiles from each of the three tissues were then pooled for further analysis ($n_{opossum}$ = 158, n_{mouse} = 156, n_{rat} = 217, n_{rhesus} = 340, n_{human} = 471) and their properties compared to circRNAs outside the 90% expression quantile. Highly expressed circRNAs are designated "1", others "0". Differences in genomic length, circRNA length, exon number and GLM model performance were assessed with a Student's t-Test; p-values are indicated in the table (ns = non-significant).

Property	Opossum	Mouse	Rat	Rhesus	Human
Genomic length	ns	ns	ns	*p = 0.0043*	*p = 0.047*
circRNA length	ns	ns	ns	ns	ns
Exon number	ns	ns	ns	ns	*p < 0.001*
% of circRNAs expressed in all 3 tissues analysed (1 = highly expressed, 0 = others); more details in **Figure 3-Figure supplement 5A**	0: 2.32% 1: 3.80%	0: 0.82% 1: 8.97%	0: 0.88% 1: 6.45%	0: 4.22% 1: 15.88%	0: 4.35% 1: 12.31%
% of circRNAs detected in a hotspot (1 = highly expressed, 0 = others); more details in **Figure 3-Figure supplement 5B**	0: 37.33% 1: 53.16%	0: 44.95% 1: 67.95%	0: 51.07% 1: 71.89%	0: 51.92% 1: 66.18%	0: 57.06% 1: 72.61%
Median number of circRNAs present in hotspots with at least 1 (= 1) or no (= 0) highly expressed circRNA	0: 3 1: 3	0: 3 1: 3	0: 3 1: 4.5	0: 3 1: 3	0: 3 1: 3
Comparison of GLM model performance between parental genes with and without a highly expressed circRNAs	*p = 0.0163* **Note:** GLM prediction values are higher (driven by a lower GC content)	*ns*	*ns*	*p = 0.05* **Note:** GLM prediction values are higher (driven by genomic length, GC content and exon count)	*p < 0.001* **Note:** GLM prediction values are higher (driven by genomic length, GC content and exon count)
Are highly expressed circRNAs more likely to be shared across species? More details in **Figure 3-Figure supplement 5C** and **Supplementary Table 10**	Yes	Yes	Yes	Yes	Yes

Supplementary Table 10: GLM for highly expressed circRNAs based on 'age groups'.

Supplementary Table 10. A generalised linear model was fitted on the complete dataset to predict the probability of parental genes of highly expressed circRNAs to be produce circRNAs in multiple species ($n_{opossum}$ = 869, n_{mouse} = 844, n_{rat} = 661, n_{rhesus} = 1,673, n_{human} = 2,016). The "sharedness" definition is based on the phylogeny of species as: present in only one species, in rodents (mouse, rat) or primates (rhesus, human), eutherian species (rodents + at least one primate, or primates + at least one rodent) and therian species (opossum + rodents + at least one primate, or opossum + primates + at least one rodents). Log-odds ratios, standard error, 95% confidence intervals (CI) and p-values are shown.

Predictor	Coefficient	Std. Error	Lower CI	Upper CI	p-value	Species
therian	0.9262	0.2171	0.4981	1.3513	2.00E-05	opossum
eutherian	1.1189	0.295	0.5526	1.7156	0.000148951	mouse
rodents	1.2415	0.3833	0.4708	1.9859	0.001199369	mouse
therian	1.7822	0.3092	1.1861	2.4045	8.22E-09	mouse
eutherian	1.1828	0.3223	0.5608	1.8324	0.000242748	rat
rodents	1.189	0.4794	0.189	2.0953	0.01312791	rat
therian	1.6279	0.359	0.9239	2.3407	5.77E-06	rat
eutherian	1.729	0.2151	1.3129	2.1582	9.11E-16	rhesus
primates	1.1084	0.2077	0.7074	1.5237	9.45E-08	rhesus
etherian	1.7435	0.2261	1.3039	2.1925	1.25E-14	rhesus
eutherian	1.3691	0.1818	1.0127	1.7266	5.08E-14	human
primates	1.1663	0.1671	0.8406	1.4966	2.97E-12	human
therian	1.782	0.1884	1.4131	2.1525	3.06E-21	human

Supplementary Table 11: Frequency and enrichment of top-5 dimers in shared and species-specific circRNA loci.

Supplementary Table 11. The total number of detected top-5 dimers in shared and species-specific circRNA loci as well as their enrichment after correction for co-occurrence in multiple RVCs (see **Material and Methods**) are shown. Loci were normalized by the number of detected genes in each category before calculating the enrichment of dimers in shared over species-specific loci. The number of parental genes in both categories is shown below the species name. For mouse, only the top-3 dimers, which are outside the 95% frequency quantile, are shown (see **Material and Methods**). For rhesus, the analysis could only be done on a subset of genes due to lifting uncertainties between the rheMac2 and the rheMac3 genome (see **Material and Methods**).

Species	Dimer	Shared loci	Species-specific loci	Enrichment
opossum n_{shared} = 224 $n_{species-specific}$ = 602	SINE1_Mdo+SINE1_Mdo	4,634	8,155	1.53
	MAR1a_Mdo+MAR1a_Mdo	535	968	1.49
	MAR1a_Mdo+MAR1b_Mdo	474	882	1.45
	SINE1_Mdo+SINE1a_Mdo	371	659	1.51
	MAR1b_Mdo+MAR1b_Mdo	154	276	1.50
mouse n_{shared} = 76 $n_{species-specific}$ = 213	B1_Mus1+B1_Mus2	275	438	1.76
	B2_Mm2+B2_Mm2	268	334	2.25
	B1_Mus1+B1_Mus1	162	274	1.66
rat n_{shared} = 80 $n_{species-specific}$ = 260	ID_Rn1+ID_Rn2	184	457	1.31
	BC1_Rn+ID_Rn2	113	248	1.49
	ID_Rn1+ID_Rn1	111	273	1.32
	BC1_Rn+ID_Rn1	108	273	1.29
	ID_Rn2+ID_Rn2	95	224	1.38
rhesus n_{shared} = 38 $n_{species-specific}$ = 86	AluSx+AluSz	33	38	1.99
	AluY+AluYRa1	32	37	1.93
	AluSx+AluYRa1	27	21	2.86

	AluSx+AluSx1	26	35	1.68
	AluSx1+AluSz	26	32	1.81
human $n_{shared} = 169$ $n_{species-specific} = 811$	AluSx+AluSx1	278	980	1.36
	AluSx1+AluY	274	883	1.49
	AluSx+AluY	269	806	1.60
	AluSx1+AluSz	259	958	1.30
	AluSx+AluSz	257	941	1.31

Figure 1-Figure supplement 1: Overview of the reconstruction pipeline.

Figure 1-Figure supplement 1. Overview of the reconstruction pipeline. CircRNA identification and transcript reconstruction. Unmapped reads from RNA-seq data were remapped and analysed with a custom pipeline. The reconstruction of circRNA transcripts was based on the junction enrichment after RNase R treatment. Further details on the pipeline are provided in the Material and Methods.

Figure 1-Figure supplement 2: Mapping summary of RNA-seq reads.

Figure 1-Figure supplement 2. Mapping summary of RNA-seq reads. Percentage of mapped, unmapped, multi-mapped and BSJ reads across all libraries in untreated and RNase R treated conditions.

Figure 1-Figure supplement 3: General circRNA properties.

Figure 1-Figure supplement 3. General circRNA properties. A: Genomic size. The genomic size (bp) of circRNAs is plotted for all species. B: Transcript size. The transcript size (nt) of circRNAs is plotted for all species. C: Exons per transcript. The number of exons in circRNAs is plotted for all species. For panel A-C, outliers are not plotted *(abbreviations: md = opossum, mm = mouse, rn = rat, rm = rhesus macaque, hs = human).* D: Biotypes of parental genes. For each species, the frequency (%) of different biotypes in the circRNA parental genes was assessed using the ensembl annotation. CircRNA loci that were not found in the annotation were marked as "unknown". E: Presence in multiple tissues. For each species, the frequency (%) of circRNAs detected in one, two or three tissues is plotted. F: Length of different intron types. Distribution of median intron length (log10-transformed) is plotted for different intron types in each gene. *Abbreviations: np = non-parental, po = parental-outside of circRNA, pf = parental-flanking of circRNA, pi = parental-inside of circRNA.*

Figure 1-Figure supplement 4: CircRNA hotspot loci by CPM (opossum, mouse, rat).

Figure 1-Figure supplement 4. CircRNA hotspot loci by CPM (opossum, mouse, rat). In grey, the proportion (%) of circRNA loci that qualify as hotspots and, in purple, the proportion (%) of circRNAs that originate from such hotspots, at three different CPM thresholds (0.01, 0.05, 0.1). The average number of circRNAs per hotspot is indicated above the purple bars.

Figure 2-Figure supplement 1: CircRNA loci overlap between species.

A: Identified clusters for overlapping circRNA loci based on "parental gene", "circRNA locus" and "start/stop exon"

B: Gain of evolutionary precision by including multiple species (based on "circRNA locus")

1. Classical mouse – human comparison to determine mammalian circRNAs

2. Adding of an additional rodent or primate species to the mouse – human comparison

3. Adding of an outgroup to the rodent – primate comparison

mammalian circRNAs: 440 (43.91%)
-> 440/1002 = 0.4391

mammalian circRNAs: 369 (36.83%)
-> 369/1002 = 0.3683
-> reduction of shared loci by 16.36%
(71 loci less / 440 = 0.1636)

mammalian circRNAs: 260 (25.95%)
-> 260/1002 = 0.2595
-> reduction of shared loci by 40.91%
(180 loci less / 440 = 0.4091)

Figure 2-Figure supplement 1. CircRNA loci overlap between species. A: Upper panel: The presence of circRNA in multiple species can be identified on the gene level (= "parental gene"), based on the location of the circRNA within the gene (= "circRNA locus") or the overlap of the first and last exons of the circRNA (= "start/stop exon"). Depending on the chosen stringency, the number of circRNA loci present in multiple species varies. For example: when considering the parental gene level (shown to the left), all four circRNAs depicted in the hypothetical example of this figure (*circRNA-A.1*, *circRNA-A.2*, *circRNA-B.1* and *circRNA-B.1*) are located in the same orthologous locus. In contrast, when looking at the start and stop exons (right), only two circRNAs (*circRNA-A.1* and *circRNA-B.1*) are generated from the same orthologous locus, whereas *circRNA-A.2* and *circRNA-B.2* - previously classified as "orthologous" - are now found in different loci and labeled as species-specific. Depending on the classification, the number of shared circRNA loci thus differs and may influence the interpretation of results. Lower panel: For each classification, orthology clusters were counted and grouped by their overlap (in purple when present in primates, rodents, eutherians or therians; in red when species-specific). Please note that in our study, we apply the definition shown in the middle panels (which are identical to main **Figure 2A**) that considers exon overlap as relevant. B: Figure shows the loss of shared circRNA loci (based on "circRNA locus" definition) by adding additional species to the classical mouse – human comparison. All comparisons are made with mouse as reference to which the other loci are compared. The reduction of loci (%) by adding additional species is indicated below each figure.

Figure 2-Figure supplement 2: Amplitude correlations.

Figure 2-Figure supplement 2. Amplitude correlations. Plotted is the correlation (Spearman's rho) between the amplitude and the GC content of introns (light brown) and exons (dark brown). *Abbreviations: np = non-parental, po = parental, outside of circRNA, pi = parental, inside of circRNA.*

Figure 3-Figure supplement 1: Replication time, gene expression steady-state levels and GHIS of human parental genes.

Figure 3-Figure supplement 1. Replication time, gene expression steady-state levels and GHIS of human parental genes. A: Replication time of parental genes. Values for the replication time were used as provided in (Koren et al., 2012). They were normalised to a mean of 0 and a standard deviation of 1. Differences between non-parental genes (n_{total} = 18,134) and parental genes (n_{total} = 2,058) were assessed by a one-tailed Mann-Whitney U test. B: Gene expression steady-state levels of parental genes. Mean steady-state expression levels were used as provided in (Pai et al., 2012). Differences between non-parental genes (n_{total} = 14,414) and parental genes (n_{total} = 2,058) were assessed by a one-tailed Mann-Whitney U test. C: GHIS of parental genes. GHIS was used as provided in (Steinberg et al., 2015). Differences between non-parental genes (n_{total} = 17,438) and parental genes (n_{total} = 1,995) were assessed by a one-tailed Mann-Whitney U test. *(Note C-D: Outliers for all panels were removed prior to plotting. Significance levels: '***' < 0.001, '**' < 0.01, '*' < 0.05, 'ns' >= 0.05).*

Figure 3-Figure supplement 2: Distribution of prediction values for non-parental and parental circRNA genes.

Figure 3-Figure supplement 2. Distribution of prediction values for non-parental and parental circRNA genes. The density of predicted values for non-parental (grey) and parental (purple) genes is plotted for each species based on the predictors identified by the GLM in each species.

Figure 3-Figure supplement 3: Properties of 'functional circRNAs' from literature.

Figure3-Figure supplement 3. Properties of 'functional circRNAs' from literature. A: Prediction values of linear regression model for human circRNA parental and non-parental genes as previously defined (**Materials and Methods**). Functional circRNAs as described in (Chen, 2020) are plotted in pink on top of the boxplot and are separated by whether they are in a non-parental or parental gene. B-D: GC content, repeat fragments (in antisense, normalized by genomic length of parental gene) and number of exons for human non-parental and parental circRNA genes; values for functional circRNAs are plotted in pink.
Parental genes of functional circRNAs listed in Chen et al. 2021, which were identified in our study: SHPRH, ZNF609, GCN1L1, HIPK2, HIKP3, ZNF91, BIRC6, FOXO3, MBNL1, ASAP1, PAN3, SMARCA5, ITCH.

Figure 3-Figure supplement 4: Validation of parental gene GLM on Werfel *et al.* dataset.

Figure 3-Figure supplement 4. Validation of parental gene GLM on Werfel *et al.* dataset. A: Mouse. To assess the parental gene properties identified by this study, the generalised model was used to predict circRNA parental genes on data from an independent study. The density plot "Prediction values" shows the predicted values for non-parental genes in both datasets (((Werfel et al., 2016) and data from this publication, n = 11,963, in grey and labeled as -/-), parental genes only present in the Werfel dataset (n = 2,843, light pink, labeled as -/+), parental genes only present in this study's underlying dataset (n = 210, dark pink, labeled as +/-) and parental genes that were present in both datasets (n = 638, purple, labeled as +/+). The plots "GC content", "Number of exons" and "Repeat fragments (as)" (the latter normalized by the genomic length of the parental gene) show the properties of circRNA parental genes (highlighted in purple) as identified by Werfel *et al*. B: Human. Same plot outline as for mouse. The number of non-parental genes in both datasets is n = 10,591; 2,724 parental genes are only present in the Werfel dataset and 356 parental genes only in our dataset. The overlap between both datasets is n = 1,666.

Figure 3-Figure supplement 5: Properties of highly expressed circRNAs.

Figure 3-Figure supplement 5. Properties of highly expressed circRNAs. A: Presence of highly expressed circRNAs in multiple tissues. Plot shows the percentage (%) of circRNAs from the 90% expression quantile ($n_{opossum}$ = 158, n_{mouse} = 156, n_{rat} = 217, n_{rhesus} = 340, n_{human} = 471), which is present in one, two or three of the tissues analysed compared to circRNAs outside the 90% expression quantile. For each species, distributions were compared using Fisher's exact test, p-values are shown above each barplot. B: Presence of highly expressed circRNAs in hotspots. Plot shows the percentage (%) of circRNAs from the 90% expression quantile, which is found in a hotspot compared to circRNAs outside the 90% expression quantile. For each species, distributions were compared using Fisher's exact test, p-values are shown above each barplot. C: Presence of highly expressed circRNAs in 'age groups'. Plot shows the percentage (%) of circRNAs from the 90% expression quantile, which is present in different 'age groups' compared to circRNAs outside the 90% expression quantile. Age groups were defined as whether circRNA is species-specific (age = 1), lineage-specific (age = 2), eutherian (age = 3) or shared across all therian species (age = 4). Log-odds ratio and significance levels (*significance levels based on p-value: '***' < 0.001, '**' < 0.01, '*' < 0.05, 'ns' >= 0.05*) were calculated using a generalised linear model (see **Supplementary Table 10**) and are shown for the respective age groups and species.

Figure 4-Figure supplement 1: Enrichment of transposable elements in flanking introns for opossum.

Figure 4-Figure supplement 1. Enrichment of transposable elements in flanking introns for opossum. The number of transposable elements was quantified in both intron groups (circRNA flanking introns and length- and GC-matched control introns). Enrichment of transposable elements is represented by colour from high (dark purple) to low (grey). The frequency distributions of TEs in background and flanking introns were compared using a Wilcoxon Signed Rank Test; p-value is shown in the upper right corner.

Figure 4-Figure supplement 2: PCA and phylogeny of opossum, rat, rhesus macaque and human repeat dimers.

Figure 4-Figure supplement 2. PCA and phylogeny of opossum, rat, rhesus macaque and human repeat dimers. A: Opossum. Panel A shows the PCA for dimer clustering based on a merged and normalised score, taking into account binding phylogenetic distance, binding capacity of TEs to each other and absolute frequency. Absolute frequency is also represented by circle size. The top- ranked dimers are indicated. Circles around the discs represent cases where the TE binds to itself. Furthermore, a phylogeny of opossum transposable elements is shown, the top-5 dimers are highlighted with purple shading. Phylogenetic trees are based on multiple alignments with Clustal-Omega. Several TE families have independent origins, which cannot be taken into account with Clustal-Omega. These cases are indicated by a grey, dotted line and TE origins - if known - have been manually added. We deemed this procedure sufficiently precise, given that the aim was to only visualise the general relationship of TEs. TEs used as outgroups, as well TEs that merged are indicated with a red line. B-D: Same analysis as in Panel A, but for rat, rhesus macaque and ruman, respectively.

Figure 5-Figure supplement 1: Contribution of species-specific repeats to the formation of shared circRNA loci.

Figure 5-Figure supplement 1. Contribution of species-specific repeats to the formation of shared circRNA loci. Dimer enrichment in shared and species-specific repeats in opossum, mouse and rhesus macaque. The frequency (number of detected dimers in a given parental gene), log2-enrichment (shared vs. species-specific) and mean age (defined as whether repeats are species-specific: age = 1, lineage-specific: age = 2, eutherian: age = 3, therian: age = 4) of the top-100 most frequent and least frequent dimers in parental genes with shared and species-specific circRNA loci in opossum, mouse and rhesus macaque were analysed and compared with a Wilcoxon Signed Rank Test. Frequencies are plotted on the x- and y-axis, point size reflects the age and point colour the enrichment (blue = decrease, red = increase). Based on the comparison between shared and species-specific dimers, the top-5 dimers defined by frequency and enrichment are highlighted and labelled in red.

Figure 5-Figure supplement 2: Repeat interaction landscape in shared vs. species-specific circRNA loci.

Figure 5-Figure supplement 2. Repeat interaction landscape in shared vs. species-specific circRNA loci. Upper left: graphical representation of possible repeat interactions (= dimers that can be formed) across RVCs. Afterwards: Frequency distribution of possible interactions of a given repeat (from the top-5 dimers, based on **Figure 5A** and **Figure 5-Figure supplement 1**) in parental genes of species-specific (red) and shared (blue) circRNA loci in opossum, mouse, rat, rhesus macaque and human. The enrichment of possible interactions (shared vs. species-specific, based on each distribution's median) is indicated above each plot.

Figure 5-Figure supplement 3: MilliDivs and MFE for dimers in shared and species-specific circRNA loci.

MilliDivs and MFEs for opossum, mouse, rat and rhesus macaque

Figure 5-Figure supplement 3: MilliDivs and MFE for dimers in shared and species-specific circRNA loci. Left panel of each species: MilliDiv values were compared between parental genes of species-specific (red) and shared (blue) circRNA loci using a Student's t-Test (alternative = "less") with corresponding p-values plotted above each boxplots. Since dimers are composed of two repeats, the mean milliDiv value between both repeats was taken. Right panel of each species: Violin Plots depicting the minimal free energy (MFE) of genomic sequences for dimers in species-specific (red) and shared (blue) circRNA loci. For each gene, the "least degraded dimer" was chosen to calculate its MFE value leading to a strong enrichment of only a few of the top-5 dimers

(see **Material and Methods**). The "maximum" MFE possible, which is based on the dimer formed by each TE's reference sequence (downloaded from RepBase (Bao et al., 2015)), is depicted with a grey line below each pair of violin plots. Each distribution's median is indicated with a grey point. MFE values between species-specific and shared circRNA loci were compared with a Student's t-Test; corresponding p-values are indicated above each pair of violin plots.

Bibliography

Bao W, Kojima KK, Kohany O. 2015. Repbase Update, a database of repetitive elements in eukaryotic genomes. *Mob DNA* **6**:11. doi:10.1186/s13100-015-0041-9

Chen L-L. 2020. The expanding regulatory mechanisms and cellular functions of circular RNAs. *Nat Rev Mol Cell Biol* **21**:475–490. doi:10.1038/s41580-020-0243-y

Koren A, Polak P, Nemesh J, Michaelson JJ, Sebat J, Sunyaev SR, McCarroll SA. 2012. Differential relationship of DNA replication timing to different forms of human mutation and variation. *Am J Hum Genet* **91**:1033–1040. doi:10.1016/j.ajhg.2012.10.018

Pai AA, Cain CE, Mizrahi-Man O, De Leon S, Lewellen N, Veyrieras J-B, Degner JF, Gaffney DJ, Pickrell JK, Stephens M, Pritchard JK, Gilad Y. 2012. The contribution of RNA decay quantitative trait loci to inter-individual variation in steady-state gene expression levels. *PLoS Genet* **8**:e1003000. doi:10.1371/journal.pgen.1003000

Steinberg J, Honti F, Meader S, Webber C. 2015. Haploinsufficiency predictions without study bias. *Nucleic Acids Res* **43**:e101. doi:10.1093/nar/gkv474

Werfel S, Nothjunge S, Schwarzmayr T, Strom T-M, Meitinger T, Engelhardt S. 2016. Characterization of circular RNAs in human, mouse and rat hearts. *J Mol Cell Cardiol* **98**:103–107. doi:10.1016/j.yjmcc.2016.07.007