The SWI/SNF chromatin remodeling assemblies BAF and PBAF differentially

regulate cell cycle exit and cellular invasion *in vivo*

Jayson J. Smith[1], Yutong Xiao[1], Nithin Parsan[2], Taylor N. Medwig-Kinney[1], Michael A. Q.

Martinez[1], Frances E. Q. Moore[1], Nicholas J. Palmisano[1], Abraham Q. Kohrman[3], Mana

Chandhok Delos Reyes[4], Rebecca C. Adikes[1], Simeiyun Liu[1,5], Sydney A. Bracht[1,6], Wan

Zhang[1], Kailong Wen[7,8], Paschalis Kratsios[7,8], David Q. Matus[1,9]

[1]Department of Biochemistry and Cell Biology, Stony Brook University, Stony Brook, NY

11794, USA

[2]Massachusetts Institute of Technology, Cambridge, MA 02139, USA

[3]Department of Molecular Biology, Princeton University, Princeton, NJ 08544, USA

[4]10 Grant St., Somerville, MA 02145, USA

[5]Molecular, Cellular and Developmental Biology, University of California Santa Cruz,

Santa Cruz, CA 95064, USA

[6]Department of Cell Biology, Johns Hopkins University, Baltimore, MD 21205, USA

[7]The Grossman Institute for Neuroscience, Quantitative Biology, and Human Behavior,

University of Chicago, Chicago, IL 60637, USA

[8]Department of Neurobiology, University of Chicago, Chicago, IL 60637, USA

[9]Lead Contact

[*]Correspondence: david.matus@stonybrook.edu

SUMMARY STATEMENT

Cellular invasion through the basement membrane by the *C. elegans* anchor cell requires both BAF and PBAF SWI/SNF assemblies to arrest the cell cycle and promote the expression of pro-invasive genes.

SUMMARY

Chromatin remodelers such as the SWI/SNF complex coordinate metazoan development through broad regulation of chromatin accessibility and transcription, ensuring normal cell cycle control and cellular differentiation in a lineage-specific and temporally restricted manner. Mutations in genes encoding the structural subunits of chromatin, such as histone subunits, and chromatin regulating factors (CRFs) are associated with a variety of disease mechanisms including cancer metastasis, in which cancer co-opts cellular invasion programs functioning in healthy cells during development. Here we utilize *Caenorhabditis elegans* anchor cell (AC) invasion as an *in vivo* model to identify the suite of chromatin agents and CRFs that promote cellular invasiveness. We demonstrate that the SWI/SNF ATP-dependent chromatin remodeling complex is a critical regulator of AC invasion, with pleiotropic effects on both G_0 cell cycle arrest and activation of invasive machinery. Using targeted protein degradation and enhanced RNA interference (RNAi) vectors, we show that SWI/SNF contributes to AC invasion in a dose-dependent fashion, with lower levels of activity in the AC corresponding to aberrant cell cycle entry and increased loss of invasion. Our data specifically implicate the SWI/SNF BAF assembly in the regulation of the G_0 cell cycle arrest in the AC, whereas the SWI/SNF PBAF assembly

promotes AC invasion via cell cycle-independent mechanisms, including attachment to

the basement membrane (BM) and activation of the pro-invasive *fos-1*/FOS gene.

Together these findings demonstrate that the SWI/SNF complex is necessary for two

essential components of AC invasion: arresting cell cycle progression and remodeling the

BM. The work here provides valuable single-cell mechanistic insight into how the

SWI/SNF assemblies differentially contribute to cellular invasion and how SWI/SNF

subunit-specific disruptions may contribute to tumorigeneses and cancer metastasis.

Keywords: Chromatin regulating factors, chromatin remodeling, SWI/SNF, BAF, PBAF,

cell invasion, cell cycle exit, *C. elegans*

INTRODUCTION

Cellular invasion through basement membranes (BMs) is a critical step in metazoan development and is important for human health and fitness. Early in hominid development, trophoblasts must invade into the maternal endometrium for proper blastocyst implantation (1). In the context of immunity, leukocytes become invasive upon injury or infection to travel between the bloodstream and interstitial tissues (2,3). Atypical activation of invasive behavior is associated with a variety of diseases, including rheumatoid arthritis wherein fibroblast-like synoviocytes adopt invasive cellular behavior, leading to the expansion of arthritic damage to previously unaffected joints (4,5). Aberrant activation of cell invasion is also one of the hallmarks of cancer metastasis (6).

A variety of *in vitro* and *in vivo* models have been developed to study the process of cellular invasion at the genetic and cellular levels. *In vitro* invasion assays typically involve 3D hydrogel lattices, such as Matrigel, through which cultured metastatic cancer cells will invade in response to chemo-attractants (7). Recently, microfluidic systems have been integrated with collagen matrices to improve these *in vitro* investigations of cellular invasion (8). While *in vitro* invasion models provide an efficient means to study the mechanical aspects of cellular invasion, they are currently unable to replicate the complex microenvironment in which cells must invade during animal development and disease. A variety of *in vivo* invasion models have been studied, including cancer xenograft models in mouse (9–11) and zebrafish (12,13), each having their respective benefits and drawbacks. Over the past ~15 years, *Caenorhabditis elegans* anchor cell (AC) invasion has emerged as a powerful alternative model due to its visually tractable single-cell nature (**Fig 1 A**) (14).

Figure 1. Summary of *C. elegans* AC invasion through the underlying BM and AC GRN. (A) Schematic depicting AC invasion in the mid-L3 stage of *C. elegans* development (left) and micrographs demonstrating the coordination of AC (magenta, *cdh-3p::PH::mCherry*) invasion through the BM (green, *laminin::GFP*) with primary vulval development. The fluorescent AC-specific membrane marker and BM marker are overlaid on DIC in each image. White arrowheads indicate AC(s), yellow arrowheads indicate boundaries of breach in BM, and white brackets indicate 1° VPCs. **(B)** Overview of the transcription factor GRN governing AC invasion (22,24), which consists of cell cycle-independent (*fos-1*) and dependent (*egl-43, hlh-2,* and *nhr-67*) subcircuits, which together with *hda-1* promote pro-invasive gene expression and maintain cell cycle arrest in the AC. In this and all subsequent figures, scale bar, 5 μm.

Previous work demonstrated a high degree of evolutionary conservation in the cell-

autonomous mechanisms underlying BM invasion (3,15), including basolateral

polarization of the F-actin cytoskeleton/cytoskeletal regulators and the expression of

matrix metalloproteinases (MMPs) (16–21). Moreover, in order to breach the BM, the AC

requires the expression of transcription factors (TFs), whose human homologs are

common to metastatic cancers, including *egl-43* (EVI1/MEL), *fos*-1 (FOS), *hlh-2*

(E/Daughterless), and *nhr-67* (TLX/Tailless) (22) (**Fig 1 B**). In addition to the expression

of pro-invasive genes, there is increasing evidence that cells must also arrest in the cell

cycle to adopt an invasive phenotype (23). Our previous work has demonstrated that the

AC must terminally differentiate and arrest in the G_0/G_1 phase of the cell cycle to invade

the BM and make contact with the underlying primary vulval precursor cells (1° VPCs)

(22,24). The regulatory mechanisms that couple G_0/G_1 cell cycle arrest with the ability of

a cell to invade the BM remain unclear.

Cell-extrinsic and cell-intrinsic factors, such as chromatin remodeling complexes

and TFs, can influence the decision to maintain a plastic cell fate or to undergo cell fate

specification and terminal differentiation. The determination of a cell to remain plastic or

specify is in part the consequence of a complex, genome-wide antagonism between

Polycomb group (PcG) transcriptional repression and Trithorax group (TrxG)

transcriptional activation (25–27). For example, the binding of pioneer transcription

factors OCT4 and SOX2 to target DNA in order to retain pluripotency in murine embryonic

stem cells is the indirect consequence of the regulation of chromatin accessibility at these

target regions (28). A recent study has shown that chromatin accessibility of enhancers

in crucial cell cycle genes which promote the G_1/S transition, including Cyclin E and E2F

transcription factor 1, is developmentally restricted to reinforce terminal differentiation and

cell cycle exit during *Drosophila melanogaster* pupal wing morphogenesis (29). In *C.*

elegans myogenesis, the SWItching defective/Sucrose Non-Fermenting (SWI/SNF) ATP-

dependent chromatin remodeling complex, a member of the TrxG family of complexes,

both regulates the expression of the MyoD transcription factor (*hlh-1*) and acts

redundantly to promote differentiation and G_0 cell cycle arrest with core cell cycle

regulators: cullin 1 (CUL1/*cul-1)*, cyclin-dependent kinase inhibitor 1 (*cki-1)*, FZR1 (*fzr*-1),

and the RB transcriptional corepressor (RBL1/*lin*-35). The importance of regulation of

chromatin states and the activity of CRFs for the acquisition and implementation of

differentiated behaviors is also reflected in the *C. elegans* AC, as previous work has

shown that the histone deacetylase *hda-1* (HDAC1/2) is required for pro-invasive gene

expression and therefore differentiated cellular behavior and invasion (24) (**Fig 1 B**).

Given these findings, a comprehensive investigation of the regulatory mechanism(s)

governing AC invasion should include a thorough description of the suite of CRFs

required for G_0/G_1 cell cycle arrest and invasive differentiation in the AC.

In this study we perform an RNA interference (RNAi) screen of *C. elegans* CRFs,

specifically focusing on genes involved in chromatin structure and remodeling or histone

modification. We identify 82 genes whose RNAi depletion phenotype resulted in a

significant AC invasion defect. Among the CRFs identified as significant regulators of AC

invasion, the SWI/SNF complex emerged as the most well-represented single chromatin

remodeling complex. RNAi knockdown of subunits specific to the SWI/SNF core (*swsn-1*

and *snfc-5/swsn-5*), and both BAF (BRG/BRM-Associated Factors; *swsn-8/let-526*) and

PBAF (Polybromo Associated BAF; *pbrm-1* and *swsn-7*) assemblies resulted in penetrant

loss of AC invasion. We generated fluorescent reporter knock-in alleles of subunits of the

core (*GFP::swsn-4*) and BAF (*swsn-8::GFP*) assembly of the SWI/SNF complex using

CRISPR/Cas9-mediated genome engineering. These alleles, used in conjunction with an

endogenously labeled PBAF (*pbrm-1::eGFP*) assembly subunit, enabled us to determine

the developmental dynamics of the SWI/SNF ATPase and assembly-specific subunits,

gauge the efficiency of various SWI/SNF knockdown strategies, and asses inter- and

intra-complex regulation. Using improved RNAi constructs and an anti-GFP nanobody

degradation strategy (30), we demonstrated that the cell autonomous contribution of the

SWI/SNF complex to AC invasion is dose dependent. This finding parallels similar studies

in cancer (31–34) and *C. elegans* mesoblast development (35). Examination using a CDK

activity sensor (36) revealed assembly-specific contributions to AC invasion: whereas

BAF promotes AC invasion in a cell cycle-dependent manner, PBAF contributes to

invasion in a cell cycle-independent manner. Finally, we utilized the auxin-inducible

degron (AID) system combined with PBAF RNAi to achieve strong PBAF subunit

depletion in the AC, which resulted in loss of both AC invasion and adhesion to the BM.

Together, these findings provide insight into how the SWI/SNF complex assemblies may

contribute to distinct aspects of proliferation and metastasis in human malignancies.

RESULTS

An RNAi screen of 269 CRFs identified SWI/SNF as a key regulator of AC invasion

To identify the suite of CRFs that, along with *hda-1,* contribute to AC invasion, we

generated an RNAi sub-library of 269 RNAi clones from the complete Vidal RNAi library

and a subset of the Ahringer RNAi library (37,38) targeting genes implicated in chromatin

state, chromatin remodeling, or histone modification (**Table S1**) (**Fig 1 B**). Because CRFs

act globally to control gene expression, we screened each RNAi clone by high-resolution

differential interference contrast (DIC) and epifluorescence microscopy in a uterine-

specific RNAi hypersensitive background containing labeled BM (*laminin::GFP*) and an

AC reporter (*cdh-3p::PH::mCherry*) (**Fig 1 A; Table S1**) (14,22,24,39). This genetic

background allowed us to limit the effect of transcriptional knockdown of CRFs to only

affect the AC and the neighboring uterine tissue, and only for a time period following the

specification of the AC (39). As the neighboring uterine cells do not contribute to the

invasion program (14), AC invasion defects following RNAi treatments are indicative of

cell autonomous pro-invasive gene function (24,39). In wild-type animals, by the time the

1° fated P6.p vulval precursor cell has divided twice (P6.p 4-cell stage), 100% of ACs

have successfully breached the underlying BMs and made contact with the P6.p grand-

daughters (14). While the AC also always invades in the uterine-specific RNAi

hypersensitive strain we utilized for our CRF screen, there is a low penetrance delay,

such that at the P6.p 4-cell stage, when we scored invasion, 2% (2/100 animals) still had

an intact BM. Thus, we used this baseline defect as a statistical reference point for this

genetic background and defined the cut-off threshold for significant defects in invasion

from RNAi depletion of CRFs in our screen as those RNAi clones that resulted in a ~13%

or greater defect in invasion (4/30 animals, Fisher's exact test = 0.0252). By this

threshold, we recovered 82 CRFs (30.5% of total CRFs screened) that significantly

regulate AC invasion, suggesting a general role for regulation of chromatin states in the

acquisition of invasive behavior (**Table S2**). Interestingly, five subunits of the broadly

conserved SWI/SNF chromatin remodeling complex were recovered as significant

regulators of AC invasion: *swsn-1*(SMARCC1/SMARCC2; 23% AC invasion defect),

swsn-5/snfc-5 (SMARCB1; 20% AC invasion defect), *swsn-7* (ARID2; 23% AC invasion

defect), *swsn-8/let-526* (ARID1A/ARID1B; 23% AC invasion defect), and *pbrm-1*

(PBRM1; 20% AC invasion defect) (**Table S2**). As such, SWI/SNF is well-represented in

our CRF screen, with representation of the core (*swsn-1* and *swsn-5*), BAF (*swsn-8*) and

PBAF (*pbrm-1* and *swsn-7*) assemblies. Given the prevalence of SWI/SNF subunits

recovered as significant regulators of AC invasion in our RNAi screen and the crucial role

SWI/SNF plays in the regulation of animal development (40–45), tumorigenesis (32,46–

48), and cell cycle control (35,49–52), we chose to focus our efforts on characterizing the

role of the SWI/SNF complex in promoting AC invasion.

To confirm our RNAi results implicating the SWI/SNF complex in the promotion of

AC invasion, we obtained two temperature sensitive hypomorphic alleles, *swsn-1(os22)*

and *swsn-4(os13)* (41)*,* and scored for defects in AC invasion in a genetic background

containing both BM (*laminin::GFP*) and AC (*cdh-3p::mCherry::moeABD*) reporters. While

we observed no defects in AC invasion in animals grown at the permissive temperature

(15°C) (**Fig S1 A**), animals containing hypomorphic alleles for core subunits *swsn-1* and

swsn-4 cultured at the restrictive temperature (25°C) displayed defects in 20% (10/50)

and 24% (12/50) of animals, respectively. These results with the *swsn-1(os22)* allele

corroborated our *swsn-1(RNAi)* data from the CRF RNAi screen. Since neither of the

RNAi libraries used to compose the CRF screen (see above) contained a *swsn-4(RNAi)*

clone, results with the *swsn-4(os13)* allele also supplement data from our CRF RNAi

screen by suggesting that AC invasion depends on the expression of the sole *C. elegans*

SWI/SNF ATPase subunit in addition to the 5 subunits identified in the screen (**Fig S1 B**).

Improved RNAi vectors revealed distinct contributions of SWI/SNF subunits to AC

invasion

Though many SWI/SNF assemblies have been described in mammalian and other

systems, including BAF, PBAF, esBAF, GBAF, nBAF, and npBAF) (53), to date, BAF and

PBAF are the only SWI/SNF assemblies that have been described in *C. elegans*. Both

assemblies consist of core subunits (SWSN-1, SWSN-4, SWSN-5) and accessory

subunits (DPFF-1, SWSN-2.1/HAM-3, SWSN-2.2, SWSN-3, SWSN-6, and PHF-10),

collectively referred to as common factors (46,54). These common factors are bound by

assembly-specific subunits in a mutually exclusive manner, which confers the distinct

character of each of the two assemblies (**Fig 2 A**). Due to the absence of thorough

biochemical investigation into the SWI/SNF complex in *C. elegans*, previous publications

have classified subunits as part of the SWI/SNF core, accessory, or BAF/PBAF

assemblies based on homology and phenotypic analyses (35,43,52,55). The prevailing

model for the two SWI/SNF assemblies in *C. elegans* is that either the SWSN-8 subunit

associates with common factors to form the BAF assembly, or the SWSN-7, SWSN-9,

and PBRM-1 subunits associate with common factors to form the PBAF assembly

(43,55,56). Prior investigations into SWI/SNF have revealed a wide array of

developmental contexts in which the BAF and PBAF assemblies have overlapping and

distinct roles in the regulation of cell cycle control, differentiation, and differentiated

behavior (35,52,55,57–61).

To investigate the contribution of individual SWI/SNF subunits to AC invasion and

to distinguish potentially distinct roles of the BAF and PBAF assemblies, we generated

improved RNAi constructs utilizing the T444T vector (62) to target representative subunits

of the core and both SWI/SNF assemblies (**Table S3**). Knockdown of SWI/SNF subunits

in whole-body RNAi sensitive animals following treatment with T444T RNAi vectors

Figure 2. Enhanced RNAi targeting SWI/SNF core, BAF, and PBAF assembly subunits results in penetrant invasion defects. (A) Schematic depicting the *C. elegans* SWI/SNF core (middle, green) and accessory (middle, gray) subunits and BAF (left, blue), and PBAF (right, orange) assemblies. **(B-D)** DIC (left), corresponding fluorescence images (middle), and fluorescence overlay (right) representing loss of AC (magenta, *cdh-3p::mCherry::moeABD*) invasion through the BM (green, *laminin::GFP*) following RNAi depletion of SWI/SNF core (*swsn-1* and *swsn-4*) **(B)**, BAF (*swsn-8*) **(C)**, and PBAF (*pbrm-1, swsn-7,* and *swsn-9*) **(D)** subunits. In cases where multiple cells expressed the AC reporter (2+ACs) in the same animal, each is indicated with a white arrowhead. In cases where multiple cells expressed the AC reporter (2+ACs), a representative image from the same treatment of a single AC that fails to breach the BM is displayed as an inset (white dashed box, bottom left). White arrowheads indicate AC(s), yellow arrowheads indicate boundaries of breach in BM, and white brackets indicate 1° VPCs. **(E)** Stacked bar chart showing the penetrance of AC invasion defects following treatment with SWI/SNF RNAi depletion, binned by AC phenotype (n≥50 animals examined for each treatment).

resulted in penetrant loss of invasion. The majority of ACs failed to invade following

treatment with RNAi targeting the core SWI/SNF ATPase subunit *swsn-4* or core subunit

swsn-1 (90% and 94%, respectively; n=50 animals; **Fig 2 B, E**)*.* RNAi-mediated

knockdown of the BAF assembly subunit *swsn-8* also resulted in significant loss of AC

invasion (32%; n=50 animals; **Fig 2 C, E**). Knockdown of the PBAF assembly subunits

with *pbrm-1(RNAi), swsn-7(RNAi)*, or *swsn-9(RNAi)* resulted in a less penetrant loss of

AC invasion (18%, 20%, and 12%, respectively; n=50 animals; **Fig 2 D, E**). Importantly,

at least one cell in the ventral uterus expressed the fluorescent AC reporter in all animals

across all treatments, suggesting that loss of the SWI/SNF complex does not preclude

the birth of an AC.

Interestingly, in addition to a single non-invasive AC phenotype, RNAi-mediated

knockdown of *swsn-1*, *swsn-4* or *swsn-8* also resulted in a second phenotype

characterized by multiple uterine cells expressing the AC reporter (*cdh-*

3p::mCherry::moeABD) which failed to invade the BM (*laminin::GFP*) (32%, 30% and 8%,

respectively; **Fig 2 B-C, E**). In all instances where more than one cell expressed the AC

reporter, no breach in the underlying BM was detected at the P6.p 4-cell stage. In

contrast, only the single non-invasive AC phenotype resulted from RNAi treatment

targeting PBAF assembly subunits (**Fig 2 D-E**). These results suggest that the SWI/SNF

assemblies BAF and PBAF may promote AC invasion through distinct mechanisms,

perhaps via regulation of both a cell cycle-dependent and -independent mechanism,

respectively.

Characterization of endogenous GFP reporter alleles and the efficacy of improved

SWI/SNF RNAi vectors

Next, to confirm expression of SWI/SNF subunits in the AC and to quantitatively

assess the potency of our enhanced SWI/SNF RNAi vectors we utilized CRISPR/Cas9

genome engineering to generate GFP-tagged alleles of *swsn-4* and *swsn-8*, inserting a

codon-optimized GFP tag into the 5' end and 3' end of the *swsn-4* and *swsn-8* loci,

respectively (**Fig S2 A**) (63). The GFP-tagged endogenous strains showed ubiquitous

and nuclear-localized expression of GFP::SWSN-4 and SWSN-8::GFP throughout the *C.*

elegans developmental life cycle (**Fig S2 B**). We also obtained a strain containing an

endogenously eGFP-labeled PBAF subunit (*pbrm-1::eGFP*) from the *Caenorhabditis*

Genetics Center (CGC). We quantified fluorescence protein expression of SWI/SNF core

ATPase (GFP::SWSN-4), BAF (SWSN-8::GFP), and PBAF (PBRM-1::eGFP) subunits in

the AC during vulval development across the L3 and early L4 stages, as defined by the

division pattern of the 1°-fated VPCs (14) (n≥28 animals per stage; **Fig S3 A-C'**).

Expression of all three subunits was enhanced in the AC relative to the neighboring

ventral uterine (VU; *swsn-4*: 18%, *swsn-8*: 21%, *pbrm-1*: 17% enhanced) and 1° VPC

(*swsn-4*: 30%, *swsn-8*: 38%, *pbrm-1*: 23% enhanced) lineages during AC invasion (P6.p

2 cell – 4 cell stage, **Fig S3 A'-C'**). Late in vulval development at the P6.p 8 cell stage,

expression of GFP::SWSN-4 and PBRM-1::eGFP increases in the 1° VPCs and is no

longer statistically separable from expression in the AC, whereas expression of SWSN-

8::GFP in the VPCs remains significantly lower than in the AC (**Fig S3 A'-C'**).

We treated SWI/SNF endogenously labeled GFP-tagged strains with our RNAi

vectors to precisely quantify the efficiency of RNAi-mediated knockdown of target

SWI/SNF complex subunits and to correlate this loss with the resulting AC phenotypes.

Treatment with either *swsn-4(RNAi)* or *swsn-8(RNAi)* vectors resulted in robust depletion

of fluorescence expression of GFP::SWSN-4 (94% depletion) and SWSN-8::GFP (81%

depletion) in the AC (**Fig S4 A-B, D**) and penetrant loss of invasion (90% and 30%,

respectively; n=30 animals for each condition; **Fig S4 E**). We also noted instances where

multiple cells expressed the AC reporter (23% and 10%, respectively; n=30 animals for

each condition; **Fig S4 E**). Treatment of the PBRM-1::eGFP strain with *pbrm-1(RNAi)*

revealed weaker but significant knockdown of PBRM-1 protein (49% depletion), and a

lower penetrance of invasion defects (17%; n=30 animals; **Fig S4 C-E**)*.* It is unclear why

PBRM-1::eGFP endogenous protein level in the AC of animals treated with enhanced

pbrm-1(RNAi) remains considerably higher compared to treatment of strains containing

swsn-4 or *swsn-8* reporter alleles with their respective RNAis. We hypothesize that this

may be the consequence of differential protein perdurance. Altogether, these results

confirm the dynamic expression of the SW/SNF core, BAF, and PBAF subunits in the AC

before, during, and after invasion and demonstrate the effectiveness of our improved

SWI/SNF-targeting RNAi vectors.

***C. elegans* SWI/SNF subunits exhibit low levels of intracomplex cross-regulation**

Work in mammalian cell culture has revealed that the mSWI/SNF complex is

assembled in a step-wise fashion, with stability of the complex as a whole and association

of individual subunits depending on the prior expression and association of other subunits

(64)**.** To date it is unknown whether in *C. elegans* individual SWI/SNF subunits activate

other SWI/SNF subunits. It is also unclear whether subunits of the two assemblies in *C.*

elegans – BAF and PBAF – stabilize the core protein subunits or vice-versa. Therefore,

we used our endogenously labeled GFP-SWI/SNF strains to ask whether transcriptional

knockdown of individual subunits of the core, BAF, or PBAF induce changes in protein

expression of other subunits at the time of AC invasion (**Fig S5**). First, to determine

whether representative subunits of the SWI/SNF assemblies promote or stabilize the

ATPase of the complex, we treated *GFP::swsn-4* animals with either *swsn-8(RNAi)* or

pbrm-1(RNAi) (**Fig S5 A**). Quantification of fluorescence expression in AC nuclei of *swsn-*

8(RNAi) treated animals at the P6.p 4 cell stage revealed significantly lower GFP::SWSN-

4 levels relative to the control group (34% GFP::SWSN-4 depletion; **Fig S5 A, D**). RNAi

knockdown of the PBAF subunit *pbrm-1* also resulted in a significant but weaker loss of

ATPase expression in the AC (11% GFP::SWSN-4 depletion; **Fig S5 A, D**). These results

suggest that individual subunits of either SWI/SNF assembly may contribute to the protein

stability and/or expression of the SWI/SNF ATPase in the *C. elegans* AC, with the BAF

complex playing a potentially dominant activating role with respect to the ATPase.

Next, we treated animals containing either the *swsn-8* or *pbrm-1* endogenous

GFP-reporters with enhanced RNAi to knockdown the expression of the SWI/SNF

ATPase or the representative subunit of the alternative SWI/SNF assembly. Interestingly,

while unaffected by knockdown of the PBAF assembly subunit *pbrm-1,* RNAi knockdown

of the ATPase *swsn-4* resulted in a 42% increase in the expression of SWSN-8::GFP in

the AC (**Fig S5 B, D**). Finally, relative to the expression of the endogenous PBAF subunit

in the ACs of animals treated with the empty vector control RNAi, AC nuclei of PBRM-

1::eGFP animals treated with *swsn-4(RNAi)* had significantly lower levels of protein

expression (38% PBRM-1::eGFP depletion), whereas ACs in *swsn-8(RNAi)* treated

animals expressed 13% more PBRM-1::eGFP (**Fig S5 C-D**).

Since knockdown of either *swsn-4* or *swsn-8* subunits resulted in two distinct

phenotypes – individual animals with single non-invasive ACs and animals with multiple

non-invasive cells expressing the AC-reporter - we next sought to determine whether

these two phenotypes were distinct with respect to SWI/SNF subunit expression. To do

this, we binned data from the intracomplex RNAi experimental series (above) into the two

non-invasive phenotypes and compared the fluorescence expression levels of the

endogenous proteins within SWI/SNF RNAi conditions. Given the infrequency of the

multi-AC phenotype, statistical comparisons were necessarily limited to treatments in

which the population of animals contained at least 10 multi non-invasive ACs. Treatment

of SWSN-8::GFP with *swsn-4(RNAi)* resulted in a total of 24 multi non-invasive ACs (53

ACs total; n=41 animals) and no significant difference was detected in SWSN-8::GFP

expression between the nuclei of the single non-invasive AC phenotype and the multi

non-invasive AC phenotype groups (**Fig S5 E**). The second statistical comparison was

made between the two phenotypes in PBRM-1::eGFP animals treated with *swsn-8(RNAi)*

(**Fig S5 E**), in which 14 multi non-invasive ACs were detected (51 ACs total; n=42

animals). Quantification of endogenous PBRM-1::eGFP fluorescence expression in this

condition revealed a slight (12%) increase in expression of the PBAF subunit in the nuclei

of ACs of the multi non-invasive phenotype group compared to the single non-invasive

phenotype (**Fig S5 E**).

Based on these results, a tentative model for epistatic interactions between the

SWI/SNF ATPase, BAF, and PBAF assembly subunits can be composed for the AC (**Fig

S5 F**). Our data indicate that some degree of SWI/SNF intra- and inter-complex regulation

occurs in the AC. We find that the most significant aspect of SWI/SNF intra-complex

regulation is exercised by the ATPase on the assembly specific subunits, where *swsn-4*

knockdown results in a significant increase in BAF/SWSN-8 and a significant decrease

PBAF/PBRM-1. SWI/SNF intercomplex regulation appears to be weaker in the AC as

knockdown of *pbrm-1* does not affect SWSN-8::GFP expression, and knockdown of

swsn-8 results in a slight increase in PBRM-1::GFP expression.

The SWI/SNF ATPase SWSN-4 provides dose-dependent regulation of AC invasion

The degree to which the SWI/SNF complex contributes to tumorigenesis in clinical

settings has been linked to the dose of functional SWI/SNF ATPase in precancerous and

transformed cells (32,34,65). Previous work in *C. elegans* has demonstrated a similar

dose dependent relationship between SWI/SNF and cell cycle control (35). To determine

whether the phenotypic dosage sensitivity seen in cancer and *C. elegans* mesodermal

(M) cell development is also characteristic of SWI/SNF in the promotion of AC invasion

(30), we modulated expression of GFP::SWSN-4 using a combination of RNAi-mediated

knockdown and AC-specific GFP-targeting nanobody technology.

Though RNAi treatment targeting the *swsn-4* subunit in the endogenously-tagged

strain resulted in significant knockdown of fluorescence expression of GFP::SWSN-4 in

the AC, loss of expression was noted in many other tissues in treated animals, including

the 1° VPCs, which contribute to AC invasion non-autonomously (14,66) (**Fig S4**)*.* Thus,

to limit loss of expression to the AC, we used an anti-GFP nanobody fused to a SOCS-

box containing a ubiquitin ligase adaptor, driven with tissue-specific promoters to achieve

lineage-restricted protein depletion (30) (**Fig 3**). To follow the expression of the anti-GFP

nanobody transgenes, we also included a fluorescent histone label separated from the

anti-GFP nanobody sequence by the p2a viral self-cleaving peptide (*ACp::antiGFP-*

nanobody::p2a::His-58::mCherry). We generated two anti-GFP nanobody constructs,

using conserved *cis*-regulatory elements from the *cdh-3* and *egl-43* promoters

(22,24,39,67,68) and introduced them into a strain containing the endogenous GFP-

tagged allele of *swsn-4* as well as background AC and BM reporters (**Fig 3 B-C**). The

Figure 3. AC invasion and cell cycle arrest depend on dosage of SWI/SNF ATPase. (A-E) Representative fluorescence images depicting expression of BM marker (*laminin::GFP*) and endogenous GFP::SWSN-4 (left), AC reporter (*cdh-3p::mCherry::moeABD*, middle), and fluorescence overlay (right) across experimental treatments. White arrowheads indicate AC(s), yellow arrowheads indicate boundaries of breach in BM, and black brackets indicate 1° VPCs. In cases where multiple cells expressed the AC reporter in the same animal, each is indicated with a single white arrowhead. Asterisk indicates anti-GFP nanobody expression in neighboring VU cell. **(F)** Quantification of mean gray values (M.G.V.) of endogenous GFP::SWSN-4 in ACs in control animals (empty vector) and across all experimental treatments normalized to mean fluorescent expression in wildtype animals (n≥40 animals per treatment, p values for Student's *t*-test comparing expression of successive knockdown are displayed on the figure). In this and all other figures, open circles and error bars denote mean±standard deviation (s.d.). n.s. not significant. **(G)** Stacked bar chart showing quantification of AC invasion defects corresponding to each treatment, binned by AC phenotype (n≥40 animals per condition; p values for Fisher's exact test comparing phenotypes of successive knockdown strategies are displayed above compared groups). Grey brackets indicate statistical significance between invasion total in each condition compared to invasion defect total. Black brackets indicate statistical significance between incidences of invasion defects with multiple ACs compared to incidences of invasion defects with single ACs. n.s. not significant.

cdh-3-driven nanobody transgene (*cdh-3p::antiGFP-nanobody::p2a::his-58::mCherry*)

resulted in a weak reduction of GFP::SWSN-4 levels with no significant difference in

fluorescence expression in the AC compared to wildtype animals (6% depletion; n=80

animals; **Fig 3 F**); however, consistent with the wildtype expression of the *cdh-3* promoter

(22,39), it expressed specifically in the AC and resulted in defective AC invasion,

suggesting partial loss of function (20.6% AC invasion defect; n=102; **Fig 3 B, G**). The

egl-43p::antiGFP-nanobody   transgene   (*egl-43p::antiGFP-nanobody::p2a::His-*

58::mCherry) expression pattern was also consistent with the wildtype expression

characterized in previous work (22,67–69), as indicated by nuclear expression of HIS-

58::mCherry in the AC and in the neighboring ventral uterine and dorsal uterine (VU/DU)

cells (**Fig 3 C**; asterisk denotes HIS-58::mCherry expression in a non-AC ventral uterine

cell) (22,39,67). Importantly, as the AC invades independent of VU/DU cells (14), anti-

GFP expression in these tissues should not affect AC invasion. Similar to animals treated

with *swsn-4(RNAi)* (**Fig 3 D**), *egl-43p::antiGFP-nanobody*-mediated protein depletion of

GFP::SWSN-4 resulted in a significant loss of fluorescence expression in the AC (71%

GFP depletion; n=80 animals; **Fig 3 C, F**) as well as a penetrant loss of invasion and

incidence of individual animals with multiple uterine cells that were in contact with the

ventral BM and expressed the AC reporter (88.2% AC invasion defect, 2.9% multiple AC

phenotype; n=101 animals; **Fig 3 G**). These results support our uterine-specific SWI/SNF

RNAi results and provide strong evidence for a cell-autonomous role for the SWI/SNF

complex in promoting cell invasion and cell cycle arrest.

To further deplete *swsn-4* expression in the AC, we treated transgenic *egl-*

43p::antiGFP-nanobody animals with *swsn-4(RNAi)* (**Fig 3 E**). Strikingly, in this

combination knockdown strategy, 100% of AC invasion was lost and the frequency of

multiple cells expressing the AC specification reporter drastically increased relative to

treatment with *swsn-4(RNAi)* or the *egl-43*-driven anti-GFP nanobody conditions alone

(83% multiple AC phenotype; n=41 animals; **Fig 3 G**). Together, these results

demonstrate a phenotypic spectrum that corresponds to successive loss of *swsn-4* in the

AC: whereas moderate loss of the ATPase results in single non-invasive ACs in animals

containing *egl-43p::antiGFP-nanobody*, strong loss of expression in the *egl-*

43p::antiGFP-nanobody background or following treatment with *swsn-4(RNAi)* results in

animals with both single and multiple non-invasive ACs; in the strongest knockdown

condition – *egl-43p::antiGFP-nanobody* animals treated with *swsn-4(RNAi)* - multiple

non-invasive ACs were present per animal with near complete penetrance. Though the

combination of *swsn-4(RNAi)* and antiGFP-nanobody-mediated depletion resulted in

robust loss of expression of the core ATPase of the SWI/SNF complex, the fluorescence

expression was not significantly different than treatment with *swsn-4(RNAi)* alone (93%

vs. 92% GFP depletion, respectively; n≥41 animals for each treatment; **Fig 3 G**); therefore

we theorize that the fluorescence values were beyond our threshold ability to quantify

based on the fluorescence detection limits of our imaging system. Altogether, our data

demonstrated that in the AC, the ATPase of the SWI/SNF complex contributes to invasion

cell-autonomously and in a dose-dependent manner: moderate loss of expression

produced non-invasive single ACs while extreme loss of expression led to a non-invasive

hyperproliferative state.

Improved *swsn-4(RNAi)* vector is sufficient to recapitulate null phenotype in M

lineage

A recent study focusing on cell cycle control of SWI/SNF throughout *C. elegans*

muscle and epithelial differentiation demonstrated tissue and lineage-specific phenotypes

following weak or strong loss of core SWI/SNF subunits (32). Within the M lineage that

gives rise to posterior body wall muscles (BWMs), coelomocytes (CCs), and reproductive

muscles or sex myoblast (SMs) descendants, different cell types responded differently to

loss of SWI/SNF. In the BWM, strong loss of SWI/SNF resulted in hyperproliferation, like

the phenotype we detect in the AC. The opposite is true in the SM lineage, where modest

knockdown of *swsn-4* resulted in hyperproliferation while complete loss of *swsn-4*

expression resulted in a null phenotype where SMs failed to divide and arrest in S phase

(35). We next sought to validate the strength of our enhanced *swsn-4(RNAi)* vector by

examining proliferative state in the SMs. To accomplish this, we treated animals

containing a lineage-restricted cyclin-dependent kinase (CDK) activity sensor (*unc-*

62p::DHB::2xmKate2) with *swsn-4(RNAi)* (**Fig S6 A**). In this genetic background, we

determined the number (**Fig S6 B**) and cell cycle state (**Fig S6 C**) of SM cells at a time

when the majority of SMs in control animals had finished cycling and subsequently

differentiated (late P6.p 8 cell stage; 16 SM cell stage). Animals treated with *swsn-*

4(RNAi) had significantly fewer SM cells than controls (mean SMs/animals = 5; n=31

animals; **Fig S6 B**) with many instances of SMs that failed to enter a single round of cell

division (n=20 single SMs out of 43 animals). Interestingly, 28% (12/43) of animals treated

with *swsn-4(RNAi)* were absent of SMs on either the left or the right side, whereas 100%

(30/30) control animals had SMs on both sides, which may indicate a defect in

specification, early cell division, and/or migration of SMs. To quantify cell cycle state, we

measured localization of an SM-specific CDK sensor, which uses a fragment of

mammalian DNA Helicase B (DHB) fused to two copies of mKate2 (36,70). In cells with

low CDK activity that are quiescent or post-mitotic, the ratiometric CDK sensor is strongly

nuclear localized (36,68,70). In cycling cells with increasing CDK activity, the CDK sensor

progressively translocates from the nucleus to the cytosoplasm, with a ratio approaching

1.0 in S phase and >1 in cells in G_2 (36). Thus, the cytoplasmic:nuclear (C/N) ratio of

DHB::2xmKate2 can serve as a proxy to identify cell cycle state. By the time the majority

of SMs in the control condition were differentiating and arrested in a G_0 cell cycle state

(mean C/N ratio=0.320; n=90 SMs; **Fig S6 C**), many animals treated with *swsn-4(RNAi)*

had single SMs that failed to divide and a mean DHB C/N ratio indicative of a long pause

or arrest in S phase (36) (Avg. C/N ratio = 0.803; n=20 SMs; **Fig S6 C**). Together, these

results suggested that the strength of our enhanced *swsn-4(RNAi)* targeting vector is

sufficient to recapitulate the *swsn-4* null condition in the SM lineage, as we detected both

the hypoproliferative phenotype and S-phase arrest that was observed using a lineage-

restricted catalytically inactive SWI/SNF ATPase (35).

The BAF assembly contributes to AC invasion via regulation of G_0 cell cycle arrest

Having established that strong depletion of the SWI/SNF complex results in a fully

penetrant defect in AC invasion with a high percentage of individual animals possessing

multiple non-invasive ACs (**Fig 3**), we next investigated whether the extra ACs observed

were the consequence of inappropriate AC proliferation (22,68). To determine whether

the SWI/SNF complex is required for G_0/G_1 cell cycle arrest in the AC, we quantified CDK

activity in the AC using a ubiquitously expressed *rps-27p::*DHB::GFP transgene paired

with AC (*cdh-3p::mCherry::moeABD*) and BM (*laminin::GFP*) reporters in live animals

following RNAi-mediated knockdown of SWI/SNF core (*swsn-4*), BAF (*swsn-8*), and

PBAF (*pbrm-1*) subunits (**Fig 4**). In wild-type invasive ACs, we observed strong nuclear

localization of the CDK sensor and quantified a cytoplasmic/nuclear (C/N) ratio indicative

of G_0/G_1 arrest (mean C/N ratio: 0.226+/-0.075, n=67 animals) (**Fig 4 A, F**). In order to

distinguish whether the wild-type AC C/N ratio is actually indicative of G_0 rather than G_1

arrest, we quantified the CDK activity in the neighboring uterine Pi cells at the P6.p 8 cell

1° VPC stage following their terminal division to establish a G_0 reference point (71,72)

(mean C/N ratio: 0.206+/-0.078, n=30 animals) (**Fig 4 E-F**). We found no significant

difference between the CDK activity of terminal Pi cells and wild-type invading ACs,

suggesting that the wild-type AC exists in a CDK^{low} G_0, pro-invasive state (**Fig 4 A, E-F**).

In animals treated with *pbrm-1(RNAi)*, the CDK sensor also localized principally in the

nucleus of ACs that failed to invade (mean C/N ratio: 0.157+/-0.063, n=41 animals) and

only a single non-invasive AC was observed per animal (**Fig 4 D, F**). In contrast, following

treatment with *swsn-8*(*RNAi*) the majority of ACs that failed to invade the BM were

in the G_1/S phases of the cell cycle (mean C/N ratio: 0.636+/-0.204, n=21 animals) (**Fig**

4 C, F). Finally, like the *swsn-8(RNAi)* condition, loss of expression of the core ATPase

of the SWI/SNF complex through treatment with *swsn-4*(*RNAi*) resulted in a broad range

of C/N ratios (C/N ratio min: 0.240, C/N ratio max: 1.140, mean C/N ratio: 0.566+/-0.205;

n=40 animals) in animals with single or multiple non-invasive ACs (**Fig 4 B, F**).

Interestingly, the *swsn-4(RNAi)* treatment resulted in a higher proportion of non-invasive

G_0 phase (C/N ratio < 0.3) ACs (14%, n=48 cells) than were present in the *swsn-8(RNAi)*

treated population (8%, n=25) (**Fig 4 B, E**). This finding reemphasizes the dependence

of both SWI/SNF BAF and PBAF assemblies on the core ATPase of the complex, as the

distribution of cell cycle states in ACs following *swsn-4(RNAi)* treatment represents both

Figure 4. CDK sensor reveals SWI/SNF contribution to G_0 arrest in the AC. Micrographs depicting DIC (left), AC (*cdh-3p::mCherry::moeABD*, center-left), DIC overlay (center-right), and DHB-based CDK activity sensor (right) in empty vector **(A)** and following treatment with SWI/SNF RNAi targeting subunits of the core **(***swsn-4*, **B)**, BAF **(***swsn-8*, **C)** and PBAF **(***pbrm-1*, **D)** assemblies. White arrowheads indicate AC(s), yellow arrowheads indicate boundaries of breach in BM, and white brackets indicate 1° VPCs. In cases where treatment resulted in multiple cells expressing the AC reporter in the same animal, representative images of both single (1AC) and mitotic (2+ACs) phenotypes are given, and each AC is indicated with a single white arrowhead. Quantification of the cytoplasmic:nuclear (C/N) ratio of DHB::GFP in ACs (white dotted outline) is listed in the bottom left of each panel. Mitotic ACs are numbered, and C/N ratios are provided for each **(B-C)**. White arrowheads indicate AC(s), yellow arrowheads indicate boundaries of breach in BM, and white brackets indicate 1° VPCs. White arrow in C indicates an AC that is out of the focal plane. **(E)** Representative single z-plane micrographs of the vulva at the P6.p 8 cell stage (left, z=1) and the terminal Pi cells (middle, z=9) in DIC, and DHB-based CDK activity sensor in Pi cells (right). Quantification of the C/N ratio of DHB::GFP in three of four Pi cells (white dotted outline) that are in the plane of the image is listed in the bottom left. **(F)** Quantification of C/N DHB::GFP ratios for wild-type terminally divided Pi cells and all ACs in empty vector control and each RNAi treatment (n≥30 animals per treatment). Statistical comparison was made between the mean C/N ratio of ACs in control (empty vector) compared to control (empty vector) Pi cells using Student's *t*-test (n≥30 for each stage and subunit; p values are displayed above compared groups). Mean C/N ratio is represented by colored open circles and correspond to numbers above the data. Gradient scale depicts cell cycle state as determined by quantification of each Pi cell or AC in all treatments (n≥30 animals per treatment), with dark/black depicting G_0 and lighter/magenta depicting G_2 cell cycle states. Dashed white line on gradient scale bar (right) corresponds to boundaries between G_0 and G_1 phases. Colored open circles on the gradient scale correspond to the mean C/N ratio in each of the same color. n.s. not significant.

482   the cell cycle-dependent and cell cycle-independent phenotypes seen in ACs deficient in

483   the *swsn-8* and *pbrm-1* subunits, respectively

Forced G$_0$ arrest through ectopic CKI-1 rescues invasive potential in BAF-deficient

but not PBAF-deficient ACs

We have previously proposed and characterized a dichotomy that exists between

invasion and proliferation in the AC (22,24). As evidence of this, loss of two of the three

TFs that function in a cell cycle-dependent manner to maintain the AC in a cell cycle-

arrested state (*nhr-67*/Tlx and *hlh-2*/Daughterless) can be rescued through induced

expression of a cyclin dependent kinase inhibitor, *cki-1* (p21/p27) (22). These results

suggest that, at least in some cases, TF activity can be bypassed completely to promote

AC invasion by maintaining G$_0$ arrest through direct cell cycle manipulation. To determine

the extent to which the BAF assembly contributes to AC invasion through regulation of

cell cycle arrest, we used a heat-shock inducible transgene to ectopically express CKI-

1::mTagBFP2 in SWI/SNF-deficient ACs (**Fig 5**). Since the heat shock inducible

transgene is ubiquitous and expresses variably between different animals and different

tissues within an individual animal, we limited our analysis to animals with ACs with

obvious mTagBFP2 fluorescence expression. While forced arrest in G$_0$ was insufficient to

significantly rescue AC invasion in animals treated with *swsn-4(RNAi)* (**Fig 5 B, B', E**) or

pbrm-1(RNAi) (**Fig 5 D, D', E**), ectopic *cki-1* (CKI-1::mTagBFP2) expression in the AC

significantly rescued cellular invasion in animals treated with *swsn-8(RNAi)* (**Fig 5 C, C',**

E). Strikingly, in 85.7% (6/7) of cases where ACs had proliferated prior to ectopic CKI-1

expression, forced G$_0$ arrest led to multiple ACs breaching the BM (**Fig 5 F**), a phenotype

we have reported previously using CKI-1 overexpression paired with loss of NHR-67. This

demonstrated that mitotic ACs maintain the capacity to invade if they are re-arrested into

Figure 5. BAF depletion is rescued by G_0 arrest. Representative micrographs depicting DIC (left), BM (*laminin::GFP*, center-left), AC (*cdh-3p::mCherry::moeABD,* center-right), and CKI-1 (*hsp::CKI-1::mTagBFP2*) expression in empty vector control **(A-A')** and treatment with SWI/SNF RNAi under standard conditions **(A-D)** and following heat shock induction of CKI-1 **(A'-D')**. CKI-1 images have been inverted for ease of visualization. White arrowheads indicate AC(s), yellow arrowheads indicate boundaries of breach in BM, and white brackets indicate 1° VPCs. **(E)** Stacked bar chart showing percentage of AC invasion defects corresponding to each RNAi treatment under standard growth conditions (control) and following heat shock induction of CKI-1 (+CKI-1), binned by AC phenotype (n≥30 animals per condition; Fisher's exact test compared CKI-1(+) animals with control, non-heat shocked animals; p value is displayed above compared groups). n.s. not significant. **(F)** Representative micrographs of invasive group of *swsn-8* deficient ACs following induction of G_0/G_1 arrest. DIC (top-left), BM (bottom-left), CKI-1 expression (top-right), AC reporter (bottom-right). Max intensity z-projection of AC and BM reporter channels (bottom). Large breach in BM is indicated by black arrow.

a G_0 state (24). To corroborate our CKI-1 heat shock data, we used an AC-specific CKI-

1 transgene (*cdh-3p::CKI-1::GFP*) to induce G_0 cell cycle arrest in *swsn-4-* and *swsn-8-*

depleted ACs (**Fig S6**). Similar to the heat shock results, lineage-restricted expression of

CKI-1::GFP failed to rescue invasion in animals deficient in *swsn-4* (**Fig S7 A-B**).

However, transgenic *cdh-3p::CKI-1::GFP* animals treated with *swsn-8(RNAi),* had

invasion defects significantly lower than control animals treated with *swsn-8(RNAi)* which

lacked the G_0 rescue transgene (**Fig S7 B**). Altogether, these data corroborate our DHB-

based CDK sensor data (**Fig 4**), suggesting that the SWI/SNF assemblies differentially

contribute to AC invasion with BAF specifically required for G_0 cell cycle arrest.

SWI/SNF chromatin remodeling promotes the invasive GRN in the AC

Previous work has demonstrated that the gene regulatory network (GRN) that

promotes AC invasion consists of both cell cycle-dependent and cell cycle-independent

TF subcircuits (22,68) (**Fig 1 B**). In the cell cycle-dependent subcircuit of the TF-GRN,

egl-43 (EVI1/MEL), *hlh-2* (E/Daughterless), and *nhr-67* (TLX/Tailless) cooperate in a type

1 coherent feed-forward loop that is reinforced via positive feedback to retain the AC in a

post-mitotic, invasive state (22,68). The cell cycle-independent subcircuit of the AC TF-

GRN is governed by the *fos-1* (FOS) TF with feedback from both *egl-43* and *hlh-2* (22).

Since transcriptional knockdown of SWI/SNF ATPase results in both single and mitotic

non-invasive AC phenotypes, we treated endogenously GFP-labeled strains for each TF

in the GRN with *swsn-4(RNAi)* to determine whether SWI/SNF chromatin remodeling

contributes to the regulation of either or both AC GRN subcircuits (**Fig 6**). In the cell cycle-

dependent subcircuit, knockdown of the SWI/SNF ATPase resulted in significant loss of

protein expression of EGL-43::GFP and NHR-67::GFP in the AC (39% and 26% GFP

depletion, respectively; n≥41 animals; **Fig 6 A,C,E**). No significant difference was

detected in the mean fluorescence expression of GFP::HLH-2 fusion protein in the AC

upon knockdown of *swsn-4,* however the range of expression was broad following *swsn-*

4(RNAi) treatment (~2% GFP increase; n≥50 animals; **Fig 6 B, E**). In the cell cycle-

Figure 6. SWI/SNF regulates endogenously tagged TFs in the AC invasion GRN. Fluorescent micrographs depicting BM (*lam::GFP*) and AC (*cdh-3p::mCherry::moeABD*) expression of endogenously tagged TFs of the cell cycle-dependent subcircuit (*egl-43::GFP::egl-43* **(A)**, *GFP::hlh-2* **(B),** and *nhr-67::GFP* **(C)**) and cell cycle-independent subcircuit (*GFP::fos-1a* **(D)**) of the AC GRN in animals treated with empty vector control (left) or *swsn-4(RNAi)* (right). White arrowheads indicate AC(s), yellow arrowheads indicate boundaries of breach in BM. **(E)** Quantification of fluorescent expression of each TF::GFP in ACs of control animals and animals treated with *swsn-4(RNAi)*. Statistical comparisons were made between the expression of each TF subunit in the AC in control and RNAi-treated animals using Student's *t*-test (n≥30 for each condition; p values are displayed above black brackets). n.s. not significant.

independent subcircuit, loss of the SWI/SNF complex following treatment of *fos-1::GFP*

animals with *swsn-4(RNAi)* resulted in a more moderate depletion of expression in the

AC (11% GFP depletion; n≥50 animals; **Fig 6 D**). These results suggest that the SWI/SNF

complex broadly remodels chromatin to promote both subcircuits of the pro-invasive AC

GRN, though we are unable to determine whether the complex does so directly in

regulatory regions of the lineage-specific pro-invasive TFs or in regulatory regions of

genes that contribute to the regulation of the AC GRN.

The PBAF assembly regulates AC contact with underlying BM

Previous investigations into SWI/SNF have demonstrated divergent roles for the PBAF

assembly in cell cycle regulation. In yeast, Remodeling the Structure of Chromatin (RSC),

the homologous complex to PBAF, is required for progression through mitosis (73,74). In

Drosophila, the homologous complex PBAP does not appear to be required for mitotic

progression; rather, cycling and G_2/M transition is solely regulated by the BAF/BAP

assembly (51). In the *C. elegans* M lineage, RNAi-mediated loss of BAF subunits results

in hyperproliferation of the developing tissue, whereas knockdown of PBAF subunits has

little effect on cell cycle control (35). Similarly, in this study, RNAi-mediated loss of PBAF

subunits *pbrm-1*, *swsn-7*, or *swsn-9* resulted exclusively in single non-invasive cells

expressing the AC reporter. However, given that the enhanced *pbrm-1(RNAi)* resulted in

much weaker endogenous protein knockdown than the enhanced RNAis targeting either

the SWI/SNF ATPase (*swsn-4*) or BAF assembly subunit (*swsn-8*) in the AC (**Fig S4**),

and the dose-dependent phenotype following loss of the core ATPase (**Fig 3**), it is

possible that we failed to observe the mitotic non-invasive AC phenotype due to

insufficient PBAF subunit knockdown. To address this, we next asked whether *strong*

loss of PBAF subunit expression contributes to the mitotic non-invasive AC phenotype.

To accomplish this, we used an auxin inducible degron (AID)-RNAi combination

knockdown strategy (75,76). We generated a strain with *pbrm-1* endogenously labeled

with mNeonGreen and an auxin inducible degron (AID) (*pbrm-1::mNG::AID*) in a genetic

background containing AC (*cdh-3p::mCherry::moeABD*) and BM (*laminin::GFP*)

reporters. We then quantified fluorescence expression in the AC in this strain. When

grown under standard conditions, 6.45% of the ACs had not invaded the BM by the P6.p

4 cell stage, suggesting a partial loss of function of *pbrm-1* (n=30; **Fig 7 A, E-F**). This

partial loss of function phenotype is likely due to the insertion of the mNG::AID tag into

the genomic locus, causing a putative hypomorphic allele. Next, we introduced a

ubiquitous, mRuby-labeled TIR1 transgene (*eft-3p::TIR1::mRuby*) into the animals and

assessed AC invasion under standard conditions (aux(-)) or in the presence of the auxin

hormone (aux(+); **Fig 7 B**). We observed no statistically significant difference in the

fluorescence expression of PBRM-1::mNG::AID protein in the AC, nor did we observe

any differences in AC invasion defects between the strains with and without the TIR1

transgene when grown on aux(-) media (TIR1+: 3% depletion, 16.67% invasion defect;

n=30; **Fig 7 E-F**). However, in both conditions, some ACs that invaded seemed to do so

only partially, as most of the ventral epidermal BM remained intact beneath ACs (**Fig 7**

B, black arrowhead). In the aux(+) condition, there was a significant reduction in PBRM-

1::mNG::AID protein level in the AC of animals containing the TIR1 transgene relative to

the same strain grown in the aux(-) condition or the strain without the TIR1 transgene

(49% and 51% depletion, respectively; n=30; **Fig 7 E**); however, there was no significant

difference in the penetrance of AC invasion defects (16.67% invasion defect; n=30; **Fig 7**

F). Like our previous results with *pbrm-1(RNAi)* treated animals, we observed no extra

cells expressing the AC reporter following loss of expression of PBAF in the AC using the

AID system.

Next, we treated *pbrm-1::mNG::AID* animals containing ubiquitous TIR1 with

pbrm-1(RNAi) in both aux(-) and aux(+) conditions. As expected, treatment of the *pbrm-*

1::mNG::AID strain with *pbrm-1(RNAi)* resulted in very low expression of the subunit in

the AC even in the absence of auxin and there was no significant difference in expression

between the Aux(-) and Aux(+) conditions (**Fig 7 C,F**). Interestingly, there was also no

significant difference in the penetrance of AC invasion defects between the *pbrm-*

1::mNG::AID strain treated with control compared to the strain treated with *pbrm-1(RNAi)*

in the presence of auxin (**Fig 7 G**). Since the combination treatment of a hypomorphic

Figure 7. PBAF promotes AC contact to the BM. Representative micrographs of BM (*lam::GFP*) and endogenous *pbrm-1::mNG::AID* (left), AC (*cdh-3p::mCherry::moeABD*, center), and fluorescent overlays (right) of animals lacking **(A)** or possessing **(B-E)** ubiquitous TIR1 expression treated with empty vector control **(B)** or RNAi targeting PBAF subunits in the absence Aux(-) (top) or presence Aux(+) (bottom) of media containing the auxin hormone **(C-E)**. PBRM-1::mNG::AID images have been inverted for ease of visualization. **(F)** Quantification of fluorescence expression (M.G.V) of PBRM-1::mNG::AID in ACs of animals in each condition (N≥30 animals in each treatment; p values for Fisher's exact test comparing strains containing TIR1 to the TIR1(-) strain, and comparing strains containing TIR1 in the Aux(-) to the Aux(+) condition, are displayed above compared groups). **(G)** Stacked bar chart showing percentage of AC invasion defects corresponding to each treatment, binned by AC phenotype (N≥30 animals per condition; Fisher's exact test compared penetrance of AC invasion defects between indicated conditions). Black brackets indicate statistical significance between invasion total in each condition compared to invasion defect total. **(H)** Max intensity z-projection of AC and BM reporter channels depicting a detached AC phenotype in *swsn-7*-deficient ACs in the Aux(+) condition. BM (left), AC (center), fluorescence overlay (right). Asterisk indicates polarized F-actin driven protrusion extending ventrally.

pbrm-1 allele, Auxin-AID-mediated depletion of endogenous PBRM-1::mNG::AID, and

pbrm-1(RNAi) does not result in a significant increase in AC invasion defects or non-

invasive mitotic ACs, these results suggest that, unlike the dose-dependent contribution

to invasion of *swsn-4*, the *pbrm-1* strong knockdown or null phenotype may be only

partial/incomplete loss of AC invasion.

Since the PBAF assembly in *C. elegans* consists of several subunits, *pbrm-1*

(PBRM1), *swsn-7* (ARID2), and *swsn-9* (BRD7/BRD9), we next investigated whether

combinatorial knockdown of PBAF subunits would enhance the penetrance of AC

invasion defects or result in the mitotic non-invasive AC phenotype. In the absence of

auxin, there was no significant difference in PBRM-1::mNG::AID expression in the AC of

animals treated with *swsn-7(RNAi)* compared to animals treated with empty vector control

(n=30; **Fig 7 E**), however there was a significant increase in loss of AC invasion (50%

invasion defect; n=30; **Fig 7 F**). Strikingly, in one case, the AC was completely detached

from the BM, as we detected no AC membrane protrusions (*cdh-3p::mCherry::moeABD*)

in contact with the ventral surface of the gonad (**Fig 7 F**). Animals treated with *swsn-*

7(RNAi) and aux(+) had significantly lower expression of PBRM-1::mNG::AID in the AC

when compared to animals treated with *swsn-7(RNAi)* in the aux(-) condition (49%

depletion; n=30; **Fig 7 E**). While no significant difference was seen in loss of AC invasion

in aux(+) (48.39% AC invasion defect), 16% (5/31) of animals in this treatment had ACs

entirely detached from the ventral BM (n=31; **Fig 7 F-G**). In contrast to treatment with

swsn-7(RNAi), in the *swsn-9(RNAi)* aux(-) condition, PBRM-1::mNG::AID expression in

ACs was significantly lower than that in the ACs of animals treated with empty vector

control aux(-) (39% depletion; n=30; **Fig 7 E**). It is unclear why transcriptional knockdown

of *swsn-9* specifically results in a decrease in PBRM-1 protein expression in the AC and

we theorize this may be the result of a potential stabilizing interaction between the SWSN-

9 and PBRM-1 proteins. We did detect a significant decrease in expression of PBRM-

1::mN G::AID in ACs in *swsn-9(RNAi)* aux(+) compared to the *swsn-9(RNAi)* aux(-)

condition (19% depletion; n=30; **Fig 7 E**), however, we saw no statistically significant

difference in penetrance of AC invasion defects between the two conditions (30% vs.

43%; n=30; **Fig 7 F**). We also noted one animal with a detached AC in the *swsn-9(RNAi)*

aux (-) condition and zero in the aux(+) condition (**Fig 7 F**). Importantly, we only observed

one AC per animal across all combinatorial treatments, supporting the hypothesis that the

PBAF assembly does not contribute to G_0 cell cycle arrest in the AC.

Detached ACs in both the *swsn-7(RNAi)* and *swsn-9(RNAi)* AID combination

knockdown conditions suggest that the PBAF assembly regulates AC contact with the

ventral epidermal BM. A previous study has shown that AC attachment is regulated by

the *fos-1/egl-43* cell cycle-independent subcircuit of the AC GRN via regulation of

lamellipodin/*mig-10b* and non-autonomously via netrin/*unc-6* signaling (77). ACs deficient

in components of this pathway are attached to the ventral epidermal BM when specified

and gradually lose contact over time, with peak loss of contact occurring at the time of AC

invasion at the P6.p 4-cell stage (77). In order to determine whether the PBAF assembly

remodels chromatin to promote activation of this subcircuit of the AC GRN, we treated

endogenously tagged *fos-1::GFP* (22) animals with *pbrm-1(RNAi)* and quantified

fluorescence expression in ACs that displayed invasion defects (**Fig S8 A-B**). Animals

treated with *pbrm-1(RNAi)* had a modest but statistically significant loss of FOS-1::GFP

protein levels in non-invasive ACs (34% depletion; n=20; **Fig S8 B**), suggesting that the

PBAF assembly partially regulates the *fos*-dependent pathway that mediates attachment

to the underlying BM.

Since depletion of the PBAF assembly resulted in moderate loss of FOS-1::GFP

in the AC, we next examined functional interactions between FOS-1 and PBRM-1. Given

that the PBRM-1::mNG::AID allele was slightly hypomorphic, with ~17% invasion defects

in backgrounds with TIR1, we used the strain containing TIR1 as a sensitized

background. We found that even without the addition of auxin, co-depletion with *fos-*

1(RNAi) resulted in almost complete loss of AC invasion (96.77% invasion defect; n=31;

Fig S8 C-D). Finally, we examined whether RNAi-mediated depletion of *pbrm-1* is

synergistic with loss of downstream targets of FOS-1, the matrix metalloproteinases

(MMPs). Previously, it has been shown that animals harboring null mutations for five of

the six MMPs encoded in the *C. elegans* genome (*zmp-1,-3,-4,-5* and *-6*), show delayed

AC invasion (21). RNAi depletion of *pbrm-1* in quintuple MMP mutants significantly and

synergistically enhanced late invasion defects (scored at the P6.p 8-cell stage) in this

background (24.24% invasion defect; n=33; **Fig S8 E-F**) as compared to loss of either

pbrm-1 (3.8%; n=52) or MMPs (0%; n=35) alone. Together, these results suggest that the

PBAF assembly functions synergistically with FOS-1 to regulate AC invasion.

DISCUSSION

A tissue-specific CRF RNAi screen identified genes critical for cellular invasion

Previous work in the *C. elegans* AC and in cancer cell invasion has emphasized

the necessity for dynamic chromatin states and chromatin regulating factors in the

promotion of cellular invasion (24,79–83). In this study, we used the *Caenorhabditis*

elegans AC invasion model as a single cell, *in vivo* system to identify a suite of CRFs that

contribute to the process of cellular invasion. We performed a tissue-specific RNAi

feeding screen to assess 269 genes implicated in chromatin binding, chromatin

remodeling complexes, or histone modification. We do not claim that genes that we failed

to identify as regulators of cellular invasion in the screen are unimportant for the process;

however, RNAi-mediated loss of the majority of CRF genes in the screen did result in

some penetrance of AC invasion defects (**Table S1**). This finding was expected, as many

of the genes we screened are global regulators of the genome and broadly contribute to

various cell biological processes. We extracted a list of the most penetrant regulators of

cell invasion from the broader list (**Table S2**). Many genes and gene classes that we

recovered as significant regulators of AC invasion are homologous to human genes that

have been previously studied in the context of cellular invasion and tumorigenesis

including *cec-6*/CBX1/CBX8 (82,84), *cfi-1*/ARID3A/ARID3C (85), *psr-1*/JMJD6 (86), *skp-*

1/SNW1 (87), and several TAFs (*taf*-1/TAF1/TAF1L, taf-*5*/TAF5/TAF5L, *taf-*

7.1/TAF7/TAF7L) (88–90). Additionally, we recovered nematode-specific genes including

nra-3, and cec-2, and genes whose human homologs have not been previously studied

in the context of cellular invasion to our knowledge, such as *cec-3* (homologous human

protein is uncharacterized) and *gna-2*/GNPNAT. Since the majority of the CRF genes we

identified as significant regulators of AC invasion have been previously studied in the

context of invasion in human development and cancer metastasis, these results

demonstrate the utility of the *C. elegans* AC invasion system as a genetically and optically

tractable *in vivo* environment to corroborate and characterize previously identified CRFs

that promote cellular invasion in human diseases such as rheumatoid arthritis and cancer.

Future studies should continue to characterize the relationship between CRFs identified

here and cellular invasion.

The dose of SWI/SNF ATPase dictates the penetrance of defects in AC invasion

For the majority of this study, we focused on characterizing the contribution of the

SWI/SNF ATP-dependent chromatin remodeling complex to cellular invasion as it was

highly represented among our list of significant regulators of AC invasion (**Table S2**) and

has been extensively studied in the context of both cellular invasion and cell cycle control

across a variety of animal models and in human cancers (31,35,40,42,48,56,59,83,91–

96). Prior whole-exome studies have determined that over 20% of human tumors harbor

mutations in one or more subunits of the SWI/SNF complex (32,48,98). Among the most

frequently mutated subunits of the chromatin remodeling complex throughout SWI/SNF-

deficient cancers is the core ATPase subunit BRG1/SMARCA4 (48,99) and the mutually

exclusive ATPase paralog to BRG (32,81,98,100).

Due to the high degree of structural similarity between BRG and BRM, mutation or

epigenetic silencing of either ATPase can be compensated by expression of the other.

Previous investigation has determined BRM to be an effective synthetic lethal target in

BRG1-deficient cancer, and vice-versa (101,102). Despite the compensatory nature of

BRG1/BRM in many tumorigenic contexts, concomitant loss of expression of the

ATPases has been described in metastatic murine models and patient-derived non-small-

cell lung cancer (NSCLC) cell lines and is associated with poor patient survival

(91,103,104). In *C elegans*, the sole SWI/SNF ATPase, *swsn-4,* has a high degree of

homology to both mammalian BRG1 and BRM, providing a unique opportunity to

accessibly model the connection between the dual loss of BRG1/BRM associated with

poor prognostic outcomes in NSCLC and cellular invasion in the AC. Additionally, a recent

study demonstrated a phenotypic dosage-sensitivity following loss of SWI/SNF

expression in relation to terminal differentiation of *C. elegans* muscle and vulval tissues

(35), suggesting that the context- and dose-dependent relationship seen in mammalian

development and SWI/SNF-deficient cancers may represent a more general behavior of

the complex.

Here we used the *C. elegans* AC invasion system as a model to investigate

whether the dose-dependent relationship between the ATPase and differentiated

phenotype extends to cellular invasion. The first indication that the functional dose of

SWI/SNF may have an instructive role in AC invasion was the relative enhancement of

all endogenously tagged subunits of the complex in the AC relative to neighboring VU

and VPC tissues (**Fig S4**). While it is tempting to interpret the enhancement of SWI/SNF

subunit expression in the AC as evidence for the dependence of cellular invasion on

SWI/SNF activity, it is also possible that this difference in expression is a consequence

of terminal differentiation, since at the time of invasion, the AC is terminally differentiated

unlike the VU and VPC). Further investigation is required to determine whether

endogenously tagged SWI/SNF subunits are enhanced in the AC relative other terminally

differentiated cells.

Additionally, we find that some degree of SWI/SNF intra- and inter-complex

regulation exists in the AC at the time of invasion with both SWI/SNF assemblies

cooperating to activate expression of the ATPase (**Fig S5 B**). It is possible that this added

level of complex autoregulation contributes to an "optimal" dose of the ATPase in a cell-

and context-specific manner. It is important to note that the model proposed in Fig S5 F

is incomplete and is included to describe the translational consequence of individual

SWI/SNF subunits after transcriptional knockdown of other individual SWI/SNF subunits.

However, these data can be meaningfully used to inform interpretations of results for our

enhanced SWI/SNF RNAi experiments in the AC.

To directly correlate SWI/SNF dose and invasive phenotype in the AC, we used a

combination of an enhanced *swsn-4(RNAi)* vector with a tissue-specific antiGFP-

targeting nanobody construct. These combined technologies allowed us to titrate the dose

of ATPase in the AC at the time of invasion. By assessing AC invasion phenotypes at

wildtype levels of SWSN-4 and in moderate and severe ATPase knockdown conditions,

results indicate that cellular invasion and cell cycle control depends on the dose of

functional SWI/SNF present in the AC.

In addition to reflecting the dose-dependent nature of the SWI/SNF ATPase in

cancer, our data in the AC is consistent with work done in *C. elegans* early mesoblast

development where complete loss of the *swsn-4* ATPase using a catalytically dead

mutant and lineage-specific knockout strategy results in loss of cell cycle arrest (35).

Although we cannot be sure that combining *swsn-4(RNAi)* with an antiGFP-targeting

nanobody to deplete the SWI/SNF ATPase results in complete loss of protein expression,

we show that treatment with the improved *swsn-4(RNAi)* vector alone is sufficient to

phenocopy the null phenotype previously reported in late mesoblast (SM) development

(**Fig S6**). Additionally, enhancement of the AC mitotic phenotype statistically tracked with

a progressive step down in mean expression of the ATPase in the AC across our

experiments. Using endogenous fluorescent reporters for conserved pro-invasive genes,

we also provide evidence that the SWI/SNF complex is involved broadly with the

regulation of the cell cycle- dependent and -independent subcircuits of the AC GRN (**Fig**

6). Altogether, this data supports the hypothesis that SWI/SNF cell-autonomously

contributes to cell cycle control in a dose-dependent manner and provides the first line of

evidence to link SWI/SNF ATPase dosage to the dichotomy between invasion and

proliferation (**Fig 8**).

The SWI/SNF BAF assembly promotes cellular invasion through induction of G_0

cell cycle arrest

While previous work in our lab, based on localization of a DNA licensing factor,

CDT-1, has demonstrated indirectly that ACs must arrest in a G_0/G_1 cell cycle state

(22,24), we lacked a sensitive enough tool to distinguish between these two interphase

states. From our recent work utilizing a CDK sensor to examine the proliferation-

quiescence decision in *C. elegans*, we can distinguish between pre-terminal cells in the

somatic gonad in G_1 (mean C/N ratio: 0.67+/-0.10) as compared to terminally

differentiated G_0 uterine cells (mean C/N ratio: 0.30+/-0.11) (36). Here, we compare CDK

activity measurements in the ACs of control animals with that of the terminal Pi lineage to

provide the first quantitative demonstration that ACs arrest in a CDK^{low} G_0 state to invade

(**Fig 4 A, E, F**). Furthermore, by combining the CDK sensor with loss of SWI/SNF

subunits, our data indicate that the SWI/SNF BAF assembly is specifically responsible for

regulation of G_0 cell cycle arrest in the AC.

Figure 8. SWI/SNF complex promotes AC invasion. Schematic summary of the how the SWI/SNF ATPase (S-4*, swsn-4*), PBAF (orange – S-7, swsn-*7*; S-9, *swsn-9*; P, *pbrm-1*), and BAF (blue – S-8, *swsn-8*) assemblies contribute to AC invasion.

The SWI/SNF core complex is composed of an ATPase, core, and accessory

factors, which are thought to collectively provide a platform of common factors that is

bound by assembly-specific subunits in a mutually exclusive manner. In *C. elegans*, the

association of SWSN-8 (BAF250a/ BAF250b/ ARID1A/ ARID1B) with the common factors

forms the SWI/SNF BAF assembly. Since we identified representative subunits from the

core of the complex and both SWI/SNF BAF and PBAF assemblies in our CRF RNAi

screen, we took a cell biological and genetic approach to investigate the role of the each

SWI/SNF assembly in promoting cellular invasion. Here, using a DHB-based CDK sensor

(36,70), we show that loss of either core or BAF assembly subunits specifically results in

mitotic ACs that failed to invade the BM. Our cell cycle sensor data first establishes that

wild-type AC invades in a G_0 CDK^{low} state, and second, that a major contribution of the

BAF assembly to AC invasion is through maintenance of this G_0 arrest, as many ACs that

failed to invade the BM had increasing CDK activity, indicative of cells cycling in G_1, S or

G_2. Alternatively, 14% of ACs that failed to invade the BM following loss of *swsn-*

4/ATPase of the complex had CDK activity ratios indicative of G_0 cell cycle arrest,

suggesting a cell cycle-independent defect. In support of this, forced arrest of BAF-

deficient ACs in G_0 was sufficient to significantly rescue invasion, whereas CKI-1

induction failed to rescue invasion in ACs with RNAi-mediated loss of *swsn-4*/ATPase.

Altogether, our results indicate that the SWI/SNF complex contributes to AC invasion

through regulation of G_0 cell cycle arrest via the BAF assembly. Further investigation will

require biochemical techniques to identify cell cycle regulators and TF targets of the BAF

assembly to provide a mechanistic explanation for how exactly BAF regulates the

chromatin landscape to promote invasion (**Fig 8**, blue arrow). Targeted DNA adenine

methyltransferase identification (TaDa) is an attractive biochemical approach that may be

adaptable to the AC invasion system, as this approach has been characterized as an

effective, tissue-specific method to identify TF-target sequence interactions in the *C.*

elegans epidermis (106).

807   The PBAF assembly regulates AC invasion and attachment to the BM

Based on homology and phenotypic characterization in this study and previous

publications, the *C. elegans* PBAF assembly consists of the PBRM-1 (PBRM1/BAF180),

SWSN-7 (BAF200/ARID2), and SWSN-9 (BRD7/BRD9) in association with the SWI/SNF

common factors. Previous work in *C. elegans* has not revealed a connection between

the PBAF assembly and cell cycle arrest. Our initial experiments with improved RNAi

vectors targeting PBAF subunits resulted in a lower penetrance of AC invasion defects

relative to loss of core or BAF subunits. Additionally, our CDK sensor data suggested that

non-invasive ACs deficient in *pbrm-1* remain in a G_0 cell cycle state. Thus, our data shows

no PBAF contribution to cell cycle control in the AC. To confirm this, we used the auxin

inducible degron (AID) system to robustly deplete the PBAF assembly through combined

loss of endogenous PBRM-1::mNG::AID with RNAi-mediated knockdown of either of the

other two PBAF assembly subunits, *swsn-7* or *swsn-9*. This combination knockdown

strategy corroborated our previous results as we saw no significant penetrance of extra

ACs. Rather, here we associate a striking AC detachment phenotype with strong

combined knockdown of the PBAF assembly subunits. We also note aberrant BM

morphology in some ACs deficient in PBAF subunits, with only one of the two BMs

removed, suggesting that this assembly regulates attachment and extracellular matrix

(ECM) remodeling in wild-type ACs to promote invasion. We hypothesize that the PBAF

assembly is regulating ventral BM attachment and ECM remodeling potentially through

the regulation of HIM-4/Hemicentin, an extracellular immunoglobulin-like matrix protein

that functions in the AC to fuse the two BMs through the formation of a novel BM-BM

adhesion, the B-LINK (107). Finally, although RNAi-mediated transcriptional knockdown

of PBAF assembly subunits only partially depleted levels of FOS-1::GFP, a key TF

responsible for the expression of MMPs and other pro-invasive targets, we detected

significant enhancement of invasion defects when depleting *fos-1* in a putative

hypomorphic *pbrm-1* background. Reciprocally, depletion of *pbrm-1* enhanced the

invasion defect of a quintuple MMP mutant. Since we noted multiple instances of AC-BM

detachment following PBAF assembly subunit depletion, we propose that PBAF functions

in part with FOS-1 to facilitating activating chromatin states at the regulatory regions of

pro-invasive genes required for BM attachment. Future studies should include an

investigation of PBAF assembly interactions with other FOS effectors and interactors at

the transcriptional level. Additionally, given the broad genomic regulation exhibited by

chromatin remodelers, biochemical techniques such as chromatin immunoprecipitation

(ChIP) and single cell RNA sequencing will be crucial going forward to generate a

comprehensive roster of direct and indirect targets of each SWI/SNF assembly in the AC.

Collectively, these results reveal a distinct contribution for each SWI/SNF assembly to the

process of cellular invasion at the phenotypic level and reiterate the dependence of each

assembly on the functional dose of ATPase in a cell.

Smith, et al. (2021)

ACKNOWLEDGEMENTS

We are thankful to David Gray, Ed Luk, Laura Mathies, Benjamin Martin, Robert Morabito,

Valerie Reinke, Courtney Tello, and Gerald Thomsen for advice and comments on this

manuscript. We would also like to thank Thom Geer of Nobska Imaging for advice and

'scientific enabling'. Some *C. elegans* strains were provided by the CGC, which is funded

by NIH Office of Research Infrastructure Programs (P40 OD010440).

AUTHOR CONTRIBUTIONS

J.J.S. and D.Q.M. designed the experiments. J.J.S., Y.X., M.A.Q.M., and T.N.M.-K.

performed the experiments. J.J.S., A.Q.K., and M.C. organized the CRF RNAi screen.

J.J.S., M.C., N.P., and Y.X. performed the RNAi screen. J.J.S., A.Q.K., S.L., T.N.M.-K.,

Y.X., K.W. and P.K generated strains. J.J.S performed the data analysis and prepared

the manuscript with feedback from other authors. J.J.S. and D.Q.M acquired funding for

the completion of this project.

DECLARATION OF INTERESTS

The authors declare no competing interests.

FUNDING

This work was funded by the National Institute of General Medical Sciences (NIGMS)

[1R01GM121597-01 to D.Q.M.]. D.Q.M. is also a Damon Runyon-Rachleff Innovator

supported (in part) by the Damon Runyon Cancer Research Foundation [DRR-47-17].

J.J.S was supported by an NIH/NIGMS Diversity Supplement [3R01GM121597-02S1]

and the W. Burghardt Turner Fellowship. M.A.Q.M. [3R01GM121597-03S1] and F.E.Q.M.

[3R01GM121597-04S1] are supported by an NIH/NIGMS Diversity Supplement. A.Q.K.

[F31GM128319], R.C.A [1F31GM1283190], and T.N.M.-K. [F31HD100091-01] were

supported by the NIH. N.J.P [132969-PF-18-226-01-CSM] was supported by the

American Cancer Society (ACS). P.K. [R01NS118078] is supported by the National

Institute of Neurological Disorders and Stroke.

MATERIALS & METHODS

C. elegans **strains and culture conditions**

All animals were maintained under standard conditions and cultured at 25°C, except

strains containing temperature-sensitive alleles *swsn-1(os22), swsn-4(os13)*, and the

uterine-specific RNAi hypersensitive strain used in the chromatin remodeler screen

containing the *rrf-3(pk1426)* allele, which were maintained at either 15°C or 20°C (108).

The heat shock inducible *cki-1::mTagBFP2* transgene was expressed via incubating

animals at 32°C for 2-3 hours in a water bath starting at the P6.p 2-cell VPC stage.

Animals were synchronized for experiments through alkaline hypochlorite treatment of

gravid adults to isolate eggs (109). In the text and figures, we designate linkage to a

promoter through the use of a (p) and fusion of a proteins via a (::) annotation.

Molecular biology and microinjection

SWI/SNF subunits *swsn-4* and *swsn-8* were tagged at their endogenous loci using

CRISPR/Cas9 genome editing via microinjection into the early adult hermaphrodite

syncytial gonad (110,111). Repair templates were generated as synthetic DNAs from

either Integrated DNA Technologies (IDT) as gene blocks (gBlocks) or Twist Biosciences

as DNA fragments and cloned into *ccdB* compatible sites in pDD282 by New England

Biolabs Gibson assembly (112). Homology arms ranged from 690-1200 bp (see **Tables**

S5 for additional details). sgRNAs were constructed by EcoRV and NheI digestion of the

plasmid pDD122. A 230 bp amplicon was generated replacing the sgRNA targeting

sequence from pDD122 with a new sgRNA and NEB Gibson assembly was used to

generate new sgRNA plasmids (see **Table S5** for additional details). Hermaphrodite

adults were co-injected with guide plasmid (50 ng/μL), repair plasmid (50 ng/μL), and an

extrachromosomal array marker (pCFJ90, 2.5 ng/μL), and incubated at 25 ºC for several

days before screening and floxing protocols associated with the SEC system (112).

RNA interference (RNAi)

All 269 RNAi clones assessed in the chromatin remodeler screen were derived from the

commercially available Vidal or Ahringer RNAi libraries. Presence of inserts into the

L4440 RNAi vector was confirmed via colony PCR amplification of all L4440 vectors used

in the chromatin remodeler screen. Vectors which resulted in penetrant loss of invasion

(see **Table S2**) were also sequenced to confirm the identity of the insert targeting

chromatin remodeler genes in the L4440 vector using Sanger sequencing at the

Genomics Core Facility at Stony Brook University. An RNAi sub-library of SWI/SNF

subunits was constructed by cloning 950-1000 bp of synthetic DNA based on cDNA

sequences available on WormBase (www.wormbase.org) into the highly efficient T444T

RNAi vector (113,114). Synthetic DNAs were generated by Twist Biosciences as DNA

fragments and cloned into restriction digested T444T using NEB Gibson Assembly (see

Tables S6 for additional details). For all experiments, synchronized L1 stage animals

were directly exposed to RNAi through feeding with bacteria expressing dsRNA (115).

Auxin-mediated degradation

To combine RNAi with the depletion of AID-tagged proteins, 1 mM K-NAA was used, and

its effects were analyzed as previously described(116). Briefly, L1 animals were first

synchronized by sodium hypochlorite treatment and transferred to NGM plates seeded

with the RNAi vector of interest. At the P6.p 1-cell stage, a time in development where

the AC has already undergone specification, animals were transferred to RNAi-seeded

plates treated with K-NAA. Animals were staged by DIC.

Live cell microscopy

All micrographs included in this manuscript were collected on a Hamamatsu Orca EM-

CCD camera mounted on an upright Zeiss AxioImager A2 with a Borealis-modified

CSU10 Yokagawa spinning disk scan head using 405nm, 488 nm, and 561 nm Vortran

lasers in a VersaLase merge and a Plan-Apochromat 100x/1.4 (NA) Oil DIC objective.

MetaMorph software (Molecular Devices) was used for microscopy automation. Several

experiments and all RNAi screening were scored using epifluorescence visualized on a

Zeiss Axiocam MRM camera, also mounted on an upright Zeiss AxioImager A2 and a

Plan-Apochromat 100x/1.4 (NA) Oil DIC objective. Animals were mounted into a drop of

M9 on a 5% Noble agar pad containing approximately 10 mM sodium azide anesthetic

and topped with a coverslip.

Assessment of AC invasion

Both for the purposes of the CRF RNAi screen and all other experiments AC invasion

was scored at the P6.p 4-cell stage, when 100% of wild-type animals exhibit a breach in

the BM (14). In strains with the laminin::GFP transgene, an intact green fluorescent barrier

under the AC was used to assess invasion. Wild-type invasion is defined as a breach as

wide as the basolateral surface of the AC (14). Raw scoring data is available in **Tables**

S1 and S4.

Image quantification and statistical analyses

Images were processed using Fiji/ImageJ (v.2.1.0/1.53c) (117). Expression levels of

GFP::SWSN-4, SWSN-8::GFP, PBRM-1::eGFP, and PBRM-1::mNG::AID were

measured by quantifying the mean gray value of AC nuclei, defined as somatic gonad

cells near the primary vulva expressing the *cdh-3p::mCherry::moeABD* transgene.

Background subtraction was performed by rolling ball background subtraction (size=50).

For characterization of experiments involving SWI/SNF endogenous tags and AC GRN

TFs::GFP treated with *SWI/SNF(RNAi)* and GFP-targeting nanobody the L3 stage, only

animals exhibiting defects in invasion were included in the analysis. Data was normalized

to negative control (empty vector) values for the plots in **Fig 3** and **Fig S8**. Quantification

of either CDK cell cycle sensor (either DHB::GFP or DHB::2xmKate2) was performed by

hand, as previously described (36). Images were overlaid and figures were assembled

using Adobe Photoshop 2020 (v. 21.1.2) and Adobe Illustrator 2020 (v. 24.1.2),

respectively. Statistical analyses and plotting of data were conducted using RStudio (v.

1.2.1335). Statistical significance was determined using either a two-tailed Student's t-

test or Fisher's exact probability test. Figure legends specify when each test was used

and the p-value cut-off.

Smith, et al. (2021)

REFERENCES

1. Rowe RG, Weiss SJ. Breaching the basement membrane: who, when and how?

Trends Cell Biol. 2008;18(11):560–74.

2. Nourshargh S, Alon R. Leukocyte Migration into Inflamed Tissues. Immunity

[Internet]. 2014;41(5):694–707. Available from:

http://dx.doi.org/10.1016/j.immuni.2014.10.008

3. Medwig TN, Matus DQ. Breaking down barriers: the evolution of cell invasion.

Curr Opin Genet Dev [Internet]. 2017;47:33–40. Available from:

http://dx.doi.org/10.1016/j.gde.2017.08.003

4. Liu Y, Pan YF, Xue YQ, Fang LK, Guo XH, Guo X, et al. UPAR promotes tumor-

like biologic behaviors of fibroblast-like synoviocytes through PI3K/Akt signaling

pathway in patients with rheumatoid arthritis. Cell Mol Immunol [Internet].

2018;15(2):171–81. Available from: http://dx.doi.org/10.1038/cmi.2016.60

5. Ye Y, Gao X, Yang N. LncRNA ZFAS1 promotes cell migration and invasion of

fibroblast-like synoviocytes by suppression of miR-27a in rheumatoid arthritis.

Hum Cell. 2018;31(1):14–21.

6. Hanahan D, Weinberg RA. Hallmarks of cancer: The next generation. Cell

[Internet]. 2011;144(5):646–74. Available from:

http://dx.doi.org/10.1016/j.cell.2011.02.013

7. Justus CR, Leffler N, Ruiz-Echevarria M, Yang L V. In vitro cell migration and

invasion assays. J Vis Exp. 2014;(88):1–8.

8. Li BB, Scott EY, Dean Chamberlain M, Duong BTV, Zhang S, Done SJ, et al. Cell

invasion in digital microfluidic microgel systems. Sci Adv. 2020;6(29).

9. Eibl G, Reber HA. A xenograft nude mouse model for perineural invasion and

recurrence in pancreatic cancer. Pancreas. 2005;31(3):258–62.

10. Busuttil RA, Liu DS, Di Costanzo N, Schröder J, Mitchell C, Boussioutas A. An

orthotopic mouse model of gastric cancer invasion and metastasis. Sci Rep.

2018;8(1):2–9.

11. Cheng B, Wang Q, Song Y, Liu Y, Liu Y, Yang S, et al. MIF inhibitor, ISO-1,

attenuates human pancreatic cancer cell proliferation, migration and invasion in

vitro, and suppresses xenograft tumour growth in vivo. Sci Rep. 2020;10(1):1–10.

12. Wen J, Yin P, Li L, Kang G, Ning G, Cao Y, et al. Knockdown of Matrix

Metallopeptidase 9 Inhibits Metastasis of Oral Squamous Cell Carcinoma Cells in

a Zebrafish Xenograft Model. Biomed Res Int. 2020;2020.

13. Guo Y, Fan Y, Pei X. Fangjihuangqi Decoction inhibits MDA-MB-231 cell invasion

in vitro and decreases tumor growth and metastasis in triple-negative breast

cancer xenografts tumor zebrafish model. Cancer Med. 2020;9(7):2564–78.

14. Sherwood DR, Sternberg PW. Anchor cell invasion into the vulval epithelium in C.

elegans. Dev Cell. 2003;5(1):21–31.

15. Kelley LC, Lohmer LL, Hagedorn EJ, Sherwood DR. Traversing the basement

membrane in vivo: A diversity of strategies. J Cell Biol. 2014;204(3):291–302.

16. Sherwood DR, Butler JA, Kramer JM, Sternberg PW. FOS-1 promotes basement-

membrane removal during anchor-cell invasion in C. elegans. Cell.

2005;121(6):951–62.

17. Hagedorn EJ, Yashiro H, Ziel JW, Ihara S, Wang Z, Sherwood DR. Integrin Acts

Upstream of Netrin Signaling to Regulate Formation of the Anchor Cell's Invasive

Membrane in C. elegans. Dev Cell [Internet]. 2009;17(2):187–98. Available from:

http://dx.doi.org/10.1016/j.devcel.2009.06.006

18.   Hagedorn EJ, Kelley LC, Naegeli KM, Wang Z, Chi Q, Sherwood DR. ADF/cofilin

promotes invadopodial membrane recycling during cell invasion in vivo. J Cell

Biol. 2014;204(7):1209–18.

19.   Morrissey MA, Jayadev R, Miley GR, Blebea CA, Chi Q, Ihara S, et al. SPARC

Promotes Cell Invasion In Vivo by Decreasing Type IV Collagen Levels in the

Basement Membrane. PLoS Genet. 2016;12(2).

20.   Hagedorn EJ, Ziel JW, Morrissey MA, Linden LM, Wang Z, Chi Q, et al. The netrin

receptor DCC focuses invadopodia-driven basement membrane transmigration in

vivo. J Cell Biol. 2013;201(6):903–13.

21.   Kelley LC, Chi Q, Cáceres R, Hastie E, Schindler AJ, Matus DQ, et al. Adaptive

F-actin polymerization and localized ATP production drive basement membrane

invasion in the absence of MMPs. Dev Cell. 2019;48(3):313–28.

22.   Medwig-Kinney TN, Smith JJ, Palmisano NJ, Tank S, Zhang W, Matus DQ. A

developmental gene regulatory network for C. elegans

anchor cell invasion. Development [Internet]. 2020 Jan 1;147(1):dev185850.

Available from: http://dev.biologists.org/content/147/1/dev185850.abstract

23.   Kohrman AQ, Matus DQ. Divide or Conquer: Cell Cycle Regulation of Invasive

Behavior. Trends Cell Biol [Internet]. 2016;27(1):12–25. Available from:

http://dx.doi.org/10.1016/j.tcb.2016.08.003

24.   Matus DQ, Lohmer LL, Kelley LC, Schindler AJ, Kohrman AQ, Barkoulas M, et al.

Invasive Cell Fate Requires G1 Cell-Cycle Arrest and Histone Deacetylase-

Mediated Changes in Gene Expression. Dev Cell [Internet]. 2015;35(2):162–74.

Available from: http://dx.doi.org/10.1016/j.devcel.2015.10.002

25.   Brand M, Nakka K, Zhu J, Dilworth FJ. Polycomb/Trithorax Antagonism: Cellular

Memory in Stem Cell Fate and Function. Cell Stem Cell [Internet].

2019;24(4):518–33. Available from: https://doi.org/10.1016/j.stem.2019.03.005

26.   Simon JA, Tamkun JW. Programming off and on states in chromatin: Mechanisms

of Polycomb and trithorax group complexes. Curr Opin Genet Dev.

2002;12(2):210–8.

27.   Kadoch C, Williams RT, Calarco JP, Miller EL, Weber CM, Braun SMG, et al.

Dynamics of BAF – Polycomb complex opposition on heterochromatin in normal

and oncogenic states. Nat Publ Gr [Internet]. 2016;49(2):213–22. Available from:

http://dx.doi.org/10.1038/ng.3734

28.   Friman ET, Deluz C, Meireles-Filho ACA, Govindan S, Gardeux V, Deplancke B,

et al. Dynamic regulation of chromatin accessibility by pluripotency transcription

factors across the cell cycle. bioRxiv. 2019;1–28.

29.   Ma Y, McKay DJ, Buttitta L. Changes in chromatin accessibility ensure robust cell

cycle exit in terminally differentiated cells. PLoS Biol. 2019;17(9):1–29.

30.   Wang S, Tang NH, Lara-Gonzalez P, Zhao Z, Cheerambathur DK, Prevo B, et al.

A toolkit for GFP-mediated tissue-specific protein degradation in C. elegans .

Development. 2017;144(14):2694–701.

31.   Bai J, Mei P, Zhang C, Chen F, Li C, Pan Z, et al. BRG1 Is a Prognostic Marker

and Potential Therapeutic Target in Human Breast Cancer. PLoS One.

2013;8(3):1–9.

32.   Kadoch C, Hargreaves DC, Hodges C, Elias L, Ho L, Ranish J, et al. a n a ly s i s

1053   Proteomic and bioinformatic analysis of mammalian SWI / SNF complexes

1054   identifies extensive roles in human malignancy. Nat Publ Gr [Internet].

1055   2013;45(6):592–601. Available from: http://dx.doi.org/10.1038/ng.2628

1056   33.   Kaufmann B, Wang B, Zhong S, Laschinger M, Patil P, Lu M, et al. BRG1

1057   promotes hepatocarcinogenesis by regulating proliferation and invasiveness.

1058   PLoS One. 2017;12(7):1–14.

1059   34.   Bögershausen N, Wollnik B. Mutational Landscapes and Phenotypic Spectrum of

1060   SWI/SNF-Related Intellectual Disability Disorders. Front Mol Neurosci.

1061   2018;11(August):1–18.

1062   35.   van der Vaart A, Godfrey M, Portegijs V, van den Heuvel S. Dose-dependent

1063   functions of SWI/SNF BAF in permitting and inhibiting cell proliferation in vivo. Sci

1064   Adv. 2020;6(21).

1065   36.   Adikes RC, Kohrman AQ, Martinez MAQ, Palmisano NJ, Smith JJ, Medwig-

1066   Kinney TN, et al. Visualizing the metazoan proliferation-quiescence decision in

1067   vivo. Elife. 2020;9:1–74.

1068   37.   Rual JF, Ceron J, Koreth J, Hao T, Nicot AS, Hirozane-Kishikawa T, et al. Toward

1069   improving Caenorhabditis elegans phenome mapping with an ORFeome-based

1070   RNAi library. Genome Res. 2004;14(10 B):2162–8.

1071   38.   Kamath RS, Ahringer J. Genome-wide RNAi screening in Caenorhabditis

1072   elegans. Methods. 2003;30(4):313–21.

1073   39.   Matus DQ, Li X-Y, Durbin S, Agarwal D, Chi Q, Weiss SJ, et al. In vivo

1074   identification of regulators of cell invasion across basement membranes. Sci

Signal [Internet]. 2010;3(120):ra35. Available from:

http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3917318&tool=pmcentr

ez&rendertype=abstract

40. W W. The SWI / SNF Family of ATP-Dependent Chromatin Remodelers : Similar

Mechanisms for Diver Function. In: Workman JL, editor. Protein Complexes that

Modify Chromatin. Springer-Verlag Berling Heidelberg; 2003. p. 144–65.

41. Sawa H, Kouike H, Okano H. Components of the SWI/SNF complex are required

for asymmetric cell division in C. elegans. Mol Cell. 2000;6(3):617–24.

42. Cui M, Fay DS, Han M. Lin-35/Rb cooperates with the SWI/SNF complex to

control Caenorhabditis elegans larval development. Genetics. 2004;167(3):1177–

85.

43. Large EE, Mathies LD. Caenorhabditis elegans SWI/SNF subunits control

sequential developmental stages in the somatic gonad. G3 (Bethesda) [Internet].

2014;4(3):471–83. Available from:

http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3962486&tool=pmcentr

ez&rendertype=abstract

44. Ertl I, Porta-De-La-Riva M, Gómez-Orte E, Rubio-Peña K, Corrales D, Cornes E,

et al. Functional interplay of two paralogs encoding SWI/SNF chromatin-

remodeling accessory subunits during Caenorhabditis elegans development.

Genetics. 2016;202(3):961–75.

45. Cenik BK, Shilatifard A. COMPASS and SWI/SNF complexes in development and

disease. Nat Rev Genet [Internet]. 2021;22(1):38–58. Available from:

http://dx.doi.org/10.1038/s41576-020-0278-0

46.   Kadoch C, Crabtree GR. Mammalian SWI/SNF chromatin remodeling complexes

and cancer: Mechanistic insights gained from human genomics. Sci Adv

[Internet]. 2015;1(5):e1500447–e1500447. Available from:

http://advances.sciencemag.org/cgi/doi/10.1126/sciadv.1500447

47.   Nickerson JA, Wu Q, Imbalzano AN. Mammalian SWI/SNF enzymes and the

epigenetics of tumor cell metabolic reprogramming. Front Oncol. 2017;7(APR):1–

9.

48.   Mittal P, Roberts CWM. The SWI/SNF complex in cancer — biology, biomarkers

and therapy. Nat Rev Clin Oncol [Internet]. 2020;17(7):435–48. Available from:

http://dx.doi.org/10.1038/s41571-020-0357-3

49.   Muchardt C, Yaniv M. When the SWI/SNF complex remodels...the cell cycle.

Oncogene. 2001;20(24):3067–75.

50.   Nagl NG, Zweitig DR, Thimmapaya B, Beck GR, Moran E. The c-myc gene is a

direct target of mammalian SWI/SNF-related complexes during differentiation-

associated cell cycle arrest. Cancer Res. 2006;66(3):1289–93.

51.   Moshkin YM, Mohrmann L, van Ijcken WFJ, Verrijzer CP. Functional

Differentiation of SWI/SNF Remodelers in Transcription and Cell Cycle Control.

Mol Cell Biol [Internet]. 2007;27(2):651–61. Available from:

http://mcb.asm.org/cgi/doi/10.1128/MCB.01257-06

52.   Ruijtenberg S, Van Den Heuvel S. G1/S Inhibitors and the SWI/SNF Complex

Control Cell-Cycle Exit during Muscle Differentiation. Cell [Internet].

2015;162(2):300–13. Available from: http://dx.doi.org/10.1016/j.cell.2015.06.013

53.   Innis SM, Cabot B. GBAF, a small BAF sub-complex with big implications: a

systematic review. Epigenetics and Chromatin [Internet]. 2020;13(1):1–18.

Available from: https://doi.org/10.1186/s13072-020-00370-8

54.   Son EY, Crabtree GR. The role of BAF (mSWI/SNF) complexes in mammalian

neural development. Am J Med Genet C Semin Med Genet. 2014;(650).

55.   Mathies LD, Blackwell GG, Austin MK, Edwards AC, Riley BP, Davies AG, et al.

SWI/SNF chromatin remodeling regulates alcohol response behaviors in

Caenorhabditis elegans and is associated with alcohol dependence in humans.

Proc Natl Acad Sci U S A [Internet]. 2015;112(10):3032–7. Available from:

/pmc/articles/PMC4364201/?report=abstract

56.   Kuzmanov A, Karina EI, Kirienko N V., Fay DS. The Conserved PBAF

Nucleosome-Remodeling Complex Mediates the Response to Stress in

Caenorhabditis elegans. Mol Cell Biol. 2014;34(6):1121–35.

57.   Hayes GD, Riedel CG, Ruvkun G. The Caenorhabditis elegans somi-1 zinc finger

protein and SWI/SNF promote regulation of development by the mir-84

microRNA. Genes Dev. 2011;25(19):2079–92.

58.   Shibata Y, Uchida M, Takeshita H, Nishiwaki K, Sawa H. Multiple functions of

PBRM-1/Polybromo- and LET-526/Osa-containing chromatin remodeling

complexes in C. Elegans development. Dev Biol [Internet]. 2012;361(2):349–57.

Available from: http://dx.doi.org/10.1016/j.ydbio.2011.10.035

59.   Weinberg P, Flames N, Sawa H, Garriga G, Hobert O. The SWI/SNF chromatin

remodeling complex selectively affects multiple aspects of serotonergic neuron

differentiation. Genetics. 2013;194(1):189–98.

60.   Riedel CG, Dowen RH, Lourenco GF, Natalia V, Heimbucher T, West JA, et al.

DAF-16/FOXO employs the chromatin remodeller SWI/SNF to promote stress

resistance and longevity. 2013;15(5):491–501.

61.   Kipreos ET, van den Heuvel S. Developmental Control of the Cell Cycle: Insights

from Caenorhabditis elegans. Genetics. 2019;211(3):797–829.

62.   Sturm Á, Saskoï É, Tibor K, Weinhardt N, Vellai T. Highly efficient RNAi and

Cas9-based auto-cloning systems for C. Elegans research. Nucleic Acids Res.

2018;46(17):1–13.

63.   Dickinson DJ, Goldstein B. CRISPR-based methods for caenorhabditis elegans

genome engineering. Genetics. 2016;202(3):885–901.

64.   Mashtalir N, D'Avino AR, Michel BC, Luo J, Pan J, Otto JE, et al. Modular

Organization and Assembly of SWI/SNF Family Chromatin Remodeling

Complexes. Cell [Internet]. 2018;175(5):1272-1288.e20. Available from:

https://doi.org/10.1016/j.cell.2018.09.032

65.   Rago F, DiMare MT, Elliott GN, Ruddy DA, Sovath S, Kerr G, et al. Degron

mediated BRM/SMARCA2 depletion uncovers novel combination partners for

treatment of BRG1/SMARCA4-mutant cancers. Biochem Biophys Res Commun

[Internet]. 2019;508(1):109–16. Available from:

https://doi.org/10.1016/j.bbrc.2018.09.009

66.   Ziel JW, Matus DQ, Sherwood DR. An expression screen for RhoGEF genes

involved in C. elegans gonadogenesis. Gene Expr Patterns [Internet].

2009;9(6):397–403. Available from:

http://www.nature.com/doifinder/10.1038/jid.2014.371

67.   Rimann I, Hajnal A. Regulation of anchor cell invasion and uterine cell fates by

the egl-43 Evi-1 proto-oncogene in Caenorhabditis elegans. Dev Biol.

2007;308(1):187–95.

68.   Deng T, Stempor P, Appert A, Daube M, Ahringer J, Hajnal A, et al. The

caenorhabditis elegans homolog of the evi1 proto-oncogene, egl-43, coordinates

G1 cell cycle arrest with pro-invasive gene expression during anchor cell invasion.

PLoS Genet [Internet]. 2020;16(3). Available from:

http://dx.doi.org/10.1371/journal.pgen.1008470

69.   Hwang BJ, Meruelo AD, Sternberg PW. C. elegans EVI1 proto-oncogene, EGL-

43, is necessary for Notch-mediated cell fate specification and regulates cell

invasion. Development. 2007;134(4):669–79.

70.   Spencer SL, Cappell SD, Tsai F, Overton KW, Wang L, Meyer T. The

Proliferation-Quiescence Decision Is Controlled by a Bifurcation in CDK2 Activity

at Mitotic Exit. 2013;155(2):369–83.

71.   Kimble J, Hirsh D. The postembryonic cell lineages of the hermaphrodite and

male gonads in Caenorhabditis elegans. Dev Biol. 1979;70(2):396–417.

72.   Newman AP, White JG, Sternberg PW. Morphogenesis of the C. elegans

hermaphrodite uterus. Development [Internet]. 1996 Nov 1;122(11):3617 LP –

3626. Available from: http://dev.biologists.org/content/122/11/3617.abstract

73.   Cao Y, Cairns BR, Kornberg RD, Laurent BC. Sfh1p, a component of a novel

chromatin-remodeling complex, is required for cell cycle progression. Mol Cell

Biol. 1997;17(6):3323–34.

74.   Angus-Hill ML, Schlichter A, Roberts D, Erdjument-Bromage H, Tempst P, Cairns

BR. A Rsc3/Rsc30 zinc cluster dimer reveals novel roles for the chromatin

1190   remodeler RSC in gene expression and cell cycle control. Mol Cell.

1191   2001;7(4):741–51.

1192 75. Zhang L, Ward JD, Cheng Z, Dernburg AF. The auxin-inducible degradation (AID)

1193   system enables versatile conditional protein depletion in C. elegans. Development

1194   [Internet]. 2015;142(24):4374–84. Available from:

1195   http://dev.biologists.org/content/142/24/4374.abstract

1196 76. Martinez MAQ, Kinney BA, Medwig-Kinney TN, Ashley G, Ragle JM, Johnson L,

1197   et al. Rapid degradation of C. elegans proteins at single-cell resolution with a

1198   synthetic auxin. G3 (Bethesda). 2020;10(January):267–80.

1199 77. Wang L, Shen W, Lei S, Matus D, Sherwood D, Wang Z. MIG-10 (Lamellipodin)

1200   stabilizes invading cell adhesion to basement membrane and is a negative

1201   transcriptional target of EGL-43 in C. elegans. Biochem Biophys Res Commun

1202   [Internet]. 2014;452(3):328–33. Available from:

1203   http://dx.doi.org/10.1016/j.bbrc.2014.08.049

1204 78. Sherwood, DR., Plastino J. Invading , Leading and Navigating Cells in

1205   Caenorhabditis elegans : Insights into Cell Movement in Vivo. In: WormBook : the

1206   online review of C elegans biology. 2018. p. 53–78.

1207 79. Ribeiro-Silva C, Vermeulen W, Lans H. SWI/SNF: Complex complexes in genome

1208   stability and cancer. DNA Repair. 2019.

1209 80. Lu P, Roberts CW. The SWI/SNF tumor suppressor complex. Nucleus [Internet].

1210   2013;4(5):374–8. Available from:

1211   http://www.tandfonline.com/doi/abs/10.4161/nucl.26654

1212 81. Hargreaves DC, Crabtree GR. ATP-dependent chromatin remodeling: Genetics,

genomics and mechanisms. Cell Res [Internet]. 2011;21(3):396–420. Available

from: http://dx.doi.org/10.1038/cr.2011.32

82.   Yang YF, Pan YH, Tian QH, Wu DC, Su SG. CBX1 Indicates Poor Outcomes and

Exerts Oncogenic Activity in Hepatocellular Carcinoma. Transl Oncol [Internet].

2018;11(5):1110–8. Available from: https://doi.org/10.1016/j.tranon.2018.07.002

83.   Bracken AP, Brien GL, Verrijzer CP. Dangerous liaisons: interplay between

SWI/SNF, NuRD, and Polycomb in chromatin regulation and cancer. Genes Dev

[Internet]. 2019;1–24. Available from:

http://genesdev.cshlp.org/lookup/doi/10.1101/gad.326066.119

84.   Wang G, Tang J, Zhan W, Zhang R, Zhang M, Liao D, et al. CBX8 suppresses

tumor metastasis via repressing snail in esophageal squamous cell carcinoma.

Theranostics. 2017;7(14):3478–88.

85.   Ma J, Zhan Y, Xu Z, Li Y, Luo A, Ding F, et al. ZEB1 induced miR-99b/let-7e/miR-

125a cluster promotes invasion and metastasis in esophageal squamous cell

carcinoma. Cancer Lett. 2017;398:37–45.

86.   Zhou D.X., Zhou D., Zhan S.Q., Wang P., Qin K., Gan W. LXF. Inhibition of

JMJD6 expression reduces the proliferation, migration and invasion of

neuroglioma stem cells. Neoplasma. 2017;64(5).

87.   Bondy-Chorney E, Baldwin RM, Didillon A, Chabot B, Jasmin BJ, Côté J. RNA

binding protein RALY promotes Protein Arginine Methyltransferase 1 alternatively

spliced isoform v2 relative expression and metastatic potential in breast cancer

cells. Int J Biochem Cell Biol [Internet]. 2017;91(March):124–35. Available from:

http://dx.doi.org/10.1016/j.biocel.2017.07.008

88. Zhong S, Yan H, Chen Z, Li Y, Shen Y, Wang Y, et al. Overexpression of TAF1L

promotes cell proliferation, migration and invasion in esophageal squamous cell

carcinoma. J Cancer. 2019;10(4):979–89.

89. Gillison ML, Akagi K, Xiao W, Jiang B, Pickard RKL, Li J, et al. Human

papillomavirus and the landscape of secondary genetic alterations in oral cancers.

Genome Res. 2019;29(1):1–17.

90. Yang L, Zeng W, Sun H, Huang F, Yang C, Cai X, et al. Bioinformatical Analysis

of Gene Expression Omnibus Database Associates TAF7/CCNB1, TAF7/CCNA2,

and GTF2E2/CDC20 Pathways with Glioblastoma Development and Prognosis.

World Neurosurg [Internet]. 2020;138:e492–514. Available from:

https://doi.org/10.1016/j.wneu.2020.02.159

91. Herpel E, Rieker RJ, Dienemann H, Muley T, Meister M, Hartmann A, et al.

SMARCA4 and SMARCA2 deficiency in non–small cell lung cancer:

immunohistochemical survey of 316 consecutive specimens. Ann Diagn Pathol

[Internet]. 2017;26:47–51. Available from:

http://dx.doi.org/10.1016/j.anndiagpath.2016.10.006

92. Shibata Y, Uchida M, Takeshita H, Nishiwaki K, Sawa H. Multiple functions of

PBRM-1/Polybromo- and LET-526/Osa-containing chromatin remodeling

complexes in C. Elegans development. Dev Biol [Internet]. 2012;361(2):349–57.

Available from: http://dx.doi.org/10.1016/j.ydbio.2011.10.035

93. Zhang Z, Wang F, Du C, Guo H, Ma L, Liu X, et al. BRM/SMARCA2 promotes the

proliferation and chemoresistance of pancreatic cancer cells by targeting

JAK2/STAT3 signaling. Cancer Lett [Internet]. 2017;402:213–24. Available from:

http://dx.doi.org/10.1016/j.canlet.2017.05.006

94.   Wu RC, Wang TL, Shih IM. The emerging roles of ARID1A in tumor suppression.

Cancer Biol Ther. 2014;15(6):655–64.

95.   Clapier CR, Iwasa J, Cairns BR, Peterson CL. Mechanisms of action and

regulation of ATP-dependent chromatin-remodelling complexes. Nat Publ Gr

[Internet]. 2017; Available from: http://dx.doi.org/10.1038/nrm.2017.26

96.   Vries RGJ, Bezrookove V, Zuijderduijn LMP, Kia SK, Houweling A, Oruetxebarria

I, et al. Cancer-associated mutations in chromatin remodeler hSNF5 promote

chromosomal instability by compromising the mitotic checkpoint. Genes Dev.

2005;19(6):665–70.

97.   Kadoch C, Copeland RA, Keilhack H. PRC2 and SWI/SNF Chromatin

Remodeling Complexes in Health and Disease. Biochemistry. 2016;55(11):1600–

14.

98.   Shain AH, Pollack JR. The Spectrum of SWI/SNF Mutations, Ubiquitous in

Human Cancers. PLoS One. 2013;8(1).

99.   Fernando TM, Piskol R, Bainer R, Sokol ES, Trabucco SE, Zhang Q, et al.

Functional characterization of SMARCA4 variants identified by targeted exome-

sequencing of 131,668 cancer patients. Nat Commun [Internet]. 2020;11(1):1–13.

Available from: http://dx.doi.org/10.1038/s41467-020-19402-8

100. Dutta A, Sardiu M, Gogol M, Gilmore J, Zhang D, Florens L, et al. Composition

and Function of Mutant Swi/Snf Complexes. Cell Rep [Internet]. 2017;18(9):2124–

34. Available from: http://linkinghub.elsevier.com/retrieve/pii/S2211124717301158

101. Hoffman GR, Rahal R, Buxton F, Xiang K, McAllister G, Frias E, et al. Functional

epigenetics approach identifies BRM/SMARCA2 as a critical synthetic lethal

target in BRG1-deficient cancers. Proc Natl Acad Sci U S A. 2014;111(8):3128–

33.

102. Sasaki M, Ogiwara H. Synthetic lethal therapy based on targeting the vulnerability

of SWI/SNF chromatin remodeling complex-deficient cancers. Cancer Sci.

2020;111(3):774–82.

103. Reisman DN, Sciarrotta J, Wang W, Funkhouser WK, Weissman BE. Loss of

BRG1/BRM in human lung cancer cell lines and primary lung cancers: Correlation

with poor prognosis. Cancer Res. 2003;63(3):560–6.

104. Marquez-Vilendrer SB, Rai SK, Gramling SJ, Lu L, Reisman DN. Loss of the

SWI/SNF atpase subunits BRM and BRG1 drives lung cancer development.

Oncoscience. 2016;3(11–12):322–36.

105. Sulston JE, Horvitz HR. Post-embryonic cell lineages of the nematode,

Caenorhabditis elegans. Dev Biol. 1977;56(1):110–56.

106. Katsanos D, Barkoulas M. Tissue-specific transcription factor target identification

in the Caenorhabditis elegans epidermis using targeted DamID. bioRxiv [Internet].

2020; Available from: https://doi.org/10.1101/2020.12.17.423252

107. Morrissey MA, Keeley DP, Hagedorn EJ, McClatchey STH, Chi Q, Hall DH, et al.

B-LINK: A Hemicentin, Plakin, and Integrin-Dependent Adhesion System that

Links Tissues by Connecting Adjacent Basement Membranes. Dev Cell.

2014;31(3):319–31.

108. Simmer F, Tijsterman M, Parrish S, Koushika SP, Nonet ML, Fire A, et al. Loss of

the putative RNA-directed RNA polymerase RRF-3 makes C. Elegans

hypersensitive to RNAi. Curr Biol. 2002;12(15):1317–9.

109. Porta-de-la-Riva M, Fontrodona L, Villanueva A, Cerón J. Basic Caenorhabditis

elegans methods: Synchronization and observation. J Vis Exp. 2012;(64):1–9.

110. Dickinson DJ, Ward JD, Reiner DJ, Goldstein B. Engineering the Caenorhabditis

elegans genome using Cas9-triggered homologous recombination. Nat Methods.

2013;10(10):1028–34.

111. Dickinson DJ, Goldstein B. CRISPR-based methods for caenorhabditis elegans

genome engineering. Genetics. 2016;202(3):885–901.

112. Dickinson DJ, Pani AM, Heppert JK, Higgins CD, Goldstein B. Streamlined

genome engineering with a self-excising drug selection cassette. Genetics.

2015;200(4):1035–49.

113. Grove C, Cain S, Chen WJ, Davis P, Harris T, Howe K, et al. Using WormBase, a

genome biology resource for Caenorhabditis elegans and related nematodes.

Methods Mol Biol. 2018;1757:399–470.

114. Sturm Á, Saskoi É, Tibor K, Weinhardt N, Vellai T. Highly efficient RNAi and

Cas9-based auto-cloning systems for C. elegans research. Nucleic Acids Res.

2018;46(17):e105.

115. Conte Jr. D, T. Macneil L, J. M. Walhout A, C. Mello C. RNA Interference in

Caenorhabditis Elegans. Current Protocol in Microbiology. 2015. 1–39 p.

116. Martinez M, Matus D. Auxin-mediated Protein Degradation in Caenorhabditis

elegans. Bio-Protocol. 2020;10(8).

117. Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, et al.

Fiji: An open-source platform for biological-image analysis. Nat Methods.

Smith, et al. (2021)

1328   2012;9(7):676–82.

1329

1330 SUPPLEMENTAL FIGURES

1331

1332 Figure S1. AC invasion is disrupted in temperature sensitive
SWI/SNF hypomorphs. Single planes of confocal z-stacks representing
AC invasion in *swsn-1(os22) and swsn-4(os13)* temperature sensitive
mutants with fluorescently labeled AC (magenta, *cdh-3>mCherry::moeABD*) and BM (green, *laminin::GFP*) scored at the
permissive temperature **(A)** and restrictive temperature **(B)**. Significant
loss of invasion was seen in both *swsn-1(os22)* (20% loss of invasion)
and *swsn-4(os13)* (24% loss of invasion) hypomorphic[ts] strains when
grown at the restrictive temperature 25°C and assessed at the P6.p 4-cell
1° VPC stage **(B)**.

A

1333

1334

Figure S2. Endogenous SWSN-4::GFP and SWSN-8::GFP are ubiquitously expressed. (A) Schematic (from http://wormweb.org/exonintron) depicting GFP insertion into the endogenous N and C termini of *swsn-4* (top) and *swsn-8* (bottom), respectively. Scale bar, 100 bp. **(B)** Expression of endogenously GFP-labeled SWI/SNF ATPase SWSN-4 protein and BM (*laminin::GFP*) in all larval stages (L1-L4), adult animals, and embryos. **(C)** Expression of endogenously GFP-labeled SWI/SNF ATPase SWSN-4 protein and BM (*laminin::GFP*) in all larval stages (L1-L4), adult animals, and embryos. Images in B-C are not to scale.

1335

1336

Figure S3. Fluorescent knock-ins express in the AC pre-, during, and post-invasion. Fluorescent micrographs depicting expression of SWSN-4::GFP **(A),** SWSN-8::**GFP (B)**, and PBRM-1::eGFP **(C)** in the AC, VU, and VPCs from the P6.p 1 cell to the P6.p 8 cell stages of development and corresponding quantifications. White arrowheads indicate AC, White brackets indicate 1° VPC stage. **(A'-C')** Quantification of endogenous GFP expression of SWI/SNF subunit in the AC, VU, and VPC over time. Statistical comparisons were made between the expression of each SWI/SNF subunit in the AC over time (magenta bracket and asterisks) or between the expression of each subunit in the AC relative to the expression of the same subunit in the neighboring VPCs or VUs at the same time (black asterisks) using Student's *t*-test (n≥30 for each stage and subunit; p values are displayed above compared groups). n.s. not significant.

1337

1338

Figure S4. Improved SWI/SNF RNAi significantly knocks down SWI/SNF **expression in the AC.** Fluorescent micrographs depicting BM (*laminin::GFP*) and expression of SWSN-4::GFP **(A)**, SWSN-8::GFP **(B)**, and PBRM-1::eGFP **(C)** in the AC in animals fed empty vector control (left) or RNAi targeting the endogenous allele (right). White arrowheads indicate AC(s), yellow arrowheads indicate boundaries of breach in BM, and white brackets indicate 1 VPCs. **(D)** Corresponding quantifications of fluorescent expression. Statistical comparisons were made between the expression of each SWI/SNF subunit in the AC in control and RNAi-treated animals using Student's *t*-test (n≥30 for each stage and subunit; p values are displayed above compared data). **(E)** Stacked bar chart showing percentage of AC invasion defects corresponding to each treatment, binned by AC phenotype (n≥30 animals per condition).

1339

1340

Figure S5. SWI/SNF subunits exhibit intra-complex and inter-assembly regulation.
(A-C) Representative fluorescence micrographs depicting endogenous GFP expression of individual SWI/SNF subunits representative of the core (*swsn-4*, **A**), BAF assembly (*swsn-8,* **B**), and PBAF assembly (*pbrm-1*, **C**) in the AC (*cdh-3p::mCherry::*moeABD) following treatment with targeting either SWI/SNF assembly **(A),** or the core ATPase and alternative SWI/SNF assembly **(B-C)**. White arrowheads indicate AC(s), yellow arrowheads indicate boundaries of breach in BM, and white brackets indicate 1 VPCs. **(D)** Quantification of fluorescence expression (mean gray value) of endogenous subunits in each condition. Statistical comparisons were made between the expression of each SWI/SNF subunit in the AC in control and RNAi-treated animals using Student's *t*-test (n≥30 for each stage and subunit; p values are displayed above compared data). n.s. not significant**. (E)** Quantification of fluorescence expression of endogenous GFP-tagged subunits of non-invasive ACs following loss of expression of alternative SWI/SNF subunits, binned per RNAi treatment by phenotype into single non-invasive AC (1AC) and mitotic non-invasive AC (2+ AC). Statistical comparisons (Student's *t*-test; p values are displayed above compared data) were limited to conditions with n>10 ACs in each phenotype. n.s. not significant. **(F)** Schematic summary of SWI/SNF core and assembly auto and cross regulation.

1341

1342

Figure S6. Improved *swsn-4* RNAi recapitulates SWI/SNF ATPase null phenotype in the sex myoblasts. (A) Single confocal z-planes depicting DIC (left) and expression of lineage-restricted CDK sensor (*unc-62>DHB::2xmKate2*, right) in the vulva and SM cells at the P6.p 8 cell stage corresponding to the stage when wild-type SM cells differentiate and exit the cell cycle. Animals were treated with empty vector control (top) or *swsn-4(RNAi)* (bottom). All representative images in each treatment are derived from the same z-stack from the same animal in the corresponding z-plane (top-left). Average or individual C/N CDK sensor ratios are listed in the bottom-right of corresponding panels. White arrowheads indicate individual SM cells. White brackets indicate 1° VPCs. **(B)** Quantification of the number of SM cells present at the P6.p 8 cell stage in control (black) and *swsn-4(RNAi)* treated animals (SMs arrive on both left and right side of vulva at the P6.p 8 cell, green; SMs arrive on either left or right side of vulva at the P6.p 8 cell stage, blue). **(C)** C/N CDK sensor ratios for SM cells in each treatment. Mean C/N ratio is represented by colored open circles and correspond to numbers of the same color. Gradient scale depicts cell cycle state as determined by quantification of each AC in all treatments (n≥30 animals per treatment), with dark-black depicting differentiation into G_0/G_1 and lighter-magenta depicting G_2 cell cycle states.

Figure S7. AC-specific expression of CKI-1 rescues invasion in BAF-depleted ACs. (A) DIC (left) and fluorescent (right) images depicting BM (*laminin::GFP*) and AC-specific CKI-1 (*cdh-3>CKI-1::GFP*) in empty vector control animals (top) and animals treated with *swsn-4(RNAi)* (middle) or *swsn-8(RNAi)* (bottom). **(B)** Stacked bar chart showing quantification of percentage of AC invasion defects corresponding to each treatment (n≥30 animals per condition, p values for Fisher's exact test comparing invasion penetrance in control animals and animals with the rescue transgene (+CKI-1::GFP) are displayed above black brackets).

1343

1344

1345

Figure S8. PBAF partially regulates the FOS-1 transcription factor. (A) Representative DIC (left) and fluorescent (right) micrographs depicting expression of endogenous GFP::FOS-1a and BM (*laminin::GFP*) in control (top) and *pbrm-1(RNAi)* treated (bottom) animals. **(B)** Quantification of GFP::FOS-1a expression in ACs of control and *pbrm-1(RNAi)* treated animals, normalized to mean expression of control group. Statistical comparisons were made between expression in the AC in control and RNAi-treated animals using Student's *t*-test (n≥20 for each condition; p value is displayed above black bracket). **(C)** DIC-Fluorescence overlay (left), and PBRM-1::mNG::AID and BM (LAM::GFP) (right), in animals treated with empty vector control (top) or *fos-1(RNAi)* (bottom). **(D)** Stacked bar chart showing percentage of AC invasion defects corresponding to each treatment and genetic background in C (n≥30 animals per condition, p values for Fisher's exact test comparing invasion defect penetrance in wild-type animals treated with *fos-1(RNAi)* and *pbrm-1::mNG::AID* animals treated with *fos-1(RNAi)* is displayed above black bracket). **(E)** Representative DIC (top-left), BM (LAM::GFP, top-right), AC (*cdh-3>PH*, bottom-left), and overlay (bottom-right) of P6.p 8 cell vulva in an MMP-deficient (-) animal treated with *pbrm-1(RNAi)*. **(F)** Stacked bar chart showing percentage of AC invasion defects corresponding to each treatment and genetic background in E (n≥30 animals per condition, p values for Fisher's exact test comparing invasion defect penetrance in wild-type animals treated with *pbrm-1(RNAi)* and MMP(-) animals treated with *pbrm-1(RNAi)* is displayed above black bracket).

SUPPLEMENTAL TABLES

Table S1. CRFs assessed for AC invasion contribution (see excel file)

270 chromatin regulating factors targeted by RNAi for AC invasion defects. n≥30 animals

for each RNAi clone. For each RNAi clone tested, the corresponding genetic sequence

name, public name, protein annotation, and human homolog (HUGO Gene

Nomenclature) from www.wormbase.com is given. Penetrance for each invasion defects

is given as the % of animals with ACs that fail to invade the BM at the P6.p 4 cell stage

out of the total number of animals assessed (Block/Invasion+Partial). Partial refers to

cases where an animal had a breach in the BM narrower than the width of the basolateral

surface of the invading AC. Genes in bold were recovered as significant regulators of AC

invasion (Table S2). Annotations were mined from the STRING consortium www.string-

db.org. Asterisks in human ortholog column denote genes with > 5 detected human

orthologs, for which only the first 5 returned orthologs were listed. N.A. denotes genes for

which no human ortholog exists. List is organized alphabetically based on genetic

sequence name.

Table S2. Significant regulators of AC invasion (see excel file)

41 chromatin and chromatin regulating factors (CRFs) identified as significant regulators

of AC invasion. For each RNAi clone listed, the corresponding genetic sequence name,

public name, and human homolog is listed. AC invasion scoring data is provided for each

clone at the P6.p 4 cell stage. Genes were determined to be significant AC invasion

regulators if RNAi targeting resulted in ≥ 20% loss of invasion at the P6.p 4-cell stage

(n≥30 animals). Genes in bold are components of the SWI/SNF complex. Asterisks

denote genes previously published to regulate *C. elegans* AC invasion. N.A. denotes

genes for which no human ortholog exists. List is organized alphabetically based on

genetic sequence name.

Table S3. Enhanced (T444T) RNAi vectors used in this study (see excel file)

Table S4. Strains used in this study (see excel file)

Table S5. CRISPR reagents (see excel file)

1375