Capturing the interplay of membrane lipids and structural transitions in human ABCA7

Le Thi My Le[1]†, James R. Thompson[1]†, Sepehr Dehghani-Ghahnaviyeh[2], Shashank Pant[2,#], Phuoc

X. Dang[1], Takahisa Kanikeyo[3], Emad Tajkhorshid[2] & Amer Alam[1,*]

[1]The Hormel Institute, University of Minnesota, Austin, Minnesota 59912

[2] Theoretical and Computational Biophysics Group, NIH Center for Macromolecular

Modeling and Bioinformatics, Beckman Institute for Advanced Science and Technology,

Department of Biochemistry, and Center for Biophysics and Quantitative Biology,

University of Illinois at Urbana-Champaign, Urbana, IL, USA

[3]Department of Neuroscience, Mayo Clinic, Jacksonville, Florida 32224

†These authors contributed equally to this work

Current affiliation: Loxo Oncology at Lilly., Louisville, CO 80027, USA

*Address correspondence to: Amer Alam, PhD, aalam@umn.edu

Abstract

Phospholipid extrusion by ABC subfamily A (ABCA) exporters is central to cellular physiology,

although the specifics of the underlying substrate interactions and transport mechanisms remain

poorly resolved at the molecular level. Here we report cryo-EM structures of lipid-embedded

human ABCA7 in an open state and a nucleotide-bound, closed state at resolutions between 3.6-

4.0 Å. The former reveals an ordered patch of bilayer lipids traversing the transmembrane domain

(TMD), while the latter reveals a lipid-free, closed TMD with a small extracellular opening. These

structures offer a structural framework for both substrate entry and exit from the ABCA7 TMD

and highlight conserved rigid-body motions that underlie the associated conformational

transitions. Combined with functional analysis and molecular dynamics (MD) simulations, our

data also shed light on lipid partitioning into the ABCA7 TMD and localized membrane

perturbations that underlie ABCA7 function and have broader implications for other ABCA family

transporters.

Introduction

ABCA family exporters mediate efflux of phospholipids and sterols from cells,

contributing to membrane homeostasis, bilayer structure and asymmetry, and the formation of

serum lipoproteins, among other key physiological processes[1]. Their dysfunction therefore

underlies several human diseases[2-4]. The molecular details governing the ABCA exporter substrate

transport cycle are not fully resolved. To fill this knowledge gap, here we present the structural

and functional analysis of human ABCA7, whose dysfunction has been strongly linked to

Alzheimer's Disease (AD)[5-12], in a lipid environment. Deficient ABCA7 activity leads to

alterations in both brain lipid profiles and fatty acid and phospholipid biosynthetic pathways[13],

impaired memory, and reduced immune responses[14,15]. Both *in vitro* lipid flipping[16] and lipid

extrusion to apolipoproteins by cells over-expressing ABCA7[17] have been demonstrated, although

the correlation between the two processes, if any, remains unclear. To date, no direct structural

information exists for ABCA7. Understanding the molecular details of the ABCA7 transport cycle

and how its dysfunction alters inflammatory and immune responses, lipid homeostasis, and

phagocytosis, which all contribute to AD progression[18-22], may therefore pave the way for novel

therapeutics for AD.

ABCA7 encodes a 2146 amino acid membrane transporter found in many tissues and

blood, hippocampal neurons, macrophages, and microglia[23,24]. Like the phospholipid and sterol

exporter ABCA1 and retinal importer ABCA4, with which it shares 54% and 59% sequence

similarity, respectively, ABCA7 comprises two halves assembled as a full transporter. Each half

consists of a TMD, with the first two transmembrane helices (TMs) of each separated by a large

extracellular domain (ECD), and a nucleotide binding domain (NBD) attached to a cytoplasmic

regulatory domain (RD). To visualize its ATP-dependent conformational cycle in a lipid

environment, we resolved the structures of human ABCA7 in multiple conformations in lipid and

detergent environments using cryo-EM and probed its lipid interactions using ATPase assays and

MD simulations. Our data allow us to directly visualize lipid partitioning into the TMDs and the

associated conformational changes in ABCA7 that provide insights into its mechanisms of

substrate entry and export that likely hold true for other members of the ABCA family.

Results

Dependence of ABCA7 ATPase activity on its lipid environment

Human ABCA7 expressed in a tetracycline inducible stable HEK293 cell line was

reconstituted in liposomes and nanodiscs comprising 80% brain polar lipids (BPL) and 20%

cholesterol (Chol) and its ATPase activity was measured (Figure 1A, Figure S1A). Although ATP

hydrolysis was slowest in nanodiscs, it followed Michaelis-Menten kinetics similar to ABCA7 in

detergent or liposomes comprising the same lipid/cholesterol composition and Michaelis constant

(K_M) values for all three were in the 0.5-0.8 mM range. ATPase rates for a hydrolysis-deficient

mutant carrying E965Q and E1951Q substitutions (ABCA7$_{EQ}$) were drastically reduced compared

to wildtype in both nanodiscs and detergents, demonstrating that the observed activity was specific

(Figure S1A).

To analyze the effect of different lipid headgroups on ATPase activity, we compared

ATPase rates of ABCA7 in BPL/Chol nanodiscs to those in which BPL was replaced by either

brain polar phosphatidylethanolamine (PE), phosphatidylserine (PS) or phosphatidylcholine (PC)

as well 100% BPL nanodiscs. As seen in Figure 1B, ATPase rates were highest in PE and PS

nanodiscs followed by 100% BPL nanodiscs lacking cholesterol and, lastly, PC nanodiscs that

showed a marginal increase. These results largely match previous findings[16], although differences

exist in extent of ATPase rate stimulation that are likely due to the divergent lipid formulations

used for reconstitution. Overall, they demonstrate that lipids modulate ABCA7 activity in a species

dependent manner and that cholesterol has an inhibitory effect on ATPase activity of ABCA7.

Cryo-EM structures of ABCA7 in nanodiscs reveal an asymmetric, open TMD

To visualize ABCA7 in a lipid environment, we determined its cryo-EM structures in

BPL/Chol nanodiscs (ABCA7$_{BPL}$) at 3.6 Å resolution (Figure 1C, Figure S1B-E). Despite the

addition of the non-hydrolysable ATP analog adenosine-5′-o-(3-thio-triphosphate) (ATPγS), the

transporter adopted an open conformation with separated NBDs and a wide open TMD pathway.

We observed density features consistent with a patch of ordered lipids from both bilayer leaflets

traversing the width of TMD as expanded upon below. A second, higher-resolution 3D class

displayed more complete density for the ECD tunnel region but comparatively lower quality of

EM density for TMD2-NBD2 and the entire RD, indicating greater conformational disorder, which

was therefore not analyzed further (Figure S1D-E). Analysis of TMD-ECD interfaces in the

ABCA7$_{BPL}$ structures revealed more extensive contacts between the ECD and TMD1 (buried

surface area (BSA) of 790 Å^2) compared to TMD2 (330 Å^2 BSA). Both TMD1 and TMD2 made

contacts with the opposite ECD subunits, leading to a domain swapped arrangement. The TMD1-

ECD2 interface (BSA 565 Å^2) was significantly larger than that of TMD1-ECD1 (BSA 223 Å^2)

and both interfaces comprised an extensive network of polar and electrostatic interactions. The RD

adopted a domain-swapped arrangement similar to that in ABCA4 structures[25], with RD1 and RD2

associated with the opposite NBDs. To see whether structural changes could rationalize the

enhanced ATPase activity observed in ABCA7$_{PE}$ compared to ABCA7$_{BPL}$, we also determined the

cryo-EM structure of the former to 4.0 Å resolution (Figure 1D, Figure S2). Interestingly, both

structures were nearly identical to each other (r.m.s.d 0.31Å), although only a single high

resolution 3D class was resolved for ABCA7$_{PE}$ (Figure S2). Compared to ABCA7$_{BPL}$, density for

TMD lipids was more homogenous , despite its comparatively lower resolution, as described

below, indicating overall decreased conformational heterogeneity. For both ABCA7$_{BPL}$ and

ABCA7$_{BPE}$, greater positional disorder was observed for TMD2-NBD2 compared to TMD1 and

NBD1 as indicated by relatively weaker EM density quality (Figure S3).

The ABCA7 TMD lumen is accessible to an ordered file of bilayer lipids

Our ABCA7$_{BPL/PE}$ structures are distinguished from available structures of ABCA1 and

ABCA4 (all resolved in a detergent environment) by the wider open TMD lumen that is almost

completely occupied by lipids. The observed lipids are continuous with the surrounding membrane

except at the cytoplasmic leaflet near residues L655 and T1646 from TM5 and TM11, respectively

(Figure 1E-F). EM density for the modeled lipids is recognizable by gaps between the extracellular

and cytoplasmic leaflets and the two acyl chains of a single file of phospholipids in the two lipid

leaflets. Towards the extracellular end, ECD residues R475, K478, R482, K1407 and TMD1

residues R544 and R548 form a cluster of positively charged side chains oriented towards the

luminal lipids (Figure 1G). Overall, the luminal lipids are more closely associated with TMD1,

with residues from TM2, TM5, and TM11 within 5 Å of the modeled acyl chains compared to only

residues from TM11 from TMD2.

To compare the architecture of ABCA7 in the presence or absence of lipids, we also

determined its structure in the detergent digitonin (Figure 2A-B, Figure S4A-E). This

$ABCA7_{DIGITONIN}$ structure revealed a similar overall conformation to that seen in structures of

ABCA1[26] and ABCA4[25,27,28] (Figure S4F) with a more complete map for the ECD compared to

$ABCA7_{BPL/PE}$. As expected, no EM density for TMD lipids was visible, and only isolated density

features attributable to detergent molecules associated with TMD1 were observed, although only

at a contour level below where much of the surrounding detergent micelle was visible (Figure

2A,C). This is in contrast to $ABCA7_{BPL/PE}$ structures where TMD lipid density was much stronger

than that of the bulk nanodisc (Figure 2D-E), further validating our assignment of bilayer lipids.

Compared to structures in nanodiscs, the TMDs in this $ABCA7_{DIGITONIN}$ structure were arranged

more symmetrically with respect to each other with a narrower lumen. Interestingly, EM density

for TMD2 was still weaker than that of TMD1, in contrast to structures of ABCA4[25,27,28] and

ABCA1[26] (Figure S4D) indicating that a more mobile TMD2-NBD2 pair may be a unique feature

of ABCA7.

127   Closed TMD lumen and exit pocket for lipid extrusion in ATP bound ABCA7

To gain insight into possible mechanisms of lipid extrusion from the ABCA7 TMD, we

used the hydrolysis-deficient $ABCA7_{EQ}$ mutant reconstituted in BPL/Chol nanodiscs and

determined its structure in a closed, ATP-bound state at 3.7 Å resolution (Figure 3A and Figure

S5). This $ABCA7_{EQ-ATP}$ structure revealed closely interacting TMD-NBD pairs and a largely

occluded TMD lumen (Figure 3B). However, at the extracellular end of the TMD lumen, we

observed a small central opening to the extracellular space/ECD that may represent an 'exit pocket'

(Figure 3C,D) akin to that seen in the yeast pleiotropic drug resistance transporter Pdr5[29] that could

likely accommodate two acyl chains, (Figure 3D). TMD closure involves formation of a 4-TM

bundle comprising TMs 2, 5, 8, and 11 that occludes the cytoplasmic bilayer leaflet. The exit

pocket is comprised largely of hydrophobic residues, except for R548, which is part of a cluster of

positively charged residues including R475, K478, R482 and R678 identified in our ABCA7$_{BPL/PE}$

structures that may aid in directing a negatively charged phospholipid headgroup towards the ECD.

Side chain density for this group of basic residues was poor compared to that in ABCA7$_{BPL/BPE}$

structures, indicating greater disorder. TMD closure is accompanied by a rearrangement of both

TMD-ECD interfaces compared to the open state structures (Figure S6). Compared to the

ABCA7$_{BPL}$ structure, both TMD1 and TMD2 interfaces with ECD1 increased from BSAs of 190

Å^2 to 331 Å^2 and from 76 Å^2 to 285 Å^2, respectively. Conversely, the BSAs of TMD1 and TMD2

with ECD2 decreased from 543 Å^2 to 275 Å^2 and from 317 Å^2 to 192 Å^2, respectively. As

expected, the ABCA7$_{EQ-ATP}$ structure shows a canonical NBD sandwich dimer with bound ATP

(Figure 3E). In contrast to the ATP bound ABCA4 structure, RD density was too weak for accurate

placement. However, contoured at lower thresholds, the available density features were more

compatible with a rigid body shift rather than a conformational rearrangement of RD, with RD2

appearing to maintain contact with NBD1, but RD1 disengaged from NBD2.

Rigid-body motions of the TMDs define ABCA7 conformational transitions

Despite significant overall conformational differences, the individual TMDs in

ABCA7$_{BPL/PE,}$ ABCA7$_{DIGITONIN}$, and ABCA7$_{EQ-ATP}$ remain largely unaltered (Figure 4A).

Moreover, the TMD-NBD pairs move as rigid body units from the fully open ABCA7$_{BPL}$ structure

to that of ABCA7$_{DIGITONIN}$, which we assert is akin to an intermediate open state between the fully

open ABCA7$_{BPL/PE}$ state and the closed ABCA7$_{EQ-ATP}$ states. The transition from open to

157 intermediate open involves a rigid-body rotation of 9° and translation of 2 Å for TMD2-NBD2.

158 The transition to the closed state involves a further 6° rotation and 15 Å translation of TMD2-

159 NBD2 (Figure 4B-C), albeit with a greater alteration in the NBDs owing to a movement of their

160 recA like domains upon ATP binding and dimerization. Rigid body movements were also observed

161 for the entire RD in all three structures. The RD of ABCA7$_{DIGITONIN}$ maintained molecular

162 interactions with both NBD1 (RD2) and NBD2 (RD1), whereas the RD of ABCA7$_{BPL/PE}$ separated

163 from NBD2 while maintaining contact with NBD1. Finally, the ECD base region transitioned

164 through a rigid body rotation of 34° and translation of 6.3 Å going from the open to closed states,

165 whereas the tunnel and lid regions displayed greater heterogeneity.

166  To establish whether the concerted TMD-NBD rigid body motions outlined above may

167 extend to other ABC transporters of the Type-V ABC transporter/Type II ABC exporter fold[30], we

168 extended our analysis of individual TMD-NBD pairs to members of the G family for which the

169 first structures of open and closed states were available, namely ABCG2 and ABCG5/G8 (Figure

170 S7A)[31,32]. As shown in Figure S7B, the TMD-NBD pairs from both the apo and ATP-bound closed

171 states of ABCG2, as well as from both halves of ABCG5/G8, all shared a similar overall

172 architecture. Despite the divergent topologies of ABCG and ABCA transporters, with the former

173 are arranged in an NBD-TMD configuration compared to TMD-NBD for the latter, the individual

174 TMD-NBD pairs from ABCA7 shared very strong structural similarities with those of ABCG2,

175 further establishing the role of rigid body movements of the TMDs to affect large scale

176 conformational changes in these transporters.

177 **Lipid partitioning in the ABCA7 TMD captured with MD simulations**

178   To gain additional molecular insight into the TMD-mediated lipid partitioning, membrane

179 perturbation, and lipid extrusion from ABCA7, we performed multi-microsecond MD simulations

180 of the open conformation of ABCA7$_{PE}$ after embedding into two distinct lipid bilayers, one

181 containing PE/Chol (4:1) and the other PC/Chol (4:1) (Figure 5, Figure S8). Each simulation was

182 performed for 2 μs using a system including four copies of the protein (Figure 5A), resulting in an

183 aggregate 8 μs sampling of lipid-protein interactions for each lipid composition. Interestingly,

184 during the simulations, we observed PE and PC phospholipids penetrating the TMD cavity in both

185 the cytoplasmic and extracellular leaflets in all four protein replicas (Figure 5A, insets). We

186 quantified the number of phospholipids within the TMD cavity by counting the lipid headgroups

187 (Figure S8C), revealing that on average, ABCA7 accommodated a larger number of PE lipids

188 compared to PC (Figure 5B). To further quantify ABCA7-mediated deformation of the lipid

189 bilayer bridging the TMDs, the average heights of phospholipid headgroups were calculated with

190 respect to the midplane of the bilayer. We captured both a tendency for lipids to move towards the

191 cytoplasmic gate and the formation of a dome-shaped phospholipid arrangement within the TMD

192 lumen (Figure 5C-E, Figure S8A-B). Interestingly, our MD simulations showed an accumulation

193 of phospholipid headgroups in the vicinity of residues R475, K478, R482, R544, and R548, with

194 R482 and R548 displaying the most frequent lipid contacts (Figure S8D-E), corroborating our

195 structural observations. Analysis of the cytoplasmic leaflet highlighted a continuous distribution

196 of phospholipids with the surrounding membrane, except near TM5 and TM11 around residues

197 L655 and T1646 (Figure 5D-E), which correlated well with the bilayer-like density in the

198 ABCA7$_{BPL}$ and ABCA7$_{PE}$ cryo-EM maps (Figure 1C-D) that we modeled with acyl chains.

199 **Discussion**

A mechanistic model derived from our results is shown in Figure 6. We establish that the

TMD lumen of ABCA7 is accessible to bilayer lipids in the open state (ABCA7$_{BPL/PE}$), providing

a basis for substrate entry from both the extracellular and cytoplasmic leaflets. These ordered lipids

are akin to those observed in the TMD cavity of AcrB, where they have been suggested to be

important for functional integrity[33]. We identify a network of positively charged residues at the

extracellular periphery of TMD1 and the base of the ECD directed towards the TMD luminal

lipids. Our MD simulations support a role for these residues in interacting with phospholipid

headgroups, consistent with our structural data. In ABCA7, R475 mutations have been identified

in AD[34] patients, and mutation of R638 in ABCA1 (equivalent to R548 in ABCA7) is associated

with reduced serum HDL[35] (Figures S9 and S10). While further studies are needed to pinpoint the

specific roles, if any of these residues, in ABCA7 function, they remain largely conserved in

ABCA1, ABCA4, and ABCA7. The upward protrusion of both bilayer leaflets within the TMD

lumen suggested by MD simulations may, upon TMD closure, aid in sequestering phospholipids

in the identified exit pocket to be primed for extrusion towards the ECD. Membrane deformations

similar to those highlighted here have also been observed in the signal peptidase complex[36], and

further evidence the influence membrane proteins have on local membrane bilayer structure.

ABCA7 could thus play a role in changing its membrane lipid environment by flipping

cytoplasmic leaflet lipids to the extracellular leaflet and also extruding them back out into the bulk

bilayer environment and/or to apolipoproteins. The reported defects in the phagocytotic activity of

cells that express dysfunctional ABCA7[37,38] could be related to ABCA7's influence on the

asymmetric distribution of bilayer lipids[39], as enrichment of PS, shown to be flipped by ABCA7[16],

at the extracellular surface is linked to phagocytosis and phagocytosis-associated proteins within

the membrane[40-42].

Despite significant progress in our mechanistic understanding of ABCA transporter

function, several questions remain open. First, it is unclear to what extent ABCA7 interacts with

apolipoproteins in a physiologically relevant manner and, more broadly, the structural basis for

apolipoprotein interactions with ABCA7 or ABCA1 is unknown. Second, while our structures

provide a potential basis for lipid entry into and extrusion from the ABCA7 TMD, the mechanism

whereby lipids get flipped, remains unresolved. Third, while our data point to a potential

preference for PE over PC for TMD partitioning, the exact mechanism whereby ABCA1 or

ABCA7 achieve lipid specificity are unknown. The observed enhancement in conformational

homogeneity, quality of lipid density, and ATPase activity of our $ABCA7_{PE}$ sample may be, in

part, due to the smaller PE headgroup, which has been shown to aid in folding and stabilization of

membrane proteins[43]. Finally, it is unknown how the opposite direction of substrate transport for

ABCA4 is achieved considering similarities in ATP bound closed and open state structures of

ABCA4 and ABCA7. Overall, our data will help devise better *in vitro* and *in silico* models to

answer these questions, which will further aid in dissecting the unique roles these proteins play in

cellular physiology.

METHODS

METHODS

Protein Purification

We utilized the Flp-In TREX system (Thermo Fisher Scientific) for tetracycline inducible

expression of human ABCA7. In short, a codon optimized synthetic gene construct

(GeneArt/Thermo Fisher Scientific) of isoform 1 of ABCA7 (Uniprot ID Q8IZY2-1), harboring a

C-terminal eYFP-Rho1D4 tag[43] with a 3C/precision protease site between the protein and

purification tags, was cloned into a PCDNA5.1 FRT/TO vector between BamHI and NotI

restriction sites and a stable cell line was generated as per manufacturer's protocol (Flp-In™ T-

Rex™ Core Kit, Thermo Fisher Scientific). The resulting HEK293 based stable cells were grown

and maintained in adherent cell culture in Dulbecco's Modified Eagle Medium (DMEM, Thermo

Fisher Scientific) supplemented with 9% Fetal Bovine Serum (FBS, Gibco) and a

penicillin/streptomycin mixture (Thermo Fisher Scientific) at 37°C with 5% carbon dioxide (CO_2)

under humidified conditions. For protein production, cells were induced with 0.6 μg ml^{-1}

tetracycline at a confluency of 80% in fresh DMEM supplemented with 2% FBS under otherwise

identical conditions for an additional 72 hours before being washed with Phosphate Buffered

Saline (PBS), harvested, and flash frozen in liquid nitrogen.

For purification, thawed cells were resuspended in a lysis buffer (Buffer L) comprising 25

mM Hepes pH 7.5, 150 mM sodium chloride (NaCl), 20% glycerol, 1 cOmplete EDTA free

protease inhibitor tablet (Roche) per 50 ml Buffer L, 800 μM phenylmethylsulfonyl fluoride

(PMSF) and 20 μg ml^{-1} soybean trypsin inhibitor (both Sigma), and mechanically cracked using

a dounce homogenizer before addition of a 0.5%/0.1% w:v mixture of dodecyl maltoside (DDM)

and cholesteryl hemisuccinate (CHS) (both Anatrace). Protein extraction was allowed to proceed

for 90 minutes at 4°C with gentle agitation, after which, the suspension was centrifuged at 48,000

r.c.f for 30 minutes and the supernatant applied to rho-1D4 antibody (University of British

Columbia) coupled Sepharose resin (Cytiva). Binding was allowed to proceed for 3 hours before

the unbound fraction was discarded and beads rinsed with 4 x 10 bed volumes (BVs) of wash

buffer (25 mM Hepes pH 7.5, 150 mM NaCl, 20% glycerol, 0.02%/0.004% w:v DDM/CHS).

Protein was eluted by incubation with 3 BVs elution buffer (wash buffer supplemented with either

3C protease (1:10 w:w 3C:ABCA7) or 0.5 mg ml^{-1} 1D4 peptide (GenScript)) for 2-18 hours.

The EQ variant of ABCA7 contained two site mutations, E965Q and E1951Q. The E965Q

site was created within the ABCA7 construct using site directed mutagenesis by PCR with the

primers: A7eq1for 5'-GGTCATCCTGGATCAACCTACAGCAGGCGTGG-3' and A7eq1rev

5'-GCCTGCTGTAGGTTGATCCAGGATGACCACC-3'. E1951Q was generated using a

synthesized dsDNA block and the enzymes NheI and BsiWI. ABCA7$_{EQ}$ with C-terminal eYFP-

Rho1D4 tag was then transferred to a pCAG vector using KpnI and NotI restriction sites. The

HEK293T cells were grown in Dulbecco's Modified Eagle Medium (DMEM, Scientific)

supplemented with 9% FBS (Gibco), penicillin/streptomycin mixture (Scientific) and antimycotic

(Gibco) at 37°C and 5% CO_2 under humidified conditions. A mixture of 37.5 µg of ABCA7$_{EQ}$

plasmid and 75 µg of polyethyleneimine (PEI, Sigma) was incubated for 15 min at room

temperature before being applied to 15 ml plates of HEK293T cells at a confluency of 60 - 80%

to initiate transfection and expression. The cells were further cultured for 72 hours before being

washed with PBS, harvested and flash frozen in liquid nitrogen. ABCA7$_{EQ}$ was purified in the

same approach of ABCA7; and eluted by incubation with 3 BVs elution buffer with either 3C

protease for ATPase assay or with 0.25 mg ml^{-1} of 1D4 peptide in an additional 2 mM ATP and

10 mM $MgCl_2$ for nanodisc reconstitution.

Membrane scaffold protein D1 (MSP1D1, addgene) and apoA1 were purified using

established protocols for MSP[44] with the following modifications: A synthetic construct of apoA1

bearing a 3C protease cleavable N-terminal deca-histidine tag (GeneArt/Thermo Fisher Scientific)

was cloned into a pET28a vector (Addgene) and transformed in *E. coli* BL21 DE3 cells (New

England Biolabs). One-liter cultures of Terrific Broth (TB) supplemented with 50 ug ml^{-1}

kanamycin were grown from 10 ml overnight cultures from single colonies grown in LB. Cells

were grown to an OD600 of 0.8 in a shaking incubator at 37 °C and induced with 1 mM isopropyl

β-d-1-thiogalactopyranoside (IPTG). Protein expression was allowed to proceed at 20°C for 12

hours. Cells were centrifuged at 12,000 r.c.f, and pellets were flash frozen in liquid nitrogen and

stored at -80°C until required. Frozen pellets were resuspended in 8 ml/ gram cell pellet

resuspension buffer comprising 25 mM Hepes pH 7.5, 150 mM NaCl and 1 mM

phenylmethylsulfonyl fluoride (PMSF) and sonicated. The suspension was spun down at 16,000

r.c.f at 4 °C for 30 min and the supernatant was applied 5 ml Ni-NTA resin (Qiagen)/ L culture

medium. After discarding the flowthrough, the resin was washed with 25 mM Hepes pH 7.5,

150 mM NaCl, 1 mM PMSF and 20 mM imidazole until a pre-established baseline A280 reading

was achieved. ApoA1 was eluted in 4 BVs of 25 mM Hepes pH 7.5, 150 mM NaCl, 1 mM PMSF

and 200 mM imidazole, concentrated using a 10 kDa molecular weight cutoff (MWCO) Amicon

filter (Millipore-Sigma) and desalted using a PD10 column (Cytiva) into 25 mM Hepes pH 7.5,

150 mM NaCl. The concentration of apoA1 was adjusted to 1 mg ml^{-1} for flash freezing in liquid

nitrogen and storage at -80°C.

ABCA7 and ABCA7$_{EQ}$ nanodisc and proteoliposome preparation

For nanodisc reconstitution, peptide eluted or 3C cleaved ABCA7 was mixed with

MSP1D1 and a mixture of BPL (brain polar lipid extract from Avanti) and cholesterol (80:20 w:w)

with 0.5%/0.1% DDM:CHS using a 1:10:350 (ABCA7:MSPD1:lipid mix) molar ratio in nanodisc

buffer (25 mM Hepes pH 7.5, 150 mM NaCl) that contained up to 4% glycerol for 30 minutes at

room temperature (RT). Nanodisc reconstitution was induced by removing detergent with 0.8 mg

ml^{-1} pre-washed Biobeads SM-2 (Bio-Rad) for 2 hours with gentle agitation at RT. For different

phospholipid compositions, BPL was replaced with brain PE, PS, or PS (all from Avanti Polar

Lipids). For structural studies, nanodisc-reconstituted ABCA7 bearing the eYFP-Rho1D4 tag was

bound to rho-1D4 resin for an additional 2 hours, washed with 4 BV of nanodisc buffer, and eluted

with 3C protease for 2 hours at 4°C. The eluted ABCA7 nanodiscs were concentrated using a 100,000 MWCO kDa Amicon filter and further purified by size exclusion chromatography using a G4000swxl column (TOSOH biosciences) equilibrated with nanodisc buffer at 4°C. Figure S1A shows a SEC chromatogram for pure ABCA7 placed into BPL/Ch nanodiscs, while a SEC chromatogram for ABCA7 in PE/Ch nanodiscs is in Figure S2A. Generally, three fractions were pooled from the main resultant peak.

ABCA7$_{EQ}$ was reconstituted in nanodiscs for cryo-EM preparation using the same approach of ABCA7; except that an additional 2 mM ATP and 10 mM MgCl$_2$ were present until the end of the purification procedure prior to grid preparation. A SEC chromatogram for ABCA7$_{EQ}$ in BPL/Ch nanodiscs is shown in Figure S5A, where the trace is affected by the additional ATP added during the run.

ABCA7 proteoliposomes were generated by mixing detergent purified ABCA7 with liposomes at a protein: liposome ratio of 1:10 w:w. Liposomes were prepared by extruding a 20 mg ml^{-1}, 80:20 w:w BPL/Ch lipid mixture 11 times using a previously described protocol [45]. Briefly, detergent purified ABCA7 and liposomes were added to 0.14% and 0.3% Triton X100 (Sigma), respectively, then incubated for 30 minutes at RT. These two samples were mixed and incubated for 60 minutes. Detergent was removed by adding 40 mg fresh Biobeads SM2 (Bio-Rad) per ml reaction mixture during five successive incubation steps, 30 minutes at RT, 60 minutes at 4°C, overnight at 4°C, and two periods of 60 minutes at 4°C with gentle agitation. The suspension was centrifuged at 80,000 rpm for 20 minutes in an ultracentrifuge. The supernatant was removed, and the liposomal pellet was washed once with reconstitution buffer containing 150 mM NaCl, 25 mM Hepes pH 7.5. The ABCA7-liposome suspension was then centrifuged to

336   remove the supernatant, and the proteoliposomes were resuspended at a final concentration of

337   0.5 - 1 mg ml^{-1} for ATPase assays.

ABCA7$_{DIGITONIN}$ preparation

For the digitonin solubilized ABCA7 purification, ABCA7 was extracted from the cell in

a lysis buffer (Buffer L) using the same approach above, and the supernatant was applied to rho-

1D4 resin for a 3-hour binding period. Then, the resin was rinsed with the 4 x 10 BVs of wash

buffer containing 25 mM Hepes pH 7.5, 150 mM NaCl, 20% glycerol (v/v), and 0.06% digitonin

(w/v). Protein was eluted by incubation with 3 BVs elution buffer, which was wash buffer

supplemented with 3C protease (1:10 w:w 3C:ABCA7). Interestingly we obtained better particle

distribution and ice quality with addition of a 1:2.5 molar excess of apoA1, prepared in house, to

3C cleaved ABCA7 prior to grid preparation. The mixture was concentrated by a 100,000 MWCO

kDa Amicon filter (Millipore) and further purified by size exclusion chromatography using a

G4000swxl column (TOSHOH biosciences) equilibrated with a buffer containing 25 mM Hepes

pH 7.5, 150 mM NaCl, 0.035% digitonin (w/v), as shown in Figure S4A. Peak fractions were

pooled and concentrated for cryo-EM grid preparation.

ATPase assays

ATPase assays were based on a molybdate based colorimetric assay[46]. Protein

concentrations used were in the range of 0.05-0.1 mg ml^{-1}. Assays were started by the addition of

either 2mM ATP, except for experiments in Figure 1B where 6.25mM ATP was used, in the

presence of 10 mM magnesium chloride (MgCl$_2$), incubated for 30 minutes at 37°C, then stopped

by addition of 6% SDS. The assay was also performed in the presence of ABCA7 inhibitors as

additives, such as 5 mM ATPγS or sodium orthovanadate. For ATP K$_M$ measurements, a range of

ATP concentrations was used. Statistical analysis was done using GraphPad Prism 9. ATPase rates

were measured using simple linear regression, and the K_M of detergent, liposome, and nanodisc

reconstituted ABCA7 were determined from the fit to the Michaelis-Menten equation of the

corresponding rates. ABCA7 concentrations were measured using gel densitometry analyzed in

ImageStudio Lite (LI-COR Biosciences) based on detergent purified ABCA7 standards with

known concentrations determined by A280 measurements. All reaction components were mixed

with ABCA7 in detergent or reconstituted in nanodiscs and liposome in the absence of ATP,

incubated for 10 minutes at 37 °C prior to addition of ATP to start the reaction.

Cryo-electron microscopy grid preparation

For ABCA7 reconstituted in nanodisc prepared in 80:20 w:w BPL/cholesterol and

PE/cholesterol, SEC purified protein was mixed with 5 mM ATPγS (TOCRIS) and 5 mM $MgCl_2$

for 20 minutes at room temperature and concentrated to 0.5 - 1.0 mg ml^{-1}. Peak fractions from SEC

already containing 2 mM ATP (Sigma) and 10 mM $MgCl_2$ were pooled and concentrated to 0.5 -

1 mg ml^{-1} for nanodisc reconstituted ABCA7$_{EQ-ATP}$. For ABCA7 in digitonin, peak fractions were

pooled and concentrated between 2 to 5 mg ml^{-1}. 4 µl samples were applied to glow discharged

Quantifoil R1.2/1.3 grids (Electron Microscopy Sciences, Hatfield, PA, USA) using a Vitrobot

Mark IV (Thermo Fisher Scientific) with a 4s blotting time and 0 blotting force under >90%

humidity at 4°C, then plunge frozen into liquid ethane. For the nanodisc reconstituted ABCA7$_{EQ-}$

$_{ATP}$ and ABCA7 in digitonin, two sample droplets were applied to glow discharged grids to obtain

more particles per hole.

Cryo-electron microscopy data collection and processing

Grids were clipped as per manufacturer guidelines and cryo-EM data was collected using

a Titan Krios electron microscope operating at 300kV and equipped with a Falcon 3EC direct

electron detector (Thermo Fisher Scientific.). Automated data collection was carried out using

EPU 2.8.0.1256REL software package (Thermo Fisher Scientific) over multiple sessions in

counting mode at a nominal magnification of 96,000x, corresponding to a calibrated pixel size of

0.895 Å for nanodisc reconstituted $ABCA7_{BPL}$. Image stacks comprising 60 frames were collected

at a defocus range of -0.6 to -2.6 µm and estimated dose rate of 1 electron/$Å^2$/frame and further

processed in Relion-3.1 (beta). Motion correction was done using Motioncor2 (Relion

implementation)[47] and contrast transfer function (CTF) correction was performed using Gctf 1.06

[48]. A summary of the overall data processing scheme and the quality was presented in Figure S1C-

E. In brief, 11802 micrographs were used for template free picking of 6725108 particles, followed

by particle extraction at a 3x binned pixel size of 2.685 Å/pix. The dataset was processed in two

batches. After 2-3 rounds of 2D classification 1259324 particles from Set 1 and 1088487 particles

from Set 2 were selected for independent 3D classification steps (number of classes (K)=8 for

both). The structure of human ABCA1 (EMDB6724) was used as a 3D reference for an initial 3D

classification of a subset of the total data to yield an initial sub-nanometer resolution map of

ABCA7 that was used as a 3D reference for the full datasets. After 1 round of 3D classification,

both sets of data yielded a similar ensemble of classes. A total of 113291 particles from similar

looking classes (black boxes) were subjected to an additional round of classification (K=3), ~80%

of which fell into a high-resolution class that yielded a 3.6 Å map after refinement and particle

polishing steps. Similarly, 124114 particles from a second set of two similar classes (red boxes in

Figure S1D) were selected for subsequent refinement, particle polishing, and post processing to

yield a 3.1 Å map. All resolution estimates were based on the gold standard 0.143 cutoff criterion

[49]. Datasets of $ABCA7_{PE}$, $ABCA7_{EQ-ATP,}$ and $ABCA7_{DIGITONIN}$ were collected at a nominal

magnification of 96,000x corresponding to a calibrated pixel size 0.889 Å, and image stacks

containing 40 frames were collected a defocus range of -0.8 to -2.6 µm with an estimated dose rate

of 1 electron/Å^2/frame and further processed using the same software versions as for the

ABCA7$_{\text{BPL}}$ dataset unless otherwise indicated.

For ABCA7$_{\text{PE}}$, a total of 2849251 particles were picked from 9218 ctf corrected (Gctf)

micrographs in Relion ver. 3.1 in two batches at a 3-fold binned pixel size of 2.667 Å/pixel (Figure

S2B-D). Of these, 783655 particles and 413330 particles were selected from Batch 1 and Batch

2, respectively and independently subjected to 3D classification (K=8) using a low pass filtered

(60 Å) map of our ABCA7$_{\text{BPL}}$ structure as a reference. A single highest resolution class from each

was selected and their particles combined and subjected to an additional round of 2D classification.

2024649 particles were subjected to another round of 3D classification (K=8). A single, highest

resolution class containing 50704 particles was refined to 5.4 Å. Particles were re-extracted using

refined coordinates and unbinned (pixel size 0.889 Å/pixel) and subjected to a round of 3D

refinement, Bayesian Polishing, and postprocessing/B-factor sharpening to yield a final map 4.0

Å resolution and its local resolution filtered variant calculated using Relion's own algorithm. Local

resolution maps are shown for ABCA7$_{\text{BPL}}$ (Map1 & Map2) and ABCA7$_{\text{PE}}$ in Figure S3.

For the ABCA7$_{\text{DIGITONIN}}$ dataset (Figure S4B-E), a total of 7437149 particles were picked

from 16213 ctf corrected and motion corrected micrographs and extracted at a 3-fold binned pixel

size of 2.667 Å/ pixel. After 2D classification, 1220497 particles were subjected to 3D

classification (K=8) using a low pass filtered (60 Å) map of our ABCA7$_{\text{BPL}}$ structure as a reference.

A single class comprising 324727 particles was refined to 5.4 Å. The refined coordinates were

then used to re-extract unbinned particles (0.889 Å/pixel), subjected to 3D refinement, and

Bayesian polishing. Further 3D classification (K=5) was performed and two similar classes

containing 149590 particles were picked for 3D refinement and postprocessing/B-factor

Sharpening to yield a 3.9 Å map.

For the ABCA7$_{EQ-ATP}$ dataset, an initial set of 2660267 particles were picked from 4914

ctf corrected micrographs (Figure S5B-E). After 2D classification, 469397 particles were subjected

to initial model building. This model was used as a 3D reference to perform 3D classification

(k=5). A single class comprising 174415 particles was refined to 5.4 Å, the corresponding particles

unbinned and re-refined. The nanodisc density was subtracted within Relion, followed by 3D

classification (k=3). The highest resolution class comprising 51780 particles was refined to 4.3

Angstroms and used for 3D reference-based particle picking for a larger data set comprising the

initial 4914 movies and a new set of 4474 ctf corrected micrographs. A total of 3773280 particles

were picked and subjected to multiple rounds of 2D classification in Relion 4.0-beta. 898916

particles were subjected to 3D classification (K=3) and a single highest resolution class consisting

of 407424 particles was refined to 5.4 Å. The refined coordinates were then used to re-extract the

respective particles without binning (0.889 Å/ pixel) and refined again before 3D classification

(K=5). A single class comprising 177230 particles was selected and subjected to 3D refinement,

Bayesian polishing, and postprocessing/B-factor sharpening to yield a final map at 3.7 Å.

Model building and refinement

Model building was done in coot 0.9.5[50] using a combination of Map 1 and its local

resolution filtered variant and Map 2. Both Map 1 and Map 2 displayed significant conformational

heterogeneity in the second half of ABCA7, with the quality of density in Map 1 allowing

placement of a TMD2 model guided in part by the homologous ABCA1 structure. Density

attributed to inter-TMD phospholipids was clearest in Map 1. Map 2 revealed very poor and

discontinuous density for TMD2-NBD2 but significantly better density for TMD1-NBD1 and the

majority of ECD1 and ECD2, allowing for de novo model building. The model for ECD was also

guided by the presence of nine glycosylation sites (N78, N98, N312, N340, N1335, N1381, N1386,

N1457, & N1518) as well as 4 disulfide bond pairs. Density for the lid region of the ECD was

missing in both maps. Model building for both NBDs was guided by structures of the homologous

transporters TM287[51], ABCG2, and ABCA1, where density features did not allow for de novo

model building. We observed extra density at the nucleotide binding sites for both NBDs despite

their open conformation. The structure of the RD was based on a homology model of the predicted

RD structure in ABCA4[25]. The quality of density for the ABCA7 RD allowed rigid body placement

for the entire domain. Restrained real space refinement of the model was carried out in Phenix

1.19.1[52] using automatically generated secondary structure restraints. Structural figures were

prepared in UCSF Chimera v. 1.13.1[53] , ChimeraX v. 1.2.5[54], and PyMOL 2.4.1 (The PyMOL

Molecular Graphics System, Version 1.8 Schrödinger, LLC).

The model for ABCA7$_{PE}$ was generated by rigid body placement of the ABCA7$_{BPL}$ model

followed by real space refinement in phenix as described above. The model for ABCA7$_{DIGITONIN}$

was generated by rigid body fitting each TMD, NBD, ECD, and RD into its postprocessed map

followed my manual adjustment of sidechains as allowed for the map. Non proteinaceous EM

density was modeled as a CHS and Digitonin (ligand ID Y01) molecule. The corrected model was

real space refined against the postprocessed map. The model for ABCA7$_{EQ\text{-}ATP}$ was built starting

with a homology model based on the ATP bound structure of ABCA4 (PDB 7LKZ), followed by

manual adjustment of the structure as required and permitted by the map. The ECD was replaced

by a rigid body fitted model of the ECD from the ABAC7$_{Digitonin}$ structure. This model was then

refined against both the postprocessed map, and its local resolution filtered counterpart.

MD simulations

We employed MD simulations to capture the arrangement and dynamics of the lipid bilayer

induced by the experimentally derived open conformation of ABCA7$_{PE}$, which was used as the

starting model in all the simulations. For system setup, a C-terminal carboxylate capping group,

an N-terminal ammonium capping group and all the hydrogen atoms were added using the PSFGEN

plugin of VMD (Visual Molecular Dynamics)[56]. The resulting all-atom (AA) model was then

converted to a coarse-grained (CG) Martini model using the MARTINIZE protocol

(http://www.cgmartini.nl/), using an elastic network on atom pairs within a 10-Å cutoff. The

Orientations of Proteins in Membranes (OPM) database[57] was used to identify and align the

transmembrane region of the protein with the membrane normal. The protein was embedded in

two distinct lipid bilayers (palmitoyl-oleoyl-phosphatidyl-ethanolamine (POPE) and cholesterol

with a molar ratio of 4:1 (POPE/Chol), and palmitoyl-oleoyl-phosphatidyl-choline (POPC) and

cholesterol with a molar ratio of 4:1 (POPC/Chol), respectively). The protein secondary structure

was defined from the AA model and was maintained throughout the CG simulations by the applied

elastic network. To increase the sampling of lipid-protein interactions and improve statistics, four

independent copies of the CG protein were placed at a distance of 200 Å in a large lipid bilayer

(400×400 Å^2). The system was then solvated and ionized with 150 mM salt using INSANE[58].

The systems were simulated employing GROMACS 2021.3[59,60]. A 20-fs timestep was

employed in all the simulations. The temperature was maintained at 310 K with a velocity-

rescaling thermostat[61] employing a coupling time constant of 1 ps. A semi-isotropic 1 bar pressure

was maintained using the Berendsen barostat [62] with a compressibility and relaxation time constant

of 3×10^{-4} bar and 5 ps, respectively. The systems were energy minimized for 1,000 steps, followed

by short equilibration runs of 18 ns, while restraints were applied to lipid bilayer headgroups and

protein backbones. During this time the restraints on bilayer headgroups were reduced gradually

from k = 200 kJ.mol^{-1}.nm^{-2} to zero, whereas the protein backbones' restraints (k = 1,000 kJ.mol^{-1}

^{1}.nm^{-2}) were kept constant. Each system was then simulated for 2 μs, with restraints only applied

to the protein backbones, resulting in an aggregate sampling of 8 μs (4 copies × 2 μs). All the

systems were simulated following the same MD protocol.

All the molecular images were generated using VMD[56]. The membrane deformation

induced by ABCA7 was quantified by calculating the z distance of the lipid phosphate moieties

(PO_4 bead type in MARTINI) with respect to the bilayer midplane, over the last 1 μs of each

trajectory. The generated histogram (binned in $2 \times 2 \ \text{Å}^2$ bins) in each leaflet illustrates the spatial

distribution of the height of the lipid head groups within each leaflet. We quantified the differential

movement of POPE and POPC within the protein lumen by calculating the number of

phospholipids located within the TMDs. If the PO_4 bead of a phospholipid was within 22.5 Å and

12.5 Å in x and y, respectively, with respect to a protein's center in the x-y plane (membrane plane),

then the phospholipid was considered to be within the TMD lumen (Figure S9C).

Acknowledgments

We would like to thank Dr. Kaspar Locher at ETH, Zurich, Switzerland, for providing the

synthetic gene construct of ABCA7. We would also like to thank the cryo-EM and shared

instruments core facilities at the Hormel Institute for help with experimental setup, and Dr.

Rhoderick Brown, Dr. Jarrod French and Dr. Jeppe Olsen for critical reading and discussion during

manuscript preparation. This work was supported in part by the Hormel Foundation (Institutional

research funds to AA), the National Institutes of Health (NIH) award 1R21-AG069180-01A1 (to

AA), and the Cure Alzheimer's fund (to TK). The computational component of the project was

supported by the NIH awards P41-GM104601 (to ET) and R01-GM123455 (to ET). We also

acknowledge computing resources provided by Blue Waters at National Center for

Supercomputing Applications (NCSA), and by eXtreme Science and Engineering Discovery

Environment (XSEDE) (grant MCA06N060 to ET), and by Microsoft Azure.

Author Contributions

AA conceived the research. LTML, JRT, and AA performed all experiments. MD simulations

were conducted and analyzed by SD, SP, and ET. TK participated in assay design. LTML, JRT,

and AA wrote the manuscript with input from all other authors.

Declaration of Interests

SP is currently an employee of Loxo Oncology @ Lilly and is a shareholder of stock in Eli Lilly

and Co. The rest of the authors declare no competing interests.

Data and materials availability:

The cryo-EM Maps have been deposited at the Electron Microscopy Databank (EMDB) under

accession codes EMD-AAAAA (ABCA7$_{BPL}$ Map 1), EMD-BBBBB (ABCA7BPL Map 2), EMD-

CCCCC (ABCA7$_{PE}$), EMD-DDDDD (ABCA7$_{EQ-ATP}$), and EMD-EEEEE (ABCA7DI$_{GITONIN}$). The

associated atomic coordinates have been deposited at the Protein Data bank (PDB) under accession

codes   1WWW   (ABCA7$_{BPL}$),   2XXX(ABCA7$_{PE}$),   3YYY,   (ABCA7$_{EQ-ATP}$),   and   4ZZZ

(ABCA7DI$_{GITONIN}$).

References

1   Albrecht, C. & Viturro, E. The ABCA subfamily--gene and protein structures, functions
and associated hereditary diseases. *Pflugers Arch* **453**, 581-589, doi:10.1007/s00424-006-
0047-8 (2007).
2   Brooks-Wilson, A. *et al.* Mutations in ABC1 in Tangier disease and familial high-density
lipoprotein deficiency. *Nat Genet* **22**, 336-345, doi:10.1038/11905 (1999).
3   Allikmets, R. *et al.* Mutation of the Stargardt disease gene (ABCR) in age-related
macular degeneration. *Science* **277**, 1805-1807, doi:10.1126/science.277.5333.1805
(1997).
4   Akiyama, M. *et al.* Mutations in lipid transporter ABCA12 in harlequin ichthyosis and
functional recovery by corrective gene transfer. *J Clin Invest* **115**, 1777-1784,
doi:10.1172/JCI24834 (2005).
5   Hollingworth, P. *et al.* Common variants at ABCA7, MS4A6A/MS4A4E, EPHA1, CD33
and CD2AP are associated with Alzheimer's disease. *Nat Genet* **43**, 429-435,
doi:10.1038/ng.803 (2011).

6   Almeida, J. F. F., Dos Santos, L. R., Trancozo, M. & de Paula, F. Updated Meta-Analysis
550   of BIN1, CR1, MS4A6A, CLU, and ABCA7 Variants in Alzheimer's Disease. *J Mol*
551   *Neurosci* **64**, 471-477, doi:10.1007/s12031-018-1045-y (2018).
552   7   Bellenguez, C. *et al.* Contribution to Alzheimer's disease risk of rare variants in TREM2,
553   SORL1, and ABCA7 in 1779 cases and 1273 controls. *Neurobiol Aging* **59**, 220 e221-
554   220 e229, doi:10.1016/j.neurobiolaging.2017.07.001 (2017).
555   8   Le Guennec, K. *et al.* ABCA7 rare variants and Alzheimer disease risk. *Neurology* **86**,
556   2134-2137, doi:10.1212/WNL.0000000000002627 (2016).
557   9   Swerdlow, R. H. Rare ABCA7 variants in Alzheimer disease: Guilt by association.
558   *Neurology* **86**, 2118-2119, doi:10.1212/WNL.0000000000002630 (2016).
559   10   Lambert, J. C. *et al.* Meta-analysis of 74,046 individuals identifies 11 new susceptibility
560   loci for Alzheimer's disease. *Nat Genet* **45**, 1452-1458, doi:10.1038/ng.2802 (2013).
561   11   Ma, F. C. *et al.* Meta-Analysis of the Association between Variants in ABCA7 and
562   Alzheimer's Disease. *J Alzheimers Dis* **63**, 1261-1267, doi:10.3233/JAD-180107 (2018).
563   12   De Roeck, A., Van Broeckhoven, C. & Sleegers, K. The role of ABCA7 in Alzheimer's
564   disease: evidence from genomics, transcriptomics and methylomics. *Acta Neuropathol*
565   **138**, 201-220, doi:10.1007/s00401-019-01994-1 (2019).
566   13   Aikawa, T. *et al.* ABCA7 Regulates Brain Fatty Acid Metabolism During LPS-Induced
567   Acute Inflammation. *Front Neurosci* **15**, 647974, doi:10.3389/fnins.2021.647974 (2021).
568   14   Serhan, C. N. Pro-resolving lipid mediators are leads for resolution physiology. *Nature*
569   **510**, 92-101, doi:10.1038/nature13479 (2014).
570   15   Astudillo, A. M., Balboa, M. A. & Balsinde, J. Selectivity of phospholipid hydrolysis by
571   phospholipase A2 enzymes in activated cells leading to polyunsaturated fatty acid
572   mobilization. *Biochim Biophys Acta Mol Cell Biol Lipids* **1864**, 772-783,
573   doi:10.1016/j.bbalip.2018.07.002 (2019).
574   16   Quazi, F. & Molday, R. S. Differential phospholipid substrates and directional transport
575   by ATP-binding cassette proteins ABCA1, ABCA7, and ABCA4 and disease-causing
576   mutants. *J Biol Chem* **288**, 34414-34426, doi:10.1074/jbc.M113.508812 (2013).
577   17   Tomioka, M. *et al.* Lysophosphatidylcholine export by human ABCA7. *Biochim Biophys*
578   *Acta Mol Cell Biol Lipids* **1862**, 658-665, doi:10.1016/j.bbalip.2017.03.012 (2017).
579   18   Tonnies, E. & Trushina, E. Oxidative Stress, Synaptic Dysfunction, and Alzheimer's
580   Disease. *J Alzheimers Dis* **57**, 1105-1121, doi:10.3233/JAD-161088 (2017).
581   19   Newcombe, E. A. *et al.* Inflammation: the link between comorbidities, genetics, and
582   Alzheimer's disease. *J Neuroinflammation* **15**, 276, doi:10.1186/s12974-018-1313-3
583   (2018).
584   20   Czubowicz, K., Jesko, H., Wencel, P., Lukiw, W. J. & Strosznajder, R. P. The Role of
585   Ceramide and Sphingosine-1-Phosphate in Alzheimer's Disease and Other
586   Neurodegenerative Disorders. *Mol Neurobiol* **56**, 5436-5455, doi:10.1007/s12035-018-
587   1448-3 (2019).
588   21   Testa, G. *et al.* Changes in brain oxysterols at different stages of Alzheimer's disease:
589   Their involvement in neuroinflammation. *Redox Biol* **10**, 24-33,
590   doi:10.1016/j.redox.2016.09.001 (2016).
591   22   Popp, J. *et al.* Cholesterol metabolism is associated with soluble amyloid precursor
592   protein production in Alzheimer's disease. *J Neurochem* **123**, 310-316,
593   doi:10.1111/j.1471-4159.2012.07893.x (2012).

23 Sleegers, K. Expression of ABCA7 in Alzheimer's disease. *Acta Neuropathol* **139**, 941-
942, doi:10.1007/s00401-020-02136-8 (2020).
24 Iwamoto, N., Abe-Dohmae, S., Sato, R. & Yokoyama, S. ABCA7 expression is regulated
by cellular cholesterol through the SREBP2 pathway and associated with phagocytosis. *J
Lipid Res* **47**, 1915-1927, doi:10.1194/jlr.M600127-JLR200 (2006).
25 Liu, F., Lee, J. & Chen, J. Molecular structures of the eukaryotic retinal importer
ABCA4. *Elife* **10**, doi:10.7554/eLife.63524 (2021).
26 Qian, H. *et al.* Structure of the Human Lipid Exporter ABCA1. *Cell* **169**, 1228-1239
e1210, doi:10.1016/j.cell.2017.05.020 (2017).
27 Scortecci, J. F. *et al.* Cryo-EM structures of the ABCA4 importer reveal mechanisms
underlying substrate binding and Stargardt disease. *Nat Commun* **12**, 5902,
doi:10.1038/s41467-021-26161-7 (2021).
28 Xie, T., Zhang, Z., Fang, Q., Du, B. & Gong, X. Structural basis of substrate recognition
and translocation by human ABCA4. *Nat Commun* **12**, 3853, doi:10.1038/s41467-021-
24194-6 (2021).
29 Harris, A. *et al.* Structure and efflux mechanism of the yeast pleiotropic drug resistance
transporter Pdr5. *Nat Commun* **12**, 5254, doi:10.1038/s41467-021-25574-8 (2021).
30 Thomas, C. *et al.* Structural and functional diversity calls for a new classification of ABC
transporters. *FEBS Lett* **594**, 3767-3775, doi:10.1002/1873-3468.13935 (2020).
31 Orlando, B. J. & Liao, M. ABCG2 transports anticancer drugs via a closed-to-open
switch. *Nat Commun* **11**, 2264, doi:10.1038/s41467-020-16155-2 (2020).
32 Manolaridis, I. *et al.* Cryo-EM structures of a human ABCG2 mutant trapped in ATP-
bound and substrate-bound states. *Nature* **563**, 426-430, doi:10.1038/s41586-018-0680-3
(2018).
33 Qiu, W. *et al.* Structure and activity of lipid bilayer within a membrane-protein
transporter. *Proc Natl Acad Sci U S A* **115**, 12985-12990, doi:10.1073/pnas.1812526115
(2018).
34 Cuyvers, E. *et al.* Mutations in ABCA7 in a Belgian cohort of Alzheimer's disease
patients: a targeted resequencing study. *Lancet Neurol* **14**, 814-822, doi:10.1016/S1474-
4422(15)00133-7 (2015).
35 Cohen, J. C. *et al.* Multiple rare alleles contribute to low plasma levels of HDL
cholesterol. *Science* **305**, 869-872, doi:10.1126/science.1099870 (2004).
36 Liaci, A. M. *et al.* Structure of the human signal peptidase complex reveals the
determinants for signal peptide cleavage. *Mol Cell* **81**, 3934-3948 e3911,
doi:10.1016/j.molcel.2021.07.031 (2021).
37 Tanaka, N., Abe-Dohmae, S., Iwamoto, N., Fitzgerald, M. L. & Yokoyama, S. Helical
apolipoproteins of high-density lipoprotein enhance phagocytosis by stabilizing ATP-
binding cassette transporter A7. *J Lipid Res* **51**, 2591-2599, doi:10.1194/jlr.M006049
(2010).
38 Abe-Dohmae, S. Y., S. . ABCA7: a potential mediator between cholesterol homeostasis
and the host defense system. *Clinical Lipidology* **7**, 677-687 (2012).
39 Zachowski, A. Phospholipids in animal eukaryotic membranes: transverse asymmetry
and movement. *Biochem J* **294 (Pt 1)**, 1-14, doi:10.1042/bj2940001 (1993).
40 Gabande-Rodriguez, E., Keane, L. & Capasso, M. Microglial phagocytosis in aging and
Alzheimer's disease. *J Neurosci Res* **98**, 284-298, doi:10.1002/jnr.24419 (2020).

41   Hirt, U. A. & Leist, M. Rapid, noninflammatory and PS-dependent phagocytic clearance
of necrotic cells. *Cell Death Differ* **10**, 1156-1164, doi:10.1038/sj.cdd.4401286 (2003).

42   Fadok, V. A., de Cathelineau, A., Daleke, D. L., Henson, P. M. & Bratton, D. L. Loss of
phospholipid asymmetry and surface exposure of phosphatidylserine is required for
phagocytosis of apoptotic cells by macrophages and fibroblasts. *J Biol Chem* **276**, 1071-
1077, doi:10.1074/jbc.M003649200 (2001).

43   Raja, M. Do small headgroups of phosphatidylethanolamine and phosphatidic acid lead
to a similar folding pattern of the K(+) channel? *J Membr Biol* **242**, 137-143,
doi:10.1007/s00232-011-9384-4 (2011).

44   Molday, L. L. & Molday, R. S. 1D4: a versatile epitope tag for the purification and
characterization of expressed membrane and soluble proteins. *Methods Mol Biol* **1177**, 1-
15, doi:10.1007/978-1-4939-1034-2_1 (2014).

45   Ritchie, T. K. *et al.* Chapter 11 - Reconstitution of membrane proteins in phospholipid
bilayer nanodiscs. *Methods Enzymol* **464**, 211-231, doi:10.1016/S0076-6879(09)64011-8
(2009).

46   Geertsma, E. R., Nik Mahmood, N. A., Schuurman-Wolters, G. K. & Poolman, B.
Membrane reconstitution of ABC transporters and assays of translocator function. *Nat*
Protoc **3**, 256-266, doi:10.1038/nprot.2007.519 (2008).

47   Chifflet, S., Torriglia, A., Chiesa, R. & Tolosa, S. A method for the determination of
inorganic phosphate in the presence of labile organic phosphate and high concentrations
of protein: application to lens ATPases. *Anal Biochem* **168**, 1-4 (1988).

48   Zheng, S. Q. *et al.* MotionCor2: anisotropic correction of beam-induced motion for
improved cryo-electron microscopy. *Nat Methods* **14**, 331-332, doi:10.1038/nmeth.4193
(2017).

49   Zhang, K. Gctf: Real-time CTF determination and correction. *J Struct Biol* **193**, 1-12,
doi:10.1016/j.jsb.2015.11.003 (2016).

50   Scheres, S. H. & Chen, S. Prevention of overfitting in cryo-EM structure determination.
Nat Methods **9**, 853-854, doi:10.1038/nmeth.2115 (2012).

51   Emsley, P., Lohkamp, B., Scott, W. G. & Cowtan, K. Features and development of Coot.
Acta Crystallogr D Biol Crystallogr **66**, 486-501, doi:10.1107/S0907444910007493
(2010).

52   Hohl, M., Briand, C., Grutter, M. G. & Seeger, M. A. Crystal structure of a heterodimeric
ABC transporter in its inward-facing conformation. *Nat Struct Mol Biol* **19**, 395-402,
doi:10.1038/nsmb.2267 (2012).

53   Adams, P. D. *et al.* PHENIX: a comprehensive Python-based system for macromolecular
structure solution. *Acta Crystallogr D Biol Crystallogr* **66**, 213-221,
doi:10.1107/S0907444909052925 (2010).

54   Pettersen, E. F. *et al.* UCSF Chimera--a visualization system for exploratory research and
analysis. *J Comput Chem* **25**, 1605-1612, doi:10.1002/jcc.20084 (2004).

55   Pettersen, E. F. *et al.* UCSF ChimeraX: Structure visualization for researchers, educators,
and developers. *Protein Sci* **30**, 70-82, doi:10.1002/pro.3943 (2021).

56   Humphrey, W., Dalke, A. & Schulten, K. VMD: visual molecular dynamics. *J Mol*
Graph **14**, 33-38, 27-38, doi:10.1016/0263-7855(96)00018-5 (1996).

57   Lomize, M. A., Lomize, A. L., Pogozheva, I. D. & Mosberg, H. I. OPM: orientations of
proteins in membranes database. *Bioinformatics* **22**, 623-625,
doi:10.1093/bioinformatics/btk023 (2006).

58 Wassenaar, T. A., Ingolfsson, H. I., Bockmann, R. A., Tieleman, D. P. & Marrink, S. J.
Computational Lipidomics with insane: A Versatile Tool for Generating Custom
Membranes for Molecular Simulations. *J Chem Theory Comput* **11**, 2144-2155,
doi:10.1021/acs.jctc.5b00209 (2015).
59 Hess, B., Kutzner, C., van der Spoel, D. & Lindahl, E. GROMACS 4: Algorithms for
Highly Efficient, Load-Balanced, and Scalable Molecular Simulation. *J Chem Theory*
Comput **4**, 435-447, doi:10.1021/ct700301q (2008).
60 de Jong, D. H. *et al.* Improved Parameters for the Martini Coarse-Grained Protein Force
Field. *J Chem Theory Comput* **9**, 687-697, doi:10.1021/ct300646g (2013).
61 Bussi, G., Donadio, D. & Parrinello, M. Canonical sampling through velocity rescaling. *J*
Chem Phys **126**, 014101, doi:10.1063/1.2408420 (2007).
62 H. J. C. Berendsen, J. P. M. P., W. F. van Gunsteren, A. DiNola, and J. R. Haak.
Molecular dynamics with coupling to an external bath. *J. Chem. Phys.* **91**,
doi:10.1063/1.448118 (1984).
699

Figures and Legends

Figure 1. Functional characterization and structure of human ABCA7 incorporated in
nanodiscs (A) ATPase activity of ABCA7 at different ATP concentrations and **(B)** different
nanodisc phospholipid/cholesterol compositions. Cholesterol (20%) presence is indicated by + or
-. Experimental replicates (n)=3 and error bars represent standard deviation (s.d.). **(C)** Cryo-EM
map of ABCA7$_{BPL}$ (yellow) at 3.6 Å resolution and **(D)** ABCA7$_{PE}$ at 4.0 Å resolution(purple).
Density for protein is shown in yellow (0.025 contour) and that for modeled lipid acyl chains
(0.035 contour) shown in pink (TMD luminal lipids) and white (peripherally associated lipids).
(E) ABCA7$_{PE}$ shown in ribbon format with Half 1 colored yellow and Half 2 purple, along with
density for TMD lipids (pink 0.025 contour) and bound nucleotide (yellow 0.035 contour). Acyl
chains and glycans are shown as pink and grey sticks, respectively. **(F)** TMD lumen with density
for TMD lipids (pink 0.025 contour) viewed from the extracellular side (top), membrane plane
(middle) and cytoplasmic side (bottom). Cα atoms for the cytoplasmic gate are shown as black
spheres. **(G)** View of the TMD-ECD interface with select residues oriented towards lipids
(transparent red spheres) shown.

Figure 2. Cryo-EM structure of ABCA7DIGITONIN. **(A)** Cryo-EM map of human ABCA7$_{Digitonin}$

at 3.9 Å with density for protein shown in green (0.013 contour) and extraneous density likely

belonging to detergent shown in pink (0.013 contour). **(B)** Cryo-EM structure of human

ABCA7$_{DIGITONIN}$ shown in ribbon format with each half colored differently (blue and green). **(C)**

TMD lumen of ABCA7$_{DIGITONIN}$ with density for bulk micelle shown in white at 0.015 contour

where detergent density inside TMDs disappears. Sticks for unmodeled detergent molecules are

shown for which density is visible at the lower contour of 0.013. **(D)** TMD lumen of ABCA7$_{PE}$

with density (0.035 contour) for TMD lipids (pink) and peripherally associated ordered lipids

(white) shown. **(E)** Same as D with higher density contour of 0.046 where density of peripherally

associated lipids is absent but that of luminal lipids remains.

Figure 3. Cryo-EM structure of ABCA7$_{EQ\text{-}ATP}$. (A) Cryo-EM map of human ABCA7$_{EQ\text{-}ATP}$ in

BPL/Chol nanodiscs at 3.7 Å resolution (orange density, 0.02 contour). **(B)** Cryo-EM structure of

ABCA7$_{EQ\text{-}ATP}$ shown in ribbon format with half 1 and half 2 colored yellow and orange,

respectively. The transparent orange oval demarcates the observed exit pocket. **(C)** Central slice

of electrostatic surface map showing surface details of exit pocket in two different orientations **(D)**

4-TM bundle forming the exit pocket. **(E)** NBDs viewed from the extracellular side with density

for bound nucleotide (blue sticks) shown in blue (0.02 contour).

Figure 4. Comparison of open and closed conformations of ABCA7 (A) Overall structural

alignment of the three ABCA7 conformations (center) along with individual alignments of rigid

body pairs TMD1-NBD1, TMD2-NBD2, ECD, and RD (**B**) Overall alignment of the three ABCA7

conformations showing only TMD1 and TMD2 viewed from the extracellular side using the

TMD1-NBD1 pair as an alignment reference. TMs lining the TMD pathway are numbered (**C**)

Same as panel B, viewed from the cytoplasmic side with Cα atoms of gate forming residues shown

as spheres (**D**) Same as panels B/C showing only RDs viewed from the cytosolic side. (**E**) Same

as panel D showing just NBDs and bound nucleotides viewed from the extracellular side. ATPγS

and ATP-Mg^{2+} are shown as purple and orange spheres, respectively. (**F**) Root mean square

deviations (RMSD) of aligned atoms of ABCA7$_{BPE}$ vs ABCA7$_{EQ-ATP}$ and ABCA7$_{DIGITONIN}$.

Figure 5. MD Simulations of ABCA7 in Lipid Bilayers. (A) Left: A representative simulation system with four copies of ABCA7 (silver), taken from the POPE/cholesterol (yellow/cyan) lipid patch. Right: the phospholipid belt (blue lines) formed in (top) the POPE/cholesterol (yellow/cyan) and (bottom) POPC/cholesterol (red/cyan) membranes (t = 2 μs). (B) The total count of phospholipids (sum of all four simulated proteins in a patch) partitioned in the TMD lumen for POPE/cholesterol and POPC/cholesterol bilayers (averaged over time ± standard deviation). (C/D) Heatmaps representing the average height (z values) of POPE headgroup with respect to the membrane midplane in the extracellular (C) and cytoplasmic (D) leaflets. Phospholipids are observed to climb the protein and form a dome-like configuration in the TMD lumen. (E) Snapshot of lipids partitioned in the TMD lumen, taken from the POPE trajectory at t = 2 μs. Polar, nonpolar, basic, and acidic residues are colored green, gray, blue, and red, respectively. TMD2 is hidden for a clearer view of the luminal dome-like lipid configuration (outlined by blue lines).

Figure 6. Proposed model of ABCA7 conformational transitions and lipid interactions. The

membrane bilayer leaflets are shown in grey. Red arrows indicate movement of lipids (black)

under the influence of TMD conformation. The ECD and RD have been omitted for simplicity as

has any reference to potential lipid flipping from the cytoplasmic to extracellular leaflets.

Figure S1. ABCA7_BPL characterization and Cryo-EM data processing. **A** ATPase data for nanodisc reconstituted ABCA7 with and without sodium orthovanadate (VO4) or ATP𝛾S. N=3 and error bars represent s.d **B** SEC peak of ABCA7_BPL sample for cryo-EM. **C** Representative cryo-EM micrograph at -2.5 μm defocus. Scale bar = 20 nm. **D** cryo-EM processing workflow. Boxes indicate 3D classes used for further refinement for both Map 1 and Map 2 (red). **E** Fourier shell correlation (FSC) curves for Map 1 (top) and Map 2 (bottom) Dotted lines indicate position 0. 143 and 0.5 cutoff criteria for resolution estimates.

Figure S2. ABCA7PE Cryo-EM data processing. A Size exclusion chromatography micrograph
of cryo-EM sample showing monodisperse ABCA7PE nanodisc (main peak). **B** Representative
micrograph at -2.5 μm defocus and 2D classes. Scale bar = 20 nm. **C** cryo-EM processing
workflow. Dashed boxes demarcate Subsets 1 and 2. Solid boxes indicate 3D classes used for
further refinement. C2D = 2D Classification, C3D=3D classification, R3D = 3D refinement **D**
Fourier shell correlation (FSC) curves for ABCA7PE Dotted lines indicate position 0.143 and 0.5
cutoff criteria for resolution estimates.

791 **Figure S3. ABCA7$_{BPL}$ and ABCA7$_{PE}$ local resolution filtered maps.** Color keys are indicated

792 on the left of each set of maps with numbers representing resolution (Å).

794
Figure S4 ABCA7$_{DIGITON}$ cryo-EM processing A SEC profile of ABCA7$_{DIGITONIN}$ and its ATPase

activity with and without ATPγS. **B** Representative micrograph at -2.5 μm defocus and 2D classes.

Scale bar = 20 nm. **C** cryo-EM processing workflow. C2D = 2D Classification, C3D=3D

classification, R3D = 3D refinement. **D** Local resolution colored EM map of ABCA7$_{DIGITONIN}$. **E**

Fourier shell correlation (FSC) curves for ABCA7$_{DIGITONIN}$. Dotted lines indicate position 0.143

and 0.5 cutoff criteria for resolution estimates. **F** Superposition of ABCA7 (green, this manuscript),

ABCA1 (grey, PDB 5XJY), and ABCA4 (red, PDB 7LKP) structures in digitonin.

Figure S5. ABCA7$_{EQ-ATP}$ cryo-EM processing. (A) SEC peak for ABCA7$_{EQ-ATP}$ in nanodiscs. **B** Representative EM micrograph (-2.5 defocus) and rep 2D classes for ABCA7$_{EQ-ATP}$ sample. **C** Cryo-EM data processing pipeline. C2D = 2D Classification, C3D=3D classification, R3D = 3D refinement. **D** FSC curves. **E** Local resolution colored EM map.

Figure S6. TMD-ECD interfaces of open and closed form ABCA7. (A) The TMD-ECD binding

interfaces of ABCA7$_{PE}$ with TMD1 and TMD2 and with Cα for residues in TMD1 and TMD2

within 5Å of either ECD or vice versa shown as spheres. **(B)** The same analysis for the TMD-ECD

binding interfaces of ABCA7$_{EQ\text{-}ATP}$ with closed cavity.

Figure S7. Conservation of ABCA and ABCG family structural elements (A) Ribbon representation of select ABCG family transporter structures and their respective PDB IDs including apo open ABCG2 (red), closed ATP bound structure of ABCG2$_{EQ}$ (black), and ABCG5/G8 (cyan and green, respectively) (**B**) Alignment of NBD-TMD pairs from open and closed conformations of ABCG2 (left), ABCG2 and ABCG5/G8 (center), and closed ABCG2 with TMD-NBD pairs from ABCA7 half 1 (gold) and half 2 (orange).

Figure S8. Lipid configuration induced by protein copies in a POPC bilayer and criteria used for identifying luminal lipids in MD simulations. (A/B) POPC headgroup height calculated for extracellular (A) and cytoplasmic (B) leaflets. **(C)** Dimension and position of the box used to calculate the number of phospholipids (yellow) partitioned in the TMD leaflets. The box is centered at the protein center and has dimensions of 50 Å and 25 Å in x and y directions, respectively. The ABCA7 transporter is hidden for clarity. **(D)** Contact map, based on the number of POPE headgroups in proximity of each residue throughout the simulation. **(E)** Accumulation of POPE headgroups (yellow spheres) in close proximity of most frequently contacted residues throughout the simulation.

Figure S9. Amino acid positions of ABCA7 with variants associated with risk of AD. Structure of ABCA7 with Cα atoms shown for amino acid with known pathogenic variants (green), protective variants (red), and ABCA7 residues conserved in ABCA1 with mutations known to disrupt the latter's binding to apoA1 and/or phospholipid translocation (blue). Cyan spheres highlight residue positions of ABCA1 mutations known to disrupt phospholipid efflux including those equivalent to the VFVNFA motif within the ABCA1 RD. Grey bars indicate the upper (extracellular) and lower (cytoplasmic) membrane bilayer leaflets (UL and LL, respectively).

```
hsABCA7  2141  TAETVL
hsABCA1  2256  VKESYV
hsABCA4        ......
```

849   **Figure S10 Sequence alignment of human (hs) ABCA7, ABCA1, and ABCA4.**

Dataset	ABCA7$_{BPL}$		ABCA7$_{PE}$	ABCA7$_{DIGITONIN}$	ABCA7$_{EQ-ATP}$
Magnification	96k		96k	96k	96k
Pixel Size (Å)	0.895		0.889	0.889	0/889
Total Dose (e/Å²)	60		40	40	40
Defocus Range (um)	-0.8 to 2.6		-0.8 to 2.6	-0.8 to 2.6	-0.8 to 2.6
Maps	Map 1	Map 2	Map3	Map4	Map5
EMDB ID	EMD-22996	EMD-22998	EMD-A	EMD-B	EMD-C
# Particles in final Class	91381	124114	50704	149590	177230
Resolution (Å) (0.143 threshold)	3.6	3.2	4.0	3.9	3.7
Sharpening B factor					
Refined Coordinates	ABCA7$_{BPL}$		ABCA7$_{PE}$	ABCA7$_{DIGITONIN}$	ABCA7$_{EQ-ATP}$
PDB ID	AAAA		BBBB	CCCC	DDDD
# Residues/Non-hydrogen Atoms	1801/14761		1846/14749	1856/4662	1871/14748
Glycans	16		16	26	16
Ligands	21		20		2
R.M.S deviations					
Bond Length (Å)	0.003		0.003	0.003	0.003
Bond Angles (°)	0.616		0.600	0.678	0.645
MolProbity Statistics					
MolProbity Score	1.68		1.72	1.83	1.77
Clashscore	8.05		8.16	11.40	9.83
Poor rotamers (%)	0.00		0.00	0.00	0.00
Ramachandran statistics					
Favored (%)	96.38		95.84	96.24	96.22
Allowed (%)	3.62		4.16	3.76	3.78
Outliers (%)	0.00		0.00	0.00	0.00

851 **Table S1 Data collection and refinement statistics.**