The Disease-Associated Proteins *Drosophila* Nab2 and Ataxin-2 Interact with Shared RNAs and Coregulate Neuronal Morphology

J. Christopher Rounds*, Edwin B. Corgiat*, Changtian Ye*, Joseph A. Behnke*, Seth M. Kelly[‡], Anita H. Corbett[†1], Kenneth H. Moberg*[1,2]

*Department of Cell Biology, Emory University School of Medicine, Atlanta, GA 30322;
[†]Department of Biology, Emory University, Atlanta, GA 30322;
[‡]Department of Biology, The College of Wooster, Wooster, OH 44691

[1]Co-corresponding authors

For correspondence:
[2] Kenneth H. Moberg
   Department of Cell Biology, Emory University School of Medicine
   615 Michael St.
   Whitehead Biomedical Research Building, Room 442
   Atlanta, GA 30322
   Phone: 404-727-3733
   Email: kmoberg@emory.edu

Running head: Nab2 and Atx2 share RNAs and functions

ABSTRACT

Nab2 encodes a conserved polyadenosine RNA-binding protein (RBP) with broad roles in post-

transcriptional regulation, including in poly(A) RNA export, poly(A) tail length control, transcription

termination, and mRNA splicing. Mutation of the *Nab2* human ortholog *ZC3H14* gives rise to an

autosomal recessive intellectual disability, but understanding of Nab2/ZC3H14 function in metazoan

nervous systems is limited, in part because no comprehensive identification of metazoan Nab2/ZC3H14-

associated RNA transcripts has yet been conducted. Moreover, many Nab2/ZC3H14 functional protein

partnerships likely remain unidentified. Here we present evidence that *Drosophila melanogaster* Nab2

interacts with the RBP Ataxin-2 (Atx2), a neuronal translational regulator, and implicate these proteins

in coordinate regulation of neuronal morphology and adult viability. We then present the first high-

throughput identifications of Nab2- and Atx2-associated RNAs in *Drosophila* brain neurons using an

RNA immunoprecipitation-sequencing (RIP-Seq) approach. Critically, the RNA interactomes of each

RBP overlap, and Nab2 exhibits high specificity in its RNA associations in neurons *in vivo*, associating

with a small fraction of all polyadenylated RNAs. The identities of shared associated transcripts (e.g.

drk, *me31B*, *stai*) and of transcripts specific to Nab2 or Atx2 (e.g. *Arpc2*, *tea*, respectively) promise

insight into neuronal functions of and interactions between each RBP. Significantly, Nab2-associated

RNAs are overrepresented for internal A-rich motifs, suggesting these sequences may partially mediate

Nab2 target selection. Taken together, these data demonstrate that Nab2 opposingly regulates neuronal

morphology and shares associated neuronal RNAs with Atx2, and that *Drosophila* Nab2 associates with

a more specific subset of polyadenylated mRNAs than its polyadenosine affinity alone may suggest.

Introduction

Intellectual disability refers to a broad group of neurodevelopmental disorders affecting approximately

1% of the world population (Maulik *et al.* 2011) and defined by significant limitations in intellectual

functioning and adaptive behavior (Tassé *et al.* 2016; Vissers *et al.* 2016). Intellectual disabilities are

etiologically diverse and in some cases genetically complex, yet many exhibit overlapping molecular

dysfunctions in a comparatively limited set of fundamental neurodevelopmental pathways (reviewed in

Chelly *et al.* 2006; van Bokhoven 2011; and Verma *et al.* 2019). Thus, monogenic intellectual

disabilities represent experimentally tractable avenues for understanding both these disorders more

broadly and neurodevelopment in general (Najmabadi *et al.* 2011; Agha *et al.* 2014). One set of such

informative monogenic intellectual disabilities is caused by mutations affecting genes encoding RNA-

binding proteins (RBPs) (reviewed in Bardoni *et al.* 2012) such as *ZC3H14* (*zinc finger CCCH-type*

containing 14). Specifically, loss-of-function mutations in *ZC3H14*, which encodes a ubiquitously

expressed polyadenosine RBP, cause a non-syndromic form of autosomal recessive intellectual

disability (Pak *et al.* 2011; Al-Nabhani *et al.* 2018). However, the molecular functions and

developmental roles of human ZC3H14 are largely unknown; defining these functions and roles

provides an opportunity to better understand intellectual disability and human neurodevelopment.

Drosophila melanogaster has proven a powerful model system to understand the molecular

functions of proteins encoded by many intellectual disability genes (Inlow and Restifo 2004; Oortveld *et*

al. 2013), and ZC3H14 is no exception—its functions have begun to be dissected in part through study

of its *Drosophila* ortholog Nab2 (Pak *et al.* 2011; Kelly *et al.* 2014). *Drosophila* Nab2, like ZC3H14, is

a polyadenosine RNA-binding protein that induces neurological defects when its expression is altered;

deletion or overexpression of *Nab2* causes neuronal morphological defects in the eye, axon projection

defects in the developing brain, and memory impairments (Pak *et al.* 2011; Kelly *et al.* 2016;

Bienkowski *et al.* 2017; Corgiat *et al.* 2020). The function of Nab2 is particularly important in

Drosophila neurons, as pan-neuronal expression of Nab2 or an isoform of human ZC3H14 is sufficient

to rescue the severe limitation in adult viability and locomotor defects caused by zygotic Nab2

deficiency (Pak *et al.* 2011; Kelly *et al.* 2014). Crucially, Nab2 physically and functionally interacts

with Fmr1, the *Drosophila* homolog of the Fragile X Syndrome RBP FMRP (Verkerk *et al.* 1991;

Ashley *et al.* 1993; Wan *et al.* 2000), to support axonal morphology and olfactory memory (Bienkowski

et al. 2017). Previous data suggest functions of *Drosophila* Nab2 in poly(A) tail length control,

translational regulation, and mRNA splicing, but mechanistic demonstrations of its molecular function

on individual, endogenous transcripts have yet to emerge (Pak *et al.* 2011; Kelly *et al.* 2014; Bienkowski

et al. 2017; Jalloh *et al.* 2020). Such demonstrations have been prevented in large part because very few

Drosophila Nab2-associated RNAs have been identified (Bienkowski *et al.* 2017; Jalloh *et al.* 2020),

and a comprehensive accounting of Nab2-associated RNAs has yet to be conducted.

While the precise molecular function of *Drosophila* Nab2 on its associated transcripts is

unknown, informed hypotheses may be drawn by synthesizing research on *Drosophila* Nab2 and

orthologs murine ZC3H14, human ZC3H14, and *S. cerevisiae* Nab2, the most well-studied

Nab2/ZC3H14 ortholog (reviewed in Fasken *et al.* 2019). In *S. cerevisiae*, Nab2 functions pervasively

across many RNAs in transcript stability and transcription termination, and it likely acts similarly

broadly in poly(A) tail length control and poly(A) RNA export (Schmid *et al.* 2015; Fasken *et al.* 2019;

Alpert *et al.* 2020). Mutation of *S. cerevisiae* Nab2 induces dramatic increases in bulk poly(A) tail

length and disrupts bulk poly(A) export from the nucleus (Green *et al.* 2002; Kelly *et al.* 2010).

Consistent with its pervasive effects on many transcripts, *S. cerevisiae* Nab2 exhibits a broad binding

target profile and is essential for cellular viability (Anderson *et al.* 1993; Tuck and Tollervey 2013). By

contrast, mutant analyses of metazoan Nab2/ZC3H14 imply increased RNA target specificity for these

proteins. Unlike Nab2 in *S. cerevisiae*, full-length ZC3H14 in mice and humans is not essential for

viability—instead, loss of ZC3H14 decreases viability in mice and causes neurological or

neurodevelopmental defects in both organisms (Pak *et al.* 2011; Rha *et al.* 2017b; Al-Nabhani *et al.*

2018). Bulk poly(A) tail lengths increase upon Nab2 loss in *Drosophila* or full-length ZC3H14 loss in

mice *in vivo*, but this increase is not observed across all mouse tissues or all individual *Drosophila*

mRNAs tested, and it is less pronounced than the effects observed in *S. cerevisiae* (Kelly *et al.* 2010;

Bienkowski *et al.* 2017; Rha *et al.* 2017b). Moreover, in *Drosophila* and mouse cells, respectively, a

pervasive nuclear poly(A) export defect is not observed upon Nab2 loss or ZC3H14 knockdown (Farny

et al. 2008; Pak *et al.* 2011; Kelly *et al.* 2014). *Drosophila* Nab2 is required for proper splicing of

individual introns and exons, but in a small, specific set of transcripts, including *Sex lethal* (Jalloh *et al.*

2020). Taken together, these data are consistent with a focused role for *Drosophila* Nab2 in regulating

poly(A) tail length, splicing, stability, and nuclear export crucial for certain transcripts, cell types, and

developmental contexts (Bienkowski *et al.* 2017; Rha *et al.* 2017b; Jalloh *et al.* 2020). Crucially

however, the theme of *Drosophila* Nab2 RNA target specificity implied by these data has not been

tested and remains an important open question, especially as the polyadenosine affinity of *Drosophila*

Nab2 (Pak *et al.* 2011) makes it theoretically capable of associating with all polyadenylated transcripts

through their poly(A) tails. Thus, a comprehensive identification of *Drosophila* Nab2-associated RNAs

is necessary to determine the potential scope of Nab2 function and provide sets of transcripts on which

the molecular consequences of Nab2-RNA association may be systematically evaluated. In the present

study, in response we define the first neuronal RNA interactome for Nab2.

Contextualizing Nab2-RNA associations requires further definition of the molecular pathways

and proteins, particularly other RBPs, that Nab2 interacts with or regulates. Notably, the *Nab2* modifier

eye screen that initially linked Nab2 and Fmr1 (Bienkowski *et al.* 2017) also recovered an allele of

Ataxin-2 (*Atx2*), which encodes a conserved RBP and regulatory partner of Fmr1 in *Drosophila* neurons

(Sudhakaran *et al.* 2014; Jiménez-López and Guzmán 2014). The shared connection of Nab2 and Atx2

with Fmr1 raised the possibility of cooperation or competition between these two proteins. Underscoring

the value of this approach, Atx2 is a protein of particular importance for human health and neuronal

function. Expansion of a polyglutamine tract within ATXN2, the human Atx2 ortholog, gives rise to the

autosomal dominant neurodegenerative disease spinocerebellar ataxia type 2 (SCA2) (Imbert *et al.*

1996; Pulst *et al.* 1996; Sanpei *et al.* 1996). Expansions of the same tract are also associated with

parkinsonism and amyotrophic lateral sclerosis (ALS) (Gwinn-Hardy *et al.* 2000; Elden *et al.* 2010; Park

et al. 2015). Functionally, *Atx2* encodes a conserved RNA-binding protein that regulates protein

translation, mRNA stability, and mRNP granule formation and plays roles in memory, cellular

metabolism, and circadian rhythms (reviewed in Ostrowski *et al.* 2017; Lee *et al.* 2018). Among the

most well-studied molecular roles of Atx2 are its contributions to regulation of mRNA translation in the

cytoplasm. Specifically, Atx2 suppresses the translation of some target RNAs through RNP granule

formation and interactions with the RNAi machinery (McCann *et al.* 2011; Sudhakaran *et al.* 2014;

Bakthavachalu *et al.* 2018) and supports the translation of other targets by promoting RNA

circularization (Lim and Allada 2013; Zhang *et al.* 2013; Lee *et al.* 2017). Intriguingly Atx2, like Nab2,

contributes to poly(A) tail length control in *S. cerevisiae*—the yeast Atx2 ortholog Pbp1 promotes

poly(A) tail length, likely by inhibiting the activity of poly(A) nuclease (PAN) (Mangus *et al.* 1998,

2004). The shared connections of Nab2 and Atx2 to Fmr1, neuronal translation, and poly(A) tail length

control emphasize the potential for and need to test whether these RBPs functionally interact beyond the

initial eye screen link.

Here, after expanding the genetic link previously identified between Nab2 and Atx2 in our

modifier screen, we used genetic and molecular approaches to probe the functional connections between

these two RBPs. We show that Nab2 and Atx2 functionally interact to control neuronal morphology of

the mushroom bodies (MBs), a learning and memory center of the *Drosophila* brain (Heisenberg 2003;

Kahsai and Zars 2011; Yagi *et al.* 2016; Takemura *et al.* 2017). We then present the first high-

throughput identification of Nab2- and Atx2-associated RNAs in *Drosophila*; in fact, such accounting

has been performed for Nab2 only in *S. cerevisiae*, not in any metazoan (Guisbert *et al.* 2005; Batisse *et*

al. 2009; Tuck and Tollervey 2013; Baejen *et al.* 2014). This approach demonstrates Nab2 and Atx2

associate with an overlapping set of RNA transcripts in fly brains and provides insight into the functions

of each protein individually and in concert with one another. Considering these data as a whole, we

propose a model in which the genetic interactions between Nab2 and Atx2 are explained by their

counterbalanced regulation of shared associated RNAs. Our data represent a valuable resource for

understanding the neuronal roles of Nab2 and Atx2 in *Drosophila* and, potentially, for understanding

links between each RBP and human disease.

Materials and Methods

***Drosophila* genetics and husbandry**

Genetic crosses of *Drosophila melanogaster* were raised on standard media and maintained at 25°C in

humidified incubators (SRI20PF, Shel Lab) with 12-hour light-dark cycles unless otherwise specified.

Cultures were often supplemented with granular yeast (Red Star Yeast) to encourage egg laying.

Parental stocks were maintained at either at room temperature (RT) or 18°C to control virgin eclosion

timing. Stocks used include $Nab2^{ex3}$ (a *Nab2* null), $Nab2^{pex41}$ (a P-element excision control serving as a

Nab2 wild type), and *UAS>Nab2-FLAG*, all first described in (Pak *et al.* 2011). Additional stocks used

include *GMR-Gal4* (on chromosome 2), $Atx2^{X1}$ (an *Atx2* null, gift of N. Bonini) (Satterfield *et al.* 2002),

and *UAS>Atx2-3xFLAG* (gift of R. Allada) (Lim and Allada 2013). Finally, stocks sourced from the

Bloomington Drosophila Stock Center (BDSC) include: *elav>Gal4* (*elav^{c155}*, BL458) (Lin and

Goodman 1994), *OK107-Gal4* (BL854) (Connolly *et al.* 1996), *Df(3R)Exel6174* (BL7653) (Parks *et al.*

2004), *UAS>Nab2* (*Nab2^{EP3716}*, BL17159) (Rørth *et al.* 1998; Bellen *et al.* 2004), and *Atx2^{DG08112}*. The

Atx2^{DG08112} stock (Huet *et al.* 2002) was mapped as part of the Gene Disruption Project (GDP) (Bellen *et*

al. 2004) and is no longer available from the BDSC; copies provided upon request.

Drosophila **eye imaging**

Drosophila eyes were imaged using a Leica MC170 HD digital camera mounted on a Nikon SMZ800N

stereo microscope at 8X magnification. To prepare subjects for imaging, flies were flash frozen (−80°C,

1 minute), fixed in place on a clear Slygard pad using minutien pins (26002-10, Fine Science Tools), and

submerged in 70% ethanol to diffuse light and reduce glare. Subjects were illuminated with a fiber optic

ring light (Dolan-Jenner) and LED illuminator (Nikon Instruments Inc.) and image acquisition was

performed using the Leica Application Suite (v4.12) for Windows under the following parameters: 140

ms exposure; automatic white balance; highest available resolution; and default values for gain,

saturation, gamma, and hue. Each subject was imaged at multiple focal planes (often ≥ 10), and these

were subsequently combined using the *Auto-Align* and *Auto-Blend* functions in Photoshop CS5.1

Extended (Adobe) to generate final, merged images in which the entire subject is in-focus. These "focus

stacking" processing steps (Patterson) combine only in-focus regions of an image series into a single,

merged image.

Immunofluorescence

For mushroom body morphology experiments, *Drosophila* brains were dissected using methods similar

to those in (Williamson and Hiesinger 2010; Kelly *et al.* 2016, 2017). Briefly, using #5 Dumont fine

forceps (Ted Pella, Inc.), for each dissection a *Drosophila* head was isolated in PBS (often supplemented

with 0.1% Triton X-100), the proboscis was removed to provide a forceps grip point, and the remaining

cuticle and trachea were peeled away from the brain within. On wet ice, dissected brains were fixed in

4% paraformaldehyde for 30 minutes and then permeabilized in 0.3% PBS-Triton (PBS-T) for 20

minutes. For both primary and secondary antibody incubations, brains were left rocking at 4°C for 1-3

nights in 0.1% PBS-T supplemented with blocking agent normal goat serum (Jackson ImmunoResearch)

at a 1:20 dilution. Immunostained brains were mounted on SuperFrost Plus slides (12-550-15, Fisher

Scientific) in Vectashield (H-1000, Vector Laboratories) using a cover slip "bridge" method (Kelly *et al.*

2017). Brains were imaged on a Zeiss LSM 510 confocal microscope. Exclusively female flies were

dissected for practicality, given that *Nab2^{ex3}* nulls were analyzed in this experiment and *Nab2^{ex3}* adult

viability skews towards females (Jalloh *et al.* 2020).

For Nab2-Atx2 localization experiments, whole animals were fixed in 4% paraformaldehyde,

0.008% PBS-T, shaking, for 3 hours at RT and then washed in PBS and stored at 4°C overnight. Brains

were dissected in 0.008% PBS-T using similar methods as described above, permeabilized by shaking in

0.5% PBS-T overnight at 4°C, and blocked by shaking in 0.5% PBS-T, 5% NGS for 2 hours at RT. For

both primary and secondary antibody/Hoechst incubations, brains were left shaking at 4°C for 2-3 nights

in 0.5% PBS-T, 5% NGS. After washing with 0.5% PBS-T followed by PBS, brains were mounted in

SlowFade Gold Antifade Mountant (S36936, Invitrogen), surrounded by an adhesive imaging spacer

(GBL654002, Sigma-Aldrich) to prevent sample compression, and finally cover-slipped and sealed with

clear nail polish. Brains were imaged on an A1R HD25 confocal microscope (Nikon) and a multi-photon

FV1000 laser-scanning microscope (Olympus).

Primary antibodies and dilutions used are as follows: mouse α-Fasciclin 2 (1:50) (1D4,

Developmental Studies Hybridoma Bank), rabbit α-GFP (1:400) (A11122, Invitrogen), and mouse α-

FLAG (1:500) (F1804, Sigma-Aldrich). Secondary antibodies and dilutions used are as follows: goat α-

mouse Cy3 (1:100) (Jackson ImmunoResearch), goat α-mouse Alexa 594 (1:400) (A11032, Invitrogen)

and goat α-rabbit Alexa 488 (1:400) (A11008, Invitrogen). To fluoresce DNA and mark nuclei in

localization experiments, brains were also incubated with a Hoechst 33342 stain (1:1,000) (H21492,

Invitrogen) during secondary antibody incubation.

Further brain image analysis and processing, including generating maximum intensity

projections and focus stacks and adjusting brightness and contrast, was performed with Photoshop

CS5.1 Extended (Adobe) and Fiji (Schindelin *et al.* 2012), a distribution of ImageJ (Schneider *et al.*

2012; Rueden *et al.* 2017).

Immunoprecipitation

This immunoprecipitation protocol was developed through optimization guided by the protocols

presented in (Yang *et al.* 2005; Banerjee *et al.* 2017; Bienkowski *et al.* 2017; Morris and Corbett 2018).

Nuclear Isolation Buffer (NIB; 10 mM Tris HCl pH 7.4, 10 mM NaCl, 3 mM $MgCl_2$, 0.5% NP-40) and

Immunoprecipitation Buffer (IP Buffer; 50 mM HEPES, 150 mM NaCl, 5 mM EDTA, 0.1% NP-40)

were prepared ahead of the experiment and stored indefinitely at 4°C. Both buffers, and the glycine and

PBS solutions below, were prepared primarily in 0.1% diethyl pyrocarbonate (DEPC)-treated and

autoclaved ultrapure Milli-Q water to limit RNase contamination. Both NIB and IP Buffer were

supplemented with an EDTA-free cOmplete protease inhibitor cocktail tablet (1 tablet/28 ml;

11873580001, Roche) and RNasin Plus RNase inhibitor (0.2%; N2615, Promega) freshly before each

experiment. Additionally, before each experiment Protein G-coupled magnetic Dynabeads (10003D,

Thermo Fisher) were conjugated to glycerol-free (Domanski *et al.* 2012) monoclonal α-FLAG (F3165,

Sigma-Aldrich) in aliquots of 1.5 mg beads/9 µg antibody by incubation for 45 minutes at room

temperature. Throughout the experiment, beads were magnetized using DynaMag-Spin magnets (e.g.

12320D, Thermo Fisher) as necessary. Exclusively female flies were used for consistency with MB

experiments and for practicality, as both *elav>Nab2-FLAG* and *elav>Atx2-3xFLAG* prohibitively

decreased relative male viability (data not shown), presumably due to deleterious effects in males likely

driven by dosage compensation of the X-chromosome-linked *elav>Gal4* construct leading to enhanced

epitope-tagged protein overexpression.

300 female *Drosophila* heads each of the genotypes *elav>Gal4* alone, *elav>Nab2-FLAG*, and

elav>Atx2-3xFLAG, previously isolated in bulk (see *Supplemental Materials and Methods*), were fixed

in 1% formaldehyde, 0.1% NP-40 in PBS for 30 minutes at 4°C. Fixation was quenched by adding

glycine to a final concentration of 250 mM and rocking for 10 minutes at 4°C. Heads were washed in

0.1% NP-40 in PBS and then manually homogenized with a smooth Teflon pestle for 5 minutes in 250

µL of NIB in a size AA glass tissue grinder at 4°C (3431D70, Thomas Scientific). Homogenates were

spun through 35 µm cell strainer caps into round-bottom tubes (352235, Falcon) to remove exoskeletal

debris, transferred, and then centrifuged for 5 minutes at 500×g at 4°C to separate an insoluble fraction.

Twenty percent of the soluble supernatant volume was isolated and defined as Input; the remaining

eighty percent was used for immunoprecipitation. Both Input and IP samples were diluted to final

concentrations of 0.8x IP Buffer to ensure comparable and efficient sample lysis. IP samples were

transferred onto the α-FLAG-conjugated magnetic Dynabeads, and both sample types were incubated,

rotating, for 10 minutes at room temperature. Next, IP sample supernatant was collected as the Unbound

fraction, and IP sample beads were washed three times in IP Buffer. Finally, IP sample beads were

resuspended in IP Buffer, transferred to clean tubes, and stored along with Input samples overnight at

4°C to allow passive hydrolysis to partially reverse formaldehyde crosslinks. This protocol was applied

for both protein co-immunoprecipitation and RNA immunoprecipitation.

For protein co-immunoprecipitation, harsh elution of protein from IP sample beads was

accomplished the next day—IP samples were diluted in modified Laemmli Sample Buffer (Laemmli

1970), incubated at 98°C for 5 minutes, centrifuged at 16,100×g for 5 minutes at room temperature, and

magnetized to collect beads. Sample supernatants were then collected as IP samples. In parallel, Input

samples were concentrated using an acetone-based method; this step was required for subsequent

immunoblot analysis. Input samples were diluted to generate 80% chilled acetone solutions, vortexed for

15 seconds, and incubated at −20°C for 60 minutes. Samples were centrifuged at 14,000×g for 10

minutes at room temperature, resulting supernatants were discarded, and most remaining acetone was

evaporated by air drying protein pellets in open tubes for 30 seconds at room temperature. To solubilize

these dried protein pellets, samples were suspended in a solution equal parts modified Laemmli Sample

Buffer (Laemmli 1970) and IP Buffer, vortexed, sonicated for 3x5 minutes in a 4°C Bioruptor

ultrasonicator (UCD-200, Diagenode), vortexed, and heated at 98°C for 10 minutes. Finally, remaining

insoluble material was collected by centrifugation at 16,100×g for 5 minutes at room temperature.

Associated supernatants were isolated as concentrated Input protein samples. For RNA

immunoprecipitation, harsh elution of RNA from IP sample beads was accomplished the next day with

Trizol—both IP and Input samples were subjected to the RNA extraction protocol detailed below.

RNA Extraction

Following immunoprecipitation, RNA was isolated from IP and Input samples using a TRIzol-column

hybrid approach adapted from (Rodriguez-Lanetty). To account for volume differences, samples were

vigorously homogenized in TRIzol reagent (15596018, Thermo Fisher) at a ratio of either 1:10 (IP

sample:TRIzol) or 1:3 (Input sample:TRIzol) and then incubated for 5 minutes at room temperature. All

homogenized samples were clarified by centrifugation at 12,000×g at 4°C for 5 minutes, IP samples

were magnetized to collect beads, and supernatant was isolated from all samples. After adding

chloroform at a ratio of 0.2:1 (choloroform:TRIzol), samples were manually shaken and incubated at

room temperature for 3 minutes. Samples were phase separated by centrifugation at 12,000×g at 4°C for

15 minutes, after which the aqueous layer was carefully isolated and mixed with an equal volume of

100% ethanol. RNA was further purified using an RNeasy Mini Kit (74106, QIAGEN) according to the

manufacturer's instructions (RNeasy Mini Handbook, 4[th] Ed., June 2012) with the following deviations:

for each sample, a final 30 µL elution was performed twice, isolating 60 µL of RNA in total into each

collection tube. An on-column DNase digestion step was also performed under the same instructions

using an RNase-Free DNase Set (79254, QIAGEN). Final RNA concentration and sample purity were

determined via a NanoDrop 1000 spectrophotometer (Thermo Fisher).

RNA Sequencing

RNA from twelve samples of 300 adult female *Drosophila* heads each was isolated via the

immunoprecipitation and extraction protocols described above, generating twelve pairs of IP and Input

samples, or twenty-four samples in total. These samples were composed of four biological replicates

each of *elav>Gal4* alone, *elav>Nab2-FLAG*, and *elav>Atx2-3xFLAG*. Once obtained, RNA samples

were transferred on dry ice to the Georgia Genomics and Bioinformatics Core at UGA for library

preparation and sequencing. There, IP samples were first concentrated using solid phase reversible

immobilization (SPRI) beads. Then, the TruSeq Stranded Total RNA Library Prep Gold kit (20020598,

Illumina) was used to deplete rRNA and prepare stranded cDNA libraries from all twenty-four samples.

These uniquely barcoded cDNA libraries were then pooled by sample type, forming one IP library pool

and one Input library pool. Each pool was sequenced on a separate NextSeq High Output Flow Cell

(Illumina) for 150 cycles to generate paired-end, 75 base-pair (bp) reads. Total non-index sequencing

yield across all IP samples was 88.49 Gbp, equivalent to about 1.2 billion reads in total and 98 million

reads per sample. Total non-index sequencing yield across all Input samples was 83.25 Gbp, equivalent

to about 1.1 billion reads in total and 93 million reads per sample. Sequencing accuracy was high;

87.83% and 91.38% of non-index reads for IP and Input samples, respectively, have a sequencing

quality (Q) score greater than or equal to 30.

RNA Sequencing Analysis—Read Mapping, Differential Expression, Visualization

Following sequencing, raw read FASTA files were transferred to Emory for bioinformatic analysis. To

start, analyses were conducted on the Galaxy web platform, specifically using the public server at

usegalaxy.org (Afgan *et al.* 2018). This analysis was supported by the BDGP6.22 release of the

Drosophila melanogaster genome (Hoskins *et al.* 2015)—both the raw sequence FASTA and the gene

annotation GTF were downloaded from release 97 of the Ensembl database (Yates *et al.* 2020) and used

as inputs in subsequent read mapping, annotation, and visualization steps. For each Galaxy tool

described below, exact parameters and version numbers used are detailed in Supplemental Table 1. For

each sample, reads from across all four NextSeq flow cell lanes were concatenated using the Galaxy

Concatenate datasets tail-to-head tool and mapped using RNA STAR (Dobin *et al.* 2013). Mapped

reads were then assigned to exons/genes and tallied using *featureCounts* (Liao *et al.* 2014). To enable

inter-sample read count comparisons, count normalization and differential expression analysis was

conducted using *DESeq2* (Love *et al.* 2014). Importantly, *DESeq2* analysis was performed twice, once

on the 12 IP samples and once on the 12 Input samples; see *Supplemental Materials and Methods* for

discussion of this sample separation method.

Outputs from all of the above tools were downloaded from Galaxy for local analysis,

computation, and visualization. Custom R scripts were written to generate the scatterplots and

hypergeometric test reported here and are available in File S3. Scripts in the R programming language

(R Core Team 2019) were written and compiled in RStudio (R Studio Team 2018). Additional R

packages used in these scripts include ggplot2 (Wickham 2016), ggrepel (Slowikowski 2019),

BiocManager (Morgan 2018), and DESeq2 (Love *et al.* 2014). Analyses were supported by bulk data

downloads along with extensive gene-level annotation, sequence information, and references provided

by Flybase (Thurmond *et al.* 2018). Principal component analysis was conducted by and reported from

the above *DESeq2* assessment on Galaxy. Mapped reads were visualized in the Integrative Genomics

Viewer (IGV) (Robinson *et al.* 2011) on the same version of the *D. melanogaster* genome used above.

Gene-by-gene one-way ANOVAs to identify significantly enriched (i.e. RBP-associated)

transcripts

Gene-by-gene ANOVAs and post-hoc tests for the 5,760 genes identified in the "testable" set, along

with bar graphs of IP/Input values, were generated in Prism 8 for Windows 64-bit (GraphPad Software).

Custom R and PRISM scripts were written to generate and label the 5,760 PRISM data tables, one per

testable gene, required for this analysis, and custom R scripts were written to extract and combine the

outputs from each test; these scripts are all available in File S3. See *Results* for a summary and below for

a further detailed discussion of the statistical testing used to define the testable transcript set and identify

significantly enriched (i.e. RBP-associated) transcripts in our RIP-Seq results.

To identify RNA targets of Nab2 and Atx2—that is, RNAs enriched in either Nab2 RIP or Atx2

RIP samples relative to control RIP—directly comparing normalized read counts between RIP samples

is insufficient. Differences in RNA expression between samples must be accounted for, as these

differences can partially or wholly explain differences in the amount of RNA isolated by IP. We

employed a common solution to this problem used in RIP- and ChIP-qPCR (Zhao *et al.* 2010; Aguilo *et*

al. 2015; Li *et al.* 2019), scaling normalized RIP reads for each gene in each sample by the

corresponding number of normalized Input reads. For clarity, we describe these values as "IP/Input"—

they are commonly referred to as "Percent Input" or "% Input." These IP/Input values could then be

compared between samples, further normalizing them to *elav-Gal4* alone controls. In this way, RIP fold

enrichment, appropriately normalized to library size/composition *and* gene expression, were calculated

for each gene in each sample. To promote the reliability of our analyses and increase our statistical

power to detect differences in fold enrichment, we limited further analyses to a testable set of 5,760

genes out of the 17,753 total genes annotated in the BDGP6.22 genome. The testable gene set was

defined as having detectable expression in all twelve Input samples and an average normalized read

count in either Nab2 or Atx2 RIP samples greater than 10. These criteria were based on those used in

(Lu *et al.* 2014; Malmevik *et al.* 2015). In this defined gene set, differences in fold enrichment were

statistically tested using gene-by-gene one-way ANOVAs (Li *et al.* 2019) in Prism 8 (GraphPad

software), applying Dunnett's post-hoc test to calculate significance *p*-values only for the comparison of

each experimental sample to the control sample (Dunnett 1955). In each case, *p*-values were adjusted to

correct for multiple hypothesis testing only within each gene-by-gene ANOVA. We identified a small,

focused set of statistically significantly enriched RNAs using this approach and concluded that

additional corrections across all genes to control type I error (i.e. false positives) are not necessary

(Rothman 1990). In fact, in the analyses above we determined that rRNA depletion during our RIP-Seq

library preparation was incomplete, resulting in comparatively low read depth. Thus, rather than failing

to adequately control type I error, we strongly suspect the RBP-associated transcripts we identified

through this approach represent an undercount, to be expanded in future studies by methods with higher

sensitivity (e.g. CLIP-Seq).

RNA Sequencing Analysis—Sequence Motif Analyses

Sequence motif analyses were conducted using the MEME Suite of software tools, accessed through the

web interface at meme-suite.org (Bailey *et al.* 2009). For each MEME Suite tool described below, exact

parameters and version numbers used are detailed in Supplemental Table 1. Within the MEME Suite, we

used MEME itself (Bailey and Elkan 1994) to scan all Nab2-associated transcripts, regardless of their

association with Atx2, to 1) identify sequence motifs shared across multiple transcripts and 2) evaluate

the frequency and statistical significance of the discovered sequence motifs. Next, FIMO (Grant *et al.*

2011) was used to quantify the frequency among 1) Nab2-associated transcripts and 2) non-Nab2

associated transcripts of user-provided sequences, specifically i) a 41-bp A-rich motif identified in

Nab2-associated transcripts by MEME, ii) A_{12}, and iii) $A_{11}G$. Non-Nab2-associated transcripts are

defined as all 5,619 transcripts in the testable set found to not be statistically significantly associated

with Nab2 by RIP-Seq. Sequence logos (i.e. visual representations of weighted sequence motifs) were

generated by MEME and by WebLogo 3.7.4, available at weblogo.threeplusone.com (Crooks *et al.*

2004).

Importantly, for any Nab2-associated or non-Nab2 associated transcripts annotated with multiple

splice variants, all variant sequences were included as inputs in our motif analyses. This inclusion

reflects an inherent limitation of standard shotgun—that is, short-read—sequencing, as most reads

cannot be unambiguously assigned to one splice variant of a given gene, only to given exon(s) encoded

by that gene. We therefore chose this inclusion strategy to avoid introducing any bias associated with

attempting to call single splice variants for RBP association, and for analytical simplicity. Full

sequences of Nab2-associated and non-Nab2 associated transcripts were obtained using the FlyBase

Sequence Downloader at flybase.org/download/sequence/batch/ (database release FB2020_04).

Data Availability

The authors affirm that all data necessary for confirming the conclusions of the article are present within

the article and associated figures, tables, supplemental materials, and database accessions. File S1

contains *Supplemental Materials and Methods*, including those focused on bulk *Drosophila* head

isolation, immunoblotting, *DESeq2*-based count normalization, and Gene Ontology analyses. File S2

contains detailed legends for all supplemental tables. File S3 contains all custom code—both R and

PRISM scripts—written to generate, analyze, or visualize data in this article and associated figures,

tables, and supplemental materials. Sequencing data, including raw reads, processed counts, and

statistical analyses for each individual RIP-Seq sample, are available at the Gene Expression Omnibus

(GEO) under accession: GSE165677. *Drosophila* stocks are available upon request. Supplemental

materials, including files, figures, and tables, are available at figshare:

https://figshare.com/s/6f28676d7119624b3105.

Results

Atx2 **loss-of-function alleles suppress Nab2 overexpression phenotypes in the adult eye**

Previous work has established a Gal4-driven Nab2 overexpression system in the *Drosophila* eye as an

effective screening platform to identify potential regulatory partners and targets of Nab2 (Pak *et al.*

2011; Bienkowski *et al.* 2017; Lee *et al.* 2020). This approach uses the *Glass Multimer Reporter* (*GMR*)

construct (Ellis *et al.* 1993; Hay *et al.* 1994) to drive expression of the *S. cerevisiae* Gal4 transcription

factor in fated eye cells (Freeman 1996). In turn, Gal4 binds to *Upstream Activating Sequence* (*UAS*)

sites within an EP-type P-element (Rørth 1996) inserted upstream of the endogenous *Nab2* gene

(*EP3716*) and induces eye-specific overexpression of endogenous Nab2 protein (a genotype hereafter

referred to as *GMR>Nab2*). *GMR>Nab2* produces a consistent array of eye morphological defects

compared to the *GMR-Gal4* transgene control (Pak *et al.* 2011; Bienkowski *et al.* 2017; Lee *et al.* 2020)

and (Figure 1A,B). Specifically, this misexpression causes loss of posterior eye pigment, sporadic

blackened patches, and disruptions to ommatidial organization lending the surface of the eye a "rough"

appearance. Notably, *GMR>Nab2*-induced pigment loss increases in severity along the anterior-to-

posterior axis of the eye, likely because *GMR* activation occurs behind the morphogenetic furrow, the

posterior-to-anterior wave of eye morphogenesis observed in the larval eye disc (Wolff and Ready 1991;

Hay *et al.* 1994). As a result, posterior *GMR>Nab2* eye cells experience the longest period of Nab2

overexpression.

Using the *GMR>Nab2* modifier screen as a foundation, we previously identified the *Drosophila*

Fragile X Syndrome RBP and neuronal translational regulator Fmr1 as a physical and functional

interactor of Nab2 (Bienkowski *et al.* 2017). An allele of the *Ataxin-2* (*Atx2*) gene, which encodes an

RNA binding protein that is a regulatory partner of Fmr1 in *Drosophila* (Sudhakaran *et al.* 2014), was

also detected in eye this screen as a candidate *GMR>Nab2* modifier (Bienkowski *et al.* 2017). To pursue

this potential Nab2-Atx2 link, we tested two *Atx2* alleles for genetic interactions with *GMR>Nab2*. The

first allele, *Atx2^{DG08112}*, is caused by the insertion of a 15.6 kb *{wHy}* P-element near the 5' end of *Atx2*

(Huet *et al.* 2002; Bellen *et al.* 2004) and is lethal *in trans* to *Df(3R)Exel6174*, a deletion that completely

removes the *Atx2* locus and nearby genes (Parks *et al.* 2004). That is, crossing balanced *Atx2^{DG08112}* and

Df(3R)Exel6174 alleles produces no *trans* heterozygotes among other F1 progeny (n=54). Based on

these data, we interpret *Atx2^{DG08112}* to be a strong hypomorph. The second *Atx2* allele, *Atx2^{X1}*, is a 1.4 kb

imprecise-excision-based deletion that removes the first 22 codons of the *Atx2* coding sequence and that

has been characterized as a null (Satterfield *et al.* 2002). In part because Nab2 loss induces some sex-

specific defects (Jalloh *et al.* 2020), we analyzed each sex individually. In adult females, heterozygosity

for either of these two loss-of-function alleles, *Atx2^{DG08112}* (Figure 1C) or *Atx2^{X1}* (Figure 1D),

dominantly suppresses the pigment loss and blackened patches caused by *GMR>Nab2*. In contrast, both

Atx2 alleles have limited impact on ommatidial organization or "roughness". In males, *GMR>Nab2*

induces morphological eye defects (Figure 1 E,F) comparable to those in females, and similarly

heterozygosity for either *Atx2^{DG08112}* (Figure 1G) or *Atx2^{X1}* (Figure 1H) dominantly suppresses the

pigment loss and blackened patch defects.

***Atx2* loss-of-function alleles suppress *Nab2* null effects on adult viability and brain morphology**

Misexpression of Nab2 induces dramatic phenotypes in domains beyond the eye; homozygosity for the

null allele *Nab2^{ex3}* causes a dramatic reduction in adult viability (Pak *et al.* 2011). Thus, to explore

whether modifying effects of *Atx2* loss-of-function alleles extend to the endogenous *Nab2* locus, we

analyzed the effect of *Atx2* heterozygosity on low adult viability in *Nab2^{ex3}* homozygotes (Supplemental

Figure 1). As in the eye, both the *Atx2^{DG08112}* and *Atx2^{X1}* alleles dominantly suppress the viability defects

observed in *Nab2^{ex3}* females, elevating adult viability from 17% to 39% and 82%, respectively (Figure

1I). The corresponding effect in males is not as penetrant; only the null *Atx2^{X1}* allele dominantly

suppresses the viability defect in *Nab2^{ex3}* males (Figure 1J). Taken together, these data establish gross

similarities in *Nab2-Atx2* genetic interactions in females and males. Thus, for practicality we focused

further experiments exclusively on female flies, given the more prohibitive impact on male viability of

changes in Nab2 expression (Jalloh *et al.* 2020 and see *Materials and Methods*).

421   That *Atx2* loss-of-function alleles improve adult viability of *Nab2^{ex3}* homozygotes suggests Atx2

and Nab2 coregulate processes or transcripts important for adult development or survival. However,

these genetic interactions do not reveal in what cell types or tissues this coregulation may occur. We

therefore focused further investigations of *Nab2-Atx2* interaction in the brain, given the established and

important roles of each protein in brain neurons (Lim and Allada 2013; Sudhakaran *et al.* 2014; Kelly *et*

al. 2016; Bienkowski *et al.* 2017). *Nab2^{ex3}* homozygous flies develop morphological defects in the axon

tracts—lobes—of the mushroom body (MB) brain structure, a principal olfactory learning and memory

center of the insect brain (Heisenberg 2003; Kahsai and Zars 2011; Yagi *et al.* 2016; Takemura *et al.*

2017). Specifically, the MBs of surviving *Nab2^{ex3}* homozygous null adults exhibit two highly penetrant

structural defects: thinning or absence of the dorsally-projecting α lobes and over-projection or "fusion"

of the medially-projecting β lobes (Kelly *et al.* 2016). We found that heterozygosity for either

Atx2^{DG08112} or *Atx2^{X1}* also causes defects in MB morphology—specifically β lobe fusion—with no

apparent effects on α lobe morphology as compared to controls (Figure 2A-C). Importantly, in the

background of *Nab2^{ex3}* nulls (Figure 2D), heterozygosity for either *Atx2^{DG08112}* (Figure 2E) or *Atx2^{X1}*

(Figure 2F) suppresses the thinning or absence of α lobes, decreasing the penetrance of this phenotype

from 62% of α lobes to 30% or 36%, respectively (Figure 2G). In contrast, neither *Atx2* allele

significantly affects the penetrance of β lobe fusion in *Nab2^ex3* nulls, demonstrating the effect of each

mutation is not additive to the effect of *Nab2^ex3* homozygosity in this context (Figure 2H). A similar α-

lobe-specific interaction occurs between alleles of *Nab2* and *Fmr1* (Bienkowski *et al.* 2017). Notably, as

α and β lobes are composed of tracts of bifurcated axons from single cells (Takemura *et al.* 2017), this

α-lobe-specific suppression by *Atx2* alleles demonstrates a *Nab2-Atx2* genetic interaction at subcellular

resolution. Moreover, that *Atx2* loss-of-function alleles suppress defects of a *Nab2* null allele implies

that Atx2 and Nab2 proteins may coregulate, but in opposing ways, pathways guiding α lobe

morphology during development.

Nab2 and Atx2 primarily localize to independent compartments in mushroom body neurons

The genetic links between *Nab2* and *Atx2* could reflect a physical interaction between their encoded

proteins (e.g. as shared components of mRNP complexes), as has been observed for both Nab2 and Atx2

with Fmr1 (Sudhakaran *et al.* 2014; Bienkowski *et al.* 2017). Alternatively, these genetic links could

reflect functional but not physical interactions between Nab2 and Atx2 on common RNAs or

neurodevelopmental processes. The latter hypothesis aligns with the localization patterns of each

protein—Nab2 localizes primarily to neuronal nuclei with a small fraction in the cytoplasm in some

contexts (Kelly *et al.* 2016; Bienkowski *et al.* 2017), while Atx2 is exclusive to the neuronal cytoplasm

except under certain pathogenic conditions (Lessing and Bonini 2008; Elden *et al.* 2010). To begin to

differentiate between these hypotheses, we evaluated the localization profiles of each protein in MBs *in*

vivo. We expressed both *UAS-Nab2-YFP* and *UAS-Atx2-3xFLAG* transgenes in adult MB Kenyon cells

using the pan-MB driver *OK107-Gal4* (Figure 3A). Similar to observations in human cerebral cortex

tissues (Huynh *et al.* 2003), Atx2 is nearly excluded from nuclei and localizes strongly to the soma

cytoplasm of MB Kenyon cells in adults *in vivo*. In contrast, Nab2 localizes predominantly to the nuclei

of these neurons *in vivo*. This distinction extends beyond the soma and into the α and β lobe axon tracts;

Atx2 localizes robustly to these cytoplasmic compartments while Nab2 does not (Supplemental Figure

2).

To more rigorously assess Nab2-Atx2 protein interactions across all cell compartments, we

expressed a FLAG-tagged Nab2 transgene (*UAS-Nab2-FLAG*) (Pak *et al.* 2011) using the pan-neuronal

driver *elav-Gal4* (Lin and Goodman 1994) and subjected brain-neuron-enriched head lysates to

immunoprecipitation with α-FLAG-conjugated beads to recover Nab2-associated proteins. Probing with

specific antibodies confirms that Fmr1 is enriched in Nab2 immunoprecipitates as previously reported

(Bienkowski *et al.* 2017), but reveals weak enrichment of Atx2 (Figure 3B). These results indicate

complexes containing Nab2 and Atx2 may form in neurons but are rare relative to Nab2-Fmr1

complexes. Taken together, these subcellular localization and biochemical data suggest Nab2 and Atx2

do not generally co-occupy the same RNA or mRNP complexes throughout the post-transcriptional life

of an RNA in adult mushroom body neurons. Therefore, we considered the possibility that *Nab2-Atx2*

genetic interactions instead reflect roles in post-transcriptional control of shared RNA targets at different

points in time or different locations in the cell.

The Nab2 and Atx2 RNA interactomes in brain neurons overlap

Neither Nab2- nor Atx2-associated RNAs have been identified by a high-throughput method in

Drosophila—such accounting has been conducted for Atx2 in human cells (Yokoshi *et al.* 2014) and for

Nab2 only in *S. cerevisiae*, not in any metazoan (Guisbert *et al.* 2005; Batisse *et al.* 2009; Tuck and

Tollervey 2013; Baejen *et al.* 2014). To test the hypothesis that Nab2 and Atx2 share RNA targets, we

identified transcripts stably associated with epitope-tagged versions of each protein in adult brain

neurons using an RNA immunoprecipitation-sequencing (RIP-Seq) approach. In this approach, protein

products of *UAS-Nab2-FLAG* or *UAS-Atx2-3xFLAG* transgenes are robustly expressed under *elav-Gal4*

control and are efficiently immunoprecipitated from adult head lysates (Figure 4A). Briefly, four

biological replicates each of *elav-Gal4*, *elav>Nab2-FLAG*, and *elav>Atx2-3xFLAG* adult female

Drosophila heads were lysed and immunoprecipitated with α-FLAG-conjugated beads. Then, RNA from

both IP and Input samples was rRNA depleted, reverse transcribed into stranded cDNA libraries, and

sequenced. Using the Galaxy web platform through the public server at usegalaxy.org (Afgan *et al.*

2018), reads were mapped using STAR (Dobin *et al.* 2013) to the BDGP6.22 release of the *Drosophila*

melanogaster genome (sourced through Ensembl, Yates *et al.* 2020), assigned to exons/genes and tallied

using *featureCounts* (Liao *et al.* 2014), and normalized for inter-library count comparisons using

DESeq2 (Love *et al.* 2014). A principal component analysis (PCA) generated as part of *DESeq2*

demonstrates the high inter-genotype reproducibility among RNA IP (RIP) samples and shows that

samples expressing Nab2-FLAG or Atx2-3xFLAG differ more from *elav-Gal4* controls than from one

another (Figure 4B).

To identify Nab2-associated and Atx2-associated RNAs, we calculated percent input (IP/Input)

enrichment values (Zhao *et al.* 2010; Aguilo *et al.* 2015; Li *et al.* 2019) for each of the 5,760 genes in

the testable set defined by 1) detectable expression in all twelve Inputs and 2) an average normalized

Nab2- or Atx2-IP read count greater than 10 (Lu *et al.* 2014; Malmevik *et al.* 2015). Fold enrichment

differences were statistically tested by performing gene-by-gene one-way ANOVAs (Li *et al.* 2019),

applying Dunnett's post-hoc test (Dunnett 1955), and calculating adjusted *p*-values corrected for

multiple hypothesis testing within each gene-by-gene ANOVA (values hereafter referred to as *Dun. Adj.*

p; see *Materials and Methods* for more detail). Using this approach, we identify 141 and 103 RNAs

significantly enriched in α-FLAG IPs of *elav>Nab2-FLAG* and *elav>Atx2-3xFLAG* female heads,

respectively (Supplemental Table 2, Supplemental Figure 3). The size and focus of these sets of

statistically significantly enriched RNAs suggests type I (i.e. false positive) error is sufficiently

controlled and additional corrections between genes are not necessary (Rothman 1990). Comparing the

Nab2- and Atx2-IP groups strongly supports our hypothesis, revealing 28 transcripts shared between

Nab2- and Atx2-associated *Drosophila* neuronal RNAs (Figure 4C). This overlap is highly significant

according to the hypergeometric test—it is extremely unlikely to occur by random selection from the

total tested gene set. The full list of transcripts associated with both Nab2 and Atx2 (Table 1) includes

multiple mRNAs that encode proteins with functions in neuronal domains in which *Nab2* and *Atx2*

genetically interact, raising the possibility that coregulation of these RNAs by Nab2 and Atx2 partially

explains these *Nab2-Atx2* genetic links. These shared transcripts include *drk* (*downstream of receptor*

kinase), *me31B* (*maternal expression at 31B*), *sm* (*smooth*), and *stai* (*stathmin*). The protein encoded by

drk is a receptor tyrosine kinase (RTK) adaptor that regulates growth and development by binding

activated RTKs, such as sevenless in R7 retinal cells (Almudi *et al.* 2010), and contributes to, among

other processes, cell survival in the eye (Schoenherr *et al.* 2012) and olfactory learning and memory in

the MB (Moressis *et al.* 2009). The protein encoded by *me31B* is a DEAD-box RNA helicase expressed

in many cellular contexts, including the MB Kenyon cells (Hillebrand *et al.* 2010) and the oocyte

(Nakamura *et al.* 2001), that physically associates with Atx2 (Lee *et al.* 2017) and serves as a central

player in miRNA-mediated translational repression (Barbee *et al.* 2006) and assembly of some RNP

granules (Eulalio *et al.* 2007). Finally, the proteins encoded by *sm* and *stai* are respectively an hnRNP

linked to defects in axon termination (Layalle *et al.* 2005) and a tubulin binding protein linked to natural

variation in the size of MB α and β lobes (Lachkar *et al.* 2010; Zwarts *et al.* 2015).

The 28 shared transcripts represent approximately 20% and 24% of the total transcripts identified

as Nab2- and Atx2-associated, respectively, underscoring that these proteins also associate with RNA

sets independent from one another. From these independent sets, we defined the top Nab2-specific and

Atx2-specific associated transcripts as the top 20 most significantly associated transcripts (by *Dun. Adj.*

p) and top 20 most strongly enriched transcripts (by IP/Input) in each set. As with shared RNAs,

multiple RBP-specific RNAs with links to *Nab2* or *Atx2* functions or mutant phenotypes are identified

among these top transcripts, raising the possibility that regulation of these RNAs by Nab2 or Atx2

partially explains the mechanism of action of these RBPs (Figure 4D,E). For example, the top Nab2-

specific associated RNAs include *Arpc2* (*Actin-related protein 2/3 complex, subunit 2*), *side-II* (*sidestep*

II), and *Cpsf160* (C*leavage and polyadenylation specificity factor 160*). These transcripts respectively

encode proteins with proposed functions in neuronal growth cone advance (Yang *et al.* 2012), synapse

formation between certain neuronal subtypes (Tan *et al.* 2015), and mRNA poly(A)-tail formation based

on orthology to mammalian *Cpsf1* (Mandel *et al.* 2008). The top Atx2-specific associated RNAs include

dj-1β, *mtm* (*myotubularin*), and *Snx16* (*Sorting nexin 16*). These transcripts respectively encode proteins

with proposed functions in ATP synthesis and motor neuron synaptic transmission (Hao *et al.* 2010;

Oswald *et al.* 2018), endosomal trafficking regulation via phosphatase activity (Velichkova *et al.* 2010;

Jean *et al.* 2012), and neuromuscular junction synaptic growth (Rodal *et al.* 2011).

Gene Ontology terms enriched in Nab2 and Atx2 RNA interactomes emphasize additional RBP-

associated transcripts

Evaluating Nab2- and Atx2-associated RNAs individually provides valuable but incomplete insight,

allowing larger trends to be missed. To complement these analyses, we holistically evaluated the shared

and specific Nab2- and Atx2-associated transcripts by subjecting each gene list to PANTHER Gene

Ontology (GO) analysis, revealing the identities and members of enriched GO terms in each transcript

set (Ashburner *et al.* 2000; Mi *et al.* 2019; The Gene Ontology Consortium 2019). Critically, GO term

enrichment was calculated by comparing term abundance between these lists and the testable set of

5,760 head-enriched genes rather than the entire genome. In this way, these analyses did not identify GO

terms as enriched simply because of their overrepresentation in *Drosophila* heads. Among shared Nab2-

and Atx2-associated RNAs, we identify overrepresented GO terms and RBP-associated transcripts

within them that highlight crucial functions and processes Nab2 and Atx2 may coregulate (Figure 4F).

Among these GO terms are 'microtubule binding', which includes *apolpp* (*apolipophorin*) and *shi*

(*shibire*); 'sensory perception of taste', which includes *Gαo* and *Gγ30A*; 'gene silencing by miRNA',

which includes *AGO2* (*Argonaute 2*) and *me31B*; and 'short-term memory', which includes *shi* and *drk*.

Survey of the associated RNAs specific to either RBP reveals overrepresented GO terms and transcripts

within them which may mediate processes Nab2 and Atx2 regulate independently of one another,

including respectively the GO terms 'exosomal secretion', which includes *Rab35* and *Rab7*; and

'regulation of ATP metabolic process', which includes *Dg* (*Dystroglycan*) and *dj-1β* (Supplemental

Figure 4).

To combine and summarize the individual transcript and GO analyses, we highlight groups of

seven transcripts found within the shared (Figure 5A) and RBP-specific (Figure 5B,C) associated

transcript sets. These highlights were selected from the combined set of transcripts 1) demonstrating a

fold enrichment (IP/Input) greater than 1.5 and/or 2) included in the most overrepresented GO terms

(fully defined in Supplemental Table 3). Beyond transcripts already described, this summary includes

the shared transcript *HmgZ* (*HMG protein Z*), Nab2-specific transcripts *fwe* (*flower*) and *SLC22A*

(*SLC22A family member*), and Atx2-specific transcripts *tea* (*telomere ends associated*) and *Xpc*

(*Xeroderma pigmentosum, complementation group C*). A group of functionally diverse transcripts in the

testable set that did not associate with either RBP is shown for comparison and as evidence of the

specificity of the RIP-Seq assay (Figure 5D).

Polyadenosine sequence motifs are enriched in Nab2-associated RNAs

The diversity of RNAs that do not associate with Nab2 and Atx2 in the RIP assay (Figure 5D)

underscores a key finding—both of these RBPs exhibit specific RNA-association patterns within brain

neurons. This observation is not surprising for Atx2 given, for example, the sequence specificity of its

human homolog in HEK293T cells (Yokoshi *et al.* 2014), but it represents a valuable insight for Nab2.

The extent of the metazoan Nab2/ZC3H14 RNA target pool has been an enduring question (Rha *et al.*

2017a), given the breadth of the *S. cerevisiae* Nab2 target pool (Batisse *et al.* 2009; Tuck and Tollervey

2013) and the ability of Nab2/ZC3H14 across eukaryotes to bind polyadenosine RNA *in vitro* (Kelly *et*

al. 2007; Pak *et al.* 2011), raising the possibility for very broad binding of mRNAs via their poly(A)

tails *in vivo*. We found a relatively focused set of RNAs co-precipitate with Nab2-FLAG from fly brain

neurons, indicating Nab2 may indeed exhibit greater specificity in *Drosophila* than would be observed if

the protein bound all or most polyadenylated transcripts via their poly(A) tails.

Thus, we sought to determine what additional RNA sequence features may drive the association

of Nab2 with its target transcripts if not only the presence of a poly(A) tail. We used MEME (Bailey and

Elkan 1994) to scan all Nab2-associated transcripts to identify shared sequence motifs that may

represent Nab2 binding sites and partially explain Nab2 specificity. Strikingly, this analysis identifies a

41-bp long, internal-A-rich stretch among the first ten 6-50-bp motifs shared among Nab2-associated

transcripts. Importantly, each of these 10 sequence motifs are shared across overlapping sets of many

but not all Nab2-associated RNAs. Using FIMO (Grant *et al.* 2011), another part of the MEME Suite

(Bailey *et al.* 2009), we quantified the frequency of close and exact matches to the consensus version of

this motif among Nab2-associated RNAs. Occurrences of this A-rich motif are significantly more

common in Nab2-associated transcripts compared to non-Nab2 associated transcripts, respectively

appearing once every 135 bases and once every 845 bases on average, a 6.3-fold enrichment (Figure

6A). The high frequency of this motif in Nab2-associated transcripts is consistent with data from *S.*

cerevisiae that Nab2 does not associate with RNAs exclusively through the poly(A) tail and also binds

to upstream UTRs and coding sequences, likely through other A-rich sequences (Guisbert *et al.* 2005;

González-Aguilera *et al.* 2011; Tuck and Tollervey 2013; Baejen *et al.* 2014; Aibara *et al.* 2017).

Importantly, that this A-rich motif is enriched in but not exclusive to Nab2-associated RNAs is

consistent with results for other RBPs—linear sequence motifs alone are generally insufficient to

explain RBP specificity (Dominguez *et al.* 2018) and RBPs do not generally occupy all of their available

binding motifs throughout the transcriptome (Li *et al.* 2010; Taliaferro *et al.* 2016).

As a complement to these analyses, we used FIMO to scan Nab2-associated RNAs for the

presence of the smallest canonical binding motifs sufficient for Nab2 association in *S. cerevisiae*—A_{12}

and $A_{11}G$ (Guisbert *et al.* 2005; Aibara *et al.* 2017). This approach reveals that in *Drosophila* brain

neurons A_{12} and $A_{11}G$ sites are significantly but moderately more common in Nab2-associated

transcripts compared to non-Nab2 associated transcripts. These A_{12} and $A_{11}G$ sites appear respectively

once every 1,553 and 687 bases on average among Nab2-associated transcripts and once every 1,901

and 935 bases on average among non-Nab2-assoicated transcripts, a 1.2- and 1.4-fold enrichment

(Figure 6B,C). Taken together, the findings that Nab2 associates with a specific subset of all RNAs with

poly(A) tails, and that these three A-rich motifs are not exclusive to Nab2-associated RNAs, strongly

argues that the polyadenosine sequence affinity of Nab2 alone is insufficient to explain Nab2-RNA

association specificity in *Drosophila* brain neurons. Other mechanisms must also contribute to Nab2

target choice, such as RNA secondary structure, protein-protein interactions, subnuclear localization,

and binding site competition. That said, the significant enrichment of a 41-bp A-rich motif, A_{12}, and

$A_{11}G$ observed in Nab2-associated RNAs suggests Nab2-RNA association is partially mediated through

these genetically encoded RNA sequence motifs as well as or instead of through the poly(A) tail.

Discussion

Mutation of either *ZC3H14 or ATXN2* gives rise to human disease, and the Nab2 and Atx2 RNA-

binding proteins encoded by their *Drosophila* orthologs are linked by a shared association with Fmr1

(Sudhakaran *et al.* 2014; Bienkowski *et al.* 2017). Here we show that *Nab2* and *Atx2* interact in multiple

contexts in *Drosophila*, specifically in fated eye cells, adult viability, and mushroom body neuronal

morphology. Notably, these interactions are dose-sensitive, as heterozygosity for *Atx2* loss-of-function

alleles is sufficient to suppress *Nab2* null phenotypes in adult viability and MB morphology. That is,

loss of Nab2 may sensitize these domains to reduced Atx2 activity, suggesting these RBPs regulate

some common processes. We find that these *Nab2-Atx2* interactions are likely not explained by

extended, simultaneous co-occupancy of Nab2 and Atx2 in common RNP complexes on shared RNA

transcripts. Each protein is concentrated in distinct subcellular compartments in adult mushroom body

neurons *in vivo*, and Nab2 and Atx2 weakly associate by co-IP from brain neurons. Thus, to explore an

alternative possibility—sequential regulation of shared RNA transcripts—we have carried out the first

high-throughput identification of Nab2- and Atx2-associated RNAs in *Drosophila*. We find these

proteins associate with overlapping sets of transcripts in *Drosophila* neurons, consistent with their

shared and distinct functions and supporting the model of sequential regulation. Identification of these

protein-transcript associations promises further insight into the functions shared between and unique to

each RBP. In addition, the identification of *Drosophila* Nab2-associated RNAs begins to address

longstanding questions about Nab2 function and the particular sensitivity of neurons to Nab2 loss,

revealing that Nab2 associates with a specific subset of polyadenylated RNAs *in vivo* despite the

theoretical potential to bind across all polyadenylated transcripts suggested by its high polyadenosine

affinity *in vitro* (Pak *et al.* 2011).

A model of opposing regulatory roles for Nab2 and Atx2

We show that Nab2 and Atx2 share associated RNAs in *Drosophila* neurons (Figures 4,5) and

that *Atx2* loss-of-function alleles suppress phenotypes of Nab2 loss (Figures 1,2). Taken together, these

findings imply that, at least for transcripts crucial for adult survival and MB α lobe morphology, Nab2

and Atx2 exert opposing regulatory roles on their shared associated RNAs. This opposing role

possibility aligns with some of the known functions of each protein. Namely, in *S. cerevisiae* Nab2

contributes to proper nuclear processing events including protection from enzymatic degradation,

poly(A) tail length control, splicing, and transcriptional termination while also facilitating poly(A) RNA

export from the nucleus (Green *et al.* 2002; Hector *et al.* 2002; Kelly *et al.* 2010; Schmid *et al.* 2015;

Soucek *et al.* 2016; Alpert *et al.* 2020). If *Drosophila* Nab2 also performs some or all of these nuclear

processing roles on its associated RNAs, then Nab2 binding should contribute to transcript stability,

nuclear export, and ultimately protein expression. Atx2, in contrast, is a key regulator of translational

efficiency in the cytoplasm, suppressing the translation of some target RNAs and activating the

translation of others (reviewed in Lee *et al.* 2018). As our data suggest Nab2 and Atx2 act in functional

opposition on a shared transcript set, we propose Atx2 primarily functions as a translational inhibitor

rather than activator on shared Nab2- and Atx2-associated RNAs. In this model (Figure 7), Nab2 and

Atx2 would act in temporal and spatial sequence to balance protein expression from their shared

associated RNAs in neurons, with Nab2 promoting proper nuclear RNA processing, stability, and export

and Atx2 inhibiting RNA translation, respectively.

This model of sequential temporal and spatial regulation aligns with evidence that Nab2 and

Atx2 primarily localize to different subcellular compartments in adult MBs at steady state and exhibit a

low level of co-precipitation from brain neurons (Figure 3). Potential explanations for the combination

of distinct localization profiles and limited physical association between Nab2 and Atx2 are found in

proposals that *S. cerevisiae* Nab2 shuttles out of the nucleus with bound RNAs during export before

releasing them and returning to the nucleus (Aitchison *et al.* 1996; Lee and Aitchison 1999; Duncan *et*

al. 2000). Thus, Nab2 and Atx2 may physically share associated RNAs briefly if neuronal *Drosophila*

Nab2 similarly shuttles and both RBPs are present during the nuclear-cytoplasmic handoff of mRNP

remodeling that follows mRNA export from the nucleus (reviewed in Müller-McNicoll and Neugebauer

2013; Chen and Shyu 2014). Functional and physical links between Nab2 and an RBP with a prominent

cytoplasmic localization pattern like Atx2 have been observed previously, specifically with Fmr1

(Bienkowski *et al.* 2017). However, the physical associations observed between Fmr1 and Nab2 are

more robust than that observed between Atx2 and Nab2 in the present study (Figure 3B)—this

distinction may be partially explained by the different localization patterns of Atx2 and Fmr1. Atx2 is

exclusively cytoplasmic in neurons except under certain pathogenic conditions (Huynh *et al.* 2003;

Lessing and Bonini 2008; Elden *et al.* 2010), while Fmr1 shuttles between the two compartments,

associating with at least some of its target RNAs in the nucleus (Tamanini *et al.* 1999; Kim *et al.* 2009).

Thus, Nab2 and Fmr1 may theoretically co-occupy and coregulate shared transcripts in both cellular

compartments while Nab2 and Atx2 sequentially regulate shared transcripts exchanged during a nuclear-

cytoplasmic handoff, representing two distinct modes of functional interaction between Nab2 and a

cytoplasmic RBP.

This model provides a firm foundation and raises many readily testable hypotheses to be

explored in future research. The model predicts that for shared Nab2- and Atx2-associated RNAs, loss

of Nab2 decreases transcript stability, impedes proper nuclear processing events including poly(A) tail

length control, and impairs poly(A) RNA export from the nucleus, ultimately leading to decreases in

protein product. Conversely, we predict partial loss of Atx2 releases translational inhibition on these

shared transcripts and induces increases in protein product. Finally, loss of both proteins would balance

these effects, resulting in steady-state levels of protein product more similar to the wild-type condition.

With the identity of Nab2- and Atx2-associated RNAs in hand, future research is enabled to test these

predictions.

Prominent Nab2- and Atx2-associated transcripts provide links to brain development and

function

Of all the RBP-associated transcripts identified here, we defined the prominent shared and RBP-specific

associated transcripts as those annotated within overrepresented GO terms (Figure 4F, Supplemental

Figure 4) and/or passing a 1.5-fold enrichment threshold. The identities and functional roles of these

prominent RBP-associated transcripts (examples in Figure 5) provide potential mechanistic explanations

for the biological roles of each RBP. For example, the effects of Nab2 and Atx2 on MB morphology

may be mediated in part through regulation of shared mRNAs *sm* and *stai*, which respectively encode an

hnRNP and a tubulin binding protein both linked to axonal morphology and development (Layalle *et al.*

2005; Lachkar *et al.* 2010; Zwarts *et al.* 2015). The effects of Nab2 and Atx2 on memory (Sudhakaran *et*

al. 2014; Kelly *et al.* 2016) may be due in part to regulation of shared transcripts *drk*, *shi*, *Gαo*, and

me31B, all of which encode proteins with roles in memory formation or retrieval (Dubnau *et al.* 2001;

Ferris *et al.* 2006; Moressis *et al.* 2009; Sudhakaran *et al.* 2014). Both Nab2 and Atx2 may be involved

in RNAi at multiple levels, regulating *me31B* RNA in neurons in addition to associating, in the case of

Atx2, with me31B protein (Lee *et al.* 2017; Bakthavachalu *et al.* 2018). Finally, the suppression of

GMR>Nab2 by *Atx2* alleles in the eye may be explained in part by the shared association of Nab2 and

Atx2 with *HmgZ* (*HMG protein Z*) RNA, which encodes a chromatin remodeler linked to survival of R7

retinal photoreceptor neurons (Kanuka *et al.* 2005; Ragab *et al.* 2006).

Among the associated RNAs specific to each RBP, we found only Nab2 associated with *fwe*

(*flower*), *Arpc2*, *side-II*, and *SLC22A* RNA, connections which may further explain the role of Nab2 in

guiding MB morphology and regulating learning and memory. These transcripts respectively encode a

transmembrane mediator of neuronal culling in development (Merino *et al.* 2013), a component of the

neuronal growth cone advance-regulating Arp2/3 complex (Hudson and Cooley 2002; Yang *et al.* 2012),

an immunoglobulin superfamily member potentially contributing to axon guidance and synapse

formation in the optic lobe (Tan *et al.* 2015), and a transmembrane acetylcholine transporter localized to

MB dendrites and involved in suppressing memory formation (Gai *et al.* 2016). On the other hand, the

association of Atx2 with Atx2-specific RNAs *Xpc* and *tea*, which respectively encode players in the

fundamental cellular processes of DNA repair and telomere protection (Henning *et al.* 1994; Goosen

2010; Zhang *et al.* 2016), may partially explain why Atx2 genomic loss, unlike Nab2 genomic loss, is

larval lethal (Satterfield *et al.* 2002). In summary, defining the potential functional impact of Nab2- and

Atx2-RNA associations like these will provide critical insight into the roles of Nab2 and Atx2 in

neurodevelopment and *Drosophila* disease models.

Nab2 associates with a more specific set of RNAs in metazoans than in *S. cerevisiae*

The degree of RNA association specificity metazoan Nab2/ZC3H14 exhibits has been a longstanding

question, in part because competing answers are suggested by the functional similarities and differences

between metazoan Nab2/ZC3H14 and the *S. cerevisiae* Nab2 ortholog. In *S. cerevisiae*, Nab2 is

essential for viability (Anderson *et al.* 1993) and is a central player in post-transcriptional regulation of

many transcripts, serving as a nuclear poly(A)-binding-protein regulating transcript stability (Schmid *et*

al. 2015), poly(A) tail length, and poly(A) RNA export from the nucleus among other processes

(reviewed in Moore 2005; Chen and Shyu 2014; and Stewart 2019). However, in metazoans Nab2 or the

full-length form of ZC3H14 is dispensable for cellular viability, and the effects of either protein on

poly(A) tail length and poly(A) RNA export from the nucleus are either less pronounced and likely

exerted on fewer transcripts than in *S. cerevisiae* or are not detected (Farny *et al.* 2008; Kelly *et al.*

2010; Wigington *et al.* 2016; Bienkowski *et al.* 2017; Rha *et al.* 2017b; Morris and Corbett 2018).

Consistently, Nab2/ZC3H14 have not been found to associate with all polyadenylated RNAs tested in

metazoans so far (Wigington *et al.* 2016; Bienkowski *et al.* 2017; Morris and Corbett 2018), but the

possibility has remained that these few identified non-Nab2/ZC3H14-associated transcripts are outliers

and metazoan Nab2/ZC3H14 associates with a large majority of polyadenylated RNAs similarly to *S.*

cerevisiae Nab2 (Tuck and Tollervey 2013), likely in part by binding poly(A) tails. Indeed, the identities

of Nab2- or ZC3H14-associated RNAs in metazoans had never previously been addressed with a

comprehensive, high-throughput method.

Our results identify a specific set of transcripts that neuronal Nab2 associates with in *Drosophila*.

Of the 5,760 transcripts tested in the RIP-Seq, only about 2.5% were found to associate with Nab2 in

Drosophila neurons (Figure 4), a much smaller percentage of the transcriptome than associates with

Nab2 in *S. cerevisiae* (Guisbert *et al.* 2005; Batisse *et al.* 2009; Tuck and Tollervey 2013). Importantly,

this likely represents an undercount of all Nab2-associated transcripts in neurons *in vivo*—some RNAs

associated with Nab2 in prior studies are absent from our Nab2-associated transcript set (Bienkowski *et*

al. 2017; Jalloh *et al.* 2020), and technical limitations impacted our sequencing read depth (see

Methods). Higher sensitivity approaches (e.g. CLIP-Seq) could reveal a broader set of Nab2-associated

transcripts in *Drosophila* than we define here. Nonetheless, in the present study the majority of both

RNAs (Figure 4) and tested polyadenosine-rich sequence motifs (Figure 6) were not found to be

associated with Nab2, strongly supporting a model in which Nab2 associates with a specific subset of

RNAs in *Drosophila* neurons. Perhaps for this more select group of transcripts Nab2 still plays a key

role in transcript stability, poly(A) tail length control, transcription termination, and poly(A) RNA

export from the nucleus, such that defects will only be observed in targeted examinations of single

transcripts and not in bulk assays—one does not always reflect the other (Kelly *et al.* 2014; Bienkowski

et al. 2017). This model of Nab2 specificity in *Drosophila* aligns well with the knowledge that

Nab2/full-length ZC3H14 is essential for cellular viability in *S. cerevisiae* (Anderson *et al.* 1993) but not

in *Drosophila* (Bienkowski *et al.* 2017), mice (Rha *et al.* 2017b), or, seemingly, humans (Pak *et al.*

2011; Al-Nabhani *et al.* 2018). This diminished global requirement for Nab2/ZC3H14 in metazoans may

be due, at least in part, to functional overlap with PABPN1, an evolutionarily distinct nuclear

polyadenosine RNA-binding protein that is absent in *S. cerevisiae* (Mangus *et al.* 2003) but controls

poly(A) tail length and is essential in *Drosophila* (Benoit *et al.* 2005), mice (Vest *et al.* 2017), and

humans (Hart *et al.* 2015).

The model of Nab2 specificity in *Drosophila* does not conflict with its affinity for

polyadenosine, which could theoretically allow Nab2 to bind all transcripts with a poly(A) tail. Even in

S. cerevisiae, the broad binding profile of Nab2 (Batisse *et al.* 2009) and central role in poly(A) tail

length control (Kelly *et al.* 2010), poly(A) RNA export from the nucleus (Green *et al.* 2002), and

protection of poly(A) RNA from degradation (Schmid *et al.* 2015) does not translate to binding the

poly(A) tails of all transcripts (Guisbert *et al.* 2005; Tuck and Tollervey 2013). More broadly, linear

sequence motifs alone are insufficient to explain RBP specificity—RBPs do not generally occupy all of

their available binding motifs throughout the transcriptome (Li *et al.* 2010; Taliaferro *et al.* 2016).

Moreover, non-paralog RBPs with substantially overlapping or identical linear target motifs still bind

distinct RNA target sets, demonstrating that linear motifs are only one of a set of RNA features that

direct RBP-RNA associations (Dominguez *et al.* 2018). Based on the present study, these general

features of RBPs hold for Nab2 as well. MEME and FIMO motif analyses reveal a long A-rich motif

and the canonical Nab2 binding motifs A_{12} and $A_{11}G$ are enriched in but not exclusive to Nab2-

associated RNAs (Figure 6). Given the behavior of other RBPs, it is consistent that *Drosophila* Nab2

exhibits this binding specificity and, given our RIP-Seq data and previous studies, likely binds some but

almost certainly does not bind not all poly(A) tails in *Drosophila* despite its high affinity for

polyadenosine RNA *in vitro* (Pak *et al.* 2011).

Taken together, these data align with the model that in metazoans Nab2/ZC3H14 is more

specific in its transcript associations than in *S. cerevisiae*. With this model in mind and the Nab2-

associated transcripts identified in this study in hand, future research will be enabled to focus on how

Nab2 functions on these particular transcripts in *Drosophila*, and why this function is so crucial for adult

viability, neuronal morphology, locomotion, and learning and memory. Given that a polyadenosine-rich

motif along with A_{12} and $A_{11}G$ motifs are correlated with but are not sufficient for Nab2-RNA

association, future research must also focus on what additional features of transcripts or their associated

proteins promote or prevent Nab2 association.

Conclusion

In sum, the data we present here identify functional interactions between Nab2 and Atx2 in *Drosophila*

brain morphology and adult viability and define a set of RNA transcripts associated with each protein in

brain neurons. Crucially, theses RNA sets overlap—some associated transcripts are shared between

Nab2 and Atx2 and some are specific to each RBP. Identifying these RBP-associated transcripts

provides potential mechanistic links between the roles in neuronal development and function their

encoded proteins perform, Nab2, and Atx2. This foundation will be especially important for Nab2, as

the exact molecular function of metazoan Nab2/ZC3H14 on the vast majority of its associated RNAs in

any cell type remains largely unknown. The identity of many *Drosophila* Nab2-associated transcripts,

now revealed, will be required to define Nab2/ZC3H14 function in metazoans and enable our

understanding of why loss of this largely nuclear polyadenosine RNA-binding protein results in

neurological or neurodevelopmental deficits in flies and mice and in intellectual disability in humans.

Acknowledgements

The authors would like to thank current and past members of the Moberg and Corbett lab groups, especially Drs. Ayan Banerjee, Rick Bienkowski, Daniel Barron, Binta Jalloh, Stephanie Jones, Annie McPherson-Davie, Milo Fasken, and Sara Leung for their support of, instruction in, and enlightening discussions of this work. We would also like to thank Drs. Bing Yao, Jingjing Yang, Michael Christopher, and Carlos Moreno for initial bioinformatics advice and the Georgia Genomics and Bioinformatics Core (GGBC) at the University of Georgia, especially Tyler James Simmonds and Dr. Magdy S. Alabady, for essential library preparation, sequencing, and assistance in sequencing experiment design and preparation.

We would like to thank Drs. Nancy Bonini and Michael Parisi for the gift of a *Atx2^{X1}* stock; Dr. Ravi Allada, Khadijah Hamid, and Dr. Satya Surabhi for the gift of a *UAS>Atx2-3xFLAG* stock; Dr. Chunghun Lim and for the gift of rabbit α-Atx2; Dr. Corey S. Goodman for the contribution of 1D4 Anti-Fas2 to the DSHB; Drs. Gary Bassell, Roger Deal, Steven Warren, James Q. Zheng, and their respective labs for assistance and use of their equipment; Laura Fox-Goharioon for confocal microscope training; Dr. Michael I. Love for extensive public online instruction in the methodology and use of *DESeq2*; Dr. Mauricio Rodriguez-Lanetty for a public TRIzol-column hybrid RNA extraction protocol; and Eileen Chow for public video instruction in bulk *Drosophila* head isolation.

The authors would also like to thank with particular enthusiasm the authors, contributors, and ongoing maintainers of the incredible public resources supporting this work and without which it would not have been possible, including Flybase (NIH U41HG000739, UK MRC MR/N030117/1), the Galaxy Project (NIH 2U41HG006620), the R Project, the Developmental Studies Hybridoma Bank (University of Iowa, NIH), and the Bloomington Drosophila Stock Center (NIH P40OD018537). This research was funded by grants from the National Institutes of Health, specifically from the National Institute of Child Health and Human Development (F31 HD088043) to J.C.R. and from the National Institute of Mental Health (R01 MH107305) to A.H.C. and K.H.M. The authors declare no conflicts of interest.

Shared Nab2- and Atx2-associated transcripts			
AGO2	*drk*	*me31B*	*shi*
apolpp	*Gαo*	*Msp300*	*sm*
CG31221	*Gat*	*mtd*	*snoRNA:Or-aca5*
CG42540	*Gγ30A*	*Rbp6*	*snoRNA:Or-CD2*
CG4360	*Gp150*	*RpL37A*	*snoRNA:Ψ18S-1854b*
CG6675	*HmgZ*	*RpS27A*	*Stai*
CG9813	*l(3)80Fg*[a]	*RpS29*	*Ulp1*

TABLE 1. Identities of the 28 transcripts overlapping between the Nab2 and Atx2 RNA interactomes. For all 5,760 genes in the RIP-Seq testable set, control-normalized IP/Input enrichment values were calculated followed by gene-by-gene one-way ANOVAs, Dunnett's post-hoc tests, and within-gene multiple hypothesis testing adjustment (*Dun. Adj. p*). All transcripts statistically significantly (*Dun. Adj. p* < 0.05) enriched in both Nab2- and Atx2-associated transcripts sets are listed here. Functional interactions between Nab2 and Atx2 in brain neurons may be explained by their coordinate regulation of these shared associated transcripts.

[a]Symbol updated from *CG40178* to current nomenclature in BDGP6.37.

Literature Cited

Afgan E., D. Baker, B. Batut, M. van den Beek, D. Bouvier, *et al.*, 2018 The Galaxy platform for accessible, reproducible and collaborative biomedical analyses: 2018 update. Nucleic Acids Res. 46: 537–544. https://doi.org/10.1093/nar/gky379

Agha Z., Z. Iqbal, M. Azam, H. Ayub, L. E. L. M. Vissers, *et al.*, 2014 Exome Sequencing Identifies Three Novel Candidate Genes Implicated in Intellectual Disability, (O. R. Bandapalli, Ed.). PLoS One 9: e112687. https://doi.org/10.1371/journal.pone.0112687

Aguilo F., F. Zhang, A. Sancho, M. Fidalgo, S. Di Cecilia, *et al.*, 2015 Coordination of m6A mRNA Methylation and Gene Transcription by ZFP217 Regulates Pluripotency and Reprogramming. Cell Stem Cell 17: 689–704. https://doi.org/10.1016/j.stem.2015.09.005

Aibara S., J. M. B. Gordon, A. S. Riesterer, S. H. McLaughlin, and M. Stewart, 2017 Structural basis for the dimerization of Nab2 generated by RNA binding provides insight into its contribution to both poly(A) tail length determination and transcript compaction in Saccharomyces cerevisiae. Nucleic Acids Res. 45: 1529–1538. https://doi.org/10.1093/nar/gkw1224

Aitchison J. D., G. Blobel, and M. P. Rout, 1996 Kap104p: A karyopherin involved in the nuclear transport of messenger RNA binding proteins. Science (80-.). 274: 624–627. https://doi.org/10.1126/science.274.5287.624

Al-Nabhani M., S. Al-Rashdi, F. Al-Murshedi, A. Al-Kindi, K. Al-Thihli, *et al.*, 2018 Reanalysis of exome sequencing data of intellectual disability samples: Yields and benefits. Clin. Genet. https://doi.org/10.1111/cge.13438

Almudi I., M. Corominas, and F. Serras, 2010 Competition between SOCS36E and Drk modulates sevenless receptor tyrosine kinase activity. J. Cell Sci. 123: 3857–3862. https://doi.org/10.1242/jcs.071134

Alpert T., K. Straube, F. Carrillo Oesterreich, and K. M. Neugebauer, 2020 Widespread Transcriptional Readthrough Caused by Nab2 Depletion Leads to Chimeric Transcripts with Retained Introns. Cell Rep. 33: 108324. https://doi.org/10.1016/j.celrep.2020.108324

Anderson J. T., S. M. Wilson, K. V Datar, and M. S. Swanson, 1993 NAB2: a yeast nuclear polyadenylated RNA-binding protein essential for cell viability. Mol. Cell. Biol. 13: 2730–2741. https://doi.org/10.1128/mcb.13.5.2730

Ashburner M., C. A. Ball, J. A. Blake, D. Botstein, H. Butler, *et al.*, 2000 Gene ontology: Tool for the unification of biology. Nat. Genet. 25: 25–29.

Ashley C. T., K. D. Wilkinson, D. Reines, and S. T. Warren, 1993 FMR1 protein: Conserved RNP family domains and selective RNA binding. Science (80-.). 262: 563–566. https://doi.org/10.1126/science.7692601

Baejen C., P. Torkler, S. Gressel, K. Essig, J. Söding, *et al.*, 2014 Transcriptome Maps of mRNP Biogenesis Factors Define Pre-mRNA Recognition. Mol. Cell 55: 745–757. https://doi.org/10.1016/j.molcel.2014.08.005

Bailey T. L., and C. Elkan, 1994 Fitting a mixture model by expectation maximization to discover motifs in biopolymers. Proceedings. Int. Conf. Intell. Syst. Mol. Biol. 2: 28–36.

Bailey T. L., M. Boden, F. A. Buske, M. Frith, C. E. Grant, *et al.*, 2009 MEME Suite: Tools for motif discovery and searching. Nucleic Acids Res. 37: W202–W208. https://doi.org/10.1093/nar/gkp335

Bakthavachalu B., J. Huelsmeier, I. P. Sudhakaran, J. Hillebrand, A. Singh, *et al.*, 2018 RNP-Granule Assembly via Ataxin-2 Disordered Domains Is Required for Long-Term Memory and Neurodegeneration Article RNP-Granule Assembly via Ataxin-2 Disordered Domains Is Required for Long-Term Memory and Neurodegeneration. Neuron 98: 754-766.e4. https://doi.org/10.1016/j.neuron.2018.04.032

Banerjee A., K. E. Vest, G. K. Pavlath, and A. H. Corbett, 2017 Nuclear poly(A) binding protein 1 (PABPN1) and Matrin3 interact in muscle cells and regulate RNA processing. Nucleic Acids Res. 45: 10706–10725. https://doi.org/10.1093/nar/gkx786

Barbee S. A., P. S. Estes, A. M. Cziko, J. Hillebrand, R. A. Luedeman, *et al.*, 2006 Staufen- and FMRP-Containing Neuronal RNPs Are Structurally and Functionally Related to Somatic P Bodies. Neuron 52: 997–1009. https://doi.org/10.1016/j.neuron.2006.10.028

Bardoni B., S. Abekhoukh, S. Zongaro, and M. Melko, 2012 Intellectual disabilities, neuronal posttranscriptional RNA metabolism, and RNA-binding proteins: Three actors for a complex scenario, pp. 29–51 in *Progress in Brain Research*, Elsevier B.V.

Batisse J., C. Batisse, A. Budd, B. Bö ttcher, and E. Hurt, 2009 Purification of Nuclear Poly(A)-binding Protein Nab2 Reveals Association with the Yeast Transcriptome and a Messenger Ribonucleoprotein Core Structure. J. Biol. Chem. 284: 34911–34917. https://doi.org/10.1074/jbc.M109.062034

Bellen H. J., R. W. Levis, G. Liao, Y. He, J. W. Carlson, *et al.*, 2004 The BDGP gene disruption project: Single transposon insertions associated with 40% of Drosophila genes. Genetics 167: 761–781. https://doi.org/10.1534/genetics.104.026427

Benoit B., G. Mitou, A. Chartier, C. Temme, S. Zaessinger, *et al.*, 2005 An essential cytoplasmic function for the nuclear poly(A) binding protein, PABP2, in poly(A) tail length control and early development in Drosophila. Dev. Cell 9: 511–522. https://doi.org/10.1016/j.devcel.2005.09.002

Bienkowski R. S., A. Banerjee, J. C. Rounds, J. Rha, O. F. Omotade, *et al.*, 2017 The Conserved, Disease-Associated RNA Binding Protein dNab2 Interacts with the Fragile X Protein Ortholog in Drosophila Neurons. Cell Rep. 20: 1372–1384. https://doi.org/10.1016/j.celrep.2017.07.038

Bokhoven H. van, 2011 Genetic and epigenetic networks in intellectual disabilities. Annu. Rev. Genet. 45: 81–104. https://doi.org/10.1146/annurev-genet-110410-132512

Chelly J., M. Khelfaoui, F. Francis, B. Chérif, and T. Bienvenu, 2006 Genetics and pathophysiology of mental retardation. Eur. J. Hum. Genet. 14: 701–713.

Chen C. Y. A., and A. Bin Shyu, 2014 Emerging mechanisms of mRNP remodeling regulation. Wiley Interdiscip. Rev. RNA 5: 713–722. https://doi.org/10.1002/wrna.1241

Connolly J. B., I. J. H. Roberts, J. D. Armstrong, K. Kaiser, M. Forte, *et al.*, 1996 Associative learning disrupted by impaired Gs signaling in Drosophila mushroom bodies. Science (80-.). 274: 2104–2107. https://doi.org/10.1126/science.274.5295.2104

Corgiat E. B., S. M. List, J. C. Rounds, A. H. Corbett, and K. H. Moberg, 2020 The RNA binding

protein Nab2 regulates the proteome of the developing Drosophila brain. bioRxiv 2020.12.10.419846. https://doi.org/10.1101/2020.12.10.419846

Crooks G. E., G. Hon, J. M. Chandonia, and S. E. Brenner, 2004 WebLogo: A sequence logo generator. Genome Res. 14: 1188–1190. https://doi.org/10.1101/gr.849004

Dobin A., C. A. Davis, F. Schlesinger, J. Drenkow, C. Zaleski, *et al.*, 2013 STAR: ultrafast universal RNA-seq aligner. Bioinformatics 29: 15–21. https://doi.org/10.1093/bioinformatics/bts635

Domanski M., K. Molloy, H. Jiang, B. T. Chait, M. P. Rout, *et al.*, 2012 Improved methodology for the affinity isolation of human protein complexes expressed at near endogenous levels. Biotechniques 0: 1–6. https://doi.org/10.2144/000113864

Dominguez D., P. Freese, M. S. Alexis, A. Su, M. Hochman, *et al.*, 2018 Sequence, Structure, and Context Preferences of Human RNA Binding Proteins. Mol. Cell 70: 854-867.e9. https://doi.org/10.1016/j.molcel.2018.05.001

Dubnau J., L. Grady, T. Kitamoto, and T. Tully, 2001 Disruption of neurotransmission in Drosophila mushroom body blocks retrieval but not acquisition of memory. Nature 411: 476–480. https://doi.org/10.1038/35078077

Duncan K., J. G. Umen, and C. Guthrie, 2000 A putative ubiquitin ligase required for efficient mRNA export differentially affects hnRNP transport. Curr. Biol. 10: 687–696. https://doi.org/10.1016/S0960-9822(00)00527-3

Dunnett C. W., 1955 A Multiple Comparison Procedure for Comparing Several Treatments with a Control. J. Am. Stat. Assoc. 50: 1096–1121. https://doi.org/10.1080/01621459.1955.10501294

Elden A. C., H.-J. Kim, M. P. Hart, A. S. Chen-Plotkin, B. S. Johnson, *et al.*, 2010 Ataxin-2 intermediate-length polyglutamine expansions are associated with increased risk for ALS. Nature 466: 1069–1075. https://doi.org/10.1038/nature09320

Ellis M. C., E. M. O'Neill, and G. M. Rubin, 1993 Expression of Drosophila glass protein and evidence for negative regulation of its activity in non-neuronal cells by another DNA-binding protein. Development 119: 855–865.

Eulalio A., I. Behm-Ansmant, D. Schweizer, and E. Izaurralde, 2007 P-Body Formation Is a Consequence, Not the Cause, of RNA-Mediated Gene Silencing. Mol. Cell. Biol. 27: 3970–3981. https://doi.org/10.1128/mcb.00128-07

Farny N. G., J. A. Hurt, and P. A. Silver, 2008 Definition of global and transcript-specific mRNA export pathways in metazoans. Genes Dev. 22: 66–78. https://doi.org/10.1101/gad.1616008

Fasken M. B., A. H. Corbett, and M. Stewart, 2019 Structure-function relationships in the Nab2 polyadenosine-RNA binding Zn finger protein family. Protein Sci. 28: 513–523. https://doi.org/10.1002/pro.3565

Ferris J., H. Ge, L. Liu, and G. Roman, 2006 G(o) signaling is required for Drosophila associative learning. Nat. Neurosci. 9: 1036–1040. https://doi.org/10.1038/nn1738

Freeman M., 1996 Reiterative use of the EGF receptor triggers differentiation of all cell types in the Drosophila eye. Cell 87: 651–660. https://doi.org/10.1016/S0092-8674(00)81385-9

Gai Y., Z. Liu, I. Cervantes-Sandoval, and R. L. Davis, 2016 Drosophila SLC22A Transporter Is a Memory Suppressor Gene that Influences Cholinergic Neurotransmission to the Mushroom Bodies. Neuron 90: 581–595. https://doi.org/10.1016/j.neuron.2016.03.017

González-Aguilera C., C. Tous, R. Babiano, J. de la Cruz, R. Luna, *et al.*, 2011 Nab2 functions in the metabolism of RNA driven by polymerases II and III. Mol. Biol. Cell 22: 2729–40. https://doi.org/10.1091/mbc.E11-01-0055

Goosen N., 2010 Scanning the DNA for damage by the nucleotide excision repair machinery. DNA Repair (Amst). 9: 593–596. https://doi.org/10.1016/j.dnarep.2010.02.015

Grant C. E., T. L. Bailey, and W. S. Noble, 2011 FIMO: Scanning for occurrences of a given motif. Bioinformatics 27: 1017–1018. https://doi.org/10.1093/bioinformatics/btr064

Green D. M., K. A. Marfatia, E. B. Crafton, X. Zhang, X. Cheng, *et al.*, 2002 Nab2p is required for poly(A) RNA export in Saccharomyces cerevisiae and is regulated by arginine methylation via Hmt1p. J. Biol. Chem. 277: 7752–7760. https://doi.org/10.1074/jbc.M110053200

Guisbert K. K., K. Duncan, H. Li, and C. Guthrie, 2005 Functional specificity of shuttling hnRNPs revealed by genome-wide analysis of their RNA binding profiles. RNA 11: 383–393. https://doi.org/10.1261/rna.7234205

Gwinn-Hardy K., J. Y. Chen, H. C. Liu, T. Y. Liu, M. Boss, *et al.*, 2000 Spinocerebellar ataxia type 2 with parkinsonism in ethnic Chinese. Neurology 55: 800–805. https://doi.org/10.1212/WNL.55.6.800

Hao L. Y., B. I. Giasson, and N. M. Bonini, 2010 DJ-1 is critical for mitochondrial function and rescues PINK1 loss of function. Proc. Natl. Acad. Sci. U. S. A. 107: 9747–9752. https://doi.org/10.1073/pnas.0911175107

Hart T., M. Chandrashekhar, M. Aregger, Z. Steinhart, K. R. Brown, *et al.*, 2015 High-Resolution CRISPR Screens Reveal Fitness Genes and Genotype-Specific Cancer Liabilities. Cell 163: 1515–1526. https://doi.org/10.1016/j.cell.2015.11.015

Hay B. A., T. Wolff, and G. M. Rubin, 1994 Expression of baculovirus P35 prevents cell death in Drosophila. Development 120: 2121–2129.

Hector R. E., K. R. Nykamp, S. Dheur, J. T. Anderson, P. J. Non, *et al.*, 2002 Dual requirement for yeast hnRNP Nab2p in mRNA poly(A) tail length control and nuclear export. EMBO J. 21: 1800–10. https://doi.org/10.1093/emboj/21.7.1800

Heisenberg M., 2003 Mushroom body memoir: From maps to models. Nat. Rev. Neurosci. 4: 266–275. https://doi.org/10.1038/nrn1074

Henning K. A., C. Peterson, R. Legerski, and E. C. Friedberg, 1994 Cloning the Drosophila homolog of the xeroderma pigmentosum complementation group C gene reveals homology between the predicted human and drosophila polypeptides and that encoded by the yeast RAD4 gene. Nucleic Acids Res. 22: 257–261. https://doi.org/10.1093/nar/22.3.257

Hillebrand J., K. Pan, A. Kokaram, S. Barbee, R. Parker, *et al.*, 2010 The Me31B DEAD-Box Helicase Localizes to Postsynaptic Foci and Regulates Expression of a CaMKII Reporter mRNA in Dendrites of Drosophila Olfactory Projection Neurons. Front. Neural Circuits 4: 121.

https://doi.org/10.3389/fncir.2010.00121

Hoskins R. A., J. W. Carlson, K. H. Wan, S. Park, I. Mendez, *et al.*, 2015 The Release 6 reference sequence of the Drosophila melanogaster genome. Genome Res. 25: 445–458. https://doi.org/10.1101/gr.185579.114

Hudson A. M., and L. Cooley, 2002 A subset of dynamic actin rearrangements in Drosophila requires the Arp2/3 complex. J. Cell Biol. 156: 677–687. https://doi.org/10.1083/jcb.200109065

Huet F., J. T. Lu, K. V. Myrick, L. R. Baugh, M. A. Crosby, *et al.*, 2002 A deletion-generator compound element allows deletion saturation analysis for genomewide phenotypic annotation. Proc. Natl. Acad. Sci. U. S. A. 99: 9948–9953. https://doi.org/10.1073/pnas.142310099

Huynh D. P., H.-T. Yang, H. Vakharia, D. Nguyen, and S. M. Pulst, 2003 Expansion of the polyQ repeat in ataxin-2 alters its Golgi localization, disrupts the Golgi complex and causes cell death. Hum. Mol. Genet. 12: 1485–1496. https://doi.org/10.1093/hmg/ddg175

Imbert G., F. Saudou, G. Yvert, D. Devys, Y. Trottier, *et al.*, 1996 Cloning of the gene for spinooerebellar ataxia 2 reveals a locus with high sensitivity to expanded CAG/glutamine repeats. Nat. Genet. 14: 285–291. https://doi.org/10.1038/ng1196-285

Inlow J. K., and L. L. Restifo, 2004 Molecular and Comparative Genetics of Mental Retardation. Genetics 166: 835–881. https://doi.org/10.1534/genetics.166.2.835

Jalloh B., J. C. Rounds, B. E. Brown, I. J. Kremsky, A. Banerjee, *et al.*, 2020 The Nab2 RNA binding protein promotes sex-specific splicing of Sex lethal in Drosophila neuronal tissue. bioRxiv 2020.11.13.382168. https://doi.org/10.1101/2020.11.13.382168

Jean S., S. Cox, E. J. Schmidt, F. L. Robinson, and A. Kiger, 2012 Sbf/MTMR13 coordinates PI(3)P and Rab21 regulation in endocytic control of cellular remodeling. Mol. Biol. Cell 23: 2723–2740. https://doi.org/10.1091/mbc.E12-05-0375

Jiménez-López D., and P. Guzmán, 2014 Insights into the evolution and domain structure of ataxin-2 proteins across eukaryotes. BMC Res. Notes 7: 453. https://doi.org/10.1186/1756-0500-7-453

Kahsai L., and T. Zars, 2011 Learning and memory in drosophila: Behavior, genetics, and neural systems, pp. 139–167 in *International Review of Neurobiology*, Academic Press Inc.

Kanuka H., T. Hiratou, T. Igaki, H. Kanda, E. Kuranaga, *et al.*, 2005 Gain-of-function screen identifies a role of the Sec61α translocon in Drosophila postmitotic neurotoxicity. Biochim. Biophys. Acta - Gen. Subj. 1726: 225–237. https://doi.org/10.1016/j.bbagen.2005.06.020

Kelly S. M., S. A. Pabit, C. M. Kitchen, P. Guo, K. A. Marfatia, *et al.*, 2007 Recognition of polyadenosine RNA by zinc finger proteins. Proc. Natl. Acad. Sci. U. S. A. 104: 12306–12311. https://doi.org/10.1073/pnas.0701244104

Kelly S. M., S. W. Leung, L. H. Apponi, A. M. Bramley, E. J. Tran, *et al.*, 2010 Recognition of polyadenosine RNA by the zinc finger domain of nuclear poly(A) RNA-binding protein 2 (Nab2) is required for correct mRNA 3'-end formation. J. Biol. Chem. 285: 26022–32. https://doi.org/10.1074/jbc.M110.141127

Kelly S. M., S. W. Leung, C. Pak, A. Banerjee, K. H. Moberg, *et al.*, 2014 A conserved role for the zinc

finger polyadenosine RNA binding protein, ZC3H14, in control of poly(A) tail length. RNA 20: 681–8. https://doi.org/10.1261/rna.043984.113

Kelly S. M., R. Bienkowski, A. Banerjee, D. J. Melicharek, Z. A. Brewer, *et al.*, 2016 The Drosophila ortholog of the Zc3h14 RNA binding protein acts within neurons to pattern axon projection in the developing brain. Dev. Neurobiol. 76: 93–106. https://doi.org/10.1002/dneu.22301

Kelly S. M., A. Elchert, and M. Kahl, 2017 Dissection and immunofluorescent staining of mushroom body and photoreceptor neurons in adult Drosophila melanogaster brains. J. Vis. Exp. 2017: e56174. https://doi.org/10.3791/56174

Kim M., M. Bellini, and S. Ceman, 2009 Fragile X Mental Retardation Protein FMRP Binds mRNAs in the Nucleus. Mol. Cell. Biol. 29: 214–228. https://doi.org/10.1128/mcb.01377-08

Lachkar S., M. Lebois, M. O. Steinmetz, A. Guichet, N. Lal, *et al.*, 2010 Drosophila stathmins bind tubulin heterodimers with high and variable stoichiometries. J. Biol. Chem. 285: 11667–11680. https://doi.org/10.1074/jbc.M109.096727

Laemmli U. K., 1970 Cleavage of structural proteins during the assembly of the head of bacteriophage T4. Nature 227: 680–685. https://doi.org/10.1038/227680a0

Layalle S., E. Coessens, A. Ghysen, and C. Dambly-Chaudière, 2005 Smooth, a hnRNP encoding gene, controls axonal navigation in Drosophila. Genes to Cells 10: 119–125. https://doi.org/10.1111/j.1365-2443.2005.00822.x

Lee D. C. Y., and J. D. Aitchison, 1999 Kap104p-mediated nuclear import. Nuclear localization signals in mRNA- binding proteins and the role of Ran and RNA. J. Biol. Chem. 274: 29031–29037. https://doi.org/10.1074/jbc.274.41.29031

Lee J., E. Yoo, H. Lee, K. Park, J.-H. Hur, *et al.*, 2017 LSM12 and ME31B/DDX6 Define Distinct Modes of Posttranscriptional Regulation by ATAXIN-2 Protein Complex in Drosophila Circadian Pacemaker Neurons. Mol. Cell 66: 129-140.e7. https://doi.org/10.1016/j.molcel.2017.03.004

Lee J., M. Kim, T. Q. Itoh, and C. Lim, 2018 Ataxin-2: A versatile posttranscriptional regulator and its implication in neural function. Wiley Interdiscip. Rev. RNA 9: e1488. https://doi.org/10.1002/wrna.1488

Lee W.-H., E. B. Corgiat, J. C. Rounds, Z. Shepherd, A. H. Corbett, *et al.*, 2020 A Genetic Screen Links the Disease-Associated Nab2 RNA-Binding Protein to the Planar Cell Polarity Pathway in Drosophila melanogaster. G3 GENES, GENOMES, Genet. https://doi.org/10.1534/g3.120.401637

Lessing D., and N. M. Bonini, 2008 Polyglutamine Genes Interact to Modulate the Severity and Progression of Neurodegeneration in Drosophila, (H. Y. Zoghbi, Ed.). PLoS Biol. 6: e29. https://doi.org/10.1371/journal.pbio.0060029

Li X., G. Quon, H. D. Lipshitz, and Q. Morris, 2010 Predicting in vivo binding sites of RNA-binding proteins using mRNA secondary structure. RNA 16: 1096–1107. https://doi.org/10.1261/rna.2017210

Li J., Y. Chen, X. Xu, J. Jones, M. Tiwari, *et al.*, 2019 HNRNPK maintains epidermal progenitor function through transcription of proliferation genes and degrading differentiation promoting mRNAs. Nat. Commun. 10: 1–14. https://doi.org/10.1038/s41467-019-12238-x

Liao Y., G. K. Smyth, and W. Shi, 2014 featureCounts: an efficient general purpose program for assigning sequence reads to genomic features. Bioinformatics 30: 923–930. https://doi.org/10.1093/bioinformatics/btt656

Lim C., and R. Allada, 2013 ATAXIN-2 Activates PERIOD Translation to Sustain Circadian Rhythms in Drosophila. Science (80-.). 340. https://doi.org/10.1126/science.1234785

Lin D. M., and C. S. Goodman, 1994 Ectopic and increased expression of fasciclin II alters motoneuron growth cone guidance. Neuron 13: 507–523. https://doi.org/10.1016/0896-6273(94)90022-1

Love M. I., W. Huber, and S. Anders, 2014 Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. Genome Biol. 15: 550. https://doi.org/10.1186/s13059-014-0550-8

Lu Z., X. Guan, C. A. Schmidt, and A. G. Matera, 2014 RIP-seq analysis of eukaryotic Sm proteins identifies three major categories of Sm-containing ribonucleoproteins. Genome Biol. 15: 1–23. https://doi.org/10.1186/gb-2014-15-1-r7

Malmevik J., R. Petri, T. Klussendorf, P. Knauff, M. Åkerblom, *et al.*, 2015 Identification of the miRNA targetome in hippocampal neurons using RIP-seq. Sci. Rep. 5: 12609. https://doi.org/10.1038/srep12609

Mandel C. R., Y. Bai, and L. Tong, 2008 Protein factors in pre-mRNA 3′-end processing. Cell. Mol. Life Sci. 65: 1099–1122.

Mangus D. A., N. Amrani, and A. Jacobson, 1998 Pbp1p, a Factor Interacting withSaccharomyces cerevisiae Poly(A)-Binding Protein, Regulates Polyadenylation. Mol. Cell. Biol. 18: 7383–7396. https://doi.org/10.1128/mcb.18.12.7383

Mangus D. A., M. C. Evans, and A. Jacobson, 2003 Poly(A)-binding proteins: Multifunctional scaffolds for the post-transcriptional control of gene expression. Genome Biol. 4: 1–14. https://doi.org/10.1186/gb-2003-4-7-223

Mangus D. A., M. M. Smith, J. M. McSweeney, and A. Jacobson, 2004 Identification of Factors Regulating Poly(A) Tail Synthesis and Maturation. Mol. Cell. Biol. 24: 4196–4206. https://doi.org/10.1128/mcb.24.10.4196-4206.2004

Maulik P. K., M. N. Mascarenhas, C. D. Mathers, T. Dua, and S. Saxena, 2011 Prevalence of intellectual disability: A meta-analysis of population-based studies. Res. Dev. Disabil. 32: 419–436.

McCann C., E. E. Holohan, S. Das, A. Dervan, A. Larkin, *et al.*, 2011 The Ataxin-2 protein is required for microRNA function and synapse-specific long-term olfactory habituation. Proc. Natl. Acad. Sci. U. S. A. 108: E655-62. https://doi.org/10.1073/pnas.1107198108

Merino M. M., C. Rhiner, M. Portela, and E. Moreno, 2013 "Fitness fingerprints" mediate physiological culling of unwanted neurons in drosophila. Curr. Biol. 23: 1300–1309. https://doi.org/10.1016/j.cub.2013.05.053

Mi H., A. Muruganujan, D. Ebert, X. Huang, and P. D. Thomas, 2019 PANTHER version 14: more genomes, a new PANTHER GO-slim and improvements in enrichment analysis tools. Nucleic Acids Res. 47: 419–426. https://doi.org/10.1093/nar/gky1038

Moore M. J., 2005 From birth to death: The complex lives of eukaryotic mRNAs. Science (80-.). 309:

1514–1518. https://doi.org/10.1126/science.1111443

Moressis A., A. R. Friedrich, E. Pavlopoulos, R. L. Davis, and E. M. C. Skoulakis, 2009 A dual role for the adaptor protein DRK in Drosophila olfactory learning and memory. J. Neurosci. 29: 2611–2625. https://doi.org/10.1523/JNEUROSCI.3670-08.2009

Morgan M., 2018 BiocManager: Access the Bioconductor Project Package Repository

Morris K. J., and A. H. Corbett, 2018 The polyadenosine RNA-binding protein ZC3H14 interacts with the THO complex and coordinately regulates the processing of neuronal transcripts. Nucleic Acids Res. 46: 6561–6575. https://doi.org/10.1093/nar/gky446

Müller-McNicoll M., and K. M. Neugebauer, 2013 How cells get the message: Dynamic assembly and function of mRNA-protein complexes. Nat. Rev. Genet. 14: 275–287. https://doi.org/10.1038/nrg3434

Najmabadi H., H. Hu, M. Garshasbi, T. Zemojtel, S. S. Abedini, *et al.*, 2011 Deep sequencing reveals 50 novel genes for recessive cognitive disorders. Nature 478: 57–63. https://doi.org/10.1038/nature10423

Nakamura A., R. Amikura, K. Hanyu, and S. Kobayashi, 2001 Me31B silences translation of oocyte-localizing RNAs through the formation of cytoplasmic RNP complex during Drosophila oogenesis. Development 128: 3233–3242.

Oortveld M. A. W., S. Keerthikumar, M. Oti, B. Nijhof, A. C. Fernandes, *et al.*, 2013 Human Intellectual Disability Genes Form Conserved Functional Modules in Drosophila, (J. Flint, Ed.). PLoS Genet. 9: e1003911. https://doi.org/10.1371/journal.pgen.1003911

Ostrowski L. A., A. C. Hall, and K. Mekhail, 2017 Ataxin-2: From RNA control to human health and disease. Genes (Basel). 8: 2–21.

Oswald M. C. W., P. S. Brooks, M. F. Zwart, A. Mukherjee, R. J. H. West, *et al.*, 2018 Reactive oxygen species regulate activity- dependent neuronal plasticity in Drosophila. Elife 7. https://doi.org/10.7554/eLife.39393

Pak C., M. Garshasbi, K. Kahrizi, C. Gross, L. H. Apponi, *et al.*, 2011 Mutation of the conserved polyadenosine RNA binding protein, ZC3H14/dNab2, impairs neural function in Drosophila and humans. Proc. Natl. Acad. Sci. U. S. A. 108: 12390–5. https://doi.org/10.1073/pnas.1107103108

Park H., H. J. Kim, and B. S. Jeon, 2015 Parkinsonism in spinocerebellar ataxia. Biomed Res. Int. 2015.

Parks A. L., K. R. Cook, M. Belvin, N. A. Dompe, R. Fawcett, *et al.*, 2004 Systematic generation of high-resolution deletion coverage of the Drosophila melanogaster genome. Nat. Genet. 36: 288–292. https://doi.org/10.1038/ng1312

Patterson S., How To Focus Stack Images In Photoshop. photoshopessentials.com.

Pulst S. M., A. Nechiporuk, T. Nechiporuk, S. Gispert, X. N. Chen, *et al.*, 1996 Moderate expansion of a normally biallelic trinucleotide repeat in spinooerebellar ataxia type. Nat. Genet. 14: 269–276. https://doi.org/10.1038/ng1196-269

R Core Team, 2019 R: A Language and Environment for Statistical Computing

R Studio Team, 2018 RStudio: Integrated Development Environment for R

Ragab A., E. C. Thompson, and A. A. Travers, 2006 High mobility group proteins HMGD and HMGZ interact genetically with the Brahma chromatin remodeling complex in Drosophila. Genetics 172: 1069–1078. https://doi.org/10.1534/genetics.105.049957

Rha J., S. K. Jones, and A. H. Corbett, 2017a ZC3H14, pp. 1–7 in *Encyclopedia of Signaling Molecules*, edited by Choi S. Springer New York, New York, NY.

Rha J., S. K. Jones, J. Fidler, A. Banerjee, S. W. Leung, *et al.*, 2017b The RNA-binding protein, ZC3H14, is required for proper poly(A) tail length control, expression of synaptic proteins, and brain function in mice. Hum. Mol. Genet. 26: 3663–3681. https://doi.org/10.1093/hmg/ddx248

Robinson J. T., H. Thorvaldsdóttir, W. Winckler, M. Guttman, E. S. Lander, *et al.*, 2011 Integrative genomics viewer. Nat. Biotechnol. 29: 24–26.

Rodal A. A., A. D. Blunk, Y. Akbergenova, R. A. Jorquera, L. K. Buhl, *et al.*, 2011 A presynaptic endosomal trafficking pathway controls synaptic growth signaling. J. Cell Biol. 193: 201–217. https://doi.org/10.1083/jcb.201009052

Rodriguez-Lanetty M., *Trizol/RNeasy hybrid RNA extraction protocol*.

Rørth P., 1996 A modular misexpression screen in Drosophila detecting tissue-specific phenotypes. Proc. Natl. Acad. Sci. U. S. A. 93: 12418–12422. https://doi.org/10.1073/pnas.93.22.12418

Rørth P., K. Szabo, A. Bailey, T. Laverty, J. Rehm, *et al.*, 1998 Systematic gain-of-function genetics in Drosophila. Development 125: 1049–1057.

Rothman K. J., 1990 No adjustments are needed for multiple comparisons. Epidemiology 1: 43–6.

Rueden C. T., J. Schindelin, M. C. Hiner, B. E. DeZonia, A. E. Walter, *et al.*, 2017 ImageJ2: ImageJ for the next generation of scientific image data. BMC Bioinformatics 18: 529. https://doi.org/10.1186/s12859-017-1934-z

Sanpei K., H. Takano, S. Igarashi, T. Sato, M. Oyake, *et al.*, 1996 Identification of the spinocerebellar ataxia type 2 gene using a direct identification of repeat expansion and cloning technique, DIRECT. Nat. Genet. 14: 277–284. https://doi.org/10.1038/ng1196-277

Satterfield T. F., S. M. Jackson, and L. J. Pallanck, 2002 A Drosophila Homolog of the Polyglutamine Disease Gene SCA2 Is a Dosage-Sensitive Regulator of Actin Filament Formation. Genetics 162: 1687–1702.

Schindelin J., I. Arganda-Carreras, E. Frise, V. Kaynig, M. Longair, *et al.*, 2012 Fiji: An open-source platform for biological-image analysis. Nat. Methods 9: 676–682.

Schmid M., P. Olszewski, V. Pelechano, I. Gupta, L. M. Steinmetz, *et al.*, 2015 The Nuclear PolyA-Binding Protein Nab2p Is Essential for mRNA Production. Cell Rep. 12: 128–139. https://doi.org/10.1016/j.celrep.2015.06.008

Schneider C. A., W. S. Rasband, and K. W. Eliceiri, 2012 NIH Image to ImageJ: 25 years of image analysis. Nat. Methods 9: 671–675.

Schoenherr J. A., J. M. Drennan, J. S. Martinez, M. R. Chikka, M. C. Hall, *et al.*, 2012 Drosophila

Activated Cdc42 Kinase Has an Anti-Apoptotic Function, (H. Steller, Ed.). PLoS Genet. 8: e1002725. https://doi.org/10.1371/journal.pgen.1002725

Slowikowski K., 2019 ggrepel: Automatically Position Non-Overlapping Text Labels with "ggplot2"

Soucek S., Y. Zeng, D. L. Bellur, M. Bergkessel, K. J. Morris, *et al.*, 2016 The Evolutionarily-conserved Polyadenosine RNA Binding Protein, Nab2, Cooperates with Splicing Machinery to Regulate the Fate of pre-mRNA. Mol. Cell. Biol. 36: 2697–2714. https://doi.org/10.1128/MCB.00402-16

Stewart M., 2019 Polyadenylation and nuclear export of mRNAs. J. Biol. Chem. 294: 2977–2987. https://doi.org/10.1074/jbc.REV118.005594

Sudhakaran I. P., J. Hillebrand, A. Dervan, S. Das, E. E. Holohan, *et al.*, 2014 FMRP and Ataxin-2 function together in long-term olfactory habituation and neuronal translational control. Proc Natl Acad Sci U S A 111: E99–E108. https://doi.org/10.1073/pnas.1309543111

Takemura S. ya, Y. Aso, T. Hige, A. Wong, Z. Lu, *et al.*, 2017 A connectome of a learning and memory center in the adult Drosophila brain. Elife 6. https://doi.org/10.7554/eLife.26975

Taliaferro J. M., N. J. Lambert, P. H. Sudmant, D. Dominguez, J. J. Merkin, *et al.*, 2016 RNA Sequence Context Effects Measured In Vitro Predict In Vivo Protein Binding and Regulation. Mol. Cell 64: 294–306. https://doi.org/10.1016/j.molcel.2016.08.035

Tamanini F., C. Bontekoe, C. E. Bakker, L. Van Unen, B. Anar, *et al.*, 1999 Different targets for the fragile X-related proteins revealed by their distinct nuclear localizations. Hum. Mol. Genet. 8: 863–869. https://doi.org/10.1093/hmg/8.5.863

Tan L., K. X. Zhang, M. Y. Pecot, S. Nagarkar-Jaiswal, P. T. Lee, *et al.*, 2015 Ig Superfamily Ligand and Receptor Pairs Expressed in Synaptic Partners in Drosophila. Cell 163: 1756–1769. https://doi.org/10.1016/j.cell.2015.11.021

Tassé M. J., R. Luckasson, and R. L. Schalock, 2016 The relation between intellectual functioning and adaptive behavior in the diagnosis of intellectual disability. Intellect. Dev. Disabil. 54: 381–390. https://doi.org/10.1352/1934-9556-54.6.381

The Gene Ontology Consortium, 2019 The Gene Ontology Resource: 20 years and still GOing strong. Nucleic Acids Res. 47: D330–D338. https://doi.org/10.1093/nar/gky1055

Thurmond J., J. L. Goodman, V. B. Strelets, H. Attrill, L. S. Gramates, *et al.*, 2018 FlyBase 2.0: the next generation. Nucleic Acids Res. 47: 759–765. https://doi.org/10.1093/nar/gky1003

Tuck A. C., and D. Tollervey, 2013 A transcriptome-wide atlas of RNP composition reveals diverse classes of mRNAs and lncRNAs. Cell 154: 996–1009. https://doi.org/10.1016/j.cell.2013.07.047

Velichkova M., J. Juan, P. Kadandale, S. Jean, I. Ribeiro, *et al.*, 2010 Drosophila Mtm and class II PI3K coregulate a PI(3)P pool with cortical and endolysosomal functions. J. Cell Biol. 190: 407–425. https://doi.org/10.1083/jcb.200911020

Verkerk A. J. M. H., M. Pieretti, J. S. Sutcliffe, Y. H. Fu, D. P. A. Kuhl, *et al.*, 1991 Identification of a gene (FMR-1) containing a CGG repeat coincident with a breakpoint cluster region exhibiting length variation in fragile X syndrome. Cell 65: 905–914. https://doi.org/10.1016/0092-8674(91)90397-H

Verma V., A. Paul, A. A. Vishwanath, B. Vaidya, and J. P. Clement, 2019 Understanding intellectual disability and utism spectrum disorders from common mouse models: Synapses to behaviour. Open Biol. 9. https://doi.org/10.1098/rsob.180265

Vest K. E., B. L. Phillips, A. Banerjee, L. H. Apponi, E. B. Dammer, *et al.*, 2017 Novel mouse models of oculopharyngeal muscular dystrophy (OPMD) reveal early onset mitochondrial defects and suggest loss of PABPN1 may contribute to pathology. Hum. Mol. Genet. 26: 3235–3252. https://doi.org/10.1093/hmg/ddx206

Vissers L. E. L. M., C. Gilissen, and J. A. Veltman, 2016 Genetic studies in intellectual disability and related disorders. Nat. Rev. Genet. 17: 9–18.

Wan L., T. C. Dockendorff, T. A. Jongens, and G. Dreyfuss, 2000 Characterization of dFMR1, a Drosophila melanogaster Homolog of the Fragile X Mental Retardation Protein. Mol. Cell. Biol. 20: 8536–8547. https://doi.org/10.1128/mcb.20.22.8536-8547.2000

Wickham H., 2016 *ggplot2: Elegant Graphics for Data Analysis*. Springer-Verlag New York.

Wigington C. P., K. J. Morris, L. E. Newman, and A. H. Corbett, 2016 The Polyadenosine RNA-binding Protein, Zinc Finger Cys 3 His Protein 14 (ZC3H14), Regulates the Pre-mRNA Processing of a Key ATP Synthase Subunit mRNA * □ S. Publ. JBC Pap. Press. https://doi.org/10.1074/jbc.M116.754069

Williamson W. R., and P. R. Hiesinger, 2010 Preparation of developing and adult Drosophila brains and retinae for live imaging. J. Vis. Exp. e1936. https://doi.org/10.3791/1936

Wolff T., and D. F. Ready, 1991 *The beginning of pattern formation in the Drosophila compound eye: the morphogenetic furrow and the second mitotic wave*.

Yagi R., Y. Mabuchi, M. Mizunami, and N. K. Tanaka, 2016 Convergence of multimodal sensory pathways to the mushroom body calyx in Drosophila melanogaster. Sci. Rep. 6. https://doi.org/10.1038/srep29481

Yang Z., H. J. Edenberg, and R. L. Davis, 2005 Isolation of mRNA from specific tissues of Drosophila by mRNA tagging. Nucleic Acids Res. 33: e148–e148. https://doi.org/10.1093/nar/gni149

Yang Q., X. F. Zhang, T. D. Pollard, and P. Forscher, 2012 Arp2/3 complex-dependent actin networks constrain myosin II function in driving retrograde actin flow. J. Cell Biol. 197: 939–956. https://doi.org/10.1083/jcb.201111052

Yates A. D., P. Achuthan, W. Akanni, J. Allen, J. Allen, *et al.*, 2020 Ensembl 2020. Nucleic Acids Res. 48. https://doi.org/10.1093/nar/gkz966

Yokoshi M., Q. Li, M. Yamamoto, H. Okada, Y. Suzuki, *et al.*, 2014 Direct Binding of Ataxin-2 to Distinct Elements in 3′ UTRs Promotes mRNA Stability and Protein Expression. Mol. Cell 55: 186–198. https://doi.org/10.1016/J.MOLCEL.2014.05.022

Zhang Y., J. Ling, C. Yuan, R. Dubruille, and P. Emery, 2013 A Role for Drosophila ATX2 in Activation of PER Translation and Circadian Behavior. Science (80-.). 340.

Zhang Y., L. Zhang, X. Tang, S. R. Bhardwaj, J. Ji, *et al.*, 2016 MTV, an ssDNA Protecting Complex Essential for Transposon-Based Telomere Maintenance in Drosophila, (G. Bosco, Ed.). PLOS

Genet. 12: e1006435. https://doi.org/10.1371/journal.pgen.1006435

Zhao J., T. K. Ohsumi, J. T. Kung, Y. Ogawa, D. J. Grau, *et al.*, 2010 Genome-wide Identification of Polycomb-Associated RNAs by RIP-seq. Mol. Cell 40: 939–953. https://doi.org/10.1016/j.molcel.2010.12.011

Zwarts L., L. Vanden Broeck, E. Cappuyns, J. F. Ayroles, M. M. Magwire, *et al.*, 2015 The genetic basis of natural variation in mushroom body size in Drosophila melanogaster. Nat. Commun. 6: 1–11. https://doi.org/10.1038/ncomms10115

Figure 1. Loss-of-function alleles of *Atx2* suppress effects of *Nab2* misexpression in female and male *Drosophila*. Compared to (A) the uniform color and regimented ommatidial structure of the *Drosophila* eye in control females expressing the fated-eye-cell-specific *GMR-Gal4* driver alone, (B) overexpression of endogenous *Nab2* with *GMR-Gal4* (*Nab2^{o/e}*) induces posterior pigment loss, sporadic blackened patches, and ommatidial disorder or "roughness". Heterozygosity for either of two *Atx2* loss-of-function alleles, (C) *Atx2^{DG08112/+}* or (D) *Atx2^{X1/+}*, dominantly suppresses the pigment loss and blackened patch phenotype, with limited impact on roughness. (E-H) These genetic relationships are also observed in eyes in males. (I, J) Flies lacking functional endogenous *Nab2*, *Nab2^{ex3}* homozygotes, demonstrate dramatically decreased adult viability, as quantified by the percentage of flies reaching pupal eclosion and adulthood out of the amount expected by Mendelian inheritance. (I) In females, both loss-of-function alleles of *Atx2* significantly suppress this effect, partially rescuing viability; (J) in males, only *Atx2^{X1/+}* suppresses. Sample sizes (n) are reported in each bar and include all F1 progeny scored, including genetically distinct siblings of the genotype of interest used to calculate *% eclosed (relative to expected)*. Fisher's Exact Test (two-sided) was used to assess statistical significance. ns=not significant, ***=$p<0.001$.

Figure 2. Loss-of-function alleles of *Atx2* specifically suppress axon morphology defects in *Nab2^ex3^* mushroom body α, but not β, lobes. (A) In a representative *Nab2^pex41^* control brain, Fasciclin 2 (Fas2)-marked axons from some Kenyon cells of the mushroom body bifurcate and project dorsally into α lobes and medially into β lobes. Fas2 also marks mushroom body γ lobes and the ellipsoid body (eb) (white arrows). Representative images show heterozygosity for (B) *Atx2^DG08112/+^* or (C) *Atx2^X1/+^* induces over-projection or "fusion" of β lobes, while (D) homozygosity for the *Nab2* null allele *Nab2^ex3^* induces both β lobe fusion and the thinning or complete absence of α lobes. Heterozygosity for either (E) *Atx2^DG08112/+^* or (F) *Atx2^X1/+^* in combination with *Nab2^ex3^* partially restores proper α lobe morphology and, as quantified in (G), significantly suppresses the penetrance of α lobe defects compared to *Nab2^ex3^* alone. (H) By comparison, as quantified in (H), these *Atx2* alleles neither suppress nor enhance the penetrance of β lobe defects compared to *Nab2^ex3^* alone. Sample sizes (n) are reported in each bar and quantify, for each genotype, the total number of α lobes scored for defects and the total number of brains scored for β lobe fusion. Fisher's Exact Test (two-tailed) was used to assess statistical significance. ns=not significant, **=$p \leq 0.01$.

Figure 3. Nab2 and Atx2 primarily localize to different cellular compartments and show limited physical association in brain neurons. (A) To specifically assess protein localization in mushroom body neurons, tagged transgenic copies of Atx2 and Nab2 (Atx2-3xFLAG and Nab2-YFP) were expressed in female brains under the MB-specific *OK107-Gal4*. Kenyon cell soma, the cell bodies of the MBs, are shown for a representative brain. False-colored panels show fluorescence corresponding to α-FLAG (red, Atx2-3xFLAG), α-GFP (green, Nab2-YFP), Hoechst 33342 (blue, nuclei), and a merge of all three channels. Nab2 is localized primarily to the nuclei at steady state based on overlap with Hoechst 33342 signal, and Atx2 localizes primarily in the surrounding cytoplasm. (B) To test for physical association between Nab2 and Atx2 in brain neurons, lysates of female *Drosophila* heads, either *elav-Gal4* alone controls or *elav>Nab2-FLAG*, were subjected to co-immunoprecipitation using α-FLAG. For both genotypes, Input samples (TOT) represent 6.25% of assayed lysate and immunoprecipitation (IP) samples represent 25% of total samples eluted from α-FLAG beads. Samples were resolved via gel electrophoresis and analyzed by immunoblotting, probing with antibodies against FLAG, Atx2, Fmr1 (a positive control), or alpha tubulin (a negative control). Atx2 associates weakly with Nab2 based on its enrichment in IP samples; this association is less robust than that between Nab2 and positive control Fmr1.

Figure 4. RIP-Seq reveals overlapping sets of transcripts associate with Nab2-FLAG and Atx2-3xFLAG in brain neurons. (A) Lysates from heads of female adult flies expressing either pan-neuronal *elav-Gal4* alone as a control, *elav>Nab2-FLAG*, or *elav>Atx2-3xFLAG* were subjected to α-FLAG immunoprecipitation and immunoblotting to test IP efficacy. Input samples (TOT) represent ~6.25% of total assayed lysates and immunoprecipitation samples (IP) represent 25% of total samples eluted from α-FLAG beads. Both epitope-tag samples show robust immunoreactivity to α-FLAG in TOT and IP (arrowheads), indicating effective transgene expression and successful tagged-protein enrichment by IP. (B) Principal component analysis of 12 sequenced RNA IP samples reveals high intra-genotype reproducibility. Comparison of principal component 1 (PC1) and principal component 2 (PC2) demonstrates Nab-FLAG (teal) and Atx2-3xFLAG (maroon) samples differ more from Gal4 controls (gray) than from one another, as predicted. (C) Venn diagram of Nab2-enriched and Atx2-enriched RNAs identified by RIP-Seq, revealing that 28 shared transcripts associate with both RBPs, a significant overlap according to the hypergeometric test (***=p<0.001). (D-E) Scatter plot of all transcripts within the 5,760 of the testable set with positive (D) $log_2(Nab2\ Fold\ Enrichment)$ or (E) $log_2(Atx2\ Fold\ Enrichment)$ values. *Fold Enrichment* values quantify how effectively a transcript was enriched by IP and are derived by calculating IP/Input (i.e. percent input) values for control and epitope-tag samples and setting the average of control values to 1 (i.e. 0 on the logarithmic scale used here). Y-axes display results of significance testing, conducted by gene-by-gene one-way ANOVA, Dunnett's post-hoc test, and within-gene multiple hypothesis testing adjustment (*Dun. Adj. p*). Statistically significant transcripts (*Dun. Adj. p* < 0.05) are colored. On each plot, labels identify three transcripts among the "top" (see *Results* for details) RBP-specific RBP-associated transcripts and two transcripts (*drk*, *me31B*) among the shared RBP-associated transcripts. (F) The independent *Molecular Function* (red), *Biological Process* (green), and *Cellular Component* (blue) Gene Ontology (GO) terms most overrepresented among the shared Nab2- and Atx2-associated transcripts as compared to the entire testable transcript set. GO term independence was determined by "Hierarchical Selection" (see *Methods*). The number of GO term members within the shared RBP-associated transcripts and within the entire testable transcript set (*Genes enriched / in total*) are reported to the right of each bar.

Figure 5. Potential functionally important RNA targets of Nab2 and Atx2 identified by combining individual transcript and holistic GO analyses of RIP-Seq results. For transcripts that associate with *Both* Nab2 and Atx2, *Nab2 Only*, *Atx2 Only*, or *Neither* RBP by RIP-Seq, seven transcripts of particular functional interest are presented as a summary of each category. (A-C) These transcripts met one or both of two criteria: 1) inclusion in an associated overrepresented GO term 2) an *IP/Input (*i.e. *Fold Enrichment*) value > 1.5. Given the functions of proteins encoded by these transcripts, these selections represent potential phenotypically important targets of post-transcriptional regulation by Nab2 and Atx2. (D) These transcripts, as a negative control, encode a functionally diverse set of proteins and do not associate with Nab2 or Atx2 (*Neither*), affirming the specificity of the RNA interactome of each RBP. Error bars represent standard errors of the mean (SEM). Gene-by-gene one-way ANOVA, Dunnett's post-hoc test, and within-gene multiple hypothesis testing adjustment (*Dun. Adj. p*) was used to assess statistical significance. * = *Dun. Adj. p* < 0.05.

A

Input motif for FIMO scan

A-rich motif, 41-bp, identified in Nab2-associated transcripts by MEME		
Transcript Set	**Motif occurrences** ($p < 0.0001$)	**Motif frequency** (Occurrences/total bases in Set)
Nab2-associated	19,265	1 motif per 135 bases
All others	77,596	1 motif per 845 bases

***

B

Input motif for FIMO scan

A_{12}		
Transcript Set	**Motif occurrences** ($p < 0.0001$)	**Motif frequency** (Occurrences/total bases in Set)
Nab2-associated	1,674	1 motif per 1,553 bases
All others	34,465	1 motif per 1,901 bases

***

C

Input motif for FIMO scan

$A_{11}G$		
Transcript Set	**Motif occurrences** ($p < 0.0001$)	**Motif frequency** (Occurrences/total bases in Set)
Nab2-associated	3,786	1 motif per 687 bases
All others	70,103	1 motif per 935 bases

***

Figure 6. A broad A-rich motif and two specific, canonical Nab2 binding motifs are enriched in Nab2-associated RNAs. Output from transcript set scans by FIMO, which quantifies the occurrences in supplied sequence sets of motifs identical or highly similar to an input motif. Two transcript sets were scanned in each analysis: 1) all transcripts encoded by *Nab2-associated* gene models and 2) all transcripts encoded by *All others*, shorthand for all non-Nab2-associated gene models in the RIP-Seq testable set. (A) A 41-bp A-rich motif, identified by MEME as one of the first ten 6-50 bp motifs within *Nab2-associated* transcripts, was used as input for FIMO. (B) A canonical Nab2 binding motif from *S. cerevisiae*, $A_{11}G$, was used as FIMO input. (C) A simple homopolymer stretch of A's for which Nab2 would have a very high affinity, A_{12}, was used as FIMO input. In all three cases, particularly in (A), the scanned motif is significantly enriched in the *Nab2-associated* transcript set compared to the *All others* transcript set. However, none of the three input motifs are exclusive or nearly exclusive to Nab2-associated transcripts—each is still notably abundant within *All others*. Statistical significance was assessed using the chi-square test (two-sided). ***=$p<0.001$.

Figure 7. A model of opposing regulatory roles for Nab2 and Atx2 on shared associated RNA transcripts. Nab2 and Atx2 associate with a shared set of RNA transcripts in *Drosophila* brain neurons but primarily localize to separate subcellular compartments and weakly physically associate. *S. cerevisiae* Nab2 regulates nuclear processing, including transcript stability, poly(A) tail length, and export, across a broad RNA transcript set—*Drosophila* Nab2 may perform similar functions on its comparatively limited associated RNA set. Atx2 serves numerous roles in post-transcriptional regulation, including as a miRNA-machinery linked translational repressor. Taken together, these data imply the following model. (*Top*) In the wild-type condition, Nab2 protects transcripts from degradation, limits poly(A) tail length, and contributes to *Target* RNA export from the nucleus, shuttling with its associated transcripts into the cytoplasm. Nab2 and Atx2 may co-occupy the same transcripts briefly or occasionally during nuclear-cytoplasmic mRNP remodeling and prior to Nab2 recycling into the nucleus. Atx2 accompanies *Target* transcripts through transport to their destinations (e.g. synaptic terminals) and contributes to miRNA-mediated translational repression, which is released under certain conditions (e.g. synaptic activity), ultimately contributing to regulated production of wild-type levels of Target protein (black —). (*Second panel*) In *Nab2^{ex3}* nulls, *Target* mRNAs are less stable, exhibit longer poly(A) tails, and are exported less efficiently from the nucleus. As a result, less *Target* mRNA reaches its appropriate destination, resulting in a decrease in steady-state levels of Target protein (red ↓). (*Third panel*) In Atx2 loss-of-function heterozygotes (i.e. *Atx2$^{DG08112/+}$* or *Atx2$^{X1/+}$*), less Atx2 protein is expressed and available to repress *Target* translation, resulting in less responsive, higher steady-state levels of Target protein (green ↑). (*Bottom*) Effects of the complete loss of Nab2 in *Nab2^{ex3}* and the decrease of functional Atx2 in *Atx2* loss-of-function heterozygotes balance one another. While nuclear *Target* mRNA is less stable and less is exported from the nucleus successfully, these RNAs are also under less strict translational control in partial absence of Atx2, ultimately resulting in Target protein levels and corresponding phenotypes more similar to the wild-type condition (black —). This model represents a prediction from our data and the published knowledge of the functions of each protein—it must be tested in future research, a task enabled by the identification of Nab2- and Atx2-associated transcripts in the current study.