ORT: A workflow linking genome-scale metabolic models with reactive transport codes

Rebecca L. Rubinstein[1,*], Mikayla A. Borton[2], Haiyan Zhou[1], Michael Shaffer[2], David W. Hoyt[3],

James Stegen[3], Christopher S. Henry[4], Kelly C. Wrighton[2] and Roelof Versteeg[1,*]

[1]Subsurface Insights, LLC., Hanover, NH,

[2]Department of Soil and Crop Sciences, Colorado State University, Fort Collins, CO,

[3]Pacific Northwest National Laboratory, Richmond, WA,

[4]Argonne National Laboratory, Lemont, IL

*To whom correspondence should be addressed.

Abstract

Motivation: Nutrient and contaminant behavior in the subsurface are governed by multiple

coupled hydrobiogeochemical processes which occur across different temporal and spatial scales.

Accurate description of macroscopic system behavior requires accounting for the effects of

microscopic and especially microbial processes. Microbial processes mediate precipitation and

dissolution and change aqueous geochemistry, all of which impacts macroscopic system behavior.

As 'omics data describing microbial processes is increasingly affordable and available, novel

methods for using this data quickly and effectively for improved ecosystem models are needed.

Results: We propose a workflow ('Omics to Reactive Transport – ORT) for utilizing

metagenomic and environmental data to describe the effect of microbiological processes in

macroscopic reactive transport models. This workflow utilizes and couples two open-source

software packages: KBase (a software platform for systems biology) and PFLOTRAN (a reactive

transport modeling code). We describe the architecture of ORT and demonstrate an

implementation using metagenomic and geochemical data from a river system. Our demonstration

uses microbiological drivers of nitrification and denitrification to predict nitrogen cycling patterns

which agree with those provided with generalized stoichiometries. While our example uses data

from a single measurement, our workflow can be applied to spatiotemporal metagenomic datasets

to allow for iterative coupling between KBASE and PFLOTRAN.

Availability and Implementation: Interactive models available at

https://pflotranmodeling.paf.subsurfaceinsights.com/pflotran-simple-model/. Microbiological data

available at NCBI via BioProject ID PRJNA576070. ORT Python code available at

https://github.com/subsurfaceinsights/ort-kbase-to-pflotran. KBase narrative available at

https://narrative.kbase.us/narrative/71260 or static narrative (no login required) at

https://kbase.us/n/71260/258

Contact: rebecca.rubinstein@subsurfaceinsights.com or roelof.versteeg@subsurfaceinsights.com

Supplementary information: Supplementary data are available online.

1   Introduction

The critical zone (CZ) – the area between the top of the forest canopy and the bottom of

the groundwater table is essential in sustaining life (Guo and Lin, 2016). Being able to understand

and predict critical zone function is essential for both scientific and operational purposes. This

understanding and prediction requires the accurate representation of key hydrobiogeochemical

ecosystem processes which occur and interact in the critical zone. These ecosystem processes

operate at different scales and have different drivers, but at the same time are tightly

interconnected. For instance, while hydrological processes control the movement of water at

macroscopic scales and are driven by groundwater table gradients, precipitation, and

evapotranspiration whereas microbiological processes (Anantharaman *et al.*, 2016; Long *et al.*,

2016) occur at the microbe scale and are driven by microbial populations, soil properties, aqueous

geochemistry, and temperature. However, processes at these two scales influence one another in

many ways.

One well-established approach to obtaining an understanding of critical zone behavior is

through the use of reactive transport models (RTM) which can simulate coupled chemistry, flow,

and transport in hydrobiogeochemical systems. There are a variety of reactive transport codes (see

(C. I. Steefel *et al.*, 2015) for a review). These models are generally continuum scale models

which represent subsurface properties on grids, with grid volumes on the order of cubic meters. As

the earth is a porous media with grains and pores, such continuum scale models obviously do not

capture pore scale properties and dynamics. One fundamental challenge in numerical modeling is

thus how to link and couple processes and properties which happen at different scales (Battiato *et*

al., 2011; Chu *et al.*, 2012, 2013; Carl I. Steefel *et al.*, 2015). Such linking is especially required

between macroscopic system behavior and microbial processes which change aqueous

geochemistry and mediate precipitation and dissolution.

With continued decrease in 'omics data analysis costs, one promising approach for this

linking is through the incorporation of site-specific microbiological data into RTM to represent

microbe-catalyzed biogeochemical more accurately than using generalized stoichiometries. The

feasibility of using the results of microbiological data analysis to parameterize RTM has been

shown previously. For instance, Scheibe et al. demonstrated the linking of genome scale models

with a reactive transport code (in their case, HYDROGEOCHEM) to improve incorporation of

microbiological processes on in situ uranium bioremediation (Scheibe *et al.*, 2009). Specifically,

they used a genome scale model of *Geobacter sulfurreducens* to populate a lookup table spanning

reasonable expected ranges for all combinations of three key system parameters. This was then

used to predict the effects of varying concentrations of three key growth factors (acetate, Fe(III),

and ammonium) on reduction of uranium (VI) at a systems level. More recently, Song et al.

developed an enzyme-based approach for simulating microbial reaction kinetics which captured

the overall behavior of a consortium rather than rely on individual taxa within the community and

coupled it with reactive transport simulations using PFLOTRAN's Reaction Sandbox (Song and

Liu, 2015; Song *et al.*, 2017; Hammond *et al.*, 2017). This approach is based on a mechanistic

understanding of microbial processes and thus can more accurately predict microbial response to

perturbations. However, this approach substantial experimental data, such as enzyme

concentrations and kinetics data, as well as advanced microbiological knowledge to implement.

These previous efforts have demonstrated the value and feasibility of accurately

representing microbial processes in RTM. This in turn opens up the potential to integrate

microbiological, geochemical, and physical subsurface properties and processes to predict

ecosystem behavior and response (Fig. 1).

Fig. 1 – Accurately capturing the interplay between microbiology, geochemistry, and physical
subsurface properties and processes is critical to understanding and predicting ecosystem
processes.

However, each of the approaches described above requires substantial manual effort to

implement for a single site, which makes them challenging to scale. The challenge of scaling these

approaches limits the ability to rapidly develop models obtain the associated understanding for

many sites. An alternative approach (proposed and demonstrated here) is to an approach which

allows for automation.

Specifically, in our workflow we use KBase (a cloud-based software platform for systems

biology (Arkin *et al.*, 2018)), to automatically generate draft metabolic models from annotated

metagenome assembled genomes (MAGs) extracted from environmental samples. These

metabolic models can be used (still in KBase) to perform flux balance analysis (FBA) on different

media compositions. These media compositions are informed by metabolomics and other site-

specific chemistry data. The output of the FBA can be used in reactive transport models (such as

the reactive transport model PFLOTRAN (Mills *et al.*, 2009; Hammond and Lichtner, 2010;

Gardner *et al.*, 2015)), which we use in this work. PFLOTRAN is an open source, massively

parallel reactive transport code which supports multi-phase (e.g. aqueous, gaseous), multi-

component (multiple chemical species), and multi-scale (e.g. pore or macroscale) simulation of

contaminant transport in porous media, as well as includes a basic implementation of microbial

reactions modeled by Monod kinetics. One major benefit of PFLOTRAN is that users can

implement custom reactions or kinetics through the Reaction Sandbox (Hammond, 2017). Our

workflow, called 'Omics to Reactive Transport (ORT) (Fig. 2), thus captures both microbial

metabolisms based on environmental samples (using KBase) and macro-scale hydrologic and

geochemical processes (using PFLOTRAN). In the remainder of this paper we present the concept

and implementation of this workflow.

Fig. 2 - Omics to Reactive Transport (ORT) workflow couples microbe-scale and macroscale
processes using the outputs of KBase and PFLOTRAN as inputs for each other.

In addition to the scientific value of this workflow, we want to highlight three operational

attributes of interest. First, this workflow can be mostly automated, offering the potential of

rapidly generating reactive transport models from microbiological data (detailed in Section 3) with

a minimum of manual labor. Second, the resulting models can easily be shared and made

accessible to other groups. For instance, we have provided two of the models we generated

through a user-friendly web interface which allows end-users to interact with these models. Third,

while in this paper we do not include results for this, our workflow lends itself well to an iterative

approach. Specifically, it is well known that microbial processes will result in changes in

geochemistry, which in turn will influence the microbial processes. In addition to this,

macroscopically driven changes in saturation, temperature, and chemistry (e.g., resulting from

stage-driven surface water/ground water interaction) will also influence microbial processes. The

approach described here can be executed in an iterative manner to capture this two-way coupling

between microscopic and macroscopic processes.

2   System & Methods

2.1 Workflow Concept

The ORT workflow was designed with automation in mind. Specifically, it is designed in a

modular manner with a well-defined start and end points and inputs and outputs, with each

component being fully automatable (Fig. 3). The inputs to this workflow are annotated genomes,

environmental chemistry, and a PFLOTRAN model template which incorporates physical site

data. This template (which would be customized to the specific site) would be something like "0D

batch reactor" or "2D model of unsaturated soil" (where "nD" indicates the number of spatial

dimensions accounted for in the model grid).

In this workflow we import annotated genomes (Shaffer et al., 2020) and site chemistry

(e.g., available carbon sources, electron acceptors, and micronutrients based on metabolome and

any other chemical analysis at the site, synthesized into a KBase media recipe) into KBase, then

use KBase apps to generate the overall reactions. Next, these reactions, as well as the site

chemistry, physical site data, and model template are used to build the actual PFLOTRAN model.

This model can then be used to simulate macroscopic system behavior.

In the iterative implementation, the PFLOTRAN model simulates changes in physical and

chemical conditions in space and time. We can then use the simulated chemistry as a new media

composition to be used by the KBase part of the workflow and repeat the process to generate and

retrieve new resulting overall reactions and substitute them into the PFLOTRAN input file.

Fig. 3 - Flowchart of the ORT workflow where orange boxes are workflow inputs based on site characterization which are pre-processed before use, green boxes are metabolic modeling steps carried out in KBase, and blue show the resulting RTM. The horizontally-aligned boxes and arrows in the KBase workflow represent robust curation steps (discussed in Section 4), and the dashed arrow indicates the iteration path wherein the PFLOTRAN-simulated chemistry is used as a new media condition in KBase.

2.2 Workflow Implementation

The ORT workflow consists of Python scripts, KBase narratives, and PFLTORAN models.

In our implementation of ORT, KBase apps import genomes and chemical data into KBase and

use these as inputs for KBase metabolic modeling apps (process described in detail in Sections 2.4

and 2.5). After the completion of the KBase part of the workflow, the KBase API (application

programming interface) programmatically exports the KBase-predicted exchange fluxes from

KBase. These fluxes are translated by our Python script into an overall reaction string that

describes chemical uptake and secretion from each modeled organism, written in PFLOTRAN-

compatible naming conventions. The flux values are used as the stoichiometric coefficients for the corresponding chemicals in the overall reaction used in PFLOTRAN, with positive fluxes indicating reactants and negative fluxes indicating products. The summation of exchange fluxes is not a chemical reaction in the traditional sense, but represents the chemical species removed from and added to the system as a result of the microbial metabolism. Thus, this "pseudo-reaction" provides the information needed by PFLOTRAN to simulate the resulting changes in chemical concentrations.

The ORT Python script outputs a *.txt* file with the reaction strings and yield terms for use in the MICROBIAL_REACTION card in PFLOTRAN as well as a set of *.dat* files which contain compound names and details which need to be added to the PFLOTRAN geochemical database (formatted for compatibility with the database). This step can either be done programmatically or manually by substituting the content of these text files into a PFLOTRAN model input file (known as an infile). This script bridges the disconnect between KBase and PFLOTRAN illustrated in Fig. 2.

2.3 Test Case

2.3.1   System Description

To evaluate the performance of our workflow, we used environmental samples from a hyporheic zone in the Columbia River. In these zones, biological nitrogen cycling is known to occur (Triska *et al.*, 1993; Zheng *et al.*, 2016). Biological nitrification and denitrification is a classic, well-understood, and extensively studied system. We can calculate and compare models which use traditional (textbook) stoichiometries for nitrification and denitrification to the model generated from our workflow. In the remainder of this paper, we refer to these two models as the "literature based model" and "genome derived model".

Nitrification is traditionally split into two sub-processes, ammonium oxidation (NH4 →

NO2) and nitrite oxidation (NO2 → NO3), while denitrification is often represented as a complete

process (NO3 → N2), though in reality it is several sequential reactions. Within KBase, we could

implement separate models for each step for which genomes are available, but for comparison to

the traditional model we used a single model for complete denitrification in this test case. The

overall reactions used for the nitrification step were based on experimentally-determined

stoichiometries (Liu and Wang, 2012) determined by fitting data collected from bench-scale

reactors to traditional half-cell reactions (Rittmann and McCarty, 2012), as given by the following

reactions:

$1.0225\ NH_4^+ + 1.3875\ O_{2\ (aq)} + 0.09\ CO_{2\ (aq)} + 0.0225\ HCO_3^-$

187                       $\rightarrow 2H^+ + NO_2^- + 0.0225\ Biomass$

189     $1\ NO_2^- + 0.0073\ NH_4^+ + 0.4635\ O_{2\ (aq)} + 0.0292\ CO_{2\ (aq)} + 0.0073\ HCO_3^-$

190                       $\rightarrow NO_3^- + 0.0073\ Biomass$

191   The complete denitrification process stoichiometry was derived from half-cell reactions

192   (Rittmann and McCarty, 2012), scaled to one unit nitrate utilization for comparability with the first

193   two reactions:

194     $1\ NO_3^- + H^+ + 0.869\ CH_3COO^-$

195                     $\rightarrow 0.458\ N_{2\ (aq)} + 0.444\ CO_{2(aq)} + 0.869\ HCO_3^- + 0.08484\ Biomass$

In both cases, the chemical species represented are limited to classical compositions, which

in some cases may serve as analogs for a range of compounds. These stoichiometries are not

associated with any specific microbes or metabolic pathways, but rather represent the exchange

fluxes observed. While this approach is very effective for process design, it does not offer much

insight into the microbiology of a system, and may obscure finer-scale dynamics – such as less

obvious resources that may become limiting or change how the microbes process available

macronutrients, particularly in systems with complex carbon sources.

The rates determined through batch kinetics tests (Liu and Wang, 2012) were used for

ammonium oxidation and nitrite oxidation and the denitrification rate was based on rates reported

in the literature (Raboni *et al.*, 2014). The same rates (shown in Table 1) were used for both the

literature-based and genome-based models (described in Section 2.5 - 2.66) in order to directly

compare the effects of the different stoichiometries. In future enhancements, we anticipate that

reaction rates could be used as tunable parameters to fit these models to system-specific

experimental data.

Table 1 – Baseline reaction rates used in nitrogen-cycling models

Process	Rate (mol/L·s)
Ammonium Oxidation	1.0×10^{-7}
Nitrite Oxidation	8.51×10^{-8}
Nitrate Reduction	2.34×10^{-8}

2.3.2   Leveraging Existing Multiomics Data

This study made use of multiomics data from previously published work. Sediment was

collected and DNA extracted as previously described (Graham et al., 2017). To identify the

metabolites available to microorganisms in these river sediments, we performed 1H Nuclear

Magnetic Resonance (NMR) spectroscopy on 17 paired sediment pore water samples which also

had microbial DA extracted, as described previously (Tfaily *et al.*, 2019). Briefly, sediment

samples were mixed with water in a 1:1 ratio and then diluted by 10% (vol/vol) with 5 mM 2,2-

dimethyl-2-silapentane-5-sulfonate-d6 as an internal standard. The 1D 1H NMR spectra of all

samples were processed, assigned, and analyzed using Chenomx NMR Suite 8.3 with

quantification based on spectral intensities relative to the internal standard as described. To obtain

a representative bulk summary of the metabolite environment in these sediments, the

concentration of 31 of the NMR identified metabolites was averaged across the 17 sediment

samples, and this data was used as the chemical data input in our ORT workflow (data available in

Supplementary Table S1).

Purified genomic DNA was sent to the Joint Genome Institute (JGI, n=33) under

JGI/EMSL proposal 1781 and to the Genomics Shared Resource facility at The Ohio State

University (OSU, n=10), producing 43 metagenomes from 34 sediment samples with an average

sequencing depth of 3.84 (JGI) 25 Gbp (OSU) per sample. JGI and OSU sequencing was

performed as previously described in Graham et al (Graham *et al.*, 2018) and Borton et al (Borton

et al., 2018) respectively. Raw reads were processed, assembled, and binned as outlined in

previous publications (Shaffer *et al.*, 2020) or via the Wrighton Lab GitHub Page

(https://github.com/TheWrightonLab). The genomes are available on NCBI via BioProject ID

PRJNA576070.

From the sediments, we obtained metagenome assembled genomes (MAGs) from which

we selected four genomes that represented key parts of the nitrogen cycle. For each stage of the

cycle, the most complete genomes capable of filling those roles were selected. To represent

nitrification, we chose the most complete genome representatives of the ammonium oxidizing

archaea classified by GTDB-Tk (version 1.3.0, as of 1-21-21) as a member of the family

Nitrososphaeraceae within the genus TA-21 (previously within the Phylum Thaumarchaeota) and

nitrite oxidizing bacterial member of the Nitrospiraceae for nitrification. Given that the expression

and activity of nitrite reductase encoded in Nitrososphaeraceae (previously Thaumarchaeota) is

poorly understood at this time (Kuypers et al., 2018), we did not incorporate the production of

nitric oxide by Nitrososphaeraceae, and focused only on nitrite outputs from ammonification. To

represent denitrification, we selected two Gammaproteobacterial MAGs, both classified within the

family Steroidobacteraceae. Note that neither of these genomes encoded a gene to produce N_2 gas,

but the reaction to convert nitrous oxide to nitrogen gas was added to the metabolic models during

gapfilling (see Section 2.5). We selected only four genomes to maintain the simplicity of this

proof of concept, but the approach could incorporate as many as are needed to capture system

behavior. Each nitrogen-cycling genome was annotated using DRAM (Distilled and Refined

Annotation of Metabolism (Shaffer et al., 2020)) with default parameters. The raw annotations

containing an inventory of all database annotations for every gene from each input genome are

included in the online Supplementary Materials. These genomes and their annotations were

uploaded to KBase (Section 2.5) and were the basis for the KBase-derived model (Section 2.6).

2.4 Pre-Processing

Prior to executing the workflow, we need to gather and preprocess data and make several

decisions such as selecting a model template. In this section, we describe the data preprocessing

steps in generic terms, as the same steps will be required for any system. To begin our workflow,

user inputs were organized and prepared, which consisted of three broad steps:

(1) Qualitative assessment – to balance model complexity and utility, the system definition

phase began with a qualitative description of the system in terms of model type (batch,

chemostat, continuously stirred tank reactor, etc.), important processes (such as nitrification

or sulfur reduction, depending on the system), and parameters of interest (pH, specific

chemical species, etc.) that can guide model development. This step includes evaluating if

there is any "missing" data, which might render the model inaccurate or impossible, and

would need to be estimated in order to produce a viable system (for example, concentrations

of biologically necessary compounds that were not measured). These are identified through a

combination of subject matter knowledge and comparison with KBase default media recipes.

Note that this does not entail delineation of every process and parameter involved in the

system, but rather selection of those important to the specific research or application. The

goal of this step is to develop a conceptual model of the system of interest, which may be

augmented and refined as needed to accommodate new data. Because many of these models

will be similar (e.g. 0D batch models), we can build a library of model templates which can

be readily reused.

(2) Data Gathering - data describing the site may be drawn from a variety of sources, including

direct sampling at the site and public resources such as weather stations or national

databases. Biological data could come in the form of annotated genomes or metagenomes

collected from the site, or genomes for key microbes as determined using 16S rRNA gene

data or literature review could be drawn from public databases. Chemical data could include

traditional geochemical analysis as well as metabolomics and metaproteomics to provide a

more detailed picture of the chemical profile at the site. Physical data could include

temperature, soil porosity, or other parameters of that nature that would be included in the

PFLOTRAN input file to produce a more site-specific model.

(3) Translation to KBase and PFLOTRAN - the data produced by the various analyses above are

not necessarily in formats that may be directly imported to KBase and/or PFLOTRAN.

Therefore, the final step in this phase was to translate these data to forms that can be used by

the relevant tools (KBase or PFLOTRAN). Aside from managing file formats (see the

KBase documentation for details), one major consideration was accounting for any un-

measured chemical species identified in the first step of the preparation phase that needed to

be added to the KBase media composition to make it biologically viable or usable by the

metabolic models generated in KBase. Additions were limited to chemical species or

compounds known (or reasonably expected) to be present and were added in sufficient

concentration that they would not be growth-limiting. The primary check for the presence

assumption was that the experimental data indicated that the microbes used were both

present and involved in nitrogen cycling at that site. We did not investigate the assumption

that these compounds were non-limiting, as this is outside the scope of this work.

2.5 KBase Metabolic Modeling

Once pre-processing was complete, we can start the ORT workflow. Genomes were uploaded to

KBase as paired FASTA and GFF3 text files using the "Import GFF3/FASTA file as Genome

from Staging Area" app and then annotated with RASTtk using the "Annotate Microbial Genome"

app in KBase. Additional custom annotations from DRAM were uploaded as flat text files using

the beta version of "Import Annotations from Staging" app. If using DRAM annotations,

preprocessing may be carried out using the provided script at

https://github.com/subsurfaceinsights/ort-kbase-to-pflotran. Notably, both RASTtk and DRAM are

available as apps in KBase, allowing users to functionally annotate genomes without high memory

computational resources. However, note that the DRAM app in KBase differs from the version

used in this example narrative (Shaffer *et al.*, 2020) as the KBase DRAM app annotates using

KOfam instead of KEGG genes and does not currently include EC reaction identifiers, so end

results may differ from the included narrative. Chemical data was uploaded as flat text files using

the "Import Media file (TSV/Excel) from Staging Area". The use of pre-processed flat text files as

inputs to the workflow significantly simplifies the process compared to using raw data, especially

for genomes, and these can be generated automatically using scripts such as the one developed for

the DRAM outputs. This first step brought all of our data in the KBase workspace in an integrated

manner.

After this step, we used all this data as inputs to the "Build Metabolic Model" app, and the

generated models were used in conjunction with the media objects as inputs to the "Run Flux

Balance Analysis" (FBA) app. Going forward, we refer to this pairing as "growing a model,"

meaning we ran the analysis to determine if biomass growth was possible under the given

chemical conditions. The output from the FBA app included the reaction and exchange fluxes for

each model grown on the corresponding media.

2.6 PFLOTRAN Reactive Transport Modeling

We used our workflow to download the FBA exchange flux values using the KBase API and

translate them from KBase objects with ModelSEED (Henry *et al.*, 2010) compound IDs to flat text

files with reaction strings written using PFLOTRAN naming conventions. We then used either the

KBase-derived reaction strings and biomass yield values or the literature-based stoichiometries

introduced in Section 2.3 to fill in the MICROBIAL_REACTION card in our 0D model template.

All parameters except the reactions and yield terms were held the same for both the literature-based

model and the genome-derived model.

3   Model behavior and General behaviors and trends

Both models exhibited sequential ammonium and nitrite oxidation followed by nitrate

reduction, ultimately producing dissolved nitrogen gas (Fig. 4). Despite using the same reaction

rates, inhibition constants, and initial nutrient concentrations, the overall progress of the system is

noticeably different. The genome-derived model exhausts the available ammonium within 1.5 hours

of the simulation start, while the literature based model does not exhaust ammonium until a little

more than 3.5 hours into the simulation. Nitrite concentration peaks earlier and at a lower level for

the genome-derived model (~18 µM at approximately 1 hr) than the literature based model (~51 µM

slightly before 3 hrs). Similarly, nitrate peaks at approximately 4 µM after 1.5 hrs for the genome-

derived model but peaks at 40 µM at the 6 hr mark for the literature based model. In the 6 hour

period shown in Fig. 4, the genome derived model has exhausted ammonium, nitrite, and nitrate,

while the literature based model is still processing nitrite and nitrate. This variance is expected since

we are comparing generic reactions (with generic substrate utilization and biomass production

reactions) to site-specific reactions based on the most dominant taxa found at our study site.

One important difference was that the microbiologically-explicit, genome-based

stoichiometry provided much greater detail on the chemistry, particularly with respect to carbon

catabolism (Fig. 4 and Fig S1). Specifically, the literature-based models relied entirely on either

carbon dioxide (nitrification) or acetate (denitrification), however, because we provided additional

carbon compounds detected from our bulk sediment metabolome, the site models used 15 to 23

unique additional carbon sources, such as betaine, leucine, and choline (see Supplementary Table

1). This greater detail allows us to evaluate more precisely the potential chemical drivers or

limiters of a system which would be entirely overlooked with traditional representations, which

presents the opportunity to probe and improve our conceptual and mechanistic understanding of

these systems and individual metabolisms.

Fig. 4 - Using our Omics to Reactive Transport (ORT) workflow allows us to not only tailor a model to a specific environmental site and system, but also provides much finer insight into the changes in chemistry driven by microbial processes. The top frame shows the steps captured by the literature-based and genome-informed models respectively. The middle frame shows graphical representations of the two sets of reaction stoichiometries. Abbreviations used in the site-specific model frame are Met for Methionine, Thr for Threonine, and SAO for S-Adenosyl-4-methylthio-2-oxobutanoate, which are compounds predicted by KBase as an output which is not part of standard literature representations. The bottom frame shows the results of using each set of reactions in a 0D PFLOTRAN simulation of nitrogen cycling.

361   Instead of generic bacterial enzymatic reactions, we can determine which site specific bacterial –

362   or archaeal – reactions are drivers in the system. Instead of pre-set stoichiometries, our 'Omics to

363   Reactive Transport workflow uses chemistry determined based on metabolomics – using these

364   data to describe the initial chemistry rather than generic or simplified chemistry. For example,

365   even with the same rate constants, we can see that the genome-informed model utilizes a higher

proportion of ammonium in the first step of nitrification, resulting in more rapid depletion of ammonium in the system and earlier generation of nitrite. As a result, subsequent steps begin earlier, resulting in an overall accelerated process. At the same time, both versions exhibit the expected cycling of ammonium to nitrite to nitrate and finally to nitrogen gas. Since PFLOTRAN relies on user-defined chemistry (as opposed to automatically generating reactions), this allowed us to incorporate more realistic, mechanism-driven reactions.

The genome-based model also allows for greater chemical breadth. The nitrogen cycling reactions are modulated by a wider range of carbon sources. Additionally, the by-products of this carbon and nitrogen metabolism also resulted in more complex chemical outputs in some cases, such as L-Threonine or L-Methionine. These inferred reactions could be further refined by using gene expression data (e.g., metatranscriptomics or metaproteomics data) to calibrate the models (by way of reaction rates, saturation constants, etc.) to a particular set of environmental conditions. Again, this presents an opportunity to test and enhance our understanding of the metabolic processes involved.

Readers can explore and interact with both of these models (without sign-in) through Subsurface Insights' web-based PFLOTRAN interface at https://pflotranmodeling.paf.subsurfaceinsights.com/pflotran-simple-model/. For the literature-based model, we have made the input concentrations of ammonium, bicarbonate, and acetate accessible to web users using sliders. For the Hanford 300 Area-specific version of the model, we have made accessible the reaction rate for each of the steps modeled. There is no limit to the number of parameters that may be exposed this way, but for the sake of a user-friendly and un-cluttered demonstration, we limited our selections to three per model. We selected the parameters we did both because the effects of varying them are significant and to highlight the power and flexibility provided by this approach.

4 Discussion

We demonstrated an Omics to Reactive Transport (ORT) workflow for creating site specific reactive transport models that include local chemical and biological content. The ORT workflow was applied to a well-understood system, and the results agree generally with expected behavior in a nitrogen cycling system. We interpret the differences in magnitude and timing to be due to the difference between generic, simplified reactions and metabolism-informed reactions, as KBase-derived stoichiometries made it possible to capture microbial metabolism in much greater detail than conventional approaches allow.

While the model predictions are borne out by comparison to traditional models, we would need extensive new data which currently is not available to comprehensively validate our modeling results. Specifically, we would need high resolution time series data. Such data was not available in this effort, but is a component of ongoing work, and is in general becoming increasingly available as technology improves and cost per sample decreases. Given similar data types, the same workflow could be applied to build and tune a model for other sites.

Much of the future work on this workflow will be focused on enhancing and expanding automation and on making it more robust in several ways. One capability which would be highly beneficial to our workflow is automated metabolic model curation. In our effort, curation was carried out manually using two different approaches: metabolism-based and media-based. The former is labor intensive and requires substantial subject-matter expertise to carry out. The latter is more straightforward and relies on a more general system understanding, but still requires manual iteration to obtain reasonable results. Partially or fully automated model curation will eventually be needed for full automation. This is a topic of active effort by both the KBase core team and other groups, and we will leverage their efforts. Additional work will be in expanding PFLOTRAN models to include processes such as temperature mediated biological processes and

material recycling. While these are currently not part of the core PFLOTRAN capabilities, these can be implemented using the PFLOTRAN sandbox.

While previous researchers have demonstrated the feasibility of coupling genome-scale metabolic models with reactive transport simulations, our work is different in some fundamental ways. First, our workflow, lends itself to automation and rapid model generation from 'omics data. As 'omics data becomes increasingly affordable, the ability to rapidly translate this data into information on its the implications for macroscopic system behavior will be needed, and our workflow provides a path towards that. Second, our workflow lends itself to easy incorporation of more realistic microbial reaction kinetics (e.g., based on temperature or soil conditions). Third, our workflow lends itself to iteration, which allows us to couple microscopic and macroscopic processes in either direction. Finally, our workflow provides an easy way to couple two powerful and complex software packages which typically are used by scientist in different domains, and allows these scientists a path to generate 'omics informed reactive transport models.

Funding

This work has been supported by the SBIR Award DE-SC0019619, Integrated Management and Analysis Platform for Multi Domain Site Data (program manager Paul Bayer) from the DOE Biological and Environmental Research program. A portion of the metagenomic sequencing for this research was performed by the Department of Energy's Joint Genome Institute (JGI) via sequencing award no. 1781. Metabolite support was provided by Environmental Molecular Sciences Laboratory (EMSL) via award no. 50334. Both JGI and EMSL facilities are sponsored by the Office of Biological and Environmental Research and operated under contract nos. DE-AC02- 05CH11231 (JGI) and DE-AC05-76RL01830 (EMSL). A portion of this work was supported by multiple grants within the Wrighton Laboratory: National Sciences Foundation

Division of Biological Infrastructure under award no. 1759874, DOE Early Career award no. DE-

SC0018020, and DOE award no. FY21.1068.001. Field sample collection and processing was part

of the Scientific Focus Area (SFA) project at PNNL, sponsored by the U.S. Department of Energy,

Office of Science, Environmental System Science (ESS) Program. This contribution originates

from the ESS Scientific Focus Area (SFA) at the Pacific Northwest National Laboratory (PNNL).

Acknowledgements

Tasya Rodzianko, Doug Johnson, and Erek Alper at Subsurface Insights work on the

cyberinfrastructure and web interface that was used in this work. Garret Smith, Pengfei Liu, and

Lindsey Solden provided additional microbiological expertise and processing. Field sample data

was collected by Evan Arntzen, Alex Crump, Brad Fritz, Dave Kennedy, Sarah Fansler, Nate

Phillips, Sadie Montgomery, Kyle Parker, and Rob Macklet at Pacific Northwest National

Laboratory. Processing of fine sediments was also performed by Ray Clayton and Chris Strickland

and cultural support was provided by Doug McFarland and Joy Ferry.

Conflict of Interest: none declared.

References

Anantharaman,K. *et al.* (2016) Thousands of microbial genomes shed light on interconnected
biogeochemical processes in an aquifer system. *Nat Commun*, **7**, 13219.
Arkin,A.P. *et al.* (2018) KBase: The United States Department of Energy Systems Biology
Knowledgebase. *Nat Biotechnol*, **36**, 566–569.
Battiato,I. *et al.* (2011) Hybrid models of reactive transport in porous and fractured media.
Advances in Water Resources, **34**, 1140–1150.
Borton,M.A. *et al.* (2018) Coupled laboratory and field investigations resolve microbial
interactions that underpin persistence in hydraulically fractured shales. *Proc Natl Acad Sci*
USA, **115**, E6585–E6594.
Chu,J. *et al.* (2012) A Multiscale Method Coupling Network and Continuum Models in Porous
Media I: Steady-State Single Phase Flow. *Multiscale Model. Simul.*, **10**, 515–549.
Chu,J. *et al.* (2013) A Multiscale Method Coupling Network and Continuum Models in Porous
Media II—Single- and Two-Phase Flows. In, Melnik,R. and Kotsireas,I.S. (eds), *Advances*
in Applied Mathematics, Modeling, and Computational Science, Fields Institute
Communications. Springer US, Boston, MA, pp. 161–185.
Gardner,W.P. *et al.* (2015) High Performance Simulation of Environmental Tracers in
Heterogeneous Domains. *Groundwater*, **53**, 71–80.

Graham,E.B. *et al.* (2018) Multi 'omics comparison reveals metabolome biochemistry, not microbiome composition or gene expression, corresponds to elevated biogeochemical function in the hyporheic zone. *Science of The Total Environment*, **642**, 742–753.

Guo,L. and Lin,H. (2016) Critical Zone Research and Observatories: Current Status and Future Perspectives. *Vadose Zone Journal*, **15**.

Hammond,G.E. *et al.* (2017) Application of a hybrid multiscale approach to simulate hydrologic and biogeochemical processes in the river-groundwater interaction zone. Sandia National Lab. (SNL-NM), Albuquerque, NM (United States).

Hammond,G.E. (2017) PFLOTRAN Reaction Sandbox: A Flexible Extensible Framework for Vetting Biogeochemical Reactions within an Open Source Subsurface Simulator.

Hammond,G.E. and Lichtner,P.C. (2010) Field-scale model for the natural attenuation of uranium at the Hanford 300 Area using high-performance computing: MODEL FOR NATURAL ATTENUATION OF URANIUM. *Water Resour. Res.*, **46**.

Henry,C.S. *et al.* (2010) High-throughput generation, optimization and analysis of genome-scale metabolic models. *Nature Biotechnology*, **28**, 977–982.

Kuypers,M.M.M. *et al.* (2018) The microbial nitrogen-cycling network. *Nat Rev Microbiol*, **16**, 263–276.

Liu,G. and Wang,J. (2012) Probing the stoichiometry of the nitrification process using the respirometric approach. *Water Research*, **46**, 5954–5962.

Long,P.E. *et al.* (2016) Microbial Metagenomics Reveals Climate-Relevant Subsurface Biogeochemical Processes. *Trends in Microbiology*, **24**, 600–610.

Mills,R.T. *et al.* (2009) Modeling subsurface reactive flows using leadership-class computing. *J. Phys.: Conf. Ser.*, **180**, 012062.

Raboni,M. *et al.* (2014) Calculating specific denitrification rates in pre-denitrification by assessing the influence of dissolved oxygen, sludge loading and mixed-liquor recycle. *Environmental Technology*, **35**, 2582–2588.

Rittmann,B.E. and McCarty,P.L. (2012) Environmental biotechnology: principles and applications Tata McGraw-Hill Education.

Scheibe,T.D. *et al.* (2009) Coupling a genome-scale metabolic model with a reactive transport model to describe in situ uranium bioremediation. *Microbial Biotechnology*, **2**, 274–286.

Shaffer,M. *et al.* (2020) DRAM for distilling microbial metabolism to automate the curation of microbiome function. *Nucleic Acids Res*, **48**, 8883–8900.

Song,H.-S. *et al.* (2017) Regulation-Structured Dynamic Metabolic Model Provides a Potential Mechanism for Delayed Enzyme Response in Denitrification Process. *Frontiers in Microbiology*, **8**.

Song,H.-S. and Liu,C. (2015) Dynamic Metabolic Modeling of Denitrifying Bacterial Growth: The Cybernetic Approach. *Industrial & Engineering Chemistry Research*, **54**, 10221–10227.

Steefel,Carl I. *et al.* (2015) Micro-Continuum Approaches for Modeling Pore-Scale Geochemical Processes. *Reviews in Mineralogy and Geochemistry*, **80**, 217–246.

Steefel,C. I. *et al.* (2015) Reactive transport codes for subsurface environmental simulation. *Computational Geosciences*, **19**, 445–478.

Tfaily,M.M. *et al.* (2019) Single-throughput Complementary High-resolution Analytical Techniques for Characterizing Complex Natural Organic Matter Mixtures. *JoVE*, 59035.

Triska,F.J. *et al.* (1993) The role of water exchange between a stream channel and its hyporheic zone in nitrogen cycling at the terrestrial—aquatic interface. In, Hillbricht-Ilkowska,A. and Pieczyńska,E. (eds), *Nutrient Dynamics and Retention in Land/Water Ecotones of Lowland, Temperate Lakes and Rivers*, Developments in Hydrobiology. Springer Netherlands, Dordrecht, pp. 167–184.

Villa,J.A. *et al.* (2020) Methane and nitrous oxide porewater concentrations and surface fluxes of a regulated river. *Science of The Total Environment*, **715**, 136920.

Zheng,L. *et al.* (2016) Temperature effects on nitrogen cycling and nitrate removal-production efficiency in bed form-induced hyporheic zones. *Journal of Geophysical Research: Biogeosciences*, **121**, 1086–1103.