Article type: MITOGENOME ANNOUNCEMENT

Title: The complete mitochondrial genome of a parthenogenetic ant *Monomorium triviale*

(Hymenoptera: Formicidae)

Authors: Naoto Idogawa[1*], Chih-Chi Lee[2], Chin-Cheng Scotty Yang[3] and Shigeto Dobata[4]

Affiliation:

[1] Laboratory of Insect Ecology, Graduate School of Agriculture, Kyoto University, Kyoto

606-8502, Japan.

[2] Institute of Evolution, Department of Evolutionary and Environmental Biology, University

of Haifa, Haifa, 31905, Israel.

[3] Department of Entomology, Virginia Polytechnic Institute and State University, Blacksburg,

VA, 24061-0131, USA.

[4] Department of General Systems Studies, Graduate School of Arts and Sciences, The

University of Tokyo, Komaba, Meguro, Tokyo 153-8902, Japan.

*Corresponding author

E-mail: idogawa.naoto.na@alumni.tsukuba.ac.jp

ORCID digit of the author

Naoto Idogawa: 0000-0003-4055-8204

Chih-Chi Lee: 0000-0002-8778-1449

Chin-Cheng Scotty Yang: 0000-0003-0967-5170

Shigeto Dobata: 0000-0003-1586-6758

ABSTRACT

Monomorium is one of the most species-rich yet taxonomically problematic ant genus. An

East Asian species, *M. triviale* Wheeler, W.M., 1906, are reproduced by obligate thelytokous

parthenogenesis and performs strict reproductive division of labor. We sequenced the *M.*

triviale mitogenome using next-generation sequencing methods. The circular mitogenome of

M. triviale was 16,290 bp in length, consisting of 13 protein-coding genes, two ribosomal

RNA genes, 22 transfer RNAs, and a single non-coding region of 568 bp. The base

composition was AT-biased (82%). Gene order rearrangements were detected and likely to be

unique to the genus *Monomorium.* We announce the *M. triviale* mitogenome as additional

genomic resources for elucidating phylogenetic and taxonomic problems of *Monomorium* and

comparative genomics of parthenogenetic ant species.

KEYWORDS: Hymenoptera, Formicidae, thelytoky, *Monomorium*, mitogenome

In the hyperdiverse ant subfamily Myrmicinae, *Monomorium* Mayr, 1855 is one of the

most species-rich genera with over 300 described species including several successful tramp

species such as the flower ant *M. floricola* and *M. salomonis*, and the pharaoh ant *M.*

pharaonis (Pontieri & Linksvayer 2019). However, recent studies (Ward 2015, Sparks 2019)

suggest polyphyly of this genus, and genomic resources are therefore essential for resolving

such a taxonomic issue.

An East Asian species, *M. triviale* Wheeler, W.M., 1906 is particularly of our interest as it

reproduces by thelytokous parthenogenesis where virgin queens produce both workers and

next-generation queens (Gotoh at al. 2012; Idogawa et al. 2021). To date, only partial

mitochondrial DNA sequences have been reported, with all of which being identical among

populations in Japan (Idogawa et al. 2021). Hence, a complete mitochondrial genome of this

species can provide additional information for further analysis. Here, we present the first

complete mitogenome for *M. triviale*.

A colony of *M. triviale* headed by a single queen was collected in Takaragaike Park,

Kyoto, Japan (35.060087 N, 135.788488 E) on Sept. 9, 2017. The queen and her

parthenogenetic offspring produced later in the laboratory (larvae and pupae, approx. 100

individuals) were fixed in 99.5% EtOH. We extracted genomic DNA from the pooled

individuals with DNeasy Blood and Tissue kit (Qiagen, Hilden, Germany). We sequenced the

pooled DNA on the HiSeq X sequencer (Illumina, San Diego, CA) at Macrogen Japan Corp.,

Tokyo. After removing adapters with Trimmomatic v0.39 (Bolger et al. 2014), we conducted

de novo mitogenome assembly based on 4,327,159 paired-end sequence reads using

NOVOPlasty v3.6 (Dierckxsens et al. 2017), with *Solenopsis invicta* mitogenome (NCBI

accession: NC_014672) as a seed. Average read coverage of the mitogenome assembly was

37,840, providing ample depth for correctness. We annotated protein-coding genes (PCGs),

rRNAs and tRNAs using GeSeq (Tillich et al. 2017), MITOS (Bernt et al. 2013), NCBI

ORF-finder (Rombel et al. 2002), and ARWEN (Laslett and Canbäck 2008). The sequence

information was deposited in the DNA Data Bank of Japan under accession number:

LC605004. A specimen was deposited at the Laboratory of Insect Ecology, Graduate School

of Agriculture, Kyoto University (http://www.insecteco.kais.kyoto-u.ac.jp; N. Idogawa; under

the voucher number Mtri_20170909_4).

The complete mitogenome of *M. triviale* was 16,920 bp in length, which is comparable to

those of other ant species. The nucleotide composition was AT-biased (82%). The mitogenome

contained 13 protein-coding genes (PCGs), two rRNAs, and 22 tRNAs, which are typical for

most metazoan animals. All PCGs used ATG or ATT as the start codon and TAA or TAG as

the stop codon. The tRNAs, ranging in size from 59 to 74 bp, were similar to those of other

ants (54 to 90 bp). The control region presumably corresponded to the single largest

non-coding AT-rich region (569 bp, A+T 94%). Gene order of *M. triviale* was identical to that

of a congener *M. pharaonis* (NC_051486.1). Notably, the gene order of the two *Monomorium*

species had two gene rearrangements: an inversion between *trn*P and *ND1* (Myrmicinae,

*ND6-CYTB-trn*S; *Monomorium*, <u>*trn*S</u>-<u>*CYTB*</u>-<u>*ND6*</u>; underlines indicate inverted genes) and

translocations between *ND3* and *trn*F (Myrmicinae common, *trn*A-*trn*R-*trn*N-*trn*S-*trn*E;

Monomorium, *trn*R-*trn*E-*trn*A-*trn*N-*trn*S). This feature was different from the common gene

order of the subfamily Myrmicinae and likely unique to *Monomorium* ants (Babbucci et al.

2014; Park 2020). This may help identify the genus *Monomorium sensu stricto*, in addition to

nucleotide substitutions.

We inferred the phylogenetic relationships of 25 ant species using the concatenated

nucleotide sequences of all 13 PCGs, with the honeybee *Apis mellifera* as an outgroup.

Sequence alignment was constructed using ClustalW (Thompson et al. 2003) implemented in

MEGA-X (Kumar et al. 2018). The GTR + I + G model was determined as a best-fit model by

ModelTest-NG v0.1.6 (Darriba et al. 2020). Both a maximum likelihood tree made by

RAxML-NG v1.0.0 (Kozlov et al. 2019) and Bayesian inference tree made by MrBayes

v3.2.7 (Ronquist et al. 2012) consistently support the current phylogenetic placement of

Monomorium in the subfamily Myrmicinae (Figure 1).

In conclusion, the newly sequenced complete mitochondrial genome of *M. triviale*

provides additional resources for further phylogenetic characterization of the taxonomically

problematic genus *Monomorium* and comparative genomics of parthenogenetic ant species.

Figure 1.

Maximum likelihood (1,000 bootstrap repeats) and Bayesian inference (100,000

generations) trees showing phylogenetic relationships among 25 ant species. A honeybee,

Apis mellifera was used as an outgroup. Phylogenetic tree was drawn based on maximum

likelihood tree. *Monomorium triviale* in bold is the result obtained in this study. The numbers

above branches indicate bootstrap support values for maximum likelihood tree and posterior

probability for Bayesian inference tree, in order.

Data availability statement

The genome sequence data that support the findings of this study are openly available in

DDBJ/GenBank at (https://www.ddbj.nig.ac.jp) under the accession no. LC605004. The

associated BioProject, SRA, and Bio-Sample numbers are PRJDB12079, DRX301164, and

SAMD00394597 respectively.

REFERENCES

1. Babbucci M, Basso A, Scupola A, et al (2014) Is it an ant or a butterfly? Convergent

evolution in the mitochondrial gene order of Hymenoptera and Lepidoptera. Genome

Biol Evol 6:3326–3343. https://doi.org/10.1093/gbe/evu265

2. Bernt M, Donath A, Jühling F, et al (2013) MITOS: Improved de novo metazoan

mitochondrial genome annotation. Mol Phylogenet Evol 69:313–319.

https://doi.org/10.1016/j.ympev.2012.08.023

3. Bolger AM, Lohse M, Usadel B (2014) Trimmomatic: A flexible trimmer for Illumina

sequence data. Bioinformatics 30:2114–2120.

https://doi.org/10.1093/bioinformatics/btu170

4. Darriba Di, Posada D, Kozlov AM, et al (2020) ModelTest-NG: A New and Scalable

Tool for the Selection of DNA and Protein Evolutionary Models. Mol Biol Evol

37:291–294. https://doi.org/10.1093/molbev/msz189

5. Dierckxsens N, Mardulyn P, Smits G (2017) NOVOPlasty: De novo assembly of

organelle genomes from whole genome data. Nucleic Acids Res 45: e18–e18.

https://doi.org/10.1093/nar/gkw955

6. Gotoh A, Billen J, Tsuji K, et al (2012) Histological study of the spermatheca in three

thelytokous parthenogenetic ant species, *Pristomyrmex punctatus*, *Pyramica*

membranifera and *Monomorium triviale* (Hymenoptera: Formicidae). Acta Zool

93:200–207. https://doi.org/10.1111/j.1463-6395.2010.00498.x

7. Idogawa N, Sasaki T, Tsuji K, Dobata S (2021) Comprehensive analysis of male-free

reproduction in *Monomorium triviale*; (Formicidae: Myrmicinae). PLoS ONE 16(4):

e0246710. https://doi.org/10.1371/journal.pone.0246710

8. Kozlov AM, Darriba D, Flouri T, et al (2019) RAxML-NG: A fast, scalable and

133 user-friendly tool for maximum likelihood phylogenetic inference. Bioinformatics

134 35:4453–4455. https://doi.org/10.1093/bioinformatics/btz305

135 9. Kumar S, Stecher G, Li M, et al (2018) MEGA X: Molecular evolutionary genetics

136 analysis across computing platforms. Mol Biol Evol 35:1547–1549.

137 https://doi.org/10.1093/molbev/msy096

138 10. Laslett D, Canbäck B (2008) ARWEN: A program to detect tRNA genes in metazoan

139 mitochondrial nucleotide sequences. Bioinformatics 24:172–175.

140 https://doi.org/10.1093/bioinformatics/btm573

141 11. Park J, Xi H, Park J (2020) The complete mitochondrial genome of *Aphaenogaster*

142 *famelica* (Smith, 1874) (Hymenoptera: Formicidae). Mitochondrial DNA Part B

143 5:492–494. https://doi.org/10.1080/23802359.2019.1698992

144 12. Pontieri L, Linksvayer TA (2019) *Monomorium*. In: Encyclopedia of Social Insects.

145 Springer International Publishing, Cham, pp 1–6.

146 13. Rombel IT, Sykes KF, Rayner S, Johnston SA (2002) ORF-FINDER: A vector for

147 high-throughput gene identification. Gene 282:33–41.

148 https://doi.org/10.1016/S0378-1119(01)00819-8

149 14. Ronquist F, Teslenko M, Van Der Mark P, et al (2012) Mrbayes 3.2: Efficient bayesian

150 phylogenetic inference and model choice across a large model space. Syst Biol

151 61:539–542. https://doi.org/10.1093/sysbio/sys029

152 15. Sparks KS, Andersen AN, Austin AD (2019) A multi-gene phylogeny of Australian

153 *Monomorium* Mayr (Hymenoptera: Formicidae) results in reinterpretation of the genus

154 and resurrection of *Chelaner* Emery. Invertebr Syst 33:225–236.

155 https://doi.org/10.1071/IS16080

156 16. Thompson JD, Gibson TJ, Higgins DG (2003) Multiple Sequence Alignment Using

ClustalW and ClustalX. Curr Protoc Bioinforma 2.3. 1-2.3. 22.

https://doi.org/10.1002/0471250953.bi0203s00

17. Tillich M, Lehwark P, Pellizzer T, et al (2017) GeSeq - Versatile and accurate

annotation of organelle genomes. Nucleic Acids Res 45: W6–W11.

https://doi.org/10.1093/nar/gkx391

18. Ward PS, Brady SG, Fisher BL, Schultz TR (2015) The evolution of myrmicine ants:

Phylogeny and biogeography of a hyperdiverse ant clade (Hymenoptera: Formicidae).

Syst Entomol 40:61–81. https://doi.org/10.1111/syen.12090

Acknowledgments

We are grateful to Kenji Matsuura who allowed us to use his laboratory.

Funding

This work was supported by a Japan Society for the Promotion of Science (JSPS)

Research Fellowship for Young Scientists to NI (19J22242) and a grant from the Secom

Science and Technology Foundation to SD.