1 **Variation in bacterial composition, diversity, and activity across different subglacial**
2 **basal ice types**
3
4

5 **Shawn M. Doyle[a] and Brent C. Christner[b]**
6
7
8 [a]Department of Oceanography, Texas A&M University, College Station, TX, USA.
9 [b]Department of Microbiology and Cell Science, University of Florida, Gainesville, FL, USA.
10
11

12 Running title: **Bacterial community structures in basal ice types**
13
14 Abstract word count: 250
15 Importance word count: 140
16 Article word count: 5485
17

*Corresponding author:
Shawn M. Doyle
Department of Oceanography
Texas A&M University
Eller O&M Bldg, Rm 716D
College Station, TX, 77843-3146
Tel: 979-845-5105
E-mail: shawndoyle@tamu.edu

26
27
28 Competing Interests: The authors declare no competing financial interests.
29

ABSTRACT

Glaciers and ice sheets possess layers of basal ice characterized by high amounts of

entrained debris that can serve as sources of nutrients and organic matter, providing

habitat for microorganisms adapted to the frozen conditions. Basal ice forms through

various mechanisms and is classified based on its ice and debris content; however, little is

known about variation in microbial composition, diversity, and activity across different

basal ice types. We investigated these parameters in four different types of basal ice from a

cold-based and temperate glacier and used a meta-analysis to compare our findings with

microbiome studies from other frozen environments. We found basal ice environments

harbor a diverse range of microbiomes whose composition and activity can vary

significantly between basal ice types, even within adjacent facies from the same glacier. In

some debris-rich basal ices, elevated ATP concentrations, isotopic gas signatures, and high

ratios of amplified sequences for 16S rRNA relative to that for 16S rRNA genes implicated

certain bacterial taxa (e.g., *Paenisporosarcina*, *Desulfocapsa*, *Syntrophus, Desulfosporosinus*)

as being potentially active, with ice temperature appearing to be an important predictor for

the diversity of taxa inferred to be active. Compared to those of other sympagic

environments, these microbiomes often resembled those found in permafrost or perennial

cave ice rather than other glacial ice environments. In contrast, debris-poor basal ices

harbored microbiomes more like those found in oligotrophic englacial ice. Collectively,

these results suggest that different basal ice types contain distinct microbiomes that are

actively structured by the diagenesis of their habitat.

IMPORTANCE

The discharge of material from the basal ice zones of glaciers and ice sheets is an important source of dissolved organic matter, nutrients, and trace metals to downstream environments. Microorganisms with the ability to remain metabolically active while frozen have the potential to alter these substrates before they are released and may thus be a vital biogeochemical component of proglacial environments. However, basal ice environments are among the least explored areas of the cryosphere and information on their microbiology is limited. Here, we examine several basal ice types that were collected from different glaciers to investigate how their microbiomes varied. Our results indicate that sediment-rich basal ice from cold-based and temperate glaciers harbor microbiomes that conduct biogeochemical transformations at subzero temperatures and raises the intriguing possibility that subsurface regions of the cryosphere may be more biogeochemically active than previously thought.

1. INTRODUCTION

At the ice-bed interface of a glacier or ice sheet, basal ice forms when subglacial debris and/or water is incorporated into the base of the ice mass through mechanisms such as freeze-on, regelation, or incorporation of pre-existing ice (1, 2) . Basal ice has a physical structure, chemistry, and microbial composition that is directly affected by its proximity to and interaction with material beneath the glacier (1, 3, 4). When frictional or geothermal heating create temperate conditions at the bed, the melting of basal ice can provide inputs of organic matter, nutrients, oxidants, and microorganisms to the subglacial environment (5). Hence, basal ice likely represents an important vector for dispersing microbes in subglacial aquatic environments that exist within the glacier's flow path (6).

Basal ice layers contain debris entrained from the underlying substrate that ranges in size from fine-grained clays and silts to coarse sands, gravels, and boulders. Basal ice formation can produce layering and stratification patterns that become deformed by the folding and shearing forces associated with glacier flow (7). These processes of entraining and restructuring of subglacial debris combine to produce distinct types of basal ice. This ranges from relatively debris-poor types containing <1% (w/v) sediment to debris-rich types composed almost entirely of sediment with only interstitial ice (1). When compared to the overlying englacial ice (i.e., derived from snowfall), the sediment-rich basal ice contains higher concentrations of potential microbial substrates that include formate, acetate, ferrous iron, and ammonia (8–11). These compounds are hypothesized to be an important source of electron donors for heterotrophic and lithotrophic microorganisms inhabiting subglacial environments, including frozen matrices within the basal ice itself (4, 12).

Although ecosystems associated with sympagic environments were first described

nearly 60 years ago (13), viable microorganisms entrapped within glacial ice have been

assumed to exist in a persistent state of metabolic dormancy (14). This view has been

challenged based on evidence from several pioneering studies that revealed anomalies in

entrapped gas concentrations within englacial (15, 16) and basal ice (17–19). The

metabolic activity of microorganisms in basal ice is now thought to play an important role

in the biogeochemistry of downstream terrestrial, freshwater, and marine environments

that receive seasonal discharges of basal ice and subglacial meltwater (20–23). In fact,

transport of basal material is one of the largest sources of labile dissolved organic carbon

and iron to marine systems receiving outflow from glaciated watersheds (24–26).

The world's ice sheets are estimated to contain ~10.2 Pg of organic carbon (27),

which represents only ~0.5% of the global soil organic carbon pool (28). However, the

sediments underlying the Greenland and Antarctic ice sheets may contain reservoirs of

~500 and ~21,000 Pg of organic carbon, respectively (29, 30). Due to accelerated rates of

melt and glacial retreat in the polar regions, the products of organic matter processing by

subglacial microorganisms may play an increasingly important role in global carbon

cycling. For example, the subglacial production of biogenic methane and its subsequent

release to the atmosphere could represent a positive feedback that accelerates further

climate warming (30). Indeed, basal ice sediments have been shown to contain viable

methanogens (8, 31) as well as methanotrophs (32, 33), indicating the presence of a

subglacial methane cycling. Microbial activities in the basal zones of the world's glaciers

and ice sheets may thus be an overlooked component of Earth's climate oscillations over

geologic timeframes. Despite their potential importance, the logistic challenges associated

with accessing these environments have contributed to the dearth of information on the

microbes and biogeochemical processes they perform in the basal zone of large ice masses.

In this study, we used 16S rRNA gene amplicon sequencing to profile the

composition and structure of microbial assemblages within basal ice types and from

different locations. To explore how these microbial assemblages compare to those in other

perennially frozen environments, we performed a meta-analysis that combined our data

with that from other studies. Many of these studies sequenced different variable regions of

the 16S rRNA gene (e.g., V4, V2, V3V4) that do not fully overlap, hindering the use of

traditional OTU-based approaches. To address this, we used the SATé-enabled phylogenetic

placement (SEPP) technique to insert exact amplicon sequence variants (ASVs) from these

heterogenous variable regions into the same reference phylogeny (34). This enables a

phylogenetically stable comparison of the microbial assemblages inhabiting these different

frozen environments. In contrast to the aeolian-source microbes in englacial ice, the

microorganisms in basal ice appear to represent "communities" in an ecological sense. We

also used RNA- and ATP-based approaches to identify potentially active taxa and regions in

basal ice types and discuss the implications of microbial processes that occur in an

important transient environment of the cryosphere.

2. MATERIALS AND METHODS

2.1 Ice sampling

Taylor Glacier

Taylor Glacier is a cold-based glacier (i.e., has basal ice that is constantly below the

pressure-melting point) that is an outlet of the East Antarctic Ice Sheet, terminating at the

western end of Taylor Valley (McMurdo Dry Valleys, Victoria Land, Antarctica). During the

austral summers of 2007 and 2009, two tunnels were excavated into the basal zone of the

northern lateral margin of the glacier (Fig. S1). Tunnel construction was initiated on fresh

ice aprons and extended ~8 m into the glacier interior. At the end of each tunnel, a vertical

shaft (2007 expedition) or large chamber (2009 expedition) was excavated to expose a

stratigraphic sequence of multiple basal ice types (4). Electric chainsaws equipped with

unlubricated carbide tipped chains were used to exhume samples of the ice sequence from

each basal ice type. Ice samples were sealed in polypropylene flat wrap, packed into

insulated containers, and shipped frozen from the field to Louisiana State University where

they were subsequently stored at -20 °C.

Matanuska Glacier

Matanuska Glacier is a 43 km temperate (i.e., englacial and basal ice constantly at

the pressure melting point), valley glacier located in south-central Alaska, approximately

140 km northeast of Anchorage. In July 2013, a horizon of debris-rich basal ice exposed

near the glacier terminus was sampled using an electric chainsaw (Fig. S2). After collection,

the ice samples were packaged in polypropylene flat wrap and shipped frozen to Louisiana

State University for storage at -20 °C.

2.2 Ice decontamination

Within a temperature-controlled environmental room (-5°C), ~2 kg of ice was cut

from larger basal ice samples using either a band saw (for the sediment-poor basal ice) or a

diamond-bladed masonry saw (sediment-rich basal ice). The surface of freshly cut samples

were then cleaned following previous methods (35). Briefly, the outermost surface was

physically removed by scraping with an autoclaved microtome blade and the newly

exposed ice surface was then washed with chilled (-5°C), 0.22 µm filtered 95% ethanol.

Microtome scraping was omitted when processing sediment-rich samples due to the

presence of coarse granules and stones embedded in the sediment-layers. Following the

ethanol wash, samples were washed extensively with ice-cold, 0.22 µm, twice-autoclaved

deionized water until a minimum of ~5 mm of the outer surface had been removed based

on the ice weight. Large, sterilized surgical forceps were used to handle all ice samples

during decontamination and were exchanged frequently during the procedure to prevent

carryover contamination. Sample weights before and after decontamination were used to

estimate the amount of material removed from each sample during processing, which was

typically between 15% and 25% of the initial mass.

The effectiveness of the decontamination procedure was monitored by coating ice

samples beforehand with a tracer solution that consisted of three components: [1] ~10^8

cells mL^{-1} of *E. coli* JM109 cells transformed with a pETBlue-2 plasmid (Novagen)

containing the gene for alcohol dehydrogenase (ADH) from *Drosophila melanogaster*, [2]

fluorescein (1000 ppm), and [3] 33% (v/v) glycerol to prevent freezing on the ice at -5°C.

The removal of the tracer was monitored after each step of the decontamination procedure

(i.e., scraping, ethanol wash, H_2O wash). The fluorescein fluorophore component was

monitored by quantifying blue-green fluorescence in 200 µL of rinse-water or meltwater

using a BioTurner 20/20n Luminometer (P/N 2030-002) equipped with the blue

fluorescence module (BioTurner 2030-041). The pETBlue-2 plasmid component was

monitored using a PCR with the following conditions: 1 U of Taq DNA polymerase

(5PRIME), 1× MasterTaq buffer, 1× TaqMaster PCR enhancer, 1.5 mM Mg^{2+}, 0.2 µM of each

primer (TF7: 5'-TAATACGACTCACTATAGGG -3'; pETBlue-DOWN: 5'-

GTTAAATTGCTAACGCAGTCA-3'), 200 μM dNTPs, and ~100 pg of template DNA. Thirty

cycles of PCR were done with a 30 s denaturation step at 94 °C, 60 s annealing step at 55 °C

and extension at 72 °C for 60 s, followed by a final extension at 72 °C for 10 min. The

presence of viable *E. coli* was monitored by spread plating rinse-water and meltwater on

agar-solidified LB media containing ampicillin (100 μg mL^{-1}) followed by overnight

incubation at 37 °C. Samples in which any component of the tracer solution was detected in

the final meltwater were discarded and reprocessed from a new subsample.

2.3 ATP analysis

Immediately after melting, samples were filtered through 0.22 μm pore size

polyethersulfone (Supor; Pall) filters using a bench-top vacuum manifold (≤ 20 kPa). Clean

ice samples were filtered through a 25 mm diameter filter while banded, solid, and

dispersed ice samples were processed through either a 47 mm or 90 mm diameter filter,

depending on total sediment content. The filters were cut into small pieces using a sterile

scalpel and extracted by vortexing for 10 s in 5 mL of boiling Tris-acetate buffer (TAB;

50mM Tris base adjusted to pH 7.75 with acetic acid). The mixture was then incubated in a

boiling water bath for 5 minutes and immediately placed on ice for 30 minutes. Following

centrifugation for 5 minutes at 4500×g to remove debris, the supernatant was collected.

The concentration of extracted ATP was measured using a modification of the firefly

luciferase-luciferin assay described by (36). Briefly, 10 μL of extract was added to an equal

volume of triethanolamine buffer (TEOA buffer; 200 mM triethanolamine pH 7.6, 2 mM

MgCl2, 240 mM KCL) and incubated at 37°C for 10 minutes. To correct for adenylate

adsorption to and luciferase inhibition by the sediment in the sample extracts, two samples

were prepared in parallel for each measurement. Following incubation at 37°C, 10 uL of

TAB supplemented with 10 µM ATP was added to one of the samples, providing an internal

standard for each measurement. All samples were kept on ice during preparation and were

equilibrated to room temperature (22 °C) prior to the addition of luciferase and

measurement of luminescence.

Weighted linear regression was used to generate calibration curves using

concentrations of ATP (in TAB) from 1 pM to 10 µM. The luciferase-luciferin cocktail was

prepared fresh for each series of measurements and consisted of 100 U ml^{-1} luciferase and

0.140 mM D-luciferin. Luminescence was quantified with a 20/20n luminometer

(Promega) using auto-injection of 100 µL of the luciferase-luciferin cocktail. Relative

luminescence units (RLU) were integrated for three seconds immediately after injection.

Sample RLU values were corrected for adsorption and inhibition using the following

formula:

$$\frac{1}{\left(\dfrac{RLU_{\text{sample+std}} - RLU_{\text{sample}}}{RLU_{\text{blank+std}}}\right)} \times RLU_{\text{sample}} = RLU_{\text{corrected}}$$

The corrected RLU values were used together with the standard calibration curve to

determine the ATP concentration in the extractant. Final ATP concentrations were

corrected for dilution and divided by the mass of the ice sampled. All values reported are

the means of replicate samples (n = 3 to 5).

2.4 Nucleic acid extraction

Decontaminated basal ice samples were placed in sterilized containers and melted

at 4°C. For samples from sediment-rich basal ice (i.e., banded, solid, and dispersed ice), the

resulting meltwater was shaken to resuspend any sediment and DNA was extracted from

the resulting meltwater-sediment slurry using MoBio PowerMax DNA Isolation kits. To

increase DNA extraction yields, two 10g extractions were combined onto a single silica spin

column for each sample. For samples from sediment-poor basal ice (i.e., clean ice),

meltwater was filtered onto 47 mm, 0.22 μm pore size Supor PES filter membranes (Pall)

with ≤20 cm Hg of vacuum. DNA was extracted from the filters using MoBio PowerWater

DNA Isolation kits as per the manufacturer's instructions with one exception: the

mechanical lysis step was performed on a BioSpec MiniBeadbeater-8 for 2 minutes at

maximum speed. All DNA samples were stored at -80°C until PCR amplification.

Due to the low cell abundances in the basal ice (≤10^4 cells g^{-1}; (3)), very large

samples (~5 kg of ice; Table S1A) were processed to recover sufficient RNA for reverse

transcription PCR (RT-PCR). To facilitate filtration of the large sample sizes, coarse

sediment was removed from sediment-rich basal ice samples after melting using low speed

centrifugation (700×g, 10 min, 4°C). The resulting supernatant containing fine clay and silt-

like particles was then concentrated onto 90 mm, 0.22 μm pore size Supor PES filter

membranes. Meltwater from clean ice samples did not require this centrifugation step and

were directly filtered onto 90mm, 0.22 μm pore size Supor PES filters.

After filtration, the filters were immediately processed for RNA extraction using a

modified phenol-chloroform extraction protocol (32). Briefly, filters were sliced into small

pieces using a sterile scalpel, immersed in 3 mL of TE buffer (1 mM EDTA, 10 mM Tris; pH

6.3) containing lysozyme (15 mg mL^{-1}), and vortexed for 30 minutes at room temperature.

Samples were then homogenized in a BioSpec Mini-Beadbeater for 2 minutes at maximum

speed using ~1 g of sterilized 0.1 mm diameter zirconia/silica beads. After bead beating,

crude extracts were amended with two volumes of chilled (4°C) denaturing buffer (4M

guanidine thiocyanate, 50 mM Tris, 10 mM EDTA, 1% w/v N-lauroylsarcosine, 1% β-

mercaptoethanol), the resulting insoluble material was pelleted via centrifugation (4500×g

for 5 min at 4°C), and the supernatant was collected. The pellet was washed with 3 mL of

chilled denaturing buffer, centrifuged again, and the resulting supernatant pooled with the

first. This pooled lysate was then extracted with an equal volume of

phenol:chloroform:isoamyl alcohol (25:24:1, pH 6.6), followed by a second extraction with

chloroform:isoamyl alcohol (24:1). Nucleic acids were purified from these extracts via an

overnight ethanol precipitation with 0.3 M sodium acetate (pH 6.0). Linear acrylamide (20

μg mL^{-1}; Ambion) was used as a coprecipitant to aid recovery. Genomic DNA was

eliminated from RNA extracts by digesting for one hour at 37°C with 4 U of TURBO DNAase

(Ambion) followed by purification with MEGAclear Transcription Clean-Up kits (Ambion).

Extracted DNA and RNA concentrations were measured using Quant-it Picogreen

and Ribogreen kits (Life Technologies), respectively, per the manufacturer's instructions.

2.5 16S rRNA amplicon sequencing

From the extracted RNA, complementary DNA (cDNA) was reverse transcribed from

~1 ng of total RNA using SuperScript II reverse transcriptase with the 806R primer (5'-

GGACTACVSGGGTATCTAAT-3') following the manufacturer's protocol. Controls lacking

RNA template or reverse transcriptase were analyzed to monitor for contaminating RNA

and the persistence of genomic DNA, respectively, in the samples.

The hyper-variable V4 region of the 16S rRNA gene was PCR amplified from the

DNA extracts and cDNA libraries using a barcoded 515F-806R primer pair (37). Each 50 μL

PCR contained ~100 pg of DNA or cDNA, 2.5 U of AmpliTaq Gold DNA polymerase LD

(Invitrogen), 1× Gold Buffer, 2.5 mM MgCl$_2$, 0.2 μM of each primer, and 0.2 μM dNTPs.

Amplification conditions included at 95°C for 9 min, followed by 30 to 40 cycles of

denaturation at 94°C for 1 min, annealing at 50°C for 30 s, and extension at 72°C for 30 s. A

final extension was performed at 72°C for 10 min.

Following amplification, amplicons were separated by electrophoresis on a 2%

agarose gel to assess relative band intensity and size. The amplicons were quantified with

Quant-it fluorometry and pooled at equimolar concentrations. Two extraction blanks were

included in the pooled libraries to serve as procedural controls (38, 39). Pooled libraries

were purified with a MoBio UltraClean PCR Cleanup Kit and sequenced on the Illumina

MiSeq platform (v2 chemistry, 2× 250 bp; Georgia Genomics Facility).

2.6 16S rRNA amplicon analysis

Sequence read curation and processing was performed using DADA2 (40) with the

following filtering parameters (maxN=0, trunQ=2, rm.phix=TRUE, maxEE=2, R1

truncLen=240, R2 truncLen=200). Error rates for the filtered and trimmed R1 and R2 reads

were calculated using the *learnErrors* function and subsequently used to denoise reads

using the DADA2 sample inference algorithm. The denoised reads were merged into

amplicon sequence variants (ASV) using a global ends-free alignment. Paired reads

containing mismatches in the overlapping region were removed from the dataset. Chimeric

ASVs were identified and removed by using the consensus method within the

removeBimeraDenovo function. A consensus taxonomy for each ASV was then assigned

using a naïve Bayesian classifier (41) trained on release 132 of the SILVA reference

database (42). The dataset was then subsampled to an even read depth across all samples

(59,562 sequences). Any ASV classified as a chloroplast, mitochondria, or from which

≥0.1% of its reads were from a procedural blank were removed. ASVs observed exclusively

in the RNA libraries were flagged as a potential contaminant and removed from the

analysis.

2.7 Meta-analyses of microbiomes of perennially frozen ecosystems

Publicly available datasets on the NCBI Sequence Read Archive (SRA) from

perennially freshwater frozen environments were identified using a search on August 8th,

2021 with the following syntax: ice metagenome[Organism], glacier

metagenome[Organism], and permafrost metagenome[Organism]. Run tables from

unambiguously labeled 16S rRNA amplicon datasets were aggregated and studies based on

Sanger sequencing were excluded; only datasets based on Illumina, 454, or IonTorrent

sequencing were considered in the analysis. Because many studies contained samples from

laboratory experiments or field manipulations (e.g., nutrient amendment, thawing, etc.),

each dataset was manually curated and cross-referenced with a publication or other data

source so that only high quality 16S rRNA surveys from genuine, unmodified permanently

frozen environments were included. As part of this effort, datasets were filtered to remove

samples from environments that were only seasonally frozen. For example, samples of an

active soil layer or transition zone from permafrost datasets were excluded. Additionally,

because of their marine origin, we also choose to exclude the wealth of sea ice sample data

available and focus the analysis on terrestrial icy environments. Unfortunately, several

datasets also had to be excluded because their sequence reads on the SRA were not

demultiplexed and barcode information was not provided in the linked metadata or

publication. A list of datasets that met these criteria and were included in the meta-analysis

are detailed in Table S1B. All runs were downloaded using the SRA Toolbox *fastq-dump*

function.

Each dataset was individually processed using DADA2 to infer ASV sequences and

produce an ASV table. For each study, the filtering parameters were adjusted to account for

variation in fragment length and the 16S rRNA variable region sequenced. Due to higher

rates of homopolymer and indel errors with 454 and IonTorrent sequencing, datasets using

these platforms used a band size parameter of 32 and homopolymer region (≥ 3 repeated

bases) gap cost of -1 during the sample inference algorithm. For Illumina datasets, the band

size parameter was set to 16 and gaps in homopolymer regions were treated as normal

gaps (gap cost of -8). Nine datasets had to be removed from the meta-analysis after these

steps due to the loss of too many sequences during quality control steps within the DADA2

pipeline. Complete parameter details and throughput statistics for each dataset in the final

meta-analysis are available in Table S1C and S1D. In datasets containing clearly labeled

replicates, read counts from replicate samples were combined and averaged before

downstream analyses.

ASV sequences from each study were merged and inserted into the 99% Greengenes

reference tree (43) using the SATé-enabled phylogenetic placement (SEPP) technique (44)

within the QIIME2 *fragment-insertion* plugin (34). The merged ASV tables from each study

were filtered to remove ASVs rejected from tree insertion by SEPP. ASVs which could not

be taxonomically classified beyond the domain rank (n=2434) or were classified as

mitochondrial sequences (n=119) were flagged as potential artifacts and removed.

The statistical significance of hierarchical clustering patterns between samples in

the meta-analysis was performed using Monte Carlo simulations of pairwise unifrac

distances as implemented in the *sigclust2* R package (45). Because parallelized, matrix-

based unifrac calculators are incompatible with *sigclust2*, we used the serial, pairwise

unifrac calculator from the *scikit-bio* python package. This calculator was embedded into R

for the *sigclust2* Monte Carlo simulations using the *reticulate* R package.

2.8 Data availability

Sequence read data generated in this study is available on NCBI SRA under

BioProject PRJNA282540.

3. RESULTS

3.1 Description of Basal Ice Samples

Based on the nomenclature of (46), we identified three distinct types of basal ice in

the basal ice profiles sampled from Taylor Glacier: clean ice, banded ice, and solid ice

(Table 1; Fig. S3; (4)). All three types were collected during the 2007 expedition, while only

clean ice and banded ice horizons were targeted during the 2009 expedition. The basal ice

recovered from Matanuska Glacier was identified as dispersed ice and was the only type

observed at the location of sampling.

3.2 Biomass in sediment-rich and sediment-poor basal ice

The quantity of DNA and RNA extracted from the basal ices was divided by the

sample mass to assess the relative amounts of microbial biomass present within each basal

ice type. Approximately 10,000-fold more DNA per gram of basal ice was extractable from

the sediment-rich basal ice types (i.e., banded, solid, and dispersed) than in the sediment-

poor clean ice (Table 2). Assuming a bacterial genome weight of 2.5 fg, the concentration of

DNA extracted from the clean Taylor Glacier basal ice corresponds to 560 genomes per

gram of ice, which agrees well with direct cell counts of samples from this ice facies (Table

2; (3, 4)). In contrast, genome abundances inferred from DNA concentrations in the

sediment-rich Taylor Glacier basal ice samples were two to three orders of magnitude

higher than direct cells counts. Among the sediment-rich basal ice types, the temperate

dispersed basal ice from Matanuska Glacier contained approximately 6-fold more DNA per

gram of ice than the banded and solid basal ices from the cold-based Taylor Glacier (-15°C).

The concentration of extractable RNA was very low in all basal ice types (between 0.7 to

3.2 pg RNA g^{-1}; Table 2). The highest RNA concentration—though only marginally—was

observed in Matanuska Glacier's dispersed basal ice. These RNA concentrations were

consistent with ATP concentration data, a proxy for viable microbial biomass, and in

general, high concentrations of ATP g^{-1} ice were observed in sediment-rich basal ice types

versus those with low sediment content.

3.3 Composition of microbial assemblages in different types of basal ice

A total of 4,227,678 paired-end reads with an average read length of 253 bp were

obtained from MiSeq sequencing. After filtration and denoising, 2,883,221 non-chimeric

sequences representing 3,310 ASVs remained. Subsampling and culling of ASVs classified

as plastids or flagged as potential contaminants produced a final, curated dataset composed

of 2,005 ASVs.

Based on average-neighbor clustering of Bray-Curtis distances, the basal ice samples

clustered into three significantly different (AMOVA; F[2,3]=5.23, *p*=0.017) groups: (1)

Taylor Glacier sediment-poor basal ice, (2) Taylor Glacier sediment-rich basal ice, and (3)

Matanuska Glacier dispersed ice. The sediment-poor clean ice samples contained microbial

assemblages overwhelmingly composed of *Firmicutes* (68%) and *Proteobacteria* (25%),

with smaller proportions of *Bacteroidetes* (5%) and *Actinobacteria* (1%). The genus

Tumebacillus was especially abundant in the clean ice samples (Fig. 1A), representing more

than half of all taxa in these samples. Assemblages within the banded and solid ice samples

were comparatively more diverse than those found in clean ice (Fig. 2), with a wider range

of abundant phyla represented: *Actinobacteria* (26%), *Firmicutes* (23%), *Bacteroidetes*

(19%), *Proteobacteria* (11%), *Atribacteria* (9%), *Tenericutes* (7%), and *Planctomycetes*

(1%). Notably, ASVs related to the class *Clostridia* composed about 20% of the total in

these samples. The microbial assemblage within Matanuska's dispersed basal ice was

distinct from those observed from Taylor Glacier. Here we observed significant amounts of

Nitrospirae (39%) and *Betaproteobacteria* (30%), of which several of the most abundant

ASVs were related to sulfur and/or iron cycling members of the genera

Thermodesulfovibrio, *Rhodoferax*, and *Thiobacillus*. Comparing assemblages from all basal

ice samples, only 21 taxa (~3%) were shared between all four basal ice types (Table S1E).

These shared taxa were generally rare members with median relative abundances ranging

between 0.03% and 0.31%.

3.4 Assessing metabolic status of bacterial taxa within different types of basal ice

Of the 2,005 ASVs observed in this study, we detected 141 (7% of total) within the

RNA-derived 16S rRNA libraries (Table S1F), suggesting they were recovered from

potentially intact cells rather than eDNA or dead cells. From this group, 80 of the ASVs had

RNA/DNA abundance log-ratios >1 and likely represent metabolically active taxa within

the basal ice. The vast majority (63 ASVs) of these high-ratio ASVs were observed in the

dispersed ice from Matanuska Glacier (Fig. 1B), the warmest and youngest basal ice

sampled in this study. Many of the Matanuska Glacier ASVs were members of the

Chloroflexi and *Deltaproteobacteria,* with some of the largest RNA/DNA abundance log-

ratios belonging to *Desulfocapsa*, *Syntrophus*, and members of the family *Anaerolineaceae.*

Within the banded basal ice from Taylor Glacier, over half of ASVs detected in the RNA-

derived libraries were members of the *Firmicutes*. Among this phylum, members of the

Paenisporosarcina, *Paenibacillaceae*, *Desulfosporosinus*, *Caldicoprobacter*, and *Virgibacillus*

had the highest activity potential based on RNA/DNA abundance log-ratios (Table S1F) and

represented >90% of the RNA reads in these samples. In comparison, the clean and solid

basal ice samples contained the least number of potentially active members—10 ASVs from

only 472 RNA total reads, only 2 of which had RNA/DNA abundance log-ratios >1.

3.5 Meta-Analysis: Microbial populations of various perennially frozen environments

For the meta-analysis, we analyzed 54.6 million 16S rRNA sequences across 256

individual samples from 24 publicly available datasets that represent five different types of

freshwater ice environments: (1) supraglacial ice, (2) englacial ice, (3) basal ice, (4)

perennial cave ice, and (6) permafrost. After sequence filtering and curation, 7 samples

were excluded from meta-analysis due to low read depth. The final meta-analysis dataset

contained 25.7 million non-chimeric sequences representing 54,170 unique ASVs.

Subsampling to equal read depth reduced this to 35,357 ASVs.

We calculated Shannon diversity indices to compare microbial alpha diversity

between these frozen environments. Within the basal ice assemblages characterized in this

study, alpha diversity was significantly higher in the sediment-rich basal ice types than the

low sediment clean ices (t(4)=3.34, p=0.03) (Fig. 2). Compared to the other frozen

environments in the meta-analysis, alpha diversity within Antarctic sediment-rich basal ice

samples was most like that observed in basal ice from various coastal glaciers in Svalbard

(Tukey HSD mean difference = 0.07, p=1) (47). The lower alpha diversity values within

Taylor Glacier's clean ice were most like those found in englacial ice samples from the

GISP2D, NEEM, Guliya Ice Cap, and WAIS Divide ice cores (Tukey HSD mean difference =

0.39, p=1) (48–51).

To parameterize microbial β-diversity between samples, we used hierarchical

cluster analyses of both unweighted (UF) and weighted UniFrac (wUF) distances (Fig. 3 and

Fig. S4). In the UF cluster analysis (Fig. 3), microbial communities largely clustered by

environment type, indicating presence/absence data alone can reasonably discriminate

between different parts of the cryosphere. This suggests microbial community composition

in permanently frozen environments is strongly influenced by founder effects, possibly due

to a population bottleneck effect occurring upon freezing. In the cluster analysis of wUF,

the overall cluster pattern was comparable but many of the branching patterns between

samples were more statistically significant, indicating the inclusion of abundance data can

help discriminate samples from the same type of environment.

Within the cluster analyses, we also found some samples displayed larger variation

and overlap in community structures than others. For example, microbial assemblages

within englacial ice were the most similar (UF: 0.47 ± 0.15; mean ± s.d; n = 42). In contrast,

perennial cave ice samples (52, 53) had the highest heterogeneity among samples (UF: 0.85

± 0.59; n=14). β-diversity of permafrost microbiomes were intermediate of these

endmembers, with relatively tight clustering observed in permafrost from the Storflaket

peat bog (northern Sweden) but wider variation in permafrost samples from University

Valley (Antarctica) (54, 55). The microbial communities within Taylor Glacier's banded and

solid basal ice facies clustered closely with perennial cave ices located in both Romania (52,

53) and California's Lava Bed National Monument (56). Those within the clean basal ice

were instead most similar to assemblages observed in englacial ice from the NEEM and

GISP2D ice cores from Greenland (48, 49)(Fig. 3).

We used an RDA model to determine how microbial lineages were distributed

across the different types of frozen environments (Fig. 4, left). The final RDA model

explained 50.7% of the variability between samples and effectively separated the samples

into three clusters, echoing patterns observed in the unifrac cluster analysis (Fig. 3). Of the

1627 microbial lineages included in the analysis, we identified 36 with loading vectors

greater than the equilibrium contribution (i.e., the proportion of variance that would be

explained by a random constrained axis), indicating they were associated with one or more

of the five types of frozen environments. Scalar projections of these lineages onto the

centroid factor of each type of frozen environment allowed us to estimate how strongly

these lineages were associated with each environment (Fig. 4, right). Unsurprisingly, many

supraglacial ice environments harbored comparatively higher proportions of

Cyanobacteria and plastid-harboring taxa than the other frozen environments. Englacial ice

environments were enriched with members of the genera *Pseudomonas*, *Methylobacterium*,

Acinetobacter, and *Polaromonas*, *Massilia,* and *Janthinobacterium*. In contrast, basal ice,

permafrost, and perennial cave ice environments all contained higher membership of

Firmicutes and *Actinobacteria* related taxa such as *Clostridium*, *Desulfosporosinus*,

Oryzihumus, and *Cryobacterium*.

3.6 Meta-Analysis of early studies based on 16S rRNA clone sequencing

Although 16S rRNA clone libraries lack the read depth to include in our primary

meta-analysis, we performed a secondary analysis to compare how many of the microbial

taxa observed in several early cryosphere microbiome studies (11, 57–60) compared to the

'indicator' lineages we identified in Fig. 4. To achieve this, we inserted full-length clone

sequences into the meta-analysis reference tree using SEPP and taxonomically classified

them using Wang's Bayesian inference method with the Silva v.132 database as a reference.

Clone sequences were then agglomerated together with meta-analysis ASVs belonging to

the same genera.

Of the 236 16S rRNA clone sequences included in this secondary analysis, all but

three belonged to a taxon that had been observed in the meta-analysis. Despite their

extremely low read depth (between 11 and 136 clones), these five studies clustered with

similar environments in a distance-based RDA ordination (Fig. 5). Remarkably, of the 36

indicator taxa we identified in the primary meta-analysis, 12 were represented by at least

one clone sequence among these five studies. Many of these were among the most

abundant sequences observed in these clone libraries (Fig. S5).

4. DISCUSSION

4.1 Microbial communities vary in size and composition across basal ice types

Of the four different types of basal ice analyzed in this study, we found that those

containing large quantities of sediment or debris contained higher quantities of extractable

nucleic acids and microbial abundances than those containing low sediment content. This

is consistent with previous microbiological investigations of the GISP2 ice core that found

higher cell concentrations in the deepest sections of the core, where clay particles were

embedded in the basal ice (10). In addition to higher numbers of total cells and biomass,

microbial diversity is also notably higher in the sediment-rich basal ice facies relative to

clean basal ice or englacial ice, implicating subglacial debris as the primary source of

microorganisms in the basal ice.

Our estimates of bacterial cell abundance based on the yield of extracted DNA ($\sim 10^6$

cells g^{-1} ice) were approximately 100-fold higher than those based on direct microscopic

counts ($\sim 10^4$ cells g^{-1} ice) in the sediment-rich banded and solid basal ice. However, in the

clean basal ice samples, these estimates were highly comparable (Table 2). This difference

likely reflects the much greater efficiency of DNA extraction versus that of cell extraction. It

may also be related to the increased prevalence of eukaryotic genomes or environmental

DNA that is associated with sediment-rich basal ices (47, 61, 62). Alternatively, Taylor

Glacier's debris-rich basal ice may contain significant amounts of necromass. This would be

consistent with previous studies that have found some deep, debris-rich horizons of basal

ice, such as those of the Greenland ice sheet, contain ancient DNA from buried organisms

(63).

We found microbial assemblages within basal ices of the cold-based (-15°C) Taylor

Glacier to contain high abundances of *Firmicutes* and *Actinobacteria* relative to the other

frozen habitats examined*.* These Gram-positive phyla are commonly observed as the

dominant taxa in a wide range of frozen environments including ground ice, permafrost,

polar saline springs, cryopegs, and sea ice (60, 64–69). One possible explanation for these

high abundances is that the physiochemical stresses faced by cells under frozen conditions

(70) favor the prevalence of microorganisms that can form highly resistant endospores

(*Firmicutes*) or spores (*Actinobacteria*). Indeed, the ability to enter a metabolically dormant

state and form an environmentally resistant spore likely represents a robust survival

strategy for microorganisms that become entrapped inside ice for extended timeframes

(71). However, multiple lines of evidence indicate this hypothesis does not adequately

explain the abundance of these spore-forming taxa in our basal ice samples. First, ASVs

related to known endospore-forming genera such as *Paenisporosarcina*, *Virgibacillus*,

Clostridium, and *Desulfosporosinus* were abundant in all our RNA libraries. Because the

rRNA content of endospores decreases relatively quickly after sporulation—on the order of

days to weeks (72, 73)—their detection in >10,000-year-old basal ice implies these taxa

existed as vegetative cells and not as endospores within the ice. Second, populations of

Paenisporosarcina sp. isolated from the Taylor Glacier banded ice do not sporulate after

being refrozen in basal meltwater at in situ temperatures (-15°C), but instead continuously

incorporate radiolabeled DNA and protein precursors into macromolecules(3). Third, we

found pasteurization of banded basal ice meltwater dramatically reduced isolate

cultivability, indicating isolates were not being recovered from heat-resistant endospores.

Finally, 16S rRNA surveys of permafrost that have selectively depleted DNA from dead or

vegetative cells have found many endospore-forming taxa, especially members of the

Clostridia, were more likely to persist as vegetative cells rather than endospores in the

permafrost (74). Collectively, these results suggest the high abundances of *Firmicutes* and

Actinobacteria in Taylor Glacier's basal ice may not only be related to their environmental

durability while dormant, but also due to properties associated with their vegetative cycle

that promote physiological activities within frozen matrices (3).

4.2 Young and warm versus old and cold basal ice

Due to the chemical and biological lability of RNA, paired RNA/DNA 16S rRNA

amplicon libraries provide a useful proxy for discerning viable and metabolically active

microorganisms in a community from those that are dead or dormant. However, there are

limitations to this analysis and over interpretation of ratios as reliable indicators of relative

metabolic activity of different ASVs should be viewed with caution (75). This may

especially be the case in frozen environments, where RNA may be more stable in dormant

cells. As such, we focus primarily on the number of detected taxa in each basal ice type,

especially those with high RNA/DNA abundance log-ratios, which are difficult to explain in

ancient ice samples without invoking metabolic activity.

On average, only about 2% of the total ASVs in each sample were also detected in

the paired RNA library. This low proportion of overlap suggests that only a small fraction of

the microorganisms preserved in the basal ice are potentially metabolically active within

the ice matrix. The dispersed basal ice from Matanuska Glacier contained a much larger

number of high-ratio ASVs than any of the basal ice facies recovered from Taylor Glacier

(Fig. 1). One plausible explanation for this observation is the substantial difference in

underlying geology, age, and temperature of basal ice between the two glaciers.

Assuming similar starting inputs of microbes, older ice would be expected to harbor

smaller numbers of potentially active populations than younger ice due to microbial

attrition over time (70). Analysis of the δD and δ^{18}O composition of meltwater from Taylor

Glacier surface ice samples estimate ages of 11,500 to 65,000 years BP (76), while recent

radiometric ^{81}Kr dating efforts have yielded age estimates near the glacial terminus of

123,500 years BP (77). In comparison, measurements of oxygen isotopes in Matanuska

Glacier's dispersed basal ice layer indicated the basal ice originated in the accumulation

area (78). Based on the velocity of Matanuska Glacier (~110 m yr^{-1}), this would represent

an approximate time of 250 years between deposition and terminal ablation. Trace

amounts of anthropogenic tritium produced by atmospheric thermonuclear weapon tests

in the 1950s and 1960s have also been detected in Matanuska Glacier's basal ice zone (79),

indicating portions of this ice were formed even more recently by freeze-on of surface-

derived meltwater (80). This indicates the dispersed ice we sampled from Matanuska is

substantially younger than any of the basal ice types collected from Taylor Glacier, with the

latter probably being between ~100- to 500-times older than the former.

In addition to age of the basal ice, the large difference in ice temperature between a

temperate and cold-based glacier would also affect the potential for metabolic activity in

the ice habitat (19, 81). Taylor Glacier is a cold-based glacier with a basal ice temperature

of -15°C (4). In contrast, the Matanuska Glacier is temperate, and the basal ice zone is at the

pressure melting point of 0°C (82). As such, cells present in the liquid habitat at grain

boundaries (81) in Taylor Glacier's basal ice are under considerably higher physiochemical

stress than ice near the melting point. For example, the predicted ionic strength of the

briny liquid at ice crystal boundaries—where the microorganisms are located— is nearly

4.6M at -15°C. Comparatively, this is 11.5-fold higher than estimates for unfrozen water in

glacial ice at -1°C (70). Hence, the wider diversity and increased number of ASVs with a

high potential for metabolic activity in the dispersed basal ice may reflect not only the

sample age, but the more favorable conditions for metabolism in temperate versus cold-

based glacial ice matrices.

4.3 Debris-rich basal ice types are hot spots for microbial activity

Multiple studies have reported observations of very low concentrations of oxygen

(17) and large excesses of CH_4 in debris-rich basal ice from the Greenland and Antarctica

ice sheets (10, 30, 83–86). Mirroring these findings, an analysis of gasses entrapped within

Taylor Glacier banded basal ice found O_2 was depleted to as low as 4% in horizons where

CO_2 concentrations were concurrently enriched as high as 20,000-fold relative to

atmospheric concentrations (4). Isotopic analysis of $\delta^{13}C-CO_2$ revealed this CO_2 was

isotopically depleted (-24‰), and therefore, was likely produced by microbial respiration

of organic matter (87). In this study, among all the RNA-based 16S rRNA libraries we

sequenced from Taylor Glacier basal ice, we found the largest reads counts and largest

number of potentially active ASVs in these same banded basal ice horizons. Together with

elevated ATP concentrations, these observations collectively suggest Taylor Glacier's

banded basal ice horizons harbor metabolically active microbiomes that alter entrapped

gas compositions by actively metabolizing organic matter to CO_2 at -15°C within the ice

matrix.

Alternatively, these trends in gas data could be the result of microbial activity that

had occurred before the basal ice formed. One way to explore this possibility is by

considering the total gas content of the ice. If the gases were dissolved in the water that

froze onto the base of the glacier, then total gas volumes as low as 0.02 to 0.08 cc/g would

be expected (88, 89). In solid facies of the Taylor Glacier basal ice profile, the lower total

gas volumes observed were indeed low (0.01 to 0.07 cc/g; (4)). Hence, although we did not

see much evidence for microbial activity in the solid basal ice, we cannot exclude largescale

melting/refreezing processes occurred during its formation. In contrast, in the banded

basal ice—where isotopic, ATP, and RNA data provided strong evidence for microbial

activity—the total gas volumes were consistent with those observed in the clean ice (0.09

to 0.11 cc/g; (4)). These volumes are more similar to those typically seen in englacial ice of

meteoric origin (0.1 cc/g; (90)). In summary, large-scale melting/refreezing cannot explain

the gas data observed across all facies, and in the banded ice, in situ microbial respiration is

the only explanation consistent with its total gas content.

This metabolic activity may simply be a consequence of substrate availability for

microbial consumption. Previous geochemical analysis of Taylor Glacier's basal ice found

dissolved organic carbon (DOC) concentrations are much higher in the debris-rich basal ice

types ($\sim$63 mg L^{-1}) than in the debris-poor types ($\sim$0.8 mg L^{-1}; (4, 87)) However, our

comparison of multiple basal ice types suggests there are likely additional, unidentified

factors besides sediment content controlling whether basal ice harbors an active

microbiome. For example, although the banded and solid basal ice samples harbored very

similar microbial communities in our DNA-based libraries, we found very little evidence for

active ASVs within the solid basal ice. This suggests—at least in polar glaciers containing

basal ice horizons well below 0°C—elevated debris and/or DOC are not alone sufficient for

supporting metabolic activity in the ice.

In many of the debris-rich basal ice types we investigated in this study, we observed

an enrichment of putatively anaerobic taxa (e.g., *Clostridia*) within the same samples where

O_2 concentrations were depleted relative to atmospheric values. One possible explanation

is that the subglacial debris within these horizons were enriched with these taxa before

being entrained into the basal ice. Alternatively, these patterns represent ecological shifts

in microbial community composition that occur inside the ice matrix as oxygen becomes

depleted. However, there is an important nuance to how these shifts probably occur.

Although there is ample evidence that some microbial taxa can remain metabolically active

while frozen (3, 91–94), there is comparatively little concrete evidence that

microorganisms can physically grow and divide while frozen inside ice. This suggests that

reproduction does not occur, and if it does, is imperceptibly slow under the conditions

within an ice matrix. Hence, the increased relative abundance of many putative anaerobic

taxa that we observed in many of the sediment rich basal ice samples likely represents a

decline of strictly aerobic taxa after oxygen has become depleted rather than a genuine

enrichment or outgrowth of putative anaerobes. This hypothesis may explain why

enrichment cultures inoculated with 100,000-year-old Greenland basal ice became turbid

substantially faster (2 to 3 weeks versus several months) when incubated anaerobically

versus aerobically (95).

4.4. Basal ice diagenesis likely controls microbiome composition and structure

The microbiomes of cold-based, debris-rich basal ice horizons such as the banded

and solid basal ice from Taylor Glacier were phylogenetically most like those in perennial

cave ice or permafrost. These similarities are likely a reflection of the comparable

physiochemical conditions found within these environments. All three are characterized by

relatively high concentrations of debris, constant subzero temperatures, and perhaps—as

suggested by the co-enrichment of several putative anaerobic taxa (e.g., *Clostridium*,

Desulfosporosinus) in some samples—depleted or decreased oxygen concentrations. In

contrast, clean basal ice appears to harbor microbiomes most like those found in englacial

ice. This is consistent with the notion that clean basal ice and englacial ice are

diagenetically very similar. Both contain very little to no debris and are typically formed

from the accumulation and compression of snow into glacial ice. In fact, except in unique

cases where new clean basal ice is formed by the accretion of subglacial lake water onto the

base of a glacier, the only real difference between clean basal and englacial ice may be

location. In other words, in many cases clean basal ice is simply deep englacial ice.

Collectively, these findings suggest the microbiomes of basal ice environments are not

simply preserved communities but rather are actively structured by the diagenesis and

physiochemical nature of their habitat.

4.5 Meta-analysis limitations

Comparing 16S rRNA amplicon datasets from different studies is hindered by

inconsistencies in the sequencing platform used and the variable region of the 16S rRNA

gene that was amplified for sequencing. Using SEPP, heterogeneous ASVs from different

studies are inserted into the same reference tree. This creates stable phylogenetic

placements that enable integration and meta-analysis of amplicon data from multiple

variable regions of the 16S rRNA gene (34). Nevertheless, it is important to note the

limitations of this approach. First, integrating previously published 16S rRNA amplicon

datasets based on DGGE profiling remains difficult as they lack sufficient read depth or

taxonomic resolution to make meaningful comparisons with more modern datasets. As a

result, our meta-analysis excluded many of the pioneering works that investigated the

microbial ecology of glacial ice. Second, there are potential biases in our meta-analysis due

to how we classified some icy environments. For example, while it is clear supraglacial ice

exists on the surface, there is no universal definition of a depth wherein glacial ice should

suddenly be considered 'englacial'. Indeed, supraglacial and englacial ice would probably

be more accurately portrayed as opposite ends of a vertical gradient through the body of a

glacier rather than discrete categories. This may explain some of the overlap in microbiome

composition and structure we observed between supraglacial and englacial environments

(Figs. 4 and 5).

Another potential source of bias in the meta-analysis to be cognizant of is how some

ice samples are collected, specifically basal ice. Due to the logistical difficulty of accessing

the basal ice zone of a glacier, most samples of basal ice included in the meta-analysis were

not collected from inside an excavated tunnel like our samples at Taylor Glacier. Instead,

most of these samples (including our sample of Matanuska's dispersed basal ice), were

collected from easily accessible horizons of basal ice that were exposed at the glacial

margin. These samples have been exposed to sunlight and seasonal temperature

fluctuations that are not found in the glacial interior. It is difficult to accurately predict how

these conditions would affect the basal ice microbiomes. However, it might explain why, for

example, phototrophic taxa (e.g., *Cyanobacteria*) are observed in several of the basal ice

samples from coastal Svalbard glaciers.

4.6 Conclusions and Implications

Our findings show that certain microbes do not persist in a state of dormancy while

frozen, and coupled with other biogeochemical data (3, 4), indicate that the sediment-rich

basal ice horizons of glaciers harbor microbial communities that actively conduct

biogeochemical cycling at subzero temperatures. Although we have limited observations,

our data further implies that a larger diversity of bacteria remain metabolically active in

temperate basal ice found beneath nonpolar glaciers and ice sheet interiors (82, 96). Given

that the basal temperature of the Antarctic and Greenland ice sheets are near or at the

pressure-melting point (97), this raises the intriguing possibility that subglacial regions of

the cryosphere may be more biogeochemically active than previously thought.

Microorganisms in the basal ice of cold-based glaciers may be important

biogeochemical actors. Although rates of microbial activity in these environments are

extremely low and much lower than those of warmer environments (3, 36, 91, 98, 99), this

may be offset by the long residence time of basal ice. Microbial processing in basal ice could

have important implications to material that is transported to and melts at the margin,

representing an important source of dissolved organic carbon, nutrients, and trace metals

to marine ecosystems (23, 26, 100, 101).

Finally, our meta-analysis demonstrates that basal ice environments harbor a

diverse range of microbial communities which can resemble those found in a wide range of

other icy environments. This is likely a reflection of the many different types of basal ice,

which vary considerable in both physical and chemical characteristics (46). Debris-rich

basal ice types, with comparatively higher concentrations of potential nutrients and redox

substrates for microbial metabolism resemble other debris-rich icy environments such as

permafrost and/or perennial cave ice. Likewise, clean basal ice microbial communities are

more like those found in the oligotrophic, debris-free englacial ice that makes up the bulk

of a glacier's mass. Broadly, these findings indicate basal ice diagenesis plays a major role

in microbiome composition and structure.

5. ACKNOWLEDGEMENTS

This study was funded by a Division of Polar Programs grant (ANT-0636828) from

the National Science Foundation. We thank Amanda Achberger, Pierre Amato, Tim Brox,

Lindsay Knippenberg, Scott Montross, and Mark Skidmore for multiple discussions and as

vital partners in tunnel excavation and sampling in Antarctica. We also thank the staff of

McMurdo station and Petroleum Helicopters International for logistical support in the field.

6. REFERENCES

1. Knight PG. 1997. The basal ice layer of glaciers and ice sheets. Quaternary Science Reviews 16:975–993.

2. Souchez R, Jouzel J, Landais A, Chappellaz J, Lorrain R, Tison J-L. 2006. Gas isotopes in ice reveal a vegetated central Greenland during ice sheet invasion. Geophysical Research Letters 33.

3. Doyle S, Montross S, Skidmore M, Christner B. 2013. Characterizing Microbial Diversity and the Potential for Metabolic Function at −15 °C in the Basal Ice of Taylor Glacier, Antarctica. Biology 2:1034–1053.

4. Montross S, Skidmore M, Christner B, Samyn D, Tison J-L, Lorrain R, Doyle S, Fitzsimons S. 2014. Debris-Rich Basal Ice as a Microbial Habitat, Taylor Glacier, Antarctica. Geomicrobiology Journal 31:76–81.

5. Siegert MJ, Ellis-Evans JC, Tranter M, Mayer C, Petit J-R, Salamatin A, Priscu JC. 2001. Physical, chemical and biological processes in Lake Vostok and other Antarctic subglacial lakes. 6864. Nature 414:603–609.

6. Achberger AM, Michaud AB, Vick-Majors TJ, Christner BC, Skidmore ML, Priscu JC, Tranter M. 2017. Microbiology of subglacial environments, p. 83–110. *In* Psychrophiles: from biodiversity to biotechnology. Springer.

7. Samyn D, Svensson A, Fitzsimons SJ. 2008. Dynamic implications of discontinuous recrystallization in cold basal ice: Taylor Glacier, Antarctica. Journal of Geophysical Research 113.

8. Skidmore ML, Foght JM, Sharp MJ. 2000. Microbial Life beneath a High Arctic Glacier. Appl Environ Microbiol 66:3214–3220.

9.   Wadham JL, Bottrell S, Tranter M, Raiswell R. 2004. Stable isotope evidence for

microbial sulphate reduction at the bed of a polythermal high Arctic glacier. Earth

and Planetary Science Letters 219:341–355.

10.   Tung H c., Price P b., Bramall N e., Vrdoljak G. 2006. Microorganisms Metabolizing on

Clay Grains in 3-Km-Deep Greenland Basal Ice. Astrobiology 6:69–86.

11.   Yde JC, Finster KW, Raiswell R, Steffensen JP, Heinemeier J, Olsen J, Gunnlaugsson

HP, Nielsen OB. 2010. Basal ice microbiology at the margin of the Greenland ice

sheet. Annals of Glaciology 51:71–79.

12.   Montross SN, Skidmore M, Tranter M, Kivimaki A-L, Parkes RJ. 2013. A microbial

driver of chemical weathering in glaciated systems. Geology 41:215–218.

13.   Bunt J. 1963. Microbiology of Antarctic sea-ice: diatoms of Antarctic sea-ice as

agents of primary production. Nature 199:1255–1257.

14.   Abyzov S, Mitskevich I, Poglazova M. 1998. The Microflora of Deep Horizons of the

Central Antarctic Ice Sheet. Mikrobiologiya 67:547–555.

15.   Sowers T. 2001. N2O record spanning the penultimate deglaciation from the Vostok

ice core. Journal of Geophysical Research: Atmospheres 106:31903–31914.

16.   Campen RK, Sowers T, Alley RB. 2003. Evidence of microbial consortia metabolizing

within a low-latitude mountain glacier. Geology 31:231.

17.   Souchez R, Janssens L, Lemmens M, Stauffer B. 1995. Very low oxygen concentration

in basal ice from Summit, central Greenland. Geophysical Research Letters 22:2001–

2004.

18.   Souchez R, Bouzette A, Clausen HB, Johnsen SJ, Jouzel J. 1998. A stacked mixing

sequence at the base of the Dye 3 Core, Greenland. Geophysical Research Letters

25:1943–1946.

19.   Price PB, Sowers T. 2004. Temperature dependence of metabolic rates for microbial

growth, maintenance, and survival. PNAS 101:4631–4636.

20.   Wadham JL, Tranter M, Skidmore M, Hodson AJ, Priscu J, Lyons WB, Sharp M, Wynn

P, Jackson M. 2010. Biogeochemical weathering under ice: Size matters: GLACIAL

BIOGEOCHEMICAL WEATHERING. Global Biogeochemical Cycles 24:n/a-n/a.

21.   Barker JD, Grottoli AG, Lyons WB. 2018. Stable isotope evidence for the

biogeochemical transformation of ancient organic matter beneath Suess Glacier,

Antarctica. Arctic, Antarctic, and Alpine Research 50:e1448643.

22.   Hopwood MJ, Carroll D, Dunse T, Hodson A, Holding JM, Iriarte JL, Ribeiro S,

Achterberg EP, Cantoni C, Carlson DF, Chierici M, Clarke JS, Cozzi S, Fransson A, Juul-

Pedersen T, Winding MS, Meire L. 2019. Review Article: How does glacier discharge

affect marine biogeochemistry and primary production in the Arctic? The

Cryosphere Discussions 1–51.

23.   Vick-Majors TJ, Michaud AB, Skidmore ML, Turetta C, Barbante C, Christner BC, Dore

JE, Christianson K, Mitchell AC, Achberger AM, Mikucki JA, Priscu JC. 2020.

Biogeochemical Connectivity Between Freshwater Ecosystems beneath the West

Antarctic Ice Sheet and the Sub-Ice Marine Environment. Global Biogeochemical

Cycles 34:e2019GB006446.

24.   Hood E, Fellman J, Spencer RGM, Hernes PJ, Edwards R, D'Amore D, Scott D. 2009.

Glaciers as a source of ancient and labile organic matter to the marine environment.

7276. Nature 462:1044–1047.

25.   Bhatia MP, Das SB, Xu L, Charette MA, Wadham JL, Kujawinski EB. 2013. Organic

carbon export from the Greenland ice sheet. Geochimica et Cosmochimica Acta

109:329–344.

26.   Hawkings JR, Skidmore ML, Wadham JL, Priscu JC, Morton PL, Hatton JE, Gardner CB,

Kohler TJ, Stibal M, Bagshaw EA, Steigmeyer A, Barker J, Dore JE, Lyons WB, Tranter

M, Spencer RGM, Team the SS. 2020. Enhanced trace element mobilization by

Earth's ice sheets. PNAS 117:31648–31659.

27.   Priscu JC, Tulaczyk S, Studinger M, Kennicutt M, Christner BC, Foreman CM. 2008.

Antarctic subglacial water: origin, evolution and ecology. Polar lakes and rivers:

limnology of Arctic and Antarctic aquatic ecosystems Oxford University Press,

Oxford 119–135.

28.   Batjes NH. 2014. Total carbon and nitrogen in the soils of the world. European

Journal of Soil Science 65:10–21.

29.   Wadham JL, Tranter M, Tulaczyk S, Sharp M. 2008. Subglacial methanogenesis: A

potential climatic amplifier? Global Biogeochemical Cycles 22.

30.   Wadham JL, Arndt S, Tulaczyk S, Stibal M, Tranter M, Telling J, Lis GP, Lawson E,

Ridgwell A, Dubnick A, Sharp MJ, Anesio AM, Butler CEH. 2012. Potential methane

reservoirs beneath Antarctica. Nature 488:633–637.

31.   Boyd ES, Skidmore M, Mitchell AC, Bakermans C, Peters JW. 2010. Methanogenesis

in subglacial sediments: Subglacial methanogenesis. Environmental Microbiology

Reports 2:685–692.

32.   Dieser M, Broemsen ELJE, Cameron KA, King GM, Achberger A, Choquette K,

Hagedorn B, Sletten R, Junge K, Christner BC. 2014. Molecular and biogeochemical

evidence for methane cycling beneath the western margin of the Greenland Ice

Sheet. The ISME Journal 8:2305–2316.

33.   Michaud AB, Dore JE, Achberger AM, Christner BC, Mitchell AC, Skidmore ML, Vick-

Majors TJ, Priscu JC. 2017. Microbial oxidation as a methane sink beneath the West

Antarctic Ice Sheet. 8. Nature Geoscience 10:582–586.

34.   Janssen S, McDonald D, Gonzalez A, Navas-Molina JA, Jiang L, Xu ZZ, Winker K, Kado

DM, Orwoll E, Manary M, Mirarab S, Knight R. 2018. Phylogenetic Placement of Exact

Amplicon Sequences Improves Associations with Clinical Information. mSystems

3:e00021-18.

35.   Christner BC, Mikucki JA, Foreman CM, Denson J, Priscu JC. 2005. Glacial ice cores: A

model system for developing extraterrestrial decontamination protocols. Icarus

174:572–584.

36.   Amato P, Christner BC. 2009. Energy Metabolism Response to Low-Temperature

and Frozen Conditions in Psychrobacter cryohalolentis. Applied and Environmental

Microbiology 75:711–718.

37.   Caporaso JG, Lauber CL, Walters WA, Berg-Lyons D, Huntley J, Fierer N, Owens SM,

Betley J, Fraser L, Bauer M, others. 2012. Ultra-high-throughput microbial

community analysis on the Illumina HiSeq and MiSeq platforms. The ISME journal

6:1621–1624.

38.   Salter SJ, Cox MJ, Turek EM, Calus ST, Cookson WO, Moffatt MF, Turner P, Parkhill J,

Loman NJ, Walker AW. 2014. Reagent and laboratory contamination can critically

impact sequence-based microbiome analyses. BMC biology 12:87.

39.   Eisenhofer R, Minich JJ, Marotz C, Cooper A, Knight R, Weyrich LS. 2019.

Contamination in Low Microbial Biomass Microbiome Studies: Issues and

Recommendations. Trends in Microbiology 27:105–117.

40.   Callahan BJ, McMurdie PJ, Rosen MJ, Han AW, Johnson AJA, Holmes SP. 2016.

DADA2: High-resolution sample inference from Illumina amplicon data. Nature

Methods 13:581–583.

41.   Wang Q, Garrity GM, Tiedje JM, Cole JR. 2007. Naïve Bayesian Classifier for Rapid

Assignment of rRNA Sequences into the New Bacterial Taxonomy. Appl Environ

Microbiol 73:5261–5267.

42.   Quast C, Pruesse E, Yilmaz P, Gerken J, Schweer T, Yarza P, Peplies J, Glockner FO.

2013. The SILVA ribosomal RNA gene database project: improved data processing

and web-based tools. Nucleic Acids Research 41:D590–D596.

43.   McDonald D, Price MN, Goodrich J, Nawrocki EP, DeSantis TZ, Probst A, Andersen

GL, Knight R, Hugenholtz P. 2012. An improved Greengenes taxonomy with explicit

ranks for ecological and evolutionary analyses of bacteria and archaea. The ISME

journal 6:610.

44.   Mirarab S, Nguyen N, Warnow T. 2012. SEPP: SATé-enabled phylogenetic placement,

p. 247–258. *In* Biocomputing 2012. World Scientific.

875   45.   Kimes PK, Liu Y, Neil Hayes D, Marron JS. 2017. Statistical significance for

876   hierarchical clustering. Biometrics 73:811–821.

877   46.   Hubbard B, Cook S, Coulson H. 2009. Basal ice facies: a review and unifying

878   approach. Quaternary Science Reviews 28:1956–1969.

879   47.   Perini L, Gostinčar C, Gunde-Cimerman N. 2019. Fungal and bacterial diversity of

880   Svalbard subglacial ice. 1. Scientific Reports 9:20230.

881   48.   Miteva V, Rinehold K, Sowers T, Sebastian A, Brenchley J. 2015. Abundance, viability

882   and diversity of the indigenous microbial populations at different depths of the

883   NEEM Greenland ice core. Polar Research 34:25057.

884   49.   Miteva V, Sowers T, Schüpbach S, Fischer H, Brenchley J. 2016. Geochemical and

885   Microbiological Studies of Nitrous Oxide Variations within the New NEEM

886   Greenland Ice Core during the Last Glacial Period. Geomicrobiology Journal 33:647–

887   660.

888   50.   Price PB, Morris J, Bay RC, Adhikari A, Giovannoni SJ, Vergin KL. 2015. Molecular

889   Fossils from Microorganisms Preserved in Glacial Ice. bioRxiv

890   https://doi.org/10.1101/019240.

891   51.   Zhong Z-P, Tian F, Roux S, Gazitúa MC, Solonenko NE, Li Y-F, Davis ME, Van Etten JL,

892   Mosley-Thompson E, Rich VI, Sullivan MB, Thompson LG. 2021. Glacier ice archives

893   nearly 15,000-year-old microbes and phages. Microbiome 9:160.

894   52.   Itcus C, Pascu MD, Lavin P, Perşoiu A, Iancu L, Purcarea C. 2018. Bacterial and

895   archaeal community structures in perennial cave ice. Sci Rep 8.

896   53.   Paun VI, Icaza G, Lavin P, Marin C, Tudorache A, Perşoiu A, Dorador C, Purcarea C.

897   2019. Total and Potentially Active Bacterial Communities Entrapped in a Late

Glacial Through Holocene Ice Core From Scarisoara Ice Cave, Romania. Frontiers in

Microbiology 10:1193.

54. Goordial J, Davila A, Lacelle D, Pollard W, Marinova MM, Greer CW, DiRuggiero J,

McKay CP, Whyte LG. 2016. Nearing the cold-arid limits of microbial life in

permafrost of an upper dry valley, Antarctica. 7. The ISME Journal 10:1613–1624.

55. Monteux S, Weedon JT, Blume-Werry G, Gavazov K, Jassey VEJ, Johansson M, Keuper

F, Olid C, Dorrepaal E. 2018. Long-term in situ permafrost thaw effects on bacterial

communities and potential aerobic respiration. The ISME Journal 12:2129–2141.

56. O'Connor BRW, Fernández-Martínez MÁ, Léveillé RJ, Whyte LG. 2021. Taxonomic

Characterization and Microbial Activity Determination of Cold-Adapted Microbial

Communities in Lava Tube Ice Caves from Lava Beds National Monument, a High-

Fidelity Mars Analogue Environment. Astrobiology 21:613–627.

57. Skidmore M, Anderson SP, Sharp M, Foght J, Lanoil BD. 2005. Comparison of

Microbial Community Compositions of Two Subglacial Environments Reveals a

Possible Role for Microbes in Chemical Weathering Processes. Applied and

Environmental Microbiology 71:6986–6997.

58. Cheng SM, Foght JM. 2007. Cultivation-independent and -dependent

characterization of Bacteria resident beneath John Evans Glacier: Characterization

of subglacial Bacterial communities. FEMS Microbiology Ecology 59:318–330.

59. Katayama T, Tanaka M, Moriizumi J, Nakamura T, Brouchkov A, Douglas TA, Fukuda

M, Tomita F, Asano K. 2007. Phylogenetic Analysis of Bacteria Preserved in a

Permafrost Ice Wedge for 25,000 Years. Applied and Environmental Microbiology

73:2360–2363.

60.   Steven B, Pollard WH, Greer CW, Whyte LG. 2008. Microbial diversity and activity

through a permafrost/ground ice core profile from the Canadian high Arctic.

Environmental Microbiology 10:3388–3403.

61.   Sonjak S, Frisvad JC, Gunde-Cimerman N. 2006. Penicillium Mycobiota in Arctic

Subglacial Ice. Microbial Ecology 52:207–216.

62.   Fraser CI, Connell L, Lee CK, Cary SC. 2018. Evidence of plant and animal

communities at exposed and subglacial (cave) geothermal sites in Antarctica. Polar

Biol 41:417–421.

63.   Willerslev E, Cappellini E, Boomsma W, Nielsen R, Hebsgaard MB, Brand TB,

Hofreiter M, Bunce M, Poinar HN, Dahl-Jensen D, Johnsen S, Steffensen JP, Bennike O,

Schwenninger J-L, Nathan R, Armitage S, de Hoog C-J, Alfimov V, Christl M, Beer J,

Muscheler R, Barker J, Sharp M, Penkman KEH, Haile J, Taberlet P, Gilbert MTP,

Casoli A, Campani E, Collins MJ. 2007. Ancient Biomolecules from Deep Ice Cores

Reveal a Forested Southern Greenland. Science 317:111–114.

64.   Kochkina G, Ivanushkina N, Karasev S, Gavrish EY, Gurina L, Evtushenko L, Spirina E,

Vorob'eva E, Gilichinskii D, Ozerskaya S. 2001. Survival of micromycetes and

actinobacteria under conditions of long-term natural cryopreservation.

Microbiology 70:356–364.

65.   Steven B, Briggs G, McKay CP, Pollard WH, Greer CW, Whyte LG. 2007.

Characterization of the microbial diversity in a permafrost sample from the

Canadian high Arctic using culture-dependent and culture-independent methods:

Microbial diversity in Canadian high Arctic permafrost. FEMS Microbiology Ecology

59:513–523.

66.   Perreault NN, Greer CW, Andersen DT, Tille S, Lacrampe-Couloume G, Lollar BS,

Whyte LG. 2008. Heterotrophic and Autotrophic Microbial Populations in Cold

Perennial Springs of the High Arctic. Appl Environ Microbiol 74:6898–6907.

67.   Lacelle D, Radtke K, Clark ID, Fisher D, Lauriol B, Utting N, Whyte LG. 2011.

Geomicrobiology and occluded O2–CO2–Ar gas analyses provide evidence of

microbial respiration in ancient terrestrial ground ice. Earth and Planetary Science

Letters 306:46–54.

68.   Boetius A, Anesio AM, Deming JW, Mikucki JA, Rapp JZ. 2015. Microbial ecology of

the cryosphere: sea ice and glacial habitats. 11. Nature Reviews Microbiology

13:677–690.

69.   Spirina E, Durdenko E, Demidov N, Abramov A, Romanovsky V, Rivkina E. 2017.

Halophilic-psychrotrophic bacteria of an Alaskan cryopeg—a model for

astrobiology. Paleontological Journal 51:1440–1452.

70.   Doyle S, Dieser M, Broemsen E, Christner B. 2012. General characteristics of cold-

adapted microorganisms, p. 103–125. *In* Polar Microbiology: Life in a Deep Freeze.

American Society of Microbiology.

71.   Filippidou S, Wunderlin T, Junier T, Jeanneret N, Dorador C, Molina V, Johnson DR,

Junier P. 2016. A Combination of Extreme Environmental Conditions Favor the

Prevalence of Endospore-Forming Firmicutes. Front Microbiol 7.

72.   Segev E, Smith Y, Ben-Yehuda S. 2012. RNA Dynamics in Aging Bacterial Spores. Cell

148:139–149.

73.   Korza G, Setlow B, Rao L, Li Q, Setlow P. 2016. Changes in Bacillus Spore Small

Molecules, rRNA, Germination, and Outgrowth after Extended Sublethal Exposure to

Various Temperatures: Evidence that Protein Synthesis Is Not Essential for Spore

Germination. J Bacteriol 198:3254–3264.

74. Burkert A, Douglas TA, Waldrop MP, Mackelprang R. 2019. Changes in the Active,

Dead, and Dormant Microbial Community Structure across a Pleistocene Permafrost

Chronosequence. Appl Environ Microbiol 85:e02646-18.

75. Blazewicz SJ, Barnard RL, Daly RA, Firestone MK. 2013. Evaluating rRNA as an

indicator of microbial activity in environmental communities: limitations and uses.

11. The ISME Journal 7:2061–2068.

76. Aciego SM, Cuffey KM, Kavanaugh JL, Morse DL, Severinghaus JP. 2007. Pleistocene

ice and paleo-strain rates at Taylor Glacier, Antarctica. Quaternary Research

68:303–313.

77. Buizert C, Baggenstos D, Jiang W, Purtschert R, Petrenko VV, Lu Z-T, Müller P, Kuhl

T, Lee J, Severinghaus JP, others. 2014. Radiometric 81Kr dating identifies 120,000-

year-old ice at Taylor Glacier, Antarctica. Proceedings of the National Academy of

Sciences 111:6876–6881.

78. Lawson DE, Kulla JB. 1978. An Oxygen Isotope Investigation of the Origin of the

Basal Zone of the Matanuska Glacier, Alaska. The Journal of Geology 86:673–685.

79. Strasser JC, Lawson DE, Larson GJ, Evenson EB, Alley RB. 1996. Preliminary results

of tritium analyses in basal ice, Matanuska Glacier, Alaska, U.S.A.: evidence for

subglacial ice accretion. Annals of Glaciology 22:126–133.

80. Lawson D. 1979. Sedimentological Analysis of the Western Terminus Region of the

Matanuska Glacier, Alaska 132.

81.   Price PB. 2000. A habitat for psychrophiles in deep Antarctic ice. PNAS 97:1247–

990   1251.

991   82.   Lawson DE, Strasser JC, Evenson EB, Alley RB, Larson GJ, Arcone SA. 1998.

992   Glaciohydraulic supercooling: a freeze-on mechanism to create stratified, debris-

993   rich basal ice: I. Field evidence. Journal of Glaciology 44:547–562.

994   83.   Rhodes RH, Faïn X, Stowasser C, Blunier T, Chappellaz J, McConnell JR, Romanini D,

995   Mitchell LE, Brook EJ. 2013. Continuous methane measurements from a late

996   Holocene Greenland ice core: Atmospheric and in-situ signals. Earth and Planetary

997   Science Letters 368:9–19.

998   84.   Lamarche-Gagnon G, Wadham JL, Sherwood Lollar B, Arndt S, Fietzek P, Beaton AD,

999   Tedstone AJ, Telling J, Bagshaw EA, Hawkings JR, Kohler TJ, Zarsky JD, Mowlem MC,

Anesio AM, Stibal M. 2019. Greenland melt drives continuous export of methane

from the ice-sheet bed. 7737. Nature 565:73–77.

85.   Lee JE, Edwards JS, Schmitt J, Fischer H, Bock M, Brook EJ. 2020. Excess methane in

Greenland ice cores associated with high dust concentrations. Geochimica et

Cosmochimica Acta 270:409–430.

86.   Stibal M, Wadham JL, Lis GP, Telling J, Pancost RD, Dubnick A, Sharp MJ, Lawson EC,

Butler CEH, Hasan F, Tranter M, Anesio AM. 2012. Methanogenic potential of Arctic

and Antarctic subglacial environments with contrasting organic carbon sources.

Global Change Biology 18:3332–3345.

87.   Montross SN. 2012. Biogeochemistry of basal ice from Taylor Glacier, Antarctica.

Montana State University.

88.   Cuffey KM, Conway H, Gades AM, Hallet B, Lorrain R, Severinghaus JP, Steig EJ,

1012   Vaughn B, White JWC. 2000. Entrainment at cold glacier beds. Geology 28:351–354.

1013   89.   Sleewaegen S, Samyn D, Fitzsimons SJ, Lorrain RD. 2003. Equifinality of basal ice

1014   facies from an Antarctic cold-based glacier. Annals of Glaciology 37:257–262.

1015   90.   Paterson WSB. 1994. Physics of glaciers. Butterworth-Heinemann.

1016   91.   Panikov NS, Flanagan PW, Oechel WC, Mastepanov MA, Christensen TR. 2006.

1017   Microbial activity in soils frozen to below −39°C. Soil Biology and Biochemistry

1018   38:785–794.

1019   92.   Bakermans C, Skidmore ML. 2011. Microbial Metabolism in Ice and Brine at −5°C:

1020   Microbial metabolism in ice and brine at −5°C. Environmental Microbiology

1021   13:2269–2278.

1022   93.   Tuorto SJ, Darias P, McGuinness LR, Panikov N, Zhang T, Häggblom MM, Kerkhof LJ.

1023   2014. Bacterial genome replication at subzero temperatures in permafrost. ISME J

1024   8:139–149.

1025   94.   Segura JH, Nilsson MB, Haei M, Sparrman T, Mikkola J-P, Gräsvik J, Schleucher J,

1026   Öquist MG. 2017. Microbial mineralization of cellulose in frozen soils. Nat Commun

1027   8:1154.

1028   95.   Sheridan PP, Miteva VI, Brenchley JE. 2003. Phylogenetic Analysis of Anaerobic

1029   Psychrophilic Enrichment Cultures Obtained from a Greenland Glacier Ice Core.

1030   Applied and Environmental Microbiology 69:2153–2160.

1031   96.   Bell RE, Ferraccioli F, Creyts TT, Braaten D, Corr H, Das I, Damaske D, Frearson N,

1032   Jordan T, Rose K, Studinger M, Wolovick M. 2011. Widespread Persistent Thickening

1033   of the East Antarctic Ice Sheet by Freezing from the Base. Science 331:1592–1595.

97. Gow AJ, Ueda HT, Garfield DE. 1968. Antarctic Ice Sheet: Preliminary Results of First

Core Hole to Bedrock. Science 161:1011–1013.

98. Christner BC. 2002. Incorporation of DNA and Protein Precursors into

Macromolecules by Bacteria at -15oC. Applied and Environmental Microbiology

68:6435–6438.

99. Amato P, Doyle SM, Battista JR, Christner BC. 2010. Implications of Subzero

Metabolic Activity on Long-Term Microbial Survival in Terrestrial and

Extraterrestrial Permafrost. Astrobiology 10:789–798.

100. Rignot E, Jacobs SS. 2002. Rapid Bottom Melting Widespread near Antarctic Ice

Sheet Grounding Lines. Science 296:2020–2023.

101. Jung J, Yoo K-C, Rosenheim BE, Conway TM, Lee JI, Yoon HI, Hwang CY, Yang K, Subt

C, Kim J. 2019. Microbial Fe(III) reduction as a potential iron source from Holocene

sediments beneath Larsen Ice Shelf. Nat Commun 10:1–10.

1047

1048

1049

1050

1051

1052

1053

1054

1055

1056

TABLES

1058   **Table 1.** Description of basal ice types collected in this study

Type	Glacier	Temp (°C)	Description[c]
Clean ice	Taylor	-15[a]	debris-free ice
Banded ice	Taylor	-15[a]	finely stratified layers of debris and ice, layers range in thickness from a few mm to several cm
Solid ice	Taylor	-15[a]	composed primarily of frozen debris with only interstitial ice, no visible layering
Dispersed ice	Matanuska	~0[b]	contains scattered debris aggregated into small clusters, no visible layering

[a]Montross *et al.,* 2014. [b]Lawson *et al.,* 1998. [c]Hubbard *et al.,* 2009

1059

1060   **Table 2.** Cell counts, biomass estimates, and yield of DNA and RNA extracted from basal ice
1061   samples. Sediment-rich basal ice facies are highlighted in gray.
1062

Sample	Cell Density[a] cells g^{-1} ice	Yield (pg g^{-1} ice) DNA	RNA	Biomass Estimates genomes[b] g^{-1} ice	ATP (pg g^{-1} ice)
Clean07	$0.3 - 0.5 \times 10^3$	4.7	0.7	1.9×10^3	<0.01 - 0.04
Clean09	n.d.	1.4	n.d.	0.6×10^3	n.d.
Banded07	$1.8 - 18.2 \times 10^3$	1.7×10^4	2.3	6.8×10^6	n.d.
Banded09	n.d.	1.6×10^4	0.8	6.5×10^6	0.24 - 2.8
Solid	6.7×10^3	1.4×10^4	1.2	5.7×10^6	<0.01
Dispersed	n.d.	9.9×10^4	3.2	4.0×10^7	0.06 - 0.14

[a]Doyle et al. 2013. [b]Number of microbial genomes estimated using 2.5 fg of DNA per genome. n.d.= no data

1063

1064

1065

1066

1067

1068

1069

1070

 FIGURE LEGENDS

 Figure 1. [A] Relative abundances of microbial lineages observed in basal ice based on the

 use of DNA and RNA templates for detecting 16S rRNA sequences. The graph displays the

 highest resolution classification for the most abundant taxa and was constructed as

 follows: First, ASVs were clustered by genus and any having a relative abundance ≥20% in

 at least one of the samples were plotted. This procedure was subsequently repeated with

 the remaining unplotted ASVs at the rank of family, order, class, and finally phylum. Any

 remaining rare taxa left after this procedure were not plotted. The bottom plot was scaled

 to reduce inflated relative abundance biases introduced during curation. **[B]** Scatter plots

 of potentially active ASVs within four types of basal ice. Each point represents an individual

 ASV. The dotted-line indicates a 1:1 ratio between the RNA- and DNA-based 16S rRNA

 libraries. Gray points denote ASVs observed only in the DNA-based libraries. In the banded

 basal ice plot, circles denote ASVs from the Banded07 sample while squares denote ASVs

 from the Banded09 sample. Detected ASVs in each plot denotes the number of ASVs

 observed in both the RNA- and DNA-based 16S rRNA libraries.

 Figure 2. Microbial alpha diversity (Shannon) within various perennially frozen

 environments.

 Figure 3. Hierarchical cluster analysis of UF distances between samples using Ward's

 minimum variance method. The significance of each node was calculated with a Monte

 Carlo simulation using the 'sigclust2' R package. Family-wise error rate (FWER) was

1093 controlled across the dendrogram using a sequential testing procedure starting at the

1094 central node.

1095

Figure 4. [Left] RDA triplot of the distribution of 1627 microbial lineages across five

different perennially frozen environments. Colored circles represent individual samples.

Centroids of each environment type are indicated with an "×". Plus signs (+) represent

microbial lineages. The purple circle denotes the equilibrium contribution of the overall

model and delineates lineages which substantially constrained the overall ordination.

Those lineages are highlighted in red. **[Right]** Magnitude of the scalar projections of these

36 lineages onto the centroids for each ice types approximates that lineage's association

with that frozen environment. Scalar projections smaller than the equilibrium contribution

were considered inconclusive and marked gray. Positive and negative associations are

black and red, respectively.

1106

1107 **Figure 5.** Distance-based redundancy analysis of Aitchison distances between different

1108 perennially frozen environments. Colored circles represent centroids of each study's

1109 samples. Black triangles represent individual early studies of basal ice and permafrost

1110 based on Sanger sequencing of 16S rRNA clone libraries.

1111

1112

1113

1114

1115

SUPPLEMENTAL MATERIAL LEGENDS

Figure S1. [A] A subglacial tunnel was excavated into the basal zone of Taylor Glacier's

northern lateral margin. **[B]** View from inside one of the excavation tunnels. **[C]** During the

2007 expedition, a vertical shaft was excavated, and a stratigraphic sequence of multiple

basal ice types was sampled. **[D]** During the 2009 expedition, a large chamber was

excavated to sample a horizon of debris-rich banded basal ice.

1122

1123 **Figure S2.** An exposed horizon of debris-rich basal ice near the terminus of Matanuska

1124 Glacier, Alaska. The sampled area, collected using an electric chainsaw, is highlighted inside

1125 the yellow box.

1126

1127 **Figure S3.** Examples of each type of basal ice analyzed in this study. Clean ice, banded ice,

1128 and solid ice were collected from Taylor Glacier, Antarctica. Dispersed ice was collected

1129 from Matanuska Glacier, Alaska.

1130

Figure S4. Hierarchical cluster analysis of wUF distances between samples using Ward's

minimum variance method. The significance of each node was calculated with a Monte

Carlo simulation using the 'sigclust2' R package. Family-wise error rate (FWER) was

controlled across the dendrogram using a sequential testing procedure starting at the

central node. The three dashed branches in the dendogram were lengthened slightly to

improve legibility and are thus not to scale.

1137

1138 **Figure S5.** Heatmap of the 40 taxa identified in five early 16S rRNA clone library studies of

1139 icy environments. Red stars denote 'indicator' microbial taxa which were identified in the

1140 overall meta-analysis (Fig. 4). Taxa abundances were transformed with a Hellinger

1141 transformation to facilitate a clearer visual comparison between abundant and sparse taxa.

1142

1143 **Table S1A.** Amount of basal ice, after decontamination, used for either DNA or RNA

1144 extraction.

1145

1146 **Table S1B.** Description of each sample included in the meta-analysis.

1147

1148 **Table S1C.** DADA2 filtering and curation parameters during meta-analysis.

1149

1150 **Table S1D.** DADA2 pipeline statistics for each sample in the meta-analysis.

1151

1152 **Table S1E.** Shared ASVs between all four basal ice types.

1153

1154 **Table S1F.** Potentially active ASVs within each type of basal ice.

Perennial Cave Ice
- Scărișoara Cave (Romania)
- Lava Beds National Monument (California)

Permafrost
- University Valley (McMurdo Dry Valleys)
- Storflaket Permafrost Plateau (Sweden)
- Mammoth Mountain (East Siberia)
- Axel Heiberg Island (Canada)
- Alaska Peatland Experiment (APEX)
- Spitsbergen Island (Svalbard)

Basal Ice
- Clean Ice (Taylor Glacier)
- Banded Ice (Taylor Glacier)
- Solid Ice (Taylor Glacier)
- Dispersed Ice (Matanuska Glacier)
- Devon Ice Cap (Canada)
- Various coastal glaciers (Svalbard)

Supraglacial Ice
- Greenland Ice Sheet (Kangerlussuaq)
- Issunguata Sermia Glacier (Greenland)
- Lyngmarksbræ Glacier (Greenland)
- Various alpine glaciers (West China)
- Anuchin Glacier (Antarctica)
- Volcanic crater glaciers (Mexico)

Nodes
- significant ($p<0.05$)
- untested by FWER
- not significant
- cluster too small

Englacial Ice
- Matanuska Glacier (Valkyrie)
- Ice Core: NEEM
- Ice Core: GISP2D
- Ice Core: WAIS Divide
- Ice Core: Guliya Ice Cap (China)
- Whillans Ice Stream (WISSARD)