**Title:** Ribosomal RNA methylation by GidB modulates discrimination of mischarged tRNA

**Authors:** Zhuo Bi[1,2†],Yu-Xiang Chen[3,4†], Iris D. Young[5,6†], Hong-Wei Su[1,7], Yuemeng Chen[1], Jia-Yao

Hong[1], James S. Fraser[5], and Babak Javid[1,3*]

**Affiliations:**

[1] Center for Global Health and Infectious Disease, Tsinghua University School of Medicine, Beijing,

China

[2] School of Life Science, Tsinghua University, Beijing, China

[3] Division of Experimental Medicine, University of California, San Francisco, CA, USA

[4]Present address: Department of Genetics, Stanford University, Stanford, CA, USA

[5] Department of Bioengineering and Therapeutic Sciences, University of California San Francisco,

San Francisco, CA, USA

[6] Molecular Biophysics and Integrated Bioimaging Division, Lawrence Berkeley National Laboratory,

Berkeley, CA, USA

[7]Present address: College of Animal Sciences, Zhejiang University, Hangzhou, China

[†]These authors contributed equally.

[*] to whom correspondence should be addressed: babak.javid@ucsf.edu

**Summary**

Despite redundant cellular pathways to minimize translational errors, errors in protein synthesis are common. Pathways and mechanisms to minimize errors are classified as pre-ribosomal or ribosomal. Pre-ribosomal pathways are primarily concerned with the appropriate charging of tRNAs with their cognate amino acid. By contrast, the ribosomal decoding centre is considered 'blind' to mischarged tRNAs since these have cognate codon•anti-codon pairing. Here, we identified that in mycobacteria, deletion of the 16S ribosomal RNA methyltransferase *gidB* led to increased ribosomal discrimination of mischarged tRNAs. Discrimination only occurred in mycobacteria enriched from environments or genetic backgrounds with high rates of mistranslation. GidB deletion was necessary but not sufficient for reducing mistranslation due to misacylation. Analysis of new cryoEM structures of the *M. smegmatis* ribosomes derived from wild-type and *gidB*-deleted strains point to the interaction between the base methylated by GidB on the 16S RNA and an asparagine on the ribosomal S12 protein that when mistranslated to aspartate may be involved in altering translational fidelity. Our data suggest a mechanism by which mycobacterial ribosomes can discriminate mischarged tRNAs and that 16S rRNA differential methylation by GidB may act to prevent catastrophic translational error.

**Introduction**

All clades of life have evolved multiple, redundant pathways to reduce translational error [1, 2]. Despite these mechanisms, errors in protein synthesis are remarkably common and are orders of magnitude more frequent than errors in DNA or RNA synthesis [3-7]. There is no one 'optimal' rate of error. Both the error rates and the dominant sources of error can be both species and even organelle specific [1, 8]. Furthermore, translational errors (mistranslation) may result in adaptive phenotypes, particularly in the context of environmental stressors [1, 5, 6, 9-20]. However, excess mistranslation can also cause protein aggregation [21, 22], organ degeneration [23, 24] and is the mechanism for the bactericidal activity of aminoglycosides [25]. Collectively, these lines of evidence suggest that no optimal balance for translational error exists and that selection favors tunable and context-specific mistranslation rates [1, 3, 5, 26]. However, the precise mechanisms by which mistranslation rates are tuned are poorly understood.

In addition, molecular mechanisms of translational error vary considerably, as do the proof-reading pathways that have evolved to reduce them. Generally, sources of error and proof-reading can be divided into pre-ribosomal and ribosomal mechanisms [1]. Pre-ribosomal errors arise from mischarging of the tRNA with a non-cognate amino acid and proof-reading mechanisms include pre- and post-transfer editing functions of aminoacyl tRNA synthetases [27]. Following aminoacyl-tRNA synthesis, *trans*-acting editing mechanisms can reject mischarged tRNAs [28, 29], and the aminoacyl-tRNA chaperone, EF-Tu optimally binds cognate aminoacyl-tRNAs compared with mischarged tRNAs [30]. These multiple and redundant pre-ribosomal proof-reading steps – solely concerned with the charging of tRNA with its cognate amino acid – have been proposed as necessary since previously described ribosomal proof-reading mechanisms ensure cognate codon·anticodon pairing [2, 31] and therefore are insensitive to the identity of the amino acid charged to the tRNA.

In mycobacteria, the dominant source of translational error is due to the pre-ribosomal indirect tRNA aminoacylation pathway [12, 32, 33]. Most bacteria, with the notable exception of a few proteo-bacteria such as *Escherichia coli,* lack either the glutaminyl- or asparaginyl-tRNA synthetases, or both

[34]. Instead, a two-step indirect tRNA synthesis pathway is required to ensure cognate charging of glutamine and asparagine tRNAs. In the first step, a non-discriminatory glutamyl or aspartyl synthetases mischarge glutaminyl-tRNA with glutamate (i.e. Glu-tRNA$^{Gln}$) and asparaginyl-tRNA with aspartate (i.e. Asp-tRNA$^{Asn}$) respectively. In the second step, the enzyme GatCAB corrects the mischarged aminoacyl moiety by transamidation [35] and see Fig. 1A. Despite its essential function, partial loss-of-function mutations in *gatCAB* can be readily selected in mycobacteria, which lacks both synthetases and mutations in *gatCAB* occur naturally in clinical isolates of pathogenic *M. tuberculosis* [12, 36, 37]. These mutant strains exhibit extremely high (up to 10%/codon) rates of the mistranslation of Gln/Asn codons to Glu/Asp, respectively and specifically [12]. Even in wild-type mycobacteria, the specific mistranslation rates of Gln→Glu and Asn→Asp, measured using similar gain-of-function reporters, are orders of magnitude higher than in *E. coli* – that has the full set of 20 aminoacyl tRNA synthetases and hence lacks GatCAB [38].

Here, we investigated whether further fidelity factors can be identified in mycobacteria. We identifed that deletion of the 16S ribosomal RNA (rRNA) methyltransferase *gidB* is necessary, but not sufficient for the discrimination of mischarged tRNAs. Deletion of *gidB* only increased discrimination of mischarged tRNAs in mycobacteria with elevated mistranslation rates – due to mutation or environmental context. Solving the structure of mycobacterial ribosomes purified from +/- *gidB* strains suggested that methylation of rRNA may interact with mistranslated ribosomal amino acids in the small subunit, implying a potential mechanism by which rRNA methylation contributes to fidelity. Collectively, these results point to an active ribosomal proof-reading of ribosomes beyond codon•anticodon pairing and suggest that nonmethylation of rRNA prevents catastrophic translational error.

**Results**

**A suppressor screen in mycobacteria identifies *gidB* as a potential translational fidelity factor**

We previously used forward genetic screens to identify strains of *Mycobacterium smegmatis* with extremely high specific rates of mistranslation – i.e. mistranslation of Asn→Asp. These screens identified mutations in the essential amidotransferase genes *gatCAB* that resulted in high rates of

mistranslation of Gln/Asn, specifically [12, 36]. We hypothesized that the mycobacterial genome may

encode for other 'fidelity factors' that could modulate mistranslation rates in the background of a

compromised indirect tRNA aminoacylation pathway. We, therefore, designed a suppressor screen

strategy in the strain HWS19 [12], which encodes a three amino acid deletion in *gatA* and,

consequently, has extremely high rates of translational error of Gln→Glu and Asn→Asp specifically

(Fig 1B). We wondered whether HWS19, with a high background mistranslation rate of Gln/Asn

codons, is more susceptible to aminoglycosides, such as streptomycin that are known to increase

ribosomal decoding errors across multiple codons [33, 39]. Plating of equivalent colony forming units

(CFU) of strain HWS19 on low-dose streptomycin-agar led to the recovery of significantly fewer

colonies compared with wild-type (Fig. S1), confirming HWS19 was hypersusceptible to

streptomycin. The basis of our suppressor screen, therefore, was as follows: plating of HWS19 onto

low-dose streptomycin-agar should select for strains with mutations that *decrease* mistranslation on

this high mistranslating background (Fig. 1B, 1C). Suppressors identified from selection on low-dose

streptomycin agar were then tested using our custom dual-luciferase mistranslation reporter system to

verify that they had decreased error rates (Fig. 1C).

To identify mutants leading to decreased mistranslation, we plated $1x10^7$ CFU onto each of 6 agar

plates containing 1 µg/mL streptomycin. The remaining survivors were transformed with dual-

luciferase reporter plasmids that measured specific mistranslation errors of asparagine for aspartate

[12, 17, 40] to identify low mistranslating candidates (Fig. 1C). We initially identified 4 suppressor

mutants with lower mistranslation rates compared with the parent HWS19 strain, and comparable to

wild-type *M. smegmatis* (Fig. 1D). Sequencing the strains revealed three of the four had mutations in

the 16S rRNA methyltransferase *gidB* (Msmeg_6940) – Table S1. The remaining suppressor (C-23)

had multiple mutations including in the gene, *tuf*, coding for the aminoacyl-tRNA chaperone EF-Tu.

(Table S1). We subsequently sequenced just the *gidB* gene of a further 10 candidate suppressors and

identified a further seven with a total of three additional independent mutations in *gidB* (Table S2).

All but one of these further suppressors also had reduced mistranslation rates (Fig. S2), strongly

supporting that mutations in *gidB* are able to suppress the high specific mistranslation rates in strain

HWS19 that arise from mischarging tRNA.

**Deletion of *gidB* increases ribosomal discrimination against misacylated tRNA in mycobacteria**

**with high mistranslation**

Loss of function mutations in GidB can cause low-level streptomycin resistance: lack of methylation

in 16S rRNA interferes with streptomycin binding [41, 42]. We were, therefore, concerned that our

candidates from the screen may simply represent selection against streptomycin. To determine

whether the observed phenotype was independent of streptomycin selection, we deleted *gidB* in both

the HWS19 and parental (wild-type) *M. smegmatis* backgrounds in the complete absence of

streptomycin selection. Deletion strains were complemented with a chromosomally integrated plasmid

expressing *gidB* from its native promoter (Fig. S3). We measured specific mistranslation rates using

two complementary reporter systems: the dual-luciferase system, used in the initial screen, as well as

a dual-fluorescent reporter system (Fig. S4, S5) that uses flow cytometry to measure relative

mistranslation rates [12]. This reporter, a GFP-mRFP fusion protein had a glutamate necessary for

GFP fluorescence mutated to glutamine, abrogating green, but not red fluorescence [12]. Increased

specific mistranslation of Gln→Glu would result in increased GFP fluorescence and hence increased

GFP/RFP ratios. Measurement of specific mistranslation rates with both reporters verified that

deletion of *gidB* significantly increased translational fidelity in strain HWS19, and this phenotype

could be readily complemented (Fig. 2A, B, left panel). Surprisingly, deletion of *gidB* had no

phenotype using the same reporters on a wild-type background (Fig. 2A, B, right panel).

**GidB is necessary for discrimination of mischarged tRNA under conditions that enrich for**

**relatively high mistranslation rates**

Our results suggested that deletion of *gidB* increased translational fidelity only in a mutant strain

(HWS19) that had basal extremely high rates of mistranslation arising from lack of quality control of

mischarged asparagine and glutamine tRNAs and that deletion of *gidB* had no phenotype in wild-type

mycobacteria grown under standard laboratory conditions. We wondered whether deletion of *gidB*

had a phenotype in wild-type mycobacteria, but only under conditions associated with increased rates

of mistranslation. We had previously shown that exposure of wild-type mycobacteria to rifampicin at

1x minimal bactericidal concentration (MBC) could select for survival and growth of *wild-type*

mycobacteria due to two reversible and non-genetic mechanisms: increased rates of specific

mistranslation [12] and a semi-heritable survival program [43], and not due to mutations in the

rifampicin's target *rpoB* or other mutations. We plated wild-type *M. smegmatis* on non-selective LB

medium or low-dose (1xMBC) rifampicin-agar (**Fig. 3A**). As before, deletion of *gidB* isolated from

non-selective medium had no phenotype (**Fig. 3B**). However, bacteria that survived and grew on

rifampicin-agar had increased rates of mistranslation as measured by the dual-fluorescent reporter, in

keeping with prior observations [12, 43]. These increased mistranslation rates were reverted by

deletion of *gidB* in fully complementable manner (**Fig. 3B**), suggesting that deletion of *gidB* increased

translational fidelity under growth conditions causing high rates of mistranslation, even in wild-type

bacteria.

The use of a dual-fluorescent reporter and flow cytometry for measurement of relative mistranslation

rates allowed us to measure the impact of *gidB* deletion on different subpopulations of mycobacteria

in the same experiment: i.e. we could selectively gate on heterogeneous subpopulations that had

relatively high or low GFP/mRFP ratios, representing high and low mistranslation rates specifically

(**Fig. 3C**). This allowed us to ask the question as to whether deletion of *gidB* was sufficient, as well as

necessary, for the discrimination of mischarged tRNA. We gated on bacteria with the highest (10%)

and lowest (10%) green/red (i.e. relative mistranslation rates) ratios from the two experimental

conditions, i.e., bacteria isolated from 'standard' non-selective LB agar, and from low-dose rifampicin

agar, which enriched for bacteria with higher average mistranslation rates.  Deletion of *gidB* had no

observable phenotype on translational fidelity in bacteria isolated from non-selective media, even

when only the cells with the highest apparent mistranslation rates were analysed (**Fig. 3D**). By

contrast, in bacteria enriched for higher mistranslation rates (rifampicin exposure) *gidB* deletion could

revert the mistranslation phenotype in a complementable manner (**Fig. 3E**). Therefore, *gidB* deletion

appears necessary, but not sufficient for translational fidelity, and it is only within environmental

contexts that enrich for high rates of mistranslation that deletion of *gidB* causes increased ribosomal

discrimination of misacylated tRNA in otherwise wild-type mycobacteria.

**Testing the generality of mischarged tRNA discrimination by deletion of *gidB* beyond**

**mischarged tRNA$^{Gln}$ and tRNA$^{Asn}$**

The previous experiments confirm that deletion of *gidB* increased discrimination of physiologically

mischarged Glu-tRNA$^{Gln}$ and Asp-tRNA$^{Asn}$. We next wanted to ask whether this discrimination of

mischarged tRNA was a general property that would allow discrimination of any mischarged tRNA.

To test this idea, we leveraged the unique properties of tRNA$^{Ala}$ and their cognate synthetase. To

ensure the accurate charging of tRNAs, most aminoacyl tRNA synthetases depend on recognizing the

anticodon of their cognate tRNAs as an identity element. For alanyl synthetase (AlaRS), the sole

identity element in all clades of life is a $G^3 \bullet U$ base-pair in the tRNA acceptor stem, and AlaRS can

charge any tRNA with this identity element, regardless of anti-codon identity, i.e. the $G^3 \bullet U$ base-pair

is both necessary and sufficient for alanine aminoacylation by AlaRS [44-46]. Therefore, the

anticodon of any alanyl-tRNA can be mutated to any triplet, and the tRNA will still be charged with

aminoacyl-alanine [44] and will mediate specific translational errors. We had previously mutated the

anticodon of a mycobacterial alanyl-tRNA to CCA, coding for tryptophan, which would result in

specific mistranslation of alanine for tryptophan at UGG codons [17] – **Fig. 4A**. To measure specific

mistranslation, we modified the dual-luciferase reporter system. We identified a critical alanine

residue in Renilla luciferase, which, when mutated to tryptophan (A214W), resulted in >20-fold

decrease in activity (Fig. S6). This reporter would thus be able to discriminate mistranslation errors

greater than 5%/codon of tryptophan to alanine. We transformed the mutant alanyl-tRNA

(tRNA$_{CCA}^{Ala}$), cloned into a tetracycline-inducible plasmid into wild-type and Δ*gidB M. smegmatis*,

along with the specific dual-luciferase reporters. Induction of tRNA$_{CCA}^{Ala}$ resulted in increased

specific mistranslation of tryptophan for alanine, as expected. Deletion of *gidB* did not increase

translational fidelity, even in this high mistranslating context (**Fig. 4B**). Next we transformed the

mutant  alanyl-tRNA and reporters into The high basal mistranslation (Gln→Glu, Asn→Asp,

specifically) strain HWS19 and measured the rates of Trp→Ala mistranslation using the new Ren-FF

dual-luciferase reporter. Deletion of *gidB* decreased rates of Trp→Ala mistranslation as measured by

rescue of Renilla activity in the HWS19 background, suggesting that in this context, ∆*gidB* ribosomes

could discriminate against even Trp→Ala mistranslation (**Fig. 4C**). Surprisingly, this phenotype is

only observed in this strain with high rates of 'physiological' mistranslation and not in the WT strain.

Taken together, these results suggested deletion of *gidB* was necessary but not sufficient for

discrimination of mischarged tRNAs and that a further environmental context, possibly generated via

excess 'physiological' mistranslation or some other stressor was also required.

**Deletion of *gidB* causes reduced tolerance to rifampicin**

We tested the physiological relevance of increased translational fidelity caused by *gidB* deletion in

rifampicin antibiotic tolerance. We had previously demonstrated that increased, specific,

mistranslation due to Glu-tRNA$^{Gln}$ and Asp-tRNA$^{Asn}$ mischarging caused increased tolerance to the

antibiotic rifampicin due to mistranslation of critical residues in the drug target, RpoB [12, 17, 36].

Since antibiotic tolerance is mediated by a subpopulation of bacteria that resist killing [47-57], we

hypothesised that deletion of *gidB*, and the subsequent increase in translational fidelity, would render

the most tolerant subpopulation susceptible to rifampicin. Deletion of *gidB* resulted in significantly

increased killing of *M. smegmatis* exposed to rifampicin in axenic culture in both the high

mistranslating strain HWS19 and, crucially, also in wild-type mycobacteria (Fig. 5A, B). This

suggests that rifampicin treatment is a sufficient condition for deletion of *gidB* to exert its effect.

**Molecular mechanism of GidB-mediated methylation in the ribosome**

To identify the molecular mechanism of changes in adaptive mistranslation, we wish to resolve the

structure of ribosomes from HWS19 and HWS19-∆*gidB* cells for analysis by cryo-electron

microscopy. Sucrose gradient ultracentrifugation and ribosome subunit profiling of ribosomes from

the strains did not reveal difference in the distribution of different ribosome species, suggesting that

deletion of *gidB* did not alter ribosome biogenesis (Fig. S8). Three dimensional reconstructions of 70S

fractions isolated from sucrose gradient and the models refined into those maps did not reveal

significant changes in structure, likely due to the limited resolution (**Fig. 6A, B**). Analysis of the

resulting models does point to a hypothesis for the modulation of mistranslation in ΔgidB ribosomes

as a function of GatCAB activity: an interaction between 16S G507 (the target of GidB methylation)

and S12 Asn46 (Fig. 6C, D). When GatCAB functions, minimizing mistranslation, as in WT *M.*

*smegmatis*, Asn46 is reliably coded as Asn; however, in the HSW19 background and other conditions

with diminished GatCAB activity, Asn46 will be coded as Asp at a much higher frequency. The

interaction between G507 and the S12 loop containing Asn46 has previously been implicated in

stabilizing tRNA-mRNA interactions during aminoacyl-tRNA selection in the decoding process in *M.*

*tuberculosis* [42]. Phenotypic misincorporation from Asn to Asp likely alters this interaction and the

conformational dynamics of this key loop. Further structural work, at higher resolution, will be

needed to unravel how the interactions between the G507 methylation, tRNA identity, mRNA

structure, and antibiotic binding affect the relative rates of adaptive mistranslation.

**Discussion**

The ubiquitous presence of multiple and redundant proof-reading mechanisms for protein synthesis

underscores the importance of accurate protein synthesis in cellular homeostasis. However, errors in

protein synthesis are both more pervasive and frequent than previously anticipated, and increasingly,

translational errors are being recognised as a mechanism for adaptation to hostile environments. One

possible resolution of these two seemingly opposed processes is the recognition that 'optimal' fidelity

is not to minimize errors to as low as possible and is context-specific [5, 6, 58]. For example,

infection by *Salmonella* mutants with both impaired and enhanced translational fidelity were less

productive than wild-type strains [26], and mycobacterial strains with extremely high mistranslation

rates due to mutations in *gatA* grew more slowly than wild-type in axenic culture, but had orders of

magnitude greater survival with rifampicin treatment [12].

In this study we sought to identify further translational fidelity factors in mycobacteria, which we had

previously shown have high, but specific rates of mistranslation due to the indirect tRNA

aminoacylation pathway required for aminoacylation of glutaminyl- and asparaginyl-tRNAs mediated

by GatCAB [10, 12, 36]. Our screen identified loss-of-function mutations in *gidB* conferring

increased fidelity to errors generated by mutants in this pathway (**Fig. 7**). We had previously shown

that deletion of *gidB* in *M. tuberculosis* increased fidelity of 'wobble' ribosomal decoding errors, but

not for mistranslation of asparagine for aspartate – one of the two specific errors in mycobacteria due

to the indirect tRNA pathway [42]. Those studies, performed with otherwise wild-type mycobacteria

under standard laboratory conditions, mimic similar conditions where in this study *gidB* deletion

under those conditions also lacked a phenotype.

Although the ribosome is known to engage in a number of proof-reading functions, these have been

confined to discrimination against non-cognate codon•anti-codon pairing, not discrimination of

mischarged tRNAs where codon•anti-codon pairing remains fully cognate [2, 59-62]. One study by

Cornish and colleagues specifically addressed the question of ribosome recognition of the aminoacyl

identity of aa-tRNA by the *E. coli* ribosome [63]. Whilst they identified few differences, there was a

subtle 2-3 fold increase in A-site sampling by mischarged tRNAs. *E. coli*, unlike mycobacteria and

indeed most bacteria, does not routinely encounter mischarged tRNAs. Hence, this may suggest that

other model systems may be better suited to investigate bacterial ribosome recognition and

discrimination of mischarged tRNA. Our prior studies with the unusual aminoglycoside kasugamycin

[10] also suggest ribosomes can discriminate against mischarged tRNAs. Kasugamycin was known to

decrease ribosomal decoding errors [64, 65] in direct contrast with the increased errors witnessed with

other aminoglycosides such as streptomycin. We showed that kasugamycin, at concentrations that did

not inhibit protein synthesis, could decrease mistranslation from mischarged tRNAs not only in live

mycobacteria, but also in a cell-free *in vitro* translation system, which had been modified to include

excess mischarged Asp-tRNA$^{Asn}$ [10].

We do not know the precise molecular mechanism by which deletion of *gidB* increases ribosomal

discrimination of mischarged tRNAs. Our limited structural analysis suggests that   increased

ribosomal conformational changes   arising from mistranslation may impact the function of the

decoding centre in the absence of 16S rRNA methylation by GidB, although further higher resolution

structures are needed for definitive mechanistic analysis. What further changes to ribosomal

composition or function are required for lack of GidB-mediated methylation to effectively

discriminate against mischarged tRNAs? We and others have recently described alternative bacterial

ribosomes that are particularly adapted to stressful environments – such environments are also

associated with relaxation of translational fidelity [32, 66, 67]. However, multiple forms of alternative

ribosomes have been identified: comprising altered stoichiometry of ribosomal protein subunits,

variant subunits or alternative rRNA composition [68, 69]. Which, if any of these alternatives

contribute to increased fidelity in absence of GidB-mediated rRNA methylation will be the subject of

ongoing study.

Deletion of *gidB* causes increased mycobacterial susceptibility to rifampicin. Since lack of *gidB*

decreases both wobble misreading [42], as well as discrimination of mischarged tRNA, which

function is responsible? Several lines of evidence suggest discrimination of mischarged tRNAs is

more likely. First, we have previously demonstrated that increased discrimination of mischarged

tRNA by kasugamycin is responsible for enhanced mycobacterial susceptibility to rifampicin *in vitro*

[10]. Secondly, the magnitude of increased susceptibility of *ΔgidB* to rifampicin is greater in strain

HWS19, in which the excess error is derived solely from mischarged tRNA. Finally, the absolute

error rates of ribosomal decoding errors in mycobacteria, including wobble mistranslation, are much

lower than for errors due to mischarged tRNA [17, 32, 33], making the latter mechanism more

parsimonious.

Emerging data suggest absolute fidelity in protein synthesis is not desirable, not only from an

efficiency perspective, but also because relaxed fidelity is associated with adaptation to hostile

environments. Although moderate mistranslation rates may be adaptive under stress, excessive

mistranslation is still harmful. Therefore, mechanisms that are permissive for moderate but not

excessive mistranslation may serve an important function in bacterial environmental adaptation. Only

mycobacteria with increased mistranslation rates – either via mutation or from an environmental

context in which higher mistranslation rates are favoured – had a high fidelity phenotype in the

absence of *gidB*. This extended to discrimination of mischarged tRNAs that are not naturally

occurring, such as Ala-tRNA$_{CCA}^{Ala}$. Ribosomal RNA methylation may be reversible [70], although

definitive evidence is not available. An analogy of the function/role of this system may be with an

"automated braking system" in cars: i.e. it only comes into play at dangerous excessive speeds.

Similarly, lack of methylation by GidB only prevents 'runaway' mistranslation that would otherwise

result in error catastrophe, but allows for the potentially adaptive functions of translational error and is

potentially tunable.

## MATERIALS AND METHODS

**Bacterial strains and culture**

Wild-type *M. smegmatis* mc$^2$-155 [71] and its derivatives were cultured in Middlebrook 7H9 Broth supplemented with 0.2% glycerol, 10% ADS (albumin-dextrose-salt) and 0.05% Tween-80 with corresponding antibiotics. Wild-type *E. coli* HK295 and its derivatives were cultured in LB Broth with corresponding antibiotics. *E. coli* DH5*α* and TOP10 were used for transformation of plasmids and amplification. LB-Agar was used for both *M. smegmatis* and *E. coli* in plating experiments. If not otherwise mentioned, bacteria were grown in 37 ℃, 220 rpm shaker or maintained in 37℃ incubator.

**Suppressor screen**.

A total of around 1x10$^7$ colony forming units (C.F.U) HWS19 were plated onto each of 6 LB agar plates containing 1μg/mL streptomycin. After 5 days, visible colonies were picked for further analysis. The *rpsL* gene was amplified by PCR and sequenced, those with wild-type *rpsL* were transformed with Renilla-Firefly dual luciferase reporter plasmids (Renilla-Firefly WT and Renilla-Firefly D214N). Specific mistranslation rates of asparagine for aspartate were measured in potential suppressor candidates to identify low mistranslating strains compared with HWS19. Genomic DNA was isolated from HWS19 and four suppressor candidates by standard methods, and were then whole-genomes sequenced and analyzed by Genewiz. The whole genome sequence data have been deposited to the NCBI short-read archive <https://www.ncbi.nlm.nih.gov/sra> with accession number: PRJNA704242. Mapping results of mutations onto genes covering over 99% of all reads in four suppressor candidates are shown as Table S1. A further 10 suppressor candidates had only *gidB* sequenced as per Table S2.

**Deletion of *gidB* and complementation in *Mycobacterium smegmatis***

The 500 bp upstream and downstream regions of *gidB* were amplified from mc$^2$-155 genomic DNA. The zeocin resistance cassette was amplified from plasmid pKM-Zeo-Lox (A kind gift from Eric J. Rubin lab). Then the zeocin resistance cassette flanking 500 bp upstream and 500 bp downstream

region of *gidB* were assembled by overlapping extension PCR for use as an allele exchange substrate

(AES), which was verified by sequencing. WT mc$^2$-155 and HWS19 transformed with

pNIT(kan)::RecET::sacB recombineering plasmid were grown to $OD_{600}$~0.4, and expression of recET

was induced with 10 μM isovaleronitrile (IVN) for 5 h, then competent cells were made by standard

methods and transformed with 2 μg of the AES. The cells were recovered for 4 h and selected on LB

agar plates containing 20 μg/mL zeocin. The recombinants were confirmed by PCR for verifying the

zeocin cassette integration and the appropriate genomic context (Figure S3) (See verification primers

sequences in Table S3). *gidB* deletion strains were then cured of the recombinase plasmid prior to

further experiments.

For *gidB* complementation construction, several plasmids were constructed. The chromosome

integrating mycobacterial plasmid pML1357 (Addgene #32378) was used as the backbone vector. The

hygromycin resistance cassette was replaced with a kanamycin resistance cassette and streptomycin

resistance cassette as pKML1357 and pSML1357 respectively. *gidB* from wild-type *M. smegmatis*

with and without native promoter were amplified and cloned into integrative plasmids pKML1357/

pSML1357 to construct pKML1357-Pnative_*gidB*, pKML1357-Psmyc_*gidB*, pSML1357-Pnative-

*gidB*. Then the plasmids were transformed into Δ*gidB* strains and selected onto LB agar with

corresponding antibiotics for *gidB* complementation strains. Δ*gidB*::Pnative_*gidB* (KanR) strains were

used for dual luciferase mistranslation assay when measuring N to D and Q to E mistranslation rates.

Δ*gidB*::Psmyc_*MsmgidB* (KanR) were used for the dual-fluorescence mistranslation assay.

Δ*gidB*::Pnative_*gidB* (StrepR) strains were used for dual-luciferase mistranslation assays when

measuring W to A mistranslation as the pACET-Renilla-Firefly construct carries a kanamycin

resistance marker. In addition, *gidB* mRNA levels in *gidB* deletion and complementation strains were

verified by RT-qPCR (Figure S3).

**Measuring mistranslation rates with Renilla-Firefly dual-luciferase reporters**

For measuring mistranslation rates in *M. smegmatis*, the shuttle plasmids pTetG-Renilla-Firefly (WT),

pTetG-Renilla-D120N-Firefly, pTetG-Renilla-E144Q-Firefly under control of a tetracycline-inducible

promoter were used as previously [17]. For measurement of tryptophan to alanine mistranslation

rates, the mycobacterial chromosome integrating plasmid pACET-Renilla-Firefly, where expression

of the reporter is under control of an acetamide-inducible promoter was used [17]. The *Renilla* gene

was mutated by site-directed mutagenesis to Renilla-A214W.

In mycobacteria, the assay was performed as previously [12]. Briefly, strains with pTetG-

Renilla-Firely were cultured to OD600>3, and diluted to OD600~0.2 into fresh 7H9 medium

supplemented with 50μg/ml hygromycin and 100 ng /ml anhydrotetracycline for inducing dual

luciferase. After 7-8 hours, cells were collected by centrifugation at 12000rpm for 3min, lysed with

40μL passive lysis buffer in 96-well white flat bottom plate for 20-30min (plate shaking at 400rpm,

room temperature), then reacted with 50μL substrate Firefly reagents, 360rpm shaked 5s followed by

measuring the Firefly luminescence by Fluoroskan Ascent FL luminometer with 1000ms integration

time. Then 50μL Stop&Glo reagent was added followed by measuring the Renilla luminescence

immediately. For strains with pACET-Renilla-Firely and pTet-tRNA$_{CCA}$$^{Ala}$ [17], the strains in

experimental group were diluted to OD600~0.2 into fresh 7H9 medium supplemented with 50μg/ml

hygromycin, 20% acetamide and 100ng/mL anhydrotetracycline for inducing dual luciferase and

tRNA$_{CCA}$$^{Ala}$ respectively. The later procedures were same as above. The corrected Renilla/Firefly

representing the mistranslation level is calculated as follows: Corrected Renilla / Firefly (N to D) =

$\frac{Renilla(D120N)/Firefly}{Renilla(WT)/Firefly}$, Corrected Renilla / Firefly (Q to E) = $\frac{Renilla(E144Q)/Firefly}{Renilla(WT)/Firefly}$, Corrected Renilla /

Firefly (W to A)= $\frac{Renilla(A214W)/Firefly}{Renilla(WT)/Firefly}$.

**404  Measuring mistranslation rates with dual-fluorescence reporters**

The pUVtetOR-GLR (green-linker-red) reporter was adapted as previously [12] with some

modifications. Briefly, a mutated green fluorescent protein (E222Q) was fused to wild-type

monomeric red fluorescent protein (mRFP) with a flexible GGSGGGGSGGGSSGG linker. A C-

terminal tag (AANDENYAAAV) targeting the protein for proteolytic degradation [74] was added by

PCR.

The procedure of measuring relative mistranslation rates by flow cytometry was as previously

described [12] with some modifications. Strains with pUVtetOR-GLR were cultured to OD600~0.2

and then 100ng/ml ATc was added for inducing dual-fluorescence. After 3 hours, cells were diluted

into 1mL PBS to OD600~0.05. The GFP and RFP signals of the sample were collected by flow

cytometry (BD LSR Fortressa), being excited/ detected by 488nm/520nm laser line and

561nm/585nm laser line respectively. For the strains after high mistranslating selection: Strains

expressing pSML1357-*Aph*_D214N and pUVtetOR-GLR were spread onto LB agar containing

50μg/ml Hygromycin and 2μg/ml Kanamycin as experimental group and LB agar containing only

50μg/ml Hygromycin as control group. After 5 days, colonies were scraped from the plates and

cultured into 7H9 with 100ng/ml ATc for 3 hours. The following procedures were same as above. For

the strains under rifampicin condition: strains with pUVtetOR-GLR were spread onto LB agar no

selection plates and LB agar containing 20μg/ml rifampicin. After 5 days, colonies were then scraped

from the plates and cultured into 7H9 with 100ng/ml ATc for 3 hours. The following procedures were

same as above. Data were analyzed by FlowJo 10.4 for Windows 10. The gating strategy was as

follows: Live bacteria were gated to eliminate debris as P1 based on SSC-A and FSC-A, then

stringent gating on single cells was applied using a tandem gating strategy based on FSC-A and FSC-

H as P2, subsequently using SSC-A and SSC-H as P3. Single cells with positive red fluorescence

were acquired as P4 based on the red fluorescence signal. See Figure S4. Relative mistranslation rates

were analysed as a histogram of GFP/ RFP ratio as P5 . The mean values of GFP/RFP ratio were

calculated by FlowJo software (Flowjo 10.4 for Windows 10).

**Rifampicin time-kill curve assay**

Wild-type and strain HWS19 *M. smegmatis* were cultured in 7H9 medium until OD600~1. Cells were

diluted to OD600~0.3 into fresh 7H9 medium containing 20μg/ml rifampicin and cultured into 37°C,

220rpm shaker. Prior to addition of rifampicin, an aliquot was removed for calculation of cell

numbers at time zero. At indicated time points, aliquots were taken, washed 3x in PBS and

resuspended in PBS prior to being plated onto LB-agar. At least 3 10-fold dilutions were plated per

time point. After 3-5 days, visible colonies were counted. The counts at different time points were

normalized by counts at time zero for analysis.

**70S ribosome purification**

HWS19 and HWS19-Δ*gidB* strains were grown in Middlebrook 7H9 Broth supplemented with 0.2%

glycerol, 10% ADS (albumin-dextrose-salt) and 0.05% Tween-80 to OD 0.8. Bacteria pellet were

collected by centrifuge at 4000rpm at 4C for 10 minutes. Pellet was washed and resuspend with

polysome buffer (50 mM Tris-actate pH=7.2, 12 mM MgCl2, 50mMNH$_4$Cl, 1mM DTT). Bacteria

resuspension was lysed by MP FastPrep-24 bead beating system and the lysate was centrifuged at

12,000rpm 4C for 10 minutes. Cleared lysate was loaded on to 10% - 50% linear sucrose gradient

with Beckman SW40Ti rotor at a speed of 39, 000rpm at 4C for 4 hours. Gradient were fractioned

with continuous A260 measurement and the correspond 70S fraction were collected, buffer exchanged

for downstream cryo-EM studies.

**CryoEM Data Collection and Map Refinement**

Samples were diluted in milliQ ddH$_2$O and deposited onto glow-discharged (EMS-100 Glow

Discharge System, Electron Microscopy Sciences, 30 s at 15 mA and 22 mbar) copper Quantifoil

(Quantifoil Micro Tools) grids, 1.2/1.3 spacing, 300-mesh, coated in 2 nm amorphous carbon. Grids

were incubated for 30 s under 100% humidity and 10ºC, blotted once at force 3 with Whatman #1

filter paper, and plunge-frozen in liquid ethane using a FEI Vitrobot Mark IV (Thermo Fisher). Grids

were screened for ice quality on an FEI Talos Arctica (Thermo Fisher, 200 kV) or FEI Titan Krios

(Thermo Fisher, 300 kV) transmission electron microscope at UCSF. WT strain non-mutant samples

were shipped by dry shipper to the National Center for CryoEM Access and Training (NCCAT) at

NYSBC. These were imaged on an FEI Titan Krios (Thermo Fisher, 300 kV) electron microscope

with a Falcon IV camera using Leginon and without an energy filter. All other samples were imaged

at UCSF on an FEI Titan Krios (Thermo Fisher, 300 kV) electron microscope with a Gatan K3 direct

electron detector, using SerialEM and with an imaging filter (20 eV slit). Datasets at UCSF were

collected with fringe-free imaging using a nine-shot beam-image shift approach with coma

compensation. Further details are reported in Table 1. For all datasets, image stacks were binned by a

factor of 2, motion corrected, and dose weighted using UCSF MotionCor2. Contrast transfer function

(CTF) initial parameters were determined using CTFFIND4 in cisTEM (development version). After

excluding images with poor CTF fits or poor ice quality, dose-weighted micrographs were subjected

to unsupervised particle picking with a soft-edged disk template. Two-dimensional classification of

the resulting particle stacks was used to select the images carried forward. Ab initio reconstructions

were carried out for each dataset to produce initial references for three-dimensional reconstruction,

which proceeded through auto refinement and manual refinement in two to three classes as necessary

to exclude non-ribosome particles and 50S subunits. Final reconstructions were generated without

sharpening from the classes yielding the 70S ribosome in the latest (highest resolution) refinement

cycle prior to the emergence of overfitting artifacts in the Fourier shell correlation (FSC). Map

resolutions are reported as particle FSC at 0.143.

**Atomic Structure Refinement**

An initial atomistic model of the 70S ribosome was generated by fitting the 5o5j and 5o60 PDB

models of the small and large ribosomal subunits in Mycobacterium smegmatis, respectively, into the

final map of the HWS19 unmutated dataset using UCSF ChimeraX. This model was then subjected to

several iterations of rigid body fitting in UCSF ChimeraX or PyMOL, molecular dynamics model

fitting in ISOLDE, model building/trimming/correction in Coot or UCSF ChimeraX, and real-space

model refinement with anisotropic displacement parameter (ADP) refinement in Phenix, not

necessarily in that order, until convergence. Restraints for residues not distributed with Phenix

libraries were generated using Schrödinger and Phenix. The final HWS19 unmutated model was re-

fitted and re-refined into the other three maps, with further model trimming as needed where map

quality was not sufficient to resolve the structure. Figures were prepared using the PyMOL Molecular

Graphics System (Schrödinger).

**Statistical analysis**

All statistical methods are described in figures legends, presented as mean with SD. The p values and

statistical significance were calculated using GraphPad prism software (Prism 8 for Mac). Two-tailed

unpaired Student's t test was used to compare means between groups. ***$p<0.001$, **$p<0.01$,

*$p<0.05$, ns $p>0.05$.

**Author contributions**

HWS optimised the conditions of and performed the initial screen.  ZB performed most experiments

with assistance from JYH and YC. YXC purified the 70S ribosomes, interpreted data and modified all

figures.  ZB and YXC analysed the results. IY and JSF solved the cryo-EM structure. BJ conceived of

and supervised the study. ZB, YXC and BJ wrote the manuscript with input from the other authors.

**Acknowledgements**

We would like to thank the Tsinghua flow cytometry core for technical assistance. We would like to

thank Dr. Peter Walter at UCSF use of the ultracentrifuge and the fractionator in his lab. We

appreciate the members from the Javid Lab and Fraser Lab for discussion and feedback of this study.

This study was in part funded by grants from the Bill & Melinda Gates Foundation (OPP1109789)

and funds from Tsinghua University School of Medicine to BJ and a Wellcome Trust Investigator

award (207487/C/17/Z) to BJ (for work performed at UCSF).

**Table 1 Cryo-EM data collection, refinement and validation statistics**

| | WT strain WT (EMDB-44092) (PDB 9B1Y) | WT strain gidB mutant (EMDB-44097) (PDB 9B24) | HWS19 strain WT (EMDB-44090) (PDB 9B1W) | HWS19 strain gidB mutant (EMDB-44091) (PDB 9B1X) |
|---|---|---|---|---|
| **Data collection and processing** | | | | |
| Magnification | 96000 | 105000 | 130000 | 130000 |
| Voltage (kV) | 300 | 300 | 300 | 300 |
| Electron exposure (e–/Å$^2$) | 60.74 | 901.165 | 72.704 | 72.704 |
| Defocus range (μm) | 400-1400 | 500-1500 | 500-1500 | 500-1500 |
| Pixel size (Å) | 0.833 | 0.833 | 0.662 | 0.664 |
| Symmetry imposed | P1 | P1 | P1 | P1 |
| Initial particle images (no.) | 704502 | 1636851 | 254392 | 417089 |
| Final particle images (no.) | 591747 | 1466881 | 108034 | 136418 |
| Map resolution (Å) | 3.47 | 2.47 | 3.26 | 3.07 |
| FSC threshold | 0.143 | 0.143 | 0.143 | 0.143 |
| Map resolution range (Å) | inf-3.47 | inf-2.47 | inf-3.26 | inf-3.07 |
| | | | | |
| **Refinement** | | | | |
| Initial model used (PDB code) | 5o6o, 5o5j | 5o6o, 5o5j | 5o6o, 5o5j | 5o6o, 5o5j |
| Model resolution (Å) | 3.32 | 3.05 | 4.22 | 3.69 |
| FSC threshold | 0.143 | 0.143 | 0.143 | 0.143 |
| Model resolution range (Å) | inf-3.47 | inf-2.47 | inf-3.26 | inf-3.07 |
| Map sharpening $B$ factor (Å$^2$) | N/A | N/A | N/A | N/A |
| Model composition | | | | |
| Non-hydrogen atoms | 146981 | 146980 | 144760 | 144818 |
| Protein residues | 5723 | 5723 | 5877 | 5877 |
| Ligands | 629 | 629 | 629 | 629 |
| $B$ factors (Å$^2$) | | | | |
| Protein | 140.30 | 197.35 | 139.43 | 191.74 |
| Ligand | 142.41 | 162.55 | 171.92 | 154.91 |
| R.m.s. deviations | | | | |
| Bond lengths (Å) | 0.014 | 0.004 | 0.003 | 0.010 |
| Bond angles (°) | 0.893 | 0.670 | 0.541 | 0.650 |
| Validation | | | | |
| MolProbity score | 2.72 | 2.08 | 1.80 | 1.70 |
| Clashscore | 59.82 | 19.66 | 9.34 | 7.56 |
| Poor rotamers (%) | 0.86 | 0.13 | 0.40 | 0.10 |
| Ramachandran plot | | | | |
| Favored (%) | 92.44 | 95.78 | 95.56 | 95.82 |
| Allowed (%) | 7.47 | 4.17 | 4.36 | 4.11 |
| Disallowed (%) | 0.09 | 0.05 | 0.09 | 0.07 |

**References**

1.    Mohler, K. and M. Ibba, *Translational fidelity and mistranslation in the cellular response to*
*stress.* Nat Microbiol, 2017. **2**: p. 17117.
2.    Zaher, H.S. and R. Green, *Fidelity at the molecular level: lessons from protein synthesis.* Cell,
2009. **136**(4): p. 746-62.
3.    Ling, J., P. O'Donoghue, and D. Soll, *Genetic code flexibility in microorganisms: novel*
*mechanisms and impact on physiology.* Nat Rev Microbiol, 2015. **13**(11): p. 707-721.
4.    Fan, Y., et al., *Heterogeneity of Stop Codon Readthrough in Single Bacterial Cells and*
*Implications for Population Fitness.* Mol Cell, 2017. **67**(5): p. 826-836 e5.
5.    Schwartz, M.H. and T. Pan, *Function and origin of mistranslation in distinct cellular contexts.*
Crit Rev Biochem Mol Biol, 2017. **52**(2): p. 205-219.
6.    Ribas de Pouplana, L., et al., *Protein mistranslation: friend or foe?* Trends Biochem Sci, 2014.
**39**(8): p. 355-62.
7.    Drummond, D.A. and C.O. Wilke, *The evolutionary consequences of erroneous protein*
*synthesis.* Nat Rev Genet, 2009. **10**(10): p. 715-24.
8.    Reynolds, N.M., et al., *Cell-specific differences in the requirements for translation quality*
*control.* Proc Natl Acad Sci U S A, 2010. **107**(9): p. 4063-8.
9.    Evans, C.R., et al., *Errors during Gene Expression: Single-Cell Heterogeneity, Stress*
*Resistance, and Microbe-Host Interactions.* mBio, 2018. **9**(4).
10.   Chaudhuri, S., et al., *Kasugamycin potentiates rifampicin and limits emergence of resistance*
*in Mycobacterium tuberculosis by specifically decreasing mycobacterial mistranslation.* Elife,
2018. **7**.
11.   Rathnayake, U.M., W.N. Wood, and T.L. Hendrickson, *Indirect tRNA aminoacylation during*
*accurate translation and phenotypic mistranslation.* Curr Opin Chem Biol, 2017. **41**: p. 114-
122.
12.   Su, H.W., et al., *The essential mycobacterial amidotransferase GatCAB is a modulator of*
*specific translational fidelity.* Nat Microbiol, 2016. **1**(11): p. 16147.
13.   Schwartz, M.H., et al., *Global tRNA misacylation induced by anaerobiosis and antibiotic*
*exposure broadly increases stress resistance in Escherichia coli.* Nucleic Acids Res, 2016.
14.   Schwartz, M.H. and T. Pan, *Temperature dependent mistranslation in a hyperthermophile*
*adapts proteins to lower temperatures.* Nucleic Acids Res, 2016. **44**(1): p. 294-303.
15.   Fan, Y., et al., *Protein mistranslation protects bacteria against oxidative stress.* Nucleic Acids
Res, 2015. **43**(3): p. 1740-8.
16.   Lee, J.Y., et al., *Promiscuous methionyl-tRNA synthetase mediates adaptive mistranslation to*
*protect cells against oxidative stress.* J Cell Sci, 2014. **127**(Pt 19): p. 4234-45.
17.   Javid, B., et al., *Mycobacterial mistranslation is necessary and sufficient for rifampicin*
*phenotypic resistance.* Proc Natl Acad Sci U S A, 2014. **111**(3): p. 1132-7.
18.   Miranda, I., et al., *Candida albicans CUG mistranslation is a mechanism to create cell surface*
*variation.* MBio, 2013. **4**(4).
19.   Li, L., et al., *Naturally occurring aminoacyl-tRNA synthetases editing-domain mutations that*
*cause mistranslation in Mycoplasma parasites.* Proc Natl Acad Sci U S A, 2011. **108**(23): p.
9378-83.
20.   Netzer, N., et al., *Innate immune and chemically triggered oxidative stress modifies*
*translational fidelity.* Nature, 2009. **462**(7272): p. 522-6.
21.   Ling, J., et al., *Protein aggregation caused by aminoglycoside action is prevented by a*
*hydrogen peroxide scavenger.* Mol Cell, 2012. **48**(5): p. 713-22.
22.   Zhang, Y.W., et al., *HspX promotes the polar localization of mycobacterial protein*
*aggregates.* Sci Rep, 2019. **9**(1): p. 14571.
23.   Lee, J.W., et al., *Editing-defective tRNA synthetase causes protein misfolding and*
*neurodegeneration.* Nature, 2006. **443**(7107): p. 50-5.

24.    Liu, Y., et al., *Deficiencies in tRNA synthetase editing activity cause cardioproteinopathy.* Proc
Natl Acad Sci U S A, 2014. **111**(49): p. 17570-5.

25.    Kohanski, M.A., et al., *Mistranslation of membrane proteins and two-component system*
*activation trigger antibiotic-mediated cell death.* Cell, 2008. **135**(4): p. 679-90.

26.    Fan, Y., et al., *Optimal translational fidelity is critical for Salmonella virulence and host*
*interactions.* Nucleic Acids Res, 2019. **47**(10): p. 5356-5367.

27.    Rubio Gomez, M.A. and M. Ibba, *Aminoacyl-tRNA synthetases.* RNA, 2020. **26**(8): p. 910-936.

28.    Chong, Y.E., X.L. Yang, and P. Schimmel, *Natural homolog of tRNA synthetase editing domain*
*rescues conditional lethality caused by mistranslation.* J Biol Chem, 2008. **283**(44): p. 30073-
8.

29.    Vargas-Rodriguez, O. and K. Musier-Forsyth, *Exclusive use of trans-editing domains prevents*
*proline mistranslation.* J Biol Chem, 2013. **288**(20): p. 14391-9.

30.    LaRiviere, F.J., A.D. Wolfson, and O.C. Uhlenbeck, *Uniform binding of aminoacyl-tRNAs to*
*elongation factor Tu by thermodynamic compensation.* Science, 2001. **294**(5540): p. 165-8.

31.    Morse, J.C., et al., *Elongation factor-Tu can repetitively engage aminoacyl-tRNA within the*
*ribosome during the proofreading stage of tRNA selection.* Proceedings of the National
Academy of Sciences, 2020. **117**(7): p. 3610-3620.

32.    Chen, Y.X., et al., *Selective translation by alternative bacterial ribosomes.* Proc Natl Acad Sci
U S A, 2020. **117**(32): p. 19487-19496.

33.    Leng, T., et al., *Translational misreading in Mycobacterium smegmatis increases in stationary*
*phase.* Tuberculosis (Edinb), 2015. **95**(6): p. 678-81.

34.    Sheppard, K. and D. Soll, *On the evolution of the tRNA-dependent amidotransferases,*
*GatCAB and GatDE.* J Mol Biol, 2008. **377**(3): p. 831-44.

35.    Curnow, A.W., et al., *Glu-tRNAGln amidotransferase: a novel heterotrimeric enzyme required*
*for correct decoding of glutamine codons during translation.* Proc Natl Acad Sci U S A, 1997.
**94**(22): p. 11819-26.

36.    Cai, R.J., et al., *Forward Genetics Reveals a gatC-gatA Fusion Polypeptide Causes*
*Mistranslation and Rifampicin Tolerance in Mycobacterium smegmatis.* Front Microbiol,
2020. **11**: p. 577756.

37.    Li, Y.Y., et al., *Clinically Relevant Mutations of Mycobacterial GatCAB Inform Regulation of*
*Translational Fidelity.* mBio, 2021. **12**(4): p. e0110021.

38.    Ruan, B., et al., *Quality control despite mistranslation caused by an ambiguous genetic code.*
Proc Natl Acad Sci U S A, 2008. **105**(43): p. 16502-7.

39.    Kramer, E.B. and P.J. Farabaugh, *The frequency of translational misreading errors in E. coli is*
*largely determined by tRNA competition.* RNA, 2007. **13**(1): p. 87-96.

40.    Chen, Y.X., et al., *Measurement of Specific Mycobacterial Mistranslation Rates with Gain-of-*
*function Reporter Systems.* J Vis Exp, 2019(146).

41.    Okamoto, S., et al., *Loss of a conserved 7-methylguanosine modification in 16S rRNA confers*
*low-level streptomycin resistance in bacteria.* Mol Microbiol, 2007. **63**(4): p. 1096-106.

42.    Wong, S.Y., et al., *Functional role of methylation of G518 of the 16S rRNA 530 loop by GidB in*
*Mycobacterium tuberculosis.* Antimicrob Agents Chemother, 2013. **57**(12): p. 6311-8.

43.    Zhu, J.H., et al., *Rifampicin can induce antibiotic tolerance in mycobacteria via paradoxical*
*changes in rpoB transcription.* Nat Commun, 2018. **9**(1): p. 4218.

44.    Chong, Y.E., et al., *Distinct ways of G:U recognition by conserved tRNA binding motifs.* Proc
Natl Acad Sci U S A, 2018. **115**(29): p. 7527-7532.

45.    Hou, Y.M. and P. Schimmel, *A simple structural feature is a major determinant of the identity*
*of a transfer RNA.* Nature, 1988. **333**(6169): p. 140-5.

46.    Swairjo, M.A., et al., *Alanyl-tRNA synthetase crystal structure and design for acceptor-stem*
*recognition.* Mol Cell, 2004. **13**(6): p. 829-41.

47.    Wang, B.W., J.H. Zhu, and B. Javid, *Clinically relevant mutations in mycobacterial LepA cause*
*rifampicin-specific phenotypic resistance.* Sci Rep, 2020. **10**(1): p. 8402.

48.  Vijay, S., et al., *Most-probable number based minimum duration of killing assay for*
*determining the spectrum of rifampicin susceptibility in clinical M. tuberculosis isolates.*
Antimicrob Agents Chemother, 2020.

49.  Safi, H., et al., *Phase variation in Mycobacterium tuberculosis glpK produces transiently*
*heritable drug tolerance.* Proc Natl Acad Sci U S A, 2019. **116**(39): p. 19665-19674.

50.  Hicks, N.D., et al., *Clinically prevalent mutations in Mycobacterium tuberculosis alter*
*propionate metabolism and mediate multidrug tolerance.* Nat Microbiol, 2018. **3**(9): p. 1032-
1042.

51.  Vilcheze, C., et al., *Enhanced respiration prevents drug tolerance and drug resistance in*
*Mycobacterium tuberculosis.* Proc Natl Acad Sci U S A, 2017. **114**(17): p. 4495-4500.

52.  Gold, B. and C. Nathan, *Targeting Phenotypically Tolerant Mycobacterium tuberculosis.*
Microbiol Spectr, 2017. **5**(1).

53.  Richardson, K., et al., *Temporal and intrinsic factors of rifampicin tolerance in mycobacteria.*
Proc Natl Acad Sci U S A, 2016. **113**(29): p. 8302-7.

54.  Brauner, A., et al., *Distinguishing between resistance, tolerance and persistence to antibiotic*
*treatment.* Nat Rev Microbiol, 2016. **14**(5): p. 320-30.

55.  Abel Zur Wiesch, P., et al., *Classic reaction kinetics can explain complex patterns of antibiotic*
*action.* Sci Transl Med, 2015. **7**(287): p. 287ra73.

56.  Aldridge, B.B., I. Keren, and S.M. Fortune, *The Spectrum of Drug Susceptibility in*
*Mycobacteria.* Microbiol Spectr, 2014. **2**(5).

57.  Wakamoto, Y., et al., *Dynamic persistence of antibiotic-stressed mycobacteria.* Science, 2013.
**339**(6115): p. 91-5.

58.  Tollerson, R., 2nd and M. Ibba, *Translational regulation of environmental adaptation in*
*bacteria.* J Biol Chem, 2020. **295**(30): p. 10434-10445.

59.  Dale, T. and O.C. Uhlenbeck, *Amino acid specificity in translation.* Trends Biochem Sci, 2005.
**30**(12): p. 659-65.

60.  Dale, T. and O.C. Uhlenbeck, *Binding of misacylated tRNAs to the ribosomal A site.* RNA,
2005. **11**(11): p. 1610-5.

61.  Pape, T., W. Wintermeyer, and M.V. Rodnina, *Complete kinetic mechanism of elongation*
*factor Tu-dependent binding of aminoacyl-tRNA to the A site of the E. coli ribosome.* EMBO J,
1998. **17**(24): p. 7490-7.

62.  Zaher, H.S. and R. Green, *Quality control by the ribosome following peptide bond formation.*
Nature, 2009. **457**(7226): p. 161-6.

63.  Effraim, P.R., et al., *Natural amino acids do not require their native tRNAs for efficient*
*selection by the ribosome.* Nat Chem Biol, 2009. **5**(12): p. 947-53.

64.  Schuwirth, B.S., et al., *Structural analysis of kasugamycin inhibition of translation.* Nat Struct
Mol Biol, 2006. **13**(10): p. 879-86.

65.  van Buul, C.P., W. Visser, and P.H. van Knippenberg, *Increased translational fidelity caused*
*by the antibiotic kasugamycin and ribosomal ambiguity in mutants harbouring the ksgA*
*gene.* FEBS Lett, 1984. **177**(1): p. 119-24.

66.  Kurylo, C.M., et al., *Endogenous rRNA Sequence Variation Can Regulate Stress Response*
*Gene Expression and Phenotype.* Cell Rep, 2018. **25**(1): p. 236-248 e6.

67.  Parks, M.M., et al., *Variant ribosomal RNA alleles are conserved and exhibit tissue-specific*
*expression.* Sci Adv, 2018. **4**(2): p. eaao0665.

68.  Emmott, E., M. Jovanovic, and N. Slavov, *Ribosome Stoichiometry: From Form to Function.*
Trends Biochem Sci, 2019. **44**(2): p. 95-109.

69.  Dinman, J.D., *Pathways to Specialized Ribosomes: The Brussels Lecture.* J Mol Biol, 2016.
**428**(10 Pt B): p. 2186-94.

70.  Sloan, K.E., et al., *Tuning the ribosome: The influence of rRNA modification on eukaryotic*
*ribosome biogenesis and function.* RNA Biology, 2017. **14**(9): p. 1138-1152.

71. Snapper, S.B., et al., *Isolation and characterization of efficient plasmid transformation mutants of Mycobacterium smegmatis.* Mol Microbiol, 1990. **4**(11): p. 1911-9.
72. Kadokura, H. and J. Beckwith, *Four cysteines of the membrane protein DsbB act in concert to oxidize its substrate DsbA.* EMBO J, 2002. **21**(10): p. 2354-63.
73. Yao, Y., et al., *A Direct RNA-to-RNA Replication System for Enhanced Gene Expression in Bacteria.* ACS Synth Biol, 2019. **8**(5): p. 1067-1078.
74. Andersen, J.B., et al., *New unstable variants of green fluorescent protein for studies of transient gene expression in bacteria.* Appl Environ Microbiol, 1998. **64**(6): p. 2240-6.
678

# Figure.1

**Figure 1. Suppressor screen identifies gidB as a potential fidelity factor.**
**A.** Indirect aminoacylation pathway for tRNA$^{Asn}$ and tRNA$^{Gln}$ in mycobacteria. **B**. Design of the suppressor screen: applying an increasing mistranslation stress to both strains makes HWS.19 more susceptible to stress and more likely to have mutation in suppressors. **C.** Workflow of suppressor screen. **D**. N to D mistranslation rate of suppressor candidates compared with WT and HWS.19 measured by Renilla-Firefly dual luciferase reporter as depicted in Figure1C. **E.** *gidB* mutation was mapped onto 3 candidates by whole genome sequencing. (*P < 0.05, **P < 0.01, ***P < 0.001, Student *t* test)

# Figure.2

**Figure 2. Deletion of GidB in mycobacteria increases translation fidelity in high mistranslation mutant.**
**A.** N to D Mistranslation rate measured by gain-of-function Renilla-Firefly dual luciferase reporter in high mistranslation HWS19 strain (Left) and wild-type strain (Right). **B.** E to Q Mistranslation rate measured by gain-of-function dual fluorescent reporter in high mistranslation HWS19 strain (Left) and wild-type strain (Right).

# Figure.3

**Figure 3. Deletion of GidB in mycobacteria increases translation fidelity under high mistranslation physiological context.**
**A.** Schematic of measuring mistranslation rate using gain-of-function dual fluorescence reporter under normal and high mistranslation physiological context **B.** Q to E mistranslation of mycobacteria scraped from LB-agar or LB-agar with low dose rifampicin. **C.** Gating strategy to gate top/bottom 10% of bacteria with high/low mistranslation rate. **D.** Q to E mistranslation of different bacteria population (no selection) gated from the highest/lowest mistranslation rate as described in C. **E.** Q to E mistranslation of different bacteria population (Rif selection) gated from the highest/lowest mistranslation rate as described in C  (*P < 0.05, **P < 0.01, ***P < 0.001, Student *t* test)

# Figure.4

**Figure 4. Deletion of GidB decreases mistranslation mediated by non-physiological 'misacylated-tRNA'.**
**A.** The anticodon of an alanine-tRNA is mutated to tryptophan codon (EMAW) result in mistranslation from tryptophan to alanine (Right) to mimic non-physiological miacylated-tRNA compared to the two natural misacylated-tRNAs (Left) in mycobacteria. Mistranslation rate of tryptophan to alanine measured by Renilla-Firefly reporter (top) in high mistranslation HWS19 strain (**B**) and wild-type strain (**C**) with or without EMAW expression. (*P < 0.05, **P < 0.01, ***P < 0.001, Student *t* test)

# Figure.5

**Figure 5. Deletion of GidB decreases rifampicin tolerance.**
Rifampicin killing curve in high mistranslation background (A) and wild-type background (B). T.test was performed between *gidB* deletion strain and the other two strains respectively. (*P < 0.05, **P < 0.01, ***P < 0.001, Student *t* test)

# Figure.6

**Fig. 6. Structures of ribosomes suggest a role for GidB methylation in modifying contacts with Asn46.** A) The overall density map for the HWS19 WT ribosome is well resolved, permitting the assessment of conformation of the methylated state of G507. B) The overall density map of the HWS19 delta-gidB ribosome is also well resolved, indicating no major structural changes to the WT ribosome and permitting the assessment of the methylated state of G507. C) A zoom in of the site of GidB methylation (G507) reveals a potential contact between S12 Asn46 and G507. D) The unmethylated state created by delta-gidB lengthens the contact. The phenotypic recoding of S12 Asn46 to Asp in this background may disrupt this contact, leading to changes in translational fidelity.

# Figure.7

**Figure 7. Proposed model of *gidB* mediated translation fidelity control.**
Under low mistranslation context (Left), wild-type ribosome and ΔGidB both have moderate translation fidelity. Under high mistranslation context (Right) due to impaired indirect aminoacylation pathway, HWS19 ribosome has a loosen ribosome conformation compared to HWS19-ΔGidB ribosome, which translation is error-prone. HWS19-ΔGidB ribosome restores structure integrity could better discriminate against misacylated-tRNA.