Adaptive ecological processes and metabolic independence drive microbial colonization and resilience in the human gut

Andrea R. Watson[1,2], Jessika Füssel[2], Iva Veseli[4], Johanna Zaal DeLongchamp[3], Marisela Silva[3], Florian Trigodet[2], Karen Lolans[2], Alon Shaiber[4], Emily Fogarty[1,2], Christopher Quince[5], Michael K. Yu[6], Arda Söylev[7], Hilary G. Morrison[8], Sonny T.M. Lee[2], David T. Rubin[2], Bana Jabri[2], Thomas Louie[3], A. Murat Eren[1,2,4,8,*]

[1] Committee on Microbiology, The University of Chicago, Chicago, IL 60637, USA. [2] Department of Medicine, The University of Chicago, Chicago, IL 60637, USA. [3] Department of Medicine, The University of Calgary, Calgary, AB T2N 1N4, Canada. [4] Biophysical Sciences Program, The University of Chicago, Chicago, IL 60637, USA. [5] Organisms and Ecosystems, Earlham Institute, Norwich, Norwich, NR4 7UZ, United Kingdom. [6] Toyota Technological Institute at Chicago, Chicago, IL 60637, USA. [7] Department of Computer Engineering, Konya Food and Agriculture University, Konya, Turkey. [8] Josephine Bay Paul Center, Marine Biological Laboratory, Woods Hole, MA 02543, USA.

* Corresponding author: meren@uchicago.edu

Running Title

Drivers of human gut colonization

Keywords

fecal microbiota transplantation; human gut microbiome; microbial colonization; microbial metabolism; metabolic competency.

Abstract

A detailed understanding of gut microbial ecology is essential to engineer effective microbial therapeutics and to model microbial community assembly and succession in health and disease. However, establishing generalizable insights into the functional determinants of microbial fitness in the human gut has been a formidable challenge. Here we employ fecal microbiota transplantation (FMT) as an *in natura* experimental model to identify determinants of microbial colonization and resilience. Our long-term sampling strategy and high-resolution multi-omics analyses of FMT donors and recipients reveal adaptive ecological processes as the main driver of microbial colonization outcomes after FMT. We also show that high-fitness donor microbial populations are significantly enriched in metabolic pathways that are responsible for the biosynthesis of nucleotides, essential amino acids, and micronutrients, independent of taxonomy. To determine whether such metabolic competence can explain the microbial ecology of human disease states, we analyzed genomes reconstructed from healthy humans and humans with inflammatory bowel disease (IBD). Our data reveal that such traits are also significantly enriched in microbial genomes recovered from IBD patients, linking presence of superior metabolic competence in bacteria to their expansion in IBD. Overall, these findings suggest that the transfer of gut microbes from a healthy donor to a disrupted recipient environment initiates an environmental filter that selects for populations that can self-sustain. Such ecological processes that select for self-sustenance under stress offer a model to explain why common yet typically rare members of healthy gut environments can become dominant in inflammatory conditions without any need for them to be causally associated with, or contribute to, such disease states.

Introduction

The human gut microbiome is associated with a wide range of diseases and disorders (Almeida et al. 2020; Durack and Lynch 2019; Lynch and Pedersen 2016). However, mechanistic underpinnings of these associations have been difficult to resolve in part due to the diversity of human lifestyles (David et al. 2014) and the limited utility of model systems to make robust causal inferences for microbially mediated human diseases (Walter et al. 2020).

Inflammatory bowel disease (IBD), a group of increasingly common intestinal disorders that cause inflammation of the gastrointestinal tract (Baumgart and Carding 2007), has been a model to study human diseases associated with the gut microbiota (Schirmer et al. 2019). The pathogenesis of IBD is attributed in part to the gut microbiome (Plichta et al. 2019), yet the microbial ecology of IBD-associated dysbiosis remains a puzzle. Despite marked changes in gut microbial community composition in IBD (Ott et al. 2004; Sokol and Seksik 2010; Joossens et al. 2011), the microbiota associated with the disease lacks traditional pathogens (Chow, Tang, and Mazmanian 2011), and microbes that are found in IBD typically also occur in healthy individuals (Clooney et al. 2021), which complicates the search for robust functional or taxonomic markers of health and disease states (Lloyd-Price et al. 2019). One of the hallmarks of IBD is reduced microbial diversity during episodes of inflammation, when the gut environment is often dominated by microbes that typically occur in lower abundances prior to inflammation (Vineis et al. 2016). The sudden increase in the relative abundance of microbes that are common to healthy individuals suggests that the harsh conditions of IBD likely act as an ecological filter that prevents the persistence of low-fitness populations. Yet, in the absence of a complete understanding of the functional drivers of microbial colonization in this habitat, critical insights into the metabolic requirements of survival in IBD remains elusive.

Understanding the determinants of microbial colonization has been one of the fundamental aims of gut microbial ecology (Costello et al. 2012; Messer et al. 2017). To overcome the difficulties of conducting well-controlled studies with humans, researchers have studied the determinants of microbial colonization of the gut in model systems, such

as germ-free mice conventionalized with individual taxa (S. M. Lee et al. 2013) or a
consortium of human microbial isolates (Feng et al. 2020). Despite their utility for
hypothesis testing, simpler models do not capture the complex ecological interactions
fostered by natural systems and thus the insights they yield do not always translate to
human gut microbial ecology (Ley et al. 2006; Finucane et al. 2014). Between the
extremes of well-controlled but simple mouse models and complex yet uncontrolled
human populations, there exists a middleground that provides a window into the microbial
ecology of complex human systems through a controlled perturbation: human fecal
microbiota transplantation (FMT), the transfer of stool from a donor into a recipient's
gastrointestinal tract (Eiseman et al. 1958).

FMT complements laboratory models of environmental perturbation by colliding two
distinct microbial ecosystems, and thus offers a powerful framework to study fundamental
questions of microbial ecology, including the determinants of microbial succession and
resilience (Schmidt, Raes, and Bork 2018). Here we use FMT as an *in natura*
experimental model to investigate the ecological and functional determinants of
successful microbial colonization of the human gut at the level of individual populations.
Our findings suggest that adaptive ecological forces are key drivers of colonization
outcomes after FMT, reveal taxonomy-independent metabolic determinants of fitness in
the human gut, and demonstrate that similar ecological principles determine resilience of
microbes in stressful and inflammatory conditions.

Results and Discussion

Our study includes 109 gut metagenomes (Supplementary Table 1) from two healthy FMT
donors (A and B) and 10 FMT recipients (five recipients per donor) who had multiply
recurrent *Clostridium difficile* infection (CDI) and received vancomycin for a minimum of
10 days to attain resolution of diarrheal illness prior to FMT. On the last day of vancomycin
treatment, a baseline fecal sample was collected from each recipient, and their bowel
contents were evacuated immediately prior to FMT. Recipients did not take any antibiotics
on the day of transplant, or during the post-FMT sampling period (Supplementary Figure
1). We also collected 24 Donor A samples over a period of 636 days and 15 Donor B

samples over a period of 532 days to establish an understanding of the long-term
microbial population dynamics within each donor microbiota. We also collected 5 to 9
samples from each recipient up to 336 days post-FMT. Deep sequencing of donor and
recipient metagenomes using Illumina paired-end (2x150) technology resulted in a total
of 7.7 billion sequences with an average of 71 million reads per metagenome (Figure 1,
Supplementary Table 1, Supplementary Table 2). We employed genome-resolved
metagenomics, pangenomics, and microbial population genetics for an in-depth
characterization of donor and recipient gut microbiota using these data, and we leveraged
publicly available gut metagenomes to benchmark our observations.

Many but not all donor microbes colonized recipients and persisted long-term

We first characterized the taxonomic composition of each donor and recipient sample by
aligning metagenomic short reads to reference genomes in the NCBI's RefSeq database
(Supplementary Table 2). The phylum-level microbial community composition of both
donors reflected those observed in healthy individuals in North America (Human
Microbiome Project Consortium 2012): a large representation of Firmicutes and
Bacteroidetes, and other taxa with relatively lower relative abundances, including
Actinobacteria, Verrucomicrobia, and Proteobacteria (Figure 1, Supplementary Table 2).
In contrast, the vast majority of the recipient pre-FMT samples were dominated by
Proteobacteria, a phylum that typically undergoes a drastic expansion in individuals
treated with vancomycin (Isaac et al. 2017). After the FMT, we observed a dramatic shift
in recipient taxonomic profiles (Supplementary Table 2, Supplementary Figure 2), a
widely documented hallmark of this procedure (Khoruts et al. 2010; Grehan et al. 2010;
Shahinas et al. 2012). Nearly all recipient samples post-FMT were dominated by
Bacteroidetes and Firmicutes as well as Actinobacteria and Verrucomicrobia in lower
abundances, resembling qualitatively, but not quantitatively, the taxonomic profiles of
their donors (Supplementary Table 2). For example, even though the median relative
abundance of Bacteroidetes populations were 5% and 17% in donors A and B, their
relative abundance in recipients post-FMT increased to 33% and 45%, respectively

(Figure 1, Supplementary Table 2). A single genus, *Bacteroides*, made up 76% and 82%

of the Bacteroidetes populations in the recipients of Donor A and B, respectively

(Supplementary Table 2). The success of the donor *Bacteroides* populations in recipients

upon FMT is not surprising given the ubiquity of this genus across human populations

throughout the globe (Wexler and Goodman 2017) and the ability of its members to

survive substantial levels of stress (Swidsinski et al. 2005; Vineis et al. 2016). This result

suggests that FMT outcomes in our dataset are unlikely random, and the study design

and resulting dataset offers a framework to study ecological principles of the human gut

microbiome.

Next, we assembled short metagenomic reads into contiguous segments of DNA

(contigs). Co-assemblies of 24 Donor A and 15 Donor B metagenomes independently

resulted in 53,891 and 54,311 contigs that were longer than 2,500 nucleotides, and

described 0.70 and 0.79 million genes occurring in 179 and 248 genomes, as estimated

by the mode of the frequency of bacterial single-copy core genes (Supplementary Table

2). One way to characterize how well a given assembly describes the DNA content of a

given metagenome is to calculate the percentage of reads it recruits from the

metagenome through read mapping. Donor contigs recruited on average 80.8% of

metagenomic reads from donor metagenomes. In contrast, they recruited 43.4% of reads

on average from pre-FMT recipient metagenomes. This number increased to 80.2% for

recipient metagenomes post-FMT (Figure 1), and the donor contigs continued to

represent 76.8% of the recipient metagenomes on average even after a year post-FMT

(Supplementary Table 2). These read recruitment results suggest that members of the

donor microbiota successfully established in recipient guts upon FMT and largely

persisted until the end of the sampling period.

Compared to metagenomic short reads, assembled contigs provide a larger genetic

context to study microbial metagenomes. However, a sole focus on contigs may yield

misleading results (Kowarsky et al. 2017) that can be ameliorated by reconstructing

microbial genomes from metagenomic assemblies (Chen et al. 2020). We reconstructed

genomes from co-assembled donor metagenomes by grouping contigs into putative bins

based on sequence composition and differential coverage signal as previously described

(Sharon et al. 2013; S. T. M. Lee et al. 2017). We retained bins that were at least 70%
complete and had no more than 10% redundancy as predicted by bacterial single-copy
core genes (Bowers et al. 2017; Chen et al. 2020) and manually refined them to improve
their quality following previously described approaches (Delmont et al. 2018; Shaiber et
al. 2020). Our binning resulted in a final list of 128 metagenome-assembled genomes
(MAGs) for Donor A and 183 MAGs for Donor B that included members of Firmicutes
(n=265), Bacteroidetes (n=20), Actinobacteria (n=14), Proteobacteria (n=7),
Verrucomicrobia (n=2), Cyanobacteria (n=2), and Patescibacteria (n=1) (Supplementary
Table 3). The taxonomy of donor-derived genomes largely reflected the taxonomic
composition of donor metagenomes as predicted by short reads (Figure 1,
Supplementary Table 2, Supplementary Table 3). While only 20 genomes (mostly of
Bacteroides and *Alistipes*) explained the entirety of the Bacteroidetes group, we
recovered 265 MAGs that represented lower abundance but diverse populations of
Firmicutes (Figure 1, Supplementary Table 2, Supplementary Table 3). We found no
difference between the delivery method of FMT for the recipients of donor A, where, on
average 45% and 43% of donor genomes emerged in recipients who received donor stool
through colonoscopy (n=3) versus pills (n=2), respectively. However, there was an
increase in the efficiency of pills for donor B, where on average 25% and 54% of donor
genomes emerged in recipients who received donor stool through colonoscopy (n=2)
versus pill (n=3) (Supplementary Figure 3).

Reconstructing genomes gave us access to microbial populations in metagenomes
through metagenomic read recruitment strategies and enabled us to characterize (1)
population-level microbial colonization dynamics before and after FMT using donor and
recipient metagenomes and (2) the distribution of each donor population across
geographically distributed humans using 1,984 publicly available human gut
metagenomes (Supplementary Table 4). As expected, we detected each donor
population in at least one donor metagenome (see Methods for 'detection' criteria). Yet,
only 16% of Donor A populations were detected in every Donor A sample, and only 44%
of Donor B MAGs were detected in every Donor B sample (Figure 1, Supplementary Table
3), in agreement with the previously documented dynamism of gut microbial community
composition over time (David et al. 2014). A marked increase in the detection of donor

populations in recipients after FMT echoed the general pattern of transfer suggested by the short-read taxonomy (Figure 1): while only 38% of Donor A and 54% of Donor B populations were detected in at least one recipient pre-FMT, these percentages increased to 96% and 96% post-FMT (Supplementary Table 3). Not every donor population colonized each recipient, but colonization events did not appear to be random: while some donor populations colonized all recipients, others colonized none (Figure 1), providing us with an opportunity to resolve colonization events and quantify colonization success for each donor population in our dataset.

Figure 1. FMT Donor genomes across recipients and publicly available gut metagenomes. In both heat maps each column represents a donor genome and each row represents a metagenome, and each data point represents the detection of a given genome in a given metagenome. Purple rows represent donor metagenomes which cover 636 days for Donor A and 532 days for Donor B. Each recipient metagenome is colored red for pre-FMT samples and blue for post-FMT samples. The three rightmost columns display for each metagenome (X) the number of metagenomic short reads in millions, (Y) the percent of metagenomic short reads recruited by genomes, and (Z) the taxonomic composition of metagenomes (based on metagenomic short reads) at the phylum level. The row Q provides the phylum-level taxonomy for each donor genome. Finally, the 11 bottom rows under each heat map show the fraction of healthy adult metagenomes from 11 different countries in which a given donor genome is detected (if a genome is detected in every individual from a country it is represented with a full bar). The dendrograms on the right-hand side of these layers organize countries based on the detection patterns of genomes (Euclidean distance and Ward clustering). Red and green shades represent the two main clusters that emerge from this analysis, where green layers are industrialized countries in which donor genomes are highly prevalent and red layers are less industrialized countries where the prevalence of donor genomes is low.

Resolving colonization events accurately is a challenging task as multiple factors may influence the ability to determine colonization outcomes unambiguously. These factors include (1) the inability to detect low-abundance populations, (2) inaccurate characterization of transient populations observed immediately after FMT as successful colonization events, (3) the reliance on relative abundance of populations to define colonization events when abundance estimates from stool do not always reflect the abundance of organisms in the GI tract (Yasuda et al. 2015; Sheth et al. 2019), and (4) the failure to distinguish between colonization by a donor population or emergence of a pre-FMT recipient population after FMT (where a low-abundance recipient population that is closely related to one or more donor populations becomes abundant after FMT and is mistaken as a bona fide colonization event). To mitigate these factors, we have (1) employed deep-sequencing of our metagenomes which averaged 71 million reads per sample, (2) implemented a longitudinal sampling strategy, that spanned 376 days on average, to observe donor populations in our recipients long after the FMT, (3) leveraged a 'detection' metric to define colonization events by presence/absence of populations rather than abundance, and (4) employed microbial population genetics to identify and resolve origins of subpopulations. We also developed an analytical approach (Supplementary Figure 4) to determine whether a given donor population has colonized a given recipient based on the detection of donor subpopulations in the transplant sample, in the recipient pre-FMT, and in the recipient post-FMT (see Materials and Methods,

Supplementary Table 5). To determine colonization outcomes, we analyzed 640

genome/recipient pairs for Donor A (128 donor genomes in 5 recipients) and identified 99

successful colonization events, 38 failed colonization events, and 503 ambiguous

colonization events (Supplementary Table 6). For Donor B, we analyzed 915

genome/recipient pairs (183 donor genomes in 5 recipients) and identified 106 successful

colonization events, 109 failed colonization events, and 700 ambiguous colonization

events (Supplementary Table 6). Our stringent criteria (see Materials and Methods,

Supplementary Figure 4) classified the vast majority of all genome/recipient pairs as

ambiguous colonization events. Nevertheless, due to the relatively large number of donor

MAGs and FMT recipients in our study, we were left with 352 MAG/recipient pairs with

unambiguous phenotypes for downstream analyses.

Adaptive ecological forces are the primary drivers of microbial colonization

The ability of a microbial population to colonize and persist in a complex ecosystem is

influenced by both neutral and adaptive forces (Maignien et al. 2014). Although which of

these is the major driver of successful colonization of the human gut remains unclear

(Smillie et al. 2018). In the context of FMT, previous studies have suggested neutral

processes to determine colonization success based on the abundance of a microbial

population in a donor stool sample (Smillie et al. 2018; Podlesny and Florian Fricke 2020).

Indeed, ecological drift may have a significant role in a system dominated by neutral

processes, where low-abundance donor populations in the transplant would be less likely

to be observed in recipients. In contrast, if the system is dominated by adaptive forces,

colonization success would be a function of the population fitness in the recipient

environment, rather than its abundance in the transplant.

To investigate the impact of neutral versus adaptive processes on colonization in our

dataset we first asked whether the prevalence of a donor population in healthy human gut

metagenomes, which we define here as a measure of its fitness, was associated with the

detection of the same population in donor or recipient metagenomes. Within both FMT

cohorts, the mean detection of each population in recipients post-FMT had a stronger

association with population fitness than mean detection in donor samples (Figure 2a).
The fitness of donor A populations explained 4.2% of the variation in mean detection of
those populations in donor samples (R^2=0.042, p=0.021) and 19% of variation in mean
detection in recipient post-FMT samples (R^2=0.19, p=2.7e-07), an increase of
approximately 4.5-fold (Figure 2a). Similarly, Donor B population fitness explained 7.3%
of the variation in mean detection in donor samples (R^2=0.073, p=2.1e-04), and 36% of
the variation in mean detection in recipient post-FMT samples (R^2=0.36, p=4.5e-19), an
increase of approximately 5-fold (Figure 2a). This suggests that fitness is a better
predictor of colonization outcome than it is of the detection of a population in the donor,
suggesting that adaptive forces are likely at play. But detecting a donor population in a
recipient post-FMT metagenome through metagenomic read recruitment does not prove
colonization, since donor genomes can recruit reads from recipient populations that are
closely related (i.e., strain variants) and that were low abundance prior to FMT. Single-
nucleotide variants in read recruitment results, however, can reveal such cases (Denef
2019) and quantify their dynamics (Quince et al. 2017). Thus, we developed an improved
model that took into consideration the presence and absence of distinct subpopulations
in our data and their origins (Supplementary Figure 4). We then used this model to test if
colonization success was correlated with population fitness or population dose, which we
define here as the relative abundance of a given population in the transplanted donor
stool sample. For Donor A populations, colonization outcome was significantly correlated
with both dose (Wald test, AUC=0.73, p=7.7e-05) and fitness (Wald test, AUC=0.76,
p=6.3e-06) (Figure 2b,c). But combining both measures as predictive variables did not
substantially improve the performance of our colonization model (AUC=0.82) (Figure 2c).
This was likely due to the small, but significant, correlation between dose and fitness in
Donor A MAGs (R^2=0.053, p=0.0070) (Figure 2d). When the fitness of a microbial
population is reflected in its relative abundance, the effect of fitness on colonization
outcome may be masked by an apparent dose effect. In contrast to Donor A, the fitness
of Donor B populations and their relative abundance in Donor B samples were not
correlated (R^2=0.0012, p=0.61) (Figure 2d), providing us with an ideal case to analyze
these two factors independently. Indeed, there was no correlation between dose of a
microbial population in Donor B transplant samples and colonization outcome in

recipients post-FMT (Wald test, AUC=0.56, p=0.09). Instead, we found a significant
correlation between the fitness of each population and the colonization outcome (Wald
test, AUC=0.70, p=9.0e-07) (Figure 2c).

Taken together, our findings suggest that fitness of a microbial population as measured
by its prevalence across global gut metagenomes can predict its colonization success
better than its abundance in the donor stool sample, giving credence to the role of
adaptive rather than neutral ecological processes in colonization. This finding contrasts
with previous studies which suggested that the abundance of a given population in the
donor sample was an important determinant of colonization (Smillie et al. 2018; Podlesny
and Florian Fricke 2020). However, these analyses included many recipient samples
collected less than one week after FMT and it is likely that their observations were
influenced by the presence of transient populations. Indeed, samples collected
immediately after FMT are more likely to inflate the number of colonization events,
whereas longitudinal sampling over a longer time course can distinguish transient
populations from those that successfully colonized the recipients. We cannot definitively
test this hypothesis as we sampled most of our recipients a week after FMT. Still, on
average 12% of the donor populations detected in our recipients a week after FMT were
no longer detected after a month (Figure 1, Supplementary Table 3). Overall, our stringent
criteria to determine colonization outcome and the extended post-FMT sampling period
likely enabled us to study the long-term engraftment of successful and potentially low-
abundance colonizers, instead of high-abundance transient populations that may be
dominant directly after FMT.

Figure 2. Relationships between dose, prevalence, and colonization outcome. Left: Donor A. Right: Donor B. a)
Linear regression models of mean detection of each MAG in either donor or recipient post-FMT samples as a function
of prevalence. b) Colonization outcome of MAG/recipient pairs as a function of MAG dose or MAG prevalence.
Significance calculated by Wald test. c) Receiver operator curves (ROCs) for logistic regression models of colonization.
d) Linear regression models of dose as a function of prevalence.

Accurately distinguishing the role of dose versus fitness in colonization success is further
compounded by the fact that microbial populations that are prevalent across human
populations may also tend to be more abundant. This is well illustrated by Donor A.
Fortunately, the abundant populations in Donor B did not reflect prevalent microbes in
healthy adult guts, which demonstrated the importance of fitness as a determinant of
colonization success compared to dose without the confounding effect of a correlation
between fitness and dose. Thus, it is a theoretical possibility that colonization success is
purely driven by adaptive forces and is not influenced by dose, at all. However, while our
data assign a larger role to adaptive forces with confidence, a more accurate
determination of the proportional influence of adaptive versus neutral processes in
colonization requires a much larger dataset.

Colonizer and resilient microbes are enriched in metabolic pathways for the biosynthesis of essential organic compounds

Fitness in a specific environment is conferred to an organism by a combination of
functional traits. In the human gut, such traits drive microbial community succession and
structure as a response to changing host diet and lifestyle (Koenig et al. 2011; Rothschild
et al. 2018). Behind successful colonization and resilience after perturbation are likely
similar functional traits that promote fitness. Building on our observation that suggests a
primary role of adaptive ecological processes in colonization outcome, we next sought to
identify genetic determinants of colonization. For this, we leveraged our access to donor
microbial population genomes and global metagenomes to investigate whether a
functional enrichment analysis could reveal predictors of success independent of
taxonomy.

To generate metabolic insights into colonization success we divided our donor
populations into 'high-fitness' and 'low-fitness' groups by considering both their
prevalence in FMT recipients and prevalence across global gut metagenomes (Materials
and Methods). The 'high-fitness' group included the microbial populations that colonized
or persisted in all FMT recipients and were the most prevalent in gut metagenomes from
Canada. We assumed that they represented a set of highly fit microbial populations as

(1) they were able to colonize human gut environments systematically, (2) they persisted
in these environments long-term regardless of the host genetics or lifestyle, and (3) they
were prevalent in gut metagenomes outside of our study. In comparison, the 'low-fitness'
group comprised microbial populations that failed to colonize or persist in at least three
FMT recipients. These populations were nevertheless viable gut microbes as not only our
long-term sampling of the donors systematically identified them but also, they sporadically
colonized some FMT recipients. Yet, unlike those in the high-fitness group, the distribution
patterns of low-fitness populations were sparse, not only within our cohort, but also within
publicly available metagenomes. In fact, low-fitness genomes were on average less
prevalent than high-fitness genomes in each of the 17 different countries we queried
(Supplementary Table 3). Overall, we conservatively categorized 20 populations in each
group for downstream analyses (Supplementary Table 7). All populations in the low-
fitness group resolved to Firmicutes. The high-fitness group was also dominated by
Firmicutes (15 of 20) but it also included five Bacteroidetes and one Actinobacteria
(Supplementary Table 7). Genome completion estimates did not differ between high and
low-fitness groups (Wilcoxon rank sum test, p=0.42) and averaged to 91% and 93%,
respectively. However, genome sizes between the two groups differed dramatically
(p=2.9e-06), where high-fitness group genomes averaged to 2.8 Mbp while low-fitness
group genomes averaged to 1.6 Mbp. These results suggest that the length difference
between genomes in high and low-fitness groups is likely to have biological relevance.
Indeed, we found a very high correspondence between the lengths of our MAGs and their
best matching reference genomes in the GTDB (r=0.88, p=5e-14) (Supplementary Table
7).

Our metabolic enrichment analysis revealed 33 KEGG pathway modules, each containing
genes that form a functional unit in a metabolic pathway. Every module that was enriched
differentially between these two groups were enriched in the high-fitness group. The lack
of any enriched modules in the low-fitness group is in line with the reduction in genome
lengths in the low-fitness group and further suggests that the reduction is associated with
the absence of metabolic modules. Of all enriched modules, 79% were modules related
to biosynthesis, which indicates an overrepresentation of biosynthetic capabilities in the
high-fitness group as KEGG modules for biosynthesis only make up 55% of all KEGG

modules (Figure 3, Supplementary Table 7). Of the 33 enriched modules, 48.5% were

associated with amino acid metabolism, 21.2% with vitamin and cofactor metabolism,

18.2% with carbohydrate metabolism, 6% with lipid metabolism and 3% with energy

metabolism (Supplementary Table 7). Metabolic modules that were enriched in the high-

fitness group included the biosynthesis of seven of the nine essential amino acids,

indicating the importance of metabolic competency to synthesize high-demand

compounds as a factor increasing fitness in colonizing new gut environments

(Supplementary Table 7). This is further supported by the enrichment of biosynthesis

pathways for the essential cofactor vitamin B12 (cobalamin), which occurred in 67.5% of

the high-fitness populations and only 12.5% of the low-fitness group (Supplementary

Table 7). Vitamin B12 is structurally highly complex and costly to produce, requiring

expression of more than 30 genes that are exclusively encoded by bacteria and archaea

(Martens et al. 2002). Thus, the competitive advantages conferred by metabolic

autonomy appear to outweigh the additional costs. In addition to the biosynthesis of

tetrahydrofolate, riboflavin, and cobalamin, the high-fitness group had a larger

representation of biosynthetic modules for vitamins including biotin, pantothenate, folate,

and thiamine (Supplementary Table 7), micronutrients that are equally important in

bacterial and human metabolism and are shown to play important roles in mediating host-

microbe interactions (Biesalski 2016). Interestingly, enriched metabolic modules in our

analysis partially overlap with those that Feng et al. identified as the determinants of

microbial fitness using metatranscriptomics and a germ-free mouse model

conventionalized with microbial isolates of human origin (Feng et al. 2020).

Even though enriched metabolic modules occurred mostly in high-fitness populations, we

did find some of these modules in the low-fitness group as well (Supplementary Table 7),

but their distribution was not uniform as they primarily occurred only in a subset of

genomes that resolved to Firmicutes (Figure 3). We then sought to identify whether the

levels of completion of these modules that occurred in both groups were identical. For

this, we matched six low-fitness genomes that encoded modules enriched in high-fitness

group genomes to six high-fitness genomes from the same phylum (marked as HF and

LF subgroups in Figure 3). Bacterial single-copy core genes estimated that genomes in

both subgroups were highly complete with a slight increase in average completion of low-

fitness genomes (93.7%) compared to high-fitness genomes (90.1%). Despite the higher
estimated genome completion for low-fitness populations, estimated metabolic module
completion values were significantly lower in the low-fitness group (Wilcoxon rank sum
test with continuity correction, V=958, p=5e-09) (Figure 3, Supplementary Table 7). This
indicates that even when modules that are associated with high-fitness were detected in
low-fitness genomes, they were systematically missing genes and were less complete
than the same modules in high-fitness genomes.

Figure 3. Distribution of metabolic modules across low and high-fitness genomes. Each data point in this
heat map shows the level of completion of a given metabolic module (rows) in a given genome (columns).
The box-plot on the right-side compares a subset of low-fitness (LF) and high-fitness (HF) genomes, where
each data point is the level of completion of a given metabolic module in a genome and shows a statistically
significant difference between the overall completion of metabolic modules between these subgroups
(Wilcoxon rank sum test, p=5.4e-09).

While gut microbial ecosystems of healthy individuals include both low- and high-fitness microbes, IBD primarily selects for high-fitness populations

436
437
438

Our results so far show that while the healthy donor environment could support both high-
fitness and low-fitness populations (Figure 1, Supplementary Table 3), challenging
microbes to colonize a new environment or withstand massive ecosystem perturbation
during FMT selects for high-fitness populations (Figure 3, Supplementary Table 7),
suggesting that metabolic competence is a more critical determinant of fitness during
stress than during homeostasis. Based on these observations, it is conceivable to
hypothesize that (1) a healthy gut environment will support a range of microbial
populations with a wide spectrum of metabolic competency, and (2) a gut environment
under stress will select for high metabolic competency in microbial populations.

To test these hypotheses, we compared genomes reconstructed from a cohort of healthy
individuals (Pasolli et al. 2019) to genomes reconstructed from a cohort of pouchitis
patients (Vineis et al. 2016), a form of IBD with similar pathology to ulcerative colitis (De
Preter et al. 2009). We conservatively excluded genomes that were less than 70%
complete from the Pasolli et al. dataset to avoid underestimating pathway completion in
genomes from healthy people. In contrast, we included any genomes from the Vineis et
al. dataset if they were more than 50% complete. The number of genomes per individual
and their level of completeness were similar between IBD and healthy individuals: we
compared 44 genomes from 4 IBD patients with an average completion of 89.2% and 264
genomes from 22 healthy individuals with an average completion of 90.4%
(Supplementary Table 8). Intriguingly, similar to the length differences between genomes
of high-fitness and low-fitness populations (2.8 Mbp versus 1.6 Mbp on average),
microbial populations associated with IBD patients had larger genomes compared to
healthy people and averaged to 3.3 Mbp versus 2.6 Mbp, respectively (Supplementary
Table 8). This suggests that despite the slightly higher completion of microbial genomes
from the healthy cohort, these genomes tended to be smaller in size than the genomes
reconstructed from IBD patients.

Next, we asked whether the completion of those metabolic modules associated with
colonization success and resilience differed between the genomes reconstructed from
healthy and IBD individuals. The level of completion of the 33 metabolic modules were
almost identical between high-fitness genomes and genomes from IBD patients
(Wilcoxon rank sum test, $p=0.5$), but they were significantly less complete in microbial
genomes from healthy individuals (Wilcoxon rank sum test, $p=5.39e-08$) (Figure 4,
Supplementary Table 8). Metabolic modules with the largest differences in completion
between genomes from healthy and IBD individuals included biosynthesis of cobalamin,
arginine, ornithine, tryptophan, isoleucine as well as the Shikimate pathway (Figure 4,
Supplementary Table 8), a seven step metabolic route bacteria use for the biosynthesis
of aromatic amino acids (phenylalanine, tyrosine, and tryptophan) (Herrmann and Weaver
1999).

Our findings show that the same set of key metabolic modules that distinguish high-fitness
and low-fitness populations in FMT were also differentially associated with populations
that occurred in healthy individuals compared to IBD patients. In particular, while healthy
individuals seem to harbor microbes with a broad range of metabolic competency, IBD
individuals appear to be enriched with organisms with metabolic autonomy. It is
conceivable that a stable gut microbial ecosystem is more likely to support low-fitness
populations through metabolic cross-feeding, where vitamins, amino acids, and
nucleotides are exchanged between microbes (D'Souza et al. 2018). In contrast, host-
mediated environmental stress in IBD likely disrupts such interactions and creates an
ecological filter that selects for metabolic competence, which subsequently leads to loss
of diversity and the dominance of organisms with large genomes that are not necessarily
abundant in states of homeostasis.

Figure 4. Distribution of metabolic modules in genomes reconstructed from healthy individuals and
individuals with IBD. The top panel shows the metabolic module completion values for (1) high- and (2) low-
fitness donor genomes identified in this study (blue and orange), (3) genomes from healthy individuals
(green), and (4) genomes from individuals with IBD. Next to group averages, shown the distribution of
metabolic modules for each individual. Each dot in a given box-plot represents one of 33 metabolic modules
that were enriched in high-fitness FMT donor populations and the y-axis indicates its estimated completion.
In the bottom panel the completion values for 10 of the 33 pathways shown as ridge-line plots.

These observations have implications for the defining features of healthy gut environments from an ecological point of view. Defining the 'healthy gut microbiome' has been a major goal of human gut microbiome research (Bäckhed et al. 2012), and remains elusive (Eisenstein 2020). Despite comprehensive investigations that considered core microbial taxa (Arumugam et al. 2011; Lloyd-Price, Abu-Ali, and Huttenhower 2016) or guilds of microbes that represent coherent functional groups (Wu et al. 2021), the search for 'biomarkers' of healthy microbiomes is ongoing (McBurney et al. 2019). Given our data we hypothesize that one of the defining features of a healthy gut environment is its ability to support a diverse community of microbes with a broad spectrum of metabolic competence, where both low-fitness and high-fitness populations live in a coherent ecosystem. Conversely, an enrichment of metabolically competent high-fitness populations would likely indicate the presence of environmental stress. Our analyses demonstrate that this is a quantifiable feature of microbial communities through genome-

resolved metagenomic surveys, although testing this hypothesis rigorously requires a
larger number of genomes reconstructed from a larger number of individuals diagnosed
with a broader set of IBDs. Our analyses have other limitations. For instance, metabolic
insights in our study have been limited to genomic potential and have considered only
well-known metabolic pathways, which, given the extent of the unknown coding space in
microbial genomes (Vanni et al. 2020), are likely far from complete. As a result, the
disproportional enrichment of biosynthetic modules in high-fitness genomes indicates that
the ability to synthesize essential biological compounds is necessary but likely insufficient
to survive environmental stress in the gut. Nevertheless, the finding that the same
metabolic modules that promote colonization success after FMT are also the hallmarks
of fitness in IBD suggests the presence of ecological principles that are shared between
these systems and warrants deeper investigation.

522 Subtle differences in key functions distinguish populations of the 523 same genus with differential colonization success

While adaptive processes that favor metabolic independence explain the determinants of
colonization and resilience for distantly related taxa, metabolic features that promote high-
fitness at this broad level may not explain differences in fitness between more closely
related taxa, such as distinct species within a single genus, which are likely to have similar
metabolic capabilities (Martiny, Treseder, and Pusch 2013) due to unifying ecological
traits in higher ranks of taxonomy (Philippot et al. 2010). We finally investigated whether
we could identify determinants of fitness across metabolically similar populations with
different levels of success in colonizing unrelated individuals.

Members of the genus *Bifidobacterium* have long been used as probiotics (Gomes and
Malcata 1999) and are prevalent occupants of the healthy human gut microbiota
(Arboleya et al. 2016). In our dataset, *Bifidobacterium* was the second most abundant
genus (14.1%) after *Bacteroides* (15.8%) in Donor A, from whom we reconstructed three
MAGs over 98% completion that resolved to three distinct species in this genus: *B.
longum*, *B. adolescentis subsp. adolescentis*, and *B. animalis subsp. lactis*
(Supplementary Table 3). While each of these *Bifidobacterium* populations occurred in

Donor A metagenomes in a relatively stable fashion, they showed vastly different
colonization efficiency upon FMT (Figure 5), enabling us to investigate determinants of
colonization among closely related taxa.

In contrast to the *B. longum* and *B. adolescentis subsp. adolescentis* (henceforth *B.*
adolescentis) populations that colonized most recipients, *B. animalis subsp. lactis*
(henceforth *B. lactis*) did not seem to have colonized any of our recipients (Supplementary
Table 3). Overall, we were able to detect *B. longum*, *B. adolescentis*, and *B. lactis*
populations in 83%, 75%, and 4% of all post-FMT recipient metagenomes, respectively
(Figure 5). Most strikingly, patterns of colonization that emerged from the analysis of FMT
recipients reflected those seen in publicly available gut metagenomes from Canada,
where *B. longum*, *B. adolescentis*, and *B. lactis* populations occurred in 74%, 39%, and
13% of the population, demonstrating a positive relationship (Pearson's correlation of 0.9,
n.s.) between the colonization efficiency upon FMT and the fitness of these populations.
Furthermore, the gut metagenomes from 17 countries confirmed the substantially
reduced fitness of *B. lactis* globally (Supplementary Table 9). Interestingly, the *B. lactis*
MAG we reconstructed from Donor A was virtually identical (with over 99.99% sequence
identity over 99.82% alignment, Supplementary Table 9) to most *B. lactis* strains that are
widely used as probiotics (Jungersen et al. 2014), revealing a disagreement between the
preferences of commercial microbial therapeutics and human gut microbial ecology.

Figure 5. Characteristics of three *Bifidobacterium species*. Top panel shows the distribution of Donor A MAGs that
represent three distinct Bifidobacterium populations across donor and recipient metagenomes before and after FMT.
The last two columns in this panel show the prevalence of these populations in post-FMT metagenomes, and publicly
available gut metagenomes from Canada. The panel below displays the distribution of KEGG orthologs across the
three Bifidobacterium MAGs along with 31 high-quality isolate genomes from the NCBI. Each item shown in concentric
circles represents a single function assigned by the database of KEGG Orthologs, and each layer is a distinct genome.
The intensity of color indicates the presence of a given function in a given genome. The most outer circle marks groups
of functions that are enriched in various groups of Bifidobacterium genomes as well as those functions that are not
enriched in any group as they are either in all genomes, or only a very small number of them.

To identify factors that may explain the differences in colonization success between *B. longum, B. adolescentis*, and *B. lactis*, we created a collection of *Bifidobacterium* genomes that, in addition to the three metagenome-assembled genomes we reconstructed, included 31 complete genomes obtained from the NCBI (within-group and across-groups average gANI of 98.9% and 77.3%, respectively) (Supplementary Table 9). All three groups of *Bifidobacterium* genomes encoded the majority of the metabolic pathways associated with the high-fitness group (63% ± 5%). However, missing pathways were not uniformly distributed across three: *B. lactis* lacked the largest fraction of these pathways (42%) compared to the more prevalent *B. adolescentis* (36%) and *B. longum* (33%) (Supplementary Table 9). *B. longum* and *B. adolescentis* carried the complete tetrahydrofolate (vitamin B9) biosynthesis pathway in agreement with previous metabolic descriptions of *Bifidobacterium* (D'Aimmo et al. 2012; Sugahara et al. 2015) which

qualifies this group as attractive probiotics (Strozzi and Mogna 2008; Pompei et al. 2007),

but this pathway was absent in *B. lactis* genomes. We also found that *B. longum* and *B.*

adolescentis genomes encoded histidine and nicotinamide adenine dinucleotide (NAD)

biosynthesis which *B. lactis* lacked (Supplementary Table 9). Finally, the average genome

lengths of *B. longum* (2.31 Mbp) and *B. adolescentis* (2.18 Mbp) were longer than the

average genome length of *B. lactis* (1.94 Mbp), which reflects the pattern we observed

previously where high-fitness populations tended to have larger genomes. In summary,

even though all *Bifidobacterium* genomes in our pangenome had a higher metabolic

overlap with one another compared to high-fitness and low-fitness genomes we have

previously studied, the reduced fitness of *B. lactis* compared to *B. longum* and *B.*

adolescentis could still be explained by the absence of a small number of metabolic

competencies associated with the high-fitness group genomes.

Next, we focused on the enrichment of individual functions across the three groups of

genomes using gene annotations from KOfam profiles (Aramaki et al. 2020) from the

Kyoto Encyclopedia of Genes and Genomes (KEGG) (M. Kanehisa and Goto 2000) and

Clusters of Orthologous Groups (COGs) from the NCBI (Galperin et al. 2021). Of all 954

unique KOfams found in our *Bifidobacterium* pangenome, 272 functions were not

common to all genomes but statistically enriched in either one or two groups. Our analysis

of these accessory functions showed that *B. longum* encoded 150 (55.5%), *B.*

adolescentis encoded 115 (42.3%), and *B. lactis* encoded 95 (34.9%) of all accessory

functions that were statistically enriched (Figure 5, Supplementary Table 9). The same

analysis with 1,286 unique COGs confirmed these observations: of all 353 COGs

enriched in any group, *B. longum* encoded 212 (60.1%), *B. adolescentis* encoded 172

(48.7%), and *B. lactis* encoded 118 (33.4%) (Supplementary Table 9). Overall, these

results reveal a striking correlation between the number of accessory functions

associated with *B. longum, B. adolescentis*, and *B. lactis*, and echo the absence of

metabolic pathways in *B. lactis* even at the level of accessory gene functions, explaining

their differential ability to colonize new individuals and distribution in global human gut

metagenomes.

We finally investigated the contents of the differentially occurring accessory functions to
speculate on whether they could be related to differences in fitness. For instance, in
contrast to all *B. longum* and *B. adolescentis* in the *Bifidobacterium* pangenome, none of
the *B. lactis* genomes encoded a phosphoenolpyruvate phosphotransferase (PEP-PTS)
system specific for the uptake of β-glucoside (Supplementary Table 9). As the genus
Bifidobacterium is characterized by a large array of genes associated with carbohydrate
uptake and metabolism (Ventura et al. 2009; Schell et al. 2002; Kleerebezem and
Vaughan 2009), *B. lactis* represents a notable exception with a lower number of genes
associated with carbohydrate metabolism, fewer genes encoding carbohydrate-specific
ABC transporters, and the absence of phosphoenolpyruvate-phosphotransferase (PEP-
PTS) systems (Barrangou et al. 2009). The absence of any other PEP-PTS system in *B.*
longum and *B. adolescentis* may indicate the catabolic niche occupied by these
populations in the human gut that is shaped by their extensive capacity for uptake and
metabolism of plant derived glycosides (Chien, Huang, and Chou 2006; Schell et al.
2002). Additional functions that exclusively occurred in *B. adolescentis* and *B. longum*
genomes included two multidrug resistance pumps of the 'multidrug and toxin extrusion'
(MATE) type, three transporters of the major facilitator superfamily (MFS) involved in bile
acid tolerance and macrolide efflux, two bile acid:natrium ion symporters, and one
proton/chloride ion antiporter conferring acid tolerance (Supplementary Table 9). The
drug defense mechanisms may act to protect these populations during periods of
inflammation and drug administration, but may also be beneficial with regard to the
common ingestion of antibiotics through various food products (Kirchhelle 2018). These
results show that in the microbial fitness landscape of the human gut, where the
determinants of success across distantly related taxa are primarily defined by the
presence or absence of a large number of metabolic pathways, there exists smaller
niches equally accessible to closely related organisms with similar metabolic potential,
among which success can be speculated by subtle differences in key functions.

Conclusions

Our study points to adaptive ecological processes as primary determinants of both long-term colonization after FMT and microbial fitness in the human gut environment through metabolic competency as conferred by biosynthesis of nucleotides, amino acids, and essential micronutrients. Even when we found these metabolic modules in low-fitness populations, they were systematically less complete compared to their high-fitness counterparts.

Our findings suggest that in a healthy gut environment high- and low-fitness populations co-occur in harmony, with their differential fitness indiscernible by taxonomy or relative abundance. However, transfer to a new gut environment through FMT, or host-mediated stress through IBD, initiates an ecological filter that selects for high-fitness populations that can self-sustain. This model offers a null hypothesis to explain how low-abundance members of healthy gut environments can come to dominate the gut microbiota under stressful conditions, while not being causally associated with disease states. If the association between particular microbial taxa and disease is solely driven by their superior metabolic competence, microbial therapies that aim to treat complex diseases by adding microbes associated with healthy individuals will be unlikely to compete with the adaptive processes that regulate complex gut microbial ecosystems.

Materials and Methods

Sample collection and storage. We used a subset of individuals who participated in a randomized clinical trial (Kao et al. 2017) and conducted a longitudinal FMT study of two human cohorts (DA and DB), each consisting of one FMT donor and 5 FMT recipients of that donor's stool. All recipients received vancomycin for a minimum of 10 days pre-FMT at a dose of 125 mg four times daily. Three DA and two DB recipients received FMT via pill, and two DA and three DB recipients received FMT via colonoscopy. All recipients had recurrent *C. difficile* infection before FMT, and two DA recipients and 1 DB recipient were also diagnosed with ulcerative colitis (UC). 24 stool samples were collected from the DA donor over a period of 636 days, and 15 stool samples were collected from the DB donor over a period of 532 days. Between 5 and 9 stool samples were collected from each recipient over periods of 187 to 404 days, with at least one sample collected pre-FMT and 2 samples collected post-FMT. This gave us a total of 109 stool samples from all donors and recipients. Samples were stored at -80°C. (Supplementary Figure 1, Supplementary Table 1)

Metagenomic short-read sequencing. We extracted the genomic DNA from frozen samples according to the centrifugation protocol outlined in MoBio PowerSoil kit with the following modifications: cell lysis was performed using a GenoGrinder to physically lyse the samples in the MoBio Bead Plates and Solution (5–10 min). After final precipitation, the DNA samples were resuspended in TE buffer and stored at −20 °C until further analysis. Sample DNA concentrations were determined by PicoGreen assay. DNA was sheared to ~400 bp using the Covaris S2 acoustic platform and libraries were constructed using the Nugen Ovation Ultralow kit. The products were visualized on an Agilent Tapestation 4200 and size-selected using BluePippin (Sage Biosciences). The final library pool was quantified with the Kapa Biosystems qPCR protocol and sequenced on the Illumina NextSeq500 in a 2 × 150 paired-end sequencing run using dedicated read indexing.

'Omics workflows. Whenever applicable, we automated and scaled our 'omics analyses using the bioinformatics workflows implemented by the program `anvi-run-workflow`

(Shaiber et al. 2020) in anvi'o (Eren et al. 2015, 2021). Anvi'o workflows implement
numerous steps of bioinformatics tasks including short-read quality filtering, assembly,
gene calling, functional annotation, hidden Markov model search, metagenomic read-
recruitment, metagenomic binning, pangenomics, and phylogenomics. Workflows use
Snakemake (Köster and Rahmann 2012) and a tutorial is available at the URL
http://merenlab.org/anvio-workflows/. The following sections detail these steps.

Taxonomic composition of metagenomes based on short reads. We used Kraken2
v2.0.8-beta (Wood, Lu, and Langmead 2019) with the NCBI's RefSeq bacterial, archaeal,
viral and viral neighbours genome databases to calculate the taxonomic composition
within short-read metagenomes.

Assembly of metagenomic short and long reads. To minimize the impact of random
sequencing errors in our downstream analyses, we used the program `iu-filter-quality-
minoche` to process short metagenomic reads, which is implemented in illumina-utils
v2.11 (Eren et al. 2013) and removes low-quality reads according to the criteria outlined
by Minoche et al. (Minoche, Dohm, and Himmelbauer 2011). IDBA_UD v1.1.2 (Peng et
al. 2012) assembled quality-filtered short reads into longer contiguous sequences
(contigs), although we needed to recompile IDBA_UD with a modified header file so it
could process 150bp paired-end reads. For the assembly of long-reads and identification
of circular contigs we used Flye v2.6 (Kolmogorov et al. 2019) with the `--meta` argument
followed by Pilon v1.23 (Walker et al. 2014) correction using the high-quality short-reads.

Processing of contigs. We use the following strategies to process both sequences we
obtained from our assemblies and those we obtained from reference genomes. Briefly,
we used (1) `anvi-gen-contigs-database` on contigs to compute k-mer frequencies and
identify open reading frames (ORFs) using Prodigal v2.6.3 (Hyatt et al. 2010), (2) `anvi-
run-hmms` to identify sets of bacterial (Campbell et al. 2013) and archaeal (Rinke et al.
2013) single-copy core genes using HMMER v3.2.1 (Eddy 2011), (3) `anvi-run-ncbi-cogs`
to annotate ORFs with functions from the NCBI's Clusters of Orthologous Groups (COGs)
(Tatusov et al. 2003), and (4) `anvi-run-kegg-kofams` to annotate ORFs with functions
from the KOfam HMM database of KEGG orthologs (KOs) (Aramaki et al. 2020; M.
Kanehisa and Goto 2000). To predict the approximate number of genomes in

metagenomic assemblies we used the program `anvi-display-contigs-stats`, which
calculates the mode of the frequency of single-copy core genes as described previously
(Delmont and Eren 2016).

Metagenomic read recruitment, reconstructing genomes from metagenomes,
determination of genome taxonomy and ANI. We recruited metagenomic short reads
to contigs using Bowtie2 v2.3.5 (Langmead and Salzberg 2012) and converted resulting
SAM files to BAM files using samtools v1.9 (Li et al. 2009). We profiled the resulting BAM
files using the program `anvi-profile` with the flag `--min-contig-length` set to 2500 to
eliminate shorter sequences to minimize noise. Once we have read recruitment results
from each metagenome is profiled to store contig coverages into single anvi'o profile
databases, `anvi-merge` combined all profiles into an anvi'o merged profile for
downstream visualization, binning, and statistical analyses. We then used `anvi-cluster-
contigs` to group contigs into 100 initial bins using CONCOCT v1.1.0 (Alneberg et al.
2014), `anvi-refine` to manually curate initial bins with conflation error based on
tetranucleotide frequency and differential coverage signal across all samples, and `anvi-
summarize` to report final summary statistics for each gene, contig, and bin. We used the
program `anvi-rename-bins` to identify bins that were more than 70% complete and less
than 10% redundant, and store them in a new collection as metagenome-assembled
genomes (MAG), discarding lower quality bins from downstream analyses. GTBD-tk
v0.3.2 (Chaumeil et al. 2019) assigned taxonomy to each of our MAG using GTDB r89
(Parks et al. 2018), but to assign species- and subspecies-level taxonomy for
`DA_MAG_00057`, `DA_MAG_00011`, `DA_MAG_00052` and `DA_MAG_00018`, we
used `anvi-get-sequences-for-hmm-hits` to recover DNA sequences for bacterial single-
copy core genes that encode ribosomal proteins, and searched them in the NCBI's
nucleotide collection (nt) database using BLAST (Altschul et al. 1990). Finally, the
program `anvi-compute-genome-similarity` calculated pairwise genomic average
nucleotide identity (gANI) of our genomes using PyANI v0.2.9 (Pritchard et al. 2016).

Criteria for MAG detection in metagenomes. Using mean coverage to assess the
occurrence of populations in a given sample based on metagenomic read recruitment can
yield misleading insights since this strategy cannot accurately distinguish reference

sequences that represent very low-abundance environmental populations from those

sequences that do not represent an environmental population in a sample yet still recruit

reads from non-target populations due to the presence of conserved genomic regions.

Thus we relied upon the 'detection' metric, which is a measure of the proportion of the

nucleotides in a given sequence that are covered by at least one short read, and

considered a population was detected in a metagenome if anvi'o reported a detection

value of at least 0.25 for its genome (whether it was a metagenome-assembled or isolate

genome). Values of detection in metagenomic read recruitment results often follow a

bimodal distribution for populations that are present and absent (see Supplementary

Figure 2 in (Utter et al. 2020)), thus 0.25 is an appropriate cutoff to eliminate false-positive

signal in read recruitment results for populations that are absent.

Identification of MAGs that represent multiple subpopulations. To identify

subpopulations of MAGs in metagenomes, we used the anvi'o command `anvi-gen-

variability-profile` with the `--quince-mode` flag which exported single-nucleotide variant

(SNV) information for all MAGs after read recruitment. We then used DESMAN v2.1.1

(Quince et al. 2017) to analyze SNVs to determine the number and distribution of

subpopulations represented by a single genome. To account for non-specific mapping

that can inflate the number of estimated subpopulations, we removed any subpopulation

that made up less than 1% of the entire population explained by a single MAG. To account

for noise due to low-coverage, we only investigated subpopulation for MAGs for which

the mean non-outlier coverage of single-copy core genes was at least 10X.

Criteria for colonization of a recipient by a MAG. We developed a method to determine

whether or not a MAG successfully colonized a recipient, and applied this method to each

MAG and each recipient within a cohort. In order to confidently assign colonization or

non-colonization phenotypes to each MAG/recipient pair, we required that the MAG be

detected in the donor sample used for transplant into the recipient. If these criteria were

met, we then determined whether the MAG was detected in any post-FMT recipient

sample taken more than 7 days after transplant. If not, the MAG/recipient pair was

considered a non-colonization event. If the MAG was detected in the recipient greater

than7 days post-FMT, we used subpopulation information to determine if any

subpopulation present in the donor and absent in the recipient pre-FMT was detected in
the recipient more than 7 days post-FMT. If this was the case, we considered this to
represent a colonization event. See Supplementary Figure 4 for a complete outline of all
possible cases.

Determination of dose and fitness for MAGs. We defined population dose as the
second and third quartile mean coverage of a population in the transplanted stool sample.
We defined fitness as the prevalence of a population in 23 healthy adult gut metagenomes
(see Materials and Methods: Criteria for MAG detection in metagenomes) from Canada,
the same country in which the FMTs were performed.

Regression analysis. To examine the association between dose and/or prevalence with
colonization outcome, we built binomial logistic regression models using the R stats `glm`
function. We used the R stats `predict` function and the R pROC `roc` function to evaluate
our models by creating receiver operating characteristic (ROC) curves and calculating the
area under the ROC curve (AUC). To determine the correlation between dose and
prevalence, we performed linear regression using the R stats `lm` function. We used the
R tidyverse package, including ggplot2, to visualize boxplots, scatterplots, and ROC
curves.

Pangenomic analysis and gANI. We used anvi'o to compute and visualize pangenomes
of MAGs and reference genomes. We stored all processed MAG and reference genome
contigs (see Contig processing methods section) in an anvi'o database using the
command `anvi-gen-genomes-storage`. To create the pangenomes, we then passed that
database to the command `anvi-pan-genome` which used NCBI's BLAST (Altschul et al.
1990) to quantify gene similarity within and between genomes and the Markov Cluster
algorithm (MCL) (Enright, Van Dongen, and Ouzounis 2002) to cluster groups of similar
genes. We set the `anvi-pan-genome` `--min-occurrence` flag to 2 to remove gene
clusters only present in one genome (singletons), and visualized pangenomes using
`anvi-display-pan`.

Phylogenomic tree construction. To concatenate and align amino acid sequences of
46 single-copy core (Campbell et al. 2013) ribosomal proteins that were present in all of

our *Bifidobacterium* MAGs and reference genomes, we ran the anvi'o command `anvi-
get-sequences-for-hmm-hits` with the `--return-best-hit`, `--get-aa-sequence` and `--
concatenate` flags, and the `--align-with` flag set to `muscle` to use MUSCLE v3.8.1551
(Edgar 2004) for alignment. We then ran `anvi-gen-phylogenomic-tree` with default
parameters to compute a phylogenomic tree using FastTree 2.1 (Price, Dehal, and Arkin
2010).

Analysis of metabolic modules and enrichment. We calculated the level of
completeness for a given KEGG module (Minoru Kanehisa et al. 2014, 2017) in our
genomes using the program `anvi-estimate-metabolism`, which leveraged previous
annotation of genes with KEGG orthologs (KOs) (see the section 'Processing of contigs').
Then, the program `anvi-compute-functional-enrichment` determined whether a given
metabolic module was enriched in based on the output from `anvi-estimate-metabolism`.
The URL https://merenlab.org/m/anvi-estimate-metabolism serves a tutorial for this
program which details the modes of usage and output file formats. The statistical
approach for enrichment analysis is defined elsewhere (Shaiber et al. 2020), but briefly it
computes enrichment scores for functions (or metabolic modules) within groups by fitting
a binomial generalized linear model (GLM) to the occurrence of each function or complete
metabolic module in each group, and then computing a Rao test statistic, uncorrected p-
values, and corrected q-values. We considered any function or metabolic module with a
q-value less than 0.05 to be 'enriched' in its associated group if it was also at least 75%
complete and in at least 50% of the group members. To display the distribution of
individual KEGG orthologs across genomes and order them based on their enrichment
scores and group affiliations we used the program `anvi-display-functions`.

Determination of high-fitness and low-fitness MAGs for metabolic enrichment
analysis. We classified MAGs as high-fitness if, in all 5 recipients, they were detected in
the donor sample used for transplantation as well as the recipient more than 7 days post-
FMT. We classified low-fitness MAGs as those that, in at least 3 recipients, were detected
in the donor sample used for FMT but were not detected in the recipient at least 7 days
post-FMT. We reduced the number of high-fitness MAGs to be the same as the number

of low-fitness MAGs for metabolic enrichment analysis by selecting only the high-fitness

MAGs which were the most prevalent in the Canadian gut metagenomes.

Ordination plots. We used the R vegan v2.4-2 package `metaMDS` function to perform

nonmetric multidimensional scaling (NMDS) with Horn-Morisita dissimilarity distance to

compare taxonomic composition between donor, recipient, and global metagenomes. We

visualized ordination plots using R ggplot2.

Data Availability

841   Raw sequencing data for donor and recipient metagenomes are stored under the NCBI

842   BioProject PRJNA701961 (see Supplementary Table 1 for accession numbers).

843   Supplementary tables are accessible via doi:10.6084/m9.figshare.14138405.

Acknowledgements

We thank Mitchell L. Sogin, Eugene B. Chang, and Howard Shuman for helpful

discussions, Ryan Moore (0000-0003-3337-8184) and Ozcan C. Esen (0000-0001-7337-

671X) for technical help. We also thank Kaiyu Wu, Robyn Louie and Linda Ward of the

IPC Research Laboratory at the University of Calgary for their help with patient

recruitment and sampling. This project was supported by the GI Research Foundation

(GIRF) and the Mutchnik Family Fund.

Author Contributions

AME, TL, BJ conceived the study. JZD, MS, TL recruited patients, performed

transplantation experiments, and collected samples. ARW, JF, AME performed primary

data analyses. IV developed research tools. KL, STML, HGM performed sample

processing and sequencing. FT, AS, EF, CQ, MKY, AY contributed to data analyses and

interpretation. DTR, BJ, TL, and AME directed research. ARW, JF, AME wrote the paper

with critical input from all authors.

Competing Interests

Authors declare no competing financial interests.

References

Almeida, Cátia, Rita Oliveira, Raquel Soares, and Pedro Barata. 2020. "Influence of Gut Microbiota Dysbiosis on Brain Function: A Systematic Review." *Porto Biomedical Journal* 5 (2). https://doi.org/10.1097/j.pbj.0000000000000059.

Alneberg, Johannes, Brynjar Smári Bjarnason, Ino de Bruijn, Melanie Schirmer, Joshua Quick, Umer Z. Ijaz, Leo Lahti, Nicholas J. Loman, Anders F. Andersson, and Christopher Quince. 2014. "Binning Metagenomic Contigs by Coverage and Composition." *Nature Methods* 11 (11): 1144–46.

Altschul, S. F., W. Gish, W. Miller, E. W. Myers, and D. J. Lipman. 1990. "Basic Local Alignment Search Tool." *Journal of Molecular Biology* 215 (3): 403–10.

Aramaki, Takuya, Romain Blanc-Mathieu, Hisashi Endo, Koichi Ohkubo, Minoru Kanehisa, Susumu Goto, and Hiroyuki Ogata. 2020. "KofamKOALA: KEGG Ortholog Assignment Based on Profile HMM and Adaptive Score Threshold." *Bioinformatics* 36 (7): 2251–52.

Arboleya, Silvia, Claire Watkins, Catherine Stanton, and R. Paul Ross. 2016. "Gut Bifidobacteria Populations in Human Health and Aging." *Frontiers in Microbiology* 7 (August): 1204.

Arumugam, Manimozhiyan, Jeroen Raes, Eric Pelletier, Denis Le Paslier, Takuji Yamada, Daniel R. Mende, Gabriel R. Fernandes, et al. 2011. "Enterotypes of the Human Gut Microbiome." *Nature* 473 (7346): 174–80.

Bäckhed, Fredrik, Claire M. Fraser, Yehuda Ringel, Mary Ellen Sanders, R. Balfour Sartor, Philip M. Sherman, James Versalovic, Vincent Young, and B. Brett Finlay. 2012. "Defining a Healthy Human Gut Microbiome: Current Concepts, Future Directions, and Clinical Applications." *Cell Host & Microbe* 12 (5): 611–22.

Barrangou, Rodolphe, Elizabeth P. Briczinski, Lindsay L. Traeger, Joseph R. Loquasto, Melissa Richards, Philippe Horvath, Anne-Claire Coûté-Monvoisin, et al. 2009. "Comparison of the Complete Genome Sequences of Bifidobacterium Animalis Subsp. Lactis DSM 10140 and Bl-04." *Journal of Bacteriology* 191 (13): 4144–51.

Baumgart, Daniel C., and Simon R. Carding. 2007. "Inflammatory Bowel Disease: Cause and Immunobiology." *The Lancet* 369 (9573): 1627–40.

Biesalski, Hans K. 2016. "Nutrition Meets the Microbiome: Micronutrients and the Microbiota." *Annals of the New York Academy of Sciences* 1372 (1): 53–64.

Bowers, Robert M., Nikos C. Kyrpides, Ramunas Stepanauskas, Miranda Harmon-Smith, Devin Doud, T. B. K. Reddy, Frederik Schulz, et al. 2017. "Minimum Information about a Single Amplified Genome (MISAG) and a Metagenome-Assembled Genome (MIMAG) of Bacteria and Archaea." *Nature Biotechnology* 35 (8): 725–31.

Campbell, James H., Patrick O'Donoghue, Alisha G. Campbell, Patrick Schwientek,

898   Alexander Sczyrba, Tanja Woyke, Dieter Söll, and Mircea Podar. 2013. "UGA Is an
899   Additional Glycine Codon in Uncultured SR1 Bacteria from the Human Microbiota."
900   *Proceedings of the National Academy of Sciences of the United States of America*
901   110 (14): 5540–45.
902 Chaumeil, Pierre-Alain, Aaron J. Mussig, Philip Hugenholtz, and Donovan H. Parks.
903   2019. "GTDB-Tk: A Toolkit to Classify Genomes with the Genome Taxonomy
904   Database." *Bioinformatics* , November.
905   https://doi.org/10.1093/bioinformatics/btz848.
906 Chen, Lin-Xing, Karthik Anantharaman, Alon Shaiber, A. Murat Eren, and Jillian F.
907   Banfield. 2020. "Accurate and Complete Genomes from Metagenomes." *Genome*
908   *Research* 30 (3): 315–33.
909 Chien, Hsiang-Lin, Hui-Yu Huang, and Cheng-Chun Chou. 2006. "Transformation of
910   Isoflavone Phytoestrogens during the Fermentation of Soymilk with Lactic Acid
911   Bacteria and Bifidobacteria." *Food Microbiology* 23 (8): 772–78.
912 Chow, Janet, Haiqing Tang, and Sarkis K. Mazmanian. 2011. "Pathobionts of the
913   Gastrointestinal Microbiota and Inflammatory Disease." *Current Opinion in*
914   *Immunology* 23 (4): 473–80.
915 Clooney, Adam G., Julia Eckenberger, Emilio Laserna-Mendieta, Kathryn A. Sexton,
916   Matthew T. Bernstein, Kathy Vagianos, Michael Sargent, et al. 2021. "Ranking
917   Microbiome Variance in Inflammatory Bowel Disease: A Large Longitudinal
918   Intercontinental Study." *Gut* 70 (3): 499–510.
919 Costello, Elizabeth K., Keaton Stagaman, Les Dethlefsen, Brendan J. M. Bohannan,
920   and David A. Relman. 2012. "The Application of Ecological Theory toward an
921   Understanding of the Human Microbiome." *Science* 336 (6086): 1255–62.
922 D'Aimmo, M. R., P. Mattarelli, B. Biavati, N. G. Carlsson, and T. Andlid. 2012. "The
923   Potential of Bifidobacteria as a Source of Natural Folate." *Journal of Applied*
924   *Microbiology* 112 (5): 975–84.
925 David, Lawrence A., Arne C. Materna, Jonathan Friedman, Maria I. Campos-Baptista,
926   Matthew C. Blackburn, Allison Perrotta, Susan E. Erdman, and Eric J. Alm. 2014.
927   "Host Lifestyle Affects Human Microbiota on Daily Timescales." *Genome Biology* 15
928   (7): R89.
929 Delmont, Tom O., and A. Murat Eren. 2016. "Identifying Contamination with Advanced
930   Visualization and Analysis Practices: Metagenomic Approaches for Eukaryotic
931   Genome Assemblies." *PeerJ* 4 (March): e1839.
932 Delmont, Tom O., Christopher Quince, Alon Shaiber, Özcan C. Esen, Sonny Tm Lee,
933   Michael S. Rappé, Sandra L. McLellan, Sebastian Lücker, and A. Murat Eren.
934   2018. "Nitrogen-Fixing Populations of Planctomycetes and Proteobacteria Are
935   Abundant in Surface Ocean Metagenomes." *Nature Microbiology* 3 (7): 804–13.
936 Denef, Vincent J. 2019. "Peering into the Genetic Makeup of Natural Microbial
937   Populations Using Metagenomics." In *Population Genomics: Microorganisms*,
938   edited by Martin F. Polz and Om P. Rajora, 49–75. Cham: Springer International
939   Publishing.
940 De Preter, Vicky, Veerle Bulteel, Peter Suenaert, Karen Paula Geboes, Gert De
941   Hertogh, Anja Luypaerts, Karel Geboes, Kristin Verbeke, and Paul Rutgeerts. 2009.
942   "Pouchitis, Similar to Active Ulcerative Colitis, Is Associated with Impaired Butyrate
943   Oxidation by Intestinal Mucosa." *Inflammatory Bowel Diseases* 15 (3): 335–40.

D'Souza, Glen, Shraddha Shitut, Daniel Preussger, Ghada Yousif, Silvio Waschina, and
Christian Kost. 2018. "Ecology and Evolution of Metabolic Cross-Feeding
Interactions in Bacteria." *Natural Product Reports* 35 (5): 455–88.
Durack, Juliana, and Susan V. Lynch. 2019. "The Gut Microbiome: Relationships with
Disease and Opportunities for Therapy." *The Journal of Experimental Medicine* 216
(1): 20–40.
Eddy, Sean R. 2011. "Accelerated Profile HMM Searches." *PLoS Computational
Biology* 7 (10): e1002195.
Edgar, Robert C. 2004. "MUSCLE: Multiple Sequence Alignment with High Accuracy
and High Throughput." *Nucleic Acids Research* 32 (5): 1792–97.
Eiseman, B., W. Silen, G. S. Bascom, and A. J. Kauvar. 1958. "Fecal Enema as an
Adjunct in the Treatment of Pseudomembranous Enterocolitis." *Surgery* 44 (5):
854–59.
Eisenstein, Michael. 2020. "The Hunt for a Healthy Microbiome." *Nature* 577 (7792):
S6–8.
Enright, A. J., S. Van Dongen, and C. A. Ouzounis. 2002. "An Efficient Algorithm for
Large-Scale Detection of Protein Families." *Nucleic Acids Research* 30 (7): 1575–
84.
Eren, A. Murat, Özcan C. Esen, Christopher Quince, Joseph H. Vineis, Hilary G.
Morrison, Mitchell L. Sogin, and Tom O. Delmont. 2015. "Anvi'o: An Advanced
Analysis and Visualization Platform for 'Omics Data." *PeerJ* 3 (October): e1319.
Eren, A. Murat, Evan Kiefl, Alon Shaiber, Iva Veseli, Samuel E. Miller, Matthew S.
Schechter, Isaac Fink, et al. 2021. "Community-Led, Integrated, Reproducible
Multi-Omics with Anvi'o." *Nature Microbiology* 6 (1): 3–6.
Eren, A. Murat, Joseph H. Vineis, Hilary G. Morrison, and Mitchell L. Sogin. 2013. "A
Filtering Method to Generate High Quality Short Reads Using Illumina Paired-End
Technology." *PloS One* 8 (6): e66643.
Feng, Lihui, Arjun S. Raman, Matthew C. Hibberd, Jiye Cheng, Nicholas W. Griffin,
Yangqing Peng, Semen A. Leyn, Dmitry A. Rodionov, Andrei L. Osterman, and
Jeffrey I. Gordon. 2020. "Identifying Determinants of Bacterial Fitness in a Model of
Human Gut Microbial Succession." *Proceedings of the National Academy of
Sciences of the United States of America* 117 (5): 2622–33.
Finucane, Mariel M., Thomas J. Sharpton, Timothy J. Laurent, and Katherine S. Pollard.
2014. "A Taxonomic Signature of Obesity in the Microbiome? Getting to the Guts of
the Matter." *PloS One* 9 (1): e84689.
Galperin, Michael Y., Yuri I. Wolf, Kira S. Makarova, Roberto Vera Alvarez, David
Landsman, and Eugene V. Koonin. 2021. "COG Database Update: Focus on
Microbial Diversity, Model Organisms, and Widespread Pathogens." *Nucleic Acids
Research* 49 (D1): D274–81.
Gomes, Ana M. P., and F. Xavier Malcata. 1999. "Bifidobacterium Spp. and
Lactobacillus Acidophilus: Biological, Biochemical, Technological and Therapeutical
Properties Relevant for Use as Probiotics." *Trends in Food Science & Technology*
10 (4): 139–57.
Grehan, Martin J., Thomas Julius Borody, Sharyn M. Leis, Jordana Campbell, Hazel
Mitchell, and Antony Wettstein. 2010. "Durable Alteration of the Colonic Microbiota
by the Administration of Donor Fecal Flora." *Journal of Clinical Gastroenterology* 44

(8): 551–61.

Herrmann, Klaus M., and Lisa M. Weaver. 1999. "THE SHIKIMATE PATHWAY." *Annual Review of Plant Physiology and Plant Molecular Biology* 50 (June): 473–503.

Human Microbiome Project Consortium. 2012. "Structure, Function and Diversity of the Healthy Human Microbiome." *Nature* 486 (7402): 207–14.

Hyatt, Doug, Gwo-Liang Chen, Philip F. Locascio, Miriam L. Land, Frank W. Larimer, and Loren J. Hauser. 2010. "Prodigal: Prokaryotic Gene Recognition and Translation Initiation Site Identification." *BMC Bioinformatics* 11 (March): 119.

Isaac, Sandrine, Jose U. Scher, Ana Djukovic, Nuria Jiménez, Dan R. Littman, Steven B. Abramson, Eric G. Pamer, and Carles Ubeda. 2017. "Short- and Long-Term Effects of Oral Vancomycin on the Human Intestinal Microbiota." *The Journal of Antimicrobial Chemotherapy* 72 (1): 128–36.

Joossens, Marie, Geert Huys, Margo Cnockaert, Vicky De Preter, Kristin Verbeke, Paul Rutgeerts, Peter Vandamme, and Severine Vermeire. 2011. "Dysbiosis of the Faecal Microbiota in Patients with Crohn's Disease and Their Unaffected Relatives." *Gut* 60 (5): 631–37.

Jungersen, Mikkel, Anette Wind, Eric Johansen, Jeffrey E. Christensen, Birgitte Stuer-Lauridsen, and Dorte Eskesen. 2014. "The Science behind the Probiotic Strain Bifidobacterium Animalis Subsp. Lactis BB-12(®)." *Microorganisms* 2 (2): 92–110.

Kanehisa, M., and S. Goto. 2000. "KEGG: Kyoto Encyclopedia of Genes and Genomes." *Nucleic Acids Research* 28 (1): 27–30.

Kanehisa, Minoru, Miho Furumichi, Mao Tanabe, Yoko Sato, and Kanae Morishima. 2017. "KEGG: New Perspectives on Genomes, Pathways, Diseases and Drugs." *Nucleic Acids Research* 45 (D1): D353–61.

Kanehisa, Minoru, Susumu Goto, Yoko Sato, Masayuki Kawashima, Miho Furumichi, and Mao Tanabe. 2014. "Data, Information, Knowledge and Principle: Back to Metabolism in KEGG." *Nucleic Acids Research* 42 (Database issue): D199–205.

Kao, Dina, Brandi Roach, Marisela Silva, Paul Beck, Kevin Rioux, Gilaad G. Kaplan, Hsiu-Ju Chang, et al. 2017. "Effect of Oral Capsule- vs Colonoscopy-Delivered Fecal Microbiota Transplantation on Recurrent Clostridium Difficile Infection: A Randomized Clinical Trial." *JAMA: The Journal of the American Medical Association* 318 (20): 1985–93.

Khoruts, Alexander, Johan Dicksved, Janet K. Jansson, and Michael J. Sadowsky. 2010. "Changes in the Composition of the Human Fecal Microbiome after Bacteriotherapy for Recurrent Clostridium Difficile-Associated Diarrhea." *Journal of Clinical Gastroenterology* 44 (5): 354–60.

Kirchhelle, Claas. 2018. "Pharming Animals: A Global History of Antibiotics in Food Production (1935–2017)." *Palgrave Communications* 4 (1): 96.

Kleerebezem, Michiel, and Elaine E. Vaughan. 2009. "Probiotic and Gut Lactobacilli and Bifidobacteria: Molecular Approaches to Study Diversity and Activity." *Annual Review of Microbiology* 63: 269–90.

Koenig, Jeremy E., Aymé Spor, Nicholas Scalfone, Ashwana D. Fricker, Jesse Stombaugh, Rob Knight, Largus T. Angenent, and Ruth E. Ley. 2011. "Succession of Microbial Consortia in the Developing Infant Gut Microbiome." *Proceedings of the National Academy of Sciences of the United States of America* 108 Suppl 1 (March): 4578–85.

Kolmogorov, Mikhail, Mikhail Rayko, Jeffrey Yuan, Evgeny Polevikov, and Pavel
Pevzner. 2019. "metaFlye: Scalable Long-Read Metagenome Assembly Using
Repeat Graphs." *bioRxiv*, 637637.
Köster, Johannes, and Sven Rahmann. 2012. "Snakemake—a Scalable Bioinformatics
Workflow Engine." *Bioinformatics* 28 (19): 2520–22.
Kowarsky, Mark, Joan Camunas-Soler, Michael Kertesz, Iwijn De Vlaminck, Winston
Koh, Wenying Pan, Lance Martin, et al. 2017. "Numerous Uncharacterized and
Highly Divergent Microbes Which Colonize Humans Are Revealed by Circulating
Cell-Free DNA." *Proceedings of the National Academy of Sciences of the United
States of America*, August. https://doi.org/10.1073/pnas.1707009114.
Langmead, Ben, and Steven L. Salzberg. 2012. "Fast Gapped-Read Alignment with
Bowtie 2." *Nature Methods* 9 (4): 357–59.
Lee, S. Melanie, Gregory P. Donaldson, Zbigniew Mikulski, Silva Boyajian, Klaus Ley,
and Sarkis K. Mazmanian. 2013. "Bacterial Colonization Factors Control Specificity
and Stability of the Gut Microbiota." *Nature* 501 (7467): 426–29.
Lee, Sonny T. M., Stacy A. Kahn, Tom O. Delmont, Alon Shaiber, Özcan C. Esen,
Nathaniel A. Hubert, Hilary G. Morrison, Dionysios A. Antonopoulos, David T.
Rubin, and A. Murat Eren. 2017. "Tracking Microbial Colonization in Fecal
Microbiota Transplantation Experiments via Genome-Resolved Metagenomics."
Microbiome 5 (1): 50.
Ley, Ruth E., Peter J. Turnbaugh, Samuel Klein, and Jeffrey I. Gordon. 2006. "Microbial
Ecology: Human Gut Microbes Associated with Obesity." *Nature* 444 (7122): 1022–
23.
Li, Heng, Bob Handsaker, Alec Wysoker, Tim Fennell, Jue Ruan, Nils Homer, Gabor
Marth, Goncalo Abecasis, Richard Durbin, and 1000 Genome Project Data
Processing Subgroup. 2009. "The Sequence Alignment/Map Format and
SAMtools." *Bioinformatics* 25 (16): 2078–79.
Lloyd-Price, Jason, Galeb Abu-Ali, and Curtis Huttenhower. 2016. "The Healthy Human
Microbiome." *Genome Medicine* 8 (1): 51.
Lloyd-Price, Jason, Cesar Arze, Ashwin N. Ananthakrishnan, Melanie Schirmer, Julian
Avila-Pacheco, Tiffany W. Poon, Elizabeth Andrews, et al. 2019. "Multi-Omics of
the Gut Microbial Ecosystem in Inflammatory Bowel Diseases." *Nature* 569 (7758):
655–62.
Lynch, Susan V., and Oluf Pedersen. 2016. "The Human Intestinal Microbiome in
Health and Disease." *The New England Journal of Medicine* 375 (24): 2369–79.
Maignien, Loïs, Emelia A. DeForce, Meghan E. Chafee, A. Murat Eren, and Sheri L.
Simmons. 2014. "Ecological Succession and Stochastic Variation in the Assembly
of Arabidopsis Thaliana Phyllosphere Communities." *mBio* 5 (1): e00682–13.
Martens, J. H., H. Barg, M. J. Warren, and D. Jahn. 2002. "Microbial Production of
Vitamin B12." *Applied Microbiology and Biotechnology* 58 (3): 275–85.
Martiny, Adam C., Kathleen Treseder, and Gordon Pusch. 2013. "Phylogenetic
Conservatism of Functional Traits in Microorganisms." *The ISME Journal* 7 (4):
830–38.
McBurney, Michael I., Cindy Davis, Claire M. Fraser, Barbara O. Schneeman, Curtis
Huttenhower, Kristin Verbeke, Jens Walter, and Marie E. Latulippe. 2019.
"Establishing What Constitutes a Healthy Human Gut Microbiome: State of the

Science, Regulatory Considerations, and Future Directions." *The Journal of Nutrition* 149 (11): 1882–95.

Messer, J. S., E. R. Liechty, O. A. Vogel, and E. B. Chang. 2017. "Evolutionary and Ecological Forces That Shape the Bacterial Communities of the Human Gut." *Mucosal Immunology* 10 (3): 567–79.

Minoche, André E., Juliane C. Dohm, and Heinz Himmelbauer. 2011. "Evaluation of Genomic High-Throughput Sequencing Data Generated on Illumina HiSeq and Genome Analyzer Systems." *Genome Biology* 12 (11): R112.

Ott, S. J., M. Musfeldt, D. F. Wenderoth, J. Hampe, O. Brant, U. R. Fölsch, K. N. Timmis, and S. Schreiber. 2004. "Reduction in Diversity of the Colonic Mucosa Associated Bacterial Microflora in Patients with Active Inflammatory Bowel Disease." *Gut* 53 (5): 685–93.

Parks, Donovan H., Maria Chuvochina, David W. Waite, Christian Rinke, Adam Skarshewski, Pierre-Alain Chaumeil, and Philip Hugenholtz. 2018. "A Standardized Bacterial Taxonomy Based on Genome Phylogeny Substantially Revises the Tree of Life." *Nature Biotechnology* 36 (10): 996–1004.

Pasolli, Edoardo, Francesco Asnicar, Serena Manara, Moreno Zolfo, Nicolai Karcher, Federica Armanini, Francesco Beghini, et al. 2019. "Extensive Unexplored Human Microbiome Diversity Revealed by Over 150,000 Genomes from Metagenomes Spanning Age, Geography, and Lifestyle." *Cell* 176 (3): 649–62.e20.

Peng, Yu, Henry C. M. Leung, S. M. Yiu, and Francis Y. L. Chin. 2012. "IDBA-UD: A de Novo Assembler for Single-Cell and Metagenomic Sequencing Data with Highly Uneven Depth." *Bioinformatics* 28 (11): 1420–28.

Philippot, Laurent, Siv G. E. Andersson, Tom J. Battin, James I. Prosser, Joshua P. Schimel, William B. Whitman, and Sara Hallin. 2010. "The Ecological Coherence of High Bacterial Taxonomic Ranks." *Nature Reviews. Microbiology* 8 (7): 523–29.

Plichta, Damian R., Daniel B. Graham, Sathish Subramanian, and Ramnik J. Xavier. 2019. "Therapeutic Opportunities in Inflammatory Bowel Disease: Mechanistic Dissection of Host-Microbiome Relationships." *Cell* 178 (5): 1041–56.

Podlesny, Daniel, and W. Florian Fricke. 2020. "Microbial Strain Engraftment, Persistence and Replacement after Fecal Microbiota Transplantation." *medRxiv*, September, 2020.09.29.20203638.

Pompei, Anna, Lisa Cordisco, Alberto Amaretti, Simona Zanoni, Diego Matteuzzi, and Maddalena Rossi. 2007. "Folate Production by Bifidobacteria as a Potential Probiotic Property." *Applied and Environmental Microbiology* 73 (1): 179–85.

Price, Morgan N., Paramvir S. Dehal, and Adam P. Arkin. 2010. "FastTree 2--Approximately Maximum-Likelihood Trees for Large Alignments." *PloS One* 5 (3): e9490.

Pritchard, Leighton, Rachel H. Glover, Sonia Humphris, John G. Elphinstone, and Ian K. Toth. 2016. "Genomics and Taxonomy in Diagnostics for Food Security: Soft-Rotting Enterobacterial Plant Pathogens." *Analytical Methods* 8 (1): 12–24.

Quince, Christopher, Tom O. Delmont, Sébastien Raguideau, Johannes Alneberg, Aaron E. Darling, Gavin Collins, and A. Murat Eren. 2017. "DESMAN: A New Tool for de Novo Extraction of Strains from Metagenomes." *Genome Biology* 18 (1): 181.

Rinke, Christian, Patrick Schwientek, Alexander Sczyrba, Natalia N. Ivanova, Iain J. Anderson, Jan-Fang Cheng, Aaron Darling, et al. 2013. "Insights into the

Phylogeny and Coding Potential of Microbial Dark Matter." *Nature* 499 (7459): 431–
37.
Rothschild, Daphna, Omer Weissbrod, Elad Barkan, Alexander Kurilshikov, Tal Korem,
David Zeevi, Paul I. Costea, et al. 2018. "Environment Dominates over Host
Genetics in Shaping Human Gut Microbiota." *Nature* 555 (7695): 210–15.
Schell, Mark A., Maria Karmirantzou, Berend Snel, David Vilanova, Bernard Berger,
Gabriella Pessi, Marie-Camille Zwahlen, et al. 2002. "The Genome Sequence of
Bifidobacterium Longum Reflects Its Adaptation to the Human Gastrointestinal
Tract." *Proceedings of the National Academy of Sciences of the United States of*
America 99 (22): 14422–27.
Schirmer, Melanie, Ashley Garner, Hera Vlamakis, and Ramnik J. Xavier. 2019.
"Microbial Genes and Pathways in Inflammatory Bowel Disease." *Nature Reviews.*
Microbiology 17 (8): 497–511.
Schmidt, Thomas S. B., Jeroen Raes, and Peer Bork. 2018. "The Human Gut
Microbiome: From Association to Modulation." *Cell* 172 (6): 1198–1215.
Shahinas, Dea, Michael Silverman, Taylor Sittler, Charles Chiu, Peter Kim, Emma
Allen-Vercoe, Scott Weese, Andrew Wong, Donald E. Low, and Dylan R. Pillai.
2012. "Toward an Understanding of Changes in Diversity Associated with Fecal
Microbiome Transplantation Based on 16S rRNA Gene Deep Sequencing." *mBio* 3
(5): e00338–12.
Shaiber, Alon, Amy D. Willis, Tom O. Delmont, Simon Roux, Lin-Xing Chen, Abigail C.
Schmid, Mahmoud Yousef, et al. 2020. "Functional and Genetic Markers of Niche
Partitioning among Enigmatic Members of the Human Oral Microbiome." *Genome*
Biology 21 (1): 292.
Sharon, Itai, Michael J. Morowitz, Brian C. Thomas, Elizabeth K. Costello, David A.
Relman, and Jillian F. Banfield. 2013. "Time Series Community Genomics Analysis
Reveals Rapid Shifts in Bacterial Species, Strains, and Phage during Infant Gut
Colonization." *Genome Research* 23 (1): 111–20.
Sheth, Ravi U., Mingqiang Li, Weiqian Jiang, Peter A. Sims, Kam W. Leong, and Harris
H. Wang. 2019. "Spatial Metagenomic Characterization of Microbial Biogeography
in the Gut." *Nature Biotechnology* 37 (8): 877–83.
Smillie, Christopher S., Jenny Sauk, Dirk Gevers, Jonathan Friedman, Jaeyun Sung,
Ilan Youngster, Elizabeth L. Hohmann, et al. 2018. "Strain Tracking Reveals the
Determinants of Bacterial Engraftment in the Human Gut Following Fecal
Microbiota Transplantation." *Cell Host & Microbe* 23 (2): 229–40.e5.
Sokol, Harry, and Philippe Seksik. 2010. "The Intestinal Microbiota in Inflammatory
Bowel Diseases: Time to Connect with the Host." *Current Opinion in*
Gastroenterology 26 (4): 327–31.
Strozzi, G. Paolo, and Luca Mogna. 2008. "Quantification of Folic Acid in Human Feces
after Administration of Bifidobacterium Probiotic Strains." *Journal of Clinical*
Gastroenterology 42 Suppl 3 Pt 2 (September): S179–84.
Sugahara, Hirosuke, Toshitaka Odamaki, Nanami Hashikura, Fumiaki Abe, and Jin-
Zhong Xiao. 2015. "Differences in Folate Production by Bifidobacteria of Different
Origins." *Bioscience of Microbiota, Food and Health* 34 (4): 87–93.
Swidsinski, Alexander, Jutta Weber, Vera Loening-Baucke, Laura P. Hale, and Herbert
Lochs. 2005. "Spatial Organization and Composition of the Mucosal Flora in

Patients with Inflammatory Bowel Disease." *Journal of Clinical Microbiology* 43 (7): 3380–89.

Tatusov, Roman L., Natalie D. Fedorova, John D. Jackson, Aviva R. Jacobs, Boris Kiryutin, Eugene V. Koonin, Dmitri M. Krylov, et al. 2003. "The COG Database: An Updated Version Includes Eukaryotes." *BMC Bioinformatics* 4 (September): 41.

Utter, Daniel R., Gary G. Borisy, A. Murat Eren, Colleen M. Cavanaugh, and Jessica L. Mark Welch. 2020. "Metapangenomics of the Oral Microbiome Provides Insights into Habitat Adaptation and Cultivar Diversity." *Genome Biology* 21 (1): 293.

Vanni, Chiara, Matthew S. Schechter, Silvia G. Acinas, Albert Barberán, Pier Luigi Buttigieg, Emilio O. Casamayor, Tom O. Delmont, et al. 2020. "Unifying the Global Coding Sequence Space Enables the Study of Genes with Unknown Function across Biomes." *Cold Spring Harbor Laboratory*. https://doi.org/10.1101/2020.06.30.180448.

Ventura, Marco, Sarah O'Flaherty, Marcus J. Claesson, Francesca Turroni, Todd R. Klaenhammer, Douwe van Sinderen, and Paul W. O'Toole. 2009. "Genome-Scale Analyses of Health-Promoting Bacteria: Probiogenomics." *Nature Reviews. Microbiology* 7 (1): 61–71.

Vineis, Joseph H., Daina L. Ringus, Hilary G. Morrison, Tom O. Delmont, Sushila Dalal, Laura H. Raffals, Dionysios A. Antonopoulos, et al. 2016. "Patient-Specific *Bacteroides* Genome Variants in Pouchitis." *mBio* 7 (6): e01713–16, /mbio/7/6/e01713–16.atom.

Walker, Bruce J., Thomas Abeel, Terrance Shea, Margaret Priest, Amr Abouelliel, Sharadha Sakthikumar, Christina A. Cuomo, et al. 2014. "Pilon: An Integrated Tool for Comprehensive Microbial Variant Detection and Genome Assembly Improvement." *PloS One* 9 (11): e112963.

Walter, Jens, Anissa M. Armet, B. Brett Finlay, and Fergus Shanahan. 2020. "Establishing or Exaggerating Causality for the Gut Microbiome: Lessons from Human Microbiota-Associated Rodents." *Cell* 180 (2): 221–32.

Wexler, Aaron G., and Andrew L. Goodman. 2017. "An Insider's Perspective: Bacteroides as a Window into the Microbiome." *Nature Microbiology*. https://doi.org/10.1038/nmicrobiol.2017.26.

Wood, Derrick E., Jennifer Lu, and Ben Langmead. 2019. "Improved Metagenomic Analysis with Kraken 2." *Genome Biology* 20 (1): 257.

Wu, Guojun, Naisi Zhao, Chenhong Zhang, Yan Y. Lam, and Liping Zhao. 2021. "Guild-Based Analysis for Understanding Gut Microbiome in Human Health and Diseases." *Genome Medicine* 13 (1): 22.

Yasuda, Koji, Keunyoung Oh, Boyu Ren, Timothy L. Tickle, Eric A. Franzosa, Lynn M. Wachtman, Andrew D. Miller, et al. 2015. "Biogeography of the Intestinal Mucosal and Lumenal Microbiome in the Rhesus Macaque." *Cell Host & Microbe* 17 (3): 385–91.

 # Supplementary Figures

Supplementary Figure 1. Timeline of stool samples collected from FMT study. Each circle represents a stool
sample collected from either an FMT donor or FMT recipient. The thicker, red vertical line at day 0 represents the FMT
event for each recipient. FMT method (pill or colonoscopy) and FMT recipient health and disease state (C. diff - chronic
recurrent *Clostridium difficile* infection, UC - ulcerative colitis) are indicated on the right.

Supplementary Figure 2. Nonmetric multidimensional scaling (NMDS) ordination of the taxonomic composition
of donor, recipient, and Canadian gut metagenomes at the genus level based on Morisita-Horn dissimilarity.
Samples from the same participant are joined by lines with the earliest time point labeled. CAN: Canadian gut
metagenomes, DA: donor A, DB: donor B, POST: recipients post-FMT, PRE: recipients pre-FMT.

1225

1226 **Supplementary Figure 3. Nonmetric multidimensional scaling (NMDS) ordination of the taxonomic composition**
1227 **of the donor and recipient metagenomes at genus level based on Morisita-Horn dissimilarity.** Samples from the
1228 same participant are joined by lines with the earliest time point labeled. DA_POST: donor A recipients post-FMT,
1229 DA_PRE: donor A recipients pre-FMT, DA: donor A, DB_POST: donor B recipients post-FMT, DB_PRE: donor B
1230 recipients pre-FMT, DB: donor B.

1231

Supplementary Figure 4. A flowchart outlining our method to assign successful colonization, failed colonization, or
undetermined colonization phenotypes to donor-derived populations in the recipients of that donor's stool.

Supplementary Tables (⌖)

Supplementary Table 1: Description of FMT study and stool samples collected. a) Description of FMT donor stool samples and SRA accession numbers. b) Description of FMT recipient samples and SRA accession numbers. c) Description of transplantation events.

Supplementary Table 2: Description of FMT metagenomes and co-assemblies. a) Metagenome SRA accession numbers and number of metagenomic short-reads sequenced and mapped to co-assemblies and MAGs. b) Phylum level taxonomic composition of metagenomes. c) Genus level taxonomic composition of metagenomes. d) Summary statistics for contigs from metagenome co-assemblies.

Supplementary Table 3: Description of MAGs. a) Summary statistics and taxonomic assignments for MAGs. b) and c) Detection of Donor A and Donor B MAGs in FMT metagenomes, respectively. d) and e) Detection of Donor A and Donor B MAGs in global gut metagenomes, respectively. f) and g) Detection summary statistics of Donor A and Donor B MAGs in global gut metagenomes, respectively.

Supplementary Table 4: Accession numbers of gut metagenomes from 17 countries.

Supplementary Table 5: MAG subpopulation information. a) and b) Number of Donor A and Donor B MAG subpopulations detected in FMT metagenomes, respectively. c) and d) Subpopulation composition of Donor A and Donor B MAGs in FMT metagenomes, respectively.

Supplementary Table 6: MAG/recipient pair colonization outcomes and MAG mean coverage in the 2nd and 3rd quartiles in stool samples used for transplantation.

Supplementary Table 7: Description of high vs. low-fitness populations. a) Taxonomic assignments and genome size estimates for high- and low-fitness populations. b) KEGG module completeness information for high- and low-fitness populations. c) Raw KEGG module enrichment information for high- and low-fitness populations. d) KEGG module enrichment and categorical information for the 33 modules enriched in high-fitness populations. e) and f) Completeness information for the 33 modules enriched in high-fitness populations in all high- and low-fitness populations.

Supplementary Table 8: a) List of genomes from healthy individuals and individuals with IBD. b) Module completion values across genomes.

Supplementary Table 9: Bifidobacteria functional analysis. a) Accession numbers for *Bifidobacteria* reference genomes. b) Summary statistics for *Bifidobacteria* MAGs and reference genomes. c) Prevalence of *Bifidobacteria* MAGs in global gut metagenomes. d) gANI percent identity between *Bifidobacteria* genomes. e) gANI percent alignment coverage between *Bifidobacteria* genomes. f) KOfams enriched in different *Bifidobacteria* species. g) KOfam presence and absence in *Bifidobacteria* genomes. h) COG functions enriched in different *Bifidobacteria* species. i) COG function presence and absence in *Bifidobacteria* genomes. j) KEGG modules enriched in different Bifidobacteria species. k) KEGG module completeness in *Bifidobacteria* genomes.