The "two-brain" approach reveals the active role of task-deactivated default

mode network in speech comprehension

Lanfang Liu [1], Hehui Li[2], Zhiting Ren[3], Qi Zhou[1], Yuxuan Zhang[1], Chunming Lu[1],

Jiang Qiu[3], Hong Chen*[3], Guosheng Ding*[1]

[1]State Key Laboratory of Cognitive Neuroscience and Learning, Beijing Normal

University & IDG/McGovern Institute for Brain Research, Beijing, 100875, P.R.

China;

[2]Center for Brain Disorders and Cognitive Sciences, Shenzhen University, Shenzhen,

518060, PR China

[3]Key Laboratory of Cognition and Personality (SWU), Ministry of Education &

Department of Psychology, Southwest University, Chongqing, 400715, P.R. China

Corresponding author: either Guosheng Ding, State Key Laboratory of Cognitive

Neuroscience and Learning, Beijing Normal University, No. 19 Xinjiekouwai Street,

Beijing 100875, China, E-mail: dinggsh@bnu.edu.cn, or Hong Chen, School of

psychology, SouthWest University, No. 2 Tiansheng Street, Beibei, Chongqing,

400715, E-mail: chenhg@swu.edu.cn

Running title: The active role of deactivated DMN in speech comprehension

Abstract

During speech comprehension, as listeners need to keep tracking the external audio streams, the default mode network (DMN) is often de-activated and anticorrelated with task-positive networks. Such a pattern has been interpreted as the suppression of the DMN to support externally-oriented cognitive processes. Taking a "two-brain" approach, the current study demonstrated that, despite exhibiting deactivation and anticorrelated with the language network and executive control network, the DMN was not suppressed but played an active role in spoken narrative comprehension. This was evidenced by significant listener-speaker neural couplings in both the posterior and anterior DMN and the positive correlation between the coupling strength and listener's speech comprehension. Moreover, we demonstrated that the functionality of posterior DMN depended on its interaction with the executive control network, rather than its level of activation. Finally, Dynamic Causal Modeling together with the two-brain results indicates the language and executive control networks, the anterior DMN, and the posterior DMN occupied the bottom, intermediate and top layers of a hierarchical system, respectively. These results suggest that the DMN may primarily serve as an internally-oriented system that cooperates with the externally-oriented networks, which may allow the transformation of external acoustic signals into internal mental representations during language comprehension.

Keywords: default mode network, neural coupling, speech comprehension,two-brain

1. Introduction

In neurolinguistic studies, verbal communication has been usually approached as a

signal transmission process, and the main attention has been paid to the neural

responses bound to and triggered by the occurrence of speech signals (Stolk et al.

2016). Within this signal-centered framework, the crucial role of a set of inferior

frontal and temporal regions (i.e., the language network, LN) in coding and decoding

the sensory-motor, syntactic and semantic properties of speech signals, as well as a set

of middle frontal and parietal regions in monitoring and coordinating these processes

(i.e., the executive control network, ECN) (Ye and Zhou 2009), have been well

recognized. Nevertheless, in real-life situations, the speech signals we receive are

usually impoverished or ambiguous. Yet, we can still decipher the meaning intended

by a speaker accurately. By contrast, current state-of-art artificial agents, such as

Apple's Siri or Microsoft's Cortana, often make communicative errors despite they

have mastered the signal decoding rules. These observations indicate that successful

speech understanding demands more than signal decoding.

Indeed, psychological models have long proposed that, in addition to the

perceptual and linguistic analyses, to successfully understand the speaker's mind

during verbal communication engages a set of extra-linguistic processes, mainly

including mentalizing, self-projection and the theory of mind (Carruthers and Smith

1996; Zwaan and Radvansky 1998; Frith and Frith 1999). These processes have been

frequently attributed to the default mode network (DMN) (Spreng et al. 2008; Spreng

and Grady 2009; Li et al. 2014), an intrinsically organized functional system

composed of the posterior cingulate cortex, precuneus, medial prefrontal cortex

(mPFC), and lateral parietal cortex.

Nevertheless, current empirical evidence regarding the functional role of DMN

in speech processing seems conflicting. On the one hand, as auditory language

processing requires participants to keep tracking external audio streams, the DMN has

been repeatedly found to be deactivated relative to low-level baselines (Wilson et al.,

2007; Szaflarski et al., 2012; Rodriguez Moreno et al., 2014; Horowitz-Kraus et al.,

2017; Cuevas et al., 2019), and anticorrelated with networks dedicated to

externally-oriented processes (including the LN and ECN) (Fox et al. 2005; Uddin et

al. 2009; Smith et al. 2012). The task-induced deactivation and network

anticorrelation seem to imply that the DMN was "suppressed" to support

externally-oriented cognitive activities (Anticevic 2012; Gauffin et al. 2013; Zhou et

al. 2016).

On the other hand, studies measuring the consistency of

blood-oxygen-level-dependent (BOLD) signal fluctuations across subjects

experiencing the same narratives (i.e., intersubject correlation, ISC) have found

significant ISC in the DMN regions (Wilson et al., 2007; Lerner et al., 2011; Simony

et al., 2016; Yeshurun et al., 2017). In addition, the ISC in the DMN was modulated

by the manipulation of story belief (Yeshurun et al., 2017) and the coherence of the

stimuli's temporal structure over minutes of time (Lerner et al., 2011; Simony et al.,

2016). These results are postulated to reflect the active role of DMN in

semantic/conceptual processing.

How to reconcile a potentially active role of the DMN as argued by ISC studies

with the longstanding observation that the DMN shows task-induced deactivation and

anticorrelated with task-positive networks? One possible explanation is that the

significant ISC (and its change) in the DMN reported in previous studies is an

epiphenomenon of task disengagement, rather than signifying a positive role. It is well

known that activities in DMN regions are systematically modulated by task difficulty

(Zwaan and Radvansky 1998; McKiernan et al. 2003; Greicius and Menon 2004;

Humphreys et al. 2015). If subjects find the same part of the narrative more or less

engaging, the DMN will be down- or up-regulated similarly across subjects, which

would result in the correlation of brain signal fluctuations across subjects (Wilson et

al. 2007). Alternatively, the DMN is indeed critically engaged in language

comprehension, but the processes (likely internally-directed) it subserves are opposite

to those (likely externally-directed) subserved by the task-positive networks. This

hypothesis is put forward mainly based on the linguistic theory that the basic design

for language faculty includes not only externally-directed modules for sensory-motor

and linguistic processing but also an internally-directed module for

conceptual-intentional representations (Berwick et al. 2013). However, it remains to

be established that the DMN showing deactivation and anticorrelation with

task-positive networks is not "suppressed" but actively involved in language

comprehension. Moreover, how the task-activated and task-deactivated networks are

organized to support speech comprehension remains largely unknown.

To address the above issues, we took a "two-brain" approach and combined

within-brain and between-brain analyses. In our experiment, a speaker telling real-life

stories and later a group of listeners hearing the stories were scanned with fMRI.

Rather than relying on the ISC across listeners to determine the involvement of DMN,

we focused on neural couplings (i.e., the correlation of brain activities over time)

between the listener and the speaker. Compared to the ISC, the inter-brain coupling

may have at least two advantages to uncover the functional role of DMN in task

processing. First, the cognitive meaning of listener-speaker neural coupling might be

clearer than the ISC. Accumulating evidence suggests that inter-brain couplings at

different cortical regions reflect the alignment between communicators at different

levels of mental representations (Menenti et al. 2012; Schoot et al. 2016; Stolk *et al.*

2016). Indeed, it has been found that stronger inter-brain coupling was associated with

better communication (Stephens et al. 2010; Liu et al. 2020; Li et al. 2021; Liu et al.

2021). Second, the neural coupling between a speaker-listener pair is likely less

susceptible to the influences of task difficulty, since speech production is typically

more cognitively demanding than speech comprehension (Hendriks 2014).

The group Independent Component Analysis (ICA) was applied to detect

large-scale neural networks and the associated network time series. Compared to

selecting regions of interest, this data-driven approach has been shown to render

better representations of neural networks and produce more reliable results (Guo et al.

2012; Yu et al. 2017). As accumulating evidence showed that the DMN is not a

homogenous network, but generally composed of two functionally heterogeneous

subsystems (Uddin *et al.* 2009; Seghier and Price 2012b), the anterior and posterior

parts, we examined the anterior DMN (aDMN) and posterior DMN (pDMN)

separately. The two externally-oriented networks, the LN and ECN, were also

investigated.

We first examined the activation pattern and network functional connectivity

within the listeners' brains to compare with previous "single-brain" studies. Next, we

examined listener-speaker neural coupling and its association with task performance.

Upon establishing the active role of DMNs in speech comprehension, we explored the

organization of network communication applying Dynamic Causal Modelling (DCM)

to gain insight into the mechanism of the DMN's contribution to language processing.

Finally, we tested whether the functionality (measured by the inter-brain coupling) of

DMN depends on its interaction with other networks or on its response amplitude to

stimuli.

2. Materials and Methods

2.1 Participants and experimental procedure

A total of 69 Chinese college students (aged 19-27 years) participated in this study,

including one female speaker and 68 listeners (35 females). We also recruited a native

Mongolian speaker (a female college student) to serve as a control condition. All

participants were right-handed and reported no history of physiological or mental

disorder. None of these listeners had learned Mongolian. The datasets of four

participants listening to the Chinese story and four participants listening to the

Mongolian story were excluded from further analysis due to excessive head

movements during the scanning (more than 3mm or 3 degrees). Another two

participants were excluded due to their extremely low comprehension scores (lower

than 3.5 standard deviation from the group mean), which indicated they probably did

not listen to the story attentively. Written informed consent was obtained from all

participants under a protocol approved by the Reviewer Board of Southwest

University in China.

We first collected the data from the two speakers telling stories based on their

personal experience (10 min for each) when undergoing fMRI scanning. A

noise-canceling microphone (FOMRI-III, Optoacoustics Ltd., Or-Yehuda, Israel) was

used to simultaneously record the speech. After further offline denoising by Adobe

Audition 3.0 (Adobe Systems Inc., USA), the audio recordings were played back to

each listener during scanning. To make the listeners attend to the stories, we informed

them beforehand that a test about the content of the story would be given after the

scanning. During the scanning, the same set of visual stimuli were presented to the

speakers and listeners, consisting of a fixation cross lasting for 20s and then an icon of

a horn lasting until the end of the scanning run (Fig. 1A). The horn was designed to

prompt the participant to start speaking or listening immediately. To control for the

effect of low-level acoustic processing, we also played the story told by the

Mongolian speaker to each listener.

2.2 Behavioral assessment

To assess the degree to which the listeners understood the speech, we conducted an interview with each listener at the end of the scan. In the interview, the listeners were required first to retell the story with as much detail as possible. Next, experimenters asked the listeners several questions concerning the part of the contents not present in their free recalls. Two independent raters then scored the listeners based on the audio recordings of the interview. Details about the scoring procedure were provided in the supplementary material. The two raters had a high agreement on their assessments ($r_{(64)} = 0.80$, by Pearson's correlation), thus their assessments were averaged to be used as the measurement for speech comprehension. We also measured participants' memory spans using a digit memory test (Wechsler 1987) to account for a potential effect of memory capacity on the story-retelling task. The number of digits correctly repeated forward and backward were used as two covariates in the following brain-behavior analysis.

2.3 MRI acquisition and preprocessing

Imaging data were collected with a 3T Siemens Trio scanner in the MRI Center of the Southwest University of China. Functional images were acquired using a gradient echo-planar imaging sequence with the following parameters: repetition time = 2000 ms, echo time = 30 ms, flip angle = 90°, field of view = 220 mm^2, matrix size = 64 × 64, 32 interleaved slice, voxel size = 3.44 × 3.44 × 3.99 mm^3. T1 structural images were acquired using a MPRAGE sequence with the following parameters: repetition

time = 2530 ms, echo time = 3.39 ms, flip angle = 7°, FOV = 256 mm^2, scan order =

interleaved, matrix size = 256 × 256, and voxel size = 1.0 × 1.0 × 1.33 mm^3.

After discarding the first three volumes to allow for the equilibration of magnetic

fields, a total of 310 volumes were collected in each run. The DPABI toolkit (Yan et al.

2016) based on SPM12 (www.fil.ion.ucl.ac.uk/spm/) was utilized for image

preprocessing. The preprocessing steps included slice-timing correction, spatial

realignment, co-registration to individual subject's anatomical images, normalization

to the Montreal Neurological Institute (MNI) space, resampling into a 3 × 3 × 3 mm^3

voxel size, and smoothing (FWHM = 7mm). The resultant images were further

detrended, nuisance variable regressed, and high-pass filtered (1/128 Hz). The

nuisance regression included the removal of five principal components of white

matter and cerebrospinal fluid within individual subjects' T1 segmentation mask

(Behzadi et al. 2007), as well as Friston's 24 motion parameters (including each of the

six motion parameters of the current and preceding volume, plus each of these values

squared) (Friston et al. 1996).

2.4 Data analysis

2.4.1 Group Independent Component Analysis and component selection

For the successful and failed communication conditions separately, a group spatial

ICA was conducted using the Group ICA of fMRI Toolbox (GIFT). Before entering

the analysis, the first 10 volumes of the preprocessed functional image corresponding

to the 20s resting (fixation) period were discarded to avoid the potential modulation of

state change on the spatial patterns of components (Calhoun et al. 2008). The

dimensionality of preprocessed images from each participant was first reduced to 30

dimensions using principal component analysis. Next, the dimension-reduced data

from each participant (including both the speaker and all listeners) were temporally

concatenated and a group dimension reduction was performed. Then, twenty

independent sources were generated from the group data with the ICA using the

infomax algorithm. We choose to separate the whole brain signals into 20 independent

components because this dimension can well detect those typical brain networks

meanwhile keeping high stability of ICA estimation (Pamilo et al. 2012). Finally,

spatial maps and associated time series were reconstructed for each participant based

on the aggregate components and original data via spatial-temporal regression

(Beckmann et al. 2009) (Fig. 1C). This dual-regression technique has been

demonstrated to show high test-retest reliability (Zuo et al. 2010) and is widely used

to explore the functional connectivity of brain networks. Since the spatial maps and

time series of components have arbitrary units after the back-reconstruction step, they

were scaled to z-scores. Those time series associated with network components were

used in the following within-brain and between-brain connectivity analyses. A

one-sample *t*-test was conducted on individual subjects' component maps to obtain

the representative spatial distribution of each component. To test the robustness of our

findings, we also separated the whole brain signals into 30 independent components

and re-performed the analyses (supplementary material).

The four networks of interest, as well as the control network, were identified

based on the spatial overlap of the component map (i.e., the *t*-statistic map) with the

meta-analytic maps generated by the Neurosynth (https://neurosynth.org). The spatial

overlap of the two types of maps was quantified by the Dice score (Dice 1945).

Details for the calculation of the Dice score were provided in the supplementary

material. The term "default mode network" was adopted to search for the

meta-analytic map of the DMN. The terms "language" and "executive control" were

adopted to search for the meta-analytic maps of LN and the ECN, respectively. The

meta-analytic map of the visual network was identified by the search term "visual

cortex". Upon identifying those components corresponding to the five networks, we

thresholded the *t*-statistic maps with a $p < 0.05$ (corrected for multiple comparisons

with family-wise error (FWE) correction) for visualization and making masks used in

the bellowing activation analyses.

2.4.2 BOLD signal change

To assess the modulation of network activity by external stimulation, we measured

percent changes in BOLD signal amplitudes during speech listening (the task phase)

relative to the 20s fixation period (the baseline phase). The task onset

was shifted forward by 6s to compensate for delays in the brain's hemodynamic

response. The BOLD signal changes were defined as [(*task-baseline*) / *baseline*],

where *task* and *baseline* denote the mean of time points within the task phase and the

baseline phase, respectively. We first computed the voxel-wise signal change for each

subject. Then for each network, the mean value across voxels within the thresholded

component map was calculated and a *t*-test was applied to examine whether the mean

signal change in the network differed from zero. In addition to using the time courses

of single voxels, we also estimated the BOLD signal changes based on representative

network time courses derived from a separately-conducted group ICA. Details about

this analysis were provided in the supplementary material.

Figure 1. Experimental design, behavioral performance and data analysis. (A) One

speaker told real-life stories while undergoing fMRI scanning. The audio recordings

of the stories were then played to a group of listeners (N=62) during fMRI scanning.

(B) The distribution of comprehension scores for the story across the listeners. (C)

270   The procedure of Group Independent Analysis.

2.4.3. Within-brain function connectivities

For each listener, functional connectivities between networks were assessed by

Pearson's correlation using the component time courses derived from the ICA. The r

values were converted to Fisher's z values and analyzed using *t*-tests.

2.4.4 Measurement for listener-speaker neural coupling

The listener-speaker neural couplings were measured by Pearson's correlation

coefficients between the ICA-derived network time courses from each listener's brain

and the time courses of the homologous network from the speaker's brain. The

r-values were transformed to z-values using Fisher's z transformation to normalize

their distribution. Previous work has demonstrated that listeners need time to process

the high-level information in speeches in order to get aligned with the speaker

(Stephens *et al.* 2010; Kuhlen et al. 2012; Liu et al. 2017; Liu *et al.* 2020). To account

for this effect, we repeated the inter-brain correlation analysis by shifting the time

courses of listener's network activity relative to that of the speaker by 0-14s with 2s

increments. At shift zero, the listener's network activity was time-locked to the

speaker's vocalization. Two-tailed *t*-tests were applied to assess whether the neural

couplings differed significantly from zero at the group level. Multiple comparisons

were corrected with an FDR q = 0.01 using the Benjamini–Hochberg procedure

(Benjamini and Hochberg 1995).

2.4.5 Brain-behavioral correlation

The Pearson's partial correlation analysis was applied to assess the correlation

between the coupling strength of each network and listeners' comprehension scores,

controlling for the potential effect of memory span. To evaluate the generalizability of

the brain-behavior relationship we may find, we further performed within-data set

cross-validation (Shen et al. 2017). Specifically, a linear regression model was built

on the data from all participants but one, taking inter-brain couplings in the networks

which showed significant brain-behavior correlations as the predictors and listener's

comprehension score as the outcome. Then the model was used to predict the

comprehension score of the left-out participant based on his or her neural coupling

with the speaker. The significance of the prediction was assessed via permutations. In

each permutation (total N = 1000), we randomized the comprehension scores across

the training sample and re-conducted the prediction. A p-value was calculated as the

percentage of permutations that showed a lower r-value than the actual value.

2.4.6 Dynamic Causal Modeling for network communication

Functional connectivities only capture the statistical dependencies between two

signals (Friston 2011). To gain insights into the organization of functional

communication among the four networks during speech comprehension, we further

conducted Dynamic Causal Modeling analysis using the DCM toolbox in SPM12.

Given previous reports that language is processed in the brain in a hierarchal manner

(Davis and Johnsrude 2003; Lerner et al. 2011; Ding et al. 2016), we presumed that

the four networks were organized into layers. Four models consisted of three layers

with varying components in those layers were tested. Model 1 hypothesized the two

externally-directed networks (the LN and ECN) lied in the same layer, having causal

interactions with the aDMN in the intermediate layer, which in turn having causal

interactions with the pDMN. Model 2 differed from model 1 in that the positions of

aDMN and pDMN were exchanged. Model 3 hypothesized the two DMN subsystems

lied in the same layer, having causal interactions with the ECN in the intermediate

layer, which in turn having causal interaction with the LN. Model 4 varied from

model 3 in that the positions of ECN and LN were exchanged. All models included

recurrent connections of each network and had no input entering the system. All

connections in these models were bidirectional. Since the speech signals presented

during the task were continuous stimuli, we fitted models based on the

frequency-domain cross-spectral density of the time courses, which is a technique

originally developed for modeling resting-state fMRI data (Friston et al. 2014). The

model with the highest probability was identified through random-effect Bayesian

Model Selection.

2.4.7 Analysis for the association between within-brain network BOLD signal

change and functional connectivity and listener-speaker neural coupling

Finally, we explored whether the functionality of DMN depends on its interaction

with other networks or on its response amplitude to stimuli. For the pDMN and

aDMN separately, we calculated the Pearson's correlation between inter-brain neural

coupling and network mean signal changes and functional connectivity within

listeners' brains.

3. Results

3.1 Behavioral results

On the story-recall task administered shortly after the scanning, participants scored on

average $91.58 \pm 8.00\%$ (Fig. 1B). For the Mongolian story (the control condition), the

interview showed that the participants had tried to comprehend the speech but all

failed.

3.2 ICA-based network detection

By measuring the spatial overlap of each ICA component with the meta-analytic maps

derived from the Neurosynth, we identified four networks of interest and a visual

network serving as a control (Fig. 2A). Two components consistently showed high

Dice scores with the meta-analytic map of DMN. The first component was mainly

comprised of the bilateral medial frontal gyrus, insular, precuneus, and angular gyrus,

peaking at the ventral mPFC (MNI coordinates: 0, 57, 3). This component matches

with the aDMN in the literature (Xu et al. 2016). The second component covered

mainly the bilateral precuneus, inferior parietal lobe, and part of the medial frontal

gyrus, peaking at the left precuneus (MNI coordinates: -3, -72, 33), which matches the

pDMN in the literature (Xu *et al.* 2016). The LN component covered mainly the

bilateral superior and middle temporal, precentral, and inferior frontal gyrus, as well

as the supplementary motor area, peaking at the right superior temporal gyrus (MNI

coordinate: 57, -24, -3). The ECN component was primarily comprised of the bilateral

middle/inferior frontal gyrus and the superior/inferior parietal lobule, peaking at the

right inferior frontal operculum (MNI coordinates: 51,15, 27). The control (visual)

network covered mainly the bilateral inferior and middle occipital gyrus, fusiform and

lingual gyrus, peaking at the left middle occipital gyrus (MNI coordinate: -24, -96, 0).

The Dice scores and the spatial maps of other components derived from the group

ICA were presented in the supplementary material (Fig. S1 and Table S1).

3.3 Within-brain activation pattern

In line with the previous studies (Seghier and Price 2012a; Rodriguez Moreno et al.

2014), t-tests on the mean signal change over voxels in each of the four networks

revealed significant activations in the LN ($t_{(61)}$= 6.21, $p < 10^{-7}$) and significant

deactivation in the pDMN ($t_{(61)}$ = -2.35, $p = 0.02$) during speech listening relative to

the fixation period. No significant signal change was observed in the aDMN ($t_{(61)}$ =

-0.05, one-tailed $p = 0.96$) or the ECN ($t_{(61)}$ =-0.46, $p = 0.64$) (Fig. 2A). Consistent

results were obtained when the signal change was assessed using the time courses

derived from the group ICA (Fig. S2).

We note the baseline used to calculate the BOLD signal change might be too

short. To assess this effect, we compared this method with a conventional General

Linear Modeling (GLM) method which included longer baselines using an

independent dataset. The results showed that the activation and deactivation patterns

obtained by the two approaches were similar, with the one obtained from the GLM

being statistically more significant (Fig. S3). Thus, a relatively short baseline does not

compromise the findings of opposite activation patterns between the pDMN and LN.

3.4 Within-brain functional connectivity

Consistent with previous studies (Uddin *et al.* 2009; Smith *et al.* 2012), we observed

negative functional connectivities of the pDMN with both the LN ($t_{(61)}$=-6.91, $p < 10^{-8}$)

and ECN ($t_{(61)}$ = -4.13, p < 10^{-3}) within the listeners' brains. The aDMN also exhibited

negative connectivity with the ECN, but positive connectivity with the LN. The two

DMN subsystems showed strong positive functional connectivity ($t_{(61)}$=13.74, $p < 10^{-9}$)

(Fig. 2B).

Figure 2. Network activation and functional connectivity pattern. (A) The spatial

distribution of networks derived from the group ICA and their activation patterns. (B).

Functional connectivities between networks. All connectivities were significant with

an FDR-corrected *p* < 0.01. The error bar denotes the 95% confidence interval of the

group mean. LN: language network; ECN: executive control network; aDMN:

anterior default mode network; pDMN: posterior default mode network.

3.5 Listener-speaker neural coupling

In all four networks, listeners' network activities were significantly coupled with the

activities of the homologous networks in the speaker's brain (Fig. 3). Consistent with

previous findings (Stephens *et al.* 2010; Liu *et al.* 2017), the coupled activity in the

listeners' brains primarily lagged by that in the speaker's brain. The temporal profile

of inter-brain coupling differed across the four networks. In the LN, the listener's

coupling responses occurred in synchronization with the speaker's vocalization (0s

delay) and peaked with a delay of 4s. In the ECN, aDMN, and pDMN, listener's

coupling responses occurred with a delay of 2s and reached the peak with a delay of

4s, 6s, and 6s, respectively. Among the four networks, the inter-brain coupling in the

pDMN was strongest and decayed most slowly. Supplementary analyses demonstrated

that the temporal pattern of inter-brain coupling was not accounted for by the

autocorrelations of brain network signals (Fig. S4). To validate that the inter-brain

coupling in DMN was not an epiphenomenon of task-disengagement, we compared

the inter-brain coupling during the first half of communication with that during the

second half. We found the coupling strength in the pDMN increased significantly with

the unfolding of the communication (Fig. S5).

In comparison, no significant inter-brain coupling was observed in the visual

network at any lag, suggesting the inter-brain coupling was specific to task-related

brain activities. When communication failed, none of the five networks presented

significant listener-speaker neural coupling at any lag, suggesting the observed

interbrain couplings were not simply driven by low-level acoustic inputs shared

between the speaker and listener. These results also mitigate a methodological

concern that the correlation of network time courses between the listener and the

speaker was simply resulted from the manipulation that the data from both sides had

been pooled together to identify the network component during the group ICA. We

note that there was significant ISC among the listeners in the visual network.

Moreover, the listeners exhibited significant ISCs in all four networks even when they

could not understand the Mongolian story, despite the strength of ISC was

significantly lower than that during successful comprehension (Supplementary Fig.

S6).

Figure 3. Listener-speaker neural couplings as a function of temporal lags. When

communication was successful, listeners' brain activities in the four networks were

significantly coupled with (i.e., correlated over time) the activities of the homologous

networks in the speaker's brain, with several temporal lags. In comparison, no

significant inter-brain coupling was found in the visual network. Those inter-brain

couplings disappeared when communication failed. Asterisks denote significant

differences from zero, with the threshold of $p < 0.01$, FDR corrected.

3.6 Listener-speaker neural coupling predicted speech comprehension

To determine the behavioral significance of listener-speaker neural coupling, we

examined its association with the listener's level of speech understanding. A

significant positive correlation was found between the coupling strength of pDMN (at

lag 6s) and listeners' comprehension scores (partial $r_{(60)} = 0.32$, p $= 0.012$, with

memory span regressed out). There was also a positive correlation between the

inter-brain coupling in LN (at lag 6s) and comprehension scores (partial $r_{(60)} = 0.31$, p

$= 0.015$, with memory span regressed out) (Fig. 4). No significant correlation was

found between the coupling strength of the aDMN or the ECN and the comprehension

scores ($p > 0.1$).

A leave-one-out cross-validation procedure was performed to evaluate the

generalizability of the brain-behavioral relation. In each fold of cross-validation, a

regression model was built which took the listener-speaker neural coupling in both the

pDMN and LN as inputs and generated predictions of comprehension scores in a

novel participant. The predicted scores were positively correlated with actual scores

($r_{(60)} = 0.28$, p $= 0.009$) (Fig. 4), with predictive power significantly better than the

results from 1000 iterations of permutation tests.

Figure 4. The strength of listener-speaker neural coupling predicts speech

comprehension. Left: significant correlations between speech understanding and

listener-speaker neural couplings in the pDMN and LN, with memory capacity

controlled. Right: Cross-validation for the brain-behavior relationship using a

451   level-one-out procedure.

3.7 Inter-network communication revealed by DCM

Results from the two-brain analyses demonstrated that the aDMN, pDMN, LN, and

the ECN were actively engaged in language comprehension. We next carried out the

DCM analysis to investigate the organization of functional communication among the

four networks within the listeners' brains. The Bayesian Model Selection revealed that

model 1 outperformed the other three models. This model hypothesized that the LN

and ECN lied in the same layer; both networks then had causal interactions with the

aDMN in the intermediate layer, which in turn causally interacted with the pDMN.

The posterior probability for model 1 was 0.39, which reflects how likely this model

generated the data of a randomly selected subject for this model. The posterior

exceedance probability was 0.85, which measured how one model is more likely than

any other model (Fig. 5). The group mean of effective connection for the selected

model and statistic information were summarized in Table S1.

Figure 5: The organization of network communication revealed by Dynamic Causal

Modeling. Model 1 was preferred by the random-effect Bayesian model selection.

3.8 Inter-brain coupling in the pDMN increased with greater pDMN-ECN

anticorrelation

To gain more insights into the functional properties of DMN, we investigated whether

the engagements of aDMN and pDMN in language comprehension (as indexed by the

listener-speaker neural coupling) dependents on their interaction with other networks

or on the response amplitudes to the speech signals. A significant negative correlation

($r_{(60)}$ = -0.36, p = 0.004) was found between the coupling strength of the pDMN and

the pDMN-ECN functional connectivity within listeners' brains. That is, listeners with

greater pDMN-ECN anticorrelation showed stronger neural coupling with the speaker

in the pDMN. This association was also found between the change (successful >

failed) of pDMN-ECN functional connectivity and the change of pDMN coupling

strength ($r_{(57)}$ = -0.32, p = 0.012). There was no significant correlation between the

inter-brain coupling and the BOLD signal change in the pDMN (Fig. 6). These results

suggest the pDMN's functionality depends on its interaction with other networks

rather than its level of activation. For the aDMN, no significant association between

the inter-brain coupling and network connectivity was found, neither between the

inter-brain coupling and BOLD signal change.

Figure 6. The functionality of pDMN depends on its interactions with other networks

rather than its level of activation. Left: Listener-speaker neural coupling in the pDMN

was not correlated significantly with the pDMN's signal change within the listeners'

brains. Right: Stronger listener-speaker neural coupling in the pDMN was associated

with greater pDMN-ECN anticorrelation. The change (successful communication >

failed) in coupling strength was also correlated with the change in pDMN-ECN

functional connectivity.

3.9 Reproducing results with group ICA of different dimensions

To evaluate the robustness of our findings, we reconducted the above analyses using

time series derived from a group ICA which separated the spatial map of the brain

into 30 independent components. The major results, including the listener-speaker

neural coupling, the positive correlation of coupling strength with speech

comprehension, the correlation between the interbrain coupling and pDMN-ECN

anticorrelation, and the organization of network communication revealed by DCM,

are largely consistent across the two cases (Supplementary Fig.S7-S9).

4. Discussion

Taking a "two-brain" approach and applying group ICA, this study investigated

whether and how the aDMN and pDMN contribute to naturalistic speech

comprehension. Consistent with previous findings (Wirth et al. 2011; Rodriguez

Moreno *et al.* 2014; Humphreys *et al.* 2015; Jackson et al. 2019), we found the DMN

exhibited largely unchanged (in the aDMN) or reduced (in the pDMN) activities while

the LN exhibited significantly increased activities during speech comprehension

relative to the low-level baseline. Besides, the network dynamics of pDMN were

anticorrelated with both the LN and ECN. However, further two-brain analyses

revealed that the dynamics of the aDMN and pDMN in the listener's brain were

tightly coupled with the dynamics of the homologous network in the speaker's brain.

Significant listener-speaker neural couplings were also observed in the LN and ECN,

but were weaker and decayed faster than that in the pDMN. Moreover, the strength of

inter-brain neural coupling in the pDMN and LN predicted the degree to which

listeners comprehended the story. In addition, the stronger inter-brain coupling in the

pDMN was associated with greater pDMN-ECN anticorrelation. Note, when

communication failed, the listener-speaker neural couplings in those networks

disappeared; however, there were still significant ISC across the listeners, suggesting

the "two-brain" measurement had a higher specificity to task-related neural process.

Finally, applying DCM we uncovered a network communication pattern wherein the

ECN and LN had direct interactions with the aDMN, which in turn interacted with the

pDMN. Implications of these findings are discussed below.

The active role of "task-deactivated" DMN in speech comprehension

According to the prevalent view, exhibiting activity decrease and anticorrelation with

the LN and ECN would indicate that the pDMN interferes with the ongoing speech

processing and is suppressed in order to support the externally-directed cognitive

processes (Anticevic 2012; Gauffin *et al.* 2013; Zhou *et al.* 2016). In contrast to this

notion, the results from the two-brain analyses suggest an active role of the pDMN in

spoken narrative comprehension. The inter-brain coupling in the pDMN was unlikely

an epiphenomenon of task disengagement, due to the three observations. First, speech

production is typically cognitively more demanding than speech comprehension

(Hendriks 2014). Therefore, the level of task disengagement in the speaker and that in

the listener were unlikely to fluctuate similarly over the course of communication.

Even the level of task disengagement covaried to some degree across the speaker and

listener, it would most likely result in synchronized rather than listener-delayed

coupling in the pDMN as observed in this study. Second, we found the inter-brain

coupling in the pDMN during the second half of the communication was significantly

stronger than that of the first half. If the pDMN was disengaged in the ongoing task, it

should not be influenced by the length of communication. Third, the individual

difference in the coupling strength of the pDMN was associated with the individual

difference in speech comprehension.

The significant inter-brain coupling and the positive association with task

performance demonstrate that the pDMN was not suppressed despite its negative

functional connectivity with the ECN and LN. Moreover, we found that the greater

pDMN-ECN anticorrelation in the listeners' brains was associated with stronger

listener-speaker neural couplings in the pDMN, indicating that the functional

segregation with the executive control system was beneficial for the pDMN's

engagement in speech comprehension. These findings provide empirical evidence

supporting the view that, instead of representing simply an antagonistic relationship,

network anticorrelations may represent a "division of labor" wherein anticorrelated

networks could sometimes work together to complete the same task (Fransson 2006).

Significant inter-brain coupling was also observed in the aDMN. Differing from

the pDMN, the aDMN showed no significant change in activity amplitudes during

speech listening relative to the fixation period and had positive functional

connectivity with the LN. Compared to the pDMN, the inter-brain coupling in the

aDMN was weaker, decayed quicker and not significantly correlated with speech

comprehension. Consistent with previous studies (e.g., Uddin *et al.* 2009; Seghier and

Price 2012b), these results support that the aDMN and pDMN are functionally

heterogeneous, possibly engaged in the different stages of language processing with

different computations.

Differences in inter-brain coupling profile among the four networks

In addition to the aDMN and pDMN, we also observed significant listener-speaker

coupling in the LN and ECN. In contrast, there was no significant inter-brain coupling

in the visual network, suggesting the coupling was specific to task-related neural

substrates. Across the four networks, only the LN exhibited inter-brain coupling

time-locked to the speaker's vocalization (i.e., with 0s-lag), probably driven by shared

acoustic and perceptual analyses conducted both by the speaker and listener. This is in

agreement with the knowledge that (among the four networks) the LN encodes

bottom-level perceptual information. Moreover, the coupling strength of the LN was

positively correlated with the comprehension score, confirming the critical role of this

network in language processing. We also found significant inter-brain coupling in the

ECN, supporting the view that successful communication requires the coordination of

attention between communicators(Richardson et al. 2007). Together, these results

demonstrate that LN, anterior and posterior DMN, and ECN were engaged in

narrative comprehension, conforming with the proposed tri-model network of

semantic processing (Xu et al. 2017).

Among the four networks, the inter-brain coupling in the pDMN was strongest

and decayed most slowly. This distinct feature indicates that the pDMN probably lies

at the top of the cortical hierarchy for language processing. Natural communication

unfolds over time: the speaker develops a concept and intention, retrieves the

lexical-syntactic forms, organizes them into sentences based on grammar principles,

and finally produce utterances; the listener analyzes the sounds, maps utterances into

meaning, integrates words into sentences, and builds a conceptual-intentional model

resembling the speaker's. Compared to the bottom-level information, the top-level

conceptual-intentional information takes relatively longer time for the speaker to

encode and for the listener to decode, resulting in the greater delay in the inter-brain

coupling of pDMN. Besides, those top-level representations (such as abstract concepts

and situation models) may vary less across speech production and comprehension

than did the bottom-level representations (such as motor versus acoustic), resulting in

stronger listener-speaker neural couplings in the pDMN than in the LN.

The organization of network communication

The DCM analysis revealed that, compared to the other three possible models, it was

more likely that the two externally-oriented networks (the LN and ECN) lied within

the same layer, having casual interactions with the aDMN in the intermediate layer,

which in turn had direct interactions with the pDMN. This interaction pattern may

reflect ordered information flow across the four networks in a hierarchical system. In

this system, the LN and ECN occupied the bottom layer, feeding information forward

to the aDMN in the intermediate layer, which then feeds information forward to the

pDMN in the top layer. Information encoded in the higher layer is also feedback to the

lower layer as a "top-down" effect. The suggestion that the pDMN occupied the top

layer of the system is in agreement with the inter-brain coupling pattern as discussed

in the above paragraph. The suggested hierarchical organization of the four networks

is also in line with previous reports on the hierarchical topology of temporal receptive

windows (TRW) (Lerner *et al.* 2011). In that study, the early auditory cortices and the

mPFC and precuneus were found to be in the low and the top of the TRW hierarchy,

respectively. Our assumption is also supported by the observation from a recent TMS

study that the nodes in the central executive network can causally regulate activities in

the medial prefrontal portion but not the posterior portion of the DMN (Chen et al.

2013). Nevertheless, further studies which directly measure information flow across

cortical regions (such as by applying the MEG) are required to test this assumption. In

addition, the DCM analysis in this study only tested four possible models of network

organization. Future studies should test more possibilities regarding the network

organization and include more extensive brain networks in the model.

The possible mechanism of the DMN's contribution to language comprehension

Currently, how the DMN may contribute to language comprehension remains elusive.

One recent proposal is that the DMN encodes information about the external stimuli

and environments that accumulated over a large time scale (Simony et al. 2016). This

perspective may still fall in the signal-centered framework. Differing from (but not

incompatible with) this view, we speculate that the DMN (mainly the posterior

subsystem) may be primarily responsible for conceptual-intentional representations

that are internal to organism, as highlighted in the theory regarding the basic design

for language faculty proposed by Berwick and colleagues (Berwick *et al.* 2013). In

that theory, those internal representations are suggested to be constructed based on the

linguistic expressions generated by a basic language module and allow for the

semantic-pragmatic interpretation, reasoning, planning, and other activities of the

internalized "mental world" (Berwick *et al.* 2013). This mechanism seems to fit well

with the observations of the current study. Operating as an internal module, the

(posterior) DMN would be highly active during task-free states and not directly

modulated by external stimulations, therefore we did not find task-induced activations

in this network during speech listening relative to the fixation periods. In addition,

computations implemented in the internal module are likely quite different from or

even opposite to those implemented in external modules, explaining the

anticorrelations of the pDMN with the LN and ECN. Finally, the aDMN in the

intermediate layer of the hierarchy may serve as a "connector" that bridges the

internal and external systems, thus this network was found to be positively correlated

with both the pDMN and LN. We speculate that such an organizational pattern of the

four networks may enable the internalization process during successful speech

comprehension in which external information is transformed into internal mental

representations.

Conclusions

This study established that, despite showing task-induced deactivation and

anticorrelation with the LN the ECN, the pDMN plays an active role in naturalistic speech comprehension, as demonstrated by the significant listener-speaker neural coupling in this network and the positive correlation of network coupling with speech comprehension. Moreover, we demonstrated that the function of DMN mainly depends on its interactions with other networks, rather than its level of activation. Finally, the DCM results together with the inter-brain coupling pattern indicate that the LN and ECN, the aDMN, and the pDMN occupied the bottom, intermediate, and top layers of a hierarchical system, respectively. We conclude that the DMN may primarily work as an internal system that cooperates with the externally-oriented networks, which may allow the transformation of external acoustic signals into internal mental representations during successful speech understanding.

Those novel findings on the functional property of DMN during task processing have important implications for pathological studies. For instance, the overactivation of DMN in elder adults may unnecessarily represent a failure to suppress task-unrelated thoughts, but a shift from a cognitive style dominated by externally-directed processing to a style with more engagement of internally-directed processing. Our study also demonstrates that the "two-brain" approach may provide a unique opportunity for researchers to understand the human brain beyond what can be captured by the "single-brain" approach.

5. Acknowledgments

This work was supported by grants from the National Natural Science Foundation of

China (NSFC:31900802, 31971036), and the Open Research Fund of the State Key

Laboratory of Cognitive Neuroscience and Learning (CNLYB1803). No conflict of

interest is declared.

6. Reference

Anticevic A. 2012. The role of default network deactivation in cognition and disease.

Trends in Cognitive Sciences 16:584-592.

Beckmann CF, Mackay CE, Filippini N, Smith SM. 2009. Group comparison of

resting-state FMRI data using multi-subject ICA and dual regression. NeuroImage

47:S148.

Behzadi Y, Restom K, Liau J, Liu TT. 2007. A component based noise correction

method (CompCor) for BOLD and perfusion based fMRI. NeuroImage 37:90-101.

Benjamini Y, Hochberg Y. 1995. Controlling the False Discovery Rate: A Practical

and Powerful Approach to Multiple Testing. Journal of the Royal Statistical

Society: Series B (Methodological) 57:289-300.

Berwick RC, Friederici AD, Chomsky N, Bolhuis JJ. 2013. Evolution, brain, and the

nature of language. Trends in Cognitive Sciences 17:89-98.

Calhoun VD, Kiehl KA, Pearlson GD. 2008. Modulation of temporally coherent brain

networks estimated using ICA at rest and during cognitive tasks. Hum Brain Mapp

29:828-838.

Carruthers P, Smith PK. 1996. Theories of theories of mind: Cambridge University

Press.

Chen AC, Oathes DJ, Chang C, Bradley T, Zhou Z-W, Williams LM, Glover GH,

Deisseroth K, Etkin A. 2013. Causal interactions between fronto-parietal central

executive and default-mode networks in humans. Proc Natl Acad Sci USA

110:19944-19949.

Davis MH, Johnsrude ISJJoN. 2003. Hierarchical processing in spoken language

comprehension. 23:3423-3431.

Dice LR. 1945. Measures of the Amount of Ecologic Association Between Species.

Ecology 26:297-302.

Ding N, Melloni L, Zhang H, Tian X, Poeppel D. 2016. Cortical tracking of

hierarchical linguistic structures in connected speech. Nat Neurosci 19:158-164.

Fox MD, Snyder AZ, Vincent JL, Corbetta M, Van Essen DC, Raichle ME. 2005. The

human brain is intrinsically organized into dynamic, anticorrelated functional

networks. Proc Natl Acad Sci U S A 102:9673-9678.

Fransson P. 2006. How default is the default mode of brain function?: Further

evidence from intrinsic BOLD signal fluctuations. Neuropsychologia

44:2836-2845.

Friston KJ. 2011. Functional and Effective Connectivity: A Review. Brain

Connectivity 1:13-36.

Friston KJ, Kahan J, Biswal B, Razi A. 2014. A DCM for resting state fMRI.

NeuroImage 94:396-407.

Friston KJ, Williams S, Howard R, Frackowiak RSJ, Turner R. 1996.

Movement-Related effects in fMRI time-series. Magn Reson Med 35:346-355.

Frith CD, Frith U. 1999. Interacting Minds--A Biological Basis. Science

286:1692-1695.

Gauffin H, van Ettinger-Veenstra H, Landtblom A-M, Ulrici D, McAllister A,

Karlsson T, Engström M. 2013. Impaired language function in generalized epilepsy:

Inadequate suppression of the default mode network. Epilepsy Behav 28:26-35.

Greicius MD, Menon V. 2004. Default-Mode Activity during a Passive Sensory Task:

Uncoupled from Deactivation but Impacting Activation. J Cogn Neurosci

16:1484-1492.

Guo CC, Kurth F, Zhou J, Mayer EA, Eickhoff SB, Kramer JH, Seeley WW. 2012.

One-year test–retest reliability of intrinsic connectivity network fMRI in older

adults. NeuroImage 61:1471-1483.

Hendriks P. 2014. Asymmetries between language production and comprehension:

Springer.

Humphreys GF, Hoffman P, Visser M, Binney RJ, Lambon Ralph MA. 2015.

Establishing task- and modality-dependent dissociations between the semantic and

default mode networks. Proc Natl Acad Sci USA 112:7857.

Jackson RL, Cloutman LL, Lambon Ralph MA. 2019. Exploring distinct default

mode and semantic networks using a systematic ICA approach. Cortex

113:279-297.

Kuhlen AK, Allefeld C, Haynes J-D. 2012. Content-specific coordination of listeners'

to speakers' EEG during communication. Front Hum Neurosci 6:266.

Lerner Y, Honey CJ, Silbert LJ, Hasson U. 2011. Topographic mapping of a hierarchy

of temporal receptive windows using a narrated story. 31:2906-2915.

Li W, Mai X, Liu C. 2014. The default mode network and social understanding of

others: what do brain connectivity studies tell us. 8.

Li Z, Li J, Hong B, Nolte G, Engel AK, Zhang D. 2021. Speaker-Listener Neural

Coupling Reveals an Adaptive Mechanism for Speech Comprehension in a Noisy

Environment. Cereb Cortex.

Liu L, Ding X, Li H, Zhou Q, Gao D, Lu C, Ding G. 2021. Reduced listener-speaker

neural coupling underlies speech understanding difficulty in older adults. Brain

Struct Funct 226:1571-1584.

Liu L, Zhang Y, Zhou Q, Garrett DD, Lu C, Chen A, Qiu J, Ding G. 2020.

Auditory–Articulatory Neural Alignment between Listener and Speaker during

Verbal Communication. Cereb Cortex 30:942-951.

Liu Y, Piazza EA, Simony E, Shewokis PA, Onaral B, Hasson U, Ayaz H. 2017.

Measuring speaker-listener neural coupling with functional near infrared

spectroscopy. Sci Rep 7:43293.

McKiernan KA, Kaufman JN, Kucera-Thompson J, Binder JR. 2003. A Parametric

Manipulation of Factors Affecting Task-induced Deactivation in Functional

Neuroimaging. J Cogn Neurosci 15:394-408.

Menenti L, Garrod S, Pickering M. 2012. Toward a neural basis of interactive

alignment in conversation. Front Hum Neurosci 6.

Pamilo S, Malinen S, Hlushchuk Y, Seppä M, Tikka P, Hari R. 2012. Functional

Subdivision of Group-ICA Results of fMRI Data Collected during Cinema

Viewing. PLoS One 7:e42000.

Richardson DC, Dale R, Kirkham NZJPs. 2007. The art of conversation is

755   coordination. 18:407-413.

756   Rodriguez Moreno D, Schiff ND, Hirsch J. 2014. Negative Blood Oxygen Level

757   Dependent Signals During Speech Comprehension. Brain Connectivity 5:232-244.

758   Schoot L, Hagoort P, Segaert K. 2016. What can we learn from a two-brain approach

759   to verbal interaction? Neurosci Biobehav Rev 68:454-459.

760   Seghier M, Price C. 2012a. Functional Heterogeneity within the Default Network

761   during Semantic Processing and Speech Production. 3.

762   Seghier ML, Price CJ. 2012b. Functional heterogeneity within the default network

763   during semantic processing and speech production. Front Psychol 3:281.

764   Shen X, Finn ES, Scheinost D, Rosenberg MD, Chun MM, Papademetris X,

765   Constable RT. 2017. Using connectome-based predictive modeling to predict

766   individual behavior from brain connectivity. Nat Protoc 12:506-518.

767   Simony E, Honey CJ, Chen J, Lositsky O, Yeshurun Y, Wiesel A, Hasson U. 2016.

768   Dynamic reconfiguration of the default mode network during narrative

769   comprehension. Nat Commun 7:12141.

770   Smith SM, Miller KL, Moeller S, Xu J, Auerbach EJ, Woolrich MW, Beckmann CF,

771   Jenkinson M, Andersson J, Glasser MF, Van Essen DC, Feinberg DA, Yacoub ES,

772   Ugurbil K. 2012. Temporally-independent functional modes of spontaneous brain

773   activity. Proc Natl Acad Sci USA 109:3131-3136.

Spreng RN, Grady CL. 2009. Patterns of Brain Activity Supporting Autobiographical

Memory, Prospection, and Theory of Mind, and Their Relationship to the Default

Mode Network. J Cogn Neurosci 22:1112-1123.

Spreng RN, Mar RA, Kim ASN. 2008. The Common Neural Basis of

Autobiographical Memory, Prospection, Navigation, Theory of Mind, and the

Default Mode: A Quantitative Meta-analysis. J Cogn Neurosci 21:489-510.

Stephens GJ, Silbert LJ, Hasson U. 2010. Speaker–listener neural coupling underlies

successful communication. Proceedings of the National Academy of Sciences

107:14425-14430.

Stolk A, Verhagen L, Toni I. 2016. Conceptual Alignment: How Brains Achieve

Mutual Understanding. Trends in Cognitive Science 20:180-191.

Uddin LQ, Clare Kelly AM, Biswal BB, Xavier Castellanos F, Milham MP. 2009.

Functional connectivity of default mode network components: Correlation,

anticorrelation, and causality. Hum Brain Mapp 30:625-637.

Wechsler D. 1987. WMS-R: Wechsler memory scale-revised: Psychological

Corporation.

Wilson SM, Molnar-Szakacs I, Iacoboni M. 2007. Beyond Superior Temporal Cortex:

Intersubject Correlations in Narrative Speech Comprehension. Cereb Cortex

18:230-242.

Wirth M, Jann K, Dierks T, Federspiel A, Wiest R, Horn H. 2011. Semantic memory

involvement in the default mode network: a functional neuroimaging study using

independent component analysis. NeuroImage 54:3057-3066.

Xu X, Yuan H, Lei X. 2016. Activation and Connectivity within the Default Mode

Network Contribute Independently to Future-Oriented Thought. Sci Rep 6:21001.

Xu Y, He Y, Bi Y. 2017. A Tri-network Model of Human Semantic Processing. Front

Psychol 8:1538.

Yan C-G, Wang X-D, Zuo X-N, Zang Y-F. 2016. DPABI: Data Processing & Analysis

for (Resting-State) Brain Imaging. Neuroinformatics 14:339-351.

Ye Z, Zhou X. 2009. Executive control in language processing. Neurosci Biobehav

Rev 33:1168-1177.

Yu Q, Du Y, Chen J, He H, Sui J, Pearlson G, Calhoun VD. 2017. Comparing brain

graphs in which nodes are regions of interest or independent components: A

simulation study. J Neurosci Methods 291:61-68.

Zhou L, Pu W, Wang J, Liu H, Wu G, Liu C, Mwansisya TE, Tao H, Chen X, Huang

X, Lv D, Xue Z, Shan B, Liu Z. 2016. Inefficient DMN Suppression in

Schizophrenia Patients with Impaired Cognitive Function but not Patients with

Preserved Cognitive Function. Sci Rep 6:21657.

Zuo X-N, Kelly C, Adelstein JS, Klein DF, Castellanos FX, Milham MP. 2010.

Reliable intrinsic connectivity networks: test–retest evaluation using ICA and dual

regression approach. NeuroImage 49:2163-2177.

Zwaan RA, Radvansky GA. 1998. Situation models in language comprehension and

memory. Psychol Bull 123:162-185.

817