The genome sequence of the cereal pest *Sitophilus oryzae*: an unprecedented transposable element content

PARISOT Nicolas[1,$], VARGAS-CHAVEZ Carlos[1,2,†,$], GOUBERT Clément[3,4,‡,$], BAA-PUYOULET Patrice[1], BALMAND Séverine[1], BERANGER Louis[1], BLANC Caroline[1], BONNAMOUR Aymeric[1], BOULESTEIX Matthieu[3], BURLET Nelly[3], CALEVRO Federica[1], CALLAERTS Patrick[5], CHANCY Théo[1], CHARLES Hubert[1,6], COLELLA Stefano[1,§], DA SILVA BARBOSA André[7], DELL'AGLIO Elisa[1], DI GENOVA Alex[3,6,8], FEBVAY Gérard[1], GABALDON Toni[9,10,11], GALVÃO FERRARINI Mariana[1], GERBER Alexandra[12], GILLET Benjamin[13], HUBLEY Robert[14], HUGHES Sandrine[13], JACQUIN-JOLY Emmanuelle[7], MAIRE Justin[1,–], MARCET-HOUBEN Marina[9], MASSON Florent[1,£], MESLIN Camille[7], MONTAGNE Nicolas[7], MOYA Andrés[2,15], RIBEIRO DE VASCONCELOS Ana Tereza[12], RICHARD Gautier[16], ROSEN Jeb[14], SAGOT Marie-France[3,6], SMIT Arian F.A.[14], STORER Jessica M.[14], VINCENT-MONEGAT Carole[1], VALLIER Agnès[1], VIGNERON Aurélien[1,#], ZAIDMAN-REMY Anna[1], ZAMOUM Waël[1], VIEIRA Cristina[3,6,*], REBOLLO Rita[1,*], LATORRE Amparo[2,15,*] and HEDDI Abdelaziz[1,*]

[1] Univ Lyon, INSA Lyon, INRAE, BF2I, UMR 203, 69621 Villeurbanne, France.

[2] Institute for Integrative Systems Biology (I2SySBio), Universitat de València and Spanish Research Council (CSIC), València, Spain.

[3] Laboratoire de Biométrie et Biologie Evolutive, UMR5558, Université Lyon 1, Université Lyon, Villeurbanne, France.

[4] Department of Molecular Biology and Genetics, 526 Campus Rd, Cornell University, Ithaca, New York 14853, USA.

[5] KU Leuven, University of Leuven, Department of Human Genetics, Laboratory of

Behavioral and Developmental Genetics, B-3000, Leuven, Belgium.

[6] ERABLE European Team, INRIA, Rhône-Alpes, France.

[7] INRAE, Sorbonne Université, CNRS, IRD, UPEC, Université de Paris, Institute of Ecology

and Environmental Sciences of Paris, Versailles, France.

[8] Instituto de Ciencias de la Ingeniería, Universidad de O'Higgins, Rancagua, Chile.

[9] Life Sciences, Barcelona Supercomputing Centre (BSC-CNS), Barcelona, Spain.

[10] Mechanisms of Disease. Institute for Research in Biomedicine (IRB), Barcelona, Spain.

[11] Institut Catalan de Recerca i Estudis Avançats (ICREA), Barcelona, Spain.

[12] Laboratório de Bioinformática, Laboratório Nacional de Computação Científica, Petrópolis,

Brazil.

[13] Institut de Génomique Fonctionnelle de Lyon (IGFL), Université de Lyon, Ecole Normale

Supérieure de Lyon, CNRS UMR 5242, Lyon, France.

[14] Institute for Systems Biology, Seattle, WA, USA.

[15] Foundation for the Promotion of Sanitary and Biomedical Research of Valencian

Community (FISABIO), València, Spain.

[16] IGEPP, INRAE, Institut Agro, Université de Rennes, Domaine de la Motte, 35653 Le Rheu,

France.

[†] Present address: Institute of Evolutionary Biology (IBE), CSIC-Universitat Pompeu Fabra,

Barcelona, Spain.

[‡] Present address: Human Genetics, McGill University, Montreal, QC, Canada.

§ Present address: LSTM, Laboratoire des Symbioses Tropicales et Méditerranéennes, IRD,

CIRAD, INRAE, SupAgro, Univ Montpellier, Montpellier, France.

£ Present address: Global Health Institute, School of Life Sciences, Ecole Polytechnique

Fédérale de Lausanne (EPFL), Lausanne 1015, Switzerland.

⁻ Present address: School of BioSciences, The University of Melbourne, Parkville, VIC 3010,

Australia.

Present address: Department of Evolutionary Ecology, Institute for Organismic and

Molecular Evolution, Johannes Gutenberg University, 55128 Mainz, Germany.

$ Authors contributed equally to this work.

* Authors contributed equally to this work.

NP: nicolas.parisot@insa-lyon.fr

CV-C: carlos.vargas@uv.es

CG: goubert.clement@gmail.com

PB-P: patrice.baa-puyoulet@inrae.fr

SB: severine.balmand@inrae.fr

LB: beranger.louis.bio@gmail.com

CB: blancbillard.caroline@gmail.com

AB: aymeric.bonnamour@gmail.com

MB: matthieu.boulesteix@univ-lyon1.fr

NB: nelly.burlet@univ-lyon1.fr

FC: federica.calevro@insa-lyon.fr

PC: patrick.callaerts@kuleuven.be

TC: theo.chancy@hotmail.fr

HC: hubert.charles@insa-lyon.fr

SC: stefano.colella@inrae.fr

ASB: andredsb.2b@gmail.com

ED: elisa.dell-aglio@insa-lyon.fr

ADG: digenova@gmail.com

GF: gerard.febvay@inrae.fr

TG: toni.gabaldon@bsc.es

MGF: mari.ferrarini@gmail.com

AG: alegerber@lncc.br

BG: benjamin.gillet@ens-lyon.fr

RH: robert.hubley@isbscience.org

SH: sandrine.hughes@ens-lyon.fr

EJ-J: emmanuelle.joly@inrae.fr

JM: justin.maire@unimelb.edu.au

MM-H: marina.marcet@bsc.es

FM: masson.ff@gmail.com

CM: camille.meslin@inrae.fr

NM: nicolas.montagne@sorbonne-universite.fr

AM: andres.moya@uv.es

ATRV: atrv@lncc.br

GR: gautier.richard@inrae.fr

JR: jeb.rosen@isbscience.org

M-FS: Marie-France.Sagot@inria.fr

AFAS: arian.smit@isbscience.org

JMS: jessica.storer@isbscience.org

CV-M: carole.monegat@insa-lyon.fr

AVa: agnes.vallier@inrae.fr

AVi: avignero@uni-mainz.de

AZ-R: anna.zaidman@insa-lyon.fr

WZ: wzamoum@gmail.com

CV: cristina.vieira@univ-lyon1.fr

RR: rita.rebollo@inrae.fr

AL: amparo.latorre@uv.es

AH: abdelaziz.heddi@insa-lyon.fr

107 Abstract

108 Background

109   Among beetles, the rice weevil *Sitophilus oryzae* is one of the most important pests causing

110   extensive damage to cereal in fields and to stored grains. *S. oryzae* has an intracellular

111   symbiotic relationship (endosymbiosis) with the Gram-negative bacterium *Sodalis*

112   *pierantonius* and is a valuable model to decipher host-symbiont molecular interactions.

113 Results

We sequenced the *Sitophilus oryzae* genome using a combination of short and long reads to

produce a 770 Mbp assembly, the best for a Curculionidae species to date. We show that *S.*

oryzae has undergone successive bursts of transposable element (TE) amplification,

representing 72% of the genome. In addition, we show that many TE families are

transcriptionally active, and changes in their expression are associated with insect symbiotic

state. *S. oryzae* has undergone a high gene expansion rate, when compared to other

beetles. Reconstruction of host-symbiont metabolic networks revealed that, despite its

recent association with cereal weevils (30 Kyear), *S. pierantonius* relies on the host for

several amino acids and nucleotides to survive and to produce vitamins and essential

amino-acids required for insect development and cuticle biosynthesis.

124 Conclusions

125 In addition to being an agricultural pest and a valuable endosymbiotic system, *S. oryzae* can
126 be a remarkable model for studying TE evolution and regulation, along with the impact of
127 TEs on eukaryotic genomes.

129 Keywords

130 Coleoptera, weevil, *Sitophilus oryzae*, genome, transposable elements, endosymbiosis,
131 immunity, evolution

133 Background

Beetles account for approximately 25% of known animals, with an estimated number of 400
000 described species [1–3]. Among them, Curculionidae (true weevils) is the largest animal
family described, comprising about 70 000 species [1,4,5]. Despite being often associated
with ecological invasion and ecosystem degradation, only three Curculionidae genomes are
publicly available to date [6–8]. Among the cereal weevils, the rice weevil *Sitophilus oryzae*
is one of the most important pests of crops of high agronomic and economic importance
(wheat, maize, rice, sorghum and barley), causing extensive quantitative and qualitative
losses in field, stored grains and grain products throughout the world [9–11]. Moreover, this
insect pest is of increasing concern due to its ability to rapidly evolve resistance to
insecticides such as phosphine, a fumigant used to protect stored grains from insect pests
[12–14].
Like other holometabolous insects, the life cycle of *S. oryzae* can be divided into four
stages: egg, larva, pupa and adult (Figure 1). Females drill a small hole in the grain, deposit
a single egg and seal it with secretions from their ovipositor. Up to six eggs can be laid daily

by each female, totaling around 400 eggs over its entire lifespan [15]. Larvae develop and

pupate within the grain kernel, metamorphose, and exit the grain as adults. The whole

process takes on average 30 days [10]. Like many insects living on nutritionally poor diets,

cereal weevils permanently associate with nutritional intracellular bacteria (endosymbionts)

that supply them with nutrients that are not readily available in the grains, thereby increasing

their fitness and invasive power. The endosymbiont of *S. oryzae*, the gamma-

proteobacterium *Sodalis pierantonius* [16,17], is housed within specialized host cells, named

bacteriocytes, that group together into an organ, the bacteriome [18]. Contrasting with most

studied symbiotic insects, the association between *Sitophilus* spp*.* and *S. pierantonius* was

established recently (less than 30 000 years ago), probably following the replacement of the

ancestor endosymbiont, *Candidatus Nardonella*, in the Dryophthorinae subfamily [19,20]. As

a result, contrary to long-lasting endosymbiotic associations, the genome of *S. pierantonius*

is GC rich (56.06%), and its size is similar to that of free-living bacteria (4.5 Mbp) [16].

Moreover, it encodes genes involved in bacterial infection, including Type Three Secretion

Systems (TTSS), as well as genes encoding Microbial Associated Molecular Patterns

(MAMPs) that trigger Pattern Recognition Receptors (PRR), and are usually absent or

reduced in bacteria involved in long-lasting associations [16,21,22]. Nevertheless, many

features indicate that the genome of *S. pierantonius* is in a process of degradation, as it

contains many pseudogenes (43% of the predicted protein-coding sequences) and a large

number of mobile elements (18% of the genome size) [16,23]. Finally, it is important to note

that no other symbionts, with the exception of the familiar *Wolbachia* endosymbiont in some

strains*,* have been described in *S. oryzae*.

In order to help unravel potential adaptive functions and features that could become the

basis for identifying novel control strategies for weevils and other major insect pests, we

have undertaken the sequencing, assembly and annotation of the genome of *S. oryzae*.

Strikingly, the repeated fraction of *S. oryzae*'s genome (repeatome), composed mostly of

transposable elements (TEs), is the largest found to date in insects. TEs, the most versatile

DNA units described to date, are sequences present in multiple copies and capable of

relocating or replicating within a genome. While most observed TE insertions evolve

neutrally or are slightly deleterious, there are a number of documented cases where TEs

may facilitate host adaptation (for reviews see [24–26]). For instance, gene families involved

in xenobiotic detoxification are enriched in TEs in *Drosophila melanogaster* [27], *Plutella*

xylostella [28], a major crop pest, and *Myzus persicae*, another phytophagous insect causing

significant agronomic losses [29]. TEs have also been frequently associated with insecticide-

resistance in *Drosophila* species [30–32]. In addition, population genetics studies suggested

that more than 84 TE copies in *D. melanogaster* may play a positive role in fitness-related

traits [33], including xenobiotic resistance [32] and immune response to Gram-negative

bacteria [34].

In eukaryotes, TE content varies drastically and contributes significantly to the size and

organization of the genome. From TE-rich genomes as maize (85% [35]), humans (≈45%

[36]), and the recently sequenced lungfish (≈90% [37]) for instance, to TE-poor genomes, as

D. melanogaster (12-15% [38]), or *Arabidopsis thaliana* (≈10% [39]) repeatomes thrive on a

high level of diversity. These drastic variations are also observed within animal clades, such

as insects, where the proportion of TE ranges from 2% in the Antarctic midge (*Belgica*

antarctica) to 65% in the migratory locust (*Locusta migratoria*) [40–42]. In addition to the

overall TE content, the number of different TE families (homogeneous groups of

phylogenetically related TE sequences), their size (number of copies per family) and

sequence diversity are also very high among insect species [43]. For instance, SINE (Short

INterspersed Element) are almost absent from most insect genomes, but many

lepidopterans harbor these elements [43]. In flies, LTR (Long Terminal Repeats

retrotransposons) are a staple of the *Drosophila* genus, but such TEs are nearly absent from

other dipteran genomes (*e.g. Glossina brevipalpis* and *Megaselia scalaris*) [43]. Recent

advances in sequencing have dramatically increased the level to which TEs can be studied

across species and reveal that such variations can persist even within recently diverged

groups, as observed within *Drosophila* species [44] or among *Heliconius* butterflies [45]. An

increasing number of insect genomes are reported with large repeatomes (*e.g. Aedes*

aegypti and *Ae. albopictus* 40-50% [46,47], *L. migratoria* 60-65% [40,41], *Dendrolimus*

punctatus 56% [48]). Here we present the genome of *S. oryzae*, with a strong focus on the

repeatome, its largest genomic compartment, spanning over ≈72% of the assembly. *S.*

oryzae represents a model system for stored grain pests, host-TE evolutionary biology, and

the study of the molecular mechanisms acting at the early steps of symbiogenesis.

Moreover, the features uncovered suggest that *S. oryzae* and its relatives have the potential

to become a platform to study the interplay between TEs, host genomes and symbionts.

212 Results and discussion

213 Genome assembly and annotation

We have sequenced and assembled the genome of the rice weevil *S. oryzae* at a base

coverage depth of 142X using a combination of short and long read strategies (see

Methods). The karyotype of *S. oryzae* comprises 22 chromosomes [49], and the genome

assembly consists of 2 025 scaffolds spanning 770 Mbp with a N50 of 2.86 Mbp,

demonstrating a high contiguity compared to other Coleopteran genomes (Table 1). The

assembly size is consistent with the genome size measured through flow cytometry (769

Mbp in females and 768 Mbp in males [49]). We assessed the completeness of the genome

assembly using BUSCO [50] (97.9% complete and 0.7% fragmented), and along with the

aforementioned statistics, *S. oryzae* is the best assembled Curculionidae genome to date

[7,51,52] (Table 1). The complete analysis of gene content and function can be found in

Additional file 1.

Table 1. Assembly statistics of *S. oryzae*'s genome in comparison to Curculionidae genomes

and *T. castaneum*

Statistics	*Sitophilus oryzae*	*Rhynchophorus ferrugineus [8]*	*Hypothenemus hampei [6]*	*Dendroctonus ponderosae [7]*	*Tribolium castaneum [52]*
Order, Family	Coleoptera, Curculionidae	Coleoptera, Curculionidae	Coleoptera, Curculionidae	Coleoptera, Curculionidae	Coleoptera, Tenebrionidae
No. chromosomes	2n=22 [49]	2n=22 [53]	2n=14 [54]	2n=24 [55]	2n=20 [56]
No. scaffolds	2,025	4,807	15,896	8,188	2,149
Total length (Mb)	770	782	151	253	166
Scaffold N50 (Kb)	2,861	64,117	39	629	4,456
GC%	32.9	30.5	27.8	38.4	35.2
Gap length (Mb)	12.6	40.6	20.9	51.0	13.5
Median coverage	142×	108×	100×	443×	-
BUSCO (% complete/partial)	98/99	92/94	97/98	96/97	99/100
No. protein-coding genes	15,057	25,567*	19,222*	13,021	12,862

228 *All genes, no NCBI RefSeq annotation report available.*

Annotation of the *Sitophilus oryzae* genome

231 Among the different pathways we were able to decipher in the genome of *S. oryzae*, we here

232 present highlights of the main annotation efforts, followed by a detailed analysis of the TE

233 content and impact on the host genome. A comprehensive analysis for each highlight is

234 presented as Supplemental Notes in Additional file 1.

 Phylome and horizontal gene transfer

Sitophilus oryzae has a high gene expansion rate when compared to other beetles. Some of

the families with the largest expansions included genes coding for proteins with DNA binding

motifs, potentially regulating functions specific to this clade. Olfactory receptors,

antimicrobial peptides (AMPs) and P450 cytochromes were expanded as well, probably in

response to their ecological niche and lifestyle. Additionally, we noticed an expansion of

plant cell wall degrading enzymes that originated from horizontal gene transfer (HGT) events

from both bacteria and fungi. Given the intimate relationship between *S. oryzae* and its

endosymbiont, including the permanent infection of the female germline, we searched for

evidence for HGT in the weevil genome possibly coming from *S. pierantonius*. Contrary to

the genome of the tsetse fly *Glossina,* where at least three HGT events from *Wolbachia*

have been reported [57], we were unable to pinpoint any HGT event from either the ancient

endosymbiont *Nardonella, Wolbachia,* or the recently acquired *S. pierantonius.* A detailed

description is reported in Additional file 1: Supplemental Note 1, and Note 3 for digestive

enzymes.

Global analysis of metabolic pathways

Using the CycADS [58] pipeline and Pathway Tools [59] we have generated BioCyc

metabolism reconstruction databases for *S. oryzae* and its endosymbiont *S. pierantonius*.

We compared *S. oryzae* metabolism to that of other arthropods present in the

ArthropodaCyc collection and we explored the metabolic exchanges between weevils and

their endosymbionts (see Additional file 1: Supplemental Note 2). The metabolic

reconstruction reveals that, despite its large genome for an endosymbiotic bacterium, *S.*

pierantonius relies on its host for several central compounds, including alanine and proline,

but also isocitrate, Inosine MonoPhosphate (IMP) and Uridine MonoPhosphate (UMP), to

produce essential molecules to weevils, including the essential amino acids tryptophan,

phenylalanine, lysine and arginine, the vitamins pantothenate, riboflavin and dihydropteroate

as a folate precursor, as well as nicotinamide adenine dinucleotide (NAD) (Additional file 1:

Supplemental Note 2). Among the amino acids listed above, phenylalanine, in particular is

an essential precursor for the cuticle synthesis in emerging adults [60]. In addition, several

studies have shown that *S. pierantonius* improves host fitness, including fertility,

developmental time and flight capacity, in part by supplying the host with vitamins and

improving its mitochondrial energy metabolism [61–63].

Development

The annotation of developmental genes revealed a high level of conservation in comparison

to the red flour beetle *Tribolium castaneum*, a model coleopteran. When compared to *D.*

melanogaster, several key coordinate group genes are absent in *T. castaneum* and *S.*

oryzae. Moreover, a number of genes with two homologs in the *Drosophila* genome are

represented by a single ortholog in *T. castaneum* and *S. oryzae*. We also observed that

homologs for several signaling pathway ligands could not be identified, which, given the

presence of conserved receptors, is probably due to divergent primary sequence of the

ligands. A detailed description is reported in Additional file 1: Supplemental Note 4.

Cuticle protein genes

Among the distinctive biological features of coleopterans is the ability to generate a hard and

thick cuticle that protects them against dehydration and represents the first physical barrier

from infections and topical insecticide penetration. The analysis of cuticle proteins (CPs)

showed that *S. oryzae* has an average number of CPs, but with an enrichment of members

of the CPAP1 family. While some members of this family are known to be involved in molting

and maintaining the integrity of the cuticle in *T. castaneum*, most are still uncharacterized

[64,65]. Thus, these proteins might be involved in the development of specific cuticular

tissues in *S. oryzae* or other weevils. The total number of CPs did not follow the taxonomy of

beetles, suggesting instead that it might be an adaptation to their diverse lifestyles. For

details see Additional file 1: Supplemental Note 5.

Innate immune system

The analysis of immunity-related genes revealed that the genome of *S. oryzae* encodes the canonical genes involved in the three main antimicrobial pathways Toll, Imd and JAK-STAT, suggesting functional conservation of these pathways in cereal weevils. The conservation of the Imd pathway in the *S. oryzae* genome is of particular interest as its degradation in other symbiotic insects (*Acyrthosiphon pisum* [66], *B. tabaci* [67], or *Rhodnius prolixus* [68] was initially correlated to their symbiotic status. The Imd pathway is not only present in *S. oryzae*, but it is also functional [69,70] and has evolved molecular features necessary for endosymbiont control [69] and host immune homeostasis [70]. Thus, not only is the Imd pathway conserved in cereal weevils, contrary to aphids and some other hemimetabolous insects, but it seems to have been evolutionary "rewired" toward additional functions in symbiotic homeostasis [69]. A detailed description can be seen in Additional file 1: Supplemental Note 6.

Detoxification and insecticide resistance

Fumigation using phosphine, hydrogen phosphide gas (PH_3), is by far the most widely used treatment for the protection of stored grains against insect pests due to its ease of use, low cost, and universal acceptance as a residue-free treatment [71,72]. However, high-level resistance to this fumigant has been reported in *S. oryzae* from different countries [13,73–80]. Hence, we searched for genes associated with detoxification and resistance to insecticide and more generally to toxins, including plant allelochemical. The *S. oryzae* repertoire of detoxification and insecticide resistance genes includes more than 300 candidates, similar to what is seen in other coleopteran genomes. For more details see Additional file 1: Supplemental Note 7.

Odorant receptors

One promising pest management strategy relies on modifying insect behavior through the use of volatile organic compounds that act on odorant receptors (ORs) [81,82]. ORs play a

significant role in many crucial behaviors in insects by mediating host-seeking behavior,

mating, oviposition, and predator avoidance [83]. Interfering with the behavior of pest insects

and modulating their ability to find suitable hosts and mates has been proven to reduce

population numbers, notably using plants that are capable of producing attractants and

repellents [84,85]. *Sitophilus* spp. are known to use kairomones for host detection [86,87], as

well as aggregation pheromones [88,89]. We annotated 100 candidate OR genes in *S.*

oryzae (named SoryORs), including the gene encoding the co-receptor Orco. Of these

genes, 46 were predicted to encode a full-length sequence. The global size of the SoryOR

gene repertoire is in the range of what has been described in other species of the

coleopteran suborder Polyphaga (between 46 in *Agrilus planipennis* and greater than 300 in

T. castaneum) and close to the number of OR genes annotated in the closely related

species *Dendroctonus ponderosae* (85 genes, [90]) (Additional file 1: Supplemental Note 8).

Massive expansion of TE copies in the genome of *S. oryzae*

Detection and annotation of the repeatome

The repeatome represents the fraction of the genome categorized as repetitive. It

encompasses TEs, satellites, tandem, and simple repeats. Eukaryotic TEs can be separated

into two classes depending on their replication mode. DNA (Class II) based elements are

able to directly move within a genome, and include terminal inverted repeat (TIR), Crypton,

rolling-circle (RC/Helitron), and large composite TIRs (Maverick) elements. Conversely,

retrotransposons (Class I) have an RNA intermediate and replicate through RNA

retrotranscription. Retrotransposons can be further divided into long terminal repeats (LTR)

elements, and non-LTR elements, including long and short interspersed nuclear repeat

elements (LINEs and SINEs). Other retrotransposons include penelope-like (PLEs) and DIR-

like elements. Each one of these TE orders can be further classified into specific

superfamilies (as for instance Copia or Gypsy LTR elements, and hAT or Tc1/Mariner DNA

elements), that may encompass hundreds of TE families, each containing thousands of

copies. The intrinsic diversity of TEs complicates TE identification and annotation, especially

in understudied species orders.

We used multiple state-of-the-art TE detection tools, including RepeatModeler2 and EDTA

[91,92] to generate consensus sequences of the TE families in *S. oryzae*. After an initial

discovery step, more than 10 000 likely redundant TE families were identified by the

dedicated programs; we combined their results using multiple sequence alignments and

clustering (see Methods and Additional file 1: Figure S1) to reduce this number to 3 399.

Due to the evolutionary distance between *S. oryzae* and other known coleopterans, the

consensus sequences obtained were further classified using a thorough combination of

sequence homology and structure (see Methods).

The *S. oryzae* genome is among the most TE-rich insect genomes to date

We uncovered 570 Mbp of non-overlapping repeat sequences, corresponding to ≈74% of

the *S. oryzae* genome: ≈2% of satellite sequences, simple or low-complexity repeats, and

≈72% of other mobile elements, including TEs, (Figure 2A, Additional file 2). Given the

limitation of the sequencing technologies, the proportion of satellites and TEs usually

abundant in the heterochromatin is likely underestimated. We took advantage of a recent

comparative analysis of TE content in 62 insect species [40] to contrast with the *S. oryzae*

TE compartment. The *S. oryzae* genome ranks among those with the highest TE fraction

observed in insects (Figure 2B and 2C). Within the largest insect order, Coleoptera, very

little is known regarding TE evolution. *T. castaneum* harbors only 6% of TEs [52],

Hypothenemus hampei contains 8.2% of TEs [6,93], while *Dichotomius schiffleri* harbors

21% [94]. The species closest to *S. oryzae*, *Rhynchophorus ferrugineus,* has a TE content of

45% [8]. Therefore, while TE content has been described to follow phylogenetic relationships

in most insects [43,44], within Coleoptera, there is a large variation among the few species

studied. It is important to note that the pipeline we used to detect and annotate TEs in *S.*

oryzae differs than the method implemented by Petersen and colleagues [40], as we

incorporated 31 manually curated TE references for *S. oryzae*, and specifically annotated

DNA/TIR elements based on their sequence structure (see Methods).

DNA elements dominate *S. oryzae*'s genome

The most striking feature of the genome of *S. oryzae* is the high abundance of DNA

elements (≈33% of the genome, ≈45% of the TE content) (Figure 2A), which is the highest

observed among all 62 insect species included in this analysis [40–42]. Indeed, the most

DNA transposon-rich genomes include mosquito *Culex quinquefasciatus,* and *Ae. aegypti,*

harboring 25% and 20% of DNA transposon content in their genome, amounting to 54% and

36% of the total TE compartment, respectively [6]. Among species of insects with the large

genomes, the *S. oryzae* genome contains the highest proportion of DNA elements reported.

For instance, the TE-rich grasshopper *L. migratoria* repeatome comprises only 14% of DNA

transposons, while LINE elements amount to 25%. Finally, among Coleoptera, a large

diversity of repeatomes is observed (Figure 2C) with *A. planipennis, Leptinotarsa*

decemlineata and *Onthophagus taurus* carrying an abundant LINE content, while *S. oryzae,*

T. castaneum and *Anoplophora glabripennis* show larger DNA transposon content.

Among the DNA elements present in *S. oryzae*, the majority has not been assigned a

known superfamily (Figure 2D). Tc Mariners make up ≈6% of DNA elements, and we also

uncovered Crypton, hAT, Academ, PIF, Maverick, P, CMC, Sola, PiggyBac, MULE-MuDR,

Ginger, Zisupton and IS3EU superfamilies, although at a lower proportion. The consensus

length distribution among DNA superfamilies shows a continuum starting at a couple of

hundred bases to around ≈20 Kbp (Maverick elements, see Additional file 1: Figure S2). We

hypothesize that most of the smaller DNA families observed are miniature inverted repeat

elements (MITEs). MITEs are non-autonomous elements, deriving from autonomous DNA

copies, comprising two TIRs flanking a unique region (sometimes absent) of variable length.

While the TE detection pipeline used was able to detect and annotate most DNA elements

based on transposase homologies, we also specifically searched for TIR sequences,

allowing the detection of DNA elements that lack protein coding regions (Additional file 1:

Figure S1). Among all DNA superfamilies, TIR length varies between tens of base pairs to

≈1 Kbp (Figure 2E). We identified short DNA elements, composed mostly of their TIR

sequences (Figure 2E), typical of MITE elements. Interestingly, the unknown DNA families

show an average size smaller than 1 Kbp, while TE families that were annotated by the TE

detection pipeline, show larger sizes (Additional file 1: Figure S3), suggesting that most

unknown families could be MITE elements. Finally, the distribution of the proportions of TIR

relative to the consensus length appears superfamily-specific (Figure 2E and Additional file

1: Figure S3), and unknown families recapitulate these patterns. In conclusion, while most

unknown families seem to be composed of MITE elements, we cannot exclude that such

consensus simply lacks enough protein hits.

The most abundant TE family detected in the *S. oryzae* genome is indeed a MITE element,

with 10 486 genomic hits (≈4 117 equivalent copies - bp per consensus size),

corresponding to 1.8% of the genome. Large fractions of MITEs were also reported in Class

II-rich genomes, such as the aforementioned mosquitoes [47,96] and the invasive *Ae.*

albopictus [46]. MITE size ranges were previously described from around 100 bp to copies

reaching more than 1 Kbp [97].

LINE elements are the second most abundant TE subclass, representing ≈11% of the *S.*

oryzae genome, among which ≈35% are assigned to RTE elements and ≈22% to I elements.

We have also uncovered R1, L2, Jockey, Dong CR1 and R2 families (Figure 2D). Among the

LINE elements present in *S. oryzae*, ≈8% have not been assigned a superfamily. No SINE

families have been detected. LTRs are rather scarce, representing only ≈3% of the genome

(Figure 2D). The vast majority of LTR families belong to the Gypsy superfamily (≈30%), and

we have also uncovered PAO, Copia, and potentially novel LTR families (unknown).

Another retrotransposon order detected are PLEs, reaching nearly 2% of *S. oryzae*'s

genome. We have also detected DIR elements (0.14% of the genome), and RC/Helitrons

(0.4% of the genome). Finally, around 22% of the repeats in *S. oryzae*'s genome could not

be assigned a known TE class by our pipeline (Figure 2D).

TE copies make up most of non-coding sequences of *S. oryzae*'s genome

S. oryzae's TE copies are present scattered around the *S. oryzae* genome, with no depletion

in gene-rich regions as seen in the largest scaffolds (size larger than 10Mb, Figure 3A). TEs

are less frequently found close to gene transcription start sites (TSS), 5' untranslated regions

(5'UTRs), and exons (Figure 3B), as expected. On the contrary, 3' UTRs seem to harbor

DNA and unknown TE copies, while introns and intergenic sequences harbor the highest TE

content (Figure 3A), suggesting that most non-coding DNA sequences in the *S. oryzae*

genome are virtually made of TEs. In order to grasp the impact of TEs on intron size, we

compared intron length in *S. oryzae* with two very well assembled genomes: *D.*

melanogaster with a very compact and small genome, and the large, TE-rich human genome

(Figure 3C). In *D. melanogaster*, introns are small and harbor few TEs, while in humans,

introns are much larger potentially due to high TE accumulation [98]. *S. oryzae* intron sizes

also seem to be due, at least partly, to TE accumulation. Interestingly, the *S. oryzae* genome

also contains a large proportion of small introns, as found in *D. melanogaster.*

TE activity inferred by evolutionary history

Within reconstructed TE families, nucleotide substitution levels (Kimura 2 Parameters, K2P)

between copies and their consensus sequences allowed estimation of the relative ages of

TE families and identified potentially active ones (Figure 4A). Such "TE landscapes" are

extremely helpful to pinpoint potential TE amplifications (modes in the distribution) and

extinctions (valleys) within the 0-30% K2P range (beyond, the increased divergence

between copies affects negatively the sensitivity of the alignments, such that TE-derived

sequences are no longer recognisable). The landscape analysis revealed a heterogeneous

distribution of the TE copy divergence to their consensus within and between the main TE

subclasses (Figure 4A). Most identified TE copies have a K2P divergence under 10, which is

often observed in insects, and strikingly distinguishes itself from TE-rich mammalian

genomes (RepeatMasker.org, [40]). LINEs and DNA transposons have the wider spectrum

of divergence levels, suggesting an aggregation of distinct dynamics for the TE families

present in *S. oryzae.* By contrast, the rare LTR copies identified appear to be the most

homogeneous within families, with only a few substitutions between copies and their

consensuses, suggesting a very recent amplification of these families. The long flat tail of the

distribution may suggest that several older TEs remain to be discovered, but have been kept

under our detection capabilities due to higher levels of divergence between copies. Finally,

unknown TEs share a large part of their K2P distribution with DNA elements, though

relatively less divergent from their consensi as a whole. A breakdown of the K2P

distributions at the superfamily level reveals specific evolutionary dynamics (Figure 4B).

Diverse superfamilies, such as Tc-Mar and hAT (DNA) or RTE (LINE), show more uniform

distributions, suggesting sustained activity throughout *S. oryzae*'s genome evolution, though

this could also indicate that these subfamilies could be subdivided further. As observed at

the class level, all three identified LTR superfamilies (Pao, Gypsy and Copia) show families

with relatively low K2P.

TEs are transcriptionally active in somatic and germline tissues

The TE K2P landscape suggests that LTR elements and some families of DNA and LINE

subclasses are young, and thus potentially active. In order to estimate the transcriptional

activity of *S. oryzae*'s TE families, we have produced somatic (midgut) and germline (ovary)

transcriptomic data. While germline tissues allow identification of potential TE families

capable of producing vertically transmitted new copies, TE derepression in somatic tissues

represents the potential mutational burden due to TEs. From 1.3 to 2.2 M reads were

successfully mapped to TE families (see Additional file 3 for complete mapping statistics).

The expression of TE families varied extensively within a class and the proportion of

transcriptionally active/inactive TE families between classes was also distinct (Figure 5A). In

total, 1 594 TE families were differentially expressed between ovary and midgut tissues

(Figure 5B, Additional file 3); of which 329 had an absolute Log2 fold change higher than 2

(71 downregulated and 258 upregulated in midgut). In total, we detected 360 TE families

downregulated in midgut when compared to ovaries: 118 DNA, 35 LINE, 50 LTR, 6 PLE, and

149 Unknown. A much larger set of upregulated TE families was detected in midgut when

compared to ovaries was detected: 1 236 (4 DIRS, 464 DNA, 184 LINE, 76 LTR, 24 PLE, 4

RC, and 476 Unknown), illustrating the tighter regulation of TE copies in germline tissues.

Moreover, the distribution of Log2 fold changes were similar between DNA, LINE, PLE, and

Unknown subclasses but different for LTRs, which had a higher proportion of upregulated TE

families in ovaries compared to other classes (Figure 5C. Kruskall and Wallis rank-sum test:

H = 36.18, $P < 0.01$; LTR vs. LINE, DNA or Unknown: Dunn's test: *P-adj* < 0.01). In

conclusion, the large TE compartment in *S. oryzae* shows abundantly expressed TE

families, and tissue-specific expression patterns.

To estimate the TE transcriptional burden imposed on *S. oryzae,* we computed the

percentage of total RNA-seq poly-A enriched reads mapping to TE consensus sequences

(Additional file 1: Figure S4). Around 5% of the midgut transcriptome corresponds to TE

sequences. We compared such transcriptional burden to a TE-poor (*D. melanogaster,*

≈12%*)* and a TE-rich (*Ae. albopictus* ≈50%*)* genome, using similar technology in equivalent

tissues (adult midgut, see Methods). It is important to note that, despite being a TE-poor

genome, *D. melanogaster* harbors many young LTR elements that have been recurrently

shown to transpose [99]. We did not detect a direct correlation between genomic TE content

and TE expression (Additional file 1: Figure S4). Indeed, *S. oryzae* bears the highest

proportion of RNAseq reads mapped against TE consensi (≈5%), followed by *D.*

melanogaster (≈1%) and *Ae. albopictus* (≈0.01%). Henceforth, not only is *S. oryzae* a TE-

rich genome, but the transcriptional burden from TEs is higher than in other TE-rich

genomes (*Ae. albopictus)*, and in genomes harboring young and active TE copies (*D.*

melanogaster, [38,100]).

Finally, it is important to note that while transcriptional activation of TE copies may have an

impact on the host genome, it does not indicate high transposition and therefore higher

mutation rates. The high transcriptional burden of *S. oryzae* compared to other species

might stem from differences in TE regulation. In insects, TEs are mainly silenced by small

RNAs and repressive chromatin marks [101]. More specifically, piwi-interacting RNAs

(piRNAs) are able to target post-transcriptional repression of TEs, and guide chromatin

silencing complexes to TE copies [101–103]. Therefore, we have annotated genes

implicated in small RNA biogenesis and found that all three pathways (piRNAs but also

microRNAs and small interfering RNAs biogenesis pathways) are complete (Additional file 1:

Supplemental Note 9). Genes involved in piRNA biosynthesis are expressed mainly in

ovaries and testis, while somatic tissues (midgut) show smaller steady-state levels

(Additional file 1: Supplemental Note 9), suggesting the piRNA pathway is potentially

functional in *S. oryzae* ovaries, and could efficiently reduce transposition.

 TE content is variable among *Sitophilus* species

Cereal weevils are part of the Dryophthoridae family that includes more than 500 species.

Very little is known about genome dynamics in this massive phylogenetic group. Because of

the unusual high TE copy number and landscape observed in *S. oryzae,* we analyzed three

other closely related species namely *Sitophilus zeamais, Sitophilus granarius* and *Sitophilus*

linearis. We produced low coverage sequencing and estimated the TE content using our

annotated *S. oryzae* TE library to investigate their overall repeat composition. Remarkably,

among *Sitophilus* species, repeat content is variable (Figure 6A), with *S. linearis* harboring

the smaller repeat load (≈54%) compared to *S. oryzae* (≈80%), *S. zeamais* (≈79%)*,* and *S.*

granarius (≈65%). Most importantly, DNA elements are nearly absent from *S. linearis*, and

no recent burst of LTR elements is observed, contrary to the other *Sitophilus* species,

suggesting different TE evolutionary histories (Figure 6B). It is important to note that our

analysis is biased towards *S. oryzae,* as the library used to annotate the TEs in the other

Sitophilus species stems from automatic and manual annotation of the *S. oryzae* genome.

Finally, the relatively higher dnaPipeTE [46] estimations of the LTR content in *S. oryzae*

compared to the assembled genome supports the hypothesis that LTR elements have seen

a recent burst of transposition, as young elements tend to collapse in genome assemblies

and eventually diminish their estimated copy number.

Overall, the comparison of TE content in closely related species highlights the complexity

of TE dynamics. In addition to the regulation mechanisms that strongly contribute to TE

amount and variation, TE accumulation is conditioned by the drift/selection balance in

populations. Indeed, effective population size has been suggested to be a major variable

influencing TE content, as small, inbred or expanding populations suffer drift, allowing

detrimental insertions to stay in the gene pool and thus favor TE fixation [104]. Such

hypotheses should be addressed in the future, especially on recently sequenced TE-rich but

rather small (<1 Gbp) genomes such as *S. oryzae.*

The four *Sitophilus* species studied have different ecologies. *S. oryzae* and *S. zeamais* infest

field cereals and silos, while *S. granarius* is mainly observed in cereal-containing silos. *S.*

linearis, however, lives in a richer environment, *i.e.* tamarind seeds. In association with their

diets, the interaction of *Sitophilus* species with endosymbiotic bacteria differs: the cereal

weevils (*S. oryzae, S. zeamais* and *S. granarius*) harbor the intracellular gram-negative

bacteria *S. pierantonius*, albeit at very different loads. While *S. oryzae* and *S. zeamais* show

high bacterial load, *S. granarius* has a smaller bacterial population [60]. In contrast, *S.*

linearis has no nutritional endosymbionts, in correlation with its richer diet. We wondered

whether the presence of intracellular bacteria impacts TE regulation, and took advantage of

artificially obtained aposymbiotic *S. oryzae* animals to search for TE families differentially

expressed in symbiotic versus aposymbiotic ovaries. There were 50 TE families upregulated

in symbiotic ovaries compared to artificially obtained aposymbiotic ones, while 15 families

were downregulated (Figure 7 and Additional file 3). Only three families presented an

absolute Log2 fold change higher than 2: one LINE and two LTR/Gypsy elements. The three

of them were upregulated both in symbiotic *versus* aposymbiotic ovaries and in ovaries

versus midgut (Additional file 5), suggesting that such elements have tissue specificity, and

are impacted by the presence of intracellular bacteria. Such TE families would be ideal

candidates to further study the impact of insects and intracellular bacteria on TE

transcriptional regulation.

557   Conclusion

The success of obtaining a TE-rich genome assembly complete enough to understand

genome architecture and regulatory networks relies on the use of multiple sequencing

platforms [105]. Here, we describe the first assembly of the TE-rich (72%) *S. oryzae*

genome, based on a combination of long and short read sequencing, and a new assembly

method, WENGAN [106]. While this first assembly reaches quality standards similar to other

coleopteran species (Table 1), it is important to stress that new sequencing methods have

emerged in order to improve genome assemblies, including linked-reads and optical

mapping [105].

We uncovered around 74% of repeated sequences in the *S. oryzae* genome, mostly TE

families. While the TE landscape is marked by a wealth of DNA elements, specially the non-

autonomous MITE elements, 22% of the genome is composed of unknown repeats. Large

duplicated gene families can be present in such unknown repeats, but it is tempting to

speculate that the majority is composed of novel DNA elements. Unknown and DNA

elements share the same K2P landscapes, and many DNA elements have only been

detected through an inverted repeat search for TIRs, and not proteins, excluding therefore

DNA copies old enough that TIRs are too divergent to be recognized. Moreover, we have

shown that many TE families in *S. oryzae* are transcriptionally active, imposing a

transcriptional burden compared to other species. How such TE families are able to escape

host silencing remains unknown. It seems obvious today that insect models such as *D.*

melanogaster only represent a small window on the complex biology and evolution of TEs,

and the sequencing and annotation of species with high TE content -- while challenging

[107] -- is key to understanding how genomes, their size, their structure and their function

evolve. In conclusion, *S. oryzae* constitutes an excellent model to understand TE dynamics

and regulation and the impact on genome function.

Sitophilus species not only differ in their TE landscape, but also in their ecology and as a

consequence, their association with intracellular bacteria. Comparison of TE content within

the *Sitophilus* genus shows variable TE amount and diversity. In addition, intracellular

bacterium impacts transcription of specific TE families in ovaries. The molecular

mechanisms behind the co-evolution between an insect, its endosymbiotic bacterium and

TEs remain elusive. The impact of intracellular bacteria on host genomes is poorly studied,

and the *Sitophilus* genus offers a simpler experimental setting, with a single intracellular

bacteria present within specific host cells [19,61], and a well established knowledge of host-

bacteria interaction [60,69,70,108,109].

Methods

DNA extraction and high-throughput sequencing

Individuals of both sexes of *S. oryzae* were reared on wheat grains at 27.5 °C with 70%

relative humidity. The aposymbiotic strain was obtained by treating the symbiotic strain

during one month at 35 °C and 90% relative humidity as previously described [110]. This

strain is viable, fertile and was raised in the same conditions as the symbiotic strain. The

aposymbiotic status was confirmed by PCR and histology. Male and female adults of *S.*

oryzae were used for DNA extraction. Only the gonads were used to minimize DNA

contamination from its diet, which could be still present in the gut. The reproductive organs

were obtained from aposymbiotic adults and a DNA extraction protocol specific for *Sitophilus*

weevils was performed. DNA extractions were performed using a STE buffer (100 mM NaCl,

1 mM Na$_2$EDTA pH 8, 10 mM Tris HCI pH 8). Tissues were homogenized in STE buffer, then

treated successively by SDS 10%, proteinase K and RNase. Briefly, genomic DNA was

purified by two successive extractions with phenol:chloroform:isoamyl alcohol (25/24/1)

followed by extraction with 1 vol of chloroform:isoamyl alcohol (24/1). Genomic DNA was

then precipitated by 0.7 vol isopropanol. After washing the pellet with 70% ethanol, genomic

DNA was recovered in TE (1 mM EDTA, 10 mM Tris HCI pH8) buffer. Using this protocol, we

obtained six different DNA samples: four from males and two from females. Each sample

corresponds to the genomic DNA from 20 individuals. Five additional DNA samples were

obtained using a high molecular weight DNA extraction protocol consisting of a single

phenol:chloroform:isoamyl alcohol (25/24/1) extraction step from the genomic DNA of 100

males. The DNA concentration in each of these samples was quantified using a NanoDrop

spectrophotometer (ThermoFisher Scientific, Waltham, MA, USA).

Sequencing was performed using a combination of Illumina, PacBio and Nanopore

technologies (Additional file 4). For each sex, two Illumina libraries were generated: one

paired-end library with an average fragment size of 500 bp and one mate pair library with an

average fragment size of 5 Kbp. The libraries were sequenced using an Illumina HiSeq 2000

platform with the V3 chemistry and a read size of 101 bp; the paired-end (PE) libraries were

sequenced at the "Génomique & Microgénomique" service from ProfileXpert (Lyon, France)

while the mate paired (MP) were sequenced at Macrogen (Seoul, South Korea). Two male

samples were used to build (i) an Illumina library with an average fragment size of 200 bp

which was sequenced on a HiSeq 2500 instrument using the V4 chemistry and a read size

of 125 bp, and (ii) a PacBio library sequenced on seven SMRT cells using the P6-C4

chemistry. These two libraries were sequenced at KeyGene (Wageningen, The

Netherlands). Finally, five male samples were used to build Nanopore libraries with the SQK-

LSK109 kit and without DNA fragmentation step. The libraries were independently

sequenced on five MinION R9.4 flow cells. These libraries were built and sequenced at the

sequencing platform of the IGFL (Institut de Génomique Fonctionnelle de Lyon, Ecole

Normale Supérieure de Lyon, France). Statistics and accession numbers from all the

sequencing runs are listed in the Additional file 3.

Genome assembly and annotation

First, the Illumina reads were error-corrected using BFC release 181 [111]. The PacBio and

Nanopore reads were error-corrected using LORDEC v0.9 [112] with the error-corrected

Illumina overlapping PE reads, a k-mer size of 19 and solidity threshold of 3. Overlapping

reads were then merged using FLASH2 v2.2 [113]. Based on the merged Illumina reads, a

first short-read assembly was produced using a modified version of MINIA v3.2.1 [114] with

a k-mer length of 211. A hybrid assembly was then performed using WENGAN v0.1 [106] on

the MINIA short-read assembly and the raw Nanopore reads. The resulting assembly was

polished using two rounds of PILON v1.23 [115] using the error-corrected Illumina

overlapping PE reads and the --diploid option. A first scaffolding was then performed with

two rounds of FAST-SG v06/2019 [116] and SCAFFMATCH v0.9 [117] with the error-

corrected Illumina MP, Illumina PE, PacBio and Nanopore libraries. The LR_GAPCLOSER

algorithm v06/2019 [118] was used for the gap-filling step using the error-corrected PacBio

and Nanopore libraries. An additional scaffolding step was performed using RASCAF v1.0.2

[119] with the available RNA-seq libraries from the Sequence Read Archive (SRX1034967-

SRX1034972 and SRX3721133-SRX3721138). The resulting scaffolds were then gap-filled

using a new round of LR_GAPCLOSER as previously described followed by two rounds of

SEALER v2.1.5 [120] using the error-corrected Illumina overlapping PE reads and k-mer

sizes of 64 and 96. Two rounds of PILON, as previously described, were performed to

produce the final assembly. Quality of the assembly was assessed by computing several

metrics using i) QUAST v5.0.2 [121] with a minimal contig size of 100 bp and the --large and

-k options, ii) BUSCO v4.0.5 [50] using the Insecta ODB10 database and the -geno option,

and iii) KMC v3.0.0 [122] to evaluate the percentage of shared 100-mers between the

assembly and the merged Illumina reads.

Three contaminant scaffolds corresponding to the mitochondrial genome and an artefact

were removed from the assembly prior to the annotation step. The 'NCBI *Sitophilus oryzae*

Annotation Release 100' was produced using the NCBI Eukaryotic Genome Annotation

Pipeline v8.2.

Low-coverage genome sequencing of other *Sitophilus* species

Twenty pairs of ovaries were dissected from *S. oryzae, S. zeamais, S. granarius and S.*

linearis females. Ovaries were homogenized in 100 mM NaCl, 1 mM EDTA pH 8, 10 mM

Tris-HCl pH 8 using a small piston. Proteinase K digestion followed in the presence of SDS

for 2 h at 55 °C with shaking and for 1 h at 37 °C with RNAse A. A typical phenol chloroform

extraction was then performed and genomic DNA was isopropanol precipitated. Eight whole

genome sequencing libraries with a median insert size of 550 bp were constructed using the

Illumina TruSeq DNA PCR-free sample preparation kit (Illumina, San Diego, CA, USA),

according to manufacturer's protocols. Briefly, 2 µg of each gDNA were sheared using a

Covaris M220 Focused-ultrasonicator (Covaris, Inc. Woburn, MA, USA), end-repaired, A-

tailed, and adapter ligated. Library quality control was performed using the 2100 Bioanalyzer

System with the Agilent High Sensitivity DNA Kit (Agilent Technologies, Santa Clara, CA,

USA). The libraries were individually quantified via qPCR using a KAPA Library

Quantification Kits (Kapa Biosystems, Wilmington, MA, USA) for Illumina platforms, then

they were pooled together in equimolar quantities and sequenced in a MiSeq sequencing

system. 2x300 paired-end reads were obtained using a MiSeq Reagent Kits (600-cycles).

TE library construction

In order to annotate the *S. oryzae* repeatome, we collected and combined cutting-edge

bioinformatic tools to (i) create and (ii) classify a non-redundant library of repeated elements

(Additional file 1: Figure S1). First, we separately ran RepeatModeler2 [91] and EDTA [92]

on the assembled genome. Together, these programs include most of the recent and long-

trusted tools used to detect generic repeats, but also include specific modules, such as for

LTR and TIR elements. Preliminary analyses of the *S. oryzae* genome with RepeatModeler1

[123] and dnaPipeTE [46] suggested a rather large fraction of Class II DNA elements with

terminal inverted repeats (TIRs). Thus, MITE-Tracker [124] was incorporated in our pipeline

and ran independently on the genome assembly using 1- and 2-Kbp size cutoffs to detect

Class II elements harboring TIRs with high sensitivity. Following this initial step, 15 510

consensi obtained from RM2, EDTA and the two runs of MITE-tracker were successively

clustered using MAFFT [125]*,* Mothur [126]*,* and Refiner [123] to reduce redundancy in the

repeat library to a total of 2 754 consensus sequences (Additional file 1: Figure S1A,

https://github.com/clemgoub/So2). Then, we inspected the quality of the raw library by

calculating the genomic coverage of each consensus. We ran the library against the genome

using RepeatMasker (52) and implemented a simple algorithm "TE-trimmer.sh" to trim or

split a consensi wherever the genomic support drops below 5% of the average consensus

coverage (Additional file 1: Figure S1A, https://github.com/clemgoub/So2). To mitigate any

redundancy generated by the splitting, the newly trimmed library was clustered before being

re-quantified using RepeatMasker [123]. At this step, we removed any consensus under 200

bp and represented by less than the equivalent of two full-length copies (in total bp). In

addition, TAREAN [127] was used to detect and quantify candidate satellite repeats. We

obtained an *ab-initio* repeat library of 3 950 consensi automatically generated (Additional file

1: Figure S1A).

To refine and improve the quality of the TE consensi, we then turned it over to DFAM who

processed the ab initio library. First, any sequences mostly composed of tandem repeats

were removed using a custom script to remove any sequences that were greater than 80%

masked and/or had a sequence less than 100 bp. To generate seed alignments for each

consensus, the consensi were used as a search library for RepeatMasker to collect

interspersed repeats. Seed alignments in the form of stockholm files were generated using

the RepeatMasker output. To extend potentially truncated elements, the instances in the

stockholm file for each model were extended into neighboring flanking sequences until the

alignment was below a threshold. Following extension, the new consensi were collected and

consensi greater than 80% similar for 80% of their length were considered duplicates and

only one consensus was kept.

Upon completion, we used RepeatMasker to quantify the improved library. We selected

the top 50 elements (by abundance in the genome) represented in each of the "LTR",

"LINE", "DNA" and "Unknown" classes for manual inspection (these categories represent the

4 most abundant classes of repeats in the *S. oryzae* genome). While most consensi where

correctly extended and annotated (200) we noticed some cases of over-extension with LTR

(consensus doubled in size) and flagged others with non-supported fragments for further

trimming (Additional file 2 | tab 1). Once our quality check completed and the sequences

curated, we removed fragments with 100% identity against a previously established

consensus (Additional file 2 | tab 2). The final TE library contains 3 399 sequences to

classify.

The classification of the final repeat library was done in successive rounds combining

homology and structure methods (Additional file 1: Figure S1B). Before the final TE library

was completed, we manually curated and annotated the sequences of 31 transposable

elements and satellites among the most represented in *S. oryzae*. These high-confidence

references were added to the default libraries used by the following programs and Repbase

v.2017 [128]. We searched for nucleotide homology using RepeatMasker (V.4.1.1 [123]) with

-s "-slow" search settings. Best hits were chosen based on the highest score at the

superfamily level allowing non-overlapping hits of related families to contribute to the same

hit. In addition we used blastx [129] to query each consensus against a curated collection of

TE proteins (available with RepeatMasker), as well as those identified in the 31 manual

consensi. We kept the best protein hit based on the blastx score. Based on the 200 consensi

manually inspected (see above), we set a hit length / consensus size threshold of 0.08

(RepeatMasker) and 0.03 (blastx) to keep a hit. In our hands, these thresholds were

conservative to automate the classification. As an alternate homology-based method, we

also ran RepeatClassifier (RepeatModeler2). Finally, because DNA elements are often

represented by non-autonomous copies (unidentifiable or absent transposase) we further

used einverted to flag terminal inverted repeats located less than 100 bp of the ends of each

sequence. The complete library of 3 399 consensi was first annotated at the subclass level

(see DFAM taxonomy: https://dfam.org/classification/tree) if two out of RepeatMasker,

RepeatClassifier and blastx annotations agreed. Further, the same rule was applied for the

superfamilies if possible. At this stage, consensi without annotation by homology but with

TIRs as flagged by einverted, were classified as DNA and all other sequences classified as

Unknown. Finally, the classifications automatically given as "Unknown" to 16/274 manually

inspected consensi were replaced to match the manually reported classification.

Estimation of the repeat content

The total repeat content of the *S. oryzae* genome was analyzed using RepeatMasker

(v.4.1.1) and our classified library of 3 390 consensus sequences and the following

parameters: -s -gccalc -no_is

-cutoff 200. The subsequent alignments were parsed with the script "parseRM.pl" [130]

https://github.com/4ureliek/Parsing-RepeatMasker-Outputs) to remove hits overlap and

statistically analyzed with R version 4.0.2.

Genomic distribution of TE copies

The distribution of TE copies across the *S. oryzae* genome was assessed using two different

approaches over six different genomic regions namely TSS ± 3 Kbp, 5' UTRs, exons,

introns, 3' UTRs and intergenic regions. Briefly, the coverage of each TE subclass/subtype

(DNA, LINE, LTR, PLE, DIRS, RC, Unknown) was computed over a sliding window of 100

bp across the whole genome sequence using the makewindows and coverage tools from the

bedtools package [131] and the bedGraphToBigWig UCSC gtfToGenePred tool. Then the

different genomic regions were retrieved from the *S. oryzae* annotation file (GFF format)

using the gencode_regions script (https://github.com/saketkc/gencode_regions) and the

UCSC gtfToGenePred tool (https://github.com/ENCODE-DCC/kentUtils). A matrix containing

the coverage of each TE subclass per genomic region was generated using the

computeMatrix tool from deepTools [132] and used to generate metaplots using the

plotProfile tool.

Additionally, to assess the landscape of the diverse TE subclasses, we performed circos

ideograms [133] for the scaffolds larger than 10 Mbp (7 scaffolds) of each TE subclass as

well as exon density as reference. The methodology was adapted from Zavallo et al. 2020

[134].

TE landscapes

The relative age of the different TE families identified in the genome assembly was drawn

performing a "TE-landscape" analysis on the RepeatMasker outputs. Briefly, the different

copies of one TE family identified by RepeatMasker are compared to their consensus

sequence and the divergence (Kimura substitution level, CpG adjusted, see RepeatMasker

webpage: http://repeatmasker.org/webrepeatmaskerhelp.html) is calculated. The TE

landscape consists of the distribution of these divergence levels. In the end, the relative age

of a TE family can be seen as its distribution within the landscape graph: "older" TE families

tend to have wider and flatter distribution spreading to the right (higher substitution levels)

than the "recent" TE families, which are found on the left of the graph and have a narrower

distribution. TE landscapes were drawn from the RepeatMasker output parsed with the

options -l of "parseRM.pl". We report here the TE landscape at the level of the TE subclass

(LINE, LTR, DNA, DIRS, PLE, RC and Unknown).

dnaPipeTE comparative analysis in *Sitophilus* species

To compare the TE content of *S. oryzae* to four related species of *Sitophilus* (*S. granarius,*

S. zeamais, S. linearis) we used dnaPipeTE v.1.3 [46]. dnaPipeTE allows unbiased

estimation and comparison of the total repeat content across different species by assembling

and quantifying TE from unassembled reads instead of a linear genome assembly. Reads

for *Sitophilus* species were produced as described above. Using our new classified library (3

390 consensus) as TE database in dnaPipeTE, we were further able to identify the

phylogenetic depth of the repeat identified in *S. oryzae*.

RNA sequencing and TE expression analysis

Adapter sequences and low quality reads were filtered out with Trimmomatic (v0.36) [135]

and clean reads were aligned to the *S. oryzae* genome with STAR aligner (v2.5.4b, [136])

and featureCounts from subread package [137] to obtain gene counts. We also used the

STAR aligner to map the clean reads against all TE copies extracted from the genome with

the following options: --outFilterMultimapNmax 100 --winAnchorMultimapNmax 100 --

outMultimapperOrder Random --outSAMmultNmax 1. The mapped bam files were used as

input to TEtools software [138] to determine TE family expression. Genes and TE family

counts were used as input for DESEq2 package [139] to determine differential TE

expression between Ovary vs Gut tissues as well as Ovaries from symbiotic and

aposymbiotic weevils. Differentially expressed TEs were defined whenever the adjusted p-

value was smaller than 0.05 and Log2 fold change was higher than 1 or smaller than -1. We

used the aforementioned STAR alignment parameters to map transcriptomic sequencing

reads from midgut of *S. oryzae* (Accession: SRX1034971, and SRX1034972), *D.*

melanogaster (Accession: SRX029389, and SRX045361), and *Ae. albopictus* (Accession:

SRX1512976, SRX1898481, SRX1898483, SRX1898487, SRX3939061, and SRX3939054)

against the TE consensus sequences for each species.

Abbreviations

AMPs: AntiMicrobial Peptides

CPs: cuticle proteins

HGT: horizontal gene transfer

IMP: Inosine MonoPhosphate

K2P: Kimura 2 Parameters

LINE: long INterspersed Element

LTR: Long Terminal Repeat

MAMPs: Microbial Associated Molecular Patterns

MITEs: miniature inverted repeat elements

ORs: odorant receptors

PRR: Pattern Recognition Receptors

PLE: penelope-like

RC: rolling circle

SINE: Short INterspersed Element

TIR: terminal inverted repeat

TSS: transcription start sites

TEs: transposable elements

TTSS: Type Three Secretion Systems

UTR: untranslated regions

UMP: Uridine MonoPhosphate

Declarations

Ethics approval and consent to participate

Not applicable

Consent for publication

Not applicable

Availability of data and materials

This Whole Genome Shotgun project has been deposited at DDBJ/ENA/GenBank under the accession PPTJ00000000. The version described in this paper is version PPTJ02000000. The assembly can be visualised, along with gene models and supporting data, on a dedicated genome browser (https://bipaa.genouest.org/sp/sitophilus_oryzae/). Raw reads from low coverage genome sequencing of *S. zeamais*, *S. granarius* and *S. linearis* have been deposited at NCBI Sequence Read Archive (SRA) under the BioProject accessions PRJNA647530, PRJNA647520 and PRJNA647347 respectively. TE annotation (GFF) and consensus sequences can be found at https://dx.doi.org/10.5281/zenodo.4570415. Bisulfite-seq reads have been deposited at NCBI SRA, under the BioProject accession PRJNA681724.

Competing interests

The authors declare that they have no competing interests.

Funding

Funding for this project was provided by the Fondation de l'Institut National des Sciences Appliquées-Lyon (INSA-Lyon), the Institut National de Recherche pour l'Agriculture, l'Alimentation et l'Environnement (INRAE), the French ANR-10-BLAN-1701 (ImmunSymbArt), the French ANR-13-BSV7-0016-01 (IMetSym), the French ANR-17_CE20_0031_01 (GREEN), the French ANR-17-CE20-0015 (UNLEASH), the Santé des Plantes et Environnement (SPE) department at INRAE , the IDEX-Lyon PALSE IMPULSION initiative and a grant from la Région Rhône-Alpes. The project was also funded by European Regional Development Fund (ERDF) and Ministerio de Ciencia, Innovación y Universidades (Spain) PGC2018-099344-B-I00 to AL, and PID2019-105969GB-I00 to AM and Conselleria d'Educació, Generalitat Valenciana (Spain), grant number PROMETEO/2018/133 to AM. CV-C was a recipient of a fellowship from the Ministerio de Economía y Competitividad (Spain).

Authors' contributions

AH and AL conceived the original sequencing project and were joined by NP, RR, CG, CV-C, AM and CV who participated in the coordination of the project. AVa, CV-M, ED, JM, FM and AVi reared the inbred lines and AVa extracted genomic DNA and RNA that was used for library construction and sequencing. BG and SH produced and sequenced the Nanopore libraries. CG, AVa, MB, NB, CV, AG and ATRV produced and sequenced the low-coverage Illumina libraries. NP, CV-C, ADG and M-FS performed the genome assembly and automated gene prediction. CV-C, MM-H and TG analyzed and wrote the *phylome and horizontal gene transfer* note. PB-P, GF, SC, HC and FC analyzed and wrote *the global analysis of metabolic pathways* note. NP analyzed and wrote the *digestive enzymes* and the

detoxification and insecticide resistance notes. PC analyzed and wrote the *development*

note. CV-C analyzed and wrote the *cuticle protein genes* note. CV-M, CV-C, NP, JM, LB,

AB, WZ, FM, AVi and AZ-R analyzed and wrote the *innate immune system* note. NM, CM,

ASB and EJ-J analyzed and wrote the *odorant receptors* note. TC, CB, AVa and RR

produced the data for the *epigenetic pathways* note. TC, CB, AVa, GR, CV-C, CV and RR,

analyzed and wrote the *epigenetic pathways* note. MGF, CG, ED, RR, SB, GF, NM, CV-M

and NP produced the figures. CG, RR, JMS, JR, RH and AFAS annotated and analyzed the

TE content while MGF analyzed the TE RNAseq data. NP, CV-C, CG, CV, RR, AL and AH

wrote the manuscript. All authors read and approved the final manuscript.

Acknowledgments

The authors acknowledge supercomputing resources made available by the Rhône-Alpes

Bioinformatics center (PRABI-AMSB, http://www.prabi.fr) to perform the NGS data analyses.

The authors would like to thank Stéphanie Robin, Fabrice Legeai and Anthony Bretaudeau

at the INRAE BioInformatics Platform for Agro-ecosystems Arthropods (BIPAA)

(https://bipaa.genouest.org) for the *S. oryzae* genome integration. The authors would like to

thank Fabienne Barbet, Séverine Croze and Nicolas Nazaret from profileXpert for Illumina

sequencing of *Sitophilus oryzae* libraries. The authors thank the network REacTION and its

coordinators (N. Ponts, G. Le Trionnaire, I. Fudal, M. Jubault), funded by INRAE-SPE, for

organizing a Bisulfite-seq workshop that allowed the production of the data presented here.

The authors would like to thank Cédric Feschotte at Cornell University for the support,

insights and the occasional use of bioinformatic resources.

Figure legends

Figure 1. *Sitophilus oryzae* overview. A. Life cycle of cereal weevil *Sitophilus oryzae*. The embryo develops into a larva and pupa, and metamorphoses into a young adult, exiting the grain around 3 days after metamorphosis completion. The developmental times indicated are from a rearing condition at 27 °C and 70% relative humidity. B. Photos of adult *S. oryzae*. Lower panel shows an adult exiting the grain.

Figure 2. A. Proportion of repeat content in *S. oryzae*'s genome. The majority of repeats detected in *S. oryzae* are represented by DNA elements, LINEs, and unclassified repeats (unknown). NR: non repetitive. B. Variation of genome size and TE content in 62 insect species from [40] and *S. oryzae*. Coleopteran species are depicted in dark blue, and *S. oryzae* in light blue. *S. oryzae* is clearly a TE-rich genome. C. TE proportion across 11 insect species, including six coleoptera. In agreement with the data used for comparison [40], PLEs are included in the LINE superfamilies, DIRS in LTRs and RC in the DNA superfamilies. NR: non repetitive. *S. oryzae* harbors the largest TE content among Coleopterans and most insect species studied to date. Within Coleoptera, there is a large variation in TE content and type, with *A. planipennis, L. decemlineata* and *O. taurus* carrying an abundant LINE content, while *S. oryzae, T. castaneum* and *A. glabripennis* show larger DNA content. Cladogram based on [95]. D. Classification of the 570 Mbs of TEs present in the *S. oryzae* genome. Most DNA families detected were not classified into known superfamilies. RTE LINE and Gypsy LTR elements are the most abundant superfamilies among retrotransposons. Around 22% of repeats in *S. oryzae*'s genome were not classified by our pipeline, and remain unknown (grey). E. Distribution of TIR length sequences (right) detected by einverted, and the internal region present between both TIRs (left) for annotated DNA superfamilies (color) and unknown DNA families (grey).

Figure 3. TE distribution in *S. oryzae*'s genome. A. Circos plot of the largest scaffolds of the *S. oryzae* genome (> 10 Mbp). Tracks represent features density (percentage of bp covered), from outside to inside: exons, DNA, LTRs, LINE, LTR, PLE, DIRS, RC, and Unknown repeats (0 to 56% coverage with a 7% incremental step). B. Density of TE copies within gene regions. TE copies are depleted within TSSs, 5' UTRs and exons. 3'UTRs show an increase in DNA and unknown elements at their 3' end, and introns and intergenic regions are riddled with TEs. TSS: transcription start site, UTR: untranslated regions. C. Relationship between intron length and TE per intron in D. melanogaster (red), *H. sapiens* (blue) and *S. oryzae* (yellow). *S. oryzae* shares characteristics of both *Drosophila* with short and TE poor introns and Humans with a significant number of large, TE-packed introns.

Figure 4. A. TE divergence landscape. Distribution of the divergence (Kimura two parameters, K2P) between TE copies and their consensus, aggregated by TE class reported in percent of the genome. The less divergent superfamilies are distributed to the left and suggest recent activity. Strikingly, most of the TE copies have less than 10% divergence to their consensus, with a large number of copies under 5%. The distribution of the "unknown" class overlaps with the leftmost mode of the DNA distribution, suggesting that many more DNA families are yet to be described in *S. oryzae.* Strikingly, LTR elements are the least diverged altogether with the mode of the distribution on the 0-1% divergence bin. B. Mean K2P distributions within TE superfamilies. Left panel depicts DNA families (pink), and all other TE families are on the right panel. LTR superfamilies harbor some of the least divergent TE families, suggesting that this class may host some of the youngest TE.

Figure 5. TE family expression in midguts and ovaries from *S. oryzae*. A. Log10 normalized counts in midguts and ovaries triplicates. Normalized counts show different proportions of

transcriptionally active TE families in different TE classes. B. Log10 of base mean average

expression of TE families in ovaries and midguts from three biological replicates. Depicted in

color only TE families which had differential expression between ovary and gut tissues

(padj<0.05, |log2FC|>2). Most TE families are upregulated in midguts compared to ovaries.

C. Distribution of all significant (padj<0.05). Log2FC depicts specifically deregulated TE

classes in each tissue. LTR elements are predominantly upregulated in ovaries.

Figure 6. TE landscape across *Sitophilus* species A. Proportion of TE per species estimated

from short reads with dnaPipeTE and a custom TE library including Repbase (release 2017)

and annotated TE consensus discovered in *S. oryzae*. *S. oryzae* and *S. zeamais* harbor very

similar TE content, while *S. granarius* presents a smaller TE load, and *S. linearis* harbors the

smallest TE content and the higher proportion of unknown repeats. The proportion of

unknown repeats only found by dnaPipeTE (black) increases from *S. oryzae* to *S. linearis*

with the phylogenetic distance. B. Distribution of divergence values between raw reads and

repeats contig assembled with dnaPipeTE (blastn) across four *Sitophilus* species. *S. oryzae*

appears to share its TE landscape with *S. zeamais* and *S. granarius*, but the three species

display a distinct repeatome than *S. linearis,* in spite of their phylogenetic proximity. SO2: *S.*

oryzae's TE library produced in this analysis, DPTE: DNApipeTE TE annotation (repeats

only found by dnaPipeTE).

Figure 7. Differentially expressed TE families between symbiotic and aposymbiotic *S. oryzae*

ovaries. Log10 of base mean average expression of TE families in symbiotic vs

aposymbiotic ovaries from two biological replicates. Depicted in color only TE families which

had differential expression between both ovary types (padj<0.05, |log2FC|>2). Two LTR

elements and one LINE element are upregulated (log2FC > 2) in symbiotic ovaries.

Additional files

Additional file 1: Supplementary notes, supplementary figures, and small tables. (PDF)

Additional file 2: Transposable elements annotation tables. (XLSX)

Additional file 3: STAR and TEtools mapping statistics. (XLSX)

Additional file 4: Summary of sequencing libraries produced for *S. oryzae.* (XLSX)

Additional file 5: Large supporting tables and datasets. (XLSX)

References

1. Hunt T, Bergsten J, Levkanicova Z, Papadopoulou A, John OS, Wild R, et al. A comprehensive phylogeny of beetles reveals the evolutionary origins of a superradiation. Science. 2007;318:1913–6.

2. Stork NE, McBroom J, Gely C, Hamilton AJ. New approaches narrow global species estimates for beetles, insects, and terrestrial arthropods. Proc Natl Acad Sci U S A. 2015;112:7519–23.

3. Hammond P. Species Inventory. In: Groombridge B, editor. Global biodiversity: Status of the Earth's living resources. 1992. Chapman and Hall, London. p. 17–39.

4. McKenna DD, Sequeira AS, Marvaldi AE, Farrell BD. Temporal lags and overlap in the diversification of weevils and flowering plants. Proc Natl Acad Sci U S A. 2009;106:7083–8.

5. Oberprieler RG, Marvaldi AE, Anderson RS. Weevils, weevils, weevils everywhere*. Zootaxa. 2007;1668:491–520.

6. Vega FE, Brown SM, Chen H, Shen E, Nair MB, Ceja-Navarro JA, et al. Draft genome of the most devastating insect pest of coffee worldwide: the coffee berry borer, *Hypothenemus hampei*. Sci Rep. 2015;5:12525.

7. Keeling CI, Yuen MM, Liao NY, Roderick Docking T, Chan SK, Taylor GA, et al. Draft genome of the mountain pine beetle, *Dendroctonus ponderosae* Hopkins, a major forest

pest. Genome Biol. 2013;14:R27.

8. Hazzouri KM, Sudalaimuthuasari N, Kundu B, Nelson D, Al-Deeb MA, Le Mansour A, et al. The genome of pest *Rhynchophorus ferrugineus* reveals gene families important at the plant-beetle interface. Commun Biol. 2020;3:1–14.

9. Zunjare R, Hossain F, Muthusamy V, Jha SK, Kumar P, Sekhar JC, et al. Genetic variability among exotic and indigenous maize inbreds for resistance to stored grain weevil (*Sitophilus oryzae* L.) infestation. Cogent Food Agric. 2016;2:1137156.

10. Longstaff BC. Biology of the grain pest species of the genus *Sitophilus* (Coleoptera: Curculionidae): a critical review. Prot Ecol. 1981;3:83–130.

11. Grenier A-M, Mbaiguinam M, Delobel B. Genetical analysis of the ability of the rice weevil *Sitophilus oryzae* (Coleoptera, Curculionidae) to breed on split peas. Heredity. 1997;79:15–23.

12. Champ BR, Dyte CE. FAO global survey of pesticide susceptibility of stored grain pests. FAO Plant Protec Bull. 1977;25(2):49-67.

13. Nguyen TT, Collins PJ, Ebert PR. Inheritance and characterization of strong resistance to phosphine in *Sitophilus oryzae* (L.). PLoS One. 2015;10:e0124335.

14. Mills KA. Phosphine resistance: Where to now? In: Donahaye, EJ, Navarro, S and Leesch JG, editors. Proceeding International Conference on Controlled Atmosphere and Fumigation in Stored Products; 2000 Oct 29-Nov 3; Fresno, USA. 2000:583–91.

15. Campbell JF. Fitness consequences of multiple mating on female *Sitophilus oryzae* L. (Coleoptera: Curculionidae). Environ Entomol. 2005;34:833–43.

16. Oakeson KF, Gil R, Clayton AL, Dunn DM, von Niederhausern AC, Hamil C, et al. Genome degeneration and adaptation in a nascent stage of symbiosis. Genome Biol Evol. 2014;6:76–93.

17. Heddi A, Charles H, Khatchadourian C, Bonnot G, Nardon P. Molecular characterization of the principal symbiotic bacteria of the weevil *Sitophilus oryza*e: a peculiar G + C content of an endocytobiotic DNA. J Mol Evol. 1998;47:52–61.

18. Heddi A, Charles H, Khatchadourian C. Intracellular bacterial symbiosis in the genus

Sitophilus: the 'biological individual' concept revisited. Res Microbiol. 2001;152:431–7.

19. Lefèvre C, Charles H, Vallier A, Delobel B, Farrell B, Heddi A. Endosymbiont

phylogenesis in the Dryophthoridae weevils: evidence for bacterial replacement. Mol Biol

Evol. 2004;21:965–73.

20. Clayton AL, Oakeson KF, Gutin M, Pontes A, Dunn DM, Niederhausern AC von, et al. A

novel human-infection-derived bacterium provides insights into the evolutionary origins of

mutualistic insect–bacterial symbioses. PLoS Genet. 2012;8:e1002990.

21. Akman L, Yamashita A, Watanabe H, Oshima K, Shiba T, Hattori M, et al. Genome

sequence of the endocellular obligate symbiont of tsetse flies, *Wigglesworthia glossinidia*.

Nat Genet. 2002;32:402–7.

22. Shigenobu S, Watanabe H, Hattori M, Sakaki Y, Ishikawa H. Genome sequence of the

endocellular bacterial symbiont of aphids *Buchnera sp.* APS. Nature. 2000;407:81–6.

23. Gil R, Belda E, Gosalbes MJ, Delaye L, Vallier A, Vincent-Monégat C, et al. Massive

presence of insertion sequences in the genome of SOPE, the primary endosymbiont of the

rice weevil *Sitophilus oryzae*. Int Microbiol Off J Span Soc Microbiol. 2008;11:41–8.

24. Rebollo R, Romanish MT, Mager DL. Transposable elements: an abundant and natural

source of regulatory sequences for host genes. Annu Rev Genet. 2012;46:21–42.

25. Bourque G, Burns KH, Gehring M, Gorbunova V, Seluanov A, Hammell M, et al. Ten

things you should know about transposable elements. Genome Biol. 2018;19:199.

26. Chuong EB, Elde NC, Feschotte C. Regulatory activities of transposable elements: from

conflicts to benefits. Nat Rev Genet. 2017;18:71–86.

27. Chen S, Li X. Transposable elements are enriched within or in close proximity to

xenobiotic-metabolizing cytochrome P450 genes. BMC Evol Biol. 2007;7:46.

28. You M, Yue Z, He W, Yang X, Yang G, Xie M, et al. A heterozygous moth genome

provides insights into herbivory and detoxification. Nat Genet. 2013;45:220–5.

29. Singh KS, Troczka BJ, Duarte A, Balabanidou V, Trissi N, Paladino LZC, et al. The

genetic architecture of a host shift: An adaptive walk protected an aphid and its

endosymbiont from plant chemical defenses. Sci Adv. 2020;6:eaba1070.

30. Carareto CMA, Hernandez EH, Vieira C. Genomic regions harboring insecticide
resistance-associated Cyp genes are enriched by transposable element fragments carrying
putative transcription factor binding sites in two sibling *Drosophila* species. Gene.
2014;537:93–9.

31. Rostant WG, Wedell N, Hosken DJ. Chapter 2 - Transposable elements and insecticide
resistance. In: Goodwin SF, Friedmann T, Dunlap JC, editors. Adv Genet. Academic Press;
2012. p. 169–201.

32. Mateo L, Ullastres A, González J. A transposable element insertion confers xenobiotic
resistance in *Drosophila*. PLoS Genet. 2014;10:e1004560.

33. Rech GE, Bogaerts-Márquez M, Barrón MG, Merenciano M, Villanueva-Cañas JL,
Horváth V, et al. Stress response, behavior, and development are shaped by transposable
element-induced mutations in *Drosophila*. PLoS Genet. 2019;15:e1007900.

34. Ullastres A, Merenciano M, González J. Natural transposable element insertions drive
expression changes in genes underlying *Drosophila* immune response. bioRxiv.
2019;655225.

35. Schnable PS, Ware D, Fulton RS, Stein JC, Wei F, Pasternak S, et al. The B73 maize
genome: complexity, diversity, and dynamics. Science. 2009;326:1112–5.

36. Lander ES, Linton LM, Birren B, Nusbaum C, Zody MC, Baldwin J, et al. Initial
sequencing and analysis of the human genome. Nature. 2001;409:860–921.

37. Meyer A, Schloissnig S, Franchini P, Du K, Woltering JM, Irisarri I, et al. Giant lungfish
genome elucidates the conquest of land by vertebrates. Nature. 2021;1–6.

38. Adams MD, Celniker SE, Holt RA, Evans CA, Gocayne JD, Amanatides PG, et al. The
genome sequence of *Drosophila melanogaster*. Science. 2000;287:2185–95.

39. The *Arabidopsis* genome initiative. Analysis of the genome sequence of the flowering
plant *Arabidopsis thaliana*. Nature. 2000;408:796–815.

40. Petersen M, Armisén D, Gibbs RA, Hering L, Khila A, Mayer G, et al. Diversity and
evolution of the transposable element repertoire in arthropods with particular reference to
insects. BMC Evol Biol. 2019;19:11.

41. Wang X, Fang X, Yang P, Jiang X, Jiang F, Zhao D, et al. The locust genome provides

insight into swarm formation and long-distance flight. Nat Commun. 2014;5:2957.

42. Kelley JL, Peyton JT, Fiston-Lavier A-S, Teets NM, Yee M-C, Johnston JS, et al.

Compact genome of the Antarctic midge is likely an adaptation to an extreme environment.

Nat Commun. 2014;5:4611.

43. Gilbert C, Peccoud J, Cordaux R. Transposable elements and the evolution of insects.

Annu Rev Entomol. 2021;66:355-372.

44. Sessegolo C, Burlet N, Haudry A. Strong phylogenetic inertia on genome size and

transposable element content among 26 species of flies. Biol Lett. 2016;12:20160407.

45. Ray DA, Grimshaw JR, Halsey MK, Korstian JM, Osmanski AB, Sullivan KAM, et al.

Simultaneous TE analysis of 19 Heliconiine butterflies yields novel insights into rapid TE-

based genome diversification and multiple SINE births and deaths. Genome Biol Evol.

2019;11:2162–77.

46. Goubert C, Modolo L, Vieira C, ValienteMoro C, Mavingui P, Boulesteix M. De Novo

Assembly and annotation of the Asian tiger mosquito (*Aedes albopictus*) repeatome with

dnaPipeTE from raw genomic reads and comparative analysis with the Yellow fever

mosquito (*Aedes aegypti*). Genome Biol Evol. 2015;7:1192–205.

47. Nene V, Wortman JR, Lawson D, Haas B, Kodira C, Tu Z (Jake), et al. Genome

sequence of *Aedes aegypti*, a major arbovirus vector. Science. 2007;316:1718–23.

48. Zhang S, Shen S, Peng J, Zhou X, Kong X, Ren P, et al. Chromosome-level genome

assembly of an important pine defoliator, *Dendrolimus punctatus* (Lepidoptera;

Lasiocampidae). Mol Ecol Resour. 2020;20:1023–37.

49. Silva AA, Braga LS, Corrêa AS, Holmes VR, Johnston JS, Oppert B, et al. Comparative

cytogenetics and derived phylogenic relationship among *Sitophilus* grain weevils

(Coleoptera, Curculionidae, Dryophthorinae). Comp Cytogenet. 2018;12:223–45.

50. Seppey M, Manni M, Zdobnov EM. BUSCO: Assessing genome assembly and

annotation completeness. In: Kollmar M, editor. Gene Prediction. Methods Mol Biol.

2019;1962. p. 227–45.

51. McKenna DD, Scully ED, Pauchet Y, Hoover K, Kirsch R, Geib SM, et al. Genome of the

Asian longhorned beetle (*Anoplophora glabripennis*), a globally significant invasive species,

reveals key functional and evolutionary innovations at the beetle-plant interface. Genome

Biol. 2016;17:227.

52. *Tribolium* genome sequencing consortium, Richards S, Gibbs RA, Weinstock GM, Brown

SJ, Denell R, et al. The genome of the model beetle and pest *Tribolium castaneum*. Nature.

2008;452:949–55.

53. Al-Qahtani AH, Al-Khalifa MS, Al-Saleh AA. Karyotype, meiosis and sperm formation in

the red palm weevil *Rhynchophorus ferrugineus*. Cytologia. 2014;79:235–42.

54. Brun LO, Stuart J, Gaudichon V, Aronstein K, French-Constant RH. Functional

haplodiploidy: a mechanism for the spread of insecticide resistance in an important

international insect pest. Proc Natl Acad Sci. U S A. 1995;92:9861–5.

55. Lanier GN, Wood DL. Controlled mating, karyology, morphology, and sex-ratio in the

Dendroctonus ponderosae complex. Ann Entomol Soc Am. 1968;61:517–26.

56. Stuart JJ, Mocelin G. Cytogenetics of chromosome rearrangements in *Tribolium*

castaneum. Genome. 1995;38(4):673-80.

57. Initiative IGG. Genome sequence of the Tsetse fly (*Glossina morsitans*): Vector of

African trypanosomiasis. Science. 2014;344:380–6.

58. Vellozo AF, Véron AS, Baa-Puyoulet P, Huerta-Cepas J, Cottret L, Febvay G, et al.

CycADS: an annotation database system to ease the development and update of BioCyc

databases. Database. 2011;2011:bar008.

59. Karp PD, Midford PE, Billington R, Kothari A, Krummenacker M, Latendresse M, et al.

Pathway Tools version 23.0 update: software for pathway/genome informatics and systems

biology. Brief Bioinform. 2019; 22(1):109-126.

60. Vigneron A, Masson F, Vallier A, Balmand S, Rey M, Vincent-Monégat C, et al. Insects

recycle endosymbionts when the benefit is over. Curr Biol. 2014;24:2267–73.

61. Heddi A, Grenier A-M, Khatchadourian C, Charles H, Nardon P. Four intracellular

genomes direct weevil biology: Nuclear, mitochondrial, principal endosymbiont, and

Wolbachia. Proc Natl Acad Sci U S A. 1999;96:6814–9.

62. Grenier AM, Nardon C, Nardon P. The role of symbiotes in flight activity of *Sitophilus*

weevils. Entomol Exp Appl. 1994;70:201–8.

63. Rio RVM, Lefevre C, Heddi A, Aksoy S. Comparative genomics of insect-symbiotic

bacteria: influence of host environment on microbial genome composition. Appl Environ

Microbiol. 2003;69:6825–32.

64. Jasrapuria S, Arakane Y, Osman G, Kramer KJ, Beeman RW, Muthukrishnan S. Genes

encoding proteins with peritrophin A-type chitin-binding domains in *Tribolium castaneum* are

grouped into three distinct families based on phylogeny, expression and function. Insect

Biochem Mol Biol. 2010;40:214–27.

65. Jasrapuria S, Specht CA, Kramer KJ, Beeman RW, Muthukrishnan S. Gene families of

cuticular proteins analogous to peritrophins (CPAPs) in *Tribolium castaneum* have diverse

functions. PLoS One. 2012;7:e49844.

66. Gerardo NM, Altincicek B, Anselme C, Atamian H, Barribeau SM, de Vos M, et al.

Immunity and other defenses in pea aphids, *Acyrthosiphon pisum*. Genome Biol.

2010;11:R21.

67. Zhang C-R, Zhang S, Xia J, Li F-F, Xia W-Q, Liu S-S, et al. The immune strategy and

stress response of the mediterranean species of the *Bemisia tabaci* complex to an orally

delivered bacterial pathogen. PLoS One. 2014;9:e94477.

68. Salcedo-Porras N, Guarneri A, Oliveira PL, Lowenberger C. *Rhodnius prolixus*:

Identification of missing components of the IMD immune signaling pathway and functional

characterization of its role in eliminating bacteria. PLoS One. 2019;14:e0214794.

69. Maire J, Vincent-Monégat C, Masson F, Zaidman-Rémy A, Heddi A. An IMD-like

pathway mediates both endosymbiont control and host immunity in the cereal weevil

Sitophilus spp. Microbiome. 2018;6:6.

70. Maire J, Vincent-Monégat C, Balmand S, Vallier A, Hervé M, Masson F, et al. Weevil

pgrp-lb prevents endosymbiont TCT dissemination and chronic host systemic immune

activation. Proc Natl Acad Sci U S A. 2019;116:5623–32.

71. Chaudhry MQ. Phosphine resistance. Pestic Outlook. 2000;11:88–91.

72. Chaudhry MQ. A Review of the mechanisms involved in the action of phosphine as an

insecticide and phosphine resistance in stored-product insects. Pestic Sci. 1997;49:213–28.

73. Athié I, Gomes RAR, Bolonhezi S, Valentini SRT, De Castro MFPM. Effects of carbon

dioxide and phosphine mixtures on resistant populations of stored-grain insects. J Stored

Prod Res. 1998;34:27–32.

74. Rajendran S. Phosphine resistance in stored grain insect pests in India. Proc 7th Int

Work Conf Stored-Prod Prot. 1998. p. 14–19.

75. Zeng L. Development and countermeasures of phosphine resistance in stored grain

insects in Guangdong, China, 642–647. Proc Seventh Int Work Conf Stored-Prod Prot Eds J

Zuxun Quan Yongsheng T Xianchang G Lianghua14–19 Oct 1998 Beijing China Sichuan

Publ House Sci Technol Chengdu China. 1999.

76. Benhalima H, Chaudhry MQ, Mills KA, Price NR. Phosphine resistance in stored-product

insects collected from various grain storage facilities in Morocco. J Stored Prod Res.

2004;40:241–9.

77. Pimentel MAG, Faroni LRD, Silva FH da, Batista MD, Guedes RNC. Spread of

phosphine resistance among brazilian populations of three species of stored product insects.

Neotrop Entomol. 2010;39:101–7.

78. Nguyen TT, Collins PJ, Duong TM, Schlipalius DI, Ebert PR. Genetic conservation of

phosphine resistance in the rice weevil *Sitophilus oryzae* (L.). J Hered. 2016;107:228–37.

79. Holloway JC, Falk MG, Emery RN, Collins PJ, Nayak MK. Resistance to phosphine in

Sitophilus oryzae in Australia: A national analysis of trends and frequencies over time and

geographical spread. J Stored Prod Res. 2016;69:129–37.

80. Agrafioti P, Athanassiou CG, Nayak MK. Detection of phosphine resistance in major

stored-product insects in Greece and evaluation of a field resistance test kit. J Stored Prod

Res. 2019;82:40–7.

81. Carey AF, Carlson JR. Insect olfaction from model systems to disease control. Proc Natl

Acad Sci U S A. 2011;108:12987–95.

82. Andersson MN, Newcomb RD. Pest control compounds targeting insect

chemoreceptors: Another silent spring? Front Ecol Evol. 2017;5:5.

83. Leal WS. Odorant reception in insects: roles of receptors, binding proteins, and

degrading enzymes. Annu Rev Entomol. 2013;58:373–91.

84. Hassanali A, Herren H, Khan Z, Pickett J, Woodcock C. Integrated pest management:

The push-pull approach for controlling insect pests and weeds of cereals, and its potential

for other agricultural systems including animal husbandry. Philos Trans R Soc Lond B Biol

Sci. 2008;363:611–21.

85. Hatano E, Saveer AM, Borrero-Echeverry F, Strauch M, Zakir A, Bengtsson M, et al. A

herbivore-induced plant volatile interferes with host plant and mate location in moths through

suppression of olfactory signalling pathways. BMC Biol. 2015;13:75.

86. Ukeh DA, Woodcock CM, Pickett JA, Birkett MA. Identification of host kairomones from

maize, *Zea mays*, for the maize weevil, *Sitophilus zeamais*. J Chem Ecol. 2012;38:1402–9.

87. Germinara GS, De Cristofaro A, Rotundo G. Behavioral responses of adult *Sitophilus*

granarius to individual cereal volatiles. J Chem Ecol. 2008;34:523–9.

88. Phillips JK, Walgenbach CA, Klein JA, Burkholder WE, Schmuff NR, Fales HM. (R (*),S

(*))-5-hydroxy-4-methyl-3-heptanone male-produced aggregation pheromone of *Sitophilus*

oryzae (L.) and *S. zeamais* motsch. J Chem Ecol. 1985;11:1263–74.

89. Schmuff NR, Phillips JK, Burkholder WE, Fales HM, Chen C-W, Roller PP, et al. The

chemical identification of the rice weevil and maize weevil aggregation pheromone.

Tetrahedron Lett. 1984;25:1533–4.

90. Mitchell RF, Schneider TM, Schwartz AM, Andersson MN, McKenna DD. The diversity

and evolution of odorant receptors in beetles (Coleoptera). Insect Mol Biol. 2020;29:77–91.

91. Flynn JM, Hubley R, Goubert C, Rosen J, Clark AG, Feschotte C, et al. RepeatModeler2

for automated genomic discovery of transposable element families. Proc Natl Acad Sci U S

A. 2020;117:9451–7.

92. Ou S, Su W, Liao Y, Chougule K, Agda JRA, Hellinga AJ, et al. Benchmarking

transposable element annotation methods for creation of a streamlined, comprehensive

pipeline. Genome Biol. 2019;20:275.

93. Hernandez-Hernandez EM, Fernández-Medina RD, Navarro-Escalante L, Nuñez J,

Benavides-Machado P, Carareto CMA. Genome-wide analysis of transposable elements in

the coffee berry borer *Hypothenemus hampei* (Coleoptera: Curculionidae): description of

novel families. Mol Genet Genomics. 2017;292:565–83.

94. Ic A, Es M, Rc M, Gl W. Diverse mobilome of *Dichotomius* (*Luederwaldtinia*) *schiffleri*

(Coleoptera: Scarabaeidae) reveals long-range horizontal transfer events of DNA

transposons. Mol Genet Genomics. 2020;295(6):1339-1353.

95. Misof B, Liu S, Meusemann K, Peters RS, Donath A, Mayer C, et al. Phylogenomics

resolves the timing and pattern of insect evolution. Science. 2014;346:763–7.

96. Feschotte C, Mouchès C. Recent amplification of miniature inverted-repeat transposable

elements in the vector mosquito *Culex pipiens*: characterization of the Mimo family. Gene.

2000;250:109–16.

97. Feschotte C, Zhang X, Wessler SR. Miniature inverted-repeat transposable elements

and their relationship to established DNA transposons. Mob DNA II. 2002;1147–58.

98. Sela N, Kim E, Ast G. The role of transposable elements in the evolution of non-

mammalian vertebrates and invertebrates. Genome Biol. 2010;11:R59.

99. Pasyukova EG, Nuzhdin SV. Doc and copia instability in an isogenic *Drosophila*

melanogaster stock. Mol Gen Genet. 1993;240:302–6.

100. Ashburner M, Bergman CM. *Drosophila melanogaster*: a case study of a model

genomic sequence and its consequences. Genome Res. 2005;15:1661–7.

101. Czech B, Hannon GJ. One loop to rule them all: The ping-pong cycle and piRNA-

guided silencing. Trends Biochem Sci. 2016;41:324–37.

102. Sienski G, Dönertas D, Brennecke J. Transcriptional silencing of transposons by Piwi

and Maelstrom and its impact on chromatin state and gene expression. Cell. 2012;151:964–

80.

103. Andersen PR, Tirian L, Vunjak M, Brennecke J. A heterochromatin-dependent

transcription machinery drives piRNA expression. Nature. 2017;549:54–9.

104. Lynch M, Conery JS. The origins of genome complexity. Science. 2003;302:1401–4.

105. Peona V, Blom MPK, Xu L, Burri R, Sullivan S, Bunikis I, et al. Identifying the causes

and consequences of assembly gaps using a multiplatform genome assembly of a bird-of-

paradise. Mol Ecol Resour. 2021;21(1):263-286.

106. Di Genova A, Buena-Atienza E, Ossowski S, Sagot M-F. Efficient hybrid *de novo*

assembly of human genomes with WENGAN. Nat Biotechnol. 2020;1–9.

107. Platt RN II, Blanco-Berdugo L, Ray DA. Accurate transposable element annotation is

vital when analyzing new genome assemblies. Genome Biol Evol. 2016;8:403–10.

108. Maire J, Parisot N, Galvao Ferrarini M, Vallier A, Gillet B, Hughes S, et al. Spatial and

morphological reorganization of endosymbiosis during metamorphosis accommodates adult

metabolic requirements in a weevil. Proc Natl Acad Sci U S A. 2020;117:19347–58.

109. Login FH, Balmand S, Vallier A, Vincent-Monégat C, Vigneron A, Weiss-Gayet M, et al.

Antimicrobial peptides keep insect endosymbionts under control. Science. 2011;334:362–5.

110. Nardon P. Obtention d'une souche asymbiotique chez le charançon *Sitophilus sasakii*

Tak: différentes méthodes d'obtention et comparaison avec la souche symbiotique d'origine.

CR Acad Sci Paris D. 1973;277:981–4.

111. Li H. BFC: correcting Illumina sequencing errors. Bioinformatics. 2015;31:2885–7.

112. Salmela L, Rivals E. LoRDEC: accurate and efficient long read error correction.

Bioinformatics. 2014;30:3506–14.

113. Magoc T, Salzberg SL. FLASH: fast length adjustment of short reads to improve

genome assemblies. Bioinformatics. 2011;27:2957–63.

114. Chikhi R, Rizk G. Space-efficient and exact de Bruijn graph representation based on a

Bloom filter. Algorithms Mol Biol. 2013;8(1):22.

115. Walker BJ, Abeel T, Shea T, Priest M, Abouelliel A, Sakthikumar S, et al. Pilon: An

integrated tool for comprehensive microbial variant detection and genome assembly

improvement. PLoS One. 2014;9:e112963.

116. Di Genova A, Ruz GA, Sagot M-F, Maass A. Fast-SG: an alignment-free algorithm for

hybrid assembly. GigaScience. 2018;7(5):giy048.

117. Mandric I, Zelikovsky A. ScaffMatch: scaffolding algorithm based on maximum weight

matching. Bioinformatics. 2015;31:2632–8.

118. Xu G-C, Xu T-J, Zhu R, Zhang Y, Li S-Q, Wang H-W, et al. LR_Gapcloser: a tiling path-

based gap closer that uses long reads to complete genome assembly. GigaScience.

2019;8(1):giy157.

119. Song L, Shankar DS, Florea L. Rascaf: Improving genome assembly with RNA

sequencing data. Plant Genome. 2016;9:1–12.

120. Paulino D, Warren RL, Vandervalk BP, Raymond A, Jackman SD, Birol I. Sealer: a

scalable gap-closing application for finishing draft genomes. BMC Bioinformatics.

2015;16:230.

121. Mikheenko A, Prjibelski A, Saveliev V, Antipov D, Gurevich A. Versatile genome

assembly evaluation with QUAST-LG. Bioinformatics. 2018;34:i142–50.

122. Kokot M, Długosz M, Deorowicz S. KMC 3: counting and manipulating k-mer statistics.

Bioinformatics. 2017;33:2759–61.

123. Smit AF, Hubley R, Green P. RepearMasker Open-4.0. http://www.repeatmasker.org.

2013;

124. Crescente JM, Zavallo D, Helguera M, Vanzetti LS. MITE Tracker: an accurate

approach to identify miniature inverted-repeat transposable elements in large genomes.

BMC Bioinformatics. 2018;19:348.

125. Katoh K, Standley DM. MAFFT multiple sequence alignment software version 7:

improvements in performance and usability. Mol Biol Evol. 2013;30:772–80.

126. Schloss PD, Westcott SL, Ryabin T, Hall JR, Hartmann M, Hollister EB, et al.

Introducing mothur: Open-source, platform-independent, community-supported software for

describing and comparing microbial communities. Appl Environ Microbiol. 2009;75:7537–41.

127. Novák P, Ávila Robledillo L, Koblížková A, Vrbová I, Neumann P, Macas J. TAREAN: a

computational tool for identification and characterization of satellite DNA from unassembled

short reads. Nucleic Acids Res. 2017;45:e111.

128. Bao W, Kojima KK, Kohany O. Repbase Update, a database of repetitive elements in

eukaryotic genomes. Mob DNA. 2015;6:11.

129. Camacho C, Coulouris G, Avagyan V, Ma N, Papadopoulos J, Bealer K, et al. BLAST+:

architecture and applications. BMC Bioinformatics. 2009;10:421.

130. Kapusta A, Suh A. Evolution of bird genomes—a transposon's-eye view. Ann N Y Acad

Sci. 2017;1389:164–85.

131. Quinlan AR, Hall IM. BEDTools: a flexible suite of utilities for comparing genomic

features. Bioinformatics. 2010;26:841–2.

132. Ramírez F, Ryan DP, Grüning B, Bhardwaj V, Kilpert F, Richter AS, et al. deepTools2:

a next generation web server for deep-sequencing data analysis. Nucleic Acids Res.

2016;44:W160–5.

133. Krzywinski MI, Schein JE, Birol I, Connors J, Gascoyne R, Horsman D, et al. Circos: An

information aesthetic for comparative genomics. Genome Res. 2009;19(9):1639-45.

134. Zavallo D, Crescente JM, Gantuz M, Leone M, Vanzetti LS, Masuelli RW, et al.

Genomic re-assessment of the transposable element landscape of the potato genome. Plant

Cell Rep. 2020;39:1161–74.

135. Bolger AM, Lohse M, Usadel B. Trimmomatic: a flexible trimmer for Illumina sequence

data. Bioinformatics. 2014;30:2114–20.

136. Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, et al. STAR: ultrafast

universal RNA-seq aligner. Bioinformatics. 2013;29:15–21.

137. Liao Y, Smyth GK, Shi W. featureCounts: an efficient general purpose program for

assigning sequence reads to genomic features. Bioinformatics. 2014;30:923–30.

138. Lerat E, Fablet M, Modolo L, Lopez-Maestre H, Vieira C. TEtools facilitates big data

expression analysis of transposable elements and reveals an antagonism between their

activity and that of piRNA genes. Nucleic Acids Res. 2017;45:e17.

139. Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for

RNA-seq data with DESeq2. Genome Biol. 2014;15:550.